


# Aufgaben zur Topologie

Prof. Dr. C.-F. Bödigheimer

Wintersemester 2016/17

Week 9 — Long exact homology sequences

Due: 11. January 2017



Henry Poincaré and his red-nosed reindeer called Rudolph Homology.

## Exercise 9.1 (The connecting homomorphism I)

Let

$$0 \rightarrow A_\bullet \xrightarrow{f_\bullet} B_\bullet \xrightarrow{g_\bullet} C_\bullet \rightarrow 0$$

be a short exact sequence of chain complexes over  $\mathbb{K}$  and denote by  $\partial_*: H_n(C_\bullet) \rightarrow H_{n-1}(A_\bullet)$  the connecting homomorphism.

- $\partial_*$  is a homomorphism of  $\mathbb{K}$ -modules.
- $\partial_*$  is natural.

## Exercise 9.2 (Induced maps on homology in degree zero.)

Let  $X$  be a space and denote by  $\pi_0(X)$  the set of path-components of  $X$ .

- Show that there is an isomorphism  $H_0(X) \cong \mathbb{Z}\langle\pi_0(X)\rangle := \bigoplus_{\pi_0(X)} \mathbb{Z}$ , the free abelian group generated by the set  $\pi_0(X)$ .
- Show that any continuous map  $f: X \rightarrow Y$  induces a well-defined function  $\pi_0(f): \pi_0(X) \rightarrow \pi_0(Y)$ .
- If we identify  $H_0(X)$  with  $\mathbb{Z}\langle\pi_0(X)\rangle$  and  $H_0(Y)$  with  $\mathbb{Z}\langle\pi_0(Y)\rangle$  as above, then the homomorphisms  $H_0(f)$  and  $\mathbb{Z}\langle\pi_0(f)\rangle$  are equal.

## Exercise 9.3 (Relative homology in degree zero.)

Let  $A$  be a subspace of  $X$ ; we denote by  $i: A \rightarrow X$  the inclusion and by  $q: X \rightarrow X/A$  the quotient map identifying  $A$  to a point. (What does this mean, when  $A = \emptyset$ ?). For the sets of path-components we have induced functions  $\pi_0(i): \pi_0(A) \rightarrow \pi_0(X)$  and  $\pi_0(q): \pi_0(X) \rightarrow \pi_0(X/A)$ .

- Show that the relative homology group  $H_0(X, A)$  is isomorphic to  $\mathbb{Z}\langle\pi_0(X/A)\rangle$ .
- Show that  $H_0(i): H_0(A) \rightarrow H_0(X)$  is injective, that  $H_0(q): H_0(X) \rightarrow H_0(X, A)$  is surjective, and that  $H_0(X, A)$  is isomorphic to the quotient module  $H_0(X)/H_0(A)$ .

(3) Describe  $H_1(X, A)$  and the connecting homomorphism  $\partial_*: H_1(X, A) \rightarrow H_0(A)$ .

**Exercise 9.4** (Torus with one boundary curve.)


Let  $X$  be a torus with one boundary curve, as shown in the figure. We denote by  $A$  the boundary curve, by  $i: A \rightarrow X$  the inclusion of spaces, by  $S_\bullet(i): S_\bullet(A) \rightarrow S_\bullet(X)$  the inclusion of singular chain complexes, and by  $q_\bullet: S_\bullet(X) \rightarrow S_\bullet(X, A) = S_\bullet(X)/S_\bullet(A)$  the quotient homomorphism of singular chain complexes; thus we have the short exact sequence of chain complexes

$$0 \rightarrow S_\bullet(A) \xrightarrow{S_\bullet(i)} S_\bullet(X) \xrightarrow{q_\bullet} S_\bullet(X, A) \rightarrow 0$$

leading to a long exact sequence of homology groups:

$$\dots \rightarrow H_2(A) \xrightarrow{i_*} H_2(X) \xrightarrow{q_*} H_2(X, A) \xrightarrow{\partial_*} H_1(A) \xrightarrow{i_*} H_1(X) \xrightarrow{q_*} H_1(X, A) \xrightarrow{\partial_*} H_0(A) \xrightarrow{i_*} H_0(X) \xrightarrow{q_*} H_0(X, A) \rightarrow 0$$

- (1) Compute all homology groups in degrees 0 and 1 and all homomorphism between them.
- (2) Show that  $H_2(X, A)$  contains at least a summand which is isomorphic to  $\mathbb{Z}$  and generated by the relative cycle  $D = D_1 + \dots + D_6$ .
- (3) Compute for the connecting homomorphism  $\partial_*([D]) = [c_1 + c_2]$ .



A torus, written as a square with dark edges  $a$  and  $b$  identified. Some singular 1- and 2-simplices, used in Exercise 9.4, are shown.

**Exercise 9.5** (Calculating with long exact sequences of abelian groups.)

(1) Show that, if  $0 \rightarrow A \xrightarrow{\alpha} B \xrightarrow{\beta} C \xrightarrow{\gamma} D \rightarrow 0$  is an exact sequence, then we also have exact sequences:

(a)  $0 \rightarrow B/A \xrightarrow{\bar{\beta}} C \xrightarrow{\gamma} D \rightarrow 0$  and

(b)  $0 \rightarrow A \xrightarrow{\alpha} B \xrightarrow{\beta} \text{image}(\beta) \rightarrow 0$ .

(2) More generally, if we have an exact sequence  $\cdots \rightarrow A \xrightarrow{e} B \xrightarrow{f} C \xrightarrow{g} D \xrightarrow{h} E \rightarrow \cdots$ , then we may “localise” it at  $C$ , meaning that there is a short exact sequence:

$$0 \rightarrow \text{coker}(e) \xrightarrow{\bar{f}} C \xrightarrow{g} \ker(h) \rightarrow 0,$$

where  $\text{coker}(e)$  means  $B/e(A)$ .

(3) Show that, if  $0 \rightarrow A \rightarrow B \rightarrow \mathbb{Z}^k \rightarrow 0$  is exact, then  $B$  is isomorphic to  $\mathbb{Z}^k \oplus A$ .

(Hint: start by constructing a homomorphism  $\mathbb{Z}^k \rightarrow B$ , which is a right-inverse for the given homomorphism  $B \rightarrow \mathbb{Z}^k$ .)

(4) Give an example of a short exact sequence  $0 \rightarrow A \rightarrow B \rightarrow \mathbb{Z}/k\mathbb{Z}$  (for any  $k \neq -1, 0, +1$ ) for which  $B$  is not isomorphic to  $A \oplus \mathbb{Z}/k\mathbb{Z}$ .

**Exercise 9.6\*** (Connecting homomorphisms II)

Assume we have a short exact sequence

$$(*) \quad 0 \rightarrow A_{\bullet} \xrightarrow{\alpha_{\bullet}} B_{\bullet} \xrightarrow{\beta_{\bullet}} C_{\bullet} \rightarrow 0$$

of free chain complexes; we know that  $B_n \cong A_n \oplus C_n$ , since  $C_n$  is free. But the boundary operator  $\partial^B: B_n \rightarrow B_{n-1}$  may not be the direct sum of  $\partial^A$  and  $\partial^C$ , i.e., not of diagonal block form; but it must have the form

$$\partial^B = \begin{pmatrix} \partial^A & \varphi_n \\ 0 & \partial^C \end{pmatrix}$$

for some family of homomorphisms  $\varphi_n: C_n \rightarrow A_{n-1}$ .

(1) Which condition must this family satisfy, such that  $\partial^B \circ \partial^B = 0$ ?

(2) Compute the connecting homomorphism in the long exact homology sequence of (\*).