

REPRESENTATIONS OF THE KOTTWITZ GERBES

SERGEI IAKOVENKO

INTRODUCTION

Let F be a local or global field, and let K/F be a finite Galois extension of degree n . In both settings there is a notion of duality for Tate cohomology groups that reduces a computation of Brauer groups to an easy computation with characters of a (pro-)torus $\mathbb{D}_{K/F}$. Namely, when F is local, then $\mathbb{D}_{K/F}$ is simply the multiplicative group \mathbb{G}_m , and a cup product with the local canonical class $\alpha_{K/F}$

$$H^0(\mathrm{Gal}(K/F), X^*(\mathbb{G}_m)) \xrightarrow{\cup \alpha_{K/F}} H^2(\mathrm{Gal}(K/F), \mathbb{G}_m(K)) = \mathrm{Br}(K/F) \quad (0.1)$$

is an isomorphism, note that $\alpha_{K/F}$ can be identified with a generator of $\mathrm{Br}(K/F) \simeq \mathbb{Z}/n\mathbb{Z}$. The case where F is a global field, is somewhat more complicated and less explicit. First of all, $\mathbb{D}_{K/F}$ is a pro-torus defined by its module of characters $X_{K/F}^0$ which is given as the kernel of the following homomorphism of $\mathrm{Gal}(K/F)$ -modules

$$\bigoplus_{v \in V_{K/F}} \mathbb{Z} \cdot v \xrightarrow{\mathrm{deg}} \mathbb{Z}, \quad \sum_v n_v v \mapsto \sum_v n_v \quad (0.2)$$

where $V_{K/F}$ is the set of places of K over F .

In [35], Tate constructed a class $\alpha_{K/F}$ in $H^2(\mathrm{Gal}(K/F), \mathbb{D}_{K/F}(K))$ such that the cup product

$$H^0(\mathrm{Gal}(K/F), X_{K/F}^0) \xrightarrow{\cup \alpha_{K/F}} \mathrm{Br}(K/F) \quad (0.3)$$

is an isomorphism. Kottwitz used the canonical classes $\alpha_{K/F}$, or rather the extensions

$$1 \rightarrow \mathbb{D}_{K/F}(K) \rightarrow \mathrm{Kt}_{K/F} \rightarrow \mathrm{Gal}(K/F) \rightarrow 1 \quad (0.4)$$

corresponding to these classes, to define a functor $G \mapsto B(F, G)$ from linear algebraic groups to pointed sets. More precisely, he defined certain sets of algebraic 1-cocycles of $\mathrm{Kt}_{K/F}$ with values in $G(K)$, and passed to the limit over the set of finite Galois extensions K/F . Using the foundational work of Giraud on non-abelian cohomology [12] and the paper of Breen [5], we interpret both $\alpha_{K/F}$ and the extensions $\mathrm{Kt}_{K/F}$ in terms of gerbes. Then we apply Tannakian duality to obtain Tannakian categories $\mathrm{Rep}(\mathrm{Kt}_F)$ which are of particular interest.

The main question is whether one can describe $\mathrm{Rep}(\mathrm{Kt}_F)$ explicitly in terms of linear algebra, especially in the case when F is a number field.

In his work [20], Kottwitz uses purely cohomological constructions in order to obtain $B(F, G)$, but it turns out that once Tannakian duality is applied to this setting, one obtains a slightly more geometric interpretation of this functor. Namely, when F is a non-archimedean local field, $B(F, G)$ is the pointed set of isomorphism classes of isocrystals with G -structure, which is the same as isomorphism classes of tensor functors from $\mathrm{Rep}(G)$ to the category of isocrystals (see [19]).

The significance of $\mathrm{Rep}(\mathrm{Kt}_{\mathbb{Q}})$ was emphasized in the ICM talk of Scholze (see Section 9 in [31]), where he noticed the analogy between $\mathrm{Rep}(\mathrm{Kt}_{\mathbb{Q}})$ and the category of isoshtukas over curves over finite fields. The latter was introduced by Drinfeld in his work on the Langlands correspondence in the function field case (see [10]). Scholze also stated the following conjecture.

Conjecture 1 (Scholze [31]). *There is a Weil cohomology theory $H_{\text{Kt}_{\mathbb{Q}}}^i(X)$ for varieties X over $\overline{\mathbb{F}}_p$ taking values in $\text{Rep}(\text{Kt}_{\mathbb{Q}})$. Under the functor $\text{Rep}(\text{Kt}_{\mathbb{Q}}) \rightarrow \text{Rep}(\text{Kt}_{\mathbb{Q}_p})$, it maps to crystalline cohomology, and under the functor $\text{Rep}(\text{Kt}_{\mathbb{Q}}) \rightarrow \text{Rep}(\text{Kt}_{\mathbb{Q}_l})$ for $l \neq p$, it maps to étale cohomology. Under the restriction $\text{Rep}(\text{Kt}_{\mathbb{Q}}) \rightarrow \text{Rep}(\text{Kt}_{\mathbb{R}})$, it gives a Weil cohomology theory $H_{\text{Kt}_{\mathbb{R}}}^i(X)$ with values in $\text{Kt}_{\mathbb{R}}$.*

In the first sections, we recall main definitions related to gerbes and present several equivalent approaches to gerbes and their categories of representations. We also prove a general result on gerbes with a smooth band that extends several well-known comparison isomorphisms into the setting of non-abelian H^2 , namely we show that $H_{\text{sm}}^2(F, G) = H_{\text{fppf}}^2(F, G)$, where G is a smooth algebraic group.

In Section 5, we prove that $\text{Rep}(\text{Kt}_{\mathbb{Q}})$ has a fiber functor in \mathbb{Q}^{cycl} . It is however unclear at the moment how one can describe this category explicitly, although there is a hope to give a linear algebraic description of $\text{Rep}(\text{Kt}_{K/\mathbb{Q}})$ for finite cyclic extensions K/\mathbb{Q} .

In Section 7, we describe $\text{Rep}(\text{Kt}_F)$ in the non-archimedean local cases, re-establishing the equivalence between $\text{Rep}(\text{Kt}_F)$ and the category of isocrystals in a very explicit fashion compared to a somewhat more abstract comparison that existed in the literature. In the subsequent section, we show that $\text{Rep}(\text{Kt}_{\mathbb{R}})$ is equivalent to the category of \mathbb{Z} -graded finite dimensional vector spaces V over \mathbb{C} with a semi-linear graded isomorphism $\rho : V \rightarrow V$, such that ρ^2 acts by $(-1)^m$ on the graded component of degree m .

One of the main results of this work is Theorem 8. We show that if F is a global function field, then $\text{Rep}(\text{Kt}_F)$ is equivalent to the category of Drinfeld φ -spaces (or isoshtukas). The latter is defined as a category of finite-dimensional $\mathbb{F}_q(T) \otimes_{\mathbb{F}_q} \overline{\mathbb{F}}_q$ -vector spaces with a Frob-linear automorphism. We use Drinfeld's classification of φ -spaces to show that this category is banded by the correct pro-torus. However, it is quite tricky to recover the class in H^2 that corresponds to the category of isoshtukas. We use ideas of Milne from [25] to reduce everything to local computations, namely, we prove a 'local-global principle' for $H^2(F_{\text{fpqc}}, \varprojlim \mathbb{D}_{K/F})$ (see Section 9.11) and make use of the section on localization of $B(F, G)$ from the Kottwitz paper [20].

As a consequence, we also prove that in the function field case $B(F, G)$ is isomorphic to the set of isomorphism classes of isoshtukas with G -structure (see Theorem 9).

As it is often the case for seemingly analogous results for function fields and number fields, the study of the Kottwitz gerbes and their representations over number fields is admittedly more complicated. One can study $\text{Rep}(\text{Kt}_{\mathbb{Q}})$ via its subcategories given by the categories of representations of algebraic quotients of the Kottwitz gerbe $\text{Kt}_{\mathbb{Q}}$. Recall that a gerbe is called algebraic if it is locally equivalent to a stack of torsors for an algebraic group.

In Section 10, we prove the following refinement of the results on the existence of fibre functors. Let $\text{Kt}'_{\mathbb{Q}}$ be an algebraic quotient $\text{Kt}_{K,S}$ of $\text{Kt}_{\mathbb{Q}}$, where K/\mathbb{Q} is a finite Galois extension and S is a finite set of places of K satisfying Kalaetha-Kottwitz-Tate conditions (see Section 3.2). Then there exists infinitely many rational primes q , such that $\text{Rep}(\text{Kt}'_{\mathbb{Q}})$ has a fibre functor in finite-dimensional vector spaces over the q th cyclotomic field $\mathbb{Q}(\zeta_q)$.

The main advantage here is that we are reduced to finite cyclic Galois extensions that are ramified at only one finite prime, namely at q . This makes the descent data on $\text{Rep}(\text{Kt}'_{\mathbb{Q}})$ more concrete and easier to handle, as compared to the situation of Theorem 2. However, it raises several questions about explicit computation of cohomology of certain non-split tori over $\mathbb{Q}(\zeta_q)$.

Finally, in Section 13, we show that Scholze's conjecture follows from the Tate's conjecture over finite fields. The key ingredient is that assuming the Tate's conjecture, we have an explicit description of $\text{Mot}(\mathbb{F})_{\text{num}}$ obtained Langlands and Rapoport in [21] and in subsequent works of Milne [26], [25].

CONTENTS

Introduction	1
1. Gerbes: from abstract definitions to group extensions	3
2. Gerbes with a smooth band and a comparison isomorphism for H^2	6
3. Definition of Kottwitz gerbes	10
4. The category of representations of the Kottwitz gerbes	13
5. Fibre functors	15
6. Representations of the Kottwitz gerbes with G -structure	18
7. Representations over non-archimedean local fields	20
8. Archimedean cases	23
9. Local-global principle	23
10. φ -spaces	26
11. Comparison over function fields	29
12. More on fiber functors	33
13. Motives over \mathbb{F} and $\text{Rep}(\text{Kt}_{\mathbb{Q}})$	37
References	40

1. GERBES: FROM ABSTRACT DEFINITIONS TO GROUP EXTENSIONS

A gerbe \mathcal{E} over a site S is defined as a stack fibred in groupoids that satisfies the following extra conditions:

- There exists a cover $U \in \text{Ob}(S)$ such that the fibre groupoid \mathcal{E}_U is not empty.
- Any two objects $x, y \in \mathcal{E}_U$ are locally isomorphic, i.e. there exists a cover $U' \xrightarrow{p} U$ such that $p^*x \simeq p^*y$.

In this work we consider étale, fppf and fpqc sites over a field F . Gerbes form a 2-subcategory of the 2-category of stacks.

We refer the reader to [12] for generalities on non-abelian cohomology and gerbes.

1.1. Local description of gerbes. For the sake of completeness we recall the results of Breen on étale gerbes (see Section 4 in [5]). Firstly, we sketch the main steps in describing a gerbe via local data consisting of a concrete bitorsor satisfying several compatibility conditions, we proceed further by describing 1- and 2-morphisms of such gerbes via local data.

Let \mathcal{E} be a gerbe over $F_{\text{ét}}$, assume that \mathcal{E} is split by a finite Galois extension K/F . Choosing a cartesian section x of \mathcal{E} over $U = \text{Spec}(K)$, we obtain an equivalence

$$\Phi : \mathcal{E} \times U \rightarrow \text{TORS}(U, G) \quad (1.1)$$

that sends an object y of the fiber groupoid \mathcal{E}_V over $p : V \rightarrow U$ to the torsor $\text{Isom}(p^*x, y)$. In this situation, we say that $G = \text{Aut}(x)$ is the band of \mathcal{E} , or that \mathcal{E} is banded by G . When G is a (smooth) algebraic group scheme, \mathcal{E} is called a (smooth) algebraic gerbe.

After pulling back to $U \times U$ the equivalence (1.1), we get an equivalence of stacks over U^2

$$\varphi : p_1^*\Phi \circ (p_2^*\Phi)^{-1} : \text{TORS}(U^2, p_1^*G) \rightarrow \text{TORS}(U^2, p_2^*G), \quad (1.2)$$

which restricts to the identity on the diagonal.

Using the definition of Φ and the fact that the category of equivalences between $\text{TORS}(U \times U, p_1^*G)$ and $\text{TORS}(U \times U, p_2^*G)$ is equivalent to the category of (p_2^*G, p_1^*G) -bitorsors (see Chap. IV, Prop. 5.2.5 in [12]), we may pass to the (p_2^*G, p_1^*G) -bitorsor

$$E = \text{Isom}(p_1^*x, p_2^*x) \quad (1.3)$$

over U^2 , whose set of sections over $V \xrightarrow{(f,g)} U^2$ is given by the set $\text{Isom}(f^*x, g^*x)$ of arrows in the groupoid \mathcal{E}_V , moreover, E restricts to $G = \text{Aut}(x)$ on the diagonal.

In order to define the descent data on $\text{TORS}(U, G)$, the equivalence (1.2) and therefore the bitorsor E must satisfy several compatibility conditions after pulling back to U^3 and U^4 ; namely, the condition over U^3 translates into an isomorphism of (p_1^*G, p_3^*G) -bitorsors

$$\psi : p_{12}^*E \wedge p_{23}^*E \xrightarrow{\sim} p_{13}^*E \quad (1.4)$$

that is given by composing $f \in \text{Isom}(p_2^*x, p_1^*x)$ with $g \in \text{Isom}(p_3^*x, p_2^*x)$. The condition over U^4 boils down to the equality of the two possible induced maps

$$p_{12}^*E \wedge p_{23}^* \wedge p_{34}^*E \rightarrow p_{14}^*E. \quad (1.5)$$

1.2. Local description of morphisms between gerbes. Given a gerbe \mathcal{F} such that its pull-back to U is equivalent to $\text{TORS}(U, H)$, where H is an affine group, we let φ' and ψ' denote the attached to \mathcal{F} equivalence analogous to (1.2) and the cocycle isomorphism (1.4) respectively. One can describe a morphism $\mathcal{E} \rightarrow \mathcal{F}$ via the local data. Namely, giving a morphism $f : \mathcal{E} \rightarrow \mathcal{F}$ is equivalent to giving a morphism $f_U : \text{TORS}(U, G) \rightarrow \text{TORS}(U, H)$ and an isomorphism

$$\alpha : p_2^*f_U \circ \varphi \xrightarrow{\sim} \varphi' \circ p_1^*f_U \quad (1.6)$$

that is compatible with the cocycle isomorphisms ψ and ψ' over U^3 .

Let F be a (p_1^*H, p_2^*H) -bitorsor attached to \mathcal{F} . To translate everything into the language of bitorsors we note that the category of morphisms described above is equivalent to the category of morphisms of bitorsors $E \rightarrow F$ over U^2 compatible with the cocycle isomorphisms and whose restrictions to the diagonal define an affine morphism $G \rightarrow H$.

The proof is completely formal, but we sketch the main points here. The functor in one direction is defined by noticing that applying the isomorphism (1.6) to p_1^*G considered as a trivial p_1^*G -torsor and noticing that $E = \varphi(p_1^*G)$ is a (p_1^*G, p_2^*G) -bitorsor. Consequently, $\alpha \circ p_1^*f_U$ defines a morphism from E to $\varphi' \circ p_1^*f_U(p_1^*G)$. Without loss of generality we can assume that the latter is given by F , since we can compose $p_1^*f_U$ with twisting by the image of p_1^*G ; thus we obtain the desired morphism of bitorsors. Going in the other direction, we recover a functor between the stacks $\text{TORS}(U, G)$ and $\text{TORS}(U, H)$ from the given morphism $G \rightarrow H$ of restrictions of the bitorsors E and F to the diagonal. Then we can easily recover an isomorphism (1.6) by using the equivalence stated above and the fact that all the morphisms in question can be described explicitly in terms of contracted products.

Note that so far we have not used the assumption that $U = \text{Spec}(K)$, where K/F is a finite Galois extension, which will be crucial in Section 1.4.

1.3. Framed gerbes over the étale site. Let S be the étale site over a field F . We introduce a 2-category of framed gerbes over S . A framed gerbe is a pair (\mathcal{E}, x) , where \mathcal{E} is a gerbe over S and $x \in \mathcal{E}_U$, where $U \in \text{Ob}(S)$ is an étale cover. A 1-morphism between (\mathcal{E}, x) and (\mathcal{F}, y) is a pair (γ, ξ) , where γ is a functor between gerbes $\gamma : \mathcal{E} \rightarrow \mathcal{F}$ and ξ is an isomorphism $\gamma(p^*(x)) \xrightarrow{\sim} p^*(y)$ in \mathcal{F}_U . A 2-morphism between two framed morphisms of gerbes is just a 2-morphism between the corresponding 1-morphisms of gerbes.

Note that an isomorphism between $\gamma(p^*(x))$ and $p^*(y)$ gives rise to a morphism of sheaves

$$\text{Aut}(p^*(x)) \rightarrow \text{Aut}(p^*(y)) \quad (1.7)$$

which is not in general induced by a morphism of affine schemes (see Remark 2.7 in [5]). We say that a morphism of framed gerbes φ respects affine structure if the morphism of sheaves (1.7) is induced by a morphism of schemes.

Additionally, a framed gerbe (\mathcal{E}, x) is called affine, algebraic or smooth, when the sheaf $\text{Aut}(x)$ is representable by an affine, algebraic or smooth group scheme, respectively.

1.4. Galois gerbes. Let K/F be a finite Galois extension, and let Γ_F denote the absolute Galois group $\text{Gal}(F^{\text{sep}}/F)$. Here we introduce a category of K/F -Galois gerbes, which is more explicit than the category of framed gerbes over $F^{\text{ét}}$ defined in Section 1.3.

Firstly, we introduce the objects. We define a K/F -Galois gerbe (E, φ, j) by the following data

- A short exact sequence of groups:

$$1 \rightarrow \mathcal{G} \rightarrow E \xrightarrow{\varphi} \Gamma_F \rightarrow 1 \quad (1.8)$$

- A splitting $j : \Gamma_K \rightarrow E$ of φ over the open normal subgroup Γ_K of Γ_F .

We say that a K/F -Galois gerbe is smooth if the following condition is satisfied

- The étale sheaf G over $\text{Spec}(K)$ defined by the Γ_K -group structure on \mathcal{G} given by conjugation by j , is representable by a smooth affine group scheme over $\text{Spec}(K)$.

Let K' be a finite Galois extension of F , such that $F \subset K \subset K'$. A $(1-)K'$ -morphism between two K/F -Galois gerbes (E, φ, j) and (F, φ', j') is a homomorphism $\gamma : E \rightarrow F$, which is compatible with the splittings j and j' over $\Gamma_{K'}$ and induces identity on Γ_F . Let \mathcal{H} denote the kernel of γ , and let H be the corresponding smooth affine group scheme. We say that γ respects affine structure if its restriction to the kernels is induced by a morphism of schemes $G \rightarrow H$.

We also define morphisms between K' -morphisms of K/F -Galois gerbes. Note that the definition replicates the equivalence relation Kottwitz used to define $B(F, G)$ (see Section 2.4 in [20]). Namely, two K' -morphisms $\gamma, \delta : (E, \varphi, j) \rightarrow (F, \varphi', j')$ are connected by a 2-morphism if there exists an element $h \in H_{K'}$ such that $\gamma(u) = h\delta(u)u(h)^{-1}$ for any section u of E , where $u(h)$ denotes the action of u on $H(F^{\text{sep}})$ via the projection to Γ_F (see also formula (2.8.3) in [5]).

Finally, we define the category of Galois gerbes by taking the colimit of categories of K/F -Galois gerbes over the directed set of finite Galois extensions K/F contained in a fixed separable closure F^{sep} .

1.5. From framed gerbes to Galois gerbes. We claim that the category of framed gerbes is 2-equivalent to the category of Galois gerbes.

Let $x \in \mathcal{E}_U$, with $U = \text{Spec}(K)$ and K/F finite Galois. The passage from a framed étale gerbe (\mathcal{E}, x) to a Galois gerbe (E, φ, j) is somewhat tedious and is explained in detail in Section 4.1 in [5]. It is worth mentioning that E has a very concrete description in terms of semi-linear isomorphisms between pullbacks of x to the F^{sep} . Namely, let $x_{F^{\text{sep}}}$ be a pullback of x along the morphism $\text{Spec}(F^{\text{sep}}) \rightarrow \text{Spec}(K)$. Then the fibre E_σ over $\sigma \in \Gamma_F$ of the Galois gerbe E is given by the sheaf $\text{Isom}(\sigma^*(x_{F^{\text{sep}}}), x_{F^{\text{sep}}})$. In particular, the fibre over $1 \in \Gamma_F$ corresponds to $\mathcal{G} = \text{Aut}(x_{F^{\text{sep}}})$ (see 1.8). Additionally, multiplication in E is translated into the twisted composition map

$$\text{Isom}(\sigma^*(x_{F^{\text{sep}}}), x_{F^{\text{sep}}}) \times \text{Isom}(\tau^*(x_{F^{\text{sep}}}), x_{F^{\text{sep}}}) \rightarrow \text{Isom}((\sigma\tau)^*(x_{F^{\text{sep}}}), x_{F^{\text{sep}}}) \quad (1.9)$$

which sends a pair of sections (u, v) to $u \circ \sigma^*(v)$. Moreover, $j(\sigma)$ for $\sigma \in \Gamma_K$ provides us with the descent data needed to describe x in terms of its pullback $x_{F^{\text{sep}}}$, and also defines a K -structure on \mathcal{G} .

Let K' be a finite Galois extension of K , also denote by $x_{K'}$ and $y_{K'}$ the inverse images of the trivializing objects in $\mathcal{E}_{K'}$ and $\mathcal{F}_{K'}$ respectively. Let (φ, γ) be a morphism of framed gerbes (see Section 1.3) and assume that γ is defined over K' , that is

$$\gamma : \varphi(x'_{K'}) \xrightarrow{\sim} y_{K'} \text{ in } \mathcal{F}_{K'}. \quad (1.10)$$

Note that the pair (φ, γ) defines a $\Gamma_{K'}$ -equivariant homomorphism $\rho_{(\varphi, \gamma)} : E \rightarrow F$ between the corresponding Galois gerbes, which restricts to a $\Gamma_{K'}$ -equivariant homomorphism $\rho_{(\varphi, \gamma)} : G(F^{\text{sep}}) \rightarrow H(F^{\text{sep}})$. More precisely, $\rho_{(\varphi, \gamma)}$ sends a section $u \in \text{Isom}(\sigma^*(x_{F^{\text{sep}}}), x_{F^{\text{sep}}})$ to the section of $\text{Isom}(\sigma^*(y_{F^{\text{sep}}}), y_{F^{\text{sep}}})$ given by the formula $\gamma\varphi(u)(\sigma^*(\gamma))^{-1}$.

Any 2-morphism $f : (\varphi, \gamma) \rightarrow (\varphi', \gamma')$ between a pair of framed morphisms defined over K' is given explicitly by twisted conjugation by a $\Gamma_{K'}$ -equivariant element $h \in \text{Aut}(x_{F^{\text{sep}}})$, that is defines a 2-morphism in the category of Galois gerbes.

This way, we obtain a 2-functor from framed gerbes over $F_{\text{ét}}$ to Galois gerbes. It follows from general arguments of Sections 1.1 and 1.2 and Galois descent that this functor is a 2-equivalence.

1.6. Kottwitz' notion of gerbes. We note that the notion of a K/F -Galois gerbe from Section 1.4 is slightly different from the notion of a Galois gerbe used by Kottwitz in [20]. Given a finite Galois extension K/F and a short exact sequence

$$1 \rightarrow G(K) \rightarrow E \xrightarrow{\varphi} \text{Gal}(K/F) \rightarrow 1 \quad (1.11)$$

where G is an algebraic group over K , we obtain a K/F -Galois gerbe in the sense of the description in Section 1.4 after considering the inflation of (1.11) along the homomorphisms $\Gamma_F \rightarrow \text{Gal}(K/F)$ and $G(K) \rightarrow G(F^{\text{sep}})$, moreover, a partial splitting j as in the righthand side of (1.8) is defined by the homomorphism of Galois groups $\Gamma_K \rightarrow \Gamma_F$. In fact, the two notions are equivalent, the inverse map is obtained by taking the invariants of E under the action of Γ that is given by the the partial splitting j in (1.8) and that restricts to descent data on \mathcal{G} by definition of K/F -Galois gerbe.

In a similar fashion, a morphism between two extensions of the form (1.11) defines a K -morphism between the corresponding K/F -Galois gerbes. In general, not every morphism between smooth algebraic gerbes $\rho : \mathcal{E} \rightarrow \mathcal{H}$ that are neutralized over K can be presented by a K -morphism of the corresponding Galois gerbes, however, the only obstruction is the existence of γ in (1.10). This problem can be addressed as follows.

We see that $\text{Isom}(z, y_{K'})$ for any $z \in \mathcal{F}_{K'}$ is an $\text{Aut}(y_{K'})$ -torsor, which splits over an étale extension of K' , since $\text{Aut}(y_{K'})$ is smooth. Thus, there exists a finite Galois extension L/F such that the inverse images of $y_{K'}$ and z become isomorphic in \mathcal{F}_L . Consequently, the category obtained by passing to the colimit of categories of K/F -gerbes in the sense of Kottwitz is 2-equivalent to the category of Galois gerbes.

Notice that the argument above shows also that the obvious transition functors between categories of K/F -Galois gerbes and L/F -Galois gerbes are not fully faithful in general.

2. GERBES WITH A SMOOTH BAND AND A COMPARISON ISOMORPHISM FOR H^2

In this section, we establish a comparison isomorphism for non-abelian H^2 that would be applied in Section 6. Namely, we prove the following

Theorem 1. *Let F be a field, and let G be a smooth group scheme over F . Then we have an isomorphism of pointed sets*

$$H^2(F_{\text{ét}}, G) = H^2(F_{\text{fppf}}, G) \quad (2.1)$$

classifying G -gerbes over the corresponding sites.

The proof of this theorem will occupy the rest of the section.

2.1. Inverse and direct images of stacks. Let us first recall the definitions of the direct and inverse image 2-functors defined for stacks.

Let S and S' be two sites. Let $f^{-1} : S \rightarrow S'$ be an adjoint functor to a morphism of sites

$$f : S' \rightarrow S. \quad (2.2)$$

Note that we are primarily interested in the case where $S = \text{Spec}(F)_{\text{fpqc}}$ and $S' = \text{Spec}(K)_{\text{fpqc}}$, and K/F is a finite field extension.

Definition 1. For any stack E' over S' its direct image $f_*(E')$ is given by the stack $E' \times_{S'} S$ obtained from E' via base change along f^{-1} . Namely, the fibre of $f_*(E')$ over $U \in \text{Ob}(S)$ is given by the fibre $E'(f^{-1}(U))$.

Definition 2. A pair (E', φ) where E' is a stack over S' and $\varphi : E' \rightarrow f_*(E')$ a morphism of stacks over S is called an inverse image of E if for any stack G' over S' the functor

$$\text{Cart}_{S'}(E', G') \rightarrow \text{Cart}_S(E, f_*(G')) \quad (2.3)$$

is an equivalence of categories.

In particular, the sites S and S' can be considered as stacks in groupoids over themselves, in this case the couple (S', id_S) is an inverse image of S .

Given a stack E over S and a morphism of sites $S' \rightarrow S$ one can construct a pair $(f^*(E), \varphi)$ passing to the underlying fibred category and then stackifying the inverse image of E as a fibred category, note that $\varphi : F \rightarrow f_* f^* F$ is a natural functor defined by 2-adjunction of direct image and inverse image functors of underlying fibred categories and the stackification.

2.2. Inverse and direct images of gerbes. If E is a gerbe over S then its inverse image f^*E is a gerbe over S' (see III. Corollaire 2.1.5.6 in [12]), thus gerbes are well-behaved under f^* , however, given a gerbe E' over S' , its direct image $f_*(E')$ is not a gerbe in general.

Note that Giraud doesn't assume that stacks are fibered in groupoids, however for a given stack E one can construct a stack $E^{\text{cart}} \subseteq E$, that has the same set of objects as E and the morphisms are morphisms in E that are cartesian. In addition, any functor $E \rightarrow E'$ induces a functor $E^{\text{cart}} \rightarrow (E')^{\text{cart}}$.

Let K/F be a finite field extension. From now on, we assume that E is an algebraic gerbe over S banded by an algebraic group G , where $S = F_{\text{ppf}}$, such that its inverse image $(E', \varphi : E' \rightarrow f_* E')$ along $f : S' \rightarrow S$, where $S' = K_{\text{ppf}}$, has a section, that is there exists a cartesian functor $S' \rightarrow E'$. Thus E' is equivalent to a neutral gerbe $\text{TORS}(S', G')$, where $G' = f^*(G)$ is the inverse image of the sheaf of groups associated to G .

We would like to deduce some properties of $f_*(\text{TORS}(S', G'))$, to do this we apply theory developed by Giraud in V.3 of [12]. For a sheaf of groups G' over S' he defines a sheaf $R^1 f_*(G')$ on S that is associated to a presheaf

$$V \in \text{Ob}(S) \rightsquigarrow H^1(f^{-1}(V), G'). \quad (2.4)$$

There is a canonical isomorphism of sheaves between $R^1 f_*(G')$ and the sheaf of maximal subgerbes of $f_*(E')$ that is denoted by $\text{Ger}(E')$. This implies that $f_* E'$ admits a fully faithful functor $H \rightarrow f_*(E')$ where H is a gerbe over S , if and only if $R^1 f_*(G')$ has a section.

There is an exact sequence of pointed sets (see V. Proposition 3.1.3 in [12])

$$0 \longrightarrow H^1(S, f_*(G')) \longrightarrow H^1(S', G') \xrightarrow{\gamma} H^0(S, R^1 f_*(G')). \quad (2.5)$$

Note that for any finite extension of fields K/F and G' smooth over K , the Shapiro-Oesterlé lemma gives an isomorphism

$$H^1(S, \text{Res}_{K/F}(G')) \simeq H^1(S', G') \quad (2.6)$$

where $\text{Res}_{K/F}(G') = f_*(G')$.

In a way, one can also extend the exact sequence (2.5). Recall that for any G' -torsor P over S' its *adjoint group* $\text{ad}(P)$ is defined as a group of G' -automorphisms of P , and there is an isomorphism of sheaves

$$R^1 f_*(G') \simeq R^1 f_*(\text{ad}(P)). \quad (2.7)$$

Let $c \in H^0(S, R^1 f_*(G'))$. We define $D(c)$ as a fibred category over S with a fiber over $U \in \text{Ob}(S)$ given by the category of G' -torsors over $f^{-1}(U)$ such that the class $p \in H^1(f^{-1}(U), G')$ is obtained by restriction of c via the natural map $H^1(f^{-1}(U), G') \rightarrow R^1 f_*(G')(U)$.

- $D(c)$ is a gerbe over S . It is neutral if and only if c belongs to the image of $H^1(S', G') \rightarrow R^1 f_*(G')$ (the map from (2.5)).
- If c is an image of a class $p \in H^1(S', G')$ that is represented by a torsor P , there is an equivalence of gerbes

$$D(c) \simeq \text{TORS}(S, f_*(\text{ad}(P))) \quad (2.8)$$

- There is a natural morphism $D(c) \subset f_*(\text{TORS}(S', G'))$ and its adjoint $f^*D(c) \rightarrow \text{TORS}(S', G')$.

Combining this with the Shapiro-Oesterlé lemma (2.6) we see that $D(c)$ is a neutral gerbe over S if and only if c is a marked point in $R^1 f_*(G')$, since $\gamma(H^1(S', G')) = 0$ in (2.5).

2.3. Vanishing of $R^1 f_*(G')$ for smooth G' and finite f . Our goal at this point is to show that the natural functor $E \rightarrow f_* f^*(E)$ factors through a neutral subgerbe of $f_* f^*(E)$, which would be given by $\text{TORS}(S, \text{Res}_{K/F} G')$ (see (2.8)). The proof follows Exp. VIII, par. 5 in [15] where the derived direct images of abelian sheaves in étale topology are treated.

We claim that $R^1 f_*(G') = \{*\}$ as an fppf-sheaf in the case of a smooth group G' . Recall that the fppf-site over a scheme has enough points (see [34, Tag 06VW]), that is there exists a conservative family of points $\{p_i\}_{i \in I}$ such that a morphism of sheaves is an isomorphism whenever it holds fiberwise for every p_i . In our case the family of S_{fppf} -local S -schemes defines such a conservative family of fibre functors (see Theorem 2.3 in [11]). Since fppf topology is finer than étale topology we see that any S_{fppf} -local scheme $\text{Spec}(R) \rightarrow S$ is in particular strictly henselian; moreover, all its residue fields are algebraically closed.

Thus we are reduced to the computation of fibres of $R^1 f_*(G')$ at points given by fppf-local schemes \bar{S} . We let \bar{S}' denote the fibre product $S' \times_S \bar{S}$ and let \bar{G}' denote the pullback of G' to \bar{S}' . Then we have an isomorphism of sheaves of pointed sets

$$R^1 f_*(G')_{\bar{S}} \simeq H_{\text{fppf}}^1(\bar{S}', \bar{G}') \simeq H_{\text{ét}}^1(\bar{S}', \bar{G}') \quad (2.9)$$

where the second isomorphism follows from the smoothness of \bar{G}' over \bar{S}' . Applying the finiteness of $S' \rightarrow S$ we see that $\bar{S}' \rightarrow \bar{S}$ is finite, thus \bar{S}' is a finite product of strictly henselian schemes, and over such schemes any torsor under a smooth group is trivial. Therefore

$$R^1 f_*(G')_{\bar{S}} = \{*\} \text{ for any } \bar{S} \text{ fppf-local,} \quad (2.10)$$

which proves the claim.

As a consequence, we see that there is a unique maximal fppf-subgerbe $D(c) \subset f_*(\text{TORS}(S', G'))$ and $D(c) \simeq \text{TORS}(S, \text{Res}_{K/F}(G'))$. By maximality of the latter, we obtain a functor

$$\theta : E \rightarrow \text{TORS}(F_{\text{fppf}}, \text{Res}_{K/F}(G')) \quad (2.11)$$

over F_{fppf} that is given by sending an object $P \in \text{Ob}(E_U)$ to $f_* f^* P$ which is an object of $\text{TORS}(F_{\text{fppf}}, \text{Res}_{K/F}(G'))_U$.

2.4. Comparison isomorphism. Any gerbe over F_{fpqc} banded by an algebraic group L can be neutralized by a finite extension K/F (see section Cohomology in the Appendix of [6]), here we assume for simplicity that L is a band that comes from a sheaf of groups. It is a consequence of the comparison isomorphism for $H^2(F_{\text{fpqc}}, L)$ and $H^2(F_{\text{fppf}}, L)$ and the fact that $H^2(F_{\text{fppf}}, L)$ is a pseudotorsor under $H^2(F_{\text{fppf}}, C(L))$, where $C(L)$ is the centre of L , and the latter is the usual fppf-cohomology of $C(L)$.

We claim that any algebraic gerbe with a smooth band can be neutralized by a finite Galois extension of F .

Note the discussion above reduces this statement to the case $\text{char}(F) > 0$. Moreover, we may assume that K/F is purely inseparable.

Let E be a gerbe banded by L , and let $f : S' = K_{\text{fppf}} \rightarrow S = F_{\text{fppf}}$ be a morphism of sites. Assume that $E' = f^*(E)$ is a neutral gerbe, i.e. $\text{Cart}_{S'}(S', f^*(E))$ is not empty. Thus we are in the situation of the previous section.

Consider the fibre product

$$\begin{array}{ccc} E \times_{E_*} S & \longrightarrow & S \\ \downarrow & & \downarrow \\ E & \xrightarrow{\theta} & E_* = \text{TORS}(S, \text{Res}_{K/F}(G')) \end{array}$$

We note that the vertical arrow provides us with a morphism $E \times_{E_*} F \rightarrow E$ and we claim that $E \times_{E_*} F$ is representable by a smooth scheme over F . To show this we observe that the bottom arrow θ is representable by a smooth scheme.

Indeed, one can check this after pulling back both gerbes to K , and thus reducing to the case of a morphism between neutral gerbes

$$\theta_K : \text{TORS}(S', G) \rightarrow \text{TORS}(S', \text{Res}_{K/F}(G')). \quad (2.12)$$

By definition of representability, we need to check that for any point $u : U \rightarrow \text{TORS}(S', \text{Res}_{K/F}(G'))$, where U is a scheme over $\text{Spec}(K)$, the fibre product

$$\begin{array}{ccc} E \times_{E_*} U & \longrightarrow & U \\ \downarrow & & \downarrow u \\ \text{TORS}(S', G') & \xrightarrow{\theta_K} & \text{TORS}(S', f^* \text{Res}_{K/F}(G')) \end{array} \quad (2.13)$$

is representable by a smooth scheme.

The fibre is given by an $f^* \text{Res}_{K/F}(G')$ -torsor Q over U that lies in the image of θ_K . Since $f^* \text{Res}_{K/F}(G')$ is a smooth group scheme, Q becomes trivial over an étale cover of K , therefore, we may assume that Q is a trivial torsor $f^* \text{Res}_{K/F}(G')$. We can describe the functor θ_K using the following remark.

Remark 1. If $u : A \rightarrow B$ is a monomorphism of groups, then there is a natural functor $\text{TORS}(S, A) \rightarrow \text{TORS}(S, B)$ given by $P \mapsto {}^u P = P \wedge^A B$. There is a natural morphism from P to ${}^u P$ that can be composed with with the quotient by A , i.e. we have a morphism

$$P \rightarrow {}^u P \rightarrow {}^u P/A \quad (2.14)$$

which gives us a section $q : e \rightarrow {}^u P/A$ since the morphism above factors through P/A (locally isomorphic to e , the final object of the site we are working on). Moreover, $\text{Tors}(S, A)$ is equivalent to the category of pairs (Q, q) where Q is a B -torsor, and $q \in \text{Hom}(e, Q/A)$.

Note that θ_K is defined by adjunction

$$f^* P \mapsto f^* f_* f^* P \quad (2.15)$$

where P is an object of a fibre of E , and $f^* P$ is the associated G' -torsor. On the other hand, we also have a natural monomorphism of sheaves of sets

$$f^* P \rightarrow f^* f_* f^* P \quad (2.16)$$

that is compatible with the monomorphism of sheaves of groups $G' \rightarrow f^* \text{Res}_{K/F}(G')$. In this case we have an isomorphism of $f^* \text{Res}_{K/F}(G')$ -torsors (see [12] Prop. III.1.4.6(iii))

$$f^* f_* f^* P \simeq f^* P \wedge^G f^* \text{Res}_{K/F}(G'), \quad (2.17)$$

which allows us to apply the remark above. It follows that the fibre product (2.13) is given by the quotient of a smooth group scheme $f^* \text{Res}_{K/F}(G')$ by G' , and therefore, is representable by a smooth scheme.

We obtain a cartesian functor from the fppf site over the smooth scheme $\text{Res}_{K/F}(G')/G)_{\text{fppf}}$ to the gerbe E , furthermore, $\text{Res}_{K/F}(G')/G)_{\text{fppf}}$ has sections étale locally, that gives us an étale section of E .

The previous argument shows that any G -gerbe E over F_{fppf} trivializes over an étale extension K/F . Taking the inverse image of E along $F \rightarrow K$ we get the category of $G \times_F K$ -torsors over F_{fppf} , which is equivalent to the category of $G \times_F K$ -torsors over $F_{\text{ét}}$. Therefore, any two local sections of E are étale-locally isomorphic.

Remark 2. There is a remark in a paper by Conrad [7] saying that any gerbe E banded by an abelian fppf S -group scheme G is actually an Artin stack, thus it would imply that E can be neutralized by an étale extension by noticing that $H_{\text{fppf}}^2(S, G)$ is a pseudo-torsor under $H_{\text{fppf}}^2(S, C(G))$.

3. DEFINITION OF KOTTWITZ GERBES

3.1. Local fields. Let F be a local field and let K/F be a finite Galois extension with the Galois group $\text{Gal}(K/F)$. Then there is an isomorphism between modified cohomology groups

$$H^0(\text{Gal}(K/F), \mathbb{Z}) \xrightarrow{\sim} H^2(\text{Gal}(K/F), K^\times) \quad (3.1)$$

given by the cup product with a class $\alpha_{K/F} \in H^2(\text{Gal}(K/F), \text{Hom}(\mathbb{Z}, K^\times))$. Using that the latter cohomology group also classifies the extensions of $\text{Gal}(K/F)$ by K^\times , we define $\text{Kt}_{K/F}$ as an extension

$$1 \rightarrow K^\times \rightarrow \text{Kt}_{K/F} \rightarrow \text{Gal}(K/F) \rightarrow 1 \quad (3.2)$$

corresponding to the local canonical class $\alpha_{K/F}$. Hilbert's Theorem 90 ensures that $\text{Kt}_{K/F}$ is essentially well-defined, namely, it is defined up to the conjugation by an element of K^\times .

In order to define Kt_F , Kottwitz considered the limit of $\text{Kt}_{K/F}$ over finite Galois extensions K/F , more accurately, consider $F \subset K \subset L$ such that L is finite Galois over F , then the canonical class $\alpha_{K/F}$ can be considered as an element of $H^2(\text{Gal}(L/F), L^\times)$ via inflation, to which one associates a gerbe $\text{Kt}_{K/F}^{\text{inf}}$ (see Section 2.10 in [20]). Moreover, there is a relation

$$\text{inf}(\alpha_{K/F}) = [L : K] \alpha_{L/F}, \quad (3.3)$$

which implies the existence of a homomorphism $\text{Kt}_{L/F} \rightarrow \text{Kt}_{K/F}^{\text{inf}}$ that is given by the map $p_{L/K} : x \mapsto x^{[L:K]}$ when restricted to L^\times .

Therefore, Kt_F corresponds to the element $\alpha_F \in \varprojlim_{K/F \text{ fin. Galois}} H^2(\text{Gal}(K/F), K^\times)$ defined by the limit of canonical classes $\alpha_{K/F}$.

Applying Hilbert's Theorem 90, we see that inflation $H^2(\text{Gal}(K/F), K^\times) \rightarrow H^2(F_{\text{ét}}, \mathbb{G}_m)$ is injective, this way α_F defines a unique class in $\varprojlim H^2(F, \mathbb{G}_m)$, where the transition maps on cohomology are induced by the maps $p_{L/K}$. Finally, again applying Hilbert's Theorem 90 we obtain that α_F can be considered as an element of $H^2(F_{\text{fpqc}}, \widetilde{\mathbb{G}}_m)$, where $\widetilde{\mathbb{G}}_m$ is the pro-torus over F whose module of characters is given by \mathbb{Q} .

Consequently, the construction of Kottwitz defines an equivalence class of gerbes over F_{fpqc} banded by $\widetilde{\mathbb{G}}_m$.

Remark 3. Note that $H^1(F_{\text{fpqc}}, \widetilde{\mathbb{G}}_m) \neq 0$, despite the fact that $H^1(F_{\text{fpqc}}, \mathbb{G}_m) = 0$, namely this cohomology group is given by $\varprojlim^1 F^\times$ with the transition maps given by raising to n -th powers for any natural n . As a rule, such transition maps are not surjective and the corresponding inverse system would fail Mittag-Leffler condition.

3.2. Global fields. Let F be a global field and let K/F be a finite Galois extension. Here we define the global Kottwitz classes and explain that the construction of Kottwitz gives us a well-defined class in $H^2(F_{\text{fpqc}}, \mathbb{D}_F)$, that is an equivalence class of gerbes banded by a pro-torus \mathbb{D}_F defined below (see Section 3.4). Although the situation is quite similar to the local case, there are several differences, for instance, one is forced to prove several vanishing theorems for the first cohomology of various (pro-)tori, that are in general no longer split over F .

Let us first recall Tate's construction from [35]. Let K/F be a finite Galois extension. Let $S \subset V_F$ be a finite subset of the set of places of F , we define S_K as the set of places in K lying over S , and $\dot{S}_K \subset S_K$ denotes a set of lifts for the places in S (that is, over each $v \in S$ there is a unique $w \in \dot{S}_K$). We assume that the pair (S, \dot{S}_K) satisfies the following list of conditions

- (T1) S contains all archimedean places and all places that ramify in K .
- (T2) Every ideal class of K has an ideal with support in S_K .
- (K1) For every $w \in V_K$ there exists $w' \in S_K$ such that $\text{Stab}(w, \Gamma_{K/F}) = \text{Stab}(w', \Gamma_{K/F})$.
- (K2) For every $\sigma \in \Gamma_{K/F}$ there exists $\dot{v} \in \dot{S}_K$ such that $\sigma \dot{v} = \sigma v$.

The conditions (T1) and (T2) were introduced by Tate in [35] and are sufficient for the construction of Tate-Nakayama classes, the conditions (K1) and (K2) appeared in a paper by Kaletha (see Conditions 3.3.1 in [16]) and are sufficient for proving that $H^1(\Gamma, \mathbb{D}_{S,K}(\bar{F})) = 0$ and an injectivity statement, consequences of which we will recall in Lemma 3.1.

3.3. Tate's construction of canonical classes. Here, K/F is a finite Galois extension of global fields. Let (A) be the following exact sequence of $\Gamma_{K/F}$ -modules

$$0 \rightarrow \mathcal{O}_{K,S}^* \xrightarrow{a'} \mathbb{A}_{K,S}^* \xrightarrow{a} C_{K,S} \rightarrow 1, \quad (\text{A})$$

where

- $\mathcal{O}_{K,S}^* \subset K^*$ is the group of S -units.
- $\mathbb{A}_{K,S}^* \subset \mathbb{A}_K^*$ is the group of S -idèles of K .
- $C_{K,S}$ is the group of S -idèle classes of K .

The second short exact sequence considered by Tate is

$$0 \rightarrow \mathbb{Z}[S_K]_0 \xrightarrow{b'} \mathbb{Z}[S_K] \xrightarrow{b} \mathbb{Z} \rightarrow 0, \quad (\text{B})$$

where

- \mathbb{Z} is a trivial $\Gamma_{K/F}$ -module.
- $\mathbb{Z}[S_K]$ is the free abelian group on the finite set of places S_K , the $\Gamma_{K/F}$ -action being induced by the natural $\Gamma_{K/F}$ -action on V_K .
- $b : \sum_{v \in S_K} n_v v \mapsto \sum_{v \in S_K} n_v$.
- $\mathbb{Z}[S_K]_0$ is the kernel of b , and b' the canonical inclusion.

Tate also considers the group $\text{Hom}(B, A)$ consisting of all triples $(f_3, f_2, f_1) \in \text{Hom}(\mathbb{Z}[S_K]_0, \mathcal{O}_{K,S}^*) \times \text{Hom}(\mathbb{Z}[S_K], \mathbb{A}_{K,S}^*) \times \text{Hom}(\mathbb{Z}, C_{K,S})$ such that

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z}[S_K]_0 & \xrightarrow{b'} & \mathbb{Z}[S_K] & \xrightarrow{b} & \mathbb{Z} \longrightarrow 0 \\ & & \downarrow f_3 & & \downarrow f_2 & & \downarrow f_1 \\ 0 & \longrightarrow & \mathcal{O}_{K,S}^* & \xrightarrow{a'} & \mathbb{A}_{K,S}^* & \xrightarrow{a} & C_{K,S} \longrightarrow 0 \end{array}$$

commutes. In order to prove the existence of cohomology classes $\alpha_i^r(K, S)$ (we will usually omit the pair (K, S) from the notation throughout this section), $i = 1, 2, 3$, giving rise to the following commutative diagram

$$\begin{array}{ccccccc} \dots & \longrightarrow & H^r(\Gamma_{K/F}, \mathbb{Z}[S_K]_0) & \xrightarrow{b'} & H^r(\Gamma_{K/F}, \mathbb{Z}[S_K]) & \xrightarrow{b} & H^r(\Gamma_{K/F}, \mathbb{Z}) & \longrightarrow & \dots \\ & & \alpha_3^r \downarrow & & \alpha_2^r \downarrow & & \alpha_1^r \downarrow & & \\ \dots & \longrightarrow & H^{r+2}(\Gamma_{K/F}, \mathcal{O}_{K,S}^*) & \xrightarrow{a'} & H^{r+2}(\Gamma_{K/F}, \mathbb{A}_{K,S}^*) & \xrightarrow{a} & H^{r+2}(\Gamma_{K/F}, C_{K,S}) & \longrightarrow & \dots \end{array}$$

where all the vertical arrows are isomorphisms, one must construct single *canonical* class in $H^2(G, \text{Hom}(B, A))$, satisfying several compatibility conditions under maps u_i sending a class $\theta \in H^2(G, \text{Hom}(A, B))$ to its i th component for $i = 1, 2, 3$.

Moreover, everything boils down to finding α_1^r and α_2^r , since we have the following short exact sequence

$$0 \rightarrow \text{Hom}(B, A) \xrightarrow{(u_1, u_2)} \text{Hom}(\mathbb{Z}, C_{K,S}) \times \text{Hom}(\mathbb{Z}[S_K], \mathbb{A}_{K,S}^*) \xrightarrow{(b,1)-(1,a)} \text{Hom}(\mathbb{Z}[S_K], C_{K,S}) \rightarrow 0 \quad (3.4)$$

and corresponding long exact sequence

$$0 \rightarrow H^2(\text{Hom}(B, A)) \xrightarrow{(u_1, u_2)} H^2(C_{K,S}) \times H^2(\text{Hom}(\mathbb{Z}[S_K], \mathbb{A}_{K,S}^*)) \rightarrow H^2(\text{Hom}(\mathbb{Z}[S_K], C_{K,S})) \rightarrow \dots \quad (3.5)$$

Note that we used the fact that $H^1(\text{Hom}(\mathbb{Z}[S_K], C_{K,S}))$ vanishes, which follows from Shapiro's lemma and global class field theory; moreover this establishes the uniqueness of canonical class in $H^2(\Gamma_{K/F}, \text{Hom}(B, A))$. Tate defines α in several steps.

Firstly, $\alpha_1(S, K) \in H^2(\Gamma_{K/F}, \text{Hom}(\mathbb{Z}, C_{K,S})) \simeq H^2(\Gamma_{K/F}, C_{K,S})$ is given by the canonical class of global class field theory.

The construction of $\alpha_2 = \alpha_2(S, K)$ is a little bit more delicate. It requires a local description of the group $H^2(\Gamma_{K/F}, \text{Hom}(\mathbb{Z}[S_K], \mathbb{A}_{K,S}^*))$, namely, there is an isomorphism

$$j_P \circ \text{res} : H^2(\Gamma_{K/F}, \text{Hom}(\mathbb{Z}[S_K], \mathbb{A}_{K,S}^*)) \simeq \bigoplus_{w \in \dot{S}_K} H^2(\Gamma_{K/K^w}, \mathbb{A}_{K,S}^*). \quad (3.6)$$

Thus, α_2 can be defined by its local components, more precisely, Tate takes $\alpha_2(w)$ to be the image under $i_w : K_w^* \hookrightarrow \mathbb{A}_{S,K}^*$ of the local fundamental class $\alpha(K_w/F_w) \in H^2(\Gamma_{K/K^w}, K_w^*)$, and checks that thus defined pair (α_1, α_2) maps to 0 in $H^2(\text{Hom}(\mathbb{Z}[S_K], C_{K,S}))$, that is that (α_1, α_2) lies in the image of inclusion in (3.4), and its projection to $H^r(\Gamma_{K/F}, \text{Hom}(\mathbb{Z}[S_K]_0, \mathcal{O}_{K,S}))$ defines the canonical class $\alpha_3^r = \alpha_3^r(S, K)$.

Definition 3. The canonical class $\alpha_3^2(S, K) \in H^2(\text{Gal}(K/F), \mathbb{D}_{S,K}(K))$ is called the Kottwitz class corresponding to the triple (F, K, S) where F, K , and S are as above. If $S = V_K$, then we denote the corresponding class by $\alpha_{K/F}$.

3.4. Global pro-torus \mathbb{D}_F . Define $X_F = \varinjlim_K \mathbb{Z}[V_K]_0$, where the colimit is being taken over the directed set of finite Galois extensions K/F contained in a fixed separable closure F^{sep} of F . Then the pro-torus over F having X_F as the module of characters is denoted by \mathbb{D}_F .

3.5. Vanishing theorems. The following lemma is a consequence of Lemmata 3.1.9 and 3.1.10 in [16].

Lemma 3.1. *Assume (T1-2) and (K1-2), then we have the following vanishing results:*

- The inflation map

$$H^i(\Gamma_{K/F}, \text{Hom}(\mathbb{Z}[S_K]_0, \mathcal{O}_{S,K}^\times)) \rightarrow H^i(\Gamma, \text{Hom}(\mathbb{Z}[S_K]_0, \bar{F}^\times)) = H^i(F, \mathbb{D}_{S,K}) \quad (3.7)$$

is injective for $i = 1, 2$.

- Moreover, $H^1(F, \mathbb{D}_{S,K}) = 0$ and $H^2(F_{\text{fpqc}}, \mathbb{D}_F) = \varprojlim_{S,K} H^2(F, \mathbb{D}_{S,K})$.

In order to prove the latter statement, one uses Grothendieck spectral sequence for the composition of derived limit and derived functor of global sections (see more in Section 5.2).

3.6. Kottwitz gerbes on finite levels. Let us consider the canonical class $\alpha_{K/F}$. Then $\alpha_{K/F}$ belongs to $H^2(\text{Gal}(K/F), \mathbb{D}_{K/F}(K))$, where $\mathbb{D}_{K/F}$ is a pro-torus over F with characters $\mathbb{Z}[V_K]_0$, note that $\mathbb{D}_{K/F}$ is split over K . The Kottwitz gerbe for K/F is defined as an extension

$$1 \rightarrow \mathbb{D}_{K/F}(K) \rightarrow \text{Kt}_{K/F} \rightarrow \text{Gal}(K/F) \rightarrow 1 \quad (3.8)$$

corresponding to the class of $\alpha_{K/F}$. It follows from Lemma 3.1 that the extension $\text{Kt}_{K/F}$ is defined up to conjugation by an element of $\mathbb{D}_{K/F}(K)$. Obviously, $\text{Kt}_{K/F}$ defines a K/F -Galois gerbe in the terminology of Breen introduced in Section 1.4.

3.7. An F_{fpqc} -gerbe attached to the Kottwitz class. Passing to the inverse limit in Lemma 3.1, we obtain the injectivity of the map

$$\varprojlim H^2(\Gamma_{K/F}, \text{Hom}(\mathbb{Z}[S_K]_0, \mathcal{O}_{S,K}^*)) \rightarrow H^2(F, \mathbb{D}_F), \quad (3.9)$$

which shows that the Kottwitz class defined as a projective limit of the Tate-Nakayama classes $\alpha_3(S, K)$ gives rise to a well-defined class in $H^2(F_{\text{fpqc}}, \mathbb{D}_F)$, thus can be represented by an fpqc-gerbe which we denote by Kt_F . Note that the gerbe Kt_F itself is only well-defined up to non-unique isomorphism, since $H^1(F_{\text{fpqc}}, \mathbb{D}_F) \neq 0$. However, equivalent gerbes in the class of Kt_F give rise to equivalent Tannakian categories $\text{Rep}(\text{Kt}_F)$.

4. THE CATEGORY OF REPRESENTATIONS OF THE KOTTWITZ GERBES

4.1. General definition. Let S be a scheme and let $\text{QCOH}(S)$ denote the stack of quasicohherent modules over S_{fpqc} , i.e. $\text{QCOH}(S)_U = \text{Qcoh}_R$ for $R = \Gamma(U, \mathcal{O}_U)$, if U is affine. If \mathcal{E} is a gerbe over S_{fpqc} , we define $\text{REP}(\mathcal{E})$ as a stack of cartesian functors

$$\mathcal{E} \rightarrow \text{QCOH}(S), \quad (4.1)$$

we define $\text{Rep}_S(\mathcal{E})$ or simply $\text{Rep}(\mathcal{E})$ as a groupoid $\text{REP}(\mathcal{E})_S$.

If \mathcal{E} is a neutral gerbe $\text{TORS}(G)$ where G is an affine group, we have an equivalence of categories

$$\text{Rep}(\mathcal{E}) \simeq \text{Rep}(G) \quad (4.2)$$

which is given by sending every cartesian functor $\Phi : \mathcal{E}_U \rightarrow \text{QCOH}(S)_U$ to $\Phi(G_d)$, where G_d is a trivial G -torsor. Since $\text{Aut}(G_d) \simeq G$, we see that G acts on the quasicohherent module $\Phi(G_d)$ by functoriality of Φ .

4.2. Explicit definition. Let K/F be a finite Galois extension. We define the category of representations of the Kottwitz gerbe $\text{Kt}_{K/F}$ as a category of pairs (V, ρ) , where V is as above, and ρ is a homomorphism $\text{Kt}_{K/F} \rightarrow \mathcal{G}_V = \text{GL}(V) \rtimes \text{Gal}(K/F)$, such that

- ρ induces the identity on $\text{Gal}(K/F)$.
- the restriction of ρ to the kernels is induced by an algebraic morphism of K -group schemes $\nu : \mathbb{D}_{K/F} \rightarrow \text{GL}(V)$.

Let $F \subset K \subset L$, where L and K are finite Galois extensions. Following Kottwitz one can also define L/F -Galois gerbes $\text{Kt}_{K/F}^{\text{inf}}$ that correspond to inflated canonical classes $\text{inf}(\alpha_{K/F}) \in H^2(\text{Gal}(L/F), \mathbb{D}_{K/F}(L))$. It follows by descent that $\text{Rep}(\text{Kt}_{K/F}) \simeq \text{Rep}(\text{Kt}_{K/F}^{\text{inf}})$. It is shown in

Section 8 in [20] that there exists a homomorphism $\text{Kt}_{L/F} \rightarrow \text{Kt}_{K/F}^{\text{inf}}$, this gives rise to the functor $\text{Rep}(\text{Kt}_{K/F}^{\text{inf}}) \rightarrow \text{Rep}(\text{Kt}_{L/F})$. Consequently we obtain a functor

$$\text{Rep}(\text{Kt}_{K/F}) \rightarrow \text{Rep}(\text{Kt}_{L/F}). \quad (4.3)$$

It follows formally that (4.3) is fully faithful, hence we define

$$\text{Rep}(\text{Kt}_F) = \varinjlim_{K/F \text{ fin. Galois}} \text{Rep}(\text{Kt}_{K/F}). \quad (4.4)$$

Claim 1. *The definitions above define are equivalent.*

4.3. Algebraic 1-cocycles and morphisms of gerbes. Let K be a finite Galois extension of F , let G be a linear algebraic group over F , and let $\mathbb{D}_{K/F}$ be the band of the gerbe $\text{Kt}_{K/F}$. In section 2.3 of [20] Kottwitz defined the set $Z^1(\text{Kt}_{K/F}, G(K))$ as a set of pairs (x, ν) , where x is an abstract 1-cocycle of $\text{Kt}_{K/F}$ with values in $G(K)$, i.e. x is a map $w \mapsto x_w$ satisfying

$$x_{w_1 w_2} = x_{w_1} w_1(x_{w_2}), \quad (4.5)$$

where $\text{Kt}_{K/F}$ acts on $G(K)$ via projection to $\text{Gal}(K/F)$, and $\nu : \mathbb{D}_{K/F}(K) \rightarrow G(K)$ is an algebraic homomorphism defined over K such that $x_d = \nu(d)$ for any $d \in \mathbb{D}_{K/F}$.

Note that $\text{Gal}(K/F)$ acts on the set of homomorphisms $\mathbb{D}_{K/F}(K) \rightarrow G(K)$ via the formula $\sigma(\nu)(d) = \sigma(\nu(\sigma^{-1}(d)))$. By $\text{Int}(x)$ for $x \in G(K)$ we denote the inner automorphism defined by $g \mapsto xgx^{-1}$.

The cocycle condition (4.5) then implies that $\text{Int}(x_w) \circ \sigma(\nu) = \nu$ whenever w maps to σ .

There is an obvious action of $G(K)$ on the set of algebraic 1-cocycles: $g \in G(K)$ transforms (x, ν) into $(w \mapsto gx_w w(g)^{-1}, \text{Int}(g) \circ \nu)$. Then a pointed set $H_{\text{alg}}^1(\text{Kt}_{K/F}, G(K))$ is defined as the quotient of $Z^1(\text{Kt}_{K/F}, G(K))$ by the action of $G(K)$.

We define \mathcal{G} as a neutral gerbe $\text{TORS}(S, G)$. In this case, the category of representations of \mathcal{G} is equivalent to $\text{Rep}_F(G)$.

Moreover, for any finite Galois extension K/F the gerbe \mathcal{G} can be represented by a K/F -Galois gerbe given by a split extension

$$1 \longrightarrow G(K) \longrightarrow \mathcal{G}_{K/F} \longrightarrow \text{Gal}(K/F) \longrightarrow 1 \quad (4.6)$$

that is $\mathcal{G}_{K/F} \simeq G(K) \rtimes \text{Gal}(K/F)$ with the natural action of $\text{Gal}(K/F)$ on the set of K -points of G .

Recall the product formula for $G(K) \rtimes \text{Gal}(K/F)$

$$(g, \sigma) \bullet (h, \tau) = (g\sigma(h), \sigma\tau), \quad (4.7)$$

which also gives the formula for the action of $G(K)$ by conjugation, namely

$$\text{Int}(g, 1) \circ (h, \tau) = (gh\tau(g)^{-1}, \tau). \quad (4.8)$$

We deduce from (4.5) and the definition of $\text{Kt}_{K/F}$ that an algebraic 1-cocycle (x, ν) defines a homomorphism of extensions

$$\begin{array}{ccccccc} 1 & \longrightarrow & \mathbb{D}_{K/F}(K) & \longrightarrow & \text{Kt}_{K/F} & \longrightarrow & \text{Gal}(K/F) \longrightarrow 1 \\ & & \downarrow \nu & & \downarrow x & & \parallel \\ 1 & \longrightarrow & G(K) & \longrightarrow & \mathcal{G}_{K/F} & \longrightarrow & \text{Gal}(K/F) \longrightarrow 1 \end{array} \quad (4.9)$$

Therefore an algebraic 1-cocycle in the sense of Kottwitz defines a K -homomorphism of the corresponding K/F -Galois gerbes, and thus a morphism between $\text{Kt}_{K/F}$ and \mathcal{G} . Additionally, the formula (4.8) shows that if $x, x' \in Z_{\text{alg}}^1(\text{Kt}_{K/F}, G(K))$ become equivalent in $H_{\text{alg}}^1(\text{Kt}_{K/F}, G(K))$,

then the morphisms between gerbes defined by x and x' are connected by a 2-morphism, and thus define the same element in the pointed set $\mathrm{Hom}(\mathrm{Kt}_{K/F}, \mathcal{G}) / \mathrm{Hom}^2(\mathrm{Kt}_{K/F}, \mathcal{G})$.

4.4. Basic example. Consider the gerbe $\mathrm{HOM}_1(\mathrm{Kt}_F, \mathcal{G})$ (see more in Section 6.1) that is fibered in groupoids of the morphisms between $\mathrm{Kt}_F \times U$ and $\mathcal{G} \times U$ banded by the trivial morphism of bands over U , in this situation we have an equivalence of gerbes

$$\mathrm{HOM}_1(\mathrm{Kt}_F, \mathcal{G}) \simeq \mathcal{G}, \quad (4.10)$$

and the latter is equivalent to the stack $\mathrm{TORS}(S, G)$. Passing to the fibre over F , we obtain the groupoid $\mathrm{Tors}(S, G)$ of G -torsors over S . Noticing that under the equivalence (4.10) 2-morphisms between Kt_F and \mathcal{G} banded by the trivial morphism correspond to morphisms between G -torsors, we obtain an isomorphism of pointed sets

$$\mathrm{Hom}_1(\mathrm{Kt}_F, \mathcal{G}) / 2\text{-}\mathrm{Hom}_1(\mathrm{Kt}_F, \mathcal{G}) \simeq H^1(F, G). \quad (4.11)$$

5. FIBRE FUNCTORS

In this section we show that $\mathrm{Rep}(\mathrm{Kt}_F)$ have fibre functors over any extension of F of cohomological dimension ≤ 1 . More precisely, an extension \tilde{F}/F is of cohomological dimension ≤ 1 , if and only if $\mathrm{Br}(L) = 0$ for every algebraic extension of \tilde{F} . It is, in particular, equivalent to the assertion that, for every finite algebraic Galois extension L/\tilde{F} , the $\mathrm{Gal}(L/\tilde{F})$ -module L^* is cohomologically trivial (for Tate cohomology). The main result of this section can be stated as follows.

Theorem 2.

- $\mathrm{Rep}(\mathrm{Kt}_{\mathbb{Q}})$ has a fiber functor in finite-dimensional vector spaces over $\mathbb{Q}^{\mathrm{cycl}}$, where $\mathbb{Q}^{\mathrm{cycl}}$ is the extension of \mathbb{Q} obtained by adjoining all roots of unity.
- If F is the function field of an algebraic curve over \mathbb{F}_q , then $\mathrm{Rep}(\mathrm{Kt}_F)$ has a fiber functor in finite-dimensional vector spaces over $\check{F} = F \otimes_{\mathbb{F}_q} \overline{\mathbb{F}_q}$.

Recall that if F is a function field of a curve over \mathbb{F}_q , then by Tsen's theorem $F \otimes_{\mathbb{F}_q} \overline{\mathbb{F}_q}$ is of dimension ≤ 1 . If $F = \mathbb{Q}$, then any totally imaginary field such that for every ultrametric place the corresponding local degree is ∞ (as a supernatural number) is of dimension ≤ 1 (see Proposition 9 and Corollary, Chap. II, §3 in [32]). In particular, $\mathbb{Q}^{\mathrm{cycl}}$ satisfies this condition.

5.1. Rephrasing the existence of a fiber functor in cohomological terms. Gerbes over F_{fpqc} banded by B (which we assume to be an abelian affine group scheme) are classified by $H^2(F_{\mathrm{fpqc}}, B)$. Thus the existence of a fiber functor of $\mathrm{Rep}(\mathcal{G})$, where \mathcal{G} is such a gerbe, over $\tilde{F} \supseteq F$ is equivalent to the vanishing of the class of \mathcal{G} in $H^2(\tilde{F}_{\mathrm{fpqc}}, B \times \mathrm{Spec}(\tilde{F}))$. In particular, the vanishing of the latter cohomology group implies the existence of a fiber functor.

The problem we face in the case of Kottwitz gerbes for global fields is that the bands are pro-algebraic groups and we must be careful working with the cohomology groups mentioned above, since projective limits do not commute with fpqc cohomology.

In the following sections, we establish exact sequences that relate fpqc cohomology groups of a projective limit with the projective limit of fpqc-cohomology groups of its algebraic quotients; we also compare cohomology of \mathbb{D}_F with that of a simpler pro-torus \mathbb{T}_F , and deduce the vanishing of $H^2(\tilde{F}_{\mathrm{fpqc}}, \mathbb{D}_F \times \mathrm{Spec} \tilde{F})$ from these two results.

5.2. Cohomological generalities. In Section 3 of [3] Bhatt and Scholze studied replete topoi, and established several properties for limits in such topoi. We only need some of those results in a special case of the topos of fpqc sheaves on the category of schemes, that is shown to be replete in Example 3.1.7. of [3]. We let \mathcal{X} denote the mentioned topos.

Proposition 1. *If \mathcal{X} is a replete topos and $F : \mathbf{N}^{\text{op}} \rightarrow \text{Ab}(\mathcal{X})$ is a diagram with $F_{n+1} \rightarrow F_n$ surjective, then $\lim F_n \simeq \text{R}\lim F_n$.*

Also, derived limit commutes with derived functor of global sections, namely

$$\text{R}\lim \text{R}\Gamma(X, F_n) \simeq \text{R}\Gamma(X, \text{R}\lim F_n). \quad (5.1)$$

Suppose that the conditions of Proposition 1 are satisfied, then we obtain

$$\text{R}\Gamma(X, \varprojlim F_n) \simeq \text{R}\varprojlim \text{R}\Gamma(X, F_n). \quad (5.2)$$

In order to compute the right-hand side one applies Grothendieck spectral sequence:

$$\text{R}^p \varprojlim \circ \text{R}^q \Gamma(X, F_n) \Rightarrow \text{R}^{p+q}(\varprojlim \circ \Gamma)(X, F_n), \quad (5.3)$$

that in its turn combined with (5.1) produces short exact sequences

$$0 \rightarrow \text{R}^1 \varprojlim \text{R}^{i-1} \Gamma(X, F_n) \rightarrow \text{R}^i(\Gamma(X, \varprojlim F_n)) \rightarrow \varprojlim \text{R}^i(X, F_n) \rightarrow 0, \quad i \geq 1. \quad (5.4)$$

In the following we will apply the results of this section in the case $X = X_{\text{fpqc}}$, where X is a spectrum of a field and $\{F_n\}$ is a pro-system of abelian sheaves on X_{fpqc} , then the middle term in (5.4) for $i = 2$ is the desired fpqc cohomology group classifying gerbes banded by $\varprojlim_n F_n$.

We will be using freely several comparison isomorphisms due to Grothendieck and Saavedra.

Theorem 3 (Grothendieck, Saavedra Rivano). *Let X be a scheme and $p : X_{\text{fppf}} \rightarrow X_{\text{ét}}$ be the canonical morphism of sites. Let G be a smooth algebraic group on X . Then the canonical morphism*

$$G \rightarrow p_*(G) \quad (5.5)$$

is an isomorphism, and the canonical morphisms

$$H^i(X_{\text{ét}}, G) \rightarrow H^i(X_{\text{fppf}}, p_*(G)) \quad (5.6)$$

are isomorphisms for $i = 0, 1$ (for all $i \geq 0$ if G is abelian, respectively). If, moreover, $X = \text{Spec } k$ where k is a field, then the canonical morphisms

$$H^i(X_{\text{fppf}}, G) \rightarrow H^i(X_{\text{fpqc}}, G) \quad \text{for } i = 1, 2 \quad (5.7)$$

are isomorphisms.

The following statement generalizes Shapiro's lemma and will be extremely useful for us when we will deal with cohomology of $\text{Res}_{L/K} \mathbb{G}_m$ for finite separable field extension L/K .

Theorem 4. *Let $X' \rightarrow X$ be a finite étale map of schemes, and let G' be a commutative finitely presented and relatively affine X' -group scheme. Then the finitely presented and relatively affine X -group scheme $G = \text{Res}_{X'/X}(G')$ defines a sheaf for the étale topology on X , and there is a natural isomorphism*

$$H^i(X, G) \simeq H^i(X', G'). \quad (5.8)$$

5.3. Reduction to a simpler band. In the case of Kottwitz gerbes for global fields, \mathbb{D}_F is defined to be the projective limit of protori

$$\mathbb{D}_F = \varprojlim_{K/F} \mathbb{D}_{K/F} \quad (5.9)$$

taken over the directed set of finite Galois extensions of F . In a similar fashion we let \mathbb{T}_F denote $\varprojlim_{K/F} \mathbb{T}_{K/F}$.

Using duality for groups of multiplicative type, we obtain a short exact sequence of pro-tori

$$0 \rightarrow \varprojlim_K \mathbb{G}_m \rightarrow \varprojlim_K \mathbb{T}_{K/F} \rightarrow \varprojlim_K \mathbb{D}_{K/F} \rightarrow 0, \quad (C)$$

note that $\varprojlim_K^1 \mathbb{G}_m$ vanishes since the transition maps are surjective. Then we extend scalars to \tilde{F} and consider the long exact sequence of fpqc-cohomology

$$\dots \rightarrow H^i(\tilde{F}, \varprojlim_K \mathbb{G}_m) \rightarrow H^i(\tilde{F}, \varprojlim_K \mathbb{T}_{K/F}) \rightarrow H^i(\tilde{F}, \varprojlim_K \mathbb{D}_{K/F}) \rightarrow H^{i+1}(\tilde{F}, \varprojlim_K \mathbb{G}_m) \rightarrow \dots, \quad (5.10)$$

where we omitted the base change for corresponding sheaves of abelian groups.

We apply short exact sequence (5.4) to $H^i(\tilde{F}, \varprojlim_K \mathbb{G}_m)$ with $i = 1, 2, 3$. It follows immediately that $H^1(\tilde{F}, \varprojlim_K \mathbb{G}_m) = \varprojlim^1 H^0(\tilde{F}, \mathbb{G}_m)$ since $\varprojlim H^1(\tilde{F}, \mathbb{G}_m)$ vanishes by Hilbert's Theorem 90; $H^2(\tilde{F}, \varprojlim_K \mathbb{G}_m) = 0$ since $\varprojlim_K H^2(\tilde{F}, \mathbb{G}_m)$ vanishes by assumption on \tilde{F} and we also have $\varprojlim^1 H^1(\tilde{F}, \mathbb{G}_m) = 0$ by Hilbert's 90; similarly, $H^3(\tilde{F}, \varprojlim_K \mathbb{G}_m)$ vanishes purely by cohomological dimension reasons.

It follows from the vanishing results above that long exact sequence (5.10) induces an isomorphism

$$H^2(\tilde{F}, \mathbb{T}_F \times \text{Spec } \tilde{F}) \simeq H^2(\tilde{F}, \mathbb{D}_F \times \text{Spec } \tilde{F}). \quad (5.11)$$

5.4. The pro-torus \mathbb{T}_F . Recall that we denoted by \mathbb{T}_F the inverse limit of the protori $\mathbb{T}_{K/F}$ defined by their character modules $\mathbb{Z}[V_K]$ with transition maps given as dual maps to the following homomorphisms of character modules

$$\mathbb{Z}[V_K] \rightarrow \mathbb{Z}[V_L] : v \in V_K \mapsto \sum_{w|v} [L_w : K_v] w, \quad F \subseteq K \subseteq L. \quad (5.12)$$

Then $X^*(\mathbb{T}_F) = \varinjlim_K \mathbb{Z}[V_K]$, the limit is being taken over the set of finite Galois extensions of F contained in the fixed separable closure \bar{F} . We can rewrite $\mathbb{Z}[V_K]$ as a direct sum of finite dimensional $\text{Gal}(K/F)$ -modules $X_{v,K}$; namely

$$X_{v,K} = \bigoplus_{w \in V_K, w|v} \mathbb{Z}w. \quad (5.13)$$

Let us denote by $\mathbb{T}_{K/F}^v$ groups of multiplicative type dual to $X_{v,K}$, thus

$$\mathbb{T}_{K/F} = \prod_{v \in V_F} \mathbb{T}_{K/F}^v \quad (5.14)$$

and the pro-torus \mathbb{T}_F can be written as a projective limit of finite products of algebraic groups of the form $\mathbb{T}_{K/F}^v$ indexed by a countable directed set I , then one applies the short exact sequence (5.4) to this limit

$$0 \rightarrow \text{R}^1 \varprojlim_{i \in I} H^1(\tilde{F}, \mathbb{T}^i \times \text{Spec}(\tilde{F})) \rightarrow H^2(\tilde{F}, \mathbb{T}_F \times \text{Spec}(\tilde{F})) \rightarrow \varprojlim_{i \in I} H^2(\tilde{F}, \mathbb{T}^i \times \text{Spec}(\tilde{F})) \rightarrow 0 \quad (5.15)$$

Using the fact that cohomology commutes with finite products, we pass to cohomology of $\mathbb{T}_{K/F}^v \times \text{Spec } \tilde{F}$.

5.5. Vanishing of H^1 and H^2 of algebraic quotients. Let w be a place of K lying over $v \in V_F$, and let K^w denote the decomposition field of $w \in V_K$, that is the field of invariants for the decomposition group G_w at the place w , which is the subgroup of $G = \text{Gal}(K/F)$ acting trivially on w .

Consider the group of characters of $\mathbb{T}_{K/F}^v$ given by G -module $X_{v,K}$ in (5.13). The set of places $w|v$ is in bijection with a system of representatives of the cosets in G/G_w , and there is an

isomorphism of G -modules

$$X^*(\mathbb{T}_{K/F}^v) = X_{v,K} \simeq \bigoplus_{\sigma \in G/G_w} \mathbb{Z}\sigma(w), \quad (5.16)$$

where the latter is endowed with the natural G -structure.

It is clear then that $\mathbb{T}_{K/F}^v$ is isomorphic to $\text{Res}_{K^w/F} \mathbb{G}_m$, the Weil restriction for \mathbb{G}_m over K^w (see Lemma 12.61 in [23]). Using the compatibility of Weil restriction with base change (see Section 7.6 in [4]), we obtain

$$\text{Res}_{K^w/F}(\mathbb{G}_m \times \text{Spec}(K^w)) \times \text{Spec}(\tilde{F}) \simeq \text{Res}_{K^w \otimes_F \tilde{F}/\tilde{F}}(\mathbb{G}_m \times \text{Spec } K^w \times \text{Spec}(\tilde{F})) \quad (5.17)$$

with all the fiber products taken over $\text{Spec}(F)$. Note that $\text{Spec}(K^w) \times_{\text{Spec}(F)} \text{Spec}(\tilde{F}) \simeq \coprod \text{Spec}(\tilde{K}_j)$, where the fields \tilde{K}_j are extensions of \tilde{F} defined by the following decomposition

$$K^w \otimes_F \tilde{F} \simeq \prod \tilde{K}_j.$$

It follows then from the universal property defining Weil restriction applied to the right-hand side of (5.17) that the latter is isomorphic to

$$\prod_j \text{Res}_{\tilde{K}_j/\tilde{F}}(\mathbb{G}_m \times_{\text{Spec}(\tilde{F})} \text{Spec}(\tilde{K}_j)). \quad (5.18)$$

Consequently, in order to prove vanishing of $H^i(\tilde{F}, \mathbb{T}_{K/F}^v)$ for $i = 1, 2$ it is enough to show this for every individual factor in (5.18).

By Theorem 4, we have

$$H^i(\tilde{F}_{\text{ét}}, \text{Res}_{\tilde{K}_j/\tilde{F}}(\mathbb{G}_m \times_{\text{Spec}(\tilde{F})} \text{Spec}(\tilde{K}_j))) \simeq H^i(\tilde{K}_{j,\text{ét}}, \mathbb{G}_m). \quad (5.19)$$

We conclude by noticing that $H^1(\tilde{K}_j, \mathbb{G}_m) = 0$ by Hilbert's 90, and $H^2(\tilde{K}_j, \mathbb{G}_m) = 0$ since \tilde{K}_j/\tilde{F} is a finite algebraic extension of a field of cohomological dimension ≤ 1 .

6. REPRESENTATIONS OF THE KOTTWITZ GERBES WITH G -STRUCTURE

Here we compare $B(G)$ and the set of isomorphism classes of tensor functors $\text{Rep}(G) \rightarrow \text{Rep}(\text{Kt}_F)$. We use Tannakian duality to express everything in terms of gerbes. Namely, let \mathcal{G} be a neutral gerbe attached to G , then the category $\text{Hom}_F^{\otimes}(\text{Rep}(G), \text{Rep}(\text{Kt}_F))$ is equivalent to the category $\text{Hom}_{F_{\text{fpqc}}}(\text{Kt}_F, \mathcal{G})$ consisting of cartesian functors between the corresponding gerbes over F_{fpqc} . In order to apply the explicit description of the morphisms between gerbes introduced in Section 1, we have to show that any functor in $\text{Hom}_{F_{\text{fpqc}}}(\text{Kt}_F, \mathcal{G})$ can be obtained by étale descent. In order to reduce this question to general cohomological statements we would invoke gerbes $\text{HOM}_{\nu}(\text{Kt}_F, \mathcal{G})$ defined in the following section.

6.1. Definition and main properties of the gerbe $\text{HOM}_{\nu}(\mathcal{E}, \mathcal{H})$. Here we work in a more abstract setting, and assume that \mathcal{E} and \mathcal{H} be algebraic gerbes over S having L and M as their bands respectively. Here S is the site of affine schemes over F , where F is field, endowed with the fppf topology.

Recall that the category of bands is defined as follows. The objects are the sheaves of groups and a morphism between two bands L and M is given by the global sections of the quotient sheaf $\text{Hom}(L, M)/M$ with M acting by inner automorphisms.

To any gerbe \mathcal{E} one can canonically attach a band L , namely

$$L|_U \xrightarrow{\sim} \text{band}(\text{Aut}(x)|_U), \quad \text{where } x \in \text{Ob}(\mathcal{E}|_U) \quad (6.1)$$

for any cover U in S .

A morphism of gerbes $f : \mathcal{E} \rightarrow \mathcal{H}$ is banded by a morphism of bands $\nu : L \rightarrow M$ if the following diagram commutes

$$\begin{array}{ccc} \text{band}(\text{Aut}(x)|_U) & \xrightarrow{f} & \text{band}(\text{Aut}(f(x))|_U) \\ \uparrow & & \uparrow \\ L|_U & \xrightarrow{\nu} & M|_U \end{array}$$

where the vertical arrows are the isomorphisms from (6.1). Furthermore, any morphism of gerbes is banded by a unique morphism of bands.

Let $\nu : L \rightarrow M$ be a morphism of bands. The category $\text{Hom}_\nu(\mathcal{E}, \mathcal{H})$ is defined as a full subcategory of $\text{Hom}(\mathcal{E}, \mathcal{H})$ whose objects are ν -morphisms of corresponding gerbes. It is also useful to define a fibred category $\text{HOM}_\nu(\mathcal{E}, \mathcal{H})$ whose objects over U are the morphisms of gerbes

$$\mathcal{E}|_U \rightarrow \mathcal{H}|_U \quad (6.2)$$

that are banded by $\nu|_U$.

Let us recall the main properties of $\text{HOM}_\nu(\mathcal{E}, \mathcal{H})$:

- $\text{HOM}_\nu(\mathcal{E}, \mathcal{H})$ is a gerbe over S .
- The band of $\text{HOM}_\nu(\mathcal{E}, \mathcal{H})$ is identified with C_ν , the centralizer of ν .
- If $L = M$ and $u = \text{id}_L$, then $\text{HOM}_\nu(\mathcal{E}, \mathcal{H})$ is banded by the centre of L .
- There exists a morphism $\mathcal{E} \rightarrow \mathcal{H}$ banded by ν if and only if $\text{HOM}_\nu(\mathcal{E}, \mathcal{H})$ is a neutral gerbe.

Recall that the category $\text{Hom}_\nu(\mathcal{E}, \mathcal{H})$ can be recovered from the gerbe $\text{HOM}_\nu(\mathcal{E}, \mathcal{H})$ as follows:

$$\text{Hom}_\nu(\mathcal{E}, \mathcal{H}) \simeq \varprojlim_{U \in \text{Ob}(S)} \text{HOM}_{\nu|_U}(\mathcal{E}|_U, \mathcal{H}|_U) \quad (6.3)$$

where the transition maps are the natural maps

$$\text{HOM}_{\nu|_U}(\mathcal{E}|_U, \mathcal{H}|_U) \rightarrow \text{HOM}_{\nu|_{U'}}(\mathcal{E}|_{U'}, \mathcal{H}|_{U'}). \quad (6.4)$$

associated to a morphism $U' \rightarrow U$ in the site S .

Note that $\text{Hom}_\nu(\mathcal{E}, \mathcal{H})$ is given by the groupoid $\text{HOM}_\nu(\mathcal{E}, \mathcal{H})_F$.

For more details see IV.2 in [12].

6.2. Reduction steps. Recall that every morphism of gerbes is banded by a unique morphism of bands, and that every morphism of bands is fpqc-locally representable by a morphism of sheaves, thus we can rewrite $\text{Hom}_{F_{\text{fpqc}}}(\text{Kt}_F, \mathcal{G})$ as a disjoint union of $\text{Hom}_\nu(\text{Kt}_F, \mathcal{G})$ over the set of representatives of the morphisms of bands over F^{alg}

$$\text{Hom}(\text{Kt}_F, \mathcal{G}) = \bigsqcup_{\nu: \mathbb{D}_F \rightarrow G \text{ over } F^{\text{alg}}} \text{Hom}_\nu(\text{Kt}_F, \mathcal{G}) \quad (6.5)$$

Remark 4. Notice that we are forced to consider morphisms $\mathbb{D}_F \rightarrow G$ over F^{alg} , since $H^1(F_{\text{fpqc}}^{\text{sep}}, \mathbb{D}_F)$ is given by the derived limit $\varprojlim_{K,S}^1 \mathbb{D}_{K,S}(F^{\text{sep}})$ and is not trivial, therefore, a description of the groupoid $\text{HOM}(\text{Kt}_F, \text{TORS}(G))_{F^{\text{sep}}}$ requires considering all the inner forms of \mathbb{D}_F .

However, any $\nu : \mathbb{D}_F \rightarrow G$ factors through an algebraic quotient of the protorus \mathbb{D}_F , that is through some $\mathbb{D}_{K,S}$ where $K = K(\nu)$ is finite Galois, and $S = S(\nu)$ is a finite set of places that can be taken sufficiently large (so that $H^1(F, \mathbb{D}_{K,S}) = 0$). Therefore, $\text{Hom}_\nu(\text{Kt}_F, \mathcal{G}) = \text{Hom}_\nu(\text{Kt}_{K,S}, \mathcal{G})$. Thus, we reduced the computation of $\text{Hom}(\text{Kt}_F, \mathcal{G})$ to the question about morphisms between algebraic gerbes $\text{Kt}_{K,S}$ and \mathcal{G} , although we are still working over F_{fpqc} .

Consider the gerbe $\mathcal{E} = \text{HOM}_\nu(\text{Kt}_{K,S}, \mathcal{G})$. As mentioned in Section 6.1, this gerbe is banded by the centralizer of ν in G , which we denote by C_ν . Since $\text{Kt}_{K,S}$ is banded by the torus $\mathbb{D}_{K,S}$ and G is a smooth group, we see that C_ν is smooth (see Corollaire 2.4 Exp. XI in [14]). Thus we are in the situation of the comparison isomorphism of Proposition 1, which shows that \mathcal{E} is a gerbe over $F_{\text{ét}}$ meaning that there exists a finite separable extension of F over which \mathcal{E} has a section, moreover, any two local sections are étale locally isomorphic. The latter translates into the statement about 2-morphisms between $\text{Kt}_{K,S}$ and \mathcal{G} .

To finish the comparison of $B(F, G)$ and $\text{Hom}(\text{Kt}_F, \mathcal{G})/\text{Hom}^2(\text{Kt}_F, \mathcal{G})$ we invoke Breen's results from Section 1 as well as Section 4.3, where we explained that algebraic 1-cocycles of Kottwitz define morphisms between gerbes and that equivalent cocycles in $Z_{\text{alg}}^1(\text{Kt}_{K/F}, G(K))$ correspond to morphisms between $\text{Kt}_{K/F} \rightarrow \mathcal{G}$ that are connected by a 2-morphism.

7. REPRESENTATIONS OVER NON-ARCHIMEDEAN LOCAL FIELDS

We write $\text{Rep}(\text{Kt}_F)$ to denote the Tannakian category defined in Section 4.2. In this section we prove the following theorem. We let \check{F} denote the maximal unramified extension of F , if F is of characteristic 0, and $\check{F} = F \otimes_{\mathbb{F}_q} \overline{\mathbb{F}_q}$, if $\text{char}(F)$ is a local field of characteristic $p > 0$ with the residue field \mathbb{F}_q . Note that in both cases $\text{Gal}(\check{F}/F)$ is a pro-cyclic group generated by the Frobenius automorphism $\text{Frob}_{\overline{\mathbb{F}_q}}$.

Theorem 5. *Let F be a non-archimedean local field. Then the category of representations of the Kottwitz gerbe $\text{Rep}(\text{Kt}_F)$ is equivalent to the category of pairs (V, ρ) , where V is a finite-dimensional vector spaces over \check{F} , and $\rho : V \xrightarrow{\sim} V$ is a $\text{Frob}_{\overline{\mathbb{F}_q}}$ -linear automorphism.*

7.1. Reduction to the unramified case. Let K and L be two finite extensions of F such that $[K : F] = [L : F] = n$, L is Galois and K is unramified. Then we have two functors

$$\begin{array}{ccc} \text{Rep}(\text{Kt}_{K/F}^{\text{inf}}) & \xrightarrow{\Theta_{K,K,L}} & \text{Rep}(\text{Kt}_{K.L/F}) , \\ & & \uparrow \Theta_{L,K,L} \\ & & \text{Rep}(\text{Kt}_{L/F}^{\text{inf}}) \end{array}$$

and it follows from local class field theory (cf. [27], Theorem (5.2)) that the cohomology groups $H^2(\text{Gal}(L/F), L^*)$ and $H^2(\text{Gal}(K/F), K^*)$ coincide when considered as subgroups of the group $H^2(\text{Gal}(K.L/F), (K.L)^*)$. More accurately, the inclusions

$$H^2(\text{Gal}(L/F), L^*), H^2(\text{Gal}(K/F), K^*) \hookrightarrow H^2(\text{Gal}(K.L/F), (K.L)^*) \quad (7.1)$$

are defined via inflations of the extensions classified by these cohomology groups. Moreover, inflated extensions corresponding to fundamental classes of K and L are equivalent. It follows that in the limit $\text{Rep}(\text{Kt}_F)$ every representation $\rho : \text{Kt}_{L/F} \rightarrow \mathcal{G}_V$ becomes equivalent to a representation of the Kottwitz gerbe of an unramified extension.

Thus it remains to describe $\text{Rep}(\text{Kt}_{K/F})$ where K is an unramified extension.

7.2. Unramified case. Let K/F be a finite unramified extension of degree n . Consider a representation of the Kottwitz gerbe of K/F , i.e. $\rho : \text{Kt}_{K/F} \rightarrow \mathcal{G}_V$, where V is a finite dimensional vector space over K , and \mathcal{G}_V is a K/F -Galois gerbe defined in Section 4.2.

The Galois group of K is naturally isomorphic to the Galois group of corresponding residue field extension, which is cyclic and generated by the Frobenius element σ_K . Consider any set-theoretic section $d^K(\sigma_K) \in \text{Kt}_{K/F}$, after composing ρ and the projection $\mathcal{G}_V \rightarrow \text{Gal}(K/F)$, we see that $\Phi := \rho(d^K(\sigma_K))$ defines a σ_K -linear automorphism of V .

Note that the following diagram commutes

$$\begin{array}{ccccccc}
 1 & \longrightarrow & \mathbb{G}_m(K) & \xrightarrow{i} & \text{Kt}_{K/F} & \xrightarrow{p} & \langle \sigma_{K/F} \rangle \longrightarrow 1 \\
 & & \downarrow \nu & & \downarrow \rho & & \parallel \\
 1 & \longrightarrow & \text{GL}(V) & \xrightarrow{i'} & \mathcal{G}_V & \xrightarrow{p'} & \langle \sigma_{K/F} \rangle \longrightarrow 1,
 \end{array}$$

and the morphism ν gives us a slope decomposition $V = \bigoplus V^m$, where V^m corresponds to the subspace on which $\mathbb{G}_m(K)$ acts through the character $\lambda \mapsto \lambda^m$.

Since $\sigma_K^n = \text{id}$, we see that

$$\Phi^n(v) = \rho(x)(v), \quad (7.2)$$

where $x = d_K^K(\sigma_K)^n \in K^*$.

It follows from the proposition below that $\Phi^n(v)|_{V^m} = \pi_F^m$.

Proposition 2. *Let K/F be a finite unramified extension of local fields of degree n , and $\sigma_K \in \text{Gal}(K/F)$ be a Frobenius automorphism. Then, for any set-theoretic section $d^K : \text{Gal}(K/F) \rightarrow \text{Kt}_{K/F}$, the element $x = d^K(\sigma_K)^n$ lies in F^* and can be identified with π_F , a uniformizer of F .*

Proof. To simplify notation, we let σ denote σ_K throughout this proof; we also set $d(\sigma) = d^K(\sigma)$.

We see that

$$d(\sigma)d(\sigma)^n = d(\sigma)x = d(\sigma)^n d(\sigma) = xd(\sigma), \quad (7.3)$$

that is $x \in F^*$.

Let $d'(\sigma) = zd(\sigma)$ with $z \in K^*$ be another section, then

$$\begin{aligned}
 x' = d'(\sigma)^n &= (zd(\sigma))^n = z\sigma(z)d(\sigma)d(\sigma)zd(\sigma) \dots = \\
 &= z\sigma(z)\sigma^2(z) \dots \sigma^{n-1}(z)d(\sigma)^n = x \prod_{0 \leq i \leq n-1} \sigma^i(z) = xN_{K/F}(z) \quad (7.4)
 \end{aligned}$$

from this we see that x is defined modulo $N_{K/F}(K^*)$.

It follows from the class field theory that the group $F^*/N_{K/F}(K^*)$ is isomorphic to the group $\pi_F^{\mathbb{Z}}/\pi_F^{[K:F]\mathbb{Z}}$; namely, it follows from the fact that $H^q(\text{Gal}(K/F), U_K) = 1$ for every unramified extension K/F and for every q , if $q = 0$ then we obtain an isomorphism $U_F \simeq N_{K/F}(U_K)$, where U_F and U_K denote the groups of units in F^* and K^* resp.

On the other hand, $\text{Kt}_{K/F}$ corresponds to the fundamental class of local class field theory, namely, to a generator of $H^2(\text{Gal}(K/F), K^*)$; thus x can be identified with π_F , the uniformizer of F . \square

7.3. Functor from $\text{Rep}(\text{Kt}_{K/F})$ to $\text{Isoc}(\bar{\mathbb{F}}_p)$. Let \check{F} denote the fraction field of the ring of Witt vectors over $\bar{\mathbb{F}}_p$, and let σ denote the induced Frobenius automorphism of \check{F} . Then after the base change every representation (V, Φ) defines an isocrystal over $\bar{\mathbb{F}}_p$, namely we have a functor

$$\mathcal{E}_K^{\check{F}} : \text{Rep}(\text{Kt}_{K/F}) \rightarrow \text{Isoc}(\bar{\mathbb{F}}_p), \quad (V, \Phi) \mapsto (V \otimes \check{F}, \Phi \otimes \sigma). \quad (7.5)$$

We claim that $\mathcal{E}_K^{\check{F}}$ is fully faithful; its faithfulness follows immediately, and we need only to show that the functor is full.

General theory of isocrystals reduces the question to studying spaces of morphisms between isoclinic isocrystals of the same slope. By Dieudonné-Manin classification, every isoclinic isocrystal over $\bar{\mathbb{F}}_p$ of slope s/r decomposes as a direct sum of simple objects $E_{s/r}$, moreover, given a module defined by relation $\Phi^n - \pi^m$, it is of dimension n as a vector space and it is isomorphic to the sum of $d = (m, n)$ copies of E^{s_0/r_0} , where $r_0 d = n$ and $s_0 d = m$. Our goal is to show that similar fact holds true for representations of the Kottwitz gerbe for K/F .

Consider a representation $\rho : \text{Kt}_{K/F} \rightarrow \mathcal{G}_V$, where V is a finite-dimensional vector space over K . Let V^m and Φ be as defined above, and let v be a non-zero vector in V^m . Then $\Phi^n(v) = \pi^m v$, by $V_{m,n}$ we denote a submodule V_m generated by $\Phi^i(v)$, where $i = 0, \dots, n-1$.

Note that we have the following decomposition

$$\Phi^n - \pi^m = (\Phi^{\frac{n}{d}} - \pi^{\frac{m}{d}})(\Phi^{\frac{n}{d}(d-1)} + \pi^{\frac{m}{d}} \Phi^{\frac{n}{d}(d-2)} + \dots + \pi^{\frac{m}{d}(d-1)}) = (\Phi^{\frac{n}{d}} - \pi^{\frac{m}{d}})\Psi, \quad d = (m, n). \quad (7.6)$$

Proposition 3. *Let $V_{m,n}$ be a module defined as above, then it decomposes as a direct sum of d invariant subspaces of the operator $\Phi^{n/d}/\pi^{m/d}$, such that their images under $\mathcal{E}_K^{\check{F}}$ are simple isocrystals of slope m/n .*

Proof. It is well known that if $d = 1$, then $V_{m,n}$ is a simple object.

Suppose $d > 1$. Then it follows from (7.6) that $\Psi(av)$ is invariant under the action of $\Phi^{n/d}/\pi^{m/d}$ for any $a \in K$. On the other hand, commuting Φ with coefficients, we get the following vector

$$\Psi(av) = \sigma_K^{\frac{n}{d}(d-1)}(a)\Phi^{\frac{n}{d}(d-1)}v + \pi^{\frac{m}{d}}\sigma_K^{\frac{n}{d}(d-2)}(a)\Phi^{\frac{n}{d}(d-2)}v + \dots + \pi^{\frac{m}{d}(d-1)}av. \quad (7.7)$$

We set $r = n/d \in \mathbb{Z}$. Since K is an unramified extension of F of degree n and $n = dr$, K contains a subfield K' which is an unramified extension of F of degree r , and $[K : K'] = d$. Let a_i with $i = 1, \dots, d$ be a basis of K over K' .

Consider the vectors $\Psi(a_i v)$, $i = 1, \dots, d$. It remains to prove that these vectors are linearly independent over K . This is equivalent to showing that the determinant

$$\begin{vmatrix} a_1 & a_2 & \dots & a_d \\ \sigma_{K'}(a_1) & \sigma_{K'}(a_2) & \dots & \sigma_{K'}(a_d) \\ \vdots & \vdots & \ddots & \vdots \\ \sigma_{K'}^{d-1}(a_1) & \sigma_{K'}^{d-1}(a_d) & \dots & \sigma_{K'}^{d-1}(a_d) \end{vmatrix}, \quad \text{where } \sigma_{K'} \text{ generates } \text{Gal}(K'/F), \quad (7.8)$$

is non-zero, which is indeed the case since it is non-zero modulo maximal ideal (its power factors as a product of all linear combinations of \bar{a}_i with coefficients in the residue field of K' , see [9], Example 2.4).

This shows that $V_{m,n}$ is isomorphic to direct sum of subspaces $V_{m,n}^i$ defined as follows

$$V_{m,n}^i = \langle \Psi(a_i v), \Phi(\Psi(a_i v)), \dots, \Phi^{n/d-1}(\Psi(a_i v)) \rangle, \quad i = 1, \dots, d \quad (7.9)$$

and every $V_{m,n}^i$ is isomorphic (after base change) to simple isocrystal $E_{s/r}$ with $s = m/d$, $r = n/d$. \square

Since every representation of $\text{Kt}_{K/F}$ is a direct sum of $V_{m,n}^i$, it is enough to show that natural injective morphism

$$\text{End}_{\text{Rep}(\text{Kt}_{K/F})}(V_{m,n}^i, \Phi) \rightarrow \text{End}_{\text{Isoc}(\bar{\mathbb{F}}_p)}(V_{m,n}^i \otimes \check{F}, \Phi \otimes \sigma) \quad (7.10)$$

is an isomorphism. Note that both sides of (7.10) define central division algebras over F and after choosing cyclic bases can be easily identified with a subalgebra of $r \times r$ matrices with coefficients in an unramified extension of F of degree r , where $r = n/(m, n)$. This follows immediately from the proof of Proposition 3 for the LHS, and it is done, for instance, in [30], Lemme 3.3.2.2 for the RHS. Thus (7.10) is an isomorphism.

7.4. Equivalence. Combining the reduction argument of subsection 7.1 and the description of $\text{Rep}(\text{Kt}_{K/F})$, where K/F is a finite unramified extension, in 7.3, we obtain a fully faithful functor from the category of representations of the Kottwitz gerbe Kt_F to the category of isocrystals over $\bar{\mathbb{F}}_p$. Moreover, this functor is essentially surjective, since for every simple object $E_{s/r}$ in $\text{Isoc}(\bar{\mathbb{F}}_p)$ there exists a representation of Kottwitz gerbe for K/F with K an unramified extension of degree r over F .

8. ARCHIMEDEAN CASES

8.1. **Representations over \mathbb{C} .** Consider the exact sequence of groups

$$1 \rightarrow \mathbb{G}_m(\mathbb{C}) \rightarrow \text{Kt}_{\mathbb{C}} \rightarrow \text{Gal}(\mathbb{C}/\mathbb{C}) \rightarrow 1, \quad (8.1)$$

since the Galois group of \mathbb{C} is trivial, we see that $\text{Kt}_{\mathbb{C}} = \mathbb{G}_m(\mathbb{C})$, thus its category of representations $\text{Rep}(\text{Kt}_{\mathbb{C}})$ is equivalent to the category \mathcal{C} of \mathbb{Z} -graded complex vector spaces (see [6] Example 2.30, [8] A.8.8 for the proof of general statement).

Note that the grading on $V = \bigoplus V^m \in \text{Ob}(\mathcal{C})$ is defined as follows:

$$V^m = \{v \in V \mid \rho(g)v = m(g)v \text{ for every } g \in \mathbb{G}_m\}, \quad (8.2)$$

where $m(g) \in \text{Hom}(\mathbb{G}_m, \mathbb{G}_m) = \mathbb{Z}$ is the character $\lambda \mapsto \lambda^m$.

8.2. **The case of the Kottwitz gerbe over \mathbb{R} .** In this case, we are given an exact sequence

$$1 \rightarrow \mathbb{G}_m(\mathbb{C}) \xrightarrow{i} \text{Kt}_{\mathbb{R}} \xrightarrow{p} \text{Gal}(\mathbb{C}/\mathbb{R}) \rightarrow 1. \quad (8.3)$$

Proposition 4. *The category of representations of real Kottwitz gerbe $\text{Rep}(\text{Kt}_{\mathbb{R}})$ is equivalent to the category \mathcal{F} of finite dimensional \mathbb{Z} -graded complex vector spaces with semi-linear automorphism α , such that α acts on V^m by $(-1)^m$.*

Proof. Firstly, we describe $\text{Kt}_{\mathbb{R}}$ explicitly.

Let $\Gamma = \text{Gal}(\mathbb{C}/\mathbb{R}) = \langle 1, \sigma \rangle$, where σ is the complex conjugation. Since the kernel \mathbb{G}_m of this extension is abelian, there is a well-defined action of Γ on \mathbb{G}_m , namely for any set-theoretic section $\omega : \Gamma \rightarrow \text{Kt}_{\mathbb{R}}$, we have the action $i(g) \mapsto \omega(\sigma)i(g)\omega(\sigma)^{-1} = \sigma(i(g)) = i(\bar{g})$.

Since $p(\omega(\sigma)) = p(\omega(\sigma^2))$, there exists an element $x \in \mathbb{C}^*$ such that $\omega(\sigma)^2 = x$, but x is known to be 2-cocycle of Γ in \mathbb{G}_m , i.e. satisfies the equation $\omega(\sigma)^2\omega(\sigma) = \omega(\sigma)(\omega(\sigma))^2$, from which we deduce that $x = \bar{x}$, i.e. $x \in \mathbb{R}$.

We have computed $Z^2(\Gamma, \mathbb{C}^*)$, in order to compute coboundary we substitute $\omega(\sigma)$ with $\omega' = z\omega(\sigma)$.

On the one hand $(\omega')^2 = x'$, where $x' \in \mathbb{R}$. On the other hand, we have $(\omega')^2 = |z|^2x$. Thus x and x' differ by the positive real number, and we have $B^2(\Gamma, \mathbb{C}^*) = \mathbb{R}_{>0} \subset \mathbb{R} = Z^2(\Gamma, \mathbb{C}^*)$. The corresponding second cohomology group can be identified with the group $\mathbb{R}/\mathbb{R}_{>0} = \{\pm 1\}$, where -1 corresponds to the class of the nontrivial extension of $\text{Gal}(\mathbb{C}/\mathbb{R})$ by \mathbb{C}^\times .

Our goal is to describe the category of representations of $\text{Kt}_{\mathbb{R}}$ in linear algebraic terms.

Given a representation $\rho : \text{Kt}_{\mathbb{R}} \rightarrow \mathcal{G}_V$ (see Section 4.2), it defines a grading on V by restricting ρ to $\mathbb{G}_m(\mathbb{C})$ and noticing that the morphism on kernels is by definition induced by an algebraic morphism $\mathbb{G}_m \rightarrow \text{GL}(V)$.

We let α denote the element of \mathcal{G}_V defined by $\rho(\omega(\sigma))$. Applying the definition (8.2) to $g = -1 = \omega(\sigma)^2$ we obtain that $\alpha^2 : V^m \rightarrow V^m$ acts by $m(-1) = (-1)^m$.

By definition we have $p' \circ \rho = p$, thus α defines a σ -linear automorphism of V .

Thus we defined a functor $\text{Rep}(\text{Kt}_{\mathbb{R}}) \rightarrow \mathcal{F}$, that is actually an equivalence, since we can recover a representation of \mathbb{G}_m from the knowledge of grading and define the image of $w(\sigma)$ to be a given semi-linear map. \square

9. LOCAL-GLOBAL PRINCIPLE

In this section F is a global field, \bar{F} denotes its separable closure, and $\Gamma_F = \text{Gal}(\bar{F}/F)$. We usually omit F from notation, when the base field is clear. For any Galois extension K/F we denote $\text{Gal}(K/F)$ by $\Gamma_{K/F}$.

9.1. Adèlic cohomology. For a finite set of primes S in F , \mathbb{A}^S is the restricted product of F_v for $v \notin S$, and for a finite extension K/F , $\mathbb{A}_K^S = \mathbb{A}^S \otimes_F K$. When S is empty, we omit it from the notation.

For a torus T over F define

$$H^r(\mathbb{A}^S, T) = \varinjlim_K H^r(\Gamma_{K/F}, T(\mathbb{A}_K^S)),$$

where the limit is taken over the finite Galois extensions L/F contained in fixed separable closure of F .

For the proof of the next result we refer to [25].

Proposition 5. *Let T be a torus over F . For all $r \geq 1$, there is a natural isomorphism*

$$H^r(\mathbb{A}^S, T) \simeq \bigoplus_{v \notin S} H^r(F_v, T). \quad (9.1)$$

Moreover, a short exact sequence of tori gives rise to a long exact sequence of adèlic cohomology.

9.2. The principle. Let K/F be a finite Galois extension of global fields, and let $S \subset V_F$ be a finite subset of the set of places of F . We define S_K as the set of all places of K lying over S , and let \dot{S}_K denote a set of lifts of elements of S to places of K .

As usual, there is an exact sequence of Galois modules

$$0 \rightarrow \mathbb{Z}[S_K]_0 \xrightarrow{b'} \mathbb{Z}[S_K] \xrightarrow{b} \mathbb{Z} \rightarrow 0, \quad (9.2)$$

and an exact sequence of algebraic tori dual to it, namely,

$$0 \rightarrow \mathbb{G}_m \xrightarrow{b} \mathbb{T}_{S,K} \xrightarrow{b'} \mathbb{D}_{S,K} \rightarrow 0. \quad (9.3)$$

Moreover, by \mathbb{D}_F we denote the inverse limit of $\mathbb{D}_{S,K}$. Applying exact sequence (5.4), we obtain

$$0 \rightarrow R^1 \varprojlim H^1(F, \mathbb{D}_{S,K}) \rightarrow H^2(F, \mathbb{D}_F) \rightarrow \varprojlim H^2(F, \mathbb{D}_{S,K}) \rightarrow 0. \quad (9.4)$$

Since $\mathbb{D}_{S,K}$ is a smooth affine algebraic group, we have a comparison isomorphism $H^i(F, \mathbb{D}_{S,K}) \simeq H^i(\Gamma, \mathbb{D}_{S,K}(\bar{F}))$.

It was shown in [17], section 3.8, that $R^i \varprojlim H^1(\Gamma, \mathbb{D}_{S,K}(\bar{F})) = 0$ for $i = 0, 1$. Therefore the sequence of cohomology groups above is reduced to the isomorphism

$$H^2(F, \mathbb{D}_F) \xrightarrow{\sim} \varprojlim H^2(F, \mathbb{D}_{S,K}). \quad (9.5)$$

Next we wish to deal with individual terms $H^2(F, \mathbb{D}_{S,K})$. First, note that (9.3) induces the long exact sequence of cohomology

$$\dots \rightarrow H^1(F, \mathbb{T}_{S,K}) \rightarrow H^1(F, \mathbb{D}_{S,K}) \rightarrow H^2(F, \mathbb{G}_m) \rightarrow H^2(F, \mathbb{T}_{S,K}) \rightarrow H^2(F, \mathbb{D}_{S,K}) \rightarrow H^3(F, \mathbb{G}_m) \rightarrow \dots \quad (9.6)$$

We can simplify this sequence using the following lemma.

Lemma 9.1. • $H^1(F, \mathbb{T}_{S,K}) = 0$.

- $H^2(F, \mathbb{T}_{S,K}) = \bigoplus_{v \in S} \text{Br}(K^{\dot{v}})$, where $\dot{v} \in S_K$ is an arbitrary lift of $v \in S$, and $K^{\dot{v}}$ is a decomposition field of \dot{v} .
- $H^3(F, \mathbb{G}_m) = 0$.
- The map $H^2(F, \mathbb{G}_m) \rightarrow H^2(F, \mathbb{T}_{S,K})$ in the sequence above coincides with the product of restrictions $\text{res}_v : \text{Br}(F) \rightarrow \text{Br}(K^{\dot{v}})$.

Proof. The first two statements follow from the étale variant of Shapiro's lemma (see Theorem 4) combined with Hilbert's Theorem 90.

The vanishing of $H^3(F, \mathbb{G}_m)$ follows from comparison of étale and Galois cohomology and Theorem 14 in [1], in which it was shown that

$$\varinjlim_K H^3(\mathrm{Gal}(K/F), K^*) = 0,$$

where the limit is being taken over the set of finite Galois extensions $F \subset K \subset \bar{F}$.

The last statement follows from Proposition 1.6.5 in [27]. \square

It follows from the lemma above that

$$H^2(F, \mathbb{D}_{S,K}) = \mathrm{Coker}\left(\bigoplus_{v \in S} \mathrm{res}_v : \mathrm{Br}(F) \rightarrow \bigoplus_{v \in S} \mathrm{Br}(K^\dot{v})\right). \quad (9.7)$$

On the other hand, applying Albert–Brauer–Hasse–Noether theorem, we obtain the following commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathrm{Br}(F) & \longrightarrow & \bigoplus_{w \in V_F} \mathrm{Br}(F_w) & \longrightarrow & \mathbb{Q}/\mathbb{Z} \longrightarrow 0 \\ & & \downarrow \bigoplus_{v \in S} \mathrm{res}_v & & \downarrow & & \downarrow \bigoplus \deg(K^\dot{v}:F) \\ 0 & \longrightarrow & \bigoplus_{\dot{v} \in \dot{S}_K} \mathrm{Br}(K^\dot{v}) & \longrightarrow & \bigoplus_{\dot{v} \in \dot{S}_K} \bigoplus_{w \in V_{K^\dot{v}}} \mathrm{Br}(K_w^\dot{v}) & \longrightarrow & \bigoplus_{v \in S} \mathbb{Q}/\mathbb{Z} \longrightarrow 0 \end{array} \quad (9.8)$$

We let $B_{S,K}$ denote $\ker(\mathbb{Q}/\mathbb{Z} \rightarrow \bigoplus_{v \in S} \mathbb{Q}/\mathbb{Z})$. It is a finite subgroup of \mathbb{Q}/\mathbb{Z} of order $\mathrm{gcd}\{\deg(K^\dot{v} : F) \mid v \in S\}$.

Remark 5. Note that if K is a cyclic extension of $F = \mathbb{Q}$ and S contains an inert prime v (that exists in this case), then $B_{S,K}$ vanishes, since $\deg(K^v : \mathbb{Q}) = 1$.

Remark 6. The middle column in the diagram (9.8) can be reinterpreted in terms of adèlic cohomology. Namely $H^2(\mathbb{A}, \mathbb{G}_m) = \bigoplus_{v \in V_F} \mathrm{Br}(F_v)$ and analogously, $H^2(\mathbb{A}, \mathbb{T}_{S,K}) = \bigoplus_{\dot{v} \in \dot{S}_K} \bigoplus_{w \in V_{K^\dot{v}}} \mathrm{Br}(K_w^\dot{v})$. Moreover, since $H^3(\mathbb{A}, \mathbb{G}_m) = 0$, we have an isomorphism

$$H^2(\mathbb{A}, \mathbb{D}_{S,K}) \simeq \mathrm{Coker}(H^2(\mathbb{A}, \mathbb{G}_m) \rightarrow H^2(\mathbb{A}, \mathbb{T}_{S,K})). \quad (9.9)$$

Let us assume for a moment that the inverse system $B_{S,K}$ satisfies the Mittag-Leffler condition and $\varprojlim B_{S,K} = 0$. We will specify the transition maps below. Snake lemma applied to the diagram above gives us an obvious exact sequence, part of which is as follows

$$\dots \rightarrow B_{S,K} \xrightarrow{\delta_{S,K}} H^2(F, \mathbb{D}_{S,K}) \xrightarrow{\eta_{S,K}} H^2(\mathbb{A}, \mathbb{D}_{S,K}) \longrightarrow \dots$$

that leads us to the short exact sequence

$$0 \rightarrow \mathrm{Coker} \delta_{S,K} \rightarrow H^2(F, \mathbb{D}_{S,K}) \rightarrow \mathrm{Im} \eta_{S,K} \rightarrow 0, \quad (9.10)$$

to which we apply the inverse limit functor and obtain $\varprojlim H^2(F, \mathbb{D}_{S,K}) \simeq \varprojlim \mathrm{Im} \eta_{S,K}$. On the other hand, $\varprojlim \mathrm{Im} \eta_{S,K} \hookrightarrow \varprojlim H^2(\mathbb{A}, \mathbb{D}_{S,K})$, from which we conclude that

$$H^2(F, \mathbb{D}_F) \simeq \varprojlim H^2(F, \mathbb{D}_{S,K}) \hookrightarrow \varprojlim H^2(\mathbb{A}, \mathbb{D}_{S,K}). \quad (9.11)$$

We refer to (9.11) as the local-global principle for Tannakian categories banded by \mathbb{D}_F . The lemma below shows that the inverse system of abelian groups $B_{S,K}$ satisfies the desired properties.

Let us first specify transition maps. If S' is a finite set of places of F containing S , the $B_{S',K} \rightarrow B_{S,K}$ is induced by the following diagram

$$\begin{array}{ccc} \mathbb{Q}/\mathbb{Z} & \xrightarrow{\bigoplus_{v \in S'} \deg(K^v:F)} & \bigoplus_{v \in S'} \mathbb{Q}/\mathbb{Z} \\ \text{id} \downarrow & & \downarrow \pi \\ \mathbb{Q}/\mathbb{Z} & \xrightarrow{\bigoplus_{v \in S} \deg(K^v:F)} & \bigoplus_{v \in S} \mathbb{Q}/\mathbb{Z} \end{array}$$

where π is an obvious projection. It is clear that $B_{S,K} = B_{S',K}$ if S is large enough, namely, if S contains all the places w corresponding to all possible degrees of decomposition fields K^w/F . Note that it is a finite condition, since the latter correspond to decomposition subgroups in $\Gamma_{K/F}$. Therefore, it is harmless to assume that S satisfies this condition.

Let E be a finite Galois extension of K , in this situation we have the set S_E and a map $\mathbb{Z}[S_K] \rightarrow \mathbb{Z}[S_E]$ defined on the basis as follows $v \in S_K \mapsto \sum_{w|v} \deg(E_w : K_v)w$, composing this with the corresponding degree maps we see that it induces the multiplication by $\deg(E : K)$ on \mathbb{Z} . Note that after passing to tori dual to these Galois modules and computing their cohomology this map corresponds simply to multiplication by $\deg(E : K)$ on $\text{Br}(F)$, that in its turn induces the map $\mathbb{Q}/\mathbb{Z} \xrightarrow{\deg(E:K)} \mathbb{Q}/\mathbb{Z}$, and we have the following commutative diagram

$$\begin{array}{ccc} B_{S,E} & \hookrightarrow & \mathbb{Q}/\mathbb{Z} \\ \downarrow & & \downarrow \cdot \deg(E:K) \\ B_{S,K} & \hookrightarrow & \mathbb{Q}/\mathbb{Z} \end{array}$$

that shows that the transition map $B_{S,E} \rightarrow B_{S,K}$ factors through the multiplication by $\deg(E : K)$.

Lemma 9.2. *Let $\{B_{S,K}\}$ be the inverse system defined above. Then $B_{S,K}$ satisfies the Mittag-Leffler condition and its inverse limit vanishes.*

Proof. From the discussion above it follows that it is sufficient to show that for any pair (S_K, K) there exists a finite Galois extension $E/K/F$ such that $\deg(E : K)$ is divisible by $|B_{S,E}|$. Suppose that $l = |B_{S,K}|$ is co-prime to the characteristic of F . We can always assume that K contains enough roots of unity and can also replace K with its Hilbert class field, then we can set $E = K(\sqrt[l]{y})$ where y is the product of divisors in \dot{S}_K , then every prime from S_K remains non-split in E and $\deg(E : K) = |B_{S,E}| = |B_{S,K}|$.

In the case $|B_{S,K}|$ is divisible by p^n where p is the characteristic of F , we may take sufficiently large unramified extension of K and consider an extension $E = K(y)$, where y is a root of the additive polynomial $x^q - x = a$, where $q = p^n$, and a is the inverted product of divisors in \dot{S}_K . Then $\deg(E : F) = p^n$ and $|B_{S,E}| = |B_{S,K}|$ since every local degree at $w \in S_K$ is equal to p^n (see Theorem 4.1 in [2]). Now the argument of the previous part can be applied to deal with the co-prime to p part of $B_{S,K}$. \square

10. φ -SPACES

10.1. Drinfeld's classification of φ -spaces. We are going to introduce the category of φ -spaces (or isoshtukas), let F be a function field with the field of constants \mathbb{F}_q .

Definition 4. Let B be an algebraic extension of \mathbb{F}_q . A φ -space over B is a finite dimensional vector space V over $F \otimes B$ with an $\text{id} \otimes \text{Frob}_B$ -linear map $\varphi : V \xrightarrow{\sim} V$. A morphism between two φ -spaces (V_1, φ_1) and (V_2, φ_2) is a $F \otimes B$ -linear map $\alpha : V_1 \rightarrow V_2$ such that $\alpha \circ \varphi_1 = \varphi_2 \circ \alpha$.

The φ -spaces over B form an F -linear Tannakian \otimes -category $\varphi\text{-Vect}(B)$. In order to prove that this category is semi-simple, when $B = \overline{\mathbb{F}}$, and to describe the set of simple objects, Drinfeld introduces the notion of a φ -pair.

Definition 5. A φ -pair is a pair (L, a) , where L is a commutative finite-dimensional F -algebra, and where $a \in L^* \otimes \mathbb{Q}$ satisfies the following property: for any proper F -subalgebra L' of L , $a \notin (L')^* \otimes \mathbb{Q}$.

Notice that the latter condition in the definition of a φ -pair ensures that the group of automorphisms of a given pair is trivial, it also implies that L is separable, and the set of isomorphisms of φ -pairs is identified with the set of orbits of $(F^{\text{sep}})^* \otimes \mathbb{Q}$ under the action of $\text{Gal}(F^{\text{sep}}/F)$. On the other hand, it follows from global class field theory that

$$(F^{\text{sep}})^* \otimes \mathbb{Q} \simeq \varinjlim_{E/F} \text{Div}^0(E) \otimes \mathbb{Q}. \quad (10.1)$$

To every φ -space (V, φ) Drinfeld associates a φ -pair $(L_{(V, \varphi)}, a_{(V, \varphi)})$, and notices that if (V, φ) is a simple object in the category of φ -spaces, then $L_{(V, \varphi)}$ is a field; in particular $a_{(V, \varphi)}$ is just an element in $L_{(V, \varphi)}^\times \otimes \mathbb{Q}$.

Let us briefly recall the construction of a φ -pair associated to a (simple) φ -space (V, φ) . Firstly, we find a pair (V', φ') , where V' is a finite-dimensional vector space defined over $F \otimes_{\mathbb{F}_q} \mathbb{F}_{q^n}$ for some natural n , and φ is a $1 \otimes \text{Frob}_{\mathbb{F}_{q^n}}$ -linear endomorphism of V' such that

$$(V, \varphi) \simeq (V', \varphi') \otimes_{\mathbb{F}_{q^n}} \overline{\mathbb{F}}_q. \quad (10.2)$$

Then since φ^n is a linear automorphism of V' , it extends to a linear automorphism of V , namely $(\varphi')^n \otimes 1$, we denote it by Π . At the same time, Π is an endomorphism of (V, φ) as a φ -space, since it obviously commutes with φ . Thus Π^N for $N \geq 1$ generates a commutative finite-dimensional algebra $L_N = F[\Pi^N] \subseteq \text{End}(V, \varphi)$.

Our next step is to consider an algebra

$$L = \bigcap_{N \geq 1} L_N, \quad (10.3)$$

which does not depend on the choice of φ' .

Remark 7. Since all L_N are finite-dimensional, there exists an index N such that $L = L_N$, moreover, since for any $l, m \geq 1$ such that $l|m$ we have an inclusion $L_m \subseteq L_l$, we see that $L_N = L_{N'}$ for any $N' \geq 1$ divisible by N .

To proceed, observe that $\Pi^N \in L^*$ is not a zero divisor, and denote by w_N its image in $L^* \otimes \mathbb{Q}$. Finally, we set

$$a = a_{(V, \varphi)} = (w_N)^{\frac{1}{n^N}}, \quad (10.4)$$

and note that a does not depend on the choices of φ' and N . The pair (L, a) is a φ -pair associated to (V, φ) .

The following theorem is proved in [10]. Note that $\deg_x(a)$ for $a \in \text{Div}^0(L) \otimes \mathbb{Q}$ denotes the product of the degree of the residue field of the place x and the multiplicity of x in the rational divisor a .

Theorem 6. (1) *The category of φ -spaces over $\overline{\mathbb{F}}_q$ is abelian and semi-simple.*

- (2) The map $(V, \varphi) \mapsto (L_{(V, \varphi)}, a_{(V, \varphi)})$ induces a bijection between the set of isomorphism classes of irreducible φ -spaces and the set of isomorphism classes of φ -pairs (L, a) , where L is a field.
- (3) Let (V, φ) be an irreducible φ -space corresponding to a pair (L, a) , then

$$\dim_{F \otimes_{\mathbb{F}_q} \overline{\mathbb{F}_q}}(V) = [L : F] \cdot d(a), \quad (10.5)$$

where $d(a)$ denotes the common denominator of $\deg_{\tilde{x}}(a) \in \mathbb{Q}$. Moreover, $\text{End}(V, \varphi)$ is a central division algebra over L of dimension $d(a)^2$ with local invariants $-\deg_{\tilde{x}}(a) \pmod{\mathbb{Z}}$.

We wish to reinterpret this theorem as a description of a Tannakian category $\varphi\text{-Vect}(\overline{\mathbb{F}_q})$. As a preparatory step we should prove the following statement

Proposition 6. *Let (V_1, φ_1) and (V_2, φ_2) be two isotypic φ -spaces over $\overline{\mathbb{F}_q}$ of types $a_1, a_2 \in (F^{\text{sep}}) \otimes \mathbb{Q}$ respectively, then their tensor product $(V_1 \otimes V_2, \varphi_1 \otimes \varphi_2)$ is an isotypic φ -space of type $a_1 a_2$.*

Proof. Since the category of φ -spaces is semi-simple, we may assume that (V_i, φ_i) are simple objects.

Let $(\tilde{V}, \tilde{\varphi})$ denote the tensor product $(V_1 \otimes V_2, \varphi_1 \otimes \varphi_2)$.

Following the construction, we define (V'_i, φ'_i) , $i = 1, 2$, to be φ -spaces over $\mathbb{F}_{q^{n_i}}$ such that

$$(V_i, \varphi_i) = (V'_i, \varphi'_i) \otimes_{\mathbb{F}_{q^{n_i}}} \overline{\mathbb{F}_q}, i = 1, 2.$$

Note that $\varphi_i^{n_i}$ is a linear bijective map $V_i \rightarrow V_i$. We can find an integer n' such that both spaces (V'_i, φ'_i) ($i = 1, 2$) are defined over $\mathbb{F}_{q^{n'}}$. We denote by (L_i, a_i) corresponding φ -pairs, where $L_i = F[\Pi_i^{N_i}]$ is a separable extension of F for $i = 1, 2$, and $\Pi_i = (\varphi'_i)^{n'} \otimes \text{id}$, we have also

$$a_i = (w_{N_i})^{\frac{1}{n'N_i}} \in L_i^* \otimes \mathbb{Q}. \quad (10.6)$$

Notice that

$$(V_1 \otimes V_2, \varphi_1 \otimes \varphi_2) \simeq (V'_1 \otimes V'_2, \varphi'_1 \otimes \varphi'_2) \otimes_{\mathbb{F}_{q^{n'}}} \overline{\mathbb{F}_q}, \quad (10.7)$$

and that $\Pi = (\varphi'_1 \otimes \varphi'_2)^n \otimes 1$ is a linear bijective endomorphism of $(\tilde{V}, \tilde{\varphi})$, let $L = L_N$ denote the intersection of finite-dimensional commutative algebras $F[\Pi^N]$ with $N \geq 1$.

Then using Remark 7, we deduce that N_1, N_2 , and N can be chosen to be equal, and $\Pi^N = \Pi_1^N \otimes \Pi_2^N$ as an element in $\text{End}(\tilde{V}, \tilde{\varphi}) \simeq \text{End}(V_1, \varphi_1) \otimes_F \text{End}(V_2, \varphi_2)$; furthermore, $a \in L^* \otimes \mathbb{Q}$ can be identified with its image in $(L_1 L_2)^* \otimes \mathbb{Q}$, which is the same as $a_1 a_2$. Thus $(\tilde{V}, \tilde{\varphi})$ is an isotypic φ -space of type $a_1 a_2$. \square

10.2. The band of the category of φ -spaces. In this section we show that the band of the category of φ -spaces can be identified with the global Kottwitz protorus \mathbb{D}_F .

Notice that in order to compute the band of a semisimple Tannakian category \mathcal{C} over F it is enough to do so for the category $\mathcal{C} \otimes \tilde{F}$ for any \tilde{F}/F such that \mathcal{C} has a fibre functor in $\text{Vect}_{\tilde{F}}$, in which case $\mathcal{C} \otimes \tilde{F}$ becomes a neutral Tannakian category over \tilde{F} , and therefore is identified with the category of representations of its band $\text{Rep}_{\tilde{F}}(B)$, where we assume B to be represented by an affine scheme over \tilde{F} . Moreover, if any simple object of $\mathcal{C} \otimes \tilde{F}$ is of rank 1, then B is identified with $D(\Sigma)$, the diagonalizable group dual to the abelian group of isomorphism classes of simple objects with the group structure given by tensor product (see Proposition 2.22 in [26]).

In the case when \mathcal{C} is the category of φ -spaces, we may take $\tilde{F} = F^{\text{sep}}$. Clearly, φ -spaces has a fibre functor over F^{sep} , and therefore φ -spaces becomes a neutral Tannakian category after tensoring with F^{sep} . We claim that every simple object in $\mathcal{C} \otimes F^{\text{sep}}$ is of rank 1, and the set of isomorphism classes of simple objects is given by the abelian group $(F^{\text{sep}})^{\times} \otimes \mathbb{Q}$.

Let (V, φ) be a simple object in \mathcal{C} , and let (L, a) be its φ -pair. Since $\text{End}(V, \varphi)$ is a central division algebra over L , we see that $\text{End}(V \otimes_F F^{\text{sep}}, \varphi \otimes \text{id})$ is a central division algebra over

$$L \otimes_F F^{\text{sep}} = \prod_{j:L \hookrightarrow K} F^{\text{sep}} \quad (10.8)$$

and therefore $(V \otimes_F F^{\text{sep}}, \varphi \otimes \text{id})$ splits into the sum of subobjects given by $(V_j, \varphi_j) = (V \otimes_{L,j} F^{\text{sep}}, \varphi \otimes (\text{id} \circ j))$. Moreover, $\text{End}(V_j, \varphi_j)$ is a central simple algebra over F^{sep} , hence it is isomorphic to $M_{d(a)}(F^{\text{sep}})$, where $d(a) = \deg_L(\text{End}(V, \varphi))$ is as in Theorem 6. On the other hand

$$\dim_L(V) = \dim_{F^{\text{sep}}}(V_j) = d(a),$$

and therefore the centre of $\text{End}(V_j, \varphi_j)$ provides us with idempotents $e_1, \dots, e_{d(a)}$ splitting (V_j, φ_j) into the sum of $d(a)$ simple isotypic objects $(e_i V_j, e_i \varphi_j)$, for $i = 1, \dots, d(a)$, of rank 1 and slope $a \in (F^{\text{sep}})^\times \otimes \mathbb{Q}$.

10.3. Drinfeld spaces and Dieudonné modules. Let u be a place of F , and let F_u denote the completion of F at u .

We have an exact and F -linear functor

$$\text{Loc}_u : (V, \varphi) \mapsto (V_u, \varphi_u) = (V \hat{\otimes}_F F_u, \varphi \hat{\otimes}_F F_u). \quad (10.9)$$

The following theorem is due to Drinfeld (see also Proposition (B.4) in [22]).

Theorem 7. *Fix a place $u \in V_F$. Let (V, φ) be an irreducible φ -space and let (L, a) be the corresponding φ -pair. For every place $w \in L$ dividing u we let*

$$(V_w, \varphi_w) = L_w \otimes_L (V, \varphi). \quad (10.10)$$

The canonical splitting $F_u \otimes_F L \simeq \prod_{w|u} L_w$ induces a splitting

$$(V_u, \varphi_u) = \bigoplus_{w|u} (V_w, \varphi_w) \quad (10.11)$$

of (V_u, φ_u) as a Dieudonné F_u -module. Then for each $w|u$, (V_w, φ_w) is non-canonically isomorphic to

$$(N_{d_w, r_w}, \psi_{d_w, r_w})^{s_w} \quad (10.12)$$

where the integers d_w , r_w , and s_w are uniquely defined from the following relations

$$\begin{cases} d_w, s_w \geq 1 \\ (d_w, r_w) = 1 \\ r_w/d_w = \deg_w(a)/\deg(L_w : F_u) \\ d_w s_w = d(a) \deg(L_w : F_u) \end{cases} \quad (10.13)$$

11. COMPARISON OVER FUNCTION FIELDS

In this section we prove the following theorem.

Theorem 8. *Let F be a function fields of a smooth algebraic curve over a finite field. Then the category of representations of the Kottwitz gerbe $\text{Rep}(\text{Kt}_F)$ is equivalent to the category of Drinfeld isoshutkas (see Section 10.1).*

11.1. **Adèlic Kottwitz class.** The definition of the Kottwitz class as an element of $H^2(F, \mathbb{D}_F)$ is quite inexplicit, but we are able to compute its local components in $\varprojlim H^2(\mathbb{A}, \mathbb{D}_{S,K})$.

It follows from the isomorphism (3.6) and Remark 6 that the natural map

$$H^i(\Gamma_{K/F}, \text{Hom}(\mathbb{Z}[S_K], \mathbb{A}_{S,K}^*)) \rightarrow H^i(\Gamma, \text{Hom}(\mathbb{Z}[S_K], \mathbb{A}_K^*)) \rightarrow H^i(\mathbb{A}, \mathbb{T}_{S,K}) \quad (11.1)$$

is injective. Therefore, the limit of semi-local classes $\alpha_2(S, K) \in H^i(\Gamma_{K/F}, \text{Hom}(\mathbb{Z}[S_K], \mathbb{A}_{S,K}^*))$, defines a unique element in $\varprojlim H^2(\mathbb{A}, \mathbb{T}_{S,K})$. Furthermore, on every finite level we have an equality of Tate-Nakayama classes $b' \alpha_2(S, K) = a' \alpha_3(S, K)$ coming from natural maps

$$\begin{array}{ccc} H^2(\Gamma_{K/F}, \text{Hom}(\mathbb{Z}[S_K], \mathbb{A}_{K,S}^*)) & & \\ & \downarrow b' & \\ H^2(\Gamma_{K/F}, \text{Hom}(\mathbb{Z}[S_K]_0, \mathcal{O}_{K,S}^*)) & \xrightarrow{a'} & H^2(\Gamma_{K/F}, \text{Hom}(\mathbb{Z}[S_K]_0, \mathbb{A}_{K,S}^*)) \end{array} \quad (11.2)$$

11.2. **Inflations of classes defined for finite extensions.** The following diagrams

$$\begin{array}{ccc} H^2(\Gamma_{K/F}, \text{Hom}(\mathbb{Z}[S_K]_0, \mathcal{O}_{K,S}^*)) & \xrightarrow{a'} & H^2(\Gamma_{K/F}, \text{Hom}(\mathbb{Z}[S_K]_0, \mathbb{A}_{K,S}^*)) \\ \downarrow & & \downarrow \\ H^2(F, \mathbb{D}_{S,K}) & \longrightarrow & H^2(\mathbb{A}, \mathbb{D}_{S,K}) \end{array}$$

and

$$\begin{array}{ccc} H^2(\Gamma_{K/F}, \text{Hom}(\mathbb{Z}[S_K], \mathbb{A}_{K,S}^*)) & \xrightarrow{b'} & H^2(\Gamma_{K/F}, \text{Hom}(\mathbb{Z}[S_K]_0, \mathbb{A}_{K,S}^*)) \\ \downarrow & & \downarrow \\ H^2(\mathbb{A}, \mathbb{T}_{S,K}) & \xrightarrow{b'} & H^2(\mathbb{A}, \mathbb{D}_{S,K}) \end{array}$$

commute. This basically follows from the same computation as Proposition 5.

11.3. **Local components of the Kottwitz class.** We note that the maps $H^2(\mathbb{A}, \mathbb{T}_{S,K}) \rightarrow H^2(\mathbb{A}, \mathbb{D}_{S,K})$ are surjective, and give rise to the surjective map between corresponding derived limits, since the kernel of every such projection is a quotient of $H^2(\mathbb{A}, \mathbb{G}_m)$, that is a direct sum of \mathbb{Q}/\mathbb{Z} and $\frac{1}{2}\mathbb{Z}$ and has vanishing $R^1 \varprojlim$, and therefore the cokernel of

$$\varprojlim H^2(\mathbb{A}, \mathbb{T}_{S,K}) \rightarrow \varprojlim H^2(\mathbb{A}, \mathbb{D}_{S,K}) \quad (11.3)$$

is zero.

We have a diagram analogous to (11.2), namely

$$\begin{array}{ccc} \varprojlim H^2(\mathbb{A}, \mathbb{T}_{S,K}) & & \\ & \downarrow b' & \\ H^2(F, \mathbb{D}_F) & \xleftarrow{a'} & \varprojlim H^2(\mathbb{A}, \mathbb{D}_{S,K}) \end{array} \quad (11.4)$$

and two uniquely defined classes $\varprojlim \alpha_2(S, K)$ and $\varprojlim \alpha_3(S, K)$, such that their images coincide in $\varprojlim H^2(\mathbb{A}, \mathbb{D}_{S,K})$.

The images of $\alpha_2(S, K)$ in $H^2(\mathbb{A}, \mathbb{T}_{S,K})$ can be described fairly easily via isomorphism (3.6), more precisely, $\alpha_2(S, K)$ corresponds to the element $(\alpha_{K^\flat/F})_{v \in S}$ where each $\alpha_{K^\flat/F}$ is the image of the local fundamental class in $H^2(\Gamma_{K/K^\flat}, K_v^\times) \simeq \frac{1}{n_v} \mathbb{Z}$ where $n_v = \deg(K : K^\flat) = \deg(K^\flat : F_v)$.

We can say a bit more about the limit of $\alpha_2(S, K)$, since

$$H^2(\mathbb{A}, \mathbb{T}_{S,K}) \simeq \bigoplus_{\dot{v} \in \dot{S}_K} \bigoplus_{w \in V_{K\dot{v}}} \mathrm{Br}(K_w^\dot{v}), \quad (11.5)$$

and we notice that $\mathrm{Br}(K_w^\dot{v}) = \mathrm{Br}(F_v)$ for any place w lying over v . Moreover, the transition maps $\mathbb{T}_{S',E} \rightarrow \mathbb{T}_{S,K}$ are defined locally and for any place $w' \in S_E$ dividing w induce the map

$$\cdot \deg(E_{w'} : K_w) : \mathrm{Br}(E_{w'}^\dot{v}) = \mathrm{Br}(F_v) \rightarrow \mathrm{Br}(K_w^\dot{v}) = \mathrm{Br}(F_v) \quad (11.6)$$

and the v -component of the inverse limit of local canonical classes with so defined transition maps coincides with the cohomology class $\alpha_v \in H^2(F_v, \widehat{\mathbb{G}}_m)$ corresponding to the category of isocrystals. Here $\widehat{\mathbb{G}}_m$ denotes the inverse limit of \mathbb{G}_m with $X^*(\widehat{\mathbb{G}}_m) = \mathbb{Q}$. This implies that $\varprojlim \alpha_2(S, K)$ equals to the image of the product of α_v under the inclusion

$$\prod_{v \in V_F} H^2(F_v, \widehat{\mathbb{G}}_m) \hookrightarrow \varprojlim H^2(\mathbb{A}, \mathbb{T}_{S,K}). \quad (11.7)$$

11.4. $\mathrm{Div}^0(K)$ and $\mathbb{Z}[V_K]_0$. Note that the degree map defined on the abelian group $\mathrm{Div}(K)$ is the following

$$\deg : \mathrm{Div}(K) \rightarrow \mathbb{Z}, \quad \sum_{P \in V_K} n_P P \mapsto \sum_{P \in V_K} \deg(P) n_P, \quad (11.8)$$

where $\deg(P)$ is the residue degree at the place P . Thus one has an isomorphism between $\mathrm{Div}(K) \otimes \mathbb{Q}$ and $\mathbb{Z}[V_K] \otimes \mathbb{Q}$ preserving kernels of the corresponding degree maps, which is given by

$$\mathrm{Div}(K) \otimes \mathbb{Q} \rightarrow \mathbb{Z}[V_K] \otimes \mathbb{Q}, \quad \deg(P)^{-1} P \mapsto w, \quad (11.9)$$

where w and P are the same places of K , and under such an identification $n_w = n_P \deg(P)$.

11.5. Local cohomology classes via duality. There is a sequence of natural maps on cohomology

$$H^2(F, \mathbb{D}_F) \xrightarrow{\sim} \varprojlim H^2(F, \mathbb{D}_{S,K}) \hookrightarrow \varprojlim \bigoplus_{u \in V_F} H^2(F_u, \mathbb{D}_{S,K}) \hookrightarrow \prod_{u \in V_F} \varprojlim H^2(F_u, \mathbb{D}_{S,K}), \quad (11.10)$$

where we omitted for brevity the obvious base change maps. Notice that the inclusion in the middle is given by the local-global principle (9.11).

It follows from Lemma 14.4 in [20] and the fact that $\mathbb{D}_{S,K} \times \mathrm{Spec}(F_u)$ is split over K_w for $w \in V_K$ lying above u , that $\varprojlim^1 H^1(F_u, \mathbb{D}_{S,K} \times \mathrm{Spec}(F_u)) = 0$, and therefore

$$H^2(F_u, \mathbb{D}_F \times \mathrm{Spec}(F_u)) \simeq \varprojlim_{S,K} H^2(F_u, \mathbb{D}_{S,K} \times \mathrm{Spec}(F_u)). \quad (11.11)$$

Taking a product of the localization functors Loc_u over the set of places of F defines a unique class in $\prod_{u \in V_F} H^2(F_u, \mathbb{D}_F)$.

Note that Tate-Nakayama duality over the local field F_u can be rephrased as a classification of Tannakian categories banded by tori. The following proposition generalizes a result of Saavedra Rivano (see VI.Proposition 3.5.3) to the case of Tannakian categories over local fields that are banded by tori that are not necessarily split over the base field.

Proposition 7. *Let F_u be a local field, and let \mathcal{C} be a Tannakian category over F_u banded by an algebraic torus T . Then the class of \mathcal{C} in $H_{\mathrm{ppf}}^2(F_u, T)$ is given by an element of $\mathrm{Hom}(X^*(T)^{\Gamma_u}, \mathrm{Br}(F))$ defined by sending every simple object E_χ , for $\chi \in X^*(T)^{\Gamma_u}$, to the class of $\mathrm{End}(E_\chi)$ in the Brauer group of F_u .*

Proof. Recall that H^2 of tori over local fields can be described fairly easily. Namely, given an algebraic torus T with the character group $X^*(T)$, the cup-product gives bilinear maps

$$\theta_i : H^i(F_u, X^*(T)) \times H^{2-i}(F_u, T) \rightarrow H^2(F_u, \mathbb{G}_m) \quad (i = 0, 1, 2) \quad (11.12)$$

such that θ_0 defines a duality between the compact group $H^0(F_u, X^*(T))^\wedge$ (the completion of the abelian group $H^0(F_u, X^*(T))$ for the topology given by subgroups of finite index) and the discrete group $H^2(F_u, T)$, in particular, θ_0 induces an isomorphism $H^2(F_u, T) \simeq \text{Hom}_{\mathbb{Z}}(X^*(T)^{\Gamma_u}, \text{Br}(F_u))$ (see II.§5 Theorem 6 in [32]). Namely, any character $\chi \in H^0(F_u, X^*(T)) = X^*(T)^{\Gamma_u}$ defines an F_u -epimorphism from T to \mathbb{G}_m , which in its turn induces the morphism on cohomology

$$\chi^* : H^2(F_u, T) \rightarrow H^2(F_u, \mathbb{G}_m), \quad (11.13)$$

such that for any $\alpha \in H^2(F_u, T)$, we have

$$\theta_0(\chi, \alpha) = \chi^*(\alpha) \in H^2(F_u, \mathbb{G}_m). \quad (11.14)$$

Let \mathcal{E}_α and $\mathcal{E}_{\chi^*(\alpha)}$ be the gerbes associated to α and $\chi^*(\alpha)$, respectively. It follows that we have a morphism of gerbes

$$\mathcal{E}_\alpha \rightarrow \mathcal{E}_{\chi^*(\alpha)} \quad (11.15)$$

that is banded by the epimorphism of bands given by χ . Applying Tannakian duality to (11.15), we obtain a fully faithful functor

$$\text{Rep}(\mathcal{E}_{\chi^*(\alpha)}) \rightarrow \text{Rep}(\mathcal{E}_\alpha) \quad (11.16)$$

that can be viewed as an embedding of the Tannakian category spanned by the simple object E_χ of $\text{Rep}(\mathcal{E}_\alpha)$ corresponding to the character χ . To finish the proof, we notice that the class of such subcategory is defined by an element in $\text{Hom}(\mathbb{Z}, \text{Br}(F_u))$ obtained naturally by computing the endomorphisms of E_χ , and in our case is given by $\chi^*(\alpha)$ (see more on Brauer groups and gerbes in Ch.V, 4.2 in [12]). \square

We apply Proposition 7 to the isomorphism (11.11) to obtain

$$H^2(F_u, \mathbb{D}_F) \simeq \varprojlim \text{Hom}(X^*(\mathbb{D}_{S,K})^{\Gamma_u}, \text{Br}(F_u)) = \text{Hom}(X^*(\mathbb{D}_F)^{\Gamma_u}, \text{Br}(F_u)). \quad (11.17)$$

Consequently, the equality of classes in $H^2(F_u, \mathbb{D}_F)$ follows from the equality of homomorphisms $X^*(\mathbb{D}_F) \rightarrow \text{Br}(F_u)$.

11.6. Localization of the Kottwitz category. This section is based on Section 7 of [20], where Kottwitz defined localization maps for finite layers of $B(F, G)$.

The localization of the Kottwitz category is defined via the sequence of maps of character modules

$$X^*(\mathbb{D}_F) \rightarrow X^*(\mathbb{T}_F) \rightarrow \mathbb{Q}, \quad (11.18)$$

that arises as a direct limit of

$$\mu'_w : X^*(\mathbb{D}_{S,K}) \xrightarrow{b'} X^*(\mathbb{T}_{S,K}) \xrightarrow{\mu_w} \mathbb{Z} = X^*(\mathbb{G}_m), \quad (11.19)$$

where the first map is an inclusion and the second is given by $\mu_w(\sum n_v v) = n_w$, where w is a chosen place in S_K lying above u . Recall that $H^2(\text{Gal}(K_w/F_u), \mathbb{G}_m(K_w)) = \frac{1}{[K_w:F_u]} \mathbb{Z}/\mathbb{Z} \hookrightarrow \mathbb{Q}/\mathbb{Z} = H^2(F_u, \mathbb{G}_m)$, consequently the canonical class $\alpha(K_w/F_u)$ corresponds to the homomorphism

$$n_w \mapsto \frac{n_w}{[K_w:F_u]} \pmod{\mathbb{Z}}.$$

Thus the homomorphism $X^*(\mathbb{D}_{S,K}) \rightarrow \text{Br}(F_u)$ is given by the composition $\alpha(K_w/F_u) \circ \mu'_w$. If $a = \sum n_v v \in X^*(\mathbb{D}_{S,K})$ then

$$\alpha(K_w/F_u) \circ \mu'_w(a) = \frac{n_w}{[K_w:F_u]} \pmod{\mathbb{Z}}. \quad (11.20)$$

11.7. The local class corresponding to φ -spaces. Let $\text{Div}^0(S, K) \otimes \mathbb{Q}$ denote a subgroup of degree 0 rational divisors with supports in S_K , and let $\widetilde{\mathbb{D}}_{S,K}$ be the dual protorus. Assume that S satisfies the condition (K1) (see section 3.2). Then local duality applies, since such a module of characters can be represented as a direct limit of $\text{Div}^0(S, K)$ with transition maps being multiplication by natural numbers. Then the same argument as we used to prove the isomorphism (11.11) shows that $H^2(F_u, \widetilde{\mathbb{D}}_{S,K}) = \text{Hom}(\text{Div}^0(S, K)^{\Gamma_u} \otimes \mathbb{Q}, \text{Br}(F_u))$.

Consequently, Theorem 7 provides us with an element in $H^2(F_u, \mathbb{D}_F)$, since the rational number r_w/d_w from its statement is a slope of the corresponding isocrystal over F_u , the endomorphism algebra of which has the invariant $r_w/d_w \pmod{\mathbb{Z}}$ in $\text{Br}(F_u)$. Strictly speaking, such maps were only defined for the subset of rational divisors of degree 0 in K that satisfy the primitivity condition imposed on φ -pairs, but they extend to the whole group of rational divisors via transition maps.

11.8. Conclusion. Recall that section 11.5 reduces the comparison of the category of Drinfeld φ -spaces over \mathbb{F}_q and the category of representations of the Kottwitz gerbe $\text{Rep}(\text{Kt}_F)$ for a global function field F to the comparison of their localizations for every place u in F . The latter is reduced to establishing the equality of maps on characters of bands.

Taking into account that the formula (11.20) and the expression for r_w/d_w in Theorem 7 are the same modulo the isomorphism described in section 11.4, we conclude that the categories in question are equivalent.

11.9. Isoshtukas with G -structure. Let G be a linear algebraic group over F . One defines an isoshtuka with G -structure as a tensor functor from $\text{Rep}(G)$ to the category of isoshtukas. Since the latter is equivalent to $\text{Rep}(\text{Kt}_F)$ we can apply results from Section 6 to obtain the following.

Theorem 9. *Let F be a global function field, then $B(F, G)$ is isomorphic to the set of isomorphism classes of isoshtukas with G -structure.*

12. MORE ON FIBER FUNCTORS

In this section, we prove the following result.

Theorem 10. *For any finite Galois extension K/\mathbb{Q} and any finite set of places of \mathbb{Q} satisfying the conditions 3.2, the class $\alpha_{S,K}$ of the Kottwitz gerbe is split by a cyclotomic extension of the form $\mathbb{Q}(\zeta_q)$, where q is a prime number.*

This is done in two steps: in Section 12.1 we reduce the question about the vanishing of restrictions of the class $\alpha_{S,K}$ to a question about the vanishing of the corresponding class in adelic cohomology, which in its turn provides us with a criterion of splitting formulated purely in terms of the local behaviour of the Galois extension to which $\alpha_{S,K}$ is restricted. The second step is performed in Section 12.3, where we show that there exists infinitely many integer primes q , such that $\mathbb{Q}(\zeta_q)$ has the desired local behaviour.

12.1. Reduction to an existence theorem. Let K/F be a finite Galois extension of number fields and let S be a finite set of places in F , satisfying the usual conditions (see 3.2). We consider the Tate-Nakayama class $\alpha_{S,K} \in H^2(F, \mathbb{D}_{S,K})$, where $\mathbb{D}_{S,K}$ is the algebraic torus over F introduced in Definition 3.

For any finite Galois extension L/F we can consider the natural restriction map

$$H^2(F, \mathbb{D}_{S,K}) \xrightarrow{\text{res}_L} H^2(L, \mathbb{D}_{S,K} \times_{\text{Spec}(F)} \text{Spec}(L)) \quad (12.1)$$

and ask whether $\text{res}_L(\alpha_{S,K}) = 0$. Since we do not have an explicit description of $\alpha_{S,K}$, but know the images of this class under localization maps, we may consider the following commutative

diagram

$$\begin{array}{ccc}
H^2(F, \mathbb{D}_{S,K}) & \xrightarrow{\text{res}_L} & H^2(L, \mathbb{D}_{S,K} \times_{\text{Spec}(F)} \text{Spec}(L)) \\
\downarrow \text{loc}_F & & \downarrow \text{loc}_L \\
H^2(\mathbb{A}_F, \mathbb{D}_{S,K}) & \xrightarrow{\text{Res}_L} & H^2(\mathbb{A}_L, \mathbb{D}_{S,K} \times_{\text{Spec}(F)} \text{Spec}(L))
\end{array} \tag{12.2}$$

where loc_F and loc_L are the natural maps to adelic cohomology (see Section 9.1), and Res is the restriction map induced on adelic cohomology.

In order to show that $\text{res}_L(\alpha_{S,K}) = 0$, it is enough to show that $\text{Res}_L(\text{loc}_F(\alpha_{S,K})) = 0$ and that $\text{res}_L(\alpha)$ lies in the kernel of loc_L if and only if it is 0. We know that the localization map is not injective in general, however we showed in Section 9.2 that the projective limit of the kernels vanishes.

Assume that $L \cap K = F$, in this case we are basically mimicking the argument from Section 9.2, namely we have an isomorphism

$$\mathbb{D}_{S,K} \times_{\text{Spec}(F)} \text{Spec} L = \left(\prod_{\dot{v} \in \dot{S}} \text{Res}_{K^{\dot{v}}L/L} \mathbb{G}_{m, K^{\dot{v}}L} \right) / \mathbb{G}_{m, L}, \tag{12.3}$$

moreover, it follows from the vanishing of $H^1(L, \mathbb{D}_{S,K})$ (see Lemma 3.1) and from the vanishing of $H^3(L, \mathbb{G}_m)$ (see Corollary I.4.21 in [24]) that $H^2(L, \mathbb{D}_{S,K}) = (\oplus_{\dot{v} \in \dot{S}} \text{Br}(K^{\dot{v}}L)) / \text{Br}(L)$. Again, applying the snake lemma to the diagram similar to (9.8), we obtain that the kernel of the localization map loc_L is a finite subgroup of \mathbb{Q}/\mathbb{Z} of order $\text{gcd}_{\dot{v} \in \dot{S}} \text{deg}(K^{\dot{v}}L/L)$. By disjointness of L and K over F , the latter is equal to $\text{gcd}_{\dot{v} \in \dot{S}} \text{deg}(K^{\dot{v}}/F)$.

Assume that $E \cap L = F$. Note that by the local-global principle, there exists a finite Galois extension $E/K/F$ and a finite set of places S' in E lying above S , such that if $\text{res}_L(\alpha_{S',E})$ is in the kernel of the localization map $H^2(L, \mathbb{D}_{S',E}) \rightarrow H^2(\mathbb{A}_L, \mathbb{D}_{S',E})$, then $\text{res}_L(\alpha_{S,K}) = \pi_{S',E}(\text{res}_L(\alpha_{S',E})) = 0$.

Consequently, after replacing (S, K) with (S', E) in the diagram (12.2), we see that it is enough to find a finite Galois extension L linearly disjoint from E and K that satisfies the following condition:

- $\text{Res}_L(\alpha_{S',E}) = 0 \in H^2(\mathbb{A}_L, \mathbb{D}_{S',E})$.

By construction of $\alpha_{S',E}$, the class $\text{loc}_F(\alpha_{S',E})$ is the image of the semi-local canonical class $\alpha_2(S', E)$ under the map $H^2(\mathbb{A}_F, \mathbb{T}_{S',E}) \rightarrow H^2(\mathbb{A}_F, \mathbb{D}_{S',E})$ (see Section 11.3). Therefore, we are reduced to proving that $\text{Res}_L(\alpha_2(S', E)) = 0$. Using the fact that $\alpha_2(S', E)$ is defined very explicitly (see Section 3.3, in particular, the isomorphism (3.6)), we obtain that $\text{Res}_L(\alpha_2(S', E))$ vanishes if $\alpha(E_{\dot{v}}/F_v) \in \text{Br}(E_{\dot{v}}/F_v)$ is killed by the map

$$H^2((E^{\dot{v}})_v, \mathbb{G}_m) = \text{Br}(F_v) \xrightarrow{\text{res}_{L_w}} \oplus_{\tilde{w} \in V_{E^{\dot{v}}L, \tilde{w}|w}} H^2((E^{\dot{v}}L)_{\tilde{w}}, \mathbb{G}_m) = \oplus_{\tilde{w}|w} \text{Br}((E^{\dot{v}}L)_{\tilde{w}}) \tag{12.4}$$

where w is a place in L , such that $w \mid v$. On the left-hand side we used that $E^{\dot{v}}$ is the decomposition field of a place lying over v . Consider an induced map $\text{Br}(F_v) \rightarrow \text{Br}((E^{\dot{v}}L)_{\tilde{w}})$, where \tilde{w} is as above. Since it comes from the restriction map res_{L_w} , it follows from local class field theory that this map can be identified with the multiplication by $\text{deg}((E^{\dot{v}}L)_{\tilde{w}}/F_v)$ on \mathbb{Q}/\mathbb{Z} .

In order to compute the degree of the extension of local fields $(E^{\dot{v}}L)_{\tilde{w}}/F_v$, we notice that $L \otimes_F E^{\dot{v}} \otimes_F F_v = (\prod_{w|v, w \in V_L} L_w)^g$, where g is the number of places of E lying over v , therefore the desired degree is given by $\text{deg}(L_w/F_v)$.

Combining all the reduction steps, we see that any extension L/F such that

- (1) $L \cap E = F$,

- (2) for any $w \in V_L$ lying above $v \in S$, the local degree $\deg(L_w/F_v)$ is divisible by $\deg(E_{v'}/F_v)$, where $v' \in S_E$ is any lift of v ,

splits the canonical class $\alpha_{S,K}$.

In the following sections, we show that infinitely many extensions $\mathbb{Q}(\zeta_q)$, where q is a prime, satisfy the second condition, therefore, the first condition poses no problem. We also notice that $\mathbb{Q}(\zeta_q)$ is a purely imaginary extension of \mathbb{Q} , thus the local degrees of $\mathbb{Q}(\zeta_q)/\mathbb{Q}$ at the infinite primes are all equal to 2. This way we may restrict ourselves to finite primes of S .

12.2. Auxiliary result: Kummer extensions of cyclotomic fields. First, we recall a result from theory of Kummer extensions of cyclotomic number fields that will be crucial in Section 12.3.

Theorem 11 (Theorem 1 in [29]). *Let K be a number field and let G be a finitely generated subgroup of K^\times . Consider a Kummer extension $K_{n,n} = K(\zeta_n, \sqrt[n]{G})$ of the n th cyclotomic extension $K_n = K(\zeta_n)$. If G is of strictly positive rank s , then there is an integer $c \geq 1$ (depending only on K and G) such that for all integers $n \geq 1$ we have*

$$\frac{n^s}{\text{Gal}(K_{n,n} : K_n)} \mid c \quad (12.5)$$

where $\frac{n^s}{\text{Gal}(K_{n,n} : K_n)}$ is a positive integer.

12.3. Existence theorem. The proof of the following theorem will occupy the rest of this section.

Theorem 12. *Let $S = \{p_1, \dots, p_s\}$ be a fixed finite set of primes in \mathbb{Q} . Then there exists a constant $c \in \mathbb{Z}$ that depends only on S such that for any integer r satisfying $c \mid r$, there exists infinitely many prime numbers $q \notin S$ with the following properties:*

- (\star) $q \equiv 1 \pmod{r}$, $\gcd(\frac{q-1}{r}, r) = c$.
- ($\star\star$) let L denote $\mathbb{Q}(\zeta_q)$, then for every $p_i \in S$, and for any prime ideal P_i in \mathcal{O}_L lying above p_i the degree of the local extension L_{P_i}/\mathbb{Q}_{p_i} is divisible by $\frac{r}{c}$.

To simplify notation we denote $r_* = \frac{r}{c}$, where c is an integral divisor of r . We will specify this constant below.

We rephrase the condition ($\star\star$) as follows. Note that since $q \notin S$, every extension L_{P_i}/\mathbb{Q}_{p_i} as above is unramified of degree f_i , which is computed as the order of p_i in $(\mathbb{Z}/q\mathbb{Z})^\times$, more concretely, f_i is the smallest natural number such that $p_i^{f_i} \equiv 1 \pmod{q}$ (see Theorem 2.13 in [36]). Thus, the degree of L_{P_i}/\mathbb{Q}_{p_i} is divisible by r_* if and only if f_i is divisible by r_* .

Focusing on the prime $p = p_i$, we see that its order in $(\mathbb{Z}/q\mathbb{Z})^\times$ can be computed by analysing the extension $\mathbb{Q}(\zeta_r, p^{1/r})/\mathbb{Q}(\zeta_r)$, in which q is unramified. Since $r \mid q - 1$, the degree of the local extension over q is equal to the degree of the extension $\mathbb{F}_q(p^{1/r})/\mathbb{F}_q$. Using Kummer theory (see Chapter IV, Theorem 3.3 in [28]), we obtain that $\text{Frob}_q \in \text{Gal}(\mathbb{F}_q(p^{1/r})/\mathbb{F}_q)$ has order equal to the order of $p^{\frac{q-1}{r}}$ in \mathbb{F}_q^\times . More precisely, in this case $\mathbb{F}_q^\times/(\mathbb{F}_q^\times)^r \simeq \mathbb{Z}/(r)$ is isomorphic to the group of characters $\text{Hom}(\text{Gal}(\overline{\mathbb{F}_q}/\mathbb{F}_q), \mu_r)$ via the map

$$p \mapsto \chi_p(\text{Frob}_q) = \frac{\text{Frob}_q(p^{1/r})}{p^{1/r}} = p^{\frac{q-1}{r}} \in \mu_r.$$

Thus, the order of p in $\mathbb{F}_q^\times/(\mathbb{F}_q^\times)^{r_*}$ is equal to the order of $\chi_p(\text{Frob}_q)$, and the latter is equal to the order of $\text{Frob}_q \in \text{Gal}(\mathbb{F}_q(p^{1/r})/\mathbb{F}_q)$. In particular, the order of $\text{Frob}_q \in \text{Gal}(\mathbb{F}_q(p^{1/r})/\mathbb{F}_q)$ divides the order of p in \mathbb{F}_q^\times .

In order to combine the local conditions for different p_i 's, we consider the following tower of extensions

$$\begin{array}{c}
\mathbb{Q}_{r^2,r} := \mathbb{Q}(\zeta_{r^2}, \sqrt[r]{p_1}, \dots, \sqrt[r]{p_s}) \\
\swarrow \quad \searrow \\
\mathbb{Q}_{r,r} := \mathbb{Q}(\zeta_r, \sqrt[r]{p_1}, \dots, \sqrt[r]{p_s}) \quad \mathbb{Q}_{r^2} := \mathbb{Q}(\zeta_{r^2}) \\
\swarrow \quad \searrow \\
E \\
\downarrow \\
\mathbb{Q}_r := \mathbb{Q}(\zeta_r) \\
\downarrow \\
\mathbb{Q}
\end{array}$$

Here E denotes the intersection of $\mathbb{Q}_{r,r}$ and \mathbb{Q}_{r^2} .

We claim that there are infinitely many rational primes q , such that $\text{Frob}_q \in \text{Gal}(\mathbb{Q}_{r^2,r}/\mathbb{Q})$ satisfies the following conditions:

- (1) $\text{Frob}_q = 1 \in \text{Gal}(\mathbb{Q}_r/\mathbb{Q})$.
- (2) Frob_q generates the subgroup $c\mathbb{Z}/r\mathbb{Z} \subseteq \text{Gal}(\mathbb{Q}_{r^2}/\mathbb{Q}_r)$.
- (3) for any $1 \leq i \leq s$ and for every projection $\text{Gal}(\mathbb{Q}_{r,r}/\mathbb{Q}_r) \rightarrow \text{Gal}(\mathbb{Q}_r(\sqrt[r]{p_i})/\mathbb{Q}_r)$ the image of Frob_q in the latter Galois group is at least of order $\frac{r}{c}$.

Note that it follows from the discussion above that the third condition implies that the orders of p_i 's in \mathbb{F}_q^\times are divisible by $\frac{r}{c}$.

First, we show that $H = \text{Gal}(\mathbb{Q}_{r,r}/E)$ contains a subgroup $(c\mathbb{Z}/r\mathbb{Z})^s$. Notice that by definition of E , the group H is isomorphic to $\text{Gal}(\mathbb{Q}_{r^2,r}/\mathbb{Q}_{r^2})$. Letting $n = r^2$ in Theorem 11, we obtain

$$\frac{r^{2s}}{\deg(\mathbb{Q}_{r^2,r^2}/\mathbb{Q}_{r^2})} = \frac{r^s}{\deg(\mathbb{Q}_{r^2,r}/\mathbb{Q}_{r^2})} \cdot \frac{r^s}{\deg(\mathbb{Q}_{r^2,r^2}/\mathbb{Q}_{r^2,r})} \mid c_S \quad (12.6)$$

Since both factors in (12.6) are integers due to the fact that the corresponding extensions are Kummer extensions, we see that $\frac{r^s}{\deg(\mathbb{Q}_{r^2,r}/\mathbb{Q}_{r^2})}$ is an integer dividing c_S . On the other hand, H is a subgroup of $(\mathbb{Z}/(r))^s$, thus any generator of the preimage of any factor $\mathbb{Z}/(r)$ is at least of order r/c_S , hence $(c_S\mathbb{Z}/r\mathbb{Z})^s \subseteq H$. Finally, we set

$$c = c_S.$$

Note that since $\mathbb{Q}_{r^2,r}$, $\mathbb{Q}_{r,r}$, and \mathbb{Q}_{r^2} are abelian extensions over \mathbb{Q}_r , the Galois group of $\mathbb{Q}_{r^2,r}$ over \mathbb{Q}_r is a fiber product of $\text{Gal}(\mathbb{Q}_{r,r}/\mathbb{Q}_r)$ and of $\text{Gal}(\mathbb{Q}_{r^2}/\mathbb{Q}_r)$ over $\text{Gal}(E/\mathbb{Q}_r)$. Next, we consider the diagram

$$\begin{array}{ccccccc}
1 & \longrightarrow & H & \longrightarrow & \text{Gal}(\mathbb{Q}_{r^2,r}/\mathbb{Q}_r) & \longrightarrow & \text{Gal}(\mathbb{Q}_{r^2}/\mathbb{Q}_r) = \mathbb{Z}/(r) \longrightarrow 1 \\
& & \parallel & & \downarrow & & \downarrow \\
1 & \longrightarrow & H & \longrightarrow & \text{Gal}(\mathbb{Q}_{r,r}/\mathbb{Q}_r) & \longrightarrow & \text{Gal}(E/\mathbb{Q}_r) \longrightarrow 1
\end{array} \quad (12.7)$$

Let σ denote a generator of $c\mathbb{Z}/r\mathbb{Z} \subseteq \text{Gal}(\mathbb{Q}_{r^2}/E)$, and let $x = (c, \dots, c) \in (c\mathbb{Z}/r\mathbb{Z})^s \subseteq H$, then both σ and x project to 1 in $\text{Gal}(E/\mathbb{Q}_r)$, since $\text{Gal}(E/\mathbb{Q}_r)$ is a quotient of $\mathbb{Z}/(r)$ and its order divides c . It follows that $(x, \sigma) \in \text{Gal}(\mathbb{Q}_{r^2,r}/\mathbb{Q}_r)$. Applying the Chebotarev density theorem to the conjugacy class of (x, σ) , we see that there are infinitely many primes $q \in \mathbb{Q}_r$ such that Frob_q lies in the chosen conjugacy class. Notice that by construction $\text{Frob}_q = 1$ in $\text{Gal}(\mathbb{Q}_r/\mathbb{Q})$.

In order to check the third condition, we need to show that for every $i = 1, \dots, s$, the Frobenius element $\text{Frob}_{q,i}$ in $\text{Gal}(\mathbb{Q}_r(\sqrt[r]{p_i})/\mathbb{Q}_r)$ has the correct order. Since q is unramified in $\mathbb{Q}_{r^2,r}$, we see

that $\text{Frob}_{q,i} = \pi_i(\text{Frob}_q)$, where π_i is defined as the sequence of natural projections

$$\pi_i : \text{Gal}(\mathbb{Q}_{r^2,r}/\mathbb{Q}_r) \rightarrow \text{Gal}(\mathbb{Q}_{r,r}/\mathbb{Q}_r) \rightarrow \text{Gal}(\mathbb{Q}_r(\sqrt[r]{p_i})/\mathbb{Q}_r). \quad (12.8)$$

By construction, $\pi_i(\text{Frob}_q)$ belongs to the conjugacy class of the element

$$c \in \text{Gal}(\mathbb{Q}_r(\sqrt[r]{p_i})/\mathbb{Q}_r) \subseteq \mathbb{Z}/r\mathbb{Z}, \quad (12.9)$$

therefore, the order of $\text{Frob}_{q,i}$ is divisible by $\frac{r}{c}$, which completes the proof of the theorem.

12.4. End of the proof of Theorem 10. We apply Theorem 12 to the conditions obtained in the end of Section 12.1. Namely, we let r be divisible by the least common multiple of local degrees $l = \text{lcm}_{v \in S}(\deg(E_{v'}/\mathbb{Q}_v))$, where $v \in S$ and $v' \in S_E$ is any lift of v . It follows that there exists infinitely many primes $q \notin S$, such that local degrees of $\mathbb{Q}(\zeta_q)$ are divisible by an integer r/c , where c does not depend on r . Therefore, we can achieve that r/c is divisible by l by simply enlarging r multiplicatively.

13. MOTIVES OVER \mathbb{F} AND $\text{Rep}(\text{Kt}_{\mathbb{Q}})$

Let \mathbb{F} be the algebraic closure of the finite field \mathbb{F}_p . In this section we show that $\text{Rep}(\text{Kt}_{\mathbb{Q}})$ has a full subcategory that under the assumption of Tate's conjecture over \mathbb{F} can be identified with $\text{Mot}(\mathbb{F})$ the category of motives over \mathbb{F} . The idea of describing motives in terms of gerbes and non-abelian cohomology goes back to Grothendieck and was pushed further in the case of $\text{Mot}(\mathbb{F})$ in the work of Langlands and Rapoport ([21]), where an explicit description of the class of $\text{Mot}(\mathbb{F})$ appeared. It was however corrected and simplified significantly in the following works of Milne (see [26], [25]).

13.1. Weil numbers and their germs. Let $q = p^n$. Recall that an algebraic number $\pi \in \mathbb{Q}^{\text{al}}$ is said to be a Weil q -number of weight m if

- for every embedding $\rho : \mathbb{Q}[\pi] \hookrightarrow \mathbb{C}$, $\rho(\pi) \cdot \overline{\rho(\pi)} = q^m$.
- for some l , $q^l \pi$ is an algebraic integer.

The set of all Weil q -numbers in \mathbb{Q}^{al} is an abelian group denoted by $W(q)$. If $q' = p^{n'}$ and $n \mid n'$, we have a group homomorphism $W(q) \rightarrow W(q')$: $\pi \mapsto \pi^{n'/n}$. Define the group of germs of Weil numbers

$$W = \varinjlim W(q). \quad (13.1)$$

There is an action of the absolute Galois group of \mathbb{Q} on W , and P is defined as a protorus over \mathbb{Q} such that

$$X^*(P) = W. \quad (13.2)$$

In order to represent P as a projective limit of algebraic tori, we rewrite W as a colimit of finitely generated Galois modules. Namely, for any CM-field K/\mathbb{Q} , we define W^K as a subset of $\pi \in W$ having a representative in K and such that for any place $v \mid p$ in K :

$$\frac{\text{ord}_v(\pi)}{\text{ord}_v(p^n)} \cdot [K_v : \mathbb{Q}_p] \in \mathbb{Z}. \quad (13.3)$$

Define P^K as a group of multiplicative type defined by $\text{Gal}(\mathbb{Q}^{\text{al}}/\mathbb{Q})$ -module W^K . Then $W = \varinjlim_K W^K$ and $P = \varprojlim_K P^K$, where the limit is being taken over the directed set of finite CM-extensions of \mathbb{Q} .

Let K^+ denote the maximal totally real subfield of K , and let $V_{p,K}$ and V_{p,K^+} denote the set of p -adic primes in K and K^+ respectively. Then there is an exact sequence of Galois modules

$$0 \rightarrow W^K \xrightarrow{\alpha} \bigoplus_{v \in V_{p,K}} \mathbb{Z} \cdot v \bigoplus \mathbb{Z} \xrightarrow{\beta} \bigoplus_{w \in V_{p,K^+}} \mathbb{Z} \cdot w \rightarrow 0, \quad (13.4)$$

where α is the map

$$\pi \mapsto \left(\sum_{v \in V_{p,K}} \frac{\text{ord}_v(\pi)}{\text{ord}_v p^n} \cdot [K_v : \mathbb{Q}_p], \text{wt}(\pi) \right), \quad (13.5)$$

and β is given by the formula

$$\left(\sum_{v \in V_{p,K}} n_v v, m \right) \mapsto \sum_{v \in V_{p,K}} n_v |_{V_{p,K^+}} - [K_v : \mathbb{Q}_p] m \sum_{w \in V_{p,K^+}} w \quad (13.6)$$

Let K' be a CM-field containing K . Then $W^K \subset W^{K'}$, and the transition map can be described naturally using the exact sequence (13.4). Namely, the transition map is induced by the map

$$v \mapsto \sum_{v'|v} [K'_{v'} : K_v] \cdot v', \quad (13.7)$$

when $v \mid p$, and the identity map on \mathbb{Z} .

The short exact sequence of Galois modules (13.4) induces the short exact sequence of the corresponding tori and provides a very useful tool for analysis of the cohomology of P^K . The following results are due to Milne.

Proposition 8. *Let K be a CM-field, and let P^K be a torus attached to the Galois module W^K , then*

- $\varprojlim_K^i H^1(\mathbb{Q}, P^K) = 0$, for $i = 0, 1$.
- $H^2(\mathbb{Q}, P) = \varprojlim_K H^2(\mathbb{Q}, P^K)$.
- $H^2(\mathbb{Q}, P^K) \rightarrow H^2(\mathbb{A}, P^K)$ is injective.
- $H^2(\mathbb{Q}, P) \rightarrow H^2(\mathbb{A}, P)$ is injective.

13.2. W^K and the characters of the Kottwitz protorus. One can also realize W^K as a submodule of the character module of the Kottwitz pro-torus $\mathbb{D}_{K/\mathbb{Q}}$.

Proposition 9. *Let K be a finite CM-extension of \mathbb{Q} . Then there is an injective homomorphism of $\text{Gal}(K/\mathbb{Q})$ modules*

$$\omega_K : W^K \hookrightarrow X^*(\mathbb{D}_{K/F}). \quad (13.8)$$

Proof. Let $\pi \in W^K$ be a representative of a Weil p^n -Number of weight m . We may apply the product formula to π , namely

$$\prod_{v \in V_K} \|\pi\|_v = 1. \quad (13.9)$$

Moreover, for any non-archimedean $v \notin V_{p,K}$, we have $\|\pi\|_v = 1$. Notice that for any $v \mid p$, we have $\|\pi\|_v = p^{-f_v \text{ord}_v(\pi)}$, where f_v is the degree of the extension of the residue field of K_v over \mathbb{Q}_p . The latter can be expressed as $[K_v : \mathbb{Q}_p]/e_v$, where e_v is the ramification index of v in K that can be computed as $\text{ord}_v(p)$. Since K is a CM-field, every archimedean place $\infty' \mid \infty$ is in fact complex, and by definition of Weil p^n -number of weight m , we have $\|\pi\|_{\infty'} = (p^n)^m$. Combining these observations, we deduce

$$\sum_{v \in V_{p,K}} \frac{\text{ord}_v \pi}{\text{ord}_v(p)} \cdot [K_v : \mathbb{Q}_p] - \sum_{\infty' \in V_{\infty,K}} nm = 0. \quad (13.10)$$

Notice that by definition of W^K and from the fact that $\pi \in W^K$, the equality (13.10) is integrally divisible by n (see (13.3)). Define

$$i_K : W^K \rightarrow \mathbb{Z}[V_K], \quad \pi \mapsto \sum_{v \in V_{p,K}} \frac{\text{ord}_v \pi}{\text{ord}_v(p^n)} \cdot [K_v : \mathbb{Q}_p] \cdot v - \sum_{\infty' \in V_{\infty,K}} m \cdot \infty' = 0. \quad (13.11)$$

Since W^K is torsion-free, this map is injective. Moreover, it follows from (13.10) that i_K induces an injective map $W^K \rightarrow X^*(\mathbb{D}_{K/\mathbb{Q}})$. \square

In order to construct the morphism $\varinjlim_K W^K \rightarrow X^*(\mathbb{D}_{\mathbb{Q}})$, we need to check that the morphisms ω_K are compatible with the transition maps $W^K \rightarrow W^{K'}$ and $X^*(\mathbb{D}_{K/\mathbb{Q}}) \rightarrow X^*(\mathbb{D}_{K'/\mathbb{Q}})$, where K'/\mathbb{Q} is a CM-extension containing K . Recall, that the transition maps for the character modules of Kottwitz protori are induced by the maps (5.12), that is by the same formulae as (13.7). We denote by ω the injective homomorphism of $\text{Gal}(\mathbb{Q}^{\text{al}}/\mathbb{Q})$ -modules $W \rightarrow X^*(\mathbb{D}_{\mathbb{Q}})$ induced by ω_K for varying CM-fields K .

13.3. Image of the Kottwitz class under ω . Let $\omega : W \hookrightarrow X^*(\mathbb{D}_{\mathbb{Q}})$ be the homomorphism defined above. By abuse of notation, we also denote by ω the epimorphism of protori $\mathbb{D}_{\mathbb{Q}} \rightarrow P$. Because $H^2(\mathbb{Q}, \mathbb{D}_{\mathbb{Q}}) = \varprojlim H^2(\mathbb{Q}, \mathbb{D}_{S,K})$ and $H^2(\mathbb{Q}, P) = \varprojlim H^2(\mathbb{Q}, P^K)$, there is a well-defined morphism

$$\omega^* : H^2(\mathbb{Q}, \mathbb{D}_{\mathbb{Q}}) \rightarrow H^2(\mathbb{Q}, P). \quad (13.12)$$

We are interested in computing the class $\omega^*(\alpha_{\mathbb{Q}})$, where $\alpha_{\mathbb{Q}}$ denotes the class of the Kottwitz gerbe in $H^2(\mathbb{Q}, \mathbb{D}_{\mathbb{Q}})$. By Proposition 8, $H^2(\mathbb{Q}, P) \hookrightarrow H^2(\mathbb{A}, P)$, therefore it suffices to describe the image of the given class locally.

Proposition 10. *The image of $\omega^*(\alpha_{\mathbb{Q}})$ in $H^2(\mathbb{A}, P)$ is given by the class $(0_{\mathbb{A}\{p,\infty\}}, \alpha_p, \alpha_{\mathbb{R}})$. Where α_p and $\alpha_{\mathbb{R}}$ are the canonical local classes.*

Proof. Let Γ_l denote the absolute Galois group of \mathbb{Q}_l . Recall that since $\lim^1 H^1(\mathbb{Q}_l, P) = 0$, we may apply arguments of Section 11.5, to describe local cohomology groups of the Weil protorus, namely

$$H^2(\mathbb{Q}_l, P) \simeq \text{Hom}_{\mathbb{Z}}(W^{\Gamma_l}, \mathbb{Q}/\mathbb{Z}), \quad (13.13)$$

where l any non-archimedean prime.

Recall also that the map $\omega : W \hookrightarrow X^*(\mathbb{D}_{\mathbb{Q}})$ induces the map of Γ_l -modules

$$\text{loc}_l : W \xrightarrow{\omega} X^*(\mathbb{D}_{\mathbb{Q}}) \hookrightarrow X^*(\mathbb{T}_{\mathbb{Q}}) \xrightarrow{\mu_l} \mathbb{Q}, \quad (13.14)$$

where μ_l is defined in Section 11.6. Namely, μ_l sends a sum of valuations with rational coefficients to the coefficient corresponding to the fixed l -adic prime of \mathbb{Q}^{al} .

Therefore, for any $l \neq p$, we have that the image of W in \mathbb{Q} the composition loc_l in (13.14) is 0. Therefore, the corresponding class in $H^2(\mathbb{Q}_l, P)$ is trivial.

If $l = p$, we see that the composition $W^{\Gamma_p} \rightarrow W \xrightarrow{\text{loc}_p} \mathbb{Q} \xrightarrow{[K_v:\mathbb{Q}_p]^{-1}} \mathbb{Q}$ is surjective. It follows from the fact that for any CM-field K and $\pi \in W^K$ we have $\text{loc}_v(\pi) = \frac{\text{ord}_v(\pi)}{\text{ord}_v(p^n)} \cdot [K_v : \mathbb{Q}_p] \in \mathbb{Z}$, where v is the p -adic place of K induced by the fixed p -adic place of \mathbb{Q}^{al} . If π is moreover Γ_v -invariant, where $\Gamma_v = \text{Gal}(K_v/\mathbb{Q}_p)$ is the decomposition group of v , then we deduce from the product formula and equality (13.10) that $\text{ord}_v(\pi)/\text{ord}_v(p^n)$ takes value $\frac{m}{2}$, where m is the weight of π . We obtain that the colimit of the cokernels of the maps

$$(W^K)^{\Gamma_v} \xrightarrow{\text{loc}_v} \mathbb{Z} \xrightarrow{[K_v:\mathbb{Q}_p]^{-1}} \frac{1}{2 \cdot [K_v : \mathbb{Q}_p]} \mathbb{Z}, \quad \pi \mapsto \frac{\text{ord}_v(\pi)}{\text{ord}_v(p^n)} \quad (13.15)$$

vanishes, since their orders are bounded by $[K_v : \mathbb{Q}_p]$, and the transition maps for $K \subset K'$, where K' is a finite CM-field over \mathbb{Q} , are given by multiplication by the local degrees $[K'_v : K_v]$, hence kill the cokernels eventually.

Consequently, the map $\text{Hom}(\mathbb{Q}, \mathbb{Q}/\mathbb{Z}) \rightarrow \text{Hom}(W^{\Gamma_p}, \mathbb{Q}/\mathbb{Z})$ is injective and maps the class of the natural projection $\mathbb{Q} \rightarrow \mathbb{Q}/\mathbb{Z}$, that is the class α_p , to the image of $\omega^*(\alpha_{\mathbb{Q}})$ in $H^2(\mathbb{Q}_p, P)$.

To complete the proof, we notice that when $l = \infty$ is the archimedean place, then $\text{loc}_\infty(\pi) = -m$. Hence the image of $\omega^*(\alpha_\mathbb{Q})$ in $H^2(\mathbb{R}, P)$ coincides with the image of the projection $\mathbb{Z} \rightarrow \mathbb{Z}/2\mathbb{Z}$ under the map $\text{Hom}(\mathbb{Z}, \text{Br}(\mathbb{R})) \rightarrow \text{Hom}(W^{\Gamma_\infty}, \text{Br}(\mathbb{R}))$, where $\Gamma_\infty \simeq \text{Gal}(\mathbb{C}/\mathbb{R})$. \square

13.4. Conjectural relation to $\text{Mot}(\mathbb{F})$. Let $\zeta(V, s)$ denote the zeta function of a variety V over \mathbb{F}_q . Let $A^r(V)$ denote the quotient of the space of algebraic cycles of codimension r on V by the subspace of cycles numerically equivalent to 0. In this section we assume the Tate conjecture.

Conjecture 2 (Tate conjecture). *For all smooth projective varieties over \mathbb{F}_q and $r \geq 0$, the dimension of $A^r(V)$ is equal to the order of the pole of $\zeta(V, s)$ at $s = r$.*

Under this assumption the category of motives over \mathbb{F} can be described fairly explicitly. Namely, it follows from theorem of Jannsen, that $\text{Mot}(\mathbb{F})$ is a semi-simple Tannakian category over \mathbb{Q} . Moreover, the classes of simple objects correspond to the orbits of the action $\text{Gal}(\mathbb{Q}^{\text{al}}/\mathbb{Q})$ on W , the abelian group of germs of Weil numbers defined in Section 13.1. Applying Tannakian duality, we see that $\text{Mot}(\mathbb{F})$ can be recovered up to equivalence from the class it defines in $H^2(\mathbb{Q}, P)$ (fpqc-cohomology). Using the fact that $H^2(\mathbb{Q}, P) \hookrightarrow H^2(\mathbb{A}, P)$, we deduce that it suffices to describe the images of the class of $\text{Mot}(\mathbb{F})$ in cohomology groups of completions of \mathbb{Q} at all places. One can apply Proposition 7 to compute the local classes via isocrystals. Namely, it turns out that the localization functor (see Theorem 1.18 in [26] for the definition)

$$\text{Mot}(\mathbb{F}) \otimes \mathbb{Q}_p \rightarrow \text{Isoc}(\mathbb{F})$$

corresponds to the morphism of gerbes induced by the map of characters $W \rightarrow \mathbb{Q}$ given by the formula

$$\pi \mapsto \frac{\text{ord}_p(\pi)}{n} \tag{13.16}$$

where ord_p is the extension of the p -adic valuation on \mathbb{Q} corresponding to the chosen embedding $\mathbb{Q}^{\text{al}} \rightarrow \mathbb{Q}_p^{\text{al}}$. The latter can be deduced from the section ‘‘The isocrystal of a motive’’ in [26]. Finally, one obtains the following description of $\text{Mot}(\mathbb{F})$.

Theorem 13 (Langlands-Rapoport, Milne (Th.3.29 in [26])). *Assume the Tate conjecture. Then the category of motives over \mathbb{F} is a semi-simple Tannakian category over \mathbb{Q} . Its band is given by the protorus P corresponding to the $\text{Gal}(\mathbb{Q}^{\text{al}}/\mathbb{Q})$ -module W . The equivalence class of $\text{Mot}(\mathbb{F})$ in $H^2(\mathbb{Q}, P)$ is identified with the class $(0_{\mathbb{A}_{\{p, \infty\}}}, \alpha_p, \alpha_\mathbb{R}) \in H^2(\mathbb{A}, P)$ via the inclusion $H^2(\mathbb{Q}, P) \hookrightarrow H^2(\mathbb{A}, P)$.*

The naturality of such a description is treated in Theorem 6.3 in [25], which combined with Proposition 10 leads us to the following statement.

Corollary 1. *Let $\omega^*(\text{Rep}(\text{Kt}_\mathbb{Q}))$ be the subcategory of $\text{Rep}(\text{Kt}_\mathbb{Q})$, defined by the image of the global Kottwitz class $\alpha_\mathbb{Q}$ along the map $\omega : W \hookrightarrow X^*(\mathbb{D}_\mathbb{Q})$. There exists an equivalence $\omega^*(\text{Rep}(\text{Kt}_\mathbb{Q})) \rightarrow \text{Mot}(\mathbb{F})$ compatible with the natural realization functors.*

REFERENCES

- [1] *E. Artin and J. Tate.* Class field theory. Benjamin, New York Amsterdam 1967
- [2] *J. F. Barreto Castaneda, F. Jarquin Zarate, M. Rzedowski Calderon, and G. Villa Salvador.* Abelian p-extensions and additive polynomials. Int. J. Math. 28 (2017), No. 14, pp. 34-65.
- [3] *B. Bhatt and P. Scholze.* The pro-étale topology for schemes. arXiv:1309.1198
- [4] *S. Bosch, W. Lütkebohmert, and M. Raynaud.* Néron Models. Ergeb. Math. Grenzgeb., 3. Folge, 21, Springer-Verlag, 1990.
- [5] *L. Breen.* Tannakian categories. Motives, Proc. Symp. Pure Math. 55 part 1, 337-376 (1994).
- [6] *P. Deligne and J. Milne.* Tannakian categories, in Hodge Cycles, Motives, and Shimura Varieties, LNM 900, 1982, pp. 101-228.

- [7] *B. Conrad* Finiteness theorems for algebraic groups over function fields. *Compositio Mathematica*, 148(2), 555–639 (2012).
- [8] *B. Conrad, O. Gabber, and G. Prasad* Pseudo-reductive groups (2nd ed.), Cambridge Univ. Press, (2015).
- [9] *P. Deligne and G. Lusztig*. Representations of reductive groups over finite fields. *Annals of Mathematics*, vol. 103, no. 1, 1976, pp. 103–161.
- [10] *V. G. Drinfeld*. The proof of peterson’s conjecture for $GL(2)$ over a global field of characteristic p . *Funct. Anal. Appl.*, 22:1 (1988), 28–43
- [11] *O. Gabber and S. Kelly*. Points in algebraic geometry. *Pure Appl. Algebr.*, Volume 219, Issue 10 (2015), pp.4667-4680
- [12] *J. Giraud*. *Cohomologie non abélienne*. Springer-Verlag, 1971.
- [13] *A. Grothendieck*. *Le groupe de Brauer III, Dix exposés sur la cohomologie des schémas*, Masson and North-Holland, Paris and Amsterdam, 1968.
- [14] *M. Demazure and A. Grothendieck*. *Schémas en Groupes (SGA 3)*, *Lect. Notes in Math.* 151-152-153, Springer-Verlag, 1970.
- [15] *Théorie des topos et cohomologie étale des schémas. Tome 2*, *Lecture Notes in Mathematics*, Vol. 270, Springer-Verlag, Berlin, 1972 (French). Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964 (SGA 4); Dirigé par M. Artin, A. Grothendieck et J. L. Verdier. Avec la collaboration de N. Bourbaki, P. Deligne et B. Saint-Donat.
- [16] *T. Kaletha*. Global rigid inner forms and multiplicities of discrete automorphic representations, *Inventiones Mathematicae* 213 (2018), no. 1, 271-369
- [17] *T. Kaletha, O. Taïbi*. Global rigid inner forms vs isocrystals, arXiv:1812.11373
- [18] *M. Kisin*. Mod- p points on Shimura varieties of abelian type, *J. Amer. Math. Soc.* 30 (2017), 819-914.
- [19] *R. E. Kottwitz*. Isocrystals with additional structure. II., *Compositio Math.* 109 (1997), no. 3, 255–339.
- [20] *R. E. Kottwitz* $B(G)$ for all local and global fields, arXiv:1401.5728
- [21] *R. Langlands, M. Rapoport*. Shimuravarietäten und Gerben
- [22] *G. Laumon, M. Rapoport, and U. Stuhler*. \mathcal{D} -elliptic sheaves and the Langlands correspondence. *Invent Math* 113, 217–338 (1993).
- [23] *J. Milne*. *Algebraic Groups: The Theory of Group Schemes of Finite Type over a Field*, Cambridge Press (2017).
- [24] *J. Milne*. *Arithmetic Duality Theorems*, Academic Press, 1986
- [25] *J. Milne*. *Gerbes and Abelian Motives*. arXiv:math/0301304
- [26] *J. Milne*. *Motives over Finite Fields*.
- [27] *J. Neukirch, A. Schmidt, and K. Winberg*. *Cohomology of number fields* (2nd ed.), *Grundlehren der Mathematischen Wissenschaften*, vol. 323, Springer-Verlag, Berlin (2008).
- [28] *J. Neukirch*. *Algebraic Number Theory*. Springer (1999).
- [29] *A. Perucca and P. Sgobba*. Kummer theory for number fields and the reductions of algebraic numbers, *International Journal of Number Theory*, Vol. 15, No. 08, pp. 1617-1633 (2019).
- [30] *N. Saavedra Rivano*. *Catégories tannakiennes. Catégories Tannakiennes. Lecture Notes in Mathematics*, Vol. 265. (1972), Springer-Verlag, Berlin.
- [31] *P. Scholze*. “ p -adic geometry”, <http://www.math.uni-bonn.de/people/scholze/Rio.pdf>
- [32] *J.-P. Serre*. *Galois Cohomology*. Springer-Verlag Berlin Heidelberg (1997).
- [33] *J.-P. Serre*. *Corps Locaux*, Hermann, Paris (1962).
- [34] *Stacks Project*
- [35] *J. Tate*. The cohomology groups of algebraic tori in finite Galois extensions of number fields. *Nagoya Math. J.* 27 (1966), 709–719.
- [36] *L. C. Washington*. *Introduction to Cyclotomic fields*, *Grad. Texts in Math.* 83, 2nd edition, Springer Verlag, 1997.