

Algebraic Geometry I

5. Exercise sheet

Exercise 1 (4 points):

Let $U_i, i = 1, 2$, be two schemes. Let $V_i \subseteq U_i$ be open subschemes and let $\varphi: V_1 \xrightarrow{\sim} V_2$ be an isomorphism.

1) Show that there exists a scheme X with open subschemes $W_i \subseteq X$ and isomorphisms $\alpha_i: U_i \xrightarrow{\sim} W_i, i = 1, 2$, such that $\alpha_i^{-1}(W_1 \cap W_2) = V_i$ and $\varphi = \alpha_2^{-1} \circ \alpha_1|_{V_1}$.

2) Let A be a ring and let X_{\pm} be the scheme obtained by glueing $U_1 = U_2 = \text{Spec}(A[T])$ along the isomorphism $\varphi_{\pm}: \text{Spec}(A[T, T^{-1}]) \rightarrow \text{Spec}(A[T, T^{-1}]), T \mapsto T^{\pm 1}$. Show that X_{\pm} are not affine schemes and that X_+ is not isomorphic to X_- .

Remark: X_+ is called the affine line over A with doubled origin, X_- is called the projective line over A .

Exercise 2 (4 points):

1) Let (Y, \mathcal{O}_Y) be a locally ringed space. Show that $(Z, \mathcal{O}_Z) \mapsto Z$ induces a bijection between open subsets of Y and equivalence classes of open immersions $(Z, \mathcal{O}_Z) \rightarrow (Y, \mathcal{O}_Y)$ of locally ringed spaces. Here, open immersions are equivalent if they are isomorphic as locally ringed spaces over (Y, \mathcal{O}_Y) .

2) Let k be an algebraically closed field. For a (classical) quasi-projective variety $X \subseteq \mathbb{P}_k^n(k)$ let \mathcal{O} be its sheaf of regular functions $U \mapsto \mathcal{O}(U)$. Let $\pi: X \rightarrow X^{\text{sob}}$ be the soberification (Sheet 3, Exercise 4). Show that $X^{\text{sch}} := (X^{\text{sob}}, \pi_*\mathcal{O})$ is a scheme over $\text{Spec}(k)$ and that $X \mapsto X^{\text{sch}}$ is a fully faithful functor from the category of quasi-projective varieties to the category of schemes over $\text{Spec}(k)$.

Hint: Reduce to the case of affine algebraic sets by glueing morphisms of locally ringed spaces.

Exercise 3 (4 points):

1) Show that the functor

$$\Phi: \{\text{schemes}\} \rightarrow \text{Fun}(\text{Rings}, \text{Sets}), X \mapsto (R \mapsto \text{Hom}_{\text{Lrs}}(\text{Spec}(R), X))$$

is fully faithful by reducing to the Yoneda lemma.

2) Show that the functors $R \mapsto F_n(R) := \{x \in R \mid x^n = 1\}$ for $n \geq 1$, and $R \mapsto G(R) := \{(x, y) \in R^2 \mid R^2 \xrightarrow{(x, y)} R \text{ surjective}\}$ lie in the essential image of Φ .

Exercise 4 (4 points):

1) Show that if a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ admits a left/right adjoint, then this adjoint is unique up to canonical isomorphism.

2) Assume that $\mathcal{C} \xrightarrow{F} \mathcal{D}, \mathcal{D} \xrightarrow{\tilde{F}} \mathcal{E}$ are left adjoint functors with right adjoints G and \tilde{G} . Show that $\tilde{F} \circ F$ is left adjoint to $G \circ \tilde{G}$.

3) Let $f: Y \rightarrow X$ be a continuous map of topological spaces. Let $f^{p,*}(-)$ resp. $f^*(-)$ be the presheaf resp. sheaf pullback, and let $(-)^{\sharp}$ denote sheafification. Show that there exists a natural isomorphism $f^{p,*}(\mathcal{F})^{\sharp} \cong f^*(\mathcal{F}^{\sharp})$ for any presheaf \mathcal{F} (of sets/abelian groups/...) on X .

To be handed in on: Thursday, 16.11.2023 (during the lecture, or via eCampus).