

On the derivative of an Eisenstein series of weight one

by

Stephen S. Kudla¹

Michael Rapoport

and

Tonghai Yang²

Introduction.

In [17], a certain family of Siegel Eisenstein series of genus g and weight $\frac{g+1}{2}$ was introduced which have an odd functional equation and hence have a natural zero at their center of symmetry ($s = 0$). It was suggested that the derivatives at $s = 0$ of such series, which we will refer to as incoherent Eisenstein series, should have some connection with arithmetical algebraic geometry, and some evidence was provided in the case of genus 2 and weight $3/2$. In that case, certain of the Fourier coefficients of the central derivative were shown to involve (parts of) the height pairing of Heegner points on Shimura curves. Additional evidence occurred earlier in the work of Gross and Keating, [12], where, implicitly, derivatives of Siegel Eisenstein series on Sp_3 of weight 2 arise. Higher dimensional cases are studied in [19] (Sp_4 , weight $5/2$) and [21] (Sp_3 , weight 2).

In the present paper, we consider the simplest possible example of an incoherent Eisenstein series and its central derivative. More precisely, let $q > 3$ be a prime congruent to 3 modulo 4. There are two types of Eisenstein series of weight 1 associated to the imaginary quadratic field $\mathbf{k} = \mathbb{Q}(\sqrt{-q})$. The first is a ‘coherent’ Eisenstein series. For $\tau = u+iv$ in the upper half plane and $s \in \mathbb{C}$ with $\text{Re}(s) > 1$, this series has the form

$$(0.1) \quad E_+(\tau, s) = v^{s/2} \sum_{\gamma \in \Gamma_\infty \backslash \Gamma} (c\tau + d)^{-1} |c\tau + d|^{-s} \Phi_q^+(\gamma),$$

where, for $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma = SL_2(\mathbb{Z})$,

$$(0.2) \quad \Phi_q^+(\gamma) = \begin{cases} \chi_q(a) & \text{if } c \equiv 0 \pmod{q}, \\ iq^{-1/2} \chi_q(c) & \text{if } c \text{ is prime to } q. \end{cases}$$

¹Partially supported by NSF Grant DMS-9622987

²Partially supported by NSF Grant DMS-9700777

After analytic continuation to the whole s plane, the normalized Eisenstein series

$$(0.3) \quad E_+^*(\tau, s) = q^{(s+1)/2} \Lambda(s+1, \chi_q) E_+(\tau, s)$$

satisfies the functional equation $E_+^*(\tau, -s) = E_+^*(\tau, s)$. Here

$$(0.4) \quad \Lambda(s, \chi_q) = \pi^{-(s+1)/2} \Gamma\left(\frac{s+1}{2}\right) L(s, \chi_q).$$

The Siegel-Weil formula (due to Hecke in this case, [16], pp.454–5 combined with p.453, (23) and p.475) yields

$$(0.5) \quad E_+^*(\tau, 0) = 2h_{\mathbf{k}} + 4 \sum_{n=1}^{\infty} \rho(n) e(n\tau),$$

where $h_{\mathbf{k}}$ is the class number of \mathbf{k} , and, for a positive integer n , $\rho(n)$ is the number of integral ideals of \mathbf{k} of norm n . The right side of (0.5) is, of course, twice the sum of Hecke's theta functions of weight 1, [16], p.440 (11),

$$(0.6) \quad \vartheta(\tau, \mathfrak{a}) = \sum_{x \in \mathfrak{a}} e\left(\frac{N(x)}{N(\mathfrak{a})} \tau\right)$$

as the fractional ideal \mathfrak{a} runs over a set of ideal class representatives.

The second type of Eisenstein series associated to the imaginary quadratic field $\mathbf{k} = \mathbb{Q}(\sqrt{-q})$, is the 'incoherent' series:

$$(0.7) \quad E_-(\tau, s) = v^{s/2} \sum_{\gamma \in \Gamma_{\infty} \setminus \Gamma} (c\tau + d)^{-1} |c\tau + d|^{-s} \Phi_q^-(\gamma),$$

where, for $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$,

$$(0.8) \quad \Phi_q^-(\gamma) = \begin{cases} \chi_q(a) & \text{if } c \equiv 0 \pmod{q}, \\ -iq^{-1/2} \chi_q(c) & \text{if } c \text{ is prime to } q. \end{cases}$$

In this case, the normalized Eisenstein series

$$(0.9) \quad E_-^*(\tau, s) = q^{(s+1)/2} \Lambda(s+1, \chi_q) E_-(\tau, s)$$

satisfies the functional equation $E_-^*(\tau, -s) = -E_-^*(\tau, s)$, so that $E_-^*(\tau, 0) = 0$.

The modular form of weight 1 of interest to us is then

$$(0.10) \quad \phi(\tau) := -\frac{\partial}{\partial s} \left\{ E_-^*(\tau, s) \right\} \Big|_{s=0} = -h_{\mathbf{k}} \cdot \frac{\partial}{\partial s} \left\{ E_-(\tau, s) \right\} \Big|_{s=0},$$

so that, up to sign, $\phi(\tau)$ is the leading term of the Laurent expansion of $E_-^*(\tau, s)$ at the central point $s = 0$.

Theorem 1. Let $e(\tau) = e^{2\pi i\tau}$. The nonholomorphic modular form ϕ has a Fourier expansion:

$$\phi(\tau) = a_0(\phi, v) + \sum_{n < 0} a_n(\phi, v) e(n\tau) + \sum_{n > 0} a_n(\phi) e(n\tau),$$

where the coefficients are as follows. Let

$$\rho(n) = |\{\mathfrak{a} \subset \mathcal{O}_{\mathfrak{k}} \mid N(\mathfrak{a}) = n\}|,$$

so that

$$\zeta_{\mathfrak{k}}(s) = \sum_{n=1}^{\infty} \rho(n) n^{-s},$$

is the Dedekind zeta function of \mathfrak{k} . The constant term of ϕ is given by

$$a_0(\phi, v) = -h_{\mathfrak{k}} \cdot \left(\log(q) + \log(v) + 2 \frac{\Lambda'(1, \chi_q)}{\Lambda(1, \chi_q)} \right).$$

For $n > 0$,

$$a_n(\phi) = 2 \log(q) (\text{ord}_q(n) + 1) \rho(n) + 2 \sum_{p \neq q} \log(p) (\text{ord}_p(n) + 1) \rho(n/p),$$

where the sum runs over primes p which are inert in \mathfrak{k} . For $n < 0$,

$$a_n(\phi, v) = 2\beta_1(4\pi|n|v) \rho(-n).$$

Here β_1 is essentially the exponential integral:

$$(0.11) \quad \beta_1(t) = -\text{Ei}(-t) = \int_1^{\infty} u^{-1} e^{-ut} du.$$

Since one of the aims of this paper is to illustrate the constructions of [17] in the simplest possible case, we work adelicly, rather than classically, and obtain both the Fourier expansion of $E(\tau, s)$ and its derivative by computing local Whittaker functions and their derivatives. This is done in sections 1–3.

Following the tradition of Hecke, we next compute the Mellin transform of ϕ , or, more precisely, of ϕ with its constant term omitted:

$$(0.12) \quad \Lambda(s, \phi) := \int_0^{\infty} \left(\phi(iv) - a_0(\phi, v) \right) v^{s-1} dv.$$

This can be done termwise. Let $\Lambda(s) = \pi^{-s/2} \Gamma(\frac{s}{2}) \zeta(s)$, and $\Lambda_{\mathfrak{k}}(s) = \Lambda(s) \Lambda(s, \chi_q)$.

Theorem 2.

$$\Lambda(s, \phi) = \Lambda_{\mathbf{k}}(s) \left[\log(q) + \frac{\Lambda'(s, \chi)}{\Lambda(s, \chi)} - \frac{\Lambda'(s)}{\Lambda(s)} \right]$$

What seems striking (to us) here is the occurrence of the *difference* of the logarithmic derivatives of $\Lambda(s, \chi)$ and $\Lambda(s)$. The computation is done in section 4.

Next, we show that the Fourier coefficients of ϕ are connected with certain 0-cycles on an arithmetic curve. This arithmetic curve is defined as the moduli scheme M over $\text{Spec}(\mathcal{O}_{\mathbf{k}})$ of elliptic curves with complex multiplication ι by the ring of integers $\mathcal{O}_{\mathbf{k}}$ in \mathbf{k} . For each positive integer n , there is a 0-cycle $Z(n)$ on M , which is the locus on which the elliptic curves have an additional endomorphism y with $y^2 = -n$, anticommuting with the action of $\mathcal{O}_{\mathbf{k}}$, i.e., with $y \cdot \iota(a) = \iota(\bar{a}) \cdot y$, for $a \in \mathcal{O}_{\mathbf{k}}$.

The following result is proved in section 5 as an application of the classical theory of complex multiplication and its further development due to Gross and Zagier, [11], [14].

Theorem 3. (i) For $n \in \mathbb{Z}_{>0}$, suppose that neither n nor n/p , for any prime p which is inert in \mathbf{k} is a norm of an integral ideal of $\mathcal{O}_{\mathbf{k}}$, i.e., suppose that $\rho(n) = 0$ and $\rho(n/p) = 0$ for all inert primes p . Then the cycle $Z(n)$ is empty and $a_n(\phi) = 0$.

(ii) If $\rho(n) \neq 0$, then $\rho(n/p) = 0$ for all inert p , and the 0-cycle $Z(n)$ is supported in the fiber at q .

(iii) If p is an inert prime and $\rho(n/p) \neq 0$, then $\rho(n) = 0$ and $\rho(n/p') = 0$ for all inert primes $p' \neq p$, and the 0-cycle $Z(n)$ is supported in the fiber at p .

(iv) In all cases,

$$\deg(Z(n)) = a_n(\phi).$$

Here the degree of a 0-cycle $Z = \text{Spec}(R)$ is given by $\deg(Z) = \log |R|$.

We obtain the following Fourier expansion:

$$(0.13) \quad \phi(\tau) = a_0(\phi, v) + \sum_{n>0} \deg(Z(n)) q^n + 2 \sum_{n>0} \rho(n) \beta_1(4\pi n v) q^{-n},$$

in which, for a moment, we write, $q^n = e(n\tau) = e^{2\pi i n \tau}$. As pointed out above, this expression provides some evidence in favor of the general proposals of [17]. In section 6, we propose a ‘modular’ definition of an Arakelov divisor $Z(t, v)$ for $t < 0$ for which

$$(0.14) \quad \deg(Z(t, v)) = a_t(\phi, v)$$

as well.

Finally, J.-B. Bost has observed that the constant term of ϕ can be expressed in terms of the Faltings height of an elliptic curve with complex multiplication by $\mathcal{O}_{\mathbf{k}}$. Let E be an elliptic curve with complex multiplication by $\mathcal{O}_{\mathbf{k}}$ defined over $\bar{\mathbb{Q}}$. Then according to Colmez, [3], p.633, the Faltings height $h_{Fal}(E)$ is given by

$$(0.15) \quad 4h_{Fal}(E) = \log(q) - 2\log(2\pi) - 2\frac{L'(0, \chi)}{L(0, \chi)}.$$

Therefore, after a short manipulation, we can rewrite the constant term as

$$(0.16) \quad a_0(\phi, v) = -h_{\mathbf{k}} \left(\log(v) - 2\log(q) + 4h_{Fal}(E) + C \right),$$

where the constant

$$(0.17) \quad \begin{aligned} C &= \log(\pi) - \frac{\Gamma'(\frac{1}{2})}{\Gamma(\frac{1}{2})} + 2\log(2\pi) \\ &= -\log(\pi) + \gamma + 4\log(2\pi) \end{aligned}$$

is independent of q . It would be nice to obtain this identity in a more conceptual way, consistent with the situation for the positive Fourier coefficients.

The most general incoherent Eisenstein series on $SL(2)$ are described in section 1, from a representation theoretic point of view. In section 6, we suggest the modifications which are necessary to extend Theorems 1 and 3 to the general case.

As far as we know, the modular form ϕ does not appear in the classical literature. It resembles, however, the modular form of weight $3/2$ considered by Zagier [28], whose Fourier expansion is

$$(0.18) \quad \phi_{Zagier}(\tau) = -\frac{1}{12} + \sum_{n>0} \deg(Z(n))q^n + v^{-1/2} \sum_{m \in \mathbb{Z}} \beta_{3/2}(4\pi m^2 v)q^{-m^2},$$

where

$$(0.19) \quad \beta_{3/2}(t) = \frac{1}{16\pi} \int_1^\infty u^{-3/2} e^{-ut} du.$$

Here we have momentarily changed notation and have denoted by $Z(n)$ certain 0-cycles on a modular curve (over \mathbb{C}), cf., for example, [1].

Zagier has observed that the coefficients of ϕ actually occur in [14]. Let $-d$ be a fundamental discriminant with $(d, q) = 1$ and with $d > 4$. Let

$$(0.20) \quad J(-q, -d) = \prod_{\substack{[\tau_1], [\tau_2] \\ disc(\tau_1) = -q \\ disc(\tau_2) = -d}} (j(\tau_1) - j(\tau_2))$$

be the integer of (1.2) of [14]. Then Theorem 1.3 and (1.4) of that paper can be written as

$$(0.21) \quad 2 \log(J(-d, -q)) = \sum_{\substack{n \\ n^2 < dq \\ n^2 \equiv dq(4)}} a_m(\phi),$$

where, in the sum, $m = (dq - n^2)/4$. Thus, (0.21) is the dq th coefficient of the modular form $\theta(\tau) \phi(4\tau)$ of weight $3/2$, where $\theta(\tau) = \sum_{n \in \mathbb{Z}} e(n^2 \tau)$.

Although the present paper was initially intended as an illustration of the results of [17] in the simplest case, the final expression of the Fourier expansion of the function ϕ suggests a reformulation of the results of [17] along the same lines. The basic questions are the following:

- I. Do the nonsingular Fourier coefficients of the central derivative of an incoherent Eisenstein series of weight $\frac{n+1}{2}$ and genus n , as defined in [17], coincide with the arithmetic degrees of certain 0-cycles on a suitable moduli scheme?
- II. Do the singular coefficients have a similar interpretation in terms of arithmetic/geometric invariants of the moduli space?

The results of [17], [19] and [21] can be reformulated as a positive answer to question I. in some cases.

This paper has its origin in a question raised by Gross at the Durham conference in the summer of 1996 when we presented results on the higher dimensional cases. We thank him for his insistence that we consider this ‘simplest’ case and for several helpful suggestions. We are indebted to Zagier for his observation about the relation of our function to his work with Gross and to Bost for his interpretation of the constant term.

Notation. We will use the following notation for elements of $SL(2)$:

$$m(a) = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}, n(b) = \begin{pmatrix} 1 & b \\ & 1 \end{pmatrix}, w = \begin{pmatrix} & 1 \\ -1 & \end{pmatrix}, k_\theta = \begin{pmatrix} \cos(\theta) & \sin(\theta) \\ -\sin(\theta) & \cos(\theta) \end{pmatrix},$$

for $\theta \in \mathbb{R}$.

Let $\mathbb{Q}_\mathbb{A}$ be the adèles of \mathbb{Q} , and let $\psi : \mathbb{Q}_\mathbb{A}/\mathbb{Q} \rightarrow \mathbb{C}^\times$ be the standard nontrivial additive character, as defined in Tate’s thesis.

§1. Incoherent Eisenstein series for $SL(2)$.

The simplest examples of incoherent Eisenstein series, as considered in [17], occurs for the group $SL(2)$. In this section, we sketch their construction.

Let χ be a nontrivial quadratic character of $\mathbb{Q}^\times \backslash \mathbb{Q}_\mathbb{A}^\times$, and let $\mathbf{k} = \mathbb{Q}(\sqrt{D})$ be the associated quadratic extension. We assume that $D < 0$ is the discriminant of the extension \mathbf{k}/\mathbb{Q} , and write $\mathcal{O}_\mathbf{k}$ for the ring of integers of \mathbf{k} .

Let V be a two dimensional vector space over \mathbb{Q} with quadratic form $x \mapsto Q(x)$. Let $\chi_V(x) = (x, -\det V)$ be the quadratic character associated to V , and assume that $\chi_V = \chi$. Then $V \simeq \mathbf{k}$ with quadratic form $Q(x) = \kappa \cdot N(x)$, $\kappa \in \mathbb{Q}^\times$, given by a multiple of the norm form. Up to isomorphism over \mathbb{Q} , the space V is determined by its collection of localizations $\{V_p\}$, for $p \leq \infty$. These localizations V_p are determined by their Hasse invariants. They have the following form, up to isomorphism:

- (i) If $p = \infty$, V_∞ has signature $(2, 0)$ or $(0, 2)$, and these spaces have Hasse invariants $\epsilon_\infty(V_\infty) = 1$ and $\epsilon_\infty(V_\infty) = -1$, respectively.
- (ii) If p is inert or ramified in \mathbf{k}/\mathbb{Q} , then $V_p = \mathbf{k}_p$ with $Q(x) = N(x)$ or $Q(x) = \kappa_p \cdot N(x)$, where $\chi_p(\kappa_p) = -1$, i.e., with $\kappa_p \notin N\mathbf{k}_p^\times$. These spaces have Hasse invariants $\epsilon_p(V_p) = 1$ and $\epsilon_p(V_p) = -1$, respectively.
- (iii) If p splits in \mathbf{k} , $V_p = V_{1,1}$, the split binary quadratic space. In this case, the Hasse invariant is $\epsilon_p(V_p) = 1$.

If $V = \mathbf{k}$ with $Q(x) = \kappa \cdot N(x)$, then $\epsilon_p(V_p) = \chi_p(\kappa) = (\kappa, D)_p$. The Hasse invariants satisfy the product formula $\prod_{p \leq \infty} \epsilon_p(V_p) = 1$.

An *incoherent collection* $\mathcal{C} = \{\mathcal{C}_p\}$ is a collection of such local quadratic spaces with the following property. For some (and hence for any) global binary quadratic space V with $\chi_V = \chi$, there is a finite set S of places, with (i) $|S|$ is odd, (ii) for any finite place $p \in S$, p does not split in \mathbf{k} , and (iii) for all $p \leq \infty$,

$$(1.1) \quad \mathcal{C}_p = \begin{cases} (V_p, \kappa_p Q) & \text{if } p \in S, \\ (V_p, Q) & \text{otherwise.} \end{cases}$$

Here κ_p is as in (ii) above and $\kappa_\infty = -1$. By the definition $\prod_{p \leq \infty} \epsilon_p(\mathcal{C}_p) = -1$, and there is no global binary quadratic space with these localizations.

Let $G = SL(2)$ over \mathbb{Q} , and let $B = TN$ be the upper triangular Borel subgroup. Associated to quadratic character χ is the global induced representation

$$I(s, \chi) = \text{Ind}_{B(\mathbb{A})}^{G(\mathbb{A})}(\chi | \cdot^s),$$

of $G(\mathbb{A})$. Here $\Phi(s) \in I(s, \chi)$ satisfies

$$(1.2) \quad \Phi(n(b)m(a)g, s) = \chi(a)|a|^{s+1}\Phi(g, s),$$

for $b \in \mathbb{A}$ and $a \in \mathbb{A}^\times$. At $s = 0$, we have a decomposition into irreducible representations of $G(\mathbb{A})$:

$$(1.3) \quad I(0, \chi) = \left(\oplus_V \Pi(V) \right) \oplus \left(\oplus_{\mathcal{C}} \Pi(\mathcal{C}) \right).$$

Here V runs over binary quadratic spaces with $\chi_V = \chi$, and \mathcal{C} runs over incoherent collections with $\chi_{\mathcal{C}} = \chi$. Let $S(V(\mathbb{A}))$ be the Schwartz space of $V(\mathbb{A})$. Recall that the irreducible subspace $\Pi(V)$ is the image of the map $\lambda_V : S(V(\mathbb{A})) \rightarrow I(0, \chi)$ defined by $\lambda_V(\varphi)(g) = (\omega(g)\varphi)(0)$, where $\omega = \omega_\psi$ denotes the action of $G(\mathbb{A})$ on $S(V(\mathbb{A}))$ via the global Weil representation determined by the fixed additive character ψ . Analogously, let $S(\mathcal{C}_{\mathbb{A}}) = \otimes_{p \leq \infty} S(\mathcal{C}_p)$ be the Schwartz space of \mathcal{C} (restricted tensor product of the spaces $S(\mathcal{C}_p)$). The group $G(\mathbb{A})$ acts on this space by the restricted product of the local Weil representations, and $\Pi(\mathcal{C})$ is the image of the equivariant map $\lambda_{\mathcal{C}} : S(\mathcal{C}_{\mathbb{A}}) \rightarrow I(0, \chi)$ defined by

$$(1.4) \quad \lambda_{\mathcal{C}}(\otimes_p \varphi_p)(g) = \otimes_p ((\omega_p(g)\varphi_p)(0)),$$

where $\omega_p = \omega_{\psi_p}$ denotes the action of $G(\mathbb{Q}_p)$ on $S(\mathcal{C}_p)$ via the local Weil representation determined by the local component ψ_p of ψ .

For $g \in G(\mathbb{A})$, write $g = n(b)m(a)k$, where $k \in K_\infty K$, $K_\infty = SO(2)$, and $K = SL_2(\hat{\mathbb{Z}})$. Here $a \in \mathbb{A}^\times$ is not uniquely determined, but the absolute value $|a(g)| := |a|_{\mathbb{A}}$ is well defined. For $\Phi \in I(0, \chi)$, let

$$(1.5) \quad \Phi(g, s) = \Phi(g, 0)|a(g)|^s$$

be its standard extension. For such a standard section $\Phi(s)$, the Eisenstein series

$$(1.6) \quad E(g, s, \Phi) = \sum_{\gamma \in B(\mathbb{Q}) \backslash G(\mathbb{Q})} \Phi(\gamma g, s)$$

converges absolutely for $\operatorname{Re}(s) > 1$ and has an entire analytic continuation.

If $\Phi = \Phi(0) \in \Pi(\mathcal{C})$ for some incoherent collection \mathcal{C} , the resulting incoherent Eisenstein series $E(g, s, \Phi)$ vanishes at $s = 0$, [17], Theorem 2.2, and [18], Theorem 3.1. In the sections which follow, we will compute the central derivative $E'(g, 0, \Phi)$ of such a series in the simplest possible case.

§2. The case of prime discriminant.

Fix a prime $q > 3$ with $q \equiv 3 \pmod{4}$, and let $\mathbf{k} = \mathbb{Q}(\sqrt{-q})$. Note that, by genus theory, the class number of \mathbf{k} is then odd. This allows for considerable simplification. Let $R = \mathcal{O}_{\mathbf{k}}$ be the ring of integers of \mathbf{k} . Let χ be the character of $\mathbb{Q}_{\mathbb{A}}^{\times}$ associated to \mathbf{k} , so that, $\chi(x) = (x, -q)_{\mathbb{A}}$, where $(\ , \)_{\mathbb{A}}$ is the global Hilbert symbol. For a prime p , let $\chi_p(x) = (x, -q)_p$, for the local Hilbert symbol $(\ , \)_p$ at p .

Let $V = \mathbf{k}$, viewed as a \mathbb{Q} -vector space with quadratic form $Q(x) = -N(x)$, and let \mathcal{C} be the incoherent collection defined by $\mathcal{C}_p = V_p$, for all finite primes p , and $\text{sig}(\mathcal{C}_{\infty}) = (2, 0)$. We identify \mathcal{C}_{∞} with \mathbf{k}_{∞} with quadratic form $N(x)$.

We will consider the incoherent Eisenstein series associated to the standard factorizable section $\Phi(s) \in I(s, \chi)$ with $\Phi(0) = \lambda(\varphi)$ for $\varphi = \otimes_p \varphi_p \in S(\mathcal{C}_{\mathbb{A}})$ and $\lambda = \lambda_{\mathcal{C}} : S(\mathcal{C}_{\mathbb{A}}) \rightarrow I(0, \chi)$, as above. We choose $\varphi = \otimes_{p \leq \infty} \varphi_p \in S(\mathcal{C}_{\mathbb{A}})$ as follows. For $p < \infty$, let φ_p be the characteristic function of $R_p = R \otimes_{\mathbb{Z}} \mathbb{Z}_p$, and let $\varphi_{\infty}(x) = \exp(-\pi N(x))$.

In the range of absolute convergence, $\text{Re}(s) > 1$, the incoherent Eisenstein series determined by $\Phi(s)$ has a Fourier expansion

$$(2.1) \quad E(g, s, \Phi) = \sum_{t \in \mathbb{Q}} E_t(g, s, \Phi),$$

with

$$(2.2) \quad E_t(g, s, \Phi) = \int_{\mathbb{Q} \setminus \mathbb{A}} E(n(b)g, s, \Phi) \psi(-tb) db.$$

A calculation similar to but simpler than that given at the end of section 7 of [13] yields the expression given in (0.1) and (0.2) for $v^{-\frac{1}{2}} E(g_{\tau}, s, \Phi)$, where, for $\tau = u + iv$,

$$g_{\tau} = \begin{pmatrix} 1 & u \\ & 1 \end{pmatrix} \begin{pmatrix} v^{\frac{1}{2}} & \\ & v^{-\frac{1}{2}} \end{pmatrix} \in G(\mathbb{R}).$$

For $t \neq 0$,

$$(2.3) \quad E_t(g, s, \Phi) = \prod_{p \leq \infty} W_{t,p}(g_p, s, \Phi_p),$$

where

$$(2.4) \quad W_{t,p}(g_p, s, \Phi_p) = \int_{\mathbb{Q}_p} \Phi_p(w_n(b)g_p, s) \psi_p(-tb) db$$

is the local Whittaker integral. Here db is the self dual measure with respect to ψ_p . Similarly,

$$(2.5) \quad E_0(g, s, \Phi) = \Phi(g, s) + M(s)\Phi(g),$$

where the global intertwining operator has a factorization $M(s)\Phi = \otimes_p M_p(s)\Phi_p$, and

$$(2.6) \quad M_p(s)\Phi(g) = W_{0,p}(g, s, \Phi).$$

To obtain an explicit formula for our particular choice of $\varphi \in S(\mathcal{C}_{\mathbb{A}})$ it is necessary, first, to determine the section $\Phi(s) = \otimes_p \Phi_p(s)$ defined by φ and then to compute the integrals $W_{t,p}(g_p, s, \Phi_p)$. For $p \neq q$, these are completely standard matters, and we will spare the reader the details of the computations and just summarize the results. Also, in the end, we will assume that, for $\tau = u + iv$ in the upper half plane, $g = g_\tau = n(u)m(v^{1/2}) \in G(\mathbb{R})$, so that we may as well assume that $g_p = 1$ for all finite places. Since our section $\Phi(s)$ has been fixed, we will omit it from the notation and simply write:

$$(2.7) \quad W_{t,p}^*(s) = L_p(s+1, \chi) W_{t,p}(e, s, \Phi_p),$$

for p a finite prime, and

$$(2.8) \quad W_{t,\infty}^*(\tau, s) = L_\infty(s+1, \chi) W_{t,\infty}(g_\tau, s, \Phi_\infty).$$

Lemma 2.1. (i) For $p \neq q, \infty$, the function φ_p is invariant under $K_p = SL_2(\mathbb{Z}_p)$, and so $\Phi_p(s)$ is the unique K_p -invariant vector in $I_p(s, \chi)$ with $\Phi_p(e, s) = 1$.

(ii) For $p = \infty$, and for $k_\theta \in SO(2) = K_\infty$, $\omega(k_\theta)\varphi_\infty = e^{i\theta} \cdot \varphi_\infty$. Thus, $\Phi_\infty(s)$ is the unique weight 1 eigenvector for K_∞ in $I_\infty(s, \chi_\infty)$ with $\Phi_\infty(e, s) = 1$.

For $p = q$, let $J_q \subset K_q = SL_2(\mathbb{Z}_q)$ be the Iwahori subgroup. The ramified character χ_q , $\chi_q(t) = (t, -q)_q$ defines a character of J_q by

$$(2.9) \quad \chi_q\left(\begin{pmatrix} a & b \\ qc & d \end{pmatrix}\right) = \chi_q(a).$$

Recall that $K_q = J_q \cup (J_q \cap N)wJ_q$. Thus, the subspace of $I_q(s, \chi)$ consisting of χ_q -eigenvectors of J_q is two dimensional, and is spanned by the ‘cell’ functions $\Phi_q^0(s)$ and $\Phi_q^1(s)$, determined by

$$(2.10) \quad \Phi_q^i(w_j, s) = \delta_{ij}, \quad \text{where } w_0 = 1 \text{ and } w_1 = w.$$

By a direct calculation, given in section 3, we find:

Lemma 2.2. For $k = \begin{pmatrix} a & b \\ qc & d \end{pmatrix} \in J_q$, $\omega(k)\varphi_q = \chi_q(a)\varphi_q$, so that $\Phi_q(s)$ lies in the 2 dimensional subspace of $I_q(s, \chi)$ consisting of χ_q eigenvectors of J_q . Explicitly $\Phi_q(s) = \Phi_q^0(s) + c_q \cdot \Phi_q^1(s)$, where $c_q = \sqrt{-1}q^{-1/2}$.

It follows that our section $\Phi(s)$ is a χ -eigenvector under the action of the group $J = J_q \prod_{p \neq q} K_p$, and hence that the Eisenstein series $E(g, s, \Phi)$ is right χ -equivariant under J .

Lemma 2.3. If $g \in G(\mathbb{R})$, then $E_t(g, s, \Phi) = 0$ for $t \notin \mathbb{Z}$.

Proof. For $t \in \mathbb{Q} - \mathbb{Z}$, take $b \in \hat{\mathbb{Z}}$ so that $\psi(tb) \neq 1$. Then, since $n(b) \in J$ with $\chi(n(b)) = 1$, we have

$$(2.11) \quad E_t(g, s, \Phi) = E_t(gn(b), s, \Phi) = E_t(n(b)g, s, \Phi) = \psi(tb) \cdot E_t(g, s, \Phi),$$

and hence $E_t(g, s, \Phi) = 0$. \square

From now on we will assume that t is integral.

Next, we compute the normalized local Whittaker functions $W_{t,p}^*(s)$, $W_{t,\infty}^*(\tau, s)$, and their derivatives.

Lemma 2.4. For a finite prime $p \neq q$, let $X = p^{-s}$. Then

$$W_{t,p}^*(s) = \sum_{r=0}^{\text{ord}_p(t)} (\chi_p(p)X)^r.$$

Here $\chi_p(p) = (p, -q)_p$ is 1 if p is split in \mathbf{k} and -1 if p is inert. If $t = 0$, so that $\text{ord}_p(t) = \infty$,

$$W_{0,p}^*(s) = (1 - \chi(p)X)^{-1},$$

and hence

$$M_p^*(s)\Phi_p^0 = L_p(s, \chi)\Phi_p^0(-s).$$

Here

$$(2.12) \quad M_p^*(s) = L_p(s+1, \chi)M_p(s).$$

Let

$$(2.13) \quad \rho_p(t) = \rho(p^{\text{ord}_p(t)}),$$

so that, for $t > 0$,

$$(2.14) \quad \rho(t) = \prod_p \rho_p(t).$$

Note that $\rho(t) = 0$ for $t < 0$.

The following facts, which follow immediately from Lemma 2.4, will be useful below.

Lemma 2.5. *At $s = 0$,*

$$W_{t,p}^*(0) = \rho_p(t).$$

If $\chi_p(p) = -1$ and $\text{ord}_p(t)$ is odd, i.e., if $\rho_p(t) = 0$, then

$$W_{t,p}^{*,\prime}(0) = \log(p) \frac{1}{2} (\text{ord}_p(t) + 1) \rho_p(t/p).$$

Note that, $\rho_p(t/p) = 1$ here.

Next consider the archimedean factor. In section 3, we indicate the proof of the following result.

Proposition 2.6. *For $\tau = u + iv$ in the upper half plane, and $g_\tau = n(u)m(v^{1/2})$, as above,*

$$W_{t,\infty}^*(\tau, s) = v^{(1-s)/2} e(tu) \frac{2i\pi^{s/2} e^{2\pi tv}}{\Gamma(s/2)} \int_{\substack{u>0 \\ u>2tv}} e^{-2\pi u} u^{s/2} (u - 2tv)^{s/2-1} du.$$

For $t \neq 0$, it extends to an entire function of s .

(i) If $t > 0$, then $W_{t,\infty}^(\tau, 0) = 2i \cdot v^{1/2} e(t\tau)$.*

(ii) If $t = 0$, then

$$W_{0,\infty}^*(\tau, s) = (M_\infty^*(s)\Phi_\infty)(g_\tau) = i \cdot v^{(1-s)/2} \cdot \pi^{-(s+1)/2} \Gamma\left(\frac{s+1}{2}\right),$$

and $W_{0,\infty}^(\tau, 0) = iv^{1/2}$.*

(iii) If $t < 0$, then $W_{t,\infty}^(\tau, 0) = 0$, and*

$$W_{t,\infty}^{*,\prime}(g_\tau, 0, \Phi_\infty^1) = i v^{1/2} e(t\tau) \cdot \beta_1(4\pi|t|v),$$

where β_1 is given by (0.6).

The above formula for $W_{t,\infty}^*(\tau, s)$ can be found in Siegel [25], eq.(14), p. 88.

Finally, in the case $p = q$, and noting that $L_q(s+1, \chi) = 1$, a direct calculation given in section 3 below yields:

Proposition 2.7. For $p = q$, let $c_q = \sqrt{-1}q^{-1/2}$, as in Lemma 2.2 above. Then

$$W_{t,q}^*(s) = (1 - \chi_q(t)q^{-s(\text{ord}_q(t)+1)}) \cdot c_q.$$

If $\chi_q(t) = -1$, then $W_{t,q}^*(0) = 2c_q \cdot \rho_q(t)$. If $\chi_q(t) = 1$, then $W_{t,q}^*(0) = 0$, and

$$W_{t,q}^{*,\prime}(0) = c_q \log(q) (\text{ord}_q(t) + 1) \rho_q(t).$$

Note that the factor $\rho_q(t) = 1$.

Allowing $\text{ord}_q(t)$ to go to infinity in $W_{t,q}^*(s)$, gives $W_{0,q}^*(s) = c_q$ and hence

$$(2.15) \quad M_q^*(s)\Phi_q = c_q \cdot \Phi_q(-s).$$

We can now assemble these facts. For $t \neq 0$, and for $\text{Re}(s) > 1$, we have

$$(2.16) \quad E_t^*(g_\tau, s, \Phi) = q^{(s+1)/2} \cdot W_{t,\infty}^*(\tau, s) \cdot W_{t,q}^*(s) \cdot \prod_{p \neq q} W_{t,p}^*(s),$$

Since the factors in the product for p with $\text{ord}_p(t) = 0$ are equal to 1, the expressions given in Lemmas 2.4, 2.6 and 2.7 provide the entire analytic continuation of the right side of (2.16). At $s = 0$ the factors on the right side vanish as follows:

- (i) $W_{t,\infty}^*(\tau, 0) = 0$, if $t < 0$, i.e., if $\chi_\infty(t) = -1$,
- (ii) $W_{t,q}^*(0) = 0$ if $\chi_q(t) = 1$, and
- (iii) $W_{t,p}^*(0) = 0$ if $\chi_p(t) = -1$ for $p \neq q, \infty$.

Note that case (iii) occurs only when p is inert in \mathbf{k} and $\text{ord}_p(t)$ is odd.

Since

$$(2.17) \quad \chi_\infty(t)\chi_q(t) \prod_{p \neq q} \chi_p(t) = \chi(t) = 1,$$

an odd number of factors in (2.16) are forced to vanish at $s = 0$, via (i)–(iii). Precisely one factor will vanish when one of the quantities $\rho(-t)$, $\rho(t)$ or $\rho(t/p)$ is nonzero. Specifically, this factor will be

- (i) $W_{t,\infty}^*(\tau, 0)$ when $\rho(-t) \neq 0$, i.e., when $\chi_\infty(t) = -1$, $\chi_q(t) = -1$ and $\chi_p(t) = 1$ for $p \neq q, \infty$;
- (ii) $W_{t,q}^*(0)$ when $\rho(t) \neq 0$, i.e., when $\chi_\infty(t) = 1$, $\chi_q(t) = 1$ and $\chi_p(t) = 1$ for $p \neq q, \infty$; and
- (iii) $W_{t,p}^*(0)$ when $\rho(t/p) \neq 0$, i.e., when $\chi_\infty(t) = 1$, $\chi_q(t) = -1$, $\chi_p(t) = -1$ and $\chi_\ell(t) = 1$ for $\ell \neq p, q, \infty$.

These are the only cases which contribute to the derivative at $s = 0$. The corresponding nonzero Fourier coefficients of $E^{*,\prime}(g_\tau, 0, \Phi)$ can then be obtained by combining the values of normalized local Whittaker functions and their derivatives described in the Lemmas above.

Proposition 2.8. (i) If $\rho(-t) \neq 0$, then

$$E_t^{*,\prime}(g_\tau, 0, \Phi) = -2v^{1/2} e(t\tau) \cdot \beta_1(4\pi|t|v) \cdot \rho(-t).$$

(ii) If $\rho(t) \neq 0$, then

$$E_t^{*,\prime}(g_\tau, 0, \Phi) = -2v^{1/2} e(t\tau) \cdot \log(q) (\text{ord}_q(t) + 1) \cdot \rho(t).$$

(iii) If $\rho(t/p) \neq 0$, then

$$E_t^{*,\prime}(g_\tau, 0, \Phi) = -2v^{1/2} e(t\tau) \cdot \log(p) (\text{ord}_p(t) + 1) \cdot \rho(t/p).$$

Since the classical Eisenstein series of the introduction is

$$(2.18) \quad E^*(\tau, s) = v^{-1/2} E^*(g_\tau, s, \Phi),$$

we obtain the expressions of Theorem 1 for the nonconstant Fourier coefficients of ϕ .

Finally, we compute the constant term and its derivative:

$$(2.19) \quad \begin{aligned} E_0^*(g_\tau, s, \Phi) &= q^{(s+1)/2} \Lambda(s+1, \chi_q) \cdot \Phi(g_\tau, s) + q^{(s+1)/2} M^*(s) \Phi(g_\tau) \\ &= q^{(s+1)/2} \Lambda(s+1, \chi_q) \cdot v^{(s+1)/2} + q^{(s+1)/2} \cdot i \cdot c_q \cdot \Lambda(s, \chi_q) \cdot v^{(1-s)/2} \\ &= q^{(s+1)/2} \Lambda(s+1, \chi_q) \cdot v^{(s+1)/2} - q^{(1-s)/2} \Lambda(1-s, \chi_q) \cdot v^{(1-s)/2}. \end{aligned}$$

Thus

$$(2.20) \quad \begin{aligned} E_0^{*,\prime}(g_\tau, 0, \Phi) &= 2 \frac{\partial}{\partial s} \left\{ q^{(s+1)/2} \Lambda(s+1, \chi_q) \cdot v^{(s+1)/2} \right\} \Big|_{s=0} \\ &= v^{1/2} \cdot h_{\mathbf{k}} \cdot \left(\log(q) + \log(v) + 2 \frac{\Lambda'(1, \chi_q)}{\Lambda(1, \chi_q)} \right) \end{aligned}$$

This gives the constant term of ϕ , and finishes the proof of Theorem 1.

§3. Some computations.

In this section, we provide the details for the computations of the values and derivatives of the Whittaker functions used in section 2, in the case $p = q$ or ∞ .

We begin with the case $p = q$.

Proof of Lemma 2.2. We compute the action of $SL_2(\mathbb{Z}_q)$ on $S(V(\mathbb{Q}_q))$ for the Weil representation. Write $J_q = (J_q \cap N_q^-)(J_q \cap M_q)(J_q \cap N_q)$. For $b \in \mathbb{Z}_q$,

$$(3.1) \quad \omega(n(b))\varphi_q(x) = \psi_q(-bN(x))\varphi_q(x) = \varphi_q(x),$$

since $x \in R_q$ implies that $N(x) \in \mathbb{Z}_q$, and ψ_q is unramified. Similarly, for $a \in \mathbb{Z}_q^\times$,

$$(3.2) \quad \omega(m(a))\varphi_q(x) = \chi_q(a)|a|\varphi_q(xa) = \chi_q(a)\varphi_q(x).$$

Finally, note that $n_-(qc) = \begin{pmatrix} 1 & \\ qc & 1 \end{pmatrix} = wn(-qc)w^{-1}$. Then

$$(3.3) \quad \begin{aligned} \omega(w^{-1})\varphi_q(x) &= -i \int_{V_q} \varphi_q(y) \psi_q(-\text{tr}(xy^\sigma)) dy \\ &= -i \int_{R_q} \psi_q(-\text{tr}(xy^\sigma)) dy \\ &= -i \text{meas}(R_q) \hat{\varphi}_q(x), \end{aligned}$$

where $\hat{\varphi}_q$ is the characteristic function of the dual lattice $\hat{R}_q = \frac{1}{\sqrt{-q}}R_q$. Then

$$(3.4) \quad \omega(n_-(qc))\omega(w^{-1})\varphi_q(x) = -i \text{meas}(R_q) \psi_q(qcN(x)) \hat{\varphi}_q(x),$$

and so

$$(3.5) \quad \begin{aligned} \omega(n_-(qc))\varphi_q(x) &= \text{meas}(R_q) \int_{V_q} \psi_q(qcN(y)) \psi_q(\text{tr}(xy^\sigma)) \hat{\varphi}_q(y) dy \\ &= \text{meas}(R_q) \int_{\hat{R}_q} \psi_q(qcN(y)) \psi_q(\text{tr}(xy^\sigma)) dy \\ &= \text{meas}(R_q) \int_{\hat{R}_q} \psi_q(\text{tr}(xy^\sigma)) dy \\ &= \text{meas}(R_q) \text{meas}(\hat{R}_q) \varphi_q(x). \end{aligned}$$

Here $\text{meas}(R_q) \text{meas}(\hat{R}_q) = 1$, since the measure dy is self dual with respect to ψ_q . The second statement is clear, since

$$(3.6) \quad \lambda_q(\varphi_q) = \lambda_q(\varphi_q)(w_0)\Phi_q^0(0) + \lambda_q(\varphi_q)(w_1)\Phi_q^1(0),$$

so that

$$(3.7) \quad \begin{aligned} c_q &= \lambda_q(\varphi_q)(w) \\ &= \omega(w)\varphi_q(0) \\ &= i \int_{V_q} \varphi_q(y) dy \\ &= i \text{meas}(R_q) \\ &= i q^{-1/2}. \end{aligned}$$

□

Proof of Proposition 2.7. Recall that $W_{t,q}^*(s) = W_{t,q}(e, s, \Phi_q)$, where, for $\operatorname{Re}(s) > 1$, the Whittaker integral $W_{t,q}(g, s, \Phi_q)$ is as in (2.4). We will actually determine this function completely. It is easy to check that

$$(3.8) \quad W_{t,q}(n(b)m(a)g, s, \Phi) = \chi_q(a)|a|^{-s+1} \psi_q(tb) W_{a^2t,q}(g, s, \Phi),$$

so that, by Lemma 2.2 and the decomposition

$$(3.9) \quad G(\mathbb{Q}_q) = N_q M_q w_0 J_q \cup N_q M_q w_1 J_q,$$

it will suffice to compute the functions

$$(3.10) \quad W_{ij}(s, t) := W_{t,q}(w_i, s, \Phi_q^j) = \int_{\mathbb{Q}_q} \Phi_q^j(w_n(b)w_i, s) \psi_q(-tb) db,$$

for $i, j \in \{0, 1\}$ and the cell functions Φ_q^j as in (2.10). Let

$$(3.11) \quad W_{ij}(s, t)^{(0)} = \int_{\mathbb{Z}_q} \Phi_q^j(w_n(b)w_i, s) \psi_q(-tb) db,$$

and

$$(3.12) \quad W_{ij}(s, t)^{(1)} = \int_{\mathbb{Q}_q - \mathbb{Z}_q} \Phi_q^j(w_n(b)w_i, s) \psi_q(-tb) db.$$

For $b \in \mathbb{Z}_q$, $w_n(b)w_i \in K_q$, so the integrand in $W_{ij}(s, t)^{(0)}$ is independent of s . Then

$$(3.13) \quad \begin{aligned} W_{0j}(s, t)^{(0)} &= \int_{\mathbb{Z}_q} \Phi_q^j(w_n(b), s) \psi_q(-tb) db \\ &= \Phi_q^j(w, s) \int_{\mathbb{Z}_q} \psi_q(-tb) db \\ &= \delta_{1j} \operatorname{char}(\mathbb{Z}_q)(t), \end{aligned}$$

where $\operatorname{char}(\mathbb{Z}_q)$ is the characteristic function of \mathbb{Z}_q . Next,

$$(3.14) \quad W_{1j}(s, t)^{(0)} = \int_{\mathbb{Z}_q} \Phi_q^j(-w_n(b)w^{-1}, s) \psi_q(-tb) db.$$

Since $w_n(qx)w^{-1} \in J_q$, the function $b \mapsto \Phi_q^j(-w_n(b)w^{-1}, s)$ depends only on b modulo $q\mathbb{Z}_q$, and thus the integral vanishes if $t \notin q^{-1}\mathbb{Z}_q$. If $t \in q^{-1}\mathbb{Z}_q$, then

$$(3.15) \quad W_{1j}(s, t)^{(0)} = q^{-1} \chi_q(-1) \sum_{b \in \mathbb{Z}/q\mathbb{Z}} \Phi_q^j(n_-(-b)) \psi_q(-tb).$$

Note that $n_-(-b) \in J_q$ if and only if $b \in q\mathbb{Z}_q$. If $b \in \mathbb{Z}_q^\times$, then write

$$(3.16) \quad n_-(-b) = n(x)wk = n(x)w \begin{pmatrix} A & B \\ qC & D \end{pmatrix}.$$

This yields

$$(3.17) \quad (0, 1)n_-(-b) = (-b, 1) = (-1, 0)k = (-A, -B),$$

and thus $A = b$ and $B = -1$. Hence

$$(3.18) \quad \Phi_q^j(n_-(-b)) = \chi_q(b) \cdot \delta_{1j},$$

and

$$(3.19) \quad \begin{aligned} W_{1j}(s, t)^{(0)} &= q^{-1}\chi_q(-1) \left(\delta_{0j} + \delta_{1j} \sum_{b \in (\mathbb{Z}/q\mathbb{Z})^\times} \chi_q(b) \psi_q(-tb) \right) \\ &= \begin{cases} q^{-1}\chi_q(-1)\delta_{0j} & \text{if } \text{ord}_q(t) \geq 0, \\ q^{-1}\chi_q(-1) \left(\delta_{0j} + \delta_{1j} \chi_q(t) \mathfrak{g}(\chi_q) \right) & \text{if } \text{ord}_q(t) = -1. \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Here

$$(3.20) \quad \mathfrak{g}(\chi_q) = \sum_{b \in (\mathbb{Z}/q\mathbb{Z})^\times} \chi_q(b) \psi(-b/q)$$

is the Gauss sum for χ_q . We note that $\chi_q(q) = (q, -q)_q = 1$ and $\chi_q(b) = \left(\frac{b}{q}\right)$ for $b \in (\mathbb{Z}/q\mathbb{Z})^\times$, so that $\chi_q(-1) = -1$.

Next consider $W_{ij}(s, t)^{(1)}$. Here, for $\text{ord}_q(b) < 0$,

$$(3.21) \quad wn(b) = \begin{pmatrix} & 1 \\ -1 & -b \end{pmatrix} = \begin{pmatrix} 1 & -b^{-1} \\ & 1 \end{pmatrix} \begin{pmatrix} -b^{-1} & \\ & -b \end{pmatrix} \begin{pmatrix} 1 & \\ b^{-1} & 1 \end{pmatrix}.$$

Thus

$$(3.22) \quad \Phi_q^j(wn(b)w_i, s) = \chi_q(-b)|b|^{-s-1}\delta_{ij},$$

since $n_-(b^{-1})w_i = w_iw_i^{-1}n_-(b^{-1})w_i$ and

$$(3.23) \quad w_i^{-1}n_-(b^{-1})w_i \in (J_q \cap N_-) \cup (J_q \cap N)$$

for either i . Therefore, recalling that $\chi_q(q) = 1$,

$$(3.24) \quad \begin{aligned} W_{ij}(s, t)^{(1)} &= \chi_q(-1) \delta_{ij} \sum_{r=1}^{\infty} q^{-rs} \int_{\mathbb{Z}_q^\times} \chi_q(b) \psi_q(-tb/q^r) db \\ &= \begin{cases} \chi_q(-1) \delta_{ij} q^{-1-s(\text{ord}_q(t)+1)} \chi_q(t) \mathfrak{g}(\chi_q) & \text{if } \text{ord}_q(t) \geq 0, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Here we use the fact that

$$(3.25) \quad \int_{\mathbb{Z}_q^\times} \chi_q(b) \psi_q(-tb/q^r) db = \begin{cases} 0 & \text{if } r \neq \text{ord}_q(t) + 1, \\ q^{-1} \chi_q(t) \mathfrak{g}(\chi_q) & \text{if } r = \text{ord}_q(t) + 1. \end{cases}$$

These results can be summarized as follows: If $\text{ord}_q(t) \geq 0$, then, writing $r = \text{ord}_q(t) + 1$,

$$(3.26) \quad \begin{pmatrix} W_{ij}(s, t) \end{pmatrix} = q^{-1} \chi_q(-1) \begin{pmatrix} q^{-sr} \chi_q(t) \mathfrak{g}(\chi_q) & q \chi_q(-1) \\ 1 & q^{-sr} \chi_q(t) \mathfrak{g}(\chi_q) \end{pmatrix}.$$

If $\text{ord}_q(t) = -1$, then

$$(3.27) \quad \begin{pmatrix} W_{ij}(s, t) \end{pmatrix} = q^{-1} \chi_q(-1) \begin{pmatrix} 0 & 0 \\ 1 & \chi_q(t) \mathfrak{g}(\chi_q) \end{pmatrix}.$$

Otherwise $(W_{ij}(s, t)) = 0$.

Finally, recall that $\Phi_q(s) = \Phi_q^0(s) + c_q \Phi_q^1(s)$, with $c_q = \sqrt{-1} q^{-1/2}$, and that $\mathfrak{g}(\chi_q) = \sqrt{-1} q^{1/2}$. Thus, if $\text{ord}_q(t) \geq 0$,

$$(3.28) \quad \begin{aligned} \begin{pmatrix} W_{t,q}(w_0, s, \Phi_q) \\ W_{t,q}(w_1, s, \Phi_q) \end{pmatrix} &= \begin{pmatrix} W_{ij}(s, t) \end{pmatrix} \cdot \begin{pmatrix} 1 \\ c_q \end{pmatrix} \\ &= (1 - \chi_q(t) q^{-s(\text{ord}_q(t)+1)}) \begin{pmatrix} c_q \\ -q^{-1} \end{pmatrix}. \end{aligned}$$

If $\text{ord}_q(t) = -1$, then

$$(3.29) \quad \begin{pmatrix} W_{t,q}(w_0, s, \Phi_q) \\ W_{t,q}(w_1, s, \Phi_q) \end{pmatrix} = (1 - \chi_q(t)) \begin{pmatrix} 0 \\ -q^{-1} \end{pmatrix}.$$

This completes the proof of Proposition 2.7, and a little more. \square

We next turn to the case $p = \infty$. Of course, these calculations are rather well known, but we include them for convenience.

Proof of Proposition 2.6. Recall that, by (2.8),

$$(3.30) \quad \begin{aligned} W_{t,\infty}^*(\tau, s) &= v^{\frac{1}{2}(1-s)} e(tu) W_{tv,\infty}^*(e, s) \\ &= v^{\frac{1}{2}(1-s)} e(tu) L_\infty(s+1, \chi) W(s, tv), \end{aligned}$$

where

$$(3.31) \quad L_\infty(s+1, \chi) = \pi^{-s/2-1} \Gamma\left(\frac{s}{2} + 1\right),$$

as in (0.4), and where we set

$$(3.32) \quad W(s, t) := W_{t, \infty}(e, s, \Phi_\infty).$$

A simple computation of the Iwasawa decomposition of $wn(b)$ yields

$$(3.33) \quad \Phi_\infty^1(wn(b), s) = i \cdot \frac{(1+ib)}{(1+b^2)^{s/2+1}}.$$

Thus

$$(3.34) \quad \begin{aligned} W(s, t) &= i \cdot \int_{-\infty}^{\infty} (1+ib)^{-s/2} (1-ib)^{-s/2-1} e(-tb) db \\ &= i \frac{\pi^{s/2+1}}{\Gamma(s/2+1)} \int_{-\infty}^{\infty} (1+ib)^{-s/2} \int_0^{\infty} e^{-\pi(1-ib)u} u^{s/2} du e(-tb) db. \end{aligned}$$

Recall that

$$(3.35) \quad \int_0^{\infty} e(-xv) e^{-xz} x^{s-1} dx = (z+2\pi iv)^{-s} \Gamma(s),$$

and hence,

$$(3.36) \quad \int_{-\infty}^{\infty} e(vx) (z+2\pi iv)^{-s} dv = \begin{cases} \frac{1}{\Gamma(s)} \cdot e^{-xz} x^{s-1} & \text{if } x > 0, \\ 0 & \text{if } x \leq 0. \end{cases}$$

Thus

$$(3.37) \quad \begin{aligned} &\int_{-\infty}^{\infty} (1+ib)^{-s/2} e(b(\frac{1}{2}u-t)) db \\ &= \begin{cases} \frac{(2\pi)^{s/2}}{\Gamma(s/2)} \cdot e^{-2\pi(\frac{1}{2}u-t)} (\frac{1}{2}u-t)^{s/2-1} & \text{if } \frac{1}{2}u-t > 0, \\ 0 & \text{if } \frac{1}{2}u-t \leq 0. \end{cases} \end{aligned}$$

This gives

$$(3.38) \quad \begin{aligned} W(s, t) &= i \frac{\pi^{s/2+1}}{\Gamma(s/2+1)} \frac{(2\pi)^{s/2}}{\Gamma(s/2)} \int_{\substack{u>0 \\ u>2t}} e^{-\pi u} u^{s/2} e^{-2\pi(\frac{1}{2}u-t)} (\frac{1}{2}u-t)^{s/2-1} du \\ &= 2i \cdot \frac{\pi^{s+1} e^{2\pi t}}{\Gamma(s/2)\Gamma(s/2+1)} \int_{\substack{u>0 \\ u>2t}} e^{-2\pi u} u^{s/2} (u-2t)^{s/2-1} du. \end{aligned}$$

This yields the first statement of Proposition 2.6.

We are interested in the behavior at $s = 0$. If $t > 0$, the substitution of $u + 2t$ for u yields

$$(3.39) \quad \begin{aligned} W(s, t) &= 2i \cdot \frac{\pi^{s+1} e^{-2\pi t}}{\Gamma(s/2)\Gamma(s/2 + 1)} \int_0^\infty e^{-2\pi u} (u + 2t)^{s/2} u^{s/2-1} du \\ &= 2i \cdot \frac{\pi^{s+1} e^{-2\pi t}}{\Gamma(s/2)\Gamma(s/2 + 1)} \left(\Gamma(s/2)(2\pi)^{-s/2}(2t)^{s/2} + O(s) \right). \end{aligned}$$

Thus, if $t > 0$, $W(0, t) = 2\pi i \cdot e^{-2\pi t}$.

If $t = 0$, we get

$$(3.40) \quad \begin{aligned} W(s, 0) &= 2i \cdot \frac{\pi^{s+1}}{\Gamma(s/2)\Gamma(s/2 + 1)} \int_0^\infty e^{-2\pi u} u^{s-1} du \\ &= 2\pi i \cdot \frac{2^{-s} \Gamma(s)}{\Gamma(s/2)\Gamma(s/2 + 1)} \\ &= i\pi^{1/2} \cdot \frac{\Gamma(\frac{s+1}{2})}{\Gamma(\frac{s+2}{2})}, \end{aligned}$$

via duplication.

Finally, when $t < 0$,

$$(3.41) \quad W(s, t) = 2i \cdot \frac{\pi^{s+1} e^{2\pi t}}{\Gamma(s/2)\Gamma(s/2 + 1)} \int_0^\infty e^{-2\pi u} u^{s/2} (u + 2|t|)^{s/2-1} du.$$

The integral here converges for $\operatorname{Re}(s) > -2$, so that the $\Gamma(s/2)$ in the denominator yields $W(0, t) = 0$ when $t < 0$. Note that

$$(3.42) \quad \begin{aligned} &\int_0^\infty e^{-2\pi u} (u + 2|t|)^{-1} du \\ &= \int_0^\infty e^{-4\pi|t|u} (2|t|u + 2|t|)^{-1} 2|t| du \\ &= \int_0^\infty \frac{e^{-4\pi|t|u}}{u + 1} du \\ &= e^{4\pi|t|} \cdot \int_0^\infty \frac{e^{-4\pi|t|(u+1)}}{u + 1} du \\ &= -e^{4\pi|t|} \cdot \operatorname{Ei}(-4\pi|t|). \end{aligned}$$

Therefore

$$(3.43) \quad \begin{aligned} \frac{\partial}{\partial s} \left\{ W(s, t) \right\} \Big|_{s=0} &= -2\pi i \cdot \frac{1}{2} \cdot e^{2\pi t} \cdot e^{4\pi|t|} \cdot \operatorname{Ei}(-4\pi|t|) \\ &= -i\pi \cdot e^{2\pi|t|} \cdot \operatorname{Ei}(-4\pi|t|), \end{aligned}$$

when $t < 0$. This completes the proof of Proposition 2.6. \square

§4. The Mellin transform.

In this section, we compute the Mellin transform $\Lambda(s, \phi)$.

Lemma 4.1. *The Mellin transform (0.7) can be computed termwise.*

Proof. It suffice to check the termwise absolute convergence of the expression

$$(4.1) \quad - \int_0^\infty \sum_{t=1}^\infty \rho(t) \cdot Ei(-4\pi tv) e^{2\pi tv} v^{s-1} dv.$$

It will turn out to be more convenient to consider the expression

$$(4.2) \quad - \sum_{t=1}^\infty \rho(t) \int_0^\infty Ei(-4\pi tv) e^{2\pi tv} v^{\sigma-1} dv,$$

with $\sigma > 1$. First, by a change of variable in each integral, we obtain

$$(4.3) \quad \begin{aligned} & - \sum_{t=1}^\infty \rho(t) (2\pi t)^{-\sigma} \int_0^\infty Ei(-2v) e^v v^{\sigma-1} dv \\ &= - \left(\sum_{t=1}^\infty \rho(t) (2\pi t)^{-\sigma} \right) \cdot \left(\int_0^\infty Ei(-2v) e^v v^{\sigma-1} dv \right) \\ &= -(2\pi)^{-\sigma} \zeta_{\mathbf{k}}(\sigma) \cdot \int_0^\infty Ei(-2v) e^v v^{\sigma-1} dv. \end{aligned}$$

Thus, it suffices to prove that that the integral is convergent when $\sigma > 1$. Note that the expression

$$(4.4) \quad -Ei(-2v) = \int_1^\infty \frac{e^{-2vu}}{u} du$$

is positive. If $v > \delta > 0$, and if $0 < \epsilon < 1$, we can write:

$$(4.5) \quad \begin{aligned} -Ei(-2v) e^v &< \int_1^\infty \frac{e^{-(1-\epsilon)vu}}{u} du e^{-\epsilon v} \\ &< \int_1^\infty \frac{e^{-(1-\epsilon)\delta u}}{u} du e^{-\epsilon v} \\ &= C_{\delta, \epsilon} \cdot e^{-\epsilon v}. \end{aligned}$$

On the other hand, if $0 < v < \delta$, we have

$$(4.6) \quad \begin{aligned} -Ei(-2v) e^v &< e^\delta \int_1^\infty \frac{e^{-2vu}}{u} du \\ &< e^\delta \int_1^\infty e^{-2vu} du \\ &= e^\delta (2v)^{-1}. \end{aligned}$$

Returning to the integral,

$$(4.7) \quad - \int_0^\infty Ei(-2v) e^v v^{\sigma-1} dv < \int_0^\delta e^\delta (2v)^{-1} v^{\sigma-1} dv + \int_\delta^\infty C_{\delta,\epsilon} \cdot e^{-\epsilon v} v^{\sigma-1} dv.$$

The second integral on the right side converges for all σ , while the first is convergent for $\text{Re}(s) > 1$, as required. \square

Computing (0.7) termwise, the contribution of the holomorphic terms is simply

$$(4.8) \quad 2(2\pi)^{-s} \Gamma(s) \sum_{t=1}^\infty \left[\log(q)(\text{ord}_q(t) + 1) \rho(t) + \sum_{p \neq q, \infty} \log(p)(\text{ord}_p(t) + 1) \rho(t/p) \right] t^{-s},$$

where the sum on p is over primes inert in \mathbf{k} . If p is such a prime, then $\rho(p^r) = 1$ if r is even and 0 if r is odd, and we have

$$(4.9) \quad \begin{aligned} \sum_{t=1}^\infty (\text{ord}_p(t) + 1) \rho(t/p) t^{-s} &= \sum_{\substack{t=1 \\ (t,p)=1}}^\infty \sum_{r=0}^\infty (r+1) \rho(p^{r-1}t) p^{-rs} t^{-s} \\ &= \left(\sum_{\substack{t=1 \\ (t,p)=1}}^\infty \rho(t) t^{-s} \right) \left(\sum_{r=0}^\infty (r+1) \rho(p^{r-1}) p^{-rs} \right) \\ &= (1 - p^{-2s}) \cdot \zeta_{\mathbf{k}}(s) \cdot \left(\sum_{r=0}^\infty (2r+2) p^{-s(2r+1)} \right) \\ &= \frac{2p^{-s}}{1 - p^{-2s}} \cdot \zeta_{\mathbf{k}}(s). \end{aligned}$$

Similarly,

$$(4.10) \quad \sum_{t=1}^\infty (\text{ord}_q(t) + 1) \rho(t) t^{-s} = (1 - q^{-s})^{-1} \cdot \zeta_{\mathbf{k}}(s).$$

Thus, the holomorphic part of the Fourier expansion contributes

$$(4.11) \quad \begin{aligned} \Lambda_{\mathbf{k}}(s) &\left[\frac{1}{1 - q^{-s}} \log(q) + \sum_{p \neq q} \frac{2p^{-s}}{1 - p^{-2s}} \log(p) \right] \\ &= \Lambda_{\mathbf{k}}(s) \left[\log(q) + \frac{L'(s, \chi)}{L(s, \chi)} - \frac{\zeta'(s)}{\zeta(s)} \right], \end{aligned}$$

as one easily checks.

The termwise transform of the nonholomorphic part of the Fourier expansion is

$$\begin{aligned}
(4.12) \quad & 2 \sum_{t=1}^{\infty} \rho(t) \int_0^{\infty} \beta_1(4\pi tv) e^{2\pi tv} v^{s-1} dv \\
& = 2(2\pi)^{-s} \sum_{t=1}^{\infty} \rho(t) t^{-s} \left(\int_0^{\infty} \beta_1(2v) e^v v^{s-1} dv \right) \\
& = 2(2\pi)^{-s} \zeta_{\mathbf{k}}(s) \Gamma(s) \left[\frac{\Gamma(\frac{s+1}{2})'}{\Gamma(\frac{s+1}{2})} - \frac{\Gamma(\frac{s}{2})'}{\Gamma(\frac{s}{2})} \right].
\end{aligned}$$

Thus, the contribution of these terms is

$$(4.13) \quad \Lambda_{\mathbf{k}}(s) \left[\frac{\Gamma(\frac{s+1}{2})'}{\Gamma(\frac{s+1}{2})} - \frac{\Gamma(\frac{s}{2})'}{\Gamma(\frac{s}{2})} \right].$$

Combining this with the contribution of the holomorphic terms, we obtain the expression of Theorem 2 for $\Lambda(s, \phi)$.

In (4.12) we have used the following identity, noting that $\Gamma(\frac{s}{2})' = \frac{1}{2}\Gamma'(\frac{s}{2})$.

Lemma 4.2.

$$\int_0^{\infty} \beta_1(2v) e^v v^{s-1} dv = \frac{1}{2} \Gamma(s) \left[\frac{\Gamma'(\frac{s+1}{2})}{\Gamma(\frac{s+1}{2})} - \frac{\Gamma'(\frac{s}{2})}{\Gamma(\frac{s}{2})} \right].$$

Proof. Recall that, [22],

$$(4.14) \quad \psi(s) := \frac{\Gamma'(s)}{\Gamma(s)} = -\gamma + \int_0^1 \frac{1-x^{s-1}}{1-x} dx.$$

Therefore

$$\begin{aligned}
(4.15) \quad \psi(s+1/2) - \psi(s) & = \int_0^1 \frac{x^{s-1} - x^{s-1/2}}{1-x} dx \\
& = \int_0^1 x^{s-1} \frac{1-x^{1/2}}{1-x} dx \\
& = \int_0^1 \frac{x^{s-1}}{1+x^{1/2}} dx \\
& = 2 \int_0^1 \frac{u^{2s-1}}{1+u} du.
\end{aligned}$$

Thus we have the useful formula:

$$(4.16) \quad 2 \int_0^1 \frac{w^{s-1}}{1+w} dw = \psi\left(\frac{s+1}{2}\right) - \psi\left(\frac{s}{2}\right).$$

But now

$$\begin{aligned}
(4.17) \quad & \int_0^\infty \beta_1(2v) e^v v^{s-1} dv \\
&= \int_0^\infty \int_1^\infty \frac{e^{-2uv}}{u} du e^v v^{s-1} dv \\
&= \Gamma(s) \int_1^\infty (2u-1)^{-s} u^{-1} du \\
&= \Gamma(s) \int_1^\infty r^{-s} (1+r)^{-1} dr \\
&= \Gamma(s) \int_0^1 \frac{w^{s-1}}{1+w} dw.
\end{aligned}$$

This gives the claimed expression. \square

§5. Algebraic-geometric aspects.

In the beginning of this section we will consider a slightly more general situation than before. We let \mathbf{k} be an imaginary quadratic field with ring of integers $\mathcal{O}_{\mathbf{k}}$. We denote by $a \mapsto \bar{a}$ the non trivial automorphism of \mathbf{k} .

We will consider the following moduli problem \mathcal{M} over $(\text{Sch}/\mathcal{O}_{\mathbf{k}})$. To $S \in (\text{Sch}/\mathcal{O}_{\mathbf{k}})$ we associate the category $\mathcal{M}(S)$ of pairs (E, ι) where

- E is an elliptic curve over S (i.e. an abelian scheme of relative dimension one over S)
- $\iota : \mathcal{O}_{\mathbf{k}} \rightarrow \text{End}_S(E)$ is a homomorphism such that the induced homomorphism

$$(5.1) \quad \text{Lie}(\iota) : \mathcal{O}_{\mathbf{k}} \rightarrow \text{End}_{\mathcal{O}_S}(\text{Lie } E) = \mathcal{O}_S$$

coincides with the structure homomorphism.

The morphisms in this category are the isomorphisms. A pair (E, ι) will be called an *elliptic curve with CM by $\mathcal{O}_{\mathbf{k}}$* . We denote by M the corresponding set valued functor of isomorphism classes of objects of \mathcal{M} .

Proposition 5.1. *a) The moduli problem \mathcal{M} is representable by an algebraic stack (in the sense of Deligne-Mumford) which is finite and étale over $\text{Spec } \mathcal{O}_{\mathbf{k}}$.*

b) The functor M has a coarse moduli scheme which is also finite and étale over $\text{Spec } \mathcal{O}_{\mathbf{k}}$.

Proof. a) Let $\tilde{\mathcal{M}}$ be the algebraic moduli stack of elliptic curves over $\text{Spec } \mathcal{O}_{\mathbf{k}}$. The forgetful morphism $i : \mathcal{M} \rightarrow \tilde{\mathcal{M}}$ is relatively representable by an unramified

morphism (rigidity theorem). Since the generic elliptic curve in any characteristic has only trivial (i.e., \mathbb{Z}) endomorphisms, \mathcal{M} has finite fibres over $\text{Spec } \mathcal{O}_{\mathfrak{k}}$. Now the finiteness of \mathcal{M} over $\text{Spec } \mathcal{O}_{\mathfrak{k}}$ follows from the valuative criterion of properness, since an elliptic curve with complex multiplication has potentially good reduction. The étaleness of \mathcal{M} over $\text{Spec } \mathcal{O}_{\mathfrak{k}}$ will follow by the Serre-Tate theorem from a study of the p -divisible group of an elliptic curve with CM by $\mathcal{O}_{\mathfrak{k}}$.

b) That M has a coarse moduli scheme follows from a) (or use level- n -structures). Let ξ be a geometric point of \mathcal{M} and $\bar{\xi}$ the corresponding geometric point of M . Let $\hat{\mathcal{O}}_{\mathcal{M},\xi}$ and $\hat{\mathcal{O}}_{M,\bar{\xi}}$ be the completions of the local rings. If ξ corresponds to (E, ι) , we have

$$(5.2) \quad \text{Aut}(\xi) = \text{Aut}(E, \iota) = \mathcal{O}_{\mathfrak{k}}^{\times} \quad ,$$

comp. cases 1 and 2 below. This group acts on $\hat{\mathcal{O}}_{\mathcal{M},\xi}$ and

$$(5.3) \quad \hat{\mathcal{O}}_{M,\bar{\xi}} = (\hat{\mathcal{O}}_{\mathcal{M},\xi})^{\text{Aut}(\xi)} \quad .$$

However, since \mathcal{M} is étale over $\text{Spec } \mathcal{O}_{\mathfrak{k}}$, we have that $\hat{\mathcal{O}}_{\mathcal{M},\xi}$ is equal to the strict completion of $\mathcal{O}_{\mathfrak{k}}$ in the prime ideal corresponding to the image of ξ in $\text{Spec } \mathcal{O}_{\mathfrak{k}}$. Hence the action of $\text{Aut}(\xi)$ on $\hat{\mathcal{O}}_{\mathcal{M},\xi}$ is trivial. Therefore the canonical morphism

$$(5.4) \quad \mathcal{M} \rightarrow M$$

is étale and the last assertion follows. \square

Corollary 5.2. *Let ξ be a geometric point of \mathcal{M} and let $\bar{\xi}$ be the corresponding point of M . Then*

$$\hat{\mathcal{O}}_{M,\bar{\xi}} = \hat{\mathcal{O}}_{\mathcal{M},\xi}.$$

Let p be a prime number and \wp a prime ideal of $\mathcal{O}_{\mathfrak{k}}$ above p . Let κ be an algebraically closed extension of the residue field $\kappa(\wp)$ of \wp . Let $(E, \iota) \in \mathcal{M}(\kappa)$, and let X be the p -divisible group of E . Then the p -adic completion $\mathcal{O}_p = \mathcal{O}_{\mathfrak{k}} \otimes \mathbb{Z}_p$ acts on X . We distinguish two cases.

Case 1: p splits in $\mathcal{O}_{\mathfrak{k}}$. Then $p = \wp \cdot \bar{\wp}$ with $\wp \neq \bar{\wp}$ and

$$(5.5) \quad \mathcal{O}_p = \mathcal{O}_{\wp} \oplus \mathcal{O}_{\bar{\wp}} = \mathbb{Z}_p \oplus \mathbb{Z}_p \quad .$$

By the condition on $\text{Lie}(\iota)$ the corresponding decomposition of X is of the form

$$(5.6) \quad X = \hat{\mathbb{G}}_m \times \mathbb{Q}_p/\mathbb{Z}_p \quad .$$

In this case it is obvious that (X, ι) deforms uniquely, i.e. the base of the universal deformation is $\mathrm{Spf} W(\kappa)$, where $W(\kappa)$ denotes the ring of Witt vectors of κ . In this case the elliptic curve E is ordinary and

$$(5.7) \quad \iota : \mathcal{O}_{\mathbf{k}} \xrightarrow{\sim} \mathrm{End}(E) \quad .$$

Case 2: p not split in $\mathcal{O}_{\mathbf{k}}$. In this case $\mathcal{O}_p = \mathcal{O}_{\wp}$ and (X, ι) is a formal \mathcal{O}_{\wp} -module of height 2 over κ , in the sense of Drinfeld [8], comp. also [11]. Let $W_{\mathcal{O}}(\kappa)$ be the ring of relative Witt vectors of κ , [2]. Then again (X, ι) deforms uniquely, i.e. the base of the universal deformation of (X, ι) is $\mathrm{Spf} W_{\mathcal{O}}(\kappa)$ ([23], comp. [11], 2.1).

In this case the elliptic curve E is supersingular. Let \mathbb{B}_p be the definite quaternion algebra over \mathbb{Q} which ramifies at p . Then $\mathrm{End}(E)$ is isomorphic to a maximal order in \mathbb{B}_p .

The main theorem of complex multiplication may be summarized as follows (comp. [24],[4])

Proposition 5.3. *Let $\bar{\mathbf{k}}$ be an algebraic closure of \mathbf{k} . The Galois group $\mathrm{Gal}(\bar{\mathbf{k}}/\mathbf{k})$ acts on $M(\bar{\mathbf{k}})$ through its maximal abelian quotient, which we identify via class field theory with $(\mathbf{k} \otimes \mathbb{A}_f)^{\times} / \mathbf{k}^{\times}$. There is a bijection*

$$M(\bar{\mathbf{k}}) = \mathbf{k}^{\times} \backslash (\mathbf{k} \otimes \mathbb{A}_f)^{\times} / (\mathcal{O}_{\mathbf{k}} \otimes \hat{\mathbb{Z}})^{\times},$$

such that an element of $\mathrm{Gal}(\bar{\mathbf{k}}/\mathbf{k})^{\mathrm{ab}}$ represented by an idèle $x \in (\mathbf{k} \otimes \mathbb{A}_f)^{\times}$ acts by translation by x on the right side.

Corollary 5.4. *Let H be the Hilbert class field of \mathbf{k} and let \mathcal{O}_H be its ring of integers. Then*

$$M \simeq \mathrm{Spec}(\mathcal{O}_H).$$

Corollary 5.5. *Fix p and a prime \wp above p . Then $\mathcal{M}(\overline{\kappa(\wp)})$ forms a single isogeny class. There is a bijection*

$$M(\overline{\kappa(\wp)}) = \mathbf{k}^{\times} \backslash (\mathbf{k} \otimes \mathbb{A}_f)^{\times} / (\mathcal{O}_{\mathbf{k}} \otimes \hat{\mathbb{Z}})^{\times}$$

The action by the Frobenius automorphism over $\kappa(\wp)$ on the left corresponds to the translation by the idèle with component a uniformizer at \wp and 1 at all other finite places on the right.

Remark 5.6. The bijection in Corollary 5.5 is given explicitly as follows. Choose a base point $(E_o, \iota_o) \in \mathcal{M}(\overline{\kappa(\wp)})$, and an identification $\mathrm{End}^0(E_o) = \mathbb{B}_p$. Let

$\mathcal{O}(E_o) \subset \mathbb{B}_p$ be the maximal order corresponding to $\text{End}(E_o)$. To $g \in (\mathbf{k} \otimes \mathbb{A}_f)^\times$ there is associated the point $(E, \iota) \in \mathcal{M}(\overline{\kappa(\wp)})$ whose Tate module $\hat{T}(E) = \prod_\ell T_\ell(E)$ (with p -component the covariant Dieudonné module) is the lattice

$$(5.8) \quad g \cdot \hat{T}(E_o) \subset \hat{T}(E_o) \otimes \mathbb{Q} .$$

In particular

$$(5.9) \quad \mathcal{O}(E) = \mathbb{B}_p \cap g(\mathcal{O}(E_o) \otimes \hat{\mathbb{Z}})g^{-1} ,$$

is a maximal order in \mathbb{B}_p .

We now turn to the definition of special cycles on the moduli space/stack \mathcal{M} .

Definition 5.7. *Let $(E, \iota) \in \mathcal{M}(S)$. A special endomorphism of (E, ι) is an element $y \in \text{End}(E)$ with*

$$(5.10) \quad y \circ \iota(a) = \iota(\bar{a}) \circ y, \quad \text{for all } a \in \mathcal{O}_{\mathbf{k}} .$$

Let $V(E, \iota)$ be the set of special endomorphisms of (E, ι) . Then $V(E, \iota)$ is an $\mathcal{O}_{\mathbf{k}}$ -module.

Let us first consider the case where $S = \text{Spec } \kappa$ with κ an algebraically closed field. If $\text{char } \kappa = 0$, then $\text{End}(E) = \mathcal{O}_{\mathbf{k}}$ and hence $V(E, \iota) = (0)$. The same is true if $\text{char } \kappa = p$ and p splits in $\mathcal{O}_{\mathbf{k}}$, cf. (5.7). Let now $\text{char } \kappa = p$ where p does not split. Then $\text{End}(E)$ can be identified with a maximal order $\mathcal{O}(E)$ in the quaternion algebra \mathbb{B}_p introduced earlier and the $\mathcal{O}_{\mathbf{k}}$ -module $V(E, \iota)$ is of rank one. Let $a \mapsto a'$ be the main involution of \mathbb{B}_p . Then $V(E, \iota)$ is stable under this involution. Let $y \in V(E, \iota)$. Since

$$(5.11) \quad y + y' \in \mathbb{Z} \cap V(E, \iota) = (0) ,$$

we obtain $y^2 = -y \cdot y' \in \mathbb{Z}$. We define a \mathbb{Z} -valued quadratic form Q on the \mathbb{Z} module $V(E, \iota)$ of rank 2 by

$$(5.12) \quad Q(y) = -y^2 = y \cdot y' .$$

The definition of this quadratic form extends to an elliptic curve with CM by $\mathcal{O}_{\mathbf{k}}$ over any *connected* scheme S .

Definition 5.8. For a positive integer t , let $\mathcal{Z}(t)$ be the moduli problem over \mathcal{M} which to $S \in (\text{Sch}/\mathcal{O}_{\mathbf{k}})$ associates the category of triples (E, ι, y) where $(E, \iota) \in \mathcal{M}(S)$ and where $y \in V(E, \iota)$ satisfies

$$(5.13) \quad Q(j) = -y^2 = t.$$

We denote by

$$(5.14) \quad \text{pr} : \mathcal{Z}(t) \longrightarrow \mathcal{M}$$

the forgetful morphism. It is obvious that pr is relatively representable, hence $\mathcal{Z}(t)$ is representable by an algebraic stack. By the rigidity theorem the morphism pr is finite and unramified. We also denote by pr the induced morphism between coarse moduli schemes

$$(5.15) \quad \text{pr} : Z(t) \longrightarrow M.$$

We note that for a geometric point $\tilde{\xi} \in \mathcal{Z}(t)$ we have $\text{Aut}(\tilde{\xi}) = \{\pm 1\}$. Indeed, if $\tilde{\xi}$ corresponds to (E, ι, y) then an automorphism is given by a unit $u \in \mathcal{O}_{\mathbf{k}}^{\times}$ with $uy = yu$, i.e. with $u = \bar{u}$.

Proposition 5.9. *The fibre of $Z(t)$ in characteristic 0 or in characteristic p where p splits in \mathbf{k} is empty. In particular, $Z(t)$ is an artinian scheme. For p not split in \mathbf{k} , let \wp be the prime ideal of $\mathcal{O}_{\mathbf{k}}$ dividing p . The fibre of pr over a geometric point $(E, \iota) \in M(\overline{\kappa(\wp)})$ is empty if t is not represented by the quadratic form on $V(E, \iota)$.*

Let p be inert or ramified in \mathbf{k} . We choose a base point $(E_o, \iota_o) \in M(\overline{\kappa(\wp)})$ and denote by $V^{(p)}$ the quadratic space $V(E_o, \iota_o) \otimes_{\mathbb{Z}} \mathbb{Q}$ over \mathbb{Q} . Then

$$(5.16) \quad \text{End}^0(E_o, \iota_o) = \mathbb{B}_p = \mathbf{k} \oplus V^{(p)},$$

where the decomposition is orthogonal with respect to the quadratic form given by the reduced norm on \mathbb{B}_p . Choosing a set of coset representative g_i , $i = 1, \dots, h_{\mathbf{k}}$ for the cosets on the right side of Corollary 5.5 and applying the construction of Remark 5.6, we obtain $\mathcal{O}_{\mathbf{k}}$ -lattices

$$(5.17) \quad V(E_i, \iota_i) = V^{(p)} \cap \mathcal{O}(E_i)$$

in $V^{(p)}$.

Corollary 5.10. *Let p be inert or ramified in \mathbf{k} and $\wp \subset \mathcal{O}_{\mathbf{k}}$ over p . Then*

$$|Z(t)(\overline{\kappa(\wp)})| = \sum_i |\{x \in V(E_i, \iota_i) \mid Q(x) = t\}|.$$

We now quote the result of Gross [11], Prop. 3.3 and 4.3. Let p be an inert or ramified prime and κ an algebraically closed extension of $\kappa(\wp)$. If $\xi \in \mathcal{M}(\kappa)$, there is an isomorphism

$$(5.18) \quad \hat{\mathcal{O}}_{\mathcal{M}, \xi} = W_{\mathcal{O}}(\kappa) \quad ,$$

cf. case 2 above. Here again $W_{\mathcal{O}}(\kappa) = W_{\mathcal{O}_{\wp}}(\kappa)$ is the ring of relative Witt vectors. We let π denote a uniformizer of \mathcal{O}_{\wp} and hence of $W_{\mathcal{O}}(\kappa)$. Let $\tilde{\xi} \in \mathcal{Z}(t)(\kappa)$ with $\text{pr}(\tilde{\xi}) = \xi$, and $\bar{\xi} \in Z(t)(\kappa)$ its image in $Z(t)$. Here pr is as in (5.14).

Theorem 5.11. (Gross): *We have*

$$\hat{\mathcal{O}}_{Z(t), \bar{\xi}} = \hat{\mathcal{O}}_{\mathcal{Z}(t), \tilde{\xi}} = W_{\mathcal{O}}(\kappa)/(\pi^{\nu})$$

where

$$\nu = \nu_p(t) = \frac{\text{ord}_p(t) + d_p - 1}{f_p} + 1,$$

$$d_p = \text{ord}_p(\text{disc}_{\mathbf{k}/\mathbb{Q}}),$$

and

$$f_p = [\kappa(\wp) : \mathbb{F}_p].$$

In particular, if p is unramified in \mathbf{k} , we have $f_p = 2$, and $d_p = 0$ and $\text{ord}_p(t) \equiv 1 \pmod{2}$, and

$$\nu_p(t) = \frac{\text{ord}_p(t) + 1}{2}.$$

If $p \neq 2$ is ramified in \mathbf{k} , then $f_p = 1$, $d_p \geq 1$ and

$$\nu_p(t) = \text{ord}_p(t) + 1.$$

Note that the length of $\hat{\mathcal{O}}_{\mathcal{Z}(t), \tilde{\xi}}$ only depends on $\text{ord}_p(t)$. Combining Corollary 5.10 and Theorem 5.11, we therefore obtain an expression for the *degree* of $Z(t)$. Here we define the degree as in [26], [10],

$$(5.19) \quad \deg(Z(t)) = \sum_{\xi \in Z(t)} \log(\#(\mathcal{O}_{Z(t), \xi})).$$

Theorem 5.12.

$$\deg(Z(t)) = \sum_p f_p \log(p) \nu_p(t) \cdot \sum_i |\{x \in V(E_i, \iota_i) \mid Q(x) = t\}|.$$

Here p ranges over the primes which are not split in \mathbf{k} .

For a fixed p which is not split in \mathbf{k} , we now describe the lattices $V(E_i, \iota_i)$ of (5.17) more explicitly. The basic technique is from [14], [7].

Let Δ be the discriminant of the field \mathbf{k} . Fix an auxillary prime $p_0 \nmid 2p\Delta$ as follows. If p is inert in \mathbf{k} , we require that

$$(5.20) \quad (\Delta, -pp_0)_v = \begin{cases} -1 & \text{if } v = p \text{ or } \infty, \text{ and} \\ 1 & \text{otherwise.} \end{cases}$$

If p is ramified in \mathbf{k} , we require that

$$(5.21) \quad (\Delta, -p_0)_v = \begin{cases} -1 & \text{if } v = p \text{ or } \infty, \text{ and} \\ 1 & \text{otherwise.} \end{cases}$$

Let

$$(5.22) \quad \kappa_p = \begin{cases} pp_0 & \text{if } p \text{ is inert in } \mathbf{k}, \text{ and} \\ p_0 & \text{if } p \text{ is ramified in } \mathbf{k}. \end{cases}$$

Then \mathbb{B}_p is the cyclic algebra $(\Delta, -\kappa_p)$ and thus can be written in the form

$$(5.23) \quad \mathbb{B}_p = \mathbf{k} \oplus \mathbf{k} \cdot y$$

for an element y with $\text{tr}(y) = 0$, $y^2 = -\kappa_p$, and $a \cdot y = y \cdot \bar{a}$ for all $a \in \mathbf{k}$.

Let $R = \mathcal{O}_{\mathbf{k}} \oplus \mathcal{O}_{\mathbf{k}} \cdot y$, so that R is an order in \mathbb{B}_p with $R \cap \mathbf{k} = \mathcal{O}_{\mathbf{k}}$. Suppose that \mathcal{O} is a maximal order of \mathbb{B}_p containing R . If $x_1 \in R$ and $x_2 \in \mathcal{O}$, then $\text{tr}(x_1 x_2) \in \mathbb{Z}$. Using the coordinates from (5.23), this gives

$$(5.24) \quad \text{tr}((\alpha_1 + \beta_2 y)(\alpha_2 + \beta_2 y)) = \text{tr}_{\mathbf{k}}(\alpha_1 \alpha_2 - \kappa_p \beta_1 \bar{\beta}_2) \in \mathbb{Z}.$$

Therefore

$$(5.25) \quad R = \mathcal{O}_{\mathbf{k}} \oplus \mathcal{O}_{\mathbf{k}} \cdot y \subset \mathcal{O} \subset \mathcal{D}^{-1} \oplus \kappa_p^{-1} \mathcal{D}^{-1} \cdot y,$$

where $\mathcal{D}^{-1} = \sqrt{\Delta}^{-1} \mathcal{O}_{\mathbf{k}}$ is the inverse different of \mathbf{k} . Setting

$$(5.26) \quad V(\mathcal{O}) = V^{(p)} \cap \mathcal{O},$$

we conclude that

$$(5.27) \quad \mathcal{O}_{\mathbf{k}} \cdot y \subset V(\mathcal{O}) \subset \kappa_p^{-1} \mathcal{D}^{-1} \cdot y.$$

The conditions imposed on p_0 imply that p_0 splits. Let $p_0 \mathcal{O}_{\mathbf{k}} = \wp_0 \bar{\wp}_0$ and let $p \mathcal{O}_{\mathbf{k}} = \wp^{e_p}$. Then, it is easily checked that

$$(5.28) \quad R' := \mathcal{O}_{\mathbf{k}} \oplus \wp_0^{-1} y \quad \text{and} \quad R'' := \mathcal{O}_{\mathbf{k}} \oplus \bar{\wp}_0^{-1} y$$

are orders in \mathbb{B}_p containing R . Note that R' and R'' cannot be contained in the same order \mathcal{O} . Indeed, if $R' + R'' \subset \mathcal{O}$, then $(\wp_0^{-1} + \bar{\wp}_0^{-1}) \cdot y = p_0^{-1} \mathcal{O}_{\mathbf{k}} \cdot y \subset \mathcal{O}$. But $Q(p_0^{-1} y) = -(p_0^{-1} y)^2 = p_0^{-2} \kappa_p \notin \mathbb{Z}$.

Suppose that \mathcal{O} is a maximal order containing $\mathcal{O}_{\mathbf{k}} \oplus \wp_0^{-1} y$. If $x = \kappa_p^{-1} \sqrt{\Delta}^{-1} \beta y \in V(\mathcal{O})$, with $\beta \in \mathcal{O}_{\mathbf{k}}$, then

$$(5.29) \quad Q(x) = -x^2 = \kappa_p^{-1} \Delta^{-1} N(\beta) \in \mathbb{Z}.$$

This forces

$$(5.30) \quad \beta \in \wp_0^{-1} \kappa_p \sqrt{\Delta} \mathcal{O}_{\mathbf{k}}, \quad \text{or} \quad \beta \in \bar{\wp}_0^{-1} \kappa_p \sqrt{\Delta} \mathcal{O}_{\mathbf{k}},$$

so that

$$(5.31) \quad x \in \wp_0^{-1} y \quad \text{or} \quad x \in \bar{\wp}_0^{-1} y.$$

Therefore,

$$(5.32) \quad V(\mathcal{O}) = \wp_0^{-1} \cdot y.$$

Fix a maximal order \mathcal{O} containing R' , and note that $\mathcal{O} \cap \mathbf{k} = \mathcal{O}_{\mathbf{k}}$, i.e., that $\mathcal{O}_{\mathbf{k}}$ is optimally embedded in \mathcal{O} , in Eichler's terminology, [9]. By the Chevalley-Hasse-Noether Theorem (comp. [8], Satz 7), it follows that if \mathcal{O}' is any maximal order in \mathbb{B}_p in which $\mathcal{O}_{\mathbf{k}}$ is optimally embedded, then there exists a finite idèle $g \in \mathbf{k}_{\mathbb{A}_f}^\times$, such that

$$(5.33) \quad \mathcal{O}' = g(\mathcal{O} \otimes_{\mathbb{Z}} \hat{\mathbb{Z}}) g^{-1} \cap \mathbb{B}_p.$$

If

$$(5.34) \quad \mathfrak{a} = (g \mathcal{O}_{\mathbf{k}} \otimes_{\mathbb{Z}} \hat{\mathbb{Z}}) \cap \mathbf{k},$$

then

$$(5.35) \quad V(\mathcal{O}') = \mathfrak{a}\bar{\mathfrak{a}}^{-1}\wp_0^{-1}y.$$

Conversely, any ideal \mathfrak{a} arises in this way.

Now suppose that $\mathcal{O}' = \text{End}(E)$ for a supersingular elliptic curve $(E, \iota) \in \mathcal{M}(\overline{\kappa_p(\wp)})$ over $\bar{\mathbb{F}}_p$ with complex multiplication by $\mathcal{O}_{\mathbf{k}}$. Note that $\mathcal{O}_{\mathbf{k}}$ is optimally embedded in \mathcal{O}' and that

$$(5.36) \quad V(E, \iota) = \mathcal{O}' \cap \mathbf{k}y = V(\mathcal{O}').$$

Proposition 5.13. *Fix an auxillary prime p_0 as above. Then there is an element y_0 in V with $Q(y_0) = \kappa_p$, and the lattices $V(E_i, \iota_i)$ in Theorem 5.12 have the form*

$$V(E_i, \iota_i) \simeq \mathfrak{a}_i \bar{\mathfrak{a}}_i^{-1} \wp_0^{-1} \cdot y_0,$$

as the \mathfrak{a}_i run over representatives for the ideal classes of \mathbf{k} . In particular, the isomorphism class of $V(E_i, \iota_i)$ runs over the genus $[[\wp_0]]$ of the ideal \wp_0 .

Remark. The genus $[[\wp_0]]$ is independent of the choice of \wp_0 and is characterized by the values of the genus characters. Recall that a basis for the characters of the group of genera are given as follows, [15]. Let q_i , $1 \leq i \leq t$ be the primes dividing Δ . Let

$$(5.37) \quad \chi_i(\mathfrak{a}) = (\Delta, N(\mathfrak{a}))_{q_i},$$

where $(\ , \)_{q_i}$ is the Hilbert symbol for \mathbb{Q}_{q_i} . Note that, if $x \in \mathbf{k}^\times$, then $(\Delta, N(x))_{q_i} = 1$, and that

$$(5.38) \quad (\Delta, N(\mathfrak{a})N(\mathfrak{b})^2)_{q_i} = (\Delta, N(\mathfrak{a}))_{q_i}.$$

Thus χ_i depends only on the ideal class of \mathfrak{b} modulo squares, i.e., defines a character of the group of genera $Cl(\mathbf{k})/2Cl(\mathbf{k})$. Any $t-1$ of the χ_i 's give a basis for the characters of this group. The conditions (5.20) and (5.21) on the auxillary prime p_0 imply that

$$(5.39) \quad \chi_i(\wp_0) = (\Delta, p_0)_{q_i} = \begin{cases} (\Delta, -p)_{q_i} & \text{if } p \text{ is inert,} \\ (\Delta, -1)_{q_i} & \text{if } p \text{ is ramified and } q_i \neq p, \text{ and} \\ -(\Delta, -1)_{q_i} & \text{if } p \text{ is ramified and } q_i = p. \end{cases}$$

The genus of \wp_0 is determined by these conditions.

Corollary 5.14. (i) The rational quadratic space $V^{(p)}$ is given by

$$(V^{(p)}, Q) \simeq (\mathbf{k}, \kappa_p N),$$

where κ_p is given by (5.22).

(ii) Let C_i be the ideal class

$$C_i = [\mathfrak{a}_i^{-1} \bar{\mathfrak{a}}_i \wp_0] = [\mathfrak{a}_i]^2 [\wp_0].$$

Then

$$|\{x \in V(E_i, \iota_i) \mid Q(x) = t\}| = |\mathcal{O}_{\mathbf{k}}^\times| \cdot |\{\mathfrak{c} \subset \mathcal{O}_{\mathbf{k}} \mid [\mathfrak{c}] = C_i, \text{ and } N(\mathfrak{c}) = p_0 t / \kappa_p\}|.$$

Note that

$$(5.40) \quad p_0 / \kappa_p = \begin{cases} p & \text{if } p \text{ is inert, and} \\ 1 & \text{if } p \text{ is ramified.} \end{cases}$$

Proof. We just note that, if $x = \beta \cdot y \in V(E_i, \iota_i)$, with $\beta \in \mathfrak{a}_i \bar{\mathfrak{a}}_i^{-1} \wp_0^{-1}$, then $t = Q(x) = \kappa_p N(\beta)$. The ideal $\mathfrak{c} = \beta \mathfrak{a}_i^{-1} \bar{\mathfrak{a}}_i \wp_0$ is integral, lies in the ideal class C_i , and has

$$(5.41) \quad N(\mathfrak{c}) = N(\beta) p_0 = t p_0 / \kappa_p.$$

□

Note that, by genus theory, as i runs from 1 to $h_{\mathbf{k}}$, $[\mathfrak{a}_i]$ runs over the ideal class group and C_i runs over $(2Cl(\mathbf{k}))[\wp_0] = [[\wp_0]]$, the genus of \wp_0 , with each class in $[[\wp_0]]$ occurring 2^{t-1} times. Here again t is the number of rational primes which ramify in \mathbf{k} .

Finally, we specialize to the case of prime discriminant considered in earlier sections of this paper, i.e., let $\mathbf{k} = \mathbb{Q}(\sqrt{-q})$ with $q > 3$ a prime congruent to 3 modulo 4. In this case, the class number of \mathbf{k} is odd, there is only one genus, and the C_i 's run over all ideal classes. Combining the results above, we obtain:

Theorem 5.15.

$$\deg(Z(t)) = 2 \log(q) \cdot (\text{ord}_q(t) + 1) \cdot \rho(t) + 2 \sum_p \log(p) \cdot (\text{ord}_p(t) + 1) \cdot \rho(t/p).$$

Here p runs over the primes which are inert in \mathbf{k} .

This is precisely the expression given in Theorem 3 of the introduction.

§6. Variants.

In this section, we will sketch a few variations and extensions of the results of the previous sections. However, none of them touch on the most tantalizing problem of establishing a *direct connection* between the Fourier coefficients and the degrees of special cycles.

The central derivatives of *all* of the incoherent Eisenstein series constructed in section 1 should have an arithmetic interpretation analogous to Theorem 3. A full description of such a result would require several extensions of what we have done above, which is specialized in several respects.

First of all, we have restricted to the case of a prime discriminant $\Delta = -q$. This eliminates complications involving genus theory, as $h_{\mathbf{k}}$ is odd. Of course, the moduli problem of section 5 has been set up to allow elliptic curves with complex multiplication by $\mathcal{O}_{\mathbf{k}}$ where $\mathbf{k} = \mathbb{Q}(\sqrt{\Delta})$ for an arbitrary fundamental discriminant $\Delta < 0$. On the analytic side, the machinery of section 1 and 2 provides corresponding incoherent Eisenstein series, whose Fourier expansions can be computed by the methods of section 2. It should be then be possible to work out the analogues of Theorems 1, 2 and 3 in this case, except that the group of genera will now play a nontrivial role. In the case of nonfundamental discriminant there is more work to be done on both the analytic and the geometric sides.

Second, in section 2, we have considered only those Eisenstein series built from sections which are defined by the characteristic functions of the completions of $\mathcal{O}_{\mathbf{k}}$. More general sections could be considered at the cost of (i) more elaborate calculations of Whittaker functions and their derivatives used in computing the Fourier expansion of the central derivative, and (ii) incorporation of a level structure in the moduli problem.

Third, in section 2, we have restricted to the simplest type of incoherent collection \mathcal{C} , i.e., a collection \mathcal{C} which is obtained by considering the global quadratic space $(V, Q) = (\mathbf{k}, -N)$ and switching the signature from $(0, 2)$ to $(2, 0)$. Thus, at each finite place, $\mathcal{C}_p = (\mathbf{k}_p, -N)$. The most general collection \mathcal{C} is obtained by switching the signature of the space $(V, Q) = (\mathbf{k}, \kappa N)$ where $\kappa \in \mathbb{Q}^\times$. In this situation, a slightly more elaborate moduli problem is necessary. We give a brief sketch of this.

Let B be a indefinite quaternion algebra over \mathbb{Q} , with a fixed embedding $i : \mathbf{k} \hookrightarrow B$. Fix a maximal order \mathcal{O}_B , in B , a positive involution $*$ of B stabilizing \mathcal{O}_B , and assume that $i(\mathcal{O}_{\mathbf{k}}) \subset \mathcal{O}_B$.

We consider polarized abelian surfaces A with an action $\iota : \mathcal{O}_B \otimes \mathcal{O}_{\mathbf{k}} \rightarrow \text{End}(A)$ compatible with the positive involution $b \otimes a \mapsto (b \otimes a)^* = b^* \otimes \bar{a}$. For such an (A, ι) , let

$$(6.1) \quad V(A, \iota) = \{y \in \text{End}(A) \mid y\iota(b \otimes a) = \iota(b \otimes \bar{a})y\}$$

be the space of special endomorphisms. Observe that

$$(6.2) \quad M_2(\mathbf{k}) \simeq B \otimes \mathbf{k} \hookrightarrow \text{End}^0(A),$$

so that A is isogenous to a product A_0^2 for an elliptic curve A_0 with complex multiplication. Thus

$$(6.3) \quad \text{End}^0(A) \simeq M_2(\text{End}^0(A_0)).$$

If A_0 is *not* a supersingular elliptic curve in characteristic p , for p not split in \mathbf{k} , then $\text{End}^0(A) \simeq M_2(\mathbf{k})$ and $V^0(A, \iota) = V(A, \iota) \otimes_{\mathbb{Z}} \mathbb{Q} = 0$. If A_0 is supersingular in characteristic p , then

$$(6.4) \quad \text{End}^0(A) \simeq M_2(\mathbb{B}_p),$$

where $\mathbb{B}_p = \text{End}^0(A_0)$ is, as before, the definite quaternion algebra ramified at infinity and p . The commutator of $\iota(B \otimes 1)$ in $\text{End}^0(A)$ is then isomorphic to B' , the definite quaternion algebra over \mathbb{Q} with

$$(6.5) \quad \text{inv}_\ell(B') = \begin{cases} \text{inv}_\ell(B) & \text{if } \ell \neq p, \text{ and} \\ -\text{inv}_\ell(B') & \text{if } \ell = p. \end{cases}$$

Note that ι yields an embedding of \mathbf{k} into B' , and we have a decomposition, analogous to (5.16),

$$(6.6) \quad B' = \mathbf{k} \oplus V^0(A, \iota),$$

where $V^0(A, \iota) = \mathbf{k} \cdot y$ for a nonzero element $y \in B'$ with $ya = \bar{a}y$ for all $a \in \mathbf{k}$. The quadratic form on $V^0(A, \iota)$ is then

$$(6.7) \quad Q(\beta y) = -(\beta y)^2 = -N(\beta)y^2,$$

so that the binary quadratic space $(V^0(A, \iota), Q)$ is isomorphic to $(\mathbf{k}, \kappa N)$ where $\kappa = -y^2$. Note that B' is then isomorphic to the cyclic algebra $(\Delta, -\kappa)$. An analysis of the corresponding integral moduli problem like that of section 5 should

be possible and give the desired generalization of the results above. The case $B = M_2(\mathbb{Q})$ will reduce precisely to the situation of section 5.

Returning to Theorem 3 in the case $\mathbf{k} = \mathbb{Q}(\sqrt{-q})$, from the point of view of Arakelov theory, it is possible to give an interpretation of the negative Fourier coefficients of ϕ as degrees of cycles.

First recall, [26], that an Arakelov divisor for $M = \text{Spec}(\mathcal{O}_H)$ is an expression of the form

$$(6.8) \quad D = \sum_{\lambda}^{h_{\mathbf{k}}} r_{\lambda} \lambda + \sum_{\mathfrak{p}} n_{\mathfrak{p}} \mathfrak{p},$$

where \mathfrak{p} runs over the nonzero prime ideals of \mathcal{O}_H , λ runs over the complex places of H , the $n_{\mathfrak{p}}$'s are integers (almost all zero), and the r_{λ} 's are real numbers. Let $\text{Div}_c(M)$ be the group of Arakelov divisors, and let $\text{Pic}_c(M)$ be its quotient by the group of principal Arakelov divisors, [26].

We can view the cycles $Z(t)$ defined in section 2 as elements of $\text{Div}_c(M)$ as follows. Fix the embedding of \mathbf{k} into \mathbb{C} , where $\sqrt{-q}$ has positive imaginary part, and let $\omega = (1 + \sqrt{-q})/2$ so that $\mathcal{O}_{\mathbf{k}} = \mathbb{Z} + \mathbb{Z}\omega$. Let

$$(6.9) \quad j_0 = j(\omega) = j(\mathcal{O}_{\mathbf{k}}) \in \mathcal{O}_H$$

be the corresponding singular value of the j -function.

Let p be a rational prime which does not split in \mathbf{k} and let $p\mathcal{O}_{\mathbf{k}} = \wp^{e_p}$. Then \wp is principal and hence splits completely in H . For any prime \mathfrak{p} of \mathcal{O}_H above \wp , the image of j_0 under the reduction map $\mathcal{O}_H \rightarrow \mathcal{O}_H/\mathfrak{p}$ is the j -invariant of an elliptic curve $E_{\mathfrak{p}}$ over $\overline{\mathbb{F}}_p$, unique up to isomorphism, with complex multiplication by $\mathcal{O}_{\mathbf{k}}$. For each such p , this gives a bijective map from the primes over \wp to the fiber $M(\overline{\mathbb{F}}_p)$. Then, for $t > 0$,

$$(6.10) \quad Z(t) = \sum_p \sum_{\mathfrak{p}|\wp} n_{\mathfrak{p}}(t) \mathfrak{p},$$

where

$$(6.11) \quad n_{\mathfrak{p}}(t) = \nu_p(t) \cdot |\{x \in V(E_{\mathfrak{p}}, \iota) \mid Q(x) = t\}|.$$

Here $V(E_{\mathfrak{p}}, \iota)$ is the lattice of special endomorphisms of $E_{\mathfrak{p}}$, as in Definition 5.7, and $\nu_p(t)$ is the integer given in Theorem 5.11. Let

$$(6.11) \quad C_{\mathfrak{p}} = [V(E_{\mathfrak{p}}, \iota)]^{-1},$$

where $[V(E_p, \iota)] \in \text{Pic}(\mathcal{O}_{\mathbf{k}}) \simeq \text{Cl}(\mathbf{k})$ is the class of the rank one $\mathcal{O}_{\mathbf{k}}$ -module $[V(E_p, \iota)]$. Using Corollary 5.14, (6.10) becomes

$$(6.12) \quad n_p = \nu_p(t) \cdot |\{\mathfrak{c} \subset \mathcal{O}_{\mathbf{k}} \mid [\mathfrak{c}] = C_p \text{ and } N(\mathfrak{c}) = t/\kappa_p\}|,$$

where

$$(6.13) \quad \kappa_p = \begin{cases} p & \text{if } p \text{ is inert in } \mathbf{k}, \text{ and} \\ 1 & \text{if } p = q. \end{cases}$$

We view $Z(t)$ as an element of $\text{Div}_c(M)$ with zero archimedean component.

To define an element $Z(t) \in \text{Div}_c(M)$ when $t < 0$, we use a construction based on an idea explained to us by Gross. For a point (E, ι) of $\mathcal{M}(\mathbb{C})$, let E^{top} be the underlying real torus. By analogy with Definition 5.7, we define the space of special endomorphisms

$$(6.14) \quad V(E, \iota) = \{y \in \text{End}(E^{\text{top}}) \mid y\iota(a) = \iota(\bar{a})y\}.$$

If we write $E = \mathbb{C}/L$ for a lattice L with a chosen basis, then

$$(6.15) \quad \text{End}(E^{\text{top}}) \simeq M_2(\mathbb{Z}),$$

and so, by analogy with the case of a supersingular elliptic curve in characteristic p , $\text{End}(E^{\text{top}})$ is isomorphic to a maximal order in the quaternion algebra $M_2(\mathbb{Q})$. It is not difficult to check that $V(E, \iota)$ is a rank one $\mathcal{O}_{\mathbf{k}}$ -module with a \mathbb{Z} -valued quadratic form given by $Q(y) = -y^2$.

For each embedding λ of $H = \mathbf{k}(j)$ into \mathbb{C} , let E_λ be the elliptic curve over \mathbb{C} with j -invariant $\lambda(j_0)$. This curve has complex multiplication by $\mathcal{O}_{\mathbf{k}}$ and space of special endomorphism, in the sense just defined, $V(E_\lambda, \iota)$. Let $C_\lambda = [V(E_\lambda, \iota)]^{-1} \in \text{Pic}(\mathcal{O}_{\mathbf{k}})$. Then, for $t \in \mathbb{Z}_{<0}$ and for $\tau = u + iv$ in the upper half plane, we associate the Arakelov divisor

$$(6.16) \quad Z(t) = Z(t, v) := \sum_{\lambda} r_{\lambda} \lambda,$$

where

$$(6.17) \quad \begin{aligned} r_{\lambda}(t) &= r_{\lambda}(t, v) = 2\beta_1(4\pi|t|v) \cdot |\{y \in V(E_{\lambda}, \iota) \mid Q(y) = t\}| \\ &= 2\beta_1(4\pi|t|v) \cdot |\{\mathfrak{c} \subset \mathcal{O}_{\mathbf{k}} \mid [\mathfrak{c}] = C_{\lambda} \text{ and } N(\mathfrak{c}) = -t\}|. \end{aligned}$$

Here β_1 is the exponential integral (0.6). Thus we have defined for all $t \neq 0$ an Arakelov divisor with

$$(6.18) \quad \deg(Z(t)) = \begin{cases} a_t(\phi) & \text{if } t > 0, \text{ and} \\ a_t(\phi, v) & \text{if } t < 0, \end{cases}$$

and hence

$$(6.19) \quad \phi(\tau) = a_0(\phi, v) + \sum_{t \neq 0} \deg(Z(t)) q^t,$$

extending (0.8).

References.

- [1] R. E. Borcherds, *The Gross-Kohnen-Zagier theorem in higher dimensions*, preprint (1997).
- [2] J.-F. Boutot and H. Carayol, *Uniformisation p -adique des courbes de Shimura: les théorèmes de Cerednik et de Drinfeld*, in: *Courbes modulaires et courbes de Shimura*, Astérisque **196–197**, 1991, pp. 45–158.
- [3] P. Colmez, *Périodes des variétés abéliennes à multiplication complexe*, *Annals of Math.* **138** (1993), 625–683.
- [4] P. Deligne, *Travaux de Shimura*. Sem. Bourbaki 389, Springer Lecture Notes in Math. **244**, 1971.
- [5] P. Deligne and D. Mumford, *The irreducibility of the space of curves of given genus*, *Pub. Math. IHES* **36** (1969), 75–109.
- [6] P. Deligne and M. Rapoport, *Les schémas de modules des courbes elliptiques*, *Modular Functions of One Variable II*, Springer Lecture Notes in Math. **349**, 1973, pp. 143–316.
- [7] D. Dorman, *Special values of the elliptic modular function and factorization formulae*, *J. Reine Angew. Math.* **383** (1988), 207–220.
- [8] V. G. Drinfeld, *Coverings of p -adic symmetric regions*, *Funct. Anal. Appl.* **10** (1977), 29–40.
- [9] M. Eichler, *Zur Zahlentheorie der Quaternionen-Algebren*, *J. Reine Angew. Math.* **195** (1955), 127–151.
- [10] H. Gillet, *Introduction to higher dimensional Arakelov theory*, *Contemporary Math.: Current trends in arithmetic algebraic geometry*, vol. 67, AMS, Providence, 1987.

- [11] B. Gross, *On canonical and quasi-canonical liftings*, *Inventiones math.* **84** (1986), 321–326.
- [12] B. Gross and K. Keating, *On the intersection of modular correspondences*, *Invent. math.* **112** (1993), 225–245.
- [13] B. Gross and S. Kudla, *Heights and the central critical values of triple product L -functions*, *Compositio Math.* **81** (1992), 143–209.
- [14] B. Gross and D. Zagier, *On singular moduli*, *J. Reine Angew. Math.* **355** (1985), 191–220.
- [15] E. Hecke, *Vorlesungen über die Theorie der algebraischen Zahlen*, Akad. Verlagsges., Leipzig, 1923.
- [16] E. Hecke, *Mathematische Werke*, Vandenhoeck & Ruprecht, Göttingen, 1983.
- [17] S. Kudla, *Central derivatives of Eisenstein series and height pairings*, *Annals of Math.* **146** (1997), 545–646.
- [18] S. Kudla and S. Rallis, *A regularized Siegel-Weil formula, the first term identity*, *Annals of Math.* **140** (1994), 1–80.
- [19] S. Kudla and M. Rapoport, *Cycles on Siegel 3-folds and derivatives of Eisenstein series*, preprint (1997).
- [20] S. Kudla and M. Rapoport, *Height pairings on Shimura curves and p -adic uniformization*, preprint (1998).
- [21] S. Kudla and M. Rapoport, *Arithmetic Hirzebruch Zagier cycles*, in preparation.
- [22] N. N. Lebedev, *Special Functions and Their Applications*, Dover, New York, 1972.
- [23] J. Lubin and J. Tate, *Formal complex multiplication in local fields*, *Annals of Math.* **81** (1965), 380–387.
- [24] J.-P. Serre, *Complex multiplication*, in: *Algebraic Number Theory*, J.W.S. Cassels and A. Fröhlich, eds., Academic Press, London, 1967, pp. 292–296.
- [25] C. L. Siegel, *Indefinite quadratische Formen und Modulfunktionen*, *Ges. Abh.*, III, Springer Verlag, Berlin, 1966, pp. 85–91.
- [26] L. Szpiro, *Degrés, intersections, hauteurs*, in: *Séminaire sur les pinceaux arithmétiques: La conjecture de Mordell*, *Astérisque* **127** (1985), 11–28.

- [27] Tonghai Yang, *An explicit formula for local densities of quadratic forms*, to appear in J. Number Theory.
- [28] D. Zagier, *Nombres de classes et formes modulaires de poids 3/2*, C.R. Acad. Sc. Paris **281** (1975), 883–886.

S. Kudla
Department of Mathematics
University of Maryland
College Park MD 20742
USA

M. Rapoport
Mathematisches Institut
der Universität zu Köln
Weyertal 86-90
50931 Köln
Germany

T. Yang
Department of Mathematics
SUNY at Stony Brook
Stony Brook, NY 11794
USA