

# THIN EQUIVALENCE RELATIONS AND INNER MODELS

PHILIPP SCHLICHT

ABSTRACT. We describe the inner models with representatives in all equivalence classes of thin equivalence relations in a given projective pointclass of even level assuming projective determinacy. These models are characterized by their correctness and the property that they correctly compute the tree from the appropriate scale. The main lemma shows that the tree from a scale can be reconstructed in a generic extension of an iterate of a mouse.

We construct models with this property as generic extensions of iterates of mice if the corresponding projective ordinal is below  $\omega_2$ .

On the way we consider several related problems, including the question when forcing does not add equivalence classes to thin projective equivalence relations. For example, we show that if every set has a sharp, then reasonable forcing does not add equivalence classes to thin provably  $\Delta_3^1$  equivalence relations, and generalize this to all projective levels.

## 1. INTRODUCTION

Definable equivalence relations are a focus of modern descriptive set theory. While the bulk of research centers around Borel equivalence relations, there has been a large amount of work on projective equivalence relations, for example Harrington and Sami [8], Hjorth [10, 12, 13], Hjorth and Kechris [14], Kechris [19], Louveau and Rosendal [24], and Silver [39], and equivalence relations in the constructible universe  $L(\mathbb{R})$  over the reals, e.g. Hjorth [11]. Inner model theory has proved to be very useful for this endeavour, in particular iterable models with Woodin cardinals (see e.g. Hjorth [12]). For the theory of iterable mice with Woodin

---

2000 *Mathematics Subject Classification.* 03E15, 03E35, 03E45, 03E55, 03E60.

*Key words and phrases.* projective equivalence relations, thin equivalence relations, inner models, projective ordinals.

cardinals see Mitchell and Steel [29] and Steel [40, 42]. Earlier approaches as in Harrington and Sami [8] and Kechris [19] use direct proofs from determinacy. It is well known that determinacy and the existence of the appropriate mice are equivalent (see [23, 27, 31, 34]).

This paper is about thin projective equivalence relations, i.e. those with no perfect set of pairwise inequivalent reals. They have been most notably studied by Harrington and Sami [8], motivated by the question about the number of equivalence classes. A starting point in this topic is Silver's theorem [39] that every thin  $\mathbf{\Pi}_1^1$  equivalence relation has countably many equivalence classes. Harrington and Sami subsequently extended this through the projective hierarchy relative to the projective ordinals. The  $n^{\text{th}}$  projective ordinal  $\delta_n^1$  is the supremum of lengths of  $\mathbf{\Delta}_n^1$  prewellorders. The number of equivalence classes of thin  $\mathbf{\Pi}_n^1$  equivalence relations is below  $\delta_n^1$  if  $n \geq 1$  is odd and at most the size of  $\delta_{n-1}^1$  if  $n \geq 2$  is even [8].

A quite different approach to this question is to look for a bound for the number of equivalence classes of *co- $\kappa$ -Suslin* equivalence relations, i.e. when the complement is the projection of a tree  $T$  on  $\omega \times \omega \times \kappa$ . Harrington and Shelah [9] showed this is at most  $\kappa$  if the complement of  $p[T]$  is an equivalence relation in a Cohen generic extension. We use this to bound the number of equivalence classes of thin  $\mathbf{\Pi}_n^1$  equivalence relations under the assumption that the pointclasses  $\mathbf{\Pi}_k^1$  are scaled for odd  $k$  and all projective sets have the Baire property. Note that scales are closely connected to Suslin representations.

Since the number of equivalence classes of thin  $\mathbf{\Pi}_n^1$  equivalence relations is bounded by a projective ordinal, we look for inner models (possibly with fewer reals than  $V$ ) which have representatives in all equivalence classes of all thin  $\mathbf{\Pi}_n^1(x)$  equivalence relations, where  $x$  is a real parameter in the inner model. Hjorth [10] showed that every inner model has this property for  $n = 1$  as a consequence of Silver's

theorem. The candidates for such inner models for  $n \geq 2$  are forcing extensions of fine structural inner models with Woodin cardinals. We will construct such models if the corresponding projective ordinal is below  $\omega_2$ .

Hjorth [10] characterized the models with this property for  $n = 2$ . Assuming all reals have sharps, the inner models with this property for  $n = 2$  are exactly the  $\Sigma_3^1$  correct inner models with the right  $\omega_1$ . We extend Hjorth's theorem to the even levels in the projective hierarchy in the main theorem. The level of correctness is adapted and instead of asking that the model has the right  $\omega_1$ , we ask that the model correctly computes the tree  $T_{2n+1}$  from the canonical  $\Pi_{2n+1}^1$ -scale, and assume the appropriate amount of determinacy. Thus these inner models are characterized in a simple and beautiful way.

The proof generalizes Hjorth's proof. In the harder direction, the main lemma shows that the tree  $T_{2n+1}^M$  from the canonical  $\Pi_{2n+1}^1$ -scale as computed in an inner model  $M$  with countably many reals can be reconstructed in an iterate of  $M_{2n}^\#$ . To do this, Woodin's genericity iteration is applied to make reals generic at local Woodin cardinals over iterates of  $M_{2n}^\#$ , and we force over the direct limit. A density argument will show that the tree can be defined. We also have to look more closely at the Harrington-Shelah result [9]. If the equivalence relation is co- $\kappa$ -Suslin, then for any real there is an infinitary formula defining a neighborhood inside its equivalence class. Combining this with Steel's result that  $M_n^\#$  is coded by a projective real, we can express the existence of a real in this neighborhood in a projective way and use this to complete the proof.

Let's look at the setting from a different perspective and suppose the universe is a forcing extension of an inner model. The issue is when a forcing introduces new equivalence classes to thin projective equivalence relations. Foreman and Magidor [4] showed that reasonable forcing of size at most  $\kappa$  does not add equivalence classes to thin  $\kappa$ -weakly homogeneously Suslin equivalence relations. Together

with the Martin-Steel theorem [28] this implies that if there are infinitely many Woodin cardinals, then reasonable forcing does not add equivalence classes to thin projective equivalence relations. We replace the large cardinal assumption with the existence of  $M_n^\#(X)$  for all  $X \in H_{\kappa^+}$  and show that reasonable forcing of size at most  $\kappa$  does not add new equivalence classes to thin provably  $\Delta_{n+2}^1$  equivalence relations. It is essential that in this situation  $M_n^\#(X)$  is absolute.

The paper is organized as follows. *Section 1* introduces the facts about thin equivalence relations, prewellorders, scales, and properties of  $M_n^\#$  which we will use.

In *section 2* we study liftings of thin projective equivalence relations to forcing extensions. We show based on an idea of Foreman and Magidor [4] that for any infinite cardinal  $\kappa$ , reasonable forcing of size at most  $\kappa$  does not introduce new equivalence classes to thin projective equivalence relations if  $M_n^\#(X)$  exists for every self-wellordered set  $X \in H_{\kappa^+}$  and every  $n$ . The argument is adapted to  $\Sigma_2^1$  c.c.c. forcings. We show that generic  $\Sigma_{n+3}^1$  absoluteness holds for these forcings from the assumption that  $M_n^\#(x)$  exists for every real  $x$ , generalizing the result for Cohen and random forcing [45], and that no new equivalence classes of thin projective equivalence relations are introduced. We further prove a lemma about absoluteness of prewellorders under Cohen forcing which we need later.

In *section 3* we present a proof of the Harrington-Shelah theorem [9] for counting the number of equivalence classes of thin co- $\kappa$ -Suslin equivalence relations, and use this to calculate the number of equivalence classes of projective equivalence relations. We further show that thin  $\Sigma_{2n}^1$  equivalence relations are  $\Pi_{2n}^1$  in any real coding  $M_{2n-1}^\#$  for  $n \geq 1$ , generalizing Hjorth's result [12] for  $n = 1$ .

In *section 4* the main theorem is proved. The main lemma shows that the tree from the  $\Pi_{2n+1}^1$  scale can be defined in an iterate of  $M_{2n}^\#$ . We then characterize the inner models which have a representative in every equivalence class of every thin

$\Pi_{2n}^1(x)$  equivalence relation defined from a real parameter  $x$  in the inner model.

We further build a transitive model with this property assuming  $\delta_{2n}^1 < \omega_2$ .

The material in this paper formed part of my dissertation. I would like to thank Ralf Schindler for many discussions and ideas related to the topic of this paper. I am further deeply indebted to Greg Hjorth for his suggestions for proving the main theorem.

## 2. THE FRAMEWORK

This section presents standard definitions and facts which are used later. We work in the theory  $\text{ZF} + \text{DC}$ . For background information we refer to Jech [16], Kanamori [17], Kechris [21], and Moschovakis [30].

**2.1. Prewellorders and scales.** In this section we discuss basic facts about prewellorders and scales.

*2.1.1. Basic definitions and facts.*  $\mathbb{R}$  as well as  ${}^\omega\omega$  denotes Baire space, the set of sequences of natural numbers with the standard topology. The elements of  $\mathbb{R}$  are called reals. A perfect set is a nonempty closed set of reals without isolated points. Clearly every perfect set has the size of the continuum.

**Definition 2.1.** *An equivalence relation  $E \subseteq \mathbb{R} \times \mathbb{R}$  is called thin if there is no perfect set of pairwise inequivalent reals.*

The corresponding notion is also defined for prewellorders. Recall that a prewellorder is a wellfounded linear preorder.

**Definition 2.2.** *A prewellorder  $\leq$  is called thin if there is no perfect set  $P \subseteq \mathbb{R}$  such that  $x < y$  or  $y < x$  for any  $x, y \in P$  with  $x \neq y$ .*

We will work with the projective pointclasses. By a pointclass we mean:

**Definition 2.3.** A (lightface) pointclass  $\Gamma$  is a set  $\emptyset \neq \Gamma \subsetneq \mathcal{P}(\mathbb{R})$  which is closed under recursive preimages and finite intersections and unions. The dual of  $\Gamma$  is defined as  $\check{\Gamma} = \{A \subseteq \mathbb{R} : \mathbb{R} - A\}$ . If  $\Gamma$  is a pointclass we write  $\Delta := \Gamma \cap \check{\Gamma}$ .

Of course for any pointclass  $\Gamma$  we have a corresponding pointclass of subsets of  $\mathbb{R}^n$  via a recursive bijection  $\mathbb{R} \rightarrow \mathbb{R}^n$ .

**Definition 2.4.** If  $\Gamma$  is a pointclass, then  $\ulcorner \Gamma$  is defined as the pointclass of all preimages of sets in  $\Gamma$  under continuous functions. A boldface pointclass is a pointclass with  $\Gamma = \ulcorner \Gamma$ .

**Definition 2.5.** If  $\Gamma$  is a pointclass, then  $< \omega - \Gamma$  denotes the pointclass of boolean combinations of sets in  $\Gamma$ , i.e. sets which are formed from sets in  $\Gamma$  by finite applications of union and complement.

Some of the relevant structural properties of pointclasses are given by norms and scales.

**Definition 2.6.** Suppose  $\Gamma$  is a pointclass and  $A \in \Gamma$ . A prewellorder  $\leq$  with domain  $\mathbb{R}$  is called a  $\Gamma$ -norm on  $A$  if  $x \leq y$  and  $y \in A$  imply  $x \in A$ , and  $\leq$  is uniformly  $\Delta$  in initial segments, i.e. there is a  $\Delta$  set  $B \subseteq \mathbb{R}^2$  with

$$\{(x, y) \in \mathbb{R}^2 : x \leq y \wedge y \in A\} = B \cap \{(x, y) \in \mathbb{R}^2 : y \in A\}.$$

Let  $\equiv$  be the equivalence relation induced by  $\leq$  and let

$$\text{rank}(x) := \text{otp}(\{y : y < x\} / \equiv)$$

for  $x \in A$  and

$$\text{rank}(x) := \infty$$

for  $x \notin A$ .

**Definition 2.7.** Suppose  $\Gamma$  is a pointclass and  $A \in \Gamma$ . A sequence  $(\leq_n : n < \omega)$  of  $\Gamma$ -norms on  $A$  with

$$\{(x, y, n) \in \mathbb{R} \times \mathbb{R} \times \omega : x \leq_n y\} \in \Gamma$$

is called a  $\Gamma$ -scale on  $A$  if there is a set  $B \in \Delta$  with

$$\{(x, y, n) \in \mathbb{R} \times \mathbb{R} \times \omega : x \leq_n y \wedge y \in A\} = B \cap \{(x, y, n) \in \mathbb{R} \times \mathbb{R} \times \omega : y \in A\},$$

and if  $(x_k : k < \omega) \in {}^\omega\mathbb{R}$  with  $x_k \rightarrow x$  and  $\text{rank}_n(x_k) \rightarrow \alpha_n$  (i.e.  $\text{rank}_n(x_k)$  is eventually constant) for all  $n$ , then  $x \in A$  and  $\text{rank}_n(x) \leq \alpha_n$ . Here  $\text{rank}_n$  denotes the rank in  $\leq_n$ . A pointclass is scaled if there is a  $\Gamma$ -scale on every  $A \in \Gamma$ .

With each scale one associates a tree, from which the scale can again be defined:

**Definition 2.8.** Suppose  $(\leq_n : n \in \omega)$  is a  $\Gamma$ -scale on  $A \in \Gamma$  where  $\Gamma$  is a pointclass. The tree from the scale is defined as

$$T = \{(x \upharpoonright n, (\text{rank}_0(x), \dots, \text{rank}_{n-1}(x))) : x \in A \wedge n < \omega\}.$$

Note that  $A = p[T]$  in the situation of the definition. Given  $x \in p[T]$ , there are ordinals  $\alpha_n$  and reals  $x_k$  such that  $x \upharpoonright k = x_k \upharpoonright k$  for all  $k \in \omega$  and  $\text{rank}_n(x_k) = \alpha_n$  for all  $n \leq k$ , so  $x \in A$  by the semicontinuity of the scale.

The projective pointclasses  $\Pi_{2n+1}^1$  and  $\Sigma_{2n+2}^1$  and their boldface versions are scaled by the second periodicity theorem [21, theorem 39.8] if  $\text{Det}(\Delta_{2n}^1)$  holds, where  $\Delta_0^1 = \Delta_\omega^0$  denotes the pointclass of arithmetical sets. Let's fix the  $\Pi_{2n+1}^1$ -complete  $\Pi_{2n+1}^1$  set and the  $\Sigma_{2n+2}^1$ -complete  $\Sigma_{2n+2}^1$  set from the proof of the second periodicity theorem for each  $n < \omega$ . We will call these sets the complete  $\Pi_{2n+1}^1$  set and the complete  $\Sigma_{2n+2}^1$  set. Let's also fix the canonical scales on these sets from the proof of this theorem.

**Definition 2.9.** *Suppose  $\text{Det}(\Delta_{2n}^1)$  holds. Then  $T_{2n+1}$  denotes the tree from the canonical  $\Pi_{2n+1}^1$ -scale on the complete  $\Pi_{2n+1}^1$  set.*

We will work with transitive models of a fragment of ZF between which wellfoundedness is absolute.

**Definition 2.10.** *A transitive set  $\mathbb{A}$  is called admissible if  $(\mathbb{A}, \in) \models \text{KP}$ . We will call  $\mathbb{A}$  strongly admissible if  $(\mathbb{A}, \in) \models \text{KP}$  and every wellfounded relation in  $\mathbb{A}$  can be collapsed in  $\mathbb{A}$  to a transitive relation (“Axiom Beta”, see [3, chapter I, section 9]).*

For a background on admissible sets see Barwise [3].

**Lemma 2.11.** *Every  $\Sigma_{2n+2}^1(x)$  set is the projection of a tree which is uniformly defined from  $T_{2n+1}$  and  $x$  in every strongly admissible set  $\mathbb{A}$  with  $T_{2n+1}, x \in \mathbb{A}$ .*

*Proof.* The tree  $T$  from the scale on the complete  $\Sigma_{2n+2}^1$  set is essentially  $T_{2n+1}$ , see [21, theorem 38.4]. Any  $\Sigma_{2n+2}^1(x)$  set  $B$  for  $x \in \mathbb{R}$  is the preimage of the complete  $\Sigma_{2n+2}^1$  set under some function  $f : \mathbb{R} \rightarrow \mathbb{R}$  recursive in  $x$ . Then the tree

$$S := \{(s, h) \in (\omega \times \text{Ord})^{<\omega} : \exists y = y_{s,h} \supset s \forall i < lh(s) (\text{rank}_i(f(y)) = h(i))\}$$

induces a  $\Sigma_{2n+2}^1(x)$ -scale on  $B$ . Now  $\text{rank}_i(f(y))$  can be calculated from  $f(y)$  and  $T$  in any strongly admissible set  $\mathbb{A}$  with  $T, f(y) \in \mathbb{A}$ . Since the existence of  $y_{s,h}$  for given  $s, h$  is absolute between strongly admissible sets,  $S$  is as required.  $\square$

2.1.2. *Prewellorders under determinacy.* Typical examples of thin equivalence relations are given by prewellorders. We will need the following facts to know that prewellorders induce thin equivalence relations under determinacy.

**Lemma 2.12.** *(Kechris [18]) Suppose  $\Gamma$  is a pointclass containing the  $\Pi_1^0$  sets and  $\text{Det}(\Gamma)$  holds. Then every  $\exists\Gamma$  set has the Baire property and there is no  $\exists\Gamma$  wellorder of the reals.*

*Proof.* To prove that every  $\mathcal{O}\Gamma$  set has the Baire property, let  $B \subseteq \mathbb{R}^2$  and

$$A = \mathcal{O}B = \{x \in \mathbb{R} : \text{player 2 wins the game for } B_x\},$$

where  $B_x := \{y \in \mathbb{R} : (x, y) \in B\}$ . Basic open subsets of  $\mathbb{R}$  and  $\mathbb{R}^2$  are denoted by

$$U_s := \{x \in \mathbb{R} : x \upharpoonright \text{dom}(s) = s\}$$

and

$$U_{s,t} := \{(x, y) \in \mathbb{R}^2 : x \upharpoonright \text{dom}(s) = s \wedge y \upharpoonright \text{dom}(t) = t\}$$

for  $s, t \in \omega^{<\omega}$ .

We first claim that the Banach-Mazur game for

$$A \cup (\mathbb{R} - U_s) = \mathcal{O}[B \cup (\mathbb{R}^2 - U_{\emptyset, s})]$$

is determined for all  $s \in \omega^{<\omega}$ . In this game two players alternate playing finite sequences  $s_0, s_1, \dots$  and player 2 wins if  $s_0 \frown s_1 \frown \dots \in A \cup (\mathbb{R} - U_s)$ . This game is equivalent to the Banach-Mazur game for  $B \cup (\mathbb{R}^2 - U_t)$  by the game formula [18, theorem 3.3.1], and hence determined.

Now let  $S$  be the set of  $s \in \omega^{<\omega}$  such that player 2 has a winning strategy in the Banach-Mazur game for  $A \cup (\mathbb{R} - U_s)$ . Then  $A$  is comeager in  $U_s$  for each  $s \in S$  by the characterization of comeager sets in [21, theorem 8.33], so  $A$  is comeager in

$$U_1 := \bigcup_{s \in S} U_s.$$

The same theorem shows that for every  $t \in \omega^{<\omega} - S$ , the set  $A$  is meager in some nonempty open subset  $U_{f(t)}$  of  $U_t$ , so  $A$  is meager in

$$U_2 := \bigcup_{t \in \omega^{<\omega} - S} U_{f(t)}.$$

Since  $(\mathbb{R} - U_1) - U_2$  is nowhere dense, this implies that  $A \Delta U_1$  is meager. Hence  $A$  has the Baire property.

Let's recall the proof that there is no wellorder of the reals with the Baire property. If  $<$  were such a wellorder, we define

$$A := \{(x, y) \in \mathbb{R}^2 : x < y\},$$

$$B := \{(x, y) \in \mathbb{R}^2 : x > y\},$$

and

$$C := \{(x, y) \in \mathbb{R}^2 : x = y\}.$$

Then both  $A$  and  $B$  are not meager, since  $C$  is nowhere dense. Hence there is some  $x \in \mathbb{R}$  such that

$$A_x := \{y \in \mathbb{R} : x < y\}$$

is not meager by the theorem of Kuratowski and Ulam. Choose  $z$  as  $<$ -minimal with this property. Again  $A \cap (A_z \times A_z)$  and  $B \cap (A_z \times A_z)$  are not meager. So there is some  $x \in A_z$  with  $A_x$  not meager, contradicting the minimality of  $z$ .  $\square$

The proof of the previous lemma shows that there is no wellorder of the reals in the  $\sigma$ -algebra generated by  $\mathfrak{D}\Gamma$ .

**Lemma 2.13.** (*Kechris [18]*) *Suppose  $\Gamma$  is a boldface pointclass containing the  $\Pi_1^0$  sets and  $\text{Det}(\Gamma)$  holds. Then every prewellorder in  $\mathfrak{D}\Gamma$  is thin.*

*Proof.* Let  $\leq$  be a prewellorder in  $\mathfrak{D}\Gamma$  and suppose  $P \subseteq \mathbb{R}$  is a perfect set so that  $x \not\leq y$  and  $y \not\leq x$  for any two distinct  $x, y \in P$ . Then  $\leq$  wellorders  $P$ . Now  $\mathfrak{D}\Gamma$  is closed under continuous preimages since  $\Gamma$  is a boldface pointclass, so any continuous injective map  $f : \mathbb{R} \rightarrow P$  induces a  $\mathfrak{D}\Gamma$  wellorder of the reals, contradicting the previous lemma.  $\square$

The next two lemmas will be important for our purposes.

**Lemma 2.14.** *Det( $\Delta_{2n}^1$ ) implies that every  $\Pi_{2n+1}^1$  norm is thin.*

*Proof.* Suppose  $P \subseteq \mathbb{R}$  is a perfect set whose elements have pairwise different norms  $\neq \infty$  and let  $f : \mathbb{R} \rightarrow P$  be a continuous injective map. Then  $f$  induces a  $\Delta_{2n+1}^1$  wellorder of the reals.

Now  $Det(\Delta_{2n}^1)$  implies  $Det(\Pi_{2n}^1)$  by [22, theorem 5.1] and further  $\partial\Pi_{2n}^1 = \Sigma_{2n+1}^1$  by [21, proposition 39.6]. So there is no  $\Sigma_{2n+1}^1$  wellorder of the reals by lemma 2.12.  $\square$

Note that the conclusion of the previous lemma follows from the Baire property or the Lebesgue measurability of all  $\Delta_{2n+1}^1$  sets alone.

**Lemma 2.15.** *Det( $\Pi_{2n+1}^1$ ) implies that every  $\Sigma_{2n+2}^1$  norm is thin.*

*Proof.* As the previous lemma; otherwise there is a  $\Delta_{2n+2}^1$  wellorder of the reals, contradicting lemma 2.12.  $\square$

It is sufficient to assume the Baire property or the Lebesgue measurability of all  $\Delta_{2n+2}^1$  sets for the previous lemma.

**Lemma 2.16.** *The following are equivalent:*

- (1) every  $\Delta_2^1$  prewellorder of the reals is thin,
- (2) there is no  $\Delta_2^1$  wellorder of the reals, and
- (3)  $L[x]$  does not contain  $\mathbb{R}$  for any  $x \in \mathbb{R}$ .

*Proof.* Condition 1 clearly implies condition 2. To show that 2 implies 1, suppose  $\leq$  is a  $\Delta_2^1$  prewellorder of the reals and  $P \subseteq \mathbb{R}$  is perfect with  $x < y$  or  $x > y$  for any two distinct  $x, y \in P$ . Then any continuous injective map  $f : \mathbb{R} \rightarrow P$  induces a  $\Delta_2^1$  wellorder of the reals.

Now condition 2 implies condition 3, since if  $\mathbb{R} \subseteq L[x]$  for some  $x \in \mathbb{R}$ , then the order of constructibility of the reals is  $\Delta_2^1(x)$ . To show that 3 implies 2, note that  $\mathbb{R} \subseteq L[x]$  by [16, theorem 25.39] if there is a  $\Delta_2^1(x)$  wellorder of the reals.  $\square$

The projective ordinals are given by

**Definition 2.17.** *The  $n^{\text{th}}$  projective ordinal  $\delta_n^1$  is the supremum of lengths of  $\Delta_n^1$  prewellorders for  $n \geq 1$ .*

We state some of their properties, since the projective ordinals play an essential role in calculating the number of equivalence classes of thin projective equivalence relations in section 3.2.

**Lemma 2.18.** *The following facts hold for the projective ordinals:*

- (1) (Martin)  $\text{ZF} + \text{PD}$  implies  $\delta_1^1 = \omega_1$  and  $\delta_n^1 \leq \omega_n$  for  $n \leq 4$ ,
- (2) (Kechris, Moschovakis)  $\text{ZF} + \text{PD}$  implies  $\delta_n^1 < \delta_{n+1}^1$  for all  $n$ ,
- (3) (Moschovakis)  $\text{ZF} + \text{AD}$  implies that each  $\delta_n^1$  is a cardinal, and
- (4) (Steel, Van Wesep [43])  $\text{ZF} + \text{AD}^{L(\mathbb{R})} + \delta_2^1 = \omega_2$  is consistent relative to  $\text{ZF} + \text{AD} + \text{AC}_{\mathbb{R}}$ .

*Proof.* The proofs for parts 1 and 2 can be found in [20, theorem 9.1]. For part 3 see [20, theorem 2.2]. Note that Jackson [15] has computed all  $\delta_n^1$  exactly under AD. For part 4 see [43]. Note that Woodin [46, theorem 3.17] proved that  $\delta_2^1 = \omega_2$  holds if  $\mathcal{P}(\omega_1)^{\#}$  exists and the nonstationary ideal on  $\aleph_1$  is  $\aleph_2$ -saturated.  $\square$

An important open question is how large the projective ordinal  $\delta_n^1$  for  $n \geq 3$  can be under  $\text{ZFC} + \text{AD}^{L(\mathbb{R})}$ . In fact, it is still open if  $\text{AD}^{L(\mathbb{R})}$  implies  $\delta_n^1 \leq \omega_n$  for all  $n$ , see [17, question 30.34].

Note that the consistency strength of  $\delta_2^1 = \omega_2$  in the presence of sharps for reals is somewhere between a strong cardinal and a Woodin cardinal with a measurable cardinal above by work of Steel and Welch [44] and Woodin [46, theorem 3.25]. While we focus on the situation that  $\text{ZF} + \text{PD}$  holds, one can consider the case that  $0^{\#}$  does not exist. Note that MA and  $\omega_1 = \omega_1^L$  already imply  $\delta_2^1 = \omega_2$ . This is because  $\omega_1 = \omega_1^L$  implies that any subset of  $\omega_1$  can be coded as  $\Delta_1^{HC}$  in a real by

c.c.c. forcing so that  $\omega_1$  many dense subsets suffice to define the real. Moreover in this situation  $\delta_3^1$  can be quite easily forced to be arbitrarily large with a forcing from Harrington [6].

**2.2. Mice with Woodin cardinals.** In this section tools for mice with Woodin cardinals are presented. The results are due to Martin, Steel, and Woodin, or folklore. For missing definitions and proofs see Martin and Steel [28], Mitchell and Steel [29], Schindler and Zeman [37], Steel [42], and Zeman [47]. Several facts about  $\omega_1 + 1$ -iterable premice are adapted to  $\omega_1$ -iterable premice. The reason is that we only want to assume PD; all one can get from PD is the existence of  $\omega_1$ -iterable premice with  $n$  Woodin cardinals for arbitrary  $n < \omega$ .

2.2.1. *Premice, comparison, and  $M_n^\#$ .*

**Definition 2.19.** *A self-wellordered (swo) set is a set which codes a wellorder of itself. The height of a self-wellordered set  $X$  is*

$$ht(X) := \sup((Ord \cap tc(X)) \cup \omega).$$

Every self-wellordered set can be coded by a set  $\sup(A) \cup A$ , where  $A$  is a set of ordinals. Recall that the first level of the  $J$ -hierarchy built over a set  $X$  is defined as  $J_0(X) = tc(\{X\})$ .

**Definition 2.20.** *A potential  $X$ -premouse is a structure*

$$\mathcal{M} = (J_\beta^{\vec{F}}(X), \in, X, \vec{F} \upharpoonright \beta, F_\beta)$$

where  $X$  is swo and  $\vec{F}$  is a fine extender sequence relative to  $X$ . An  $X$ -premouse is a potential  $X$ -premouse all of whose proper initial segments are  $\omega$ -sound; a premouse is simply a  $\emptyset$ -premouse. A boldface or relativized premouse is an  $X$ -premouse for some swo set  $X$ .  $\mathcal{M}$  is called active if  $F_\beta \neq \emptyset$ , otherwise it is passive. We write  $\vec{F}^\mathcal{M}$  for the extender sequence of  $\mathcal{M}$ .

For the definition of fine extender sequences see [42, definition 2.4] and for  $\omega$ -sound [42, definition 2.17].

**Definition 2.21.** *Suppose*

$$\mathcal{M} = (J_{\alpha}^{\vec{F}}(X), \in, X, \vec{F} \upharpoonright \alpha, F_{\alpha})$$

and

$$\mathcal{N} = (J_{\beta}^{\vec{F}}(X), \in, X, \vec{F} \upharpoonright \beta, F_{\beta})$$

are  $X$ -premise where  $X$  is swo and  $\alpha \leq \beta$  ( $\alpha < \beta$ ). Then  $\mathcal{M}$  is called a (proper) initial segment of  $\mathcal{N}$  and we write  $\mathcal{M} \sqsubseteq \mathcal{N}$  ( $\mathcal{M} \triangleleft \mathcal{N}$ ). For notation write

$$\mathcal{N} \upharpoonright \alpha := (J_{\alpha}^{\vec{F}}(X), \in, X, \vec{F} \upharpoonright \alpha, F_{\alpha})$$

and

$$\mathcal{N} \upharpoonright \alpha := (J_{\alpha}^{\vec{F}}(X), \in, X, \vec{F} \upharpoonright \alpha, \emptyset).$$

An ordinal  $\delta$  is called a cutpoint of  $\mathcal{M}$ , if for no extender  $F$  on the  $\mathcal{M}$ -sequence do we have  $\text{crit}(F) < \delta \leq \text{lh}(F)$ . For the definition of iteration trees see [42, section 3.1]; for normal iteration trees see [47, section 4.2]. An iteration tree  $\mathcal{T}$  on an  $X$ -premouse  $\mathcal{M}$  is said to live on  $\mathcal{M} \upharpoonright \alpha$  if all extenders in  $\mathcal{T}$  have length less than  $\alpha$ .

**Definition 2.22.** *Let  $k \leq \omega$  and  $\theta \in \text{Ord}$  and suppose  $X$  is swo. An  $X$ -premouse  $\mathcal{M}$  is called  $(k, \theta)$ -iterable if player 2 has a winning strategy in the iteration game  $\mathcal{G}_k(\mathcal{M}, \theta)$  described in [42, section 3.1].  $\mathcal{M}$  is called normally  $(k, \theta)$ -iterable if it is  $(k, \theta)$ -iterable with respect to normal iteration trees. It is (normally)  $(k, \theta)$ -iterable above  $\delta$  if it is (normally)  $(k, \theta)$ -iterable with respect to (normal) iteration trees all of whose extenders have critical points above  $\delta$ . It is (normally)  $\theta$ -iterable if it is (normally)  $(\omega, \theta)$ -iterable.*

We need the next two lemmas from [42, theorem 3.11] and [42, corollary 3.12].

**Lemma 2.23.** (*Comparison lemma*) *Let  $\mathcal{M}$  and  $\mathcal{N}$  be countable  $X$ -premouse, where  $X$  is swo. Let  $\delta$  be a cutpoint of both  $\mathcal{M}$  and  $\mathcal{N}$  and suppose  $\mathcal{M}$  and  $\mathcal{N}$  are normally  $\omega_1 + 1$ -iterable above  $\delta$  and  $\omega$ -sound above  $\delta$  with  $\mathcal{M}|_\delta = \mathcal{N}|_\delta$ . Then there are countable iteration trees  $\mathcal{S}$  on  $\mathcal{M}$  and  $\mathcal{T}$  on  $\mathcal{N}$  with last models  $\mathcal{M}_\alpha^{\mathcal{S}}$  and  $\mathcal{M}_\beta^{\mathcal{T}}$  so that*

- (1)  $[0, \alpha]_{\mathcal{S}}$  does not drop in model or degree and  $\mathcal{M}_\alpha^{\mathcal{S}} \trianglelefteq \mathcal{M}_\beta^{\mathcal{T}}$ , or
- (2)  $[0, \beta]_{\mathcal{T}}$  does not drop in model or degree and  $\mathcal{M}_\beta^{\mathcal{T}} \trianglelefteq \mathcal{M}_\alpha^{\mathcal{S}}$ .

**Lemma 2.24.** *Let  $\mathcal{M}$  and  $\mathcal{N}$  be countable  $X$ -premouse where  $X$  is swo. Let  $\delta$  be a cutpoint of both  $\mathcal{M}$  and  $\mathcal{N}$  and suppose  $\mathcal{M}$  and  $\mathcal{N}$  are normally  $\omega_1 + 1$ -iterable above  $\delta$  and  $\omega$ -sound above  $\delta$  with  $\rho_\omega(\mathcal{M}), \rho_\omega(\mathcal{N}) \leq \delta$  and  $\mathcal{M}|_\delta = \mathcal{N}|_\delta$ . Then  $\mathcal{M} \trianglelefteq \mathcal{N}$  or  $\mathcal{N} \trianglelefteq \mathcal{M}$ .*

*Proof.* Neither  $\mathcal{M}$  nor  $\mathcal{N}$  are moved in the coiteration, since they are  $\omega$ -sound above  $\delta$  and  $\rho_\omega(\mathcal{M}), \rho_\omega(\mathcal{N}) \leq \delta$ . □

Let's recall the definition of Woodin cardinals:

**Definition 2.25.** *Suppose  $A \subseteq V_\delta$ . An ordinal  $\kappa < \delta$  is  $A$ -reflecting in  $\delta$  if for all  $\alpha < \delta$  there is an extender  $F$  in  $V_\delta$  with  $\text{crit}(F) = \kappa$ ,  $j_F(\kappa) > \alpha$ , and*

$$j_F(A) \cap V_\alpha = A \cap V_\alpha,$$

where  $j_F : V \rightarrow \text{ult}(V, F)$  is the ultrapower embedding.

**Definition 2.26.** *A cardinal  $\delta$  is a Woodin cardinal if for every  $A \subseteq V_\delta$  there is some  $\kappa < \delta$  which is  $A$ -reflecting in  $\delta$ .*

Now  $M_n^\#$  can be defined:

**Definition 2.27.** *Let  $X$  be swo and  $n \leq \omega$ . An  $X$ -premouse  $\mathcal{M}$  is  $n$ -small above  $\delta$  if there is no extender  $F$  on the  $\mathcal{M}$ -sequence so that in  $\mathcal{M}|_{\text{crit}(F)}$  there are  $n$  Woodin cardinals above  $\delta$ .*

**Definition 2.28.** *Let  $\mathcal{M}$  be an active  $\omega_1$ -iterable  $X$ -premouse, where  $X$  is swo, such that  $\mathcal{M}$  is  $\omega$ -sound above  $ht(X)$  and  $\rho_1(\mathcal{M}) \leq ht(X)$ . Let  $F$  be the top extender of  $\mathcal{M}$  and  $n \leq \omega$ .  $\mathcal{M}$  is called  $M_n^\#(X)$  if  $\mathcal{M}|crit(F)$  is  $n$ -small and in  $\mathcal{M}$  there are  $n$  Woodin cardinals below  $crit(F)$ . Moreover  $M_n^\# := M_n^\#(\emptyset)$ .*

Note that usually  $M_n^\#$  is defined as an  $\omega_1 + 1$ -iterable premouse with the same first-order properties; thus  $M_n^\#$  is unique by the comparison lemma. The  $M_n^\#(X)$  defined here is also unique in the relevant case that  $M_n^\#(x)$  exists for every  $x \in \mathbb{R}$ . Suppose we have two candidates for  $M_n^\#(X)$  where  $X$  is swo. Let's consider the preimages of the candidates in the transitive collapse of a countable substructure of some large  $V_\lambda$ . Since these are  $\omega_1$ -iterable, they can be compared in  $M_n^\#(x)$  for some  $x \in \mathbb{R}$  by the argument in lemma 2.38 below.

The standard definition of  $M_n^\#(X)$  just states that it is countably iterable, i.e. all countable substructures are  $\omega_1 + 1$ -iterable, instead of being  $\omega_1 + 1$ -iterable itself. Everything would work if in the definition of  $M_n(X)$  we only ask that countable substructures are  $\omega_1$ -iterable.

Note that modulo Gödel numbers for first-order formulas, any  $x$ -premouse  $\mathcal{M}$  with  $x \in \mathbb{R}$  and  $\rho_{k+1}(\mathcal{M}) = \omega$  comes with a code  $z \in \mathbb{R}$  from the canonical  $\Sigma_1^{(k)}$ -definable surjection from  $\omega$  onto  $\mathcal{M}$ , see [47, section 1.6] for the definition of  $\Sigma_1^{(k)}$  formulas. Hence there is no need to distinguish between  $M_n^\#(\mathcal{M})$  and  $M_n^\#(z)$ .

**Definition 2.29.** *Let  $\mathcal{M}$  be an active  $\omega_1$ -iterable  $X$ -premouse, where  $X$  is swo, such that  $\mathcal{M}$  is  $\omega$ -sound above  $ht(X)$  and  $\rho_1(\mathcal{M}) \leq ht(X)$ . Let  $F$  be the top extender of  $\mathcal{M}$  and suppose the topmost extender  $G$  below  $F$  is total.  $\mathcal{M}$  is called  $M_n^\dagger(X)$  if  $\mathcal{M}|crit(G)$  is  $n$ -small and in  $\mathcal{M}$  there are  $n$  Woodin cardinals below  $crit(G)$ . Moreover  $M_n^\dagger := M_n^\dagger(\emptyset)$ .*

Again  $M_n^\dagger(X)$  is unique in the relevant situation that  $M_n^\#(x)$  exists for every  $x \in \mathbb{R}$ . Since we would like to prove the main theorem from PD, we will use

**Theorem 2.30.** (Harrington, Martin, Steel, Woodin, Neeman) *The following are equivalent for  $n < \omega$ :*

- (1)  $Det(\Pi_{n+1}^1)$
- (2) *there is an  $M_n^\#(x)$  for every  $x \in \mathbb{R}$*
- (3) *there is a unique  $M_n^\#(x)$  for every  $x \in \mathbb{R}$ .*

*Proof.* See [34, theorem 5.3]. Harrington [7] proved the implication from 1 to 3 for  $n = 0$ . For arbitrary  $n$  see Koellner and Woodin [23]. Martin [25] proved that 2 implies 1 for  $n = 0$ , Neeman [31] has a proof for arbitrary  $n$ . The original proof for odd  $n$  is due to Woodin. Note that lemma 2.38 below can be used to show that 2 implies 3. □

2.2.2. *Genericity iteration.* We will use a theorem of Woodin to iterate an  $\omega_1 + 1$ -iterable premouse with a Woodin cardinal so that a given real is generic over the iterate for Woodin's extender algebra.

The extender algebra is built from a set of infinitary formulas. Let's state the necessary definitions. We let  $\delta$  be an inaccessible cardinal and  $\mathcal{L}$  a language which contains at least  $\in$  and constants  $c$  for a real and  $\dot{n}$  for each  $n < \omega$ . Let  $N$  be the set of atomic formulas  $\dot{n} \in c$  for  $n < \omega$ . Now let  $\mathcal{L}_{\delta,0,N}$  be the closure of  $N$  under negations and infinitary disjunctions and conjunctions of length less than  $\delta$ . Note that one can equivalently work with the infinitary logic built over a language with propositional formulas  $p_n$  for  $n < \omega$ .

The infinitary proof calculus for this logic has the infinitary rule

$$\forall \alpha < \beta \vdash \varphi_\alpha \quad \Rightarrow \quad \vdash \bigwedge_{\alpha < \beta} \varphi_\alpha$$

in addition to the rules of first-order logic; for details see Barwise [3, chapter III, definition 5.1]. Let  $\chi$  be the  $\mathcal{L}_{\delta,0,N}$ -sentence

$$\bigwedge_{n < \omega} (\forall x \in \dot{n} \bigvee_{m < n} x = \dot{m}) \wedge (\bigwedge_{m < n} \dot{m} \in \dot{n}) \wedge c \subseteq \omega,$$

which we add as an axiom. Hence in every model,  $c$  is interpreted as a subset of  $\omega$  and each  $\dot{n}$  is interpreted as  $n$ . Moreover, the infinitary disjunction of a sequence  $\vec{\varphi} = (\varphi_\alpha : \alpha < \beta)$  of  $\mathcal{L}_{\delta,0,N}$ -formulas is denoted by  $\bigvee_{\alpha < \beta} \varphi_\alpha$  or  $\bigvee \vec{\varphi}$ .

The following is Steel's version [42, section 7.2] of Woodin's extender algebra for fine structural mice.

**Definition 2.31.** *Let  $\mathcal{M}$  be an  $X$ -premouse with Woodin cardinal  $\delta$ , where  $X$  is swo. Let  $S$  be the set of all  $\mathcal{L}_{\delta,0,N}$ -formulas*

$$\bigvee \vec{\varphi} \leftrightarrow \bigvee j_E(\vec{\varphi}) \upharpoonright \lambda$$

in  $\mathcal{M}$ , where

- (1)  $\vec{\varphi} = (\varphi_\alpha : \alpha < \kappa) \in \mathcal{M}$  is a sequence of  $\mathcal{L}_{\delta,0,N}$ -formulas with  $\kappa < \delta$ ,
- (2)  $F$  is an extender on the  $\mathcal{M}$ -sequence with  $\text{crit}(F) = \kappa \leq \lambda < \delta$ ,
- (3)  $\nu(F)$  is a cardinal in  $\mathcal{M}$ , and
- (4)  $j_F(\vec{\varphi}) \upharpoonright \lambda \in J_{\nu(F)}^{\mathcal{M}}$ .

Here  $\nu(F)$  is the natural length of  $F$ , see [42, definition 2.2]. Working in  $\mathcal{M}$ , the extender algebra  $\mathbb{W}_\delta$  over  $\delta$  is defined as the Lindenbaum algebra over  $\mathcal{L}_{\delta,0,N}$  for provability from  $S$ ; let

$$[\varphi] := \{\psi \in \mathcal{L}_{\delta,0,N} : S \vdash \varphi \leftrightarrow \psi\}$$

for  $\varphi \in \mathcal{L}_{\delta,0,N}$  and define

$$\mathbb{W}_\delta := \{[\varphi] : \varphi \in \mathcal{L}_{\delta,0,N}\}$$

with partial ordering

$$[\varphi] \leq [\psi] :\Leftrightarrow S \vdash \varphi \rightarrow \psi.$$

The extender algebra  $\mathbb{W}_\delta$  has size  $\delta$  and the  $\delta$ -c.c. in  $\mathcal{M}$  by [42, theorem 7.14]. We will heavily use the next theorem of Woodin following Steel [42]:

**Lemma 2.32.** (*Genericity iteration*) *Let  $\mathcal{M}$  be a countable  $X$ -premouse, where  $X$  is swo. Suppose  $\mathcal{M}$  is normally  $\omega_1 + 1$ -iterable above  $\gamma < \delta$  and  $\delta$  is Woodin in  $\mathcal{M}$ . Then for each  $x \in \mathbb{R}$ , there is a countable iteration tree  $\mathcal{T}$  on  $\mathcal{M}$  with iteration map  $\pi$  and last model  $\mathcal{M}_\alpha^{\mathcal{T}}$  such that  $[0, \alpha]_{\mathcal{T}}$  does not drop in model and  $x$  is  $\mathbb{W}_{\pi(\delta)}^{\mathcal{M}_\alpha^{\mathcal{T}}}$ -generic over  $\mathcal{M}_\alpha^{\mathcal{T}}$ .*

*Proof.* See [42, theorem 7.14]. The idea is to iterate away the least extender which induces an axiom false for  $x$ . A reflection argument as in the proof of the comparison lemma shows that after countably many steps  $x$  is a model of  $\pi(S)$ , where  $\pi$  is the iteration map. It follows that  $x$  is  $\mathbb{W}_{\pi(\delta)}^{\mathcal{M}_\alpha^{\mathcal{T}}}$ -generic.  $\square$

2.2.3. *The  $\mathcal{Q}$ -structure iteration strategy.* In this section we describe a partial iteration strategy based on so-called  $\mathcal{Q}$ -structures. In the relevant cases this is the unique iteration strategy.

**Definition 2.33.** *Suppose  $\mathcal{T}$  is an iteration tree of limit length  $\theta$  with models  $(\mathcal{M}_\alpha : \alpha < \theta)$  and extenders  $(F_\alpha : \alpha < \theta)$ . Define*

$$\delta(\mathcal{T}) := \sup_{\alpha < \theta} lh(F_\alpha)$$

and

$$\mathcal{M}(\mathcal{T}) := \bigcup_{\alpha < \theta} \mathcal{M}_\alpha \upharpoonright lh(F_\alpha),$$

where  $lh(F_\alpha)$  denotes the length of  $F_\alpha$ . The model  $\mathcal{M}(\mathcal{T})$  is called the common part model of  $\mathcal{T}$ .

$\mathcal{Q}$ -structures for iteration trees are defined as follows.

**Definition 2.34.** Let  $\mathcal{T}$  be an iteration tree of limit length on an  $X$ -premouse  $\mathcal{M}$ , where  $X$  is swo. A  $\mathcal{Q}$ -structure for  $\mathcal{T}$  is an  $X$ -premouse  $\mathcal{Q}$  with

- (1)  $\mathcal{M}(\mathcal{T}) \trianglelefteq \mathcal{Q}$  such that  $\delta(\mathcal{T})$  is a cutpoint of  $\mathcal{Q}$ ,
- (2)  $\mathcal{Q}$  is  $\omega_1$ -iterable above  $\delta(\mathcal{T})$ , and
- (3) the Woodin property of  $\delta(\mathcal{T})$  is destroyed definably over  $\mathcal{Q}$ , i.e. there is a  $k < \omega$  such that
  - (a)  $\mathcal{Q}$  is  $k + 1$ -sound and
  - (b) either  $\rho_{k+1}(\mathcal{Q}) < \delta(\mathcal{T})$ , or  $k$  is minimal such that there is a map  $f : \delta(\mathcal{T}) \rightarrow \delta(\mathcal{T})$  which is  $\Sigma_1^{(k)}$ -definable over  $\mathcal{Q}$  so that for no extender  $F$  on the  $\mathcal{Q}$ -sequence do we have  $i_F(f)(\text{crit}(F)) \geq \nu(F)$ .

If in the previous definition  $\mathcal{Q} = (J_{\beta}^{\vec{F}}(X), \in, X, \vec{F} \upharpoonright \beta, F_{\beta})$  is a proper initial segment of a premouse, then condition 3 simplifies to the statement that  $\beta$  is minimal with  $J_{\beta+1}^{\vec{F}}(X) \models \text{''}\delta(\mathcal{T}) \text{ is not Woodin''}$ . Based on  $\mathcal{Q}$ -structures, one builds a partial iteration strategy:

**Definition 2.35.** Let  $\mathcal{T}$  be a normal iteration tree on an  $X$ -premouse  $\mathcal{M}$ , where  $X$  is swo. Let  $\Sigma(\mathcal{T})$  be the unique cofinal branch  $b \subseteq \mathcal{T}$  such that  $\mathcal{M}_b$  is well-founded and carries a  $\mathcal{Q}$ -structure  $\mathcal{Q} \trianglelefteq \mathcal{M}_b^{\mathcal{T}}$ , if such a branch exists. Let  $\Sigma(\mathcal{T})$  be undefined if there is no such branch, or if there is one but it is not unique. This partial iteration strategy for normal iteration trees is called the  $\mathcal{Q}$ -structure iteration strategy.

We have

**Lemma 2.36.** If  $\Sigma$  is a  $\theta$ -iteration strategy for normal iteration trees on an  $X$ -premouse  $\mathcal{N}$ , where  $X$  is swo, then  $\Sigma$  is a  $\theta$ -iteration strategy for normal iteration trees on every initial segment  $\mathcal{M} \trianglelefteq \mathcal{N}$ .

Let's consider the situation that  $\mathcal{M}$  and  $\mathcal{N}$  are  $X$ -premouse which are  $\theta$ -iterable via  $\Sigma$ , where  $X$  is swo. Suppose  $\mathcal{M}|_\delta = \mathcal{N}|_\delta$  and  $\delta$  is a cutpoint of both  $\mathcal{M}$  and  $\mathcal{N}$ . Then every iteration tree  $\mathcal{T}$  according to  $\Sigma$  on  $\mathcal{M}$  living on  $\mathcal{M}|_\delta$  gives rise to an iteration tree on  $\mathcal{N}$ . In this case we say that  $\mathcal{T}$  acts on  $\mathcal{N}$ .

In the relevant situation the  $\mathcal{Q}$ -structure iteration strategy  $\Sigma$  is the unique  $\omega_1$ -iteration strategy for normal iteration trees:

**Lemma 2.37.** *Suppose  $M_n^\#(x)$  exists for every  $x \in \mathbb{R}$ . Then  $\Sigma$  is the unique  $\omega_1$ -iteration strategy for normal iteration trees on  $M_n^\#(x)$ .*

We prove a more general fact:

**Lemma 2.38.** *Suppose  $M_n^\#(x)$  exists for every  $x \in \mathbb{R}$ . Let  $\mathcal{N}$  be a countable  $X$ -premouse with  $n$  Woodin cardinals above  $\delta$  and an extender above, where  $X$  is swo. Suppose  $\mathcal{N}$  is normally  $\omega_1$ -iterable above  $\delta$ . Let  $\mathcal{M} \in \mathcal{N}$  be a  $Y$ -premouse which is countable in  $\mathcal{N}$  and normally  $\omega_1$ -iterable above  $\delta$  via an iteration strategy  $\Sigma'$ , where  $Y$  is swo. Further suppose  $\mathcal{M}$  and  $\mathcal{N}$  are  $(n+1)$ -small above  $\delta$  and  $\omega$ -sound above  $\delta$  with  $\rho_\omega(\mathcal{M}) \leq \delta$  and  $\rho_\omega(\mathcal{N}) \leq \delta$ . Then  $\mathcal{M}$  is normally  $\omega_1 + 1$ -iterable above  $\delta$  via  $\Sigma$  in  $\mathcal{N}$ . Moreover,  $\Sigma$  is the unique  $\omega_1$ -iteration strategy for normal iteration trees on  $\mathcal{M}$  in  $V$ .*

*Proof.* The proof is organized as an induction on  $n$ . We will show that

$$\Sigma'(\mathcal{T}) = \Sigma^{\mathcal{N}}(\mathcal{T}) = \Sigma(\mathcal{T})$$

for all normal iteration trees  $\mathcal{T} \in \mathcal{N}$  on  $\mathcal{M}$  above  $\delta$  of limit length  $\leq \omega_1^{\mathcal{N}}$ .

Suppose this has been proved for all  $k < n$  and let  $\mathcal{T} \in \mathcal{N}$  be a normal iteration tree on  $\mathcal{M}$  above  $\delta$  of limit length  $\leq \omega_1^{\mathcal{N}}$  according to  $\Sigma'$ . Let  $b := \Sigma'(\mathcal{T})$ . Then  $\mathcal{M}_b^{\mathcal{T}}$  carries a  $\mathcal{Q}$ -structure  $\mathcal{Q} \trianglelefteq \mathcal{M}_b^{\mathcal{T}}$ , since  $\mathcal{T}$  is a normal iteration tree and  $\rho(\mathcal{M}) \leq \delta < \delta(\mathcal{T})$ . We can inductively assume that  $\mathcal{T}$  is according to  $\Sigma'$ ,  $\Sigma$ , and

$\Sigma^{\mathcal{N}}$ . It has to be shown that  $\mathcal{Q} \in \mathcal{N}$ . It can be assumed that  $\mathcal{M}(\mathcal{T}) \triangleleft \mathcal{Q}$ , so  $\delta(\mathcal{T})$  is Woodin in  $\mathcal{Q}$ .

For  $n = 0$  there are no extenders above  $\delta(\mathcal{T})$  on the  $\mathcal{Q}$ -sequence, since  $\mathcal{Q}$  is 1-small; in this case  $\mathcal{Q} \in \mathcal{N}$  since  $ht(\mathcal{Q}) < ht(\mathcal{N})$ . Now suppose  $n > 0$ . Let  $\kappa$  be the critical point of an extender on the  $\mathcal{N}$ -sequence such that in  $\mathcal{N}$  there are  $n$  Woodin cardinals between  $\delta$  and  $\kappa$ . We do an  $L[\vec{E}]$ -construction over  $\mathcal{M}(\mathcal{T})$  in  $\mathcal{N}|\kappa$ .  $L[\vec{E}, \mathcal{M}(\mathcal{T})]^{|\mathcal{N}| \kappa}$  inherits Woodin cardinals and iterability from  $\mathcal{N}$ , see [29, chapter 11].

We first prove  $\mathcal{Q} \in \mathcal{N}$  in the special case  $\mathcal{M} = M_n^\#(x)$ . Otherwise it is not clear how to find a premouse  $\mathcal{P}$  as in case 3 of the next claim.

**Claim 2.39.** *If  $\mathcal{M} = M_n^\#(x)$  for some  $x \in \mathbb{R}$ , then  $\mathcal{Q} \in \mathcal{N}$ .*

*Proof.* We distinguish three cases.

**Case 1.**  $\delta(\mathcal{T})$  is not Woodin in  $L[\vec{E}, \mathcal{M}(\mathcal{T})]^{|\mathcal{N}| \kappa}$ .

Let  $\alpha < \kappa$  be minimal such that  $\delta(\mathcal{T})$  is not Woodin in  $J_{\alpha+1}[\vec{E}, \mathcal{M}(\mathcal{T})]^{|\mathcal{N}| \kappa}$ . Let  $\mathcal{P} := J_\alpha[\vec{E}, \mathcal{M}(\mathcal{T})]^\mathcal{N}$ . Then  $\mathcal{P}$  and  $\mathcal{Q}$  are  $n$ -small above  $\delta(\mathcal{T})$  and  $\omega$ -sound above  $\delta(\mathcal{T})$  with  $\mathcal{P}|\delta(\mathcal{T}) = \mathcal{Q}|\delta(\mathcal{T})$  and  $\rho_\omega(\mathcal{P}), \rho_\omega(\mathcal{Q}) \leq \delta(\mathcal{T})$ . So  $\mathcal{P}$  and  $\mathcal{Q}$  can be coiterated in  $M_{n-1}^\#(x)$  by the induction hypothesis, where  $x \in \mathbb{R}$  codes  $\mathcal{P}$  and  $\mathcal{Q}$ . Now  $\mathcal{P}$  cannot be a proper initial segment of  $\mathcal{Q}$  because  $\mathcal{Q}$  is a  $\mathcal{Q}$ -structure. Thus  $\mathcal{Q} \trianglelefteq \mathcal{P}$  and hence  $\mathcal{Q} \in \mathcal{N}$ .

**Case 2.**  $\delta(\mathcal{T})$  is Woodin in  $L[\vec{E}, \mathcal{M}(\mathcal{T})]^{|\mathcal{N}| \kappa}$  and there is some  $\mathcal{P}$  in the  $L[\vec{E}]$ -construction with  $\rho_\omega(\mathcal{P}) \leq \delta(\mathcal{T})$ .

Again  $\mathcal{P}$  and  $\mathcal{Q}$  can be compared in  $M_n^\#(x)$ , where  $x \in \mathbb{R}$  codes  $\mathcal{P}$  and  $\mathcal{Q}$ .

**Case 3.**  $\delta(\mathcal{T})$  is Woodin in  $L[\vec{E}, \mathcal{M}(\mathcal{T})]^{|\mathcal{N}| \kappa}$  and the last projectum never falls below  $\delta(\mathcal{T})$  in the  $L[\vec{E}]$ -construction.

Let  $\mathcal{P}$  be the first model in the  $L[\vec{E}]$ -construction with  $n - 1$  Woodin cardinals above  $\delta(\mathcal{T})$  and two extenders above. Then  $\rho_1(\mathcal{P}) = \delta(\mathcal{T})$ .  $\mathcal{P}$  and  $\mathcal{Q}$  can be coiterated in  $M_n^\#(x)$  by the induction hypothesis, where  $x \in \mathbb{R}$  codes  $\mathcal{P}$  and  $\mathcal{Q}$ . But  $\mathcal{Q}$  cannot win the coiteration, since  $\mathcal{P}$  has more large cardinals. So  $\mathcal{Q} \triangleleft \mathcal{P}$ .

In fact, this case does not occur by the proof of the following claims.  $\square$

**Claim 2.40.** *If  $\mathcal{N} = M_n^\#(x)$  for some  $x \in \mathbb{R}$ , then  $\mathcal{Q} \in \mathcal{N}$ .*

*Proof.* It suffices to show that case 3 cannot occur. Let  $\mathcal{P}$  be the premouse obtained by adding the extender with critical point  $\kappa$  on top of  $L[\vec{E}, \mathcal{M}(\mathcal{T})]^{N|\kappa}$ . Now  $\mathcal{P}$  and  $\mathcal{Q}$  can be compared, since  $\mathcal{N}$  and hence  $\mathcal{P}$  is  $\omega_1 + 1$ -iterable in  $M_n^\#(x)$  by the previous claim, where  $x \in \mathbb{R}$  codes  $\mathcal{N}$ . But neither can iterate to an initial segment of the other.  $\square$

We finally conclude:

**Claim 2.41.**  $\mathcal{Q} \in \mathcal{N}$ .

*Proof.* Again it is sufficient that case 3 does not occur. This holds because  $\mathcal{N}$  is  $\omega_1 + 1$ -iterable in  $M_n^\#(x)$  by the previous claim, where  $x \in \mathbb{R}$  codes  $\mathcal{N}$ .  $\square$

It remains to be shown that  $b = \Sigma(\mathcal{T}) \in \mathcal{N}$ . Let  $g$  be  $Col(\omega, ht(\mathcal{T}))$ -generic over  $\mathcal{N}$ . Note that one can rearrange  $\mathcal{N}[g]$  as a boldface premouse, see [36, lemma 1.4] and [41, section 3]. We can form a tree in  $\mathcal{N}[g]$  searching for cofinal branches  $b \subseteq \mathcal{T}$  with  $\mathcal{Q} \trianglelefteq \mathcal{M}_b^\mathcal{T}$ . The nodes consist of initial segments of  $b$  and partial finite  $\in$ -isomorphisms between  $\mathcal{Q}$  and the corresponding model. Since wellfoundedness is absolute between  $\mathcal{N}[g]$  and  $V$ ,  $\mathcal{N}[g]$  knows that there is a branch in this tree. But there is at most one cofinal branch in  $\mathcal{T}$  with the required property by the argument in the proof of [42, corollary 6.14]. Hence  $b \in \mathcal{N}$  by homogeneity of  $Col(\omega, ht(\mathcal{T}))$ . Thus  $\Sigma(\mathcal{T}) = \Sigma^\mathcal{N}(\mathcal{T}) = b$ .

To see that  $\Sigma$  is unique, consider a countable normal iteration tree  $\mathcal{T}$  above  $\delta$  on  $\mathcal{M}$  of limit length according to an  $\omega_1$ -iteration strategy  $\Sigma'$ . Let  $x \in \mathbb{R}$  code  $\mathcal{T}$ . Then  $\Sigma'(\mathcal{T}) = \Sigma^{M_n^\#(x)}(\mathcal{T}) = \Sigma(\mathcal{T})$ .  $\square$

2.2.4. *Tools for  $\omega_1$ -iterable premice.* In this section we adapt the tools for  $\omega_1 + 1$ -iterable premice from the previous sections to  $\omega_1$ -iterable premice.

**Lemma 2.42.** (*Comparison lemma*) *Assume  $M_n^\#(x)$  exists for every  $x \in \mathbb{R}$ . Let  $\mathcal{M}$  and  $\mathcal{N}$  be countable  $X$ -premise, where  $X$  is swo. Let  $\delta$  be a cutpoint of  $\mathcal{M}$  and  $\mathcal{N}$  such that both are  $\omega_1$ -iterable above  $\delta$  and  $\mathcal{M}|\delta = \mathcal{N}|\delta$ . Further suppose  $\mathcal{M}$  and  $\mathcal{N}$  are  $(n + 1)$ -small above  $\delta$  and  $\omega$ -sound above  $\delta$  with  $\rho_\omega(\mathcal{M}), \rho_\omega(\mathcal{N}) \leq \delta$ . Then there are countable iteration trees  $\mathcal{S}$  on  $\mathcal{M}$  and  $\mathcal{T}$  on  $\mathcal{N}$  with last models  $\mathcal{M}_\alpha^{\mathcal{S}}$  and  $\mathcal{M}_\beta^{\mathcal{T}}$  so that*

- (1)  $[0, \alpha]_{\mathcal{S}}$  does not drop in model or degree and  $\mathcal{M}_\alpha^{\mathcal{S}} \trianglelefteq \mathcal{M}_\beta^{\mathcal{T}}$ , or
- (2)  $[0, \beta]_{\mathcal{T}}$  does not drop in model or degree and  $\mathcal{M}_\beta^{\mathcal{T}} \trianglelefteq \mathcal{M}_\alpha^{\mathcal{S}}$ .

*Proof.*  $\mathcal{M}$  and  $\mathcal{N}$  are  $\omega_1 + 1$ -iterable in  $M_n^\#(x)$  by lemma 2.38 where  $x \in \mathbb{R}$  codes  $\mathcal{M}$  and  $\mathcal{N}$ . So we can coiterate them in  $M_n^\#(x)$  by lemma 2.23.  $\square$

A consequence is

**Lemma 2.43.** *Suppose  $M_n^\#(x)$  exists for every  $x \in \mathbb{R}$ . Let  $\mathcal{M}$  and  $\mathcal{N}$  be countable  $X$ -premise, where  $X$  is swo. Suppose  $\delta$  is a cutpoint of  $\mathcal{M}$  and  $\mathcal{N}$  such that both are  $\omega_1$ -iterable above  $\delta$  and  $\mathcal{M}|\delta = \mathcal{N}|\delta$ . Further suppose that both  $\mathcal{M}$  and  $\mathcal{N}$  are  $(n + 1)$ -small above  $\delta$  and  $\omega$ -sound above  $\delta$  with  $\rho_\omega(\mathcal{M}), \rho_\omega(\mathcal{N}) \leq \delta$ . Then  $\mathcal{M} \trianglelefteq \mathcal{N}$  or  $\mathcal{N} \trianglelefteq \mathcal{M}$ .*

We get a version of the genericity iteration for  $\omega_1$ -iterable premice:

**Lemma 2.44.** (*Genericity iteration*) *Assume  $M_n^\#(x)$  exists for every  $x \in \mathbb{R}$ . Let  $\mathcal{M}$  be a countable  $X$ -premouse, where  $X$  is swo, such that  $\delta$  is Woodin in  $\mathcal{M}$  and*

$\mathcal{M}$  is  $\omega_1$ -iterable above some  $\gamma < \delta$ . Further suppose  $\mathcal{M}$  is  $(n+1)$ -small above  $\gamma$  and  $\omega$ -sound above  $\gamma$  with  $\rho_\omega(\mathcal{M}) \leq \gamma$ . Then for each  $x \in \mathbb{R}$ , there is a countable iteration tree  $\mathcal{T}$  on  $\mathcal{M}$  with iteration map  $\pi$  and last model  $\mathcal{M}_\alpha^\mathcal{T}$  such that  $[0, \alpha]_\mathcal{T}$  does not drop in model and  $x$  is  $\mathbb{W}_{\pi(\delta)}^{\mathcal{M}_\alpha^\mathcal{T}}$ -generic over  $\mathcal{M}_\alpha^\mathcal{T}$ .

*Proof.* Apply lemma 2.32 inside  $M_n^\#(z)$  where  $z \in \mathbb{R}$  codes  $\mathcal{M}$  and  $x$ .  $\square$

While forcing is usually applied to models of ZF, we would like to use the forcing theorem for small forcing over relativized premeice in the next lemma. Let  $\mathcal{M}$  be a relativized premouse and  $\kappa$  the critical point of an extender on the  $\mathcal{M}$ -sequence. Note that the forcing relation for any partial order  $\mathbb{P} \in \mathcal{M}|\kappa$  is defined in  $\mathcal{M}|\kappa$  and the forcing theorem holds for  $\mathcal{M}|\kappa$ , since this is a model of ZF. We are only interested in formulas whose quantifiers range over a bounded subset of  $\mathcal{M}|\kappa$ , especially projective formulas. The forcing theorem holds for such formulas since the relevant names are in  $\mathcal{M}|\kappa$ .

A key property of  $M_n^\#$  is that it determines which  $\Sigma_{n+2}^1$  statements about reals in  $M_n^\#$  are true in  $V$ :

**Lemma 2.45.** *Let  $n \leq k < \omega$  and assume  $M_k^\#(x)$  exists for every  $x \in \mathbb{R}$ . Let  $\mathcal{M}$  be a countable  $(k+1)$ -small  $X$ -premouse with  $\rho_\omega(\mathcal{M}) \leq \gamma$  which is  $\omega_1$ -iterable above  $\gamma$ , where  $X$  is swo. Suppose that in  $\mathcal{M}$  there are  $n$  Woodin cardinals above  $\gamma$  and an extender above. Let  $\delta$  be the least Woodin cardinal above  $\gamma$  in  $\mathcal{M}$  if  $n \geq 1$ .*

- (1) *If  $n$  is even then  $\mathcal{M} \prec_{\Sigma_{n+2}^1} V$ , and*
- (2) *if  $n$  is odd then*

$$V \models \varphi(x) \Leftrightarrow \mathcal{M} \models \text{''}\exists p \in \mathbb{W}_\delta(p \Vdash_{\mathbb{W}_\delta} \varphi(\check{x})\text{''}$$

*for all  $\Sigma_{n+2}^1$  formulas  $\varphi$  and all  $x \in \mathbb{R} \cap \mathcal{M}$ .*

*Proof.* The proof is organized as an induction on  $n$ . For  $n = 0$  we iterate  $\mathcal{M}$  by an extender on the  $\mathcal{M}$ -sequence to a model of height  $\geq \omega_1$ . The conclusion follows from Shoenfield absoluteness, since this model has the same reals as  $\mathcal{M}$ .

**Case 1.**  $n$  is odd.

Suppose  $\varphi$  is a  $\Pi_{n+1}^1$  formula,  $x \in \mathbb{R}$ , and  $y \in \mathbb{R} \cap \mathcal{M}$  so that  $\varphi(x, y)$  holds. Do a genericity iteration on  $\mathcal{M}$  for  $x$  and let  $\pi : \mathcal{M} \rightarrow \mathcal{N}$  be the iteration map so that  $x$  is  $\mathbb{W}_{\pi(\delta)}$ -generic over  $\mathcal{N}$ . We get  $\mathcal{N}[x] \models \varphi(x)$  from the induction hypothesis. Hence

$$\mathcal{N} \models \text{''}\exists p \in \mathbb{W}_{\pi(\delta)} (p \Vdash_{\mathbb{W}_{\pi(\delta)}}^{\mathcal{N}} \exists x \varphi(x, \check{y}))\text{''}$$

and the claim follows from elementarity of  $\pi$ .

For the other direction suppose there is a condition  $p \in \mathbb{W}_{\delta}^{\mathcal{M}}$  which forces  $\exists x \varphi(x, \check{y})$  over  $\mathcal{M}$ . Let  $x$  be  $\mathbb{W}_{\delta} \restriction p$ -generic over  $\mathcal{M}$  in  $V$ . Then  $\varphi(x, y)$  holds by the induction hypothesis.

**Case 2.**  $n \geq 2$  is even.

Suppose  $\varphi$  is a  $\Pi_{n+1}^1$  formula and  $y \in \mathbb{R} \cap \mathcal{M}$  so that  $\exists x \varphi(x, y)$  holds. The assumptions imply  $\Pi_{n+1}^1$  uniformization via lemma 2.30. Let  $\psi$  be a  $\Pi_{n+1}^1$  formula and  $x \in \mathbb{R}$  so that  $\varphi(x, y)$  holds and  $x$  is unique with  $\psi(x, y)$ . We have to show that  $x \in \mathcal{M}$ .

Let  $\pi : \mathcal{M} \rightarrow \mathcal{N}$  be an iteration map so that  $x$  is  $\mathbb{W}_{\pi(\delta)}$ -generic over  $\mathcal{N}$ . Let  $\eta$  be the least Woodin cardinal above  $\delta$  in  $\mathcal{M}$ . We have

$$\exists p \in \mathbb{W}_{\pi(\eta)} (p \Vdash_{\mathbb{W}_{\pi(\eta)}}^{\mathcal{N}[x]} \psi(\check{x}, \check{y}))$$

by the induction hypothesis. So there is a condition  $q \in \mathbb{W}_{\pi(\delta)}$  which forces

$$\exists p \in \mathbb{W}_{\pi(\eta)} (p \Vdash_{\mathbb{W}_{\pi(\eta)}}^{\mathcal{N}[\tau]} \psi(\tau, \check{y}))$$

such that  $x$  is  $\mathbb{W}_{\pi(\delta)} \upharpoonright q$ -generic over  $\mathcal{N}$ , where  $\tau$  is a name for the  $\mathbb{W}_{\pi(\delta)}$ -generic real. Let  $z$  be  $\mathbb{W}_{\pi(\delta)}^{\mathcal{N}} \upharpoonright q$ -generic over  $\mathcal{N}[x]$  in  $V$ . Then  $\psi(z, y)$  holds and hence  $z = x$ . This implies that  $\mathbb{W}_{\pi(\delta)} \upharpoonright q$  is atomic and  $x \in \mathcal{N}$ . Then  $x \in \mathcal{M}$  since the iteration does not add reals.

For the other direction suppose  $\varphi$  is a  $\Pi_{n+1}^1$  formula and  $y$  a real in  $\mathcal{M}$  with

$$\mathcal{M} \models \text{''}\exists p \in \mathbb{W}_\delta(p \Vdash_{\mathbb{W}_\delta} \exists x \varphi(x, \check{y})\text{''}.$$

Then  $\varphi(x, y)$  holds by the induction hypothesis for any real  $x$  which witnesses this in a  $\mathbb{W}_\delta \upharpoonright p$ -generic extension of  $\mathcal{M}$ .

□

The previous lemma is also true if  $\mathcal{M}$  is uncountable. To show this one simply applies the lemma to a countable elementary substructure of  $\mathcal{M}$ . Note that the lemma also works for the forcing  $Col(\omega, \delta)$ . In fact this version of the lemma uses a weaker notion of iterability called  $n$ -iterability, see [31, definition 1.1] for a definition of  $n$ -iterability and [32, theorem 7.16] for the result.

If in the situation of the previous lemma there is an extra extender on top in  $\mathcal{M}$ , then  $\mathcal{M} \prec_{\Sigma_{n+2}^1} V$  holds for odd  $n$  as well:

**Lemma 2.46.** *Suppose  $n \leq k$  and  $M_k^\#(x)$  exists for every  $x \in \mathbb{R}$ . Suppose  $\mathcal{M}$  is a countable  $\omega$ -sound  $(k+1)$ -small  $X$ -premouse which is  $\omega_1$ -iterable above  $\delta$  with  $\rho_\omega(\mathcal{M}) \leq \delta$ , where  $X$  is swo. Suppose there are  $n$  Woodin cardinals above  $\delta$  in  $\mathcal{M}$  and at least two total extenders above. Then  $M_n^\#(x)$  is unique for every  $x \in \mathbb{R} \cap \mathcal{M}$  and is calculated correctly by  $\mathcal{M}$ .*

*Proof.* Do an  $L[\vec{E}]$ -construction over  $x$  in  $\mathcal{M}$ . Then the  $M_n^\#(x)$  of both  $\mathcal{M}$  and  $V$  occurs in the construction when one forms the core of an  $x$ -premouse with an extender above  $n$  Woodin cardinals for the first time. □

**Lemma 2.47.** *Suppose  $n \leq k$  and  $M_k^\#(x)$  exists for every  $x \in \mathbb{R}$ . Suppose  $\mathcal{M}$  is a countable  $\omega$ -sound  $(k+1)$ -small  $X$ -premouse which is  $\omega_1$ -iterable above  $\delta$  with  $\rho_\omega(\mathcal{M}) \leq \delta$ , where  $X$  is swo. Suppose there are  $n$  Woodin cardinals above  $\delta$  in  $\mathcal{M}$  and at least two total extenders above. Then  $\mathcal{M} \prec_{\Sigma_{n+2}^1} V$ .*

*Proof.* For  $n$  even this is true by lemma 2.45. Suppose  $n$  is odd and  $\varphi$  is a  $\Sigma_{n+2}^1$  formula. Let  $x \in \mathbb{R}$  and let  $\delta$  be the least Woodin cardinal in  $M_n^\#(x)$ . We know that  $\mathcal{M}$  computes  $M_n^\#(x)$  correctly by the previous lemma. So  $\varphi(x)$  holds if and only if

$$\exists p \in \mathbb{W}_\delta^{M_n^\#(x)} (p \Vdash_{\mathbb{W}_\delta^{M_n^\#(x)}} \varphi(x))$$

holds if and only if  $\mathcal{M} \models \varphi(x)$  by lemma 2.45. □

The next two lemmas show that statements about  $M_{2n}^\#(x)$  and  $M_{2n}^\dagger(x)$  are projective.

**Lemma 2.48.** *Suppose  $M_n^\#(x)$  exists for every  $x \in \mathbb{R}$ . Let  $\mathcal{M}$  and  $\mathcal{N}$  countable  $(n+1)$ -small  $X$ -premouse with the same first order properties which are  $\omega$ -sound above  $\delta$  with  $\rho_\omega(\mathcal{M}) \leq \delta$  and  $\rho_\omega(\mathcal{N}) \leq \delta$ , where  $X$  is swo. Suppose  $\mathcal{M}$  is  $\omega_1$ -iterable above  $\delta$  and  $\mathcal{N}$  is  $\Pi_{n+1}$ -iterable. Then  $\mathcal{M} = \mathcal{N}$ .*

*Proof.* The proof is organized as an induction. We sketch the proof for odd  $n$  following the proof of [40, lemma 2.2]. This proof has to be slightly modified since we don't have large cardinals in  $V$ . The case for even  $n$  can be similarly derived from the proof of [40, lemma 2.2]. The difference between the odd and even cases lies in the weak iteration game from [40].

Let  $\mathcal{P} := M_n^\#(x)$  where  $x \in \mathbb{R}$  codes  $\mathcal{M}$  and  $\mathcal{N}$ . Let further  $g$  be a  $Col(\omega, \omega_1^{\mathcal{P}})$ -generic filter over  $\mathcal{N}$  and define  $\mathcal{R} := \mathcal{P}[g]$ . Then  $\mathcal{M}$  is  $\omega_1 + 1$ -iterable in  $\mathcal{P}$  and in  $\mathcal{R}$  by lemma 2.38. Note that  $\Pi_{n+1}$ -iterability is  $\Pi_{n+2}^1$  in the codes. Since there are  $n$  Woodin cardinals in  $\mathcal{P}$  and in  $\mathcal{R}$ , it follows from lemma 2.47 that  $\mathcal{N}$  is  $\Pi_{n+1}$ -iterable in  $\mathcal{N}$  and in  $\mathcal{P}$ .

We define coiterations of  $\mathcal{M}$  and  $\mathcal{N}$  in both  $\mathcal{P}$  and  $\mathcal{R}$ . For iteration trees on  $\mathcal{M}$  of limit length choose the unique branch with a  $\mathcal{Q}$ -structure. For iteration trees  $\mathcal{T}$  on  $\mathcal{N}$  of limit length choose a cofinal branch with a wellfounded  $\Pi_n$ -iterable model. The winning position for player 2 in the weak iteration game  $\mathcal{I}(\mathcal{N}, \delta, n+1)$  produces such a branch. A coiteration argument shows that the branch is unique. Note that the coiteration is possible by the induction hypothesis.

One can show that the same branches are chosen in the coiterations in  $\mathcal{P}$  and  $\mathcal{R}$  since the forcing  $Col(\omega, \omega_1^{\mathcal{P}})$  is small. So the coiteration in  $\mathcal{P}$  is an initial segment of the coiteration in  $\mathcal{R}$ . Hence there is at least one cofinal branch in the coiteration in  $\mathcal{P}$ . This is an element of  $\mathcal{P}$  by homogeneity of  $Col(\omega, \omega_1^{\mathcal{P}})$ . Now the argument from the proof of the comparison lemma shows that the coiteration terminates after countably many steps. It follows that  $\mathcal{M} = \mathcal{N}$ .  $\square$

**Lemma 2.49.** *Suppose  $M_n^\#(x)$  exists for every  $x \in \mathbb{R}$ . Then  $M_n^\#(x)$  and  $M_n^\dagger(x)$  are coded by  $\Pi_{n+2}^1(x)$  singletons for each  $x \in \mathbb{R}$ .*

*Proof.* Let  $f : \omega \rightarrow M_n^\#(x)$  be the canonical  $\Sigma_1$ -definable surjection over  $M_n^\#(x)$ .

Let

$$(k, m) \in z :\Leftrightarrow f(k) \in f(m)$$

for  $k, m < \omega$ , so that  $(\omega, z)$  is isomorphic to  $(M_n^\#(x), \in)$ . Hence  $z \subseteq \omega \times \omega$  is the unique set which codes  $M_n^\#(x)$  and which computes itself via the canonical  $\Sigma_1$ -definable surjection computed in its transitive collapse. Now the set of  $(x, y) \in \mathbb{R}^2$  so that  $x$  codes  $M_n^\#(y)$  is a  $\Pi_{n+2}^1(x, y)$  set by the previous lemma, since for sets of reals  $\Pi_{n+1}^{HC}$  is equivalent to  $\Pi_{n+2}^1$ . Thus  $z$  is a  $\Pi_{n+2}^1(x)$  singleton. The same works for  $M_n^\dagger$ .  $\square$

## 3. LIFTING THIN EQUIVALENCE RELATIONS TO FORCING EXTENSIONS

In the situation where  $E$  is a provably  $\Delta_{n+1}^1$  equivalence relation and the forcing  $\mathbb{P}$  preserves  $\Sigma_n^1$  truth, we ask whether forcing with  $\mathbb{P}$  introduces any new equivalence classes of  $E$ . In the first section we show generic  $\Sigma_{n+3}^1$  absoluteness for reasonable forcing  $\mathbb{P}$  of size  $\kappa$  and that  $\mathbb{P}$  does not add equivalence classes to thin provably  $\Delta_{n+3}^1$  equivalence relations, assuming that  $M_n^\#(X)$  exists for every self-wellordered set  $X \in H_{\kappa^+}$ , based on an idea of Foreman and Magidor [4, section 3]. In the second section we derive analogous results for  $\Sigma_2^1$  c.c.c. forcing from projective determinacy. We further show that generic  $\Sigma_{n+1}^1$  Cohen absoluteness implies that Cohen forcing does not add equivalence classes to  $< \omega - \Pi_n^1$  prewellorders. We work in  $\text{ZF} + \text{DC}$ .

**3.1. Reasonable forcing.** We work with a weaker version of the notion of proper forcing called reasonable forcing, introduced by Foreman and Magidor [4].

**Definition 3.1.** *Let  $\mathbb{P}$  be a partial order and  $p \in \mathbb{P}$ .*

- (1) *Suppose  $N$  is a set with  $p \in N$ . Then  $p$  is called  $(N, \mathbb{P})$ -generic if for every maximal antichain  $A \subseteq \mathbb{P}$  with  $A \in N$  the set  $A \cap N$  is predense below  $p$ .*
- (2)  *$\mathbb{P}$  is called reasonable if for all  $q \in \mathbb{P}$  and for some (for all) regular  $\lambda \geq (2^{2^{\overline{\mathbb{P}}}})^+$  there exist a countable elementary substructure  $N \prec H_\lambda$  with  $q, \mathbb{P} \in N$  and an  $(N, \mathbb{P})$ -generic condition  $r \leq q$ .*

Here  $H_\lambda$  can equivalently be replaced by  $V_\lambda$ . Let  $\mathcal{P}_\kappa(\lambda) := \{X \subseteq \lambda : \overline{X} < \kappa\}$  for  $\kappa, \lambda \in \text{Ord}$ . By standard proper forcing arguments we have

**Lemma 3.2.** *(Foreman and Magidor [4]) A forcing  $\mathbb{P}$  is reasonable if and only if  $\mathcal{P}_{\omega_1}^V(\alpha)$  is stationary in  $\mathcal{P}_{\omega_1}^{V^{\mathbb{P}}}(\alpha)$  for every ordinal  $\alpha$ .*

3.1.1. *Absoluteness of  $M_n^\#$ .* We will use

**Lemma 3.3.** *Suppose  $M_n^\#(X)$  exists for all swo  $X \in H_\kappa$ , where  $\kappa$  is an uncountable cardinal. Then  $M_n^\#(X)$  is  $\kappa$ -iterable for each swo  $X \in H_\kappa$ .*

*Proof.* We can assume that  $H_\kappa$  is closed under the relevant  $\mathcal{Q}$ -structures by the induction hypothesis. A reflection argument then shows that  $M_n^\#(X)$  is  $\kappa$ -iterable.  $\square$

The results in this section are based on the absoluteness of  $M_n^\#(X)$ :

**Lemma 3.4.** *(Folklore) Let  $\mathbb{P}$  be a forcing of size  $\kappa$ , where  $\kappa$  is an infinite cardinal. Suppose  $M_n^\#(X)$  exists for every swo  $X \in H_{\kappa^+}$ . Then for every  $\mathbb{P}$ -generic filter  $G$  over  $V$*

- (1)  $M_n^\#(X)$  is normally  $\kappa^+$ -iterable in  $V[G]$  via  $\Sigma$  for every swo  $X \in H_{\kappa^+}$ ,
- (2)  $V[G] \models$  " $M_n^\#(X)$  exists for every swo  $X \in H_{\kappa^+}$  and is normally  $\kappa^+$ -iterable", and
- (3) suppose
  - (a)  $H \prec V_\eta$  is a countable substructure with  $\mathbb{P} \in H$  where  $\eta$  is a large limit ordinal,
  - (b)  $\bar{H}$  is the transitive collapse of  $H$  with uncollapsing map  $\pi : \bar{H} \rightarrow H$  and  $\pi(\bar{\mathbb{P}}) = \mathbb{P}$ ,  $\pi(\bar{\kappa}) = \kappa$ , and
  - (c)  $g$  is a  $\bar{\mathbb{P}}$ -generic filter over  $\bar{H}$  in  $V$ ,  
 then  $M_n^\#(X)$  exists in  $\bar{H}[g]$  for each swo  $X \in H_{\bar{\kappa}^+}^{\bar{H}[g]}$  and is normally  $\kappa^+$ -iterable via  $\Sigma$  in both  $\bar{H}[g]$  and  $V$ .

*Proof.* The proof works by induction on  $n$ . We get uniqueness of  $M_n^\#(X)$  for  $X \in H_{\kappa^+}$  by the argument in lemma 2.38.

1. In the case  $n = 0$  the claim holds since all iterations are linear. Let  $n \geq 1$  and suppose  $X \in H_{\kappa^+}$  is swo. Let  $\mathcal{M} := M_n^\#(X)$ . We have to show that  $\mathcal{M}$  is normally  $\kappa^+$ -iterable in  $V[G]$  via  $\Sigma$ .

Suppose not. Then in  $V[G]$  there is a normal iteration tree  $\mathcal{T}$  on  $\mathcal{M}$  of length  $< \kappa^+$  which witnesses that  $\mathcal{M}$  is not normally  $\kappa^+$ -iterable via  $\Sigma$ . I.e.  $\mathcal{T}$  is according to  $\Sigma$  and  $\mathcal{M}_{\Sigma(\mathcal{T})}$  is ill-founded or  $\Sigma(\mathcal{T})$  is undefined. Let  $\dot{\mathcal{T}}$  be a  $\mathbb{P}$ -name and  $p \in \mathbb{P}$  a condition with

$$p \Vdash \dot{\mathcal{T}} \text{ witnesses that } \check{\mathcal{M}} \text{ is not } \kappa^+ \text{-iterable via } \Sigma''.$$

Now let  $H \prec V_\eta$  be a countable substructure with  $p, \mathbb{P}, \mathcal{M}, \dot{\mathcal{T}} \in H$  for some large limit ordinal  $\eta$ . Let  $\bar{H}$  be the transitive collapse of  $H$  with uncollapsing map  $\pi : \bar{H} \rightarrow H$  and  $\pi(\bar{p}) = p$ ,  $\pi(\bar{\mathbb{P}}) = \mathbb{P}$ ,  $\pi(\bar{\mathcal{T}}) = \dot{\mathcal{T}}$ ,  $\pi(\bar{\mathcal{M}}) = \mathcal{M}$ . Then  $\bar{\mathcal{M}}$  is  $\kappa^+$ -iterable in  $V$  since  $\pi \upharpoonright \bar{\mathcal{M}} : \bar{\mathcal{M}} \rightarrow \mathcal{M}$  is an elementary embedding.

Let  $g$  be a  $\bar{\mathbb{P}}$ -generic filter over  $\bar{H}$  in  $V$  with  $p \in g$ . Then

$$\bar{H}[g] \models \bar{\mathcal{T}}^g \text{ witnesses that } \mathcal{M} \text{ is not } \kappa^+ \text{-iterable via } \Sigma''.$$

Let  $\alpha < lh(\bar{\mathcal{T}}^g)$ .  $M_{n-1}^\#(\mathcal{M}(\bar{\mathcal{T}}^g \upharpoonright \alpha))$  exists in  $\bar{H}[g]$  and is  $\kappa^+$ -iterable via  $\Sigma$  in both  $\bar{H}[g]$  and  $V$  by the induction hypothesis 3.

Let  $\mathcal{Q}(\bar{\mathcal{T}}^g \upharpoonright \alpha)$  denote the  $\mathcal{Q}$ -structure for  $\bar{\mathcal{T}}^g \upharpoonright \alpha$  in  $\bar{H}[g]$ . We can compare  $\mathcal{Q}(\bar{\mathcal{T}}^g \upharpoonright \alpha)$  and  $M_{n-1}^\#(\mathcal{M}(\bar{\mathcal{T}}^g \upharpoonright \alpha))$  in  $\bar{H}[g]$  by lemma 2.38. Hence

$$\mathcal{Q}(\bar{\mathcal{T}}^g \upharpoonright \alpha) \trianglelefteq M_{n-1}^\#(\mathcal{M}(\bar{\mathcal{T}}^g \upharpoonright \alpha)).$$

So  $\bar{\mathcal{T}}^g$  is according to  $\Sigma$  in both  $\bar{H}[g]$  and  $V$ .

Let  $b := \Sigma^V(\bar{\mathcal{T}}^g)$ . We have to show that  $b \in \bar{H}[g]$ . Let  $g'$  be  $Col(\omega, lh(\bar{\mathcal{T}}^g))$ -generic over  $\bar{H}[g]$  where  $lh(\bar{\mathcal{T}}^g)$  is the length of  $\bar{\mathcal{T}}^g$ . Now  $\mathcal{Q}(\bar{\mathcal{T}}^g) \in \bar{H}[g]$  since

$$\mathcal{Q}(\bar{\mathcal{T}}^g) \trianglelefteq M_{n-1}^\#(\mathcal{M}(\bar{\mathcal{T}}^g)).$$

One can build a tree in  $\bar{H}[g][g']$  searching for a cofinal branch  $b' \subseteq \bar{\mathcal{T}}^g$  with  $\mathcal{Q}(\bar{\mathcal{T}}^g) \trianglelefteq \mathcal{M}_{b'}^{\bar{\mathcal{T}}^g}$ . Since  $b$  is such a branch in  $V$ , there is a cofinal branch  $b' \subseteq \bar{\mathcal{T}}$

in  $\bar{H}[g][g']$  with  $\mathcal{Q}(\bar{T}^g) \trianglelefteq \mathcal{M}_{b'}^{\bar{T}}$  by absoluteness of wellfoundedness. But there can be only one such branch by the argument in [42, corollary 6.14]. So  $b = b'$  and  $b' \in \bar{H}[g]$  by homogeneity of  $Col(\omega, \bar{T}^g)$ . Hence

$$\Sigma^V(\bar{T}^g) = b = \Sigma^{\bar{H}[g]}(\bar{T}^g),$$

contradicting the assumption on  $\bar{T}^g$ .

2. Suppose  $X \in H_{\kappa^+}^{V[G]}$  is swo. Let's code  $X$  by a subset of  $\kappa$  and choose a nice  $\mathbb{P}$ -name for this set. So there is a  $\mathbb{P}$ -name  $\tau \in H_{\kappa^+}$  with  $\tau^G = X$ . We can assume that  $(\mathbb{P}, \tau)$  is swo; otherwise we work with a swo set in  $H_{\kappa^+}$  coding  $\mathbb{P}$  and  $\tau$ . Now  $G' := G \cap M_n^\#(\mathbb{P}, \tau)$  is  $\mathbb{P}$ -generic over  $M_n^\#(\mathbb{P}, \tau)$ , since  $G$  is  $\mathbb{P}$ -generic over  $V$ . Moreover

$$x = \tau^{G'} \in M_n^\#(\mathbb{P}, \tau)[G'].$$

Then  $M_n^\#(\mathbb{P}, \tau)$  is normally  $\kappa^+$ -iterable via  $\Sigma$  in  $V[G]$  by 1. Since  $\mathbb{P}$  is small compared to the critical points of the extenders on the  $M_n^\#(\mathbb{P}, \tau)$  sequence,  $M_n^\#(\mathbb{P}, \tau)[G']$  is normally  $\kappa^+$ -iterable via  $\Sigma$  in  $V[G]$  as well. Let  $F$  be the top extender of  $M_n^\#(\mathbb{P}, \tau)[G']$  and  $\kappa := crit(F)$ . Do an  $L[\vec{E}]$  construction over  $X$  in  $M_n^\#(\mathbb{P}, \tau)[G'] \upharpoonright \kappa$ . It follows from the argument of the commutativity lemma [5, lemma 3.2] that the premouse obtained by extending the  $L[\vec{E}]$  model with the restriction of  $F$  is normally  $\kappa^+$ -iterable via  $\Sigma$ . Hence this is  $M_n^\#(X)$  in  $V[G]$ .

3. Suppose  $X \in H_{\kappa^+}^{\bar{H}[g]}$  is swo. Let  $\bar{\tau} \in H_{\kappa^+}$  be a  $\bar{\mathbb{P}}$ -name for  $X$  and let  $\tau := \pi(\bar{\tau})$ . Let's assume that  $(\bar{\mathbb{P}}, \bar{\tau})$  is swo. Then  $M_n^\#(\bar{\mathbb{P}}, \bar{\tau})^{\bar{H}}$  and  $M_n^\#(\mathbb{P}, \tau)$  exist and are normally  $\kappa^+$ -iterable via  $\Sigma$  in  $\bar{H}$  and  $V$  respectively and

$$\pi(M_n^\#(\bar{\mathbb{P}}, \bar{\tau})^{\bar{H}}) = M_n^\#(\mathbb{P}, \tau).$$

Then  $M_n^\#(\bar{\mathbb{P}}, \bar{\tau})^{\bar{H}}$  is normally  $\kappa^+$ -iterable via  $\Sigma$  in  $\bar{H}[g]$  by 1.

Let  $g' := g \cap M_n^\#(\bar{\mathbb{P}}, \bar{\tau})^{\bar{H}}$ . Then  $g'$  is a  $\bar{\mathbb{P}}$ -generic filter over  $M_n^\#(\bar{\mathbb{P}}, \bar{\tau})^{\bar{H}}$  and  $x = \bar{\tau}^{g'} \in M_n^\#(\bar{\mathbb{P}}, \bar{\tau})^{\bar{H}}[g']$ . Now  $M_n^\#(\bar{\mathbb{P}}, \bar{\tau})^{\bar{H}}[g']$  is normally  $\kappa^+$ -iterable via  $\Sigma$  in  $\bar{H}[g]$ , since  $\bar{\mathbb{P}}$  is small compared to the critical points of the extenders on the  $M_n^\#(\bar{\mathbb{P}}, \bar{\tau})^{\bar{H}}$  sequence. Moreover  $M_n^\#(\bar{\mathbb{P}}, \bar{\tau})^{\bar{H}}$  is normally  $\kappa^+$ -iterable via  $\Sigma$  in  $V$ , since

$$\pi \upharpoonright M_n^\#(\bar{\mathbb{P}}, \bar{\tau})^{\bar{H}} : M_n^\#(\bar{\mathbb{P}}, \bar{\tau})^{\bar{H}} \rightarrow M_n^\#(\mathbb{P}, \tau)$$

is elementary. Hence  $M_n^\#(\bar{\mathbb{P}}, \bar{\tau})^{\bar{H}}[g']$  is normally  $\kappa^+$ -iterable via  $\Sigma$  in  $V$  as well. As in part 2 we can build a model in  $M_n^\#(\bar{\mathbb{P}}, \bar{\tau})$  via an  $L[\vec{E}]$ -construction over  $X$  which is the  $M_n^\#(X)$  of both  $\bar{H}[g]$  and  $V$ .  $\square$

Note that the lemma works for  $M_n^\dagger$  with the same proof.

3.1.2. *Absoluteness of equivalence classes.* We will need a direct consequence of lemma 2.45:

**Lemma 3.5.** *Suppose  $M$  is a transitive model of ZF which computes  $M_n^\#(x)$  correctly for every  $x \in \mathbb{R} \cap M$ . Then  $M \prec_{\Sigma_{n+2}^1} V$ .*

*Proof.* Note that in this situation  $M_n^\#(x)$  is unique by lemma 2.38.  $M_n^\#(x)$  computes the truth value of  $\Sigma_{n+2}^1$  statements by lemma 2.45.  $\square$

**Definition 3.6.**  $\Sigma_n^1$ -*absoluteness holds for a partial order  $\mathbb{P}$  if  $V \prec_{\Sigma_n^1} V[G]$  for any  $\mathbb{P}$ -generic filter  $G$  over  $V$ .*

**Lemma 3.7.** *(Martin, Solovay, Schindler) Suppose  $M_n^\#(X)$  exists for every swo  $X \in H_{\kappa^+}$ , where  $\kappa$  is an infinite cardinal. Then  $\Sigma_{n+3}^1$ -absoluteness holds for every forcing of size  $\kappa$ .*

*Proof.* We follow the proof of [35, theorem 1]. Suppose  $\exists x \varphi(x, y)$  holds in some  $\mathbb{P}$ -generic extension of  $V$ , where  $\varphi$  is a  $\Pi_{n+2}^1$  formula and  $y \in \mathbb{R}$ . Let  $\tau$  be a nice  $\mathbb{P}$ -name for a real and  $p \in \mathbb{P}$  a condition with  $p \Vdash_{\mathbb{P}} \varphi(\tau, \check{y})$ . We can assume that  $(\mathbb{P}, \tau)$  is swo. Then  $M_n^\#(\mathbb{P}, \tau)$  exists since  $\mathbb{P}, \tau \in H_{\kappa^+}$ .

Consider the tree  $T$  in  $V$  searching for a 5-tuple  $(\mathcal{M}, \pi, \bar{\mathbb{P}}, g, x)$  such that

- (1)  $\mathcal{M}$  is a countable premouse with  $y \in \mathcal{M}$ ,
- (2)  $\pi : \mathcal{M} \rightarrow M_n^\#(\mathbb{P}, \tau)$  is elementary with  $\pi(\bar{\mathbb{P}}) = \mathbb{P}$ ,
- (3)  $g$  is  $\bar{\mathbb{P}}$ -generic over  $\mathcal{M}$ , and
- (4)  $x$  is a real in  $\mathcal{M}[g]$  such that  $\mathcal{M}[g] \models \varphi(x, y)$  if  $n$  is even, and  $\Vdash_{Col(\omega, \delta)}^{\mathcal{M}[g]} \varphi(\check{x}, \check{y})$  if  $n$  is odd, where  $\delta$  is the least Woodin cardinal in  $\mathcal{M}[g]$ .

A branch in this tree defines a complete theory so that every existential statement in the theory is witnessed by a constant, giving rise to a model  $\mathcal{M}[g]$ , as well as a set of finite partial  $\in$ -isomorphisms whose union is an elementary map  $\pi : \mathcal{M} \rightarrow M_n^\#(\mathbb{P}, \tau)$ , witnessing that  $\mathcal{M}$  is wellfounded.

Now let  $G$  be  $\mathbb{P} \upharpoonright p$ -generic over  $V$ . Then  $g := G \cap M_n^\#(\mathbb{P}, \tau)$  is  $\mathbb{P}$ -generic over  $M_n^\#(\mathbb{P}, \tau)$  and we have  $x := \tau^g \in M_n^\#(\mathbb{P}, \tau)[g]$ . Since  $M_n^\#(\mathbb{P}, \tau)[g]$  is  $\kappa^+$ -iterable in  $V[G]$  by lemma 3.4, the collapse of a countable elementary substructure of  $M_n^\#(\mathbb{P}, \tau)[g]$  witnesses that  $T$  has a branch in  $V[G]$  by lemma 2.45. Then  $T$  is also ill-founded in  $V$  and hence  $V \models \exists x \varphi(x, y)$ .  $\square$

For any set  $E$  with a fixed definition we always write  $E$  for the corresponding set in any forcing extension. If further  $\mathbb{P}$  is a forcing and  $\tau$  is a  $\mathbb{P}$ -name, then in any  $\mathbb{P} \times \mathbb{P}$ -generic extension  $\tau$  defines two objects via the two  $\mathbb{P}$ -generic filters. We write  $\tau$  and  $\tau'$  for  $\mathbb{P} \times \mathbb{P}$ -names for these objects.

The idea for the next lemma and the next theorem comes from [4, theorem 3.4].

**Lemma 3.8.** *Let  $E$  be a thin  $\Pi_{n+3}^1$  equivalence relation. Suppose  $\mathbb{P}$  is a forcing of size  $\kappa$  and  $M_n^\#(X)$  exists for every swo  $X \in H_{\kappa^+}$ , where  $\kappa$  is an infinite cardinal. Let  $\tau$  be a  $\mathbb{P}$ -name for a real. Then the set*

$$D := \{p \in \mathbb{P} : (p, p) \Vdash_{\mathbb{P} \times \mathbb{P}} \tau E \tau'\}$$

*is dense.*

*Proof.* Let  $a \in \mathbb{R}$  so that  $E$  is  $\Pi_{n+3}^1(a)$ . Suppose  $D$  is not dense. Then there is a condition  $p \in \mathbb{P}$  so that for every  $q \leq p$  there are  $r, s \leq q$  with

$$(r, s) \Vdash_{\mathbb{P} \times \mathbb{P}} \neg \tau E \tau'.$$

Let  $\lambda$  be a large limit ordinal and  $H \prec V_\lambda$  a countable elementary substructure with  $a, \mathbb{P}, p, \tau, \tau' \in H$ . Let  $\bar{H}$  be the transitive collapse with uncollapsing map  $\pi : \bar{H} \rightarrow H$  and  $\pi(\bar{\mathbb{P}}) = \mathbb{P}$ ,  $\pi(\bar{p}) = p$ ,  $\pi(\bar{\tau}) = \tau$ , and  $\pi(\bar{\tau}') = \tau'$ .

Let  $(D_n : n \in \omega)$  enumerate the open dense subsets in  $\bar{H}$  of  $\bar{\mathbb{P}} \times \bar{\mathbb{P}}$ . We construct a family of conditions  $(p_s : s \in 2^{<\omega})$  in  $\bar{\mathbb{P}}$  such that

- (1)  $p_\emptyset = \bar{p}$ ,
- (2)  $p_s \leq p_t$  if  $t \subseteq s$ ,
- (3)  $(p_{s \smallfrown 0}, p_{s \smallfrown 1}) \Vdash_{\bar{\mathbb{P}} \times \bar{\mathbb{P}}} \neg \bar{\tau} E \bar{\tau}'$ ,
- (4)  $p_s$  decides  $\bar{\tau} \upharpoonright lh(s)$ , and
- (5)  $(p_s, p_t) \in D_0 \cap D_1 \cap \dots \cap D_i$  if  $s, t \in {}^i 2$  and  $s \neq t$

for all  $s, t \in 2^{<\omega}$ . When  $p_s$  is defined we choose as candidates for  $p_{s \smallfrown 0}$  and  $p_{s \smallfrown 1}$  conditions  $r, s \leq p_s$  with  $(r, s) \Vdash_{\mathbb{P} \times \mathbb{P}} \neg \tau E \tau'$ . Then one enumerates the pairs of these conditions for all  $s$  of fixed length and extends the conditions to satisfy properties 4 and 5.

Now let

$$g_x := \{q \in \bar{\mathbb{P}} : \exists n \in \omega p_{x \upharpoonright n} \leq q\}$$

for each  $x \in 2^{<\omega}$ . Then  $g_x$  and  $g_y$  are mutually  $\bar{\mathbb{P}}$ -generic over  $\bar{H}$  for  $x, y \in 2^{<\omega}$  with  $x \neq y$ , so

$$\bar{H}[g_x, g_y] \Vdash \neg \bar{\tau}^{g_x} E \bar{\tau}^{g_y}$$

by property 3. Since  $\bar{H}[g_x, g_y]$  computes  $M_n^\#(z)$  correctly for each  $z \in \mathbb{R} \cap \bar{H}[g_x, g_y]$  by 3 of lemma 3.4 we have  $\bar{H}[g_x, g_y] \prec_{\Sigma_{n+2}^1} V$  by the previous lemma. Since  $E$  is

$\Pi_{n+3}^1(a)$  this implies

$$V \models \neg \bar{\tau}^{g_x} E \bar{\tau}^{g_y}$$

for  $x \neq y$ . Since  $\bar{\tau}^{g_x}$  depends continuously on  $x$ , we get a perfect set of pairwise inequivalent reals in  $V$ . This would contradict that  $E$  is thin.  $\square$

A set is called provably  $\Delta_n^1(a)$  for  $a \in \mathbb{R}$  if there are  $\Sigma_n^1$  and  $\Pi_n^1$  formulas  $\varphi$  and  $\psi$  such that both  $\varphi(\cdot, a)$  and  $\psi(\cdot, a)$  define the set, and ZFC proves  $\forall x, y(\varphi(x, y) \leftrightarrow \psi(x, y))$ . For our purposes it will be sufficient to know that  $\forall x(\varphi(x, a) \leftrightarrow \psi(x, a))$  holds in all generic extensions of sufficiently elementary substructures of  $V$  containing  $a$ .

**Theorem 3.9.** *Let  $\mathbb{P}$  be a reasonable forcing of size  $\kappa$ , where  $\kappa$  is an infinite cardinal. Suppose  $M_n^\#(X)$  exists for every  $X \in H_{\kappa^+}$ . Then  $\mathbb{P}$  does not add equivalence classes to thin provably  $\Delta_{n+3}^1$  equivalence relations.*

*Proof.* Suppose  $E$  is a thin provably  $\Delta_{n+3}^1(a)$  equivalence relation where  $a \in \mathbb{R}$ . We use  $E$  to denote the set given by the same  $\Sigma_{n+3}^1(a)$  and  $\Pi_{n+3}^1(a)$  formulas in any  $\mathbb{P}$ -generic extension. This is an equivalence relation by lemma 3.7.

Suppose  $\tau$  is a  $\mathbb{P}$ -name for a real and  $p \in \mathbb{P}$  is a condition such that for every  $x \in \mathbb{R}$  we have  $p \Vdash_{\mathbb{P}} \neg \check{x} E \tau$ . Let  $q \leq p$  be a condition with

$$(q, q) \Vdash_{\mathbb{P} \times \mathbb{P}} \tau E \tau'$$

by the previous lemma. Since  $\mathbb{P}$  is reasonable there is a large regular  $\lambda$  and a countable substructure  $H \prec V_\eta$  with  $a, \mathbb{P}, q, \tau, \tau' \in H$  such that there is an  $(H, \mathbb{P})$ -generic condition  $r \leq q$ . Let  $\bar{H}$  be the transitive collapse of  $H$  with uncollapsing map  $\pi : \bar{H} \rightarrow H$  and  $\pi(\bar{\mathbb{P}}) = \mathbb{P}$ ,  $\pi(\bar{q}) = q$ ,  $\pi(\bar{\tau}) = \tau$ , and  $\pi(\bar{\tau}') = \tau'$ .

Let  $g_0$  be  $\bar{\mathbb{P}} \upharpoonright \bar{q}$ -generic over  $\bar{H}$  in  $V$ . Further let  $G$  be  $\mathbb{P}$ -generic over  $V$  with  $r \in G$  and define  $g_1 := \pi^{-1} G$ . Then  $\bar{q} \in g_1$ . As in the proof of lemma 3.7  $g_1$  is  $\bar{\mathbb{P}}$ -generic over  $\bar{H}$ .

Now let  $h$  be  $\bar{\mathbb{P}} \upharpoonright \bar{q}$ -generic over both  $\bar{H}[g_0]$  and  $\bar{H}[g_1]$  in  $V$ . Let  $x_0 := \bar{\tau}^{g_0}$ ,  $x_1 := \bar{\tau}^{g_1}$ , and  $y := \bar{\tau}^h$ . Then  $x_1 = \tau^G$ . Since  $(\bar{q}, \bar{q}) \Vdash_{\bar{\mathbb{P}} \times \bar{\mathbb{P}}}^{\bar{H}} \bar{\tau} E \bar{\tau}'$  we have

$$\bar{H}[g_0, h] \models x_0 E y$$

and

$$\bar{H}[g_1, h] \models x_1 E y.$$

As in the proof of lemma 3.7  $\bar{H}[g_i, h]$  computes  $M_n^\#(x)$  correctly for every  $x \in \mathbb{R} \cap \bar{H}[g_i, h]$ . Hence  $\bar{H}[g_i, h] \prec_{\Sigma_{n+2}^1} V$  by lemma 3.5.

Since  $E$  is provably  $\Delta_{n+3}^1(a)$ , this shows that  $x_0$ ,  $x_1$ , and  $y$  are equivalent with respect to  $E$ . But  $x_0 \in V$  and on the other hand we assumed that  $x_1$  is in a new equivalence class in  $V[G]$ , which is contradictory.  $\square$

**3.2. Projective c.c.c. forcing.** In this section we present versions of the results in the previous section for  $\Sigma_2^1$  c.c.c. forcing.

**3.2.1. Absoluteness of equivalence classes.** We use the notion of projective forcing from [1].

**Definition 3.10.** *Let  $a \in \mathbb{R}$ . A partially ordered set  $\mathbb{P} \subseteq \mathbb{R}$  is called a  $\Sigma_n^1(a)$  forcing if the partial order  $\leq$  and the incompatibility relation  $\perp$  are  $\Sigma_n^1(a)$  subsets of  $\mathbb{R}^2$ .*

For example Cohen forcing, random forcing, and Amoeba forcing are  $\Sigma_1^1$  c.c.c. forcings.

**Lemma 3.11.** *Let  $\mathbb{P}$  be a  $\Sigma_2^1(a)$  c.c.c. forcing where  $a \in \mathbb{R}$ . Suppose  $M_n^\#(x)$  exists for every  $x \in \mathbb{R}$ . Then for every  $\mathbb{P}$ -generic filter  $G$  over  $V$*

- (1)  $M_n^\#(x)$  is normally  $\omega_1$ -iterable in  $V[G]$  via  $\Sigma$  for every  $x \in \mathbb{R}$ ,
- (2)  $V[G] \models \text{"} M_n^\#(x) \text{ exists for every } x \in \mathbb{R}\text{"}$ , and

(3) *suppose*

(a)  $H \prec V_\eta$  is a countable substructure with  $a \in H$  where  $\eta$  is a large limit ordinal,

(b)  $\bar{H}$  is the transitive collapse of  $H$  with uncollapsing map  $\pi : \bar{H} \rightarrow H$ , and

(c)  $g$  is a  $\mathbb{P}^{\bar{H}}$ -generic filter over  $\bar{H}$  in  $V$ ,

then  $M_n^\#(x)$  exists in  $\bar{H}[g]$  for each  $x \in \mathbb{R}^{\bar{H}[g]}$  and is normally  $\omega_1$ -iterable via  $\Sigma$  in both  $\bar{H}[g]$  and  $V$ .

*Proof.* The proof works by induction on  $n$  as in lemma 3.4. We get uniqueness of  $M_n^\#(x)$  for  $x \in \mathbb{R}$  by lemma 2.38.

1. This works just as in the proof of lemma 3.4.

2. Let  $x \in \mathbb{R}^{V[G]}$  and let  $\tau$  be a nice  $\mathbb{P}$ -name with  $\tau^G = x$ . Since  $\mathbb{P} \subseteq \mathbb{R}$  is c.c.c.  $\tau$  can be coded by a real, so  $M_n^\#(\tau, a)$  exists. We can avoid working with  $M_n^\#(\mathbb{P}, \tau)$  since  $M_n^\#(\tau, a)$  has its own version of the forcing  $\mathbb{P}$  and this is absolute between  $M_n^\#(\tau, a)$  and  $V$ . We get

$$\forall y, y' \in \mathbb{P}(y \perp y' \Leftrightarrow \neg \exists z \in \mathbb{P}(z \leq y, y'))$$

in  $M_n^\#(\tau, a)$  by  $\Pi_3^1$  downwards absoluteness, where  $\mathbb{P}, \leq, \perp$  are given by their  $\Sigma_2^1(a)$  definition. Now for  $y \in \mathbb{R}$  the statement

” $y$  codes a countable subset of  $\mathbb{P}$ ”

is  $\Sigma_2^1(a)$ . Since

” $y$  codes a countable predense subset of  $\mathbb{P}$ ”

holds if and only if  $y$  codes a subset  $\{y_n : n < \omega\}$  of  $y$  and  $\forall z \in \mathbb{P} \exists n(y_n \not\leq z)$ , this is a combination of a  $\Sigma_2^1(a)$  and a  $\Pi_2^1(a)$  statement. So it is absolute between

$M_n^\#(\tau, a)$  and  $V$ . Hence

$$G' := G \cap M_n^\#(\tau, a)$$

is  $\mathbb{P}$ -generic over  $M_n^\#(\tau, a)$ . Moreover

$$x = \tau^{G'} \in M_n^\#(\tau, a)[G'].$$

Now  $M_n^\#(\tau, a)$  is normally  $\omega_1$ -iterable via  $\Sigma$  in  $V[G]$  by 1. Since  $\mathbb{P}$  is small compared to the critical points of the extenders on the  $M_n^\#(\tau, a)$  sequence,  $M_n^\#(\tau, a)[G']$  is  $\omega_1$ -iterable via  $\Sigma$  in  $V[G]$  as well. We can construct  $M_n^\#(x)$  in  $M_n^\#(\tau, a)[G']$  via an  $L[\vec{E}]$ -construction as in lemma 3.4.

3. Let  $x \in \mathbb{R}^{\bar{H}[g]}$ . Let  $\bar{\tau}$  be a nice  $\mathbb{P}^{\bar{H}[g]}$ -name with  $\bar{\tau}^g = x$  and  $\tau := \pi(\bar{\tau})$ . Then  $M_n^\#(\bar{\tau}, a)^{\bar{H}}$  and  $M_n^\#(\tau, a)$  exist and are normally  $\omega_1$ -iterable via  $\Sigma$  in  $\bar{H}$  and  $V$  respectively and

$$\pi(M_n^\#(\bar{\tau}, a)^{\bar{H}}) = M_n^\#(\tau, a).$$

So  $M_n^\#(\bar{\tau}, a)^{\bar{H}}$  is normally  $\omega_1$ -iterable via  $\Sigma$  in  $\bar{H}[g]$  by 1.

Let  $g' := g \cap M_n^\#(\bar{\tau}, a)^{\bar{H}}$ . Since the statement

” $y$  codes a countable predense subset of  $\mathbb{P}$ ”

is absolute between  $M_n^\#(\bar{\tau}, a)$  and  $V$ , we can conclude that  $g'$  is  $\bar{\mathbb{P}}$ -generic over  $M_n^\#(\bar{\tau}, a)^{\bar{H}}$ . Moreover  $x = \bar{\tau}^{g'} \in M_n^\#(\bar{\tau}, a)^{\bar{H}}[g']$ . Now  $M_n^\#(\bar{\tau}, a)^{\bar{H}}[g']$  is normally  $\omega_1$ -iterable via  $\Sigma$  in  $\bar{H}[g]$ , since  $\bar{\mathbb{P}}$  is small compared to the critical points of the extenders on the  $M_n^\#(\bar{\tau}, a)^{\bar{H}}$  sequence. Moreover  $M_n^\#(\bar{\tau}, a)^{\bar{H}}$  is normally  $\omega_1$ -iterable via  $\Sigma$  in  $V$  since

$$\pi \upharpoonright M_n^\#(\bar{\tau}, a)^{\bar{H}} : M_n^\#(\bar{\tau}, a)^{\bar{H}} \rightarrow M_n^\#(\tau, a)$$

is elementary. Hence  $M_n^\#(\bar{\tau}, a)^{\bar{H}}[g']$  is normally  $\omega_1$ -iterable via  $\Sigma$  in  $V$  as well. Finally we can construct the  $M_n^\#(x)$  of  $\bar{H}[g]$  and  $V$  in  $M_n^\#(\bar{\tau}, a)^{\bar{H}}[g']$  as in lemma 3.4.  $\square$

Note that the existence of  $M_n^\#(x)$  for every  $x \in \mathbb{R}$  is equivalent to  $\text{Det}(\mathbf{\Pi}_{n+1}^1)$  by theorem 2.30. The previous lemma can be applied to generalize

**Lemma 3.12.** (Woodin [45])  *$\text{Det}(\mathbf{\Pi}_n^1)$  implies  $\Sigma_{n+2}^1$  Cohen (random) absoluteness. In fact for odd  $n$  it is sufficient to assume*

- (1)  $\mathbf{\Pi}_n^1$  is scaled and
- (2) all  $\mathbf{\Delta}_{n+1}^1$  sets have the Baire property (are Lebesgue measurable).

*Proof.* See [45, lemma 2].  $\square$

**Lemma 3.13.** *Let  $\mathbb{P}$  be a  $\Sigma_2^1$  c.c.c. forcing and suppose  $M_n^\#(x)$  exists for every  $x \in \mathbb{R}$ . Then  $\Sigma_{n+3}^1$ -absoluteness holds for  $\mathbb{P}$ .*

*Proof.* We follow the proof of lemma 3.7. Let  $\mathbb{P}$  be a  $\Sigma_2^1(a)$  forcing. Suppose  $\exists x\varphi(x, y)$  holds in some  $\mathbb{P}$ -generic extension of  $V$ , where  $\varphi$  is a  $\mathbf{\Pi}_{n+2}^1$  formula and  $y \in \mathbb{R}$ . Let  $\tau$  be a nice name and  $p \in \mathbb{P}$  a condition with  $p \Vdash_{\mathbb{P}} \varphi(\tau, \check{y})$ . Let further  $\mathcal{M} := M_n^\#(a, y, \tau)$ .

Consider the tree  $T$  in  $V$  searching for  $g$  and  $x$  such that

- (1)  $g$  is  $\mathbb{P}^{\mathcal{M}}$ -generic over  $\mathcal{M}$  and
- (2)  $x$  is a real in  $\mathcal{M}[g]$  such that  $\mathcal{M}[g] \models \varphi(x, y)$  if  $n$  is even, and  $\Vdash_{\text{Col}(\omega, \delta)}^{\mathcal{M}[g]} \varphi(\check{x}, \check{y})$  if  $n$  is odd, where  $\delta$  is the least Woodin cardinal in  $\mathcal{M}$ .

Now let  $G$  be  $\mathbb{P} \upharpoonright p$ -generic over  $V$ . Then  $g := G \cap \mathcal{M}$  is  $\mathbb{P}^{\mathcal{M}}$ -generic over  $\mathcal{M}$  since  $\mathbb{P}$  is  $\Sigma_2^1(a)$ . Since  $\mathcal{M}[g]$  is normally  $\omega_1$ -iterable in  $V[G]$  by lemma 3.11,  $T$  has a branch in  $V[G]$ . Then  $T$  has a branch in  $V$  and hence  $V \models \exists x\varphi(x, y)$ .  $\square$

Note that one cannot prove this from  $n$  Woodin cardinals:

**Lemma 3.14.**  $\Sigma_{n+3}^1$  Cohen absoluteness does not hold in  $M_n^\#$  for even  $n < \omega$ .

*Proof.* The set of reals  $\mathbb{R} \cap M_n^\#$  is  $\Sigma_{n+1}^1$  for even  $n$  by [40, theorem 3.4]. So the statement that there is a Cohen real over  $M_n^\#$  is  $\Sigma_{n+3}^1$ . Since  $M_n^\#$  and the Cohen generic extension  $M_n^\#[g]$  are  $\Sigma_{n+2}^1$ -correct in  $V$  by lemma 2.45, this statement holds true in  $M_n^\#[g]$  but not in  $M_n^\#$ .  $\square$

We get analogues of lemma 3.8 and theorem 3.9:

**Lemma 3.15.** Let  $E$  be a thin  $\Pi_{n+3}^1$  equivalence relation. Suppose  $\mathbb{P}$  is a  $\Sigma_2^1$  c.c.c. forcing and  $M_n^\#(x)$  exists for every  $x \in \mathbb{R}$ . Let  $\tau$  be a  $\mathbb{P}$ -name for a real. Then the set

$$D := \{p \in \mathbb{P} : (p, p) \Vdash_{\mathbb{P} \times \mathbb{P}} \tau E \tau'\}$$

is dense.

**Theorem 3.16.** Let  $\mathbb{P}$  be a  $\Sigma_2^1$  c.c.c. forcing and suppose  $M_n^\#(x)$  exists for every  $x \in \mathbb{R}$ . Then  $\mathbb{P}$  does not add equivalence classes to thin provably  $\Delta_{n+3}^1$  equivalence relations.

3.2.2. *Prewellorders and generic absoluteness.* Let  $N_{\mathbb{P}}$  denote the set of nice names  $\tau = \{(p, \check{n}) : p \in A_n\}$  for reals, where each  $A_n$  is an antichain in a forcing  $\mathbb{P}$ . In case  $\mathbb{P} \subseteq \mathbb{R}$  has the c.c.c. every nice name can be coded by a real.

**Lemma 3.17.** (Bagaria, Bosch [1]) Suppose  $\mathbb{P}$  is a c.c.c.  $\Sigma_n^1(x)$  forcing and  $\varphi$  is a  $\Sigma_k^1(\Pi_k^1)$  formula where  $n \geq 1$  and  $k \geq 2$ . Then

$$R := \{(p, \tau) : \tau \in N_{\mathbb{P}} \wedge p \Vdash_{\mathbb{P}} \varphi(\tau)\}$$

is a  $\Sigma_{n+k-1}^1(x)$  ( $\Pi_{n+k-1}^1(x)$ ) set.

*Proof.* The set  $N_{\mathbb{P}}$  of nice  $\mathbb{P}$ -names for reals is a  $\Pi_n^1$  subset of  $\mathbb{R}$  by [1, fact 2.6]. We sketch the proof of the lemma from [1] for  $\Sigma_k^1$  formulas. For  $k = 2$  and a  $\Pi_1^1$

formula  $\psi$  one can express

$$p \Vdash_{\mathbb{P}} \psi(\sigma, \tau)$$

by the  $\Delta_2^1$  statement that for every (for some) countable transitive model  $M$  containing  $p, \sigma, \tau$  of a fixed finite fragment of ZFC such that the inclusion  $\mathbb{P}^M \rightarrow \mathbb{P}$  is a complete embedding, we have that  $p \Vdash_{\mathbb{P}^M}^M \psi(\sigma, \tau)$ . Now by the forcing theorem

$$p \Vdash_{\mathbb{P}} \exists y \psi(\sigma, y) \Leftrightarrow \exists \tau \in N_{\mathbb{P}} (p \Vdash_{\mathbb{P}} \psi(\sigma, \tau)),$$

so  $R$  is  $\Sigma_{n+1}^1(x)$ . The rest is a straightforward induction on  $k$ . The proof for  $\Pi_k^1$  formulas is analogous.  $\square$

We use the previous lemma to show

**Lemma 3.18.**  $\Sigma_{n+1}^1$  Cohen absoluteness implies that Cohen forcing does not add any equivalence classes to  $< \omega - \mathbf{\Pi}_n^1$  prewellorders.

*Proof.* Suppose  $\leq$  is a  $< \omega - \mathbf{\Pi}_n^1$  prewellorder. Let  $\mathbb{P}$  denote Cohen forcing and suppose  $G$  is  $\mathbb{P}$ -generic over  $V$ . We denote the relation given by the same definition in  $V[G]$  by  $\leq$  as well. This is a prewellorder in  $V[G]$  since the statement that  $\leq$  is a prewellorder is  $\Pi_{n+1}^1$ . Moreover

$$\mathbb{1} \Vdash_{\mathbb{P}} \text{'' } \leq \text{ is a prewellorder''}$$

is  $\Pi_{n+1}^1$  by lemma 3.17. Hence this holds in  $V[G]$  by  $\Sigma_{n+1}^1$ -absoluteness, so we get

$$\mathbb{1} \Vdash_{\mathbb{P} * \mathbb{P}} \text{'' } \leq \text{ is a prewellorder''}.$$

Since the two-step iteration  $\mathbb{P} * \mathbb{P}$  of Cohen forcing is equal to the product, the same is forced by  $\mathbb{P} \times \mathbb{P}$ .

Suppose  $p \in \mathbb{P}$  and  $\tau \in N_{\mathbb{P}}$  such that  $p \Vdash_{\mathbb{P}} \tau \not\leq \check{x} \vee \check{x} \not\leq \tau$  for all  $x \in \mathbb{R}$ .

**Case 1.** *There is a real  $x \in \mathbb{R}$  and a condition  $q \leq p$  with  $q \Vdash_{\mathbb{P}} \tau \leq \check{x}$ . In this case choose  $x \in \mathbb{R}$  which is  $\leq$ -minimal with this property. Then*

$$\forall y < x (p \Vdash_{\mathbb{P}} \tau \not\leq \check{y}),$$

*which is a  $\Pi_{n+1}^1$  statement by lemma 3.17, since the map  $y \mapsto \check{y}$  is Borel.*

*Let  $G$  and  $H$  be mutually  $\mathbb{P} \upharpoonright p$ -generic filters over  $V$ . Then  $\forall y < x (p \Vdash_{\mathbb{P}} \tau \not\leq \check{y})$  holds in  $V[G]$  by  $\Sigma_{n+1}^1$ -absoluteness. In particular we have*

$$p \Vdash_{\mathbb{P}}^{V[G]} \tau \not\leq \tau^G.$$

*So  $V[G \times H] \models \tau^H \not\leq \tau^G$  and by the same argument  $V[G \times H] \models \tau^G \not\leq \tau^H$ . This is contradictory, since  $\leq$  is linear in  $V[G \times H]$ .*

**Case 2.** *There is a condition  $q \leq p$  such that for every  $x \in \mathbb{R}$  we have  $q \Vdash_{\mathbb{P}} \check{x} < \tau$ . Then*

$$\forall x \in \mathbb{R} (q \Vdash_{\mathbb{P}} \check{x} < \tau).$$

*Let  $G$  and  $H$  be mutually  $\mathbb{P} \upharpoonright q$ -generic over  $V$ . Again we get  $\forall x \in \mathbb{R} (q \Vdash_{\mathbb{P}} \check{x} < \tau)$  in  $V[G]$ . In particular*

$$q \Vdash_{\mathbb{P}}^{V[G]} \tau^G < \tau$$

*so that  $V[G \times H] \models \tau^G < \tau^H$ . By the same argument  $V[G \times H] \models \tau^H < \tau^G$ , which is impossible.*

□

We conclude that

**Corollary 3.19.** *Cohen forcing does not add equivalence classes to  $< \omega - \Pi_n^1$  prewellorders if and only if  $\Sigma_{n+1}^1$  Cohen absoluteness holds, for  $n \geq 1$ .*

*Proof.* One direction is the previous lemma. For the other direction suppose we have proved  $\Sigma_k^1$  Cohen absoluteness for some  $k \leq n$ . Let  $G$  be Cohen generic over  $V$ . Suppose  $V[G] \models \exists x \varphi(x, \vec{a})$  where  $\varphi \in \Pi_k^1$  and  $\vec{a} \in \mathbb{R}^{<\omega}$ .

We define a prewellorder  $\leq$  by letting  $x \leq y$  if and only if  $\varphi(x, \vec{a}) \vee \neg\varphi(y, \vec{a})$ . Then one of the equivalence classes of the prewellorder is

$$\{x \in \mathbb{R} : \varphi(x, \vec{a})\}.$$

Since Cohen forcing does not add any equivalence classes, there is a real  $x \in \mathbb{R} \cap V$  with  $V \models \varphi(x, \vec{a})$ .  $\square$

#### 4. THE NUMBER OF EQUIVALENCE CLASSES

We give a proof of the Harrington-Shelah theorem [9] for counting the number of equivalence classes of thin  $\text{co-}\kappa$ -Suslin equivalence relations in the first section. This is applied to compute the number of equivalence classes of thin  $\Pi_n^1$  and  $\Sigma_{2n+1}^1$  equivalence relations, assuming the existence of  $\Pi_{2k+1}^1$ -scales and the Baire property of projective sets. It is finally shown that thin  $\Sigma_{2n}^1$  equivalence relations are  $\Pi_{2n}^1$  in any real coding  $M_{2n-1}^\#$  for  $n \geq 1$ . The base theory is  $\text{ZF} + \text{DC}$ .

**4.1. Co- $\kappa$ -Suslin equivalence relations.** We present a proof of the Harrington-Shelah theorem. Most of this section is due to Harrington and Shelah.

4.1.1. *A few lemmas.*

**Definition 4.1.** *Suppose  $\kappa \in \text{Ord}$ . A set  $A \subseteq \mathbb{R}^n$  is called  $\kappa$ -Suslin if  $A = p[T]$  for some tree  $T$  on  $\omega^n \times \kappa$ . It is  $\text{co-}\kappa$ -Suslin if  $\mathbb{R}^n - A$  is  $\kappa$ -Suslin.*

Note that if  $\text{AD}$  holds in  $L(\mathbb{R})$ , then the sets of reals which are  $\kappa$ -Suslin in  $L(\mathbb{R})$  for some ordinal  $\kappa$  are exactly the  $(\Sigma_1^2)^{L(\mathbb{R})}$  sets of reals, as shown by Martin and Steel [26].

In this section,  $E$  denotes an equivalence relation which is co- $\kappa$ -Suslin via  $T$ . Note that  $\mathbb{R}^2 - p[T]$  is not necessarily an equivalence relation in generic extensions. For example, if  $\mathbb{R}$  is wellorderable and

$$T := \{(s, (x, \dots, x)) \in (\omega \times \mathbb{R})^{<\omega} : s \subseteq x \wedge x \in \mathbb{R}^2 - E\},$$

then  $p[T] = \mathbb{R}^2 - E$  is the same set in every generic extension, so whenever  $\mathbb{P}$  adds reals  $\mathbb{R}^2 - p[T]$  is not an equivalence relation in  $V^{\mathbb{P}}$ .

For the application to projective equivalence relations we note the following consequence of the second periodicity theorem [17, see 30.12]:

**Lemma 4.2.** *Assume  $\text{ZF} + \text{DC}$  and  $\text{Det}(\Delta_{2n}^1)$ . Then every  $\Pi_{2n+2}^1$  set is co- $\delta_{2n+1}^1$ -Suslin via the tree from a  $\Sigma_{2n+2}^1$ -scale.*

The existence of a perfect set of pairwise inequivalent reals for  $E = \mathbb{R}^2 - p[T]$  is not absolute between  $V$  and generic extensions. We work with a stronger and absolute version, which is called strongly thick in [9].

**Definition 4.3.** *Suppose  $T$  is a tree and  $E = \mathbb{R}^2 - p[T]$  is an equivalence relation and  $S \subseteq T$ . We say  $S$  witnesses that  $E$  is not thin if there is a perfect set  $P \subseteq \mathbb{R}$  such that  $(x, y) \in p[S]$  for all  $x, y \in P$  with  $x \neq y$ .*

For a tree  $T$  and a node  $r \in T$  one says that  $r$  splits in  $T$  if there are  $s, t \in T$  with  $r \subseteq s, t$  and  $s \perp t$ .

**Lemma 4.4.** *The existence of a countable set  $S \subseteq T$  which witnesses that  $E = \mathbb{R}^2 - p[T]$  is not thin is absolute between transitive models of  $\text{ZF}$ .*

*Proof.* We define a partial order  $(X, <)$  such that a countable set  $S$  with this property exists if and only if  $(X, <)$  is ill-founded. Let  $X$  be the set of all triples  $(r, s, F)$  such that

- (1)  $r$  is a finite tree on  $\omega$ ,
- (2)  $s \subseteq T$  is finite,
- (3)  $F$  is a finite set of finite functions,
- (4) for any two  $u, v \in r$  with  $u \neq v$  there is a function  $f \in F$  with  $(u, v, f) \in s$ ,

and let  $(r, s, F) < (p, q, G)$  if

- (1)  $p \subseteq r$ ,
- (2)  $q \subseteq s$ ,
- (3) every node in  $p$  splits in  $r$ , and
- (4) for any  $u, v \in p$  with  $u \neq v$  and any  $g \in G$  with  $(w, x, g) \in q$  there are  $y, z \in r$  and a function  $f \in F$  with  $u \subseteq y, v \subseteq z, g \subseteq f$ , and  $(y, z, f) \in s$ .

If a countable set  $S \subseteq T$  and a perfect set  $P \subseteq \mathbb{R}$  witness that  $E$  is not thin, one can define  $(r_n, s_n, F_n) \in X$  such that  $(r_n, s_n, F_n) < (r_k, s_k, F_k)$  for all  $k < n < \omega$  and for any two distinct  $u, v \in r_n$  of the same length there are reals  $x, y \in P$  and a function  $f : \omega \rightarrow Ord$  with

- (1)  $u \subseteq x$ ,
- (2)  $v \subseteq y$ ,
- (3)  $(u, v, f \upharpoonright lh(u)) \in s_n$ , and
- (4)  $(x, y, f) \in [S]$ .

If on the other hand  $(X, <)$  is illfounded, let  $((r_n, s_n, F_n) : n < \omega)$  be a strictly decreasing sequence in  $(X, <)$ . We can set  $S := \{f \upharpoonright k : \exists n < \omega (f \in F_n \wedge k < \omega)\}$  and  $P := [U]$  where  $U := \bigcup_{n \in \omega} r_n$ .

This works without choice since  $X$  can be wellordered. □

We will work with the infinitary logic  $\mathcal{L}_{\infty, \omega}$  over a language  $\mathcal{L}$  [3, chapter III, definition 1.5].  $\mathcal{L}_{\infty, \omega}$ -formulas are distinguished from finitary formulas by the fact that disjunctions and conjunctions of arbitrary ordinal length are possible. Let  $\mathcal{L}$  be a language which contains at least  $\in$  and the following constants:  $c, d$

for reals,  $\dot{f}$  for a function  $f : \omega \rightarrow \kappa$ , and  $\dot{s}$  for each  $s \in tc(\{T\})$ . Let  $N$  be the set of atomic formulas  $\dot{n} \in c$  with  $n \in \omega$ . We build  $\mathcal{L}_{\infty,0,N}$  by starting with  $N$  and closing under negations and wellordered infinitary disjunctions and conjunctions. We will also write  $\varphi_d$  for the formula obtained from a formula  $\varphi \in \mathcal{L}_{\infty,0,N}$  by replacing  $c$  with  $d$ . Note that instead of  $\mathcal{L}_{\infty,0,N}$  one can equivalently work with the infinitary logic  $\mathcal{L}_{\infty,0}$  built over a language with a set of propositional formulas  $\{p_n : n < \omega\}$ , as is done in [9] and [13].

Whether a statement  $\varphi \in \mathcal{L}_{\infty,0,N}$  is true depends only on the truth value of each individual atomic statement  $\dot{n} \in c$ .

**Definition 4.5.** *Suppose  $\varphi \in \mathcal{L}_{\infty,0,N}$  and  $x \in \mathbb{R}$ . Define the truth value of  $\dot{n} \in c$  as true if and only if  $n \in x$ . This induces a truth value for  $\varphi$  by induction on the formula complexity. If this value is true we say that  $\varphi(x)$  holds and  $x$  is a model of  $\varphi$ .*

We refer to the infinitary proof calculus from [3] which has the rule

$$\forall \alpha < \beta \vdash \varphi_\alpha \quad \Rightarrow \quad \vdash \bigwedge_{\alpha < \beta} \varphi_\alpha$$

in addition to the rules of first-order logic [3, chapter III, definition 5.1]. In the following let  $\chi$  be the  $\mathcal{L}_{\infty,\omega}$ -formula

$$c \subseteq \omega \wedge d \subseteq \omega \wedge \bigwedge_{t \in tc(\{T\})} (\forall x \in t \bigvee_{s \in t} x = \dot{s}) \wedge (\bigwedge_{s \in t} \dot{s} \in t).$$

Then  $c, d$  are interpreted as reals and  $\dot{s}$  takes the value  $s$  for each  $s \in tc(\{T\})$  in any transitive model of  $\chi$ . Note that for any admissible set  $\mathbb{A}$  with  $T \in \mathbb{A}$  we have  $\chi \in \mathbb{A}$  by  $\Delta_0$ -replacement, since we can assume that  $s$  and  $\dot{s}$  are  $\Delta_0$ -definable from each other for each  $s \in tc(\{T\})$ . We write  $\vdash_\chi$  for the provability relation when  $\chi$  is used as an axiom.

A theory in  $\mathcal{L}_{\infty,\omega}$  is consistent if it is not contradictory in terms of infinitary proofs. Note that for hereditarily countable theories, this definition coincides with several other definitions of consistency:

**Lemma 4.6.** *The following are equivalent for any hereditarily countable theory  $\Sigma \subseteq \mathcal{L}_{\infty,0,N}$ :*

- (1)  $\Sigma$  is consistent
- (2)  $\Sigma$  is consistent in any admissible set  $\mathbb{A}$  with  $\Sigma \in \mathbb{A}$
- (3)  $\Sigma$  has a model
- (4) there is a model of  $\Sigma$  in some generic extension
- (5) player 1 wins the closed game  $G_\Sigma$  from [13, section 2.2].

*Proof.* We sketch the relevant part that  $\Sigma$  has a model if it is consistent. Suppose  $\Sigma$  is consistent. We can assume that negations occur only at the atomic level in formulas in  $\Sigma$ , since every  $\mathcal{L}_{\infty,0,N}$ -formula is equivalent to a formula of this form. Suppose two players play a game  $G_\Sigma$  with the rules:

- (1) if player 1 plays  $\bigvee_{\alpha < \beta} \chi_\alpha$ , then player 2 has to play  $\chi_\alpha$  for some  $\alpha < \beta$ ,  
and
- (2) both players can only play formulas which are consistent with  $\Sigma$  and the previously played moves,

where player 2 wins if the game does not stop after finitely many moves. Then player 2 has a winning strategy in  $G_\Sigma$ , since  $\Sigma$  is consistent. By letting player 1 play each disjunction in  $tc(\Sigma)$  consistent with  $\Sigma$  and the previous moves at some point in the game, the play determines a real  $x$  so that  $n \in x$  if and only if the formula  $\dot{n} \in c$  was played during this run of the game. One shows by induction on the formula complexity that  $x$  models  $\Sigma$ .

The equivalence of 1 and 2 follows from the Barwise completeness theorem [3, part III, theorem 5.5].  $\square$

**Lemma 4.7.** *A Cohen real adds a perfect set of mutually generic Cohen reals*

*Proof.* Let  $\mathbb{Q}$  be the forcing which consists of finite trees on  $\omega$ , where  $s \leq t$  if and only if  $t \subseteq s$  and every node of  $t$  splits in  $s$ . Then  $\mathbb{Q}$  is equivalent to Cohen forcing  $\mathbb{P}$  since it is countable and has no atoms. Also any two branches of the tree added by  $\mathbb{Q}$  are mutually  $\mathbb{P}$ -generic, since for every dense open set  $D \subseteq \mathbb{P} \times \mathbb{P}$  the set

$$D' := \{t \in \mathbb{P} : \forall r, s \in t (r \neq s \Rightarrow \exists r', s' \in t (r \subseteq r' \wedge s \subseteq s' \wedge (r', s') \in D))\}$$

is dense in  $\mathbb{Q}$ . □

4.1.2. *The theorem of Harrington and Shelah.* In this section we give a proof of

**Theorem 4.8.** *(Harrington-Shelah [9]) Assume ZF. Suppose  $\kappa$  is an infinite cardinal and  $T$  is a tree on  $\omega \times \omega \times \kappa$ . Let  $\mathbb{A}$  be an admissible set with  $T \in \mathbb{A}$ . Suppose  $E = \mathbb{R}^2 - p[T]$  is a thin equivalence relation such that*

- (1)  $\Vdash_{Cohen}^{L[T]}$  " $\mathbb{R}^2 - p[T]$  is transitive" or
- (2) *there is a Cohen real over  $L[T]$  in  $V$ .*

*Then for every  $x \in \mathbb{R}$  there is a formula  $\varphi \in \mathcal{L}_{\infty,0,N} \cap \mathbb{A}$  with*

- (1)  $\varphi(x)$  and
- (2)  $\vdash_x^{\mathbb{A}} (\varphi \wedge \varphi_d) \rightarrow (c, d) \notin p[\dot{T}]$ .

*Proof.* If there is a Cohen real  $x$  over  $L[T]$  in  $V$ , then  $\mathbb{R}^2 - p[T]$  is an equivalence relation in  $L[T, x]$  by absoluteness of  $p[T]$ , since  $E = \mathbb{R}^2 - p[T]$  is an equivalence relation in  $V$ . So we assume the first condition holds. Note that this condition is also true if  $\mathbb{R}^2 - p[T]$  is transitive in a Cohen generic extension of  $V$ .

We work in  $L[T]$ . If the theory

$$\Sigma := \{\neg\varphi : \varphi \in \mathcal{L}_{\infty,0,N} \cap \mathbb{A} : \vdash_x (\varphi \wedge \varphi_d) \rightarrow (c, d) \notin p[\dot{T}]\}$$

is inconsistent, then no real can satisfy every statement in  $\Sigma$ . Here  $\vdash_\chi$  can be equivalently replaced by  $\vdash_\chi^{\mathbb{A}}$  by the Barwise completeness theorem [3, chapter III, theorem 5.5]. Then for every  $x \in \mathbb{R}$  there is a formula  $\varphi$  satisfying the conditions and we are done.

Assume  $\Sigma$  is consistent and let  $A \prec \mathbb{A}$  be a countable substructure with  $\Sigma, T \in A$  and  $\bar{A}$  its transitive collapse with uncollapsing map  $\pi : \bar{A} \rightarrow A$ ,  $\pi(\bar{T}) = T$ ,  $\pi(\bar{\chi}) = \chi$ ,  $\pi(\bar{\kappa}) = \kappa$ , and  $\pi(p) = \Sigma$ . Further suppose  $\dot{T}, \dot{n} \in A$ ,  $\pi(\dot{T}) = T$ ,  $\pi(\dot{f}) = f$ , and  $\pi(\dot{n}) = n$  for all  $n \in \omega$ . Also assume that  $s$  and  $\dot{s}$  are  $\Delta_0$ -definable from each other for each  $s \in tc(\{T\})$  to ensure that  $\dot{T}$  is interpreted as  $\bar{T}$  in every model of  $\bar{\chi}$ . We will refer to provability and consistency as provability from  $\bar{\chi}$  and consistency with  $\bar{\chi}$  and denote  $\vdash_{\bar{\chi}}$  simply by  $\vdash$ .

**Claim 4.9.** *Suppose  $p \subseteq q \subseteq \mathcal{L}_{\infty,0,N} \cap \bar{A}$ ,  $q$  is consistent,  $q$  is  $\Sigma_1$  over  $\bar{A}$ . Then  $q \cup q_d \cup \{(c, d) \in p[\dot{T}]\}$  is consistent.*

*Proof.* Assume the theory is inconsistent, so it does not have a model. Now Barwise compactness [3, chapter III, theorem 5.6] implies that there is a theory  $s \in \bar{A}$  with  $\vdash q \rightarrow s$  such that  $s \cup s_d \cup \{(c, d) \in p[\dot{T}]\}$  does not have a model. Hence this theory is inconsistent by lemma 4.6. We can replace  $s$  by a the conjunction  $\varphi \in \mathcal{L}_{\infty,0,N} \cap \bar{A}$  of all formulas in  $s$  and get  $\vdash (\varphi \wedge \varphi_d) \rightarrow (c, d) \notin p[\dot{T}]$ . Then  $\neg\varphi \in p \subseteq q$  by definition of  $\Sigma$ . But this cannot happen, since  $\vdash q \rightarrow \varphi$  and  $q$  is consistent.  $\square$

Using the claim, one would like to build a perfect tree of consistent theories starting with  $p$  such that every branch defines a complete theory. Then every branch defines a unique real. At the same time one would have to ensure that  $(x, y) \in p[T]$  for reals  $x, y$  from distinct branches and this would contradict that  $E$  is thin.

However, this cannot work directly, since the assumption on Cohen forcing is necessary by remark 4.14. Instead one can construct a countable set  $S \subseteq T$  and a perfect set in a Cohen generic extension witnessing that  $\mathbb{R}^2 - p[T]$  is not thin and then apply lemma 4.4.

**Claim 4.10.** *Every consistent  $\mathcal{L}_{\infty,0,N}$ -theory  $q$  with  $p \subseteq q$  which is  $\Sigma_1$  over  $\bar{A}$  is incomplete.*

*Proof.* Suppose  $q$  is complete and consistent. Then the theory  $q \cup q_d \cup \{(c, d) \in p[\dot{T}]\}$  is consistent by claim 4.6. So

$$q \cup q_d \cup \{(c, d, \dot{f}) \in [\dot{T}]\}$$

is consistent as well. This theory has a model  $(x, y, f) \in L[T]$  since one can also apply lemma 4.6 if  $N$  is replaced by the set  $N'$  which additionally contains each formula  $\dot{f}(\dot{n}) = \dot{\alpha}$  for  $\alpha < \bar{\kappa}$  and  $n < \omega$ . Then  $x = y$  since  $x$  and  $y$  are both models of  $q$  and  $q$  is complete. Hence  $(x, x, f) \in [\bar{T}]$ . Note that  $p[\bar{T}] \subseteq p[T]$  since  $\pi''\bar{T}$  is obtained from  $T$  by omitting all ordinals not in  $A$ . But this would imply  $(x, x) \in p[T]$  and hence  $(x, x) \notin E$ .  $\square$

Let  $(\psi_n : n < \omega)$  enumerate the formulas in  $\mathcal{L}_{\infty,0,N} \cap \bar{A}$ . We can assume that negations only occur on the atomic level in all formulas.

We build a tree  $\mathbb{C}$  in  $L[T]$  whose nodes are consistent  $\mathcal{L}_{\infty,0,N}$ -theories  $q \supseteq p$  which are  $\Sigma_1$  over  $\bar{A}$ , ordered by inclusion. The root of the tree is  $p$ . We can construct the tree level by level and ensure that  $\mathbb{C}$  is isomorphic to  $2^{<\omega}$  and

- (1)  $\psi_n \in q$  or  $\neg\psi_n \notin q$  for all  $q$  on level  $n$  and
- (2) if  $\psi_n \equiv \bigvee_{\alpha < \beta} \chi_\alpha$ ,  $\psi_n \in q$ , and  $q$  is on level  $\langle n, k \rangle$  for some  $k < \omega$ , then for every  $r \supseteq q$  on level  $\langle n, k \rangle + 1$  there is some  $\alpha < \beta$  with  $\chi_\alpha \in r$ .

Then every branch in  $\mathbb{C}$  defines a consistent theory and a real which is a model of the theory. Note that there are no end nodes in  $\mathbb{C}$  by claim 4.10. We will force

with  $(\mathbb{C}, \leq)$ , where  $\leq$  denotes reverse inclusion. If  $G$  is  $\mathbb{C}$ -generic, then  $\bigcup G$  is a complete theory and defines a real  $x$ . The generic filter can also be recovered from the real  $x$  as

$$G = \{q \in \mathbb{C} : \forall \varphi \in q \varphi(x)\}.$$

Let  $N'$  be the set of atomic statements about  $c$ ,  $d$ , and  $f$ . Let  $\mathbb{P}$  be a tree of consistent  $\mathcal{L}_{\infty,0,N'}$ -theories containing  $p \cup p_d \cup \{c, d, f\} \in [\dot{T}]$  which are  $\Sigma_1$  over  $\bar{A}$ . We build  $\mathbb{P}$  in the same way as  $\mathbb{C}$  such that additionally the value of  $(c|n, d|n, f|n)$  is decided on the  $n^{\text{th}}$  level and  $\mathbb{P}$  is isomorphic to  $2^{<\omega}$ . Then every branch in  $\mathbb{P}$  defines a consistent theory containing  $p \cup p_d \cup \{(c, d, f) \in [\dot{T}]\}$  and a triple  $(x, y, f)$  which is a model of this theory. We will force with  $(\mathbb{P}, \leq)$  where  $\leq$  denotes reverse inclusion.

**Claim 4.11.** *Suppose  $G$  is  $\mathbb{P}$ -generic and  $(x, y, f)$  is a model of the corresponding theory. Then both  $x$  and  $y$  are  $\mathbb{C}$ -generic.*

*Proof.* Suppose  $D \subseteq \mathbb{C}$  is open dense. It suffices to find  $r \in D$  such that  $x$  models  $r$ . For  $q \in \mathbb{P}$  let

$$q(c) := \{\varphi \in \mathcal{L}_{\infty,0,N} \cap \bar{A} : q \vdash \varphi\}$$

be the set of statements about  $c$  which are provable from  $q$ .

We claim that

$$D' := \{q \in \mathbb{P} : q(c) \in D\}$$

is dense in  $\mathbb{P}$ . So suppose  $q \in \mathbb{P}$ . There is a condition  $r \in D$  with  $q(c) \subseteq r$  since  $D$  is dense. If  $q' := q \cup r$  was inconsistent, then by Barwise compactness and lemma 4.6 there would be a set  $s \subseteq r$  with  $s \in \bar{A}$  such that  $q \vdash \neg \bigwedge s$ . Hence  $\neg \bigwedge s \in q(c)$ . But  $\bigwedge s$  is consistent with  $q(c)$  since  $s \subseteq r$ . So  $q'$  is consistent. Since  $r \subseteq q'(c)$  we have  $q'(c) \in D$  and hence  $q' \in D'$ .

Choose a condition  $q \in G \cap D'$  and let  $r := q(c) \in D$ . Then  $x$  models  $r$ , since  $q \in G$  and  $(x, y, f)$  models  $\bigcup G$ .  $\square$

**Claim 4.12.** *If  $(x, y)$  is  $\mathbb{C} \times \mathbb{C}$ -generic over  $L[T]$ , then  $(x, y) \in p[T]$ .*

*Proof.* Suppose there are conditions  $q, q' \in \mathbb{C}$  with  $(q, q') \Vdash_{\mathbb{C} \times \mathbb{C}} (\dot{x}_0, \dot{x}_1) \notin p[\check{T}]$ , where  $\dot{x}_0$  and  $\dot{x}_1$  are names for the left and right generic reals. Then

$$s := q \cup q_d \cup \{(c, d) \in p[\check{T}]\}$$

is consistent by claim 4.9. Let  $s' \in \mathbb{P}$  be a condition with  $s' \supseteq s$ . Suppose  $G$  is Cohen-generic over  $L[T]$ . There is a  $\mathbb{C} \restriction q' \times \mathbb{P} \restriction s'$ -generic filter over  $L[T]$  in  $L[T, G]$  by lemma 4.7. Let  $x$  be the  $\mathbb{C}$ -generic real and  $y$  and  $z$  the reals from the  $\mathbb{P}$ -generic filter as in claim 4.11. Then both  $y$  and  $z$  are  $\mathbb{C} \restriction q$ -generic over  $L[T]$ . Since  $p[T]$  is absolute, we have

$$L[T, G] \models (x, z) \notin p[T]$$

and

$$L[T, G] \models (y, z) \notin p[T]$$

since this is forced by  $(q, q')$  and

$$L[T, G] \models (x, y) \in p[\bar{T}].$$

But this cannot happen, since  $p[\bar{T}] \subseteq p[T]$  and  $\mathbb{R}^2 - p[T]$  is transitive in any Cohen generic extension of  $L[T]$ .  $\square$

Hence Cohen forcing adds a perfect set of pairwise inequivalent reals by lemma 4.7 and claim 4.12.

Let  $\tau$  be a  $\mathbb{C}$ -name for a sequence of ordinals with  $\Vdash_{\mathbb{C} \times \mathbb{C}} (\dot{x}_0, \dot{x}_1, \tau) \in [\check{T}]$  by the forcing theorem, where  $\dot{x}_0$  and  $\dot{x}_1$  are names for the left and right generic reals. Since  $\mathbb{C}$  is proper, there is in fact a countable set  $S \subseteq T$  such that  $\Vdash_{\mathbb{C} \times \mathbb{C}} (\dot{x}_0, \dot{x}_1, \tau) \in [\check{S}]$ . But then there would also be a countable subset of  $T$  witnessing

that  $\mathbb{R}^2 - p[T]$  is not thin in  $L[T]$  by the absoluteness proved in lemma 4.4. This would imply that  $E$  is not thin in  $V$ .  $\square$

It is not clear whether the previous proof can be generalized to Sacks forcing, or other forcings whose conditions are trees on  $\omega$ , instead of Cohen forcing.

Note that if  $\varphi$  satisfies the conditions in the previous theorem, then the set  $\{y \in \mathbb{R} : \varphi(y)\}$  is contained in the equivalence class of  $x$ . The aim of the theorem was to show

**Corollary 4.13.** *(Harrington-Shelah [9]) Assume ZF. Suppose  $T$  is a tree on  $\omega \times \omega \times \kappa$  and  $E = \mathbb{R}^2 - p[T]$  is a thin equivalence relation such that*

- (1)  $\Vdash_{Cohen}^{L[T]}$  " $\mathbb{R}^2 - p[T]$  is transitive" or
- (2) *there is a Cohen real over  $L[T]$  in  $V$ .*

*Then the equivalence classes of  $E$  can be wellordered with order type  $\leq \kappa$ .*

*Proof.* Let  $\mathbb{A}$  be the least admissible set with  $T \in \mathbb{A}$ . Then  $\mathbb{A} = L_\alpha[T]$  for some ordinal  $\alpha$ . Then  $\overline{\mathbb{A}} = \kappa$  since  $\mathbb{A} = L_\alpha[T] = h^{\mathbb{A}}(\kappa \cup \{T\})$  by minimality of  $\mathbb{A}$ . Let  $\Phi$  be the set of  $\mathcal{L}_{\infty,0,N}$ -formulas  $\varphi \in \mathbb{A}$  satisfying the conditions in theorem 4.8 and let  $(\varphi_\beta : \beta < \gamma)$  enumerate  $\Phi$  for some  $\gamma \leq \kappa$ . Then every equivalence class is the union of sets of the form  $\{x \in \mathbb{R} : \varphi_\beta(x)\}$  with  $\beta < \gamma$ . Hence

$$f([x]_E) := \min\{\beta < \gamma : \exists y \in [x]_E \varphi_\beta(xy)\}$$

is a rank function for the equivalence classes.  $\square$

**Remark 4.14.** *(Shelah [38]) The assumption that  $\mathbb{R}^2 - p[T]$  is transitive in a Cohen generic extension cannot be eliminated from the previous corollary.*

*Proof.* Shelah [38] defines a finite support iteration of c.c.c. forcings of length  $\omega_1$  assuming  $2^{\aleph_0} = \aleph_2$ , so that in the generic extension there is a thin co- $\aleph_1$ -Suslin equivalence relation with  $2^{\aleph_0}$  equivalence classes.  $\square$

Note that if  $\kappa$  is a successor cardinal and  $T$  a tree on  $\omega \times \omega \times \kappa$ , then  $L[T]$  is not a counterexample to the conclusion of corollary 4.13, since  $L[T]$  has at most  $\kappa$  many reals by a standard argument.

The Harrington-Shelah theorem has an interesting consequence together with a result of Neeman and Zapletal:

**Theorem 4.15.** *(Neeman and Zapletal [33]) Suppose  $\kappa$  is an infinite cardinal and there are a class inner model  $M$  and a countable ordinal  $\lambda$  with*

- (1)  $M = L(V_\lambda^M)$ ,
- (2)  $M \models$  " $\lambda$  is the supremum of  $\omega$  Woodin cardinals", and
- (3)  $M$  is uniquely  $\kappa^+ + 1$ -iterable.

*If  $\mathbb{P}$  is a proper forcing of size  $\leq \kappa$  and  $G$  is a  $\mathbb{P}$ -generic filter over  $V$ , then there is an elementary embedding  $j : L(\mathbb{R})^V \rightarrow L(\mathbb{R})^{V[G]}$  which fixes the ordinals.*

**Corollary 4.16.** *Suppose there is an inner model as in the previous theorem. Then proper forcing of size  $\leq \kappa$  does not add equivalence classes to thin  $(\mathbf{\Pi}_1^2)^{L(\mathbb{R})}$  equivalence relations.*

*Proof.* The assumption implies  $\text{AD}^{L(\mathbb{R})}$  by [42, lemma 7.15] and Woodin's theorem [31, theorem 3.1]:  $\text{AD}^{L(\mathbb{R}^*)}$  holds if  $\lambda$  is a limit of Woodin cardinals and  $\mathbb{R}^*$  is the set of reals of a symmetric collapse below  $\lambda$ . Now suppose  $E$  is a thin  $(\mathbf{\Pi}_1^2)^{L(\mathbb{R})}$  equivalence relation, so  $E$  is co- $(\delta_1^2)^{L(\mathbb{R})}$ -Suslin for via the tree  $T$  from a  $(\mathbf{\Sigma}_1^2)^{L(\mathbb{R})}$  scale on the complement of  $E$ . Let  $G$  be generic over  $V$  for a proper forcing of size  $\leq \kappa$ . Then there is an elementary embedding

$$j : L(\mathbb{R}) \rightarrow L(\mathbb{R})^{V[G]}$$

which fixes the ordinals by the previous theorem, in particular  $\mathbb{R}^2 - p[T]$  is an equivalence relation in every Cohen generic extension of  $V$ . Hence there is a

wellorder of the equivalence classes of  $E$  in  $L(\mathbb{R})$  by corollary 4.13. Since  $j$  fixes the ordinals, there are no new equivalence classes in  $V[G]$ .  $\square$

**4.2. Projective equivalence relations.** In this section we use the theorem of Harrington and Shelah to determine the number of equivalence classes of thin projective equivalence relations relative to the projective ordinals, assuming PD.

4.2.1.  $\Pi_n^1$  and  $\Sigma_{2n+1}^1$  equivalence relations. Silver [39] proved

**Lemma 4.17.** *Assume ZF. Then every thin  $\Pi_1^1$  equivalence relation has countably many equivalence classes.*

Harrington's simpler proof of this result can be found in Jech [16, theorem 32.1]. The lemma follows from corollary 4.13, since  $\Sigma_1^1$  sets are  $\aleph_0$ -Suslin and the statement that a  $\Pi_1^1$  formula defines an equivalence relation is  $\Pi_2^1$  and hence absolute. Burgess proved

**Lemma 4.18.** *Assume ZF. Then every thin  $\Sigma_1^1$  equivalence relation has at most  $\aleph_1$  many equivalence classes.*

For a proof see Jech [16, theorem 32.9]. This result is a consequence of corollary 4.13, since  $\Sigma_2^1$  sets are  $\aleph_1$ -Suslin and the Shoenfield tree projects to the complete  $\Sigma_2^1$  set in any Cohen generic extension. For the same reason one has

**Lemma 4.19.** *Assume ZF and*

- (1) *there is a Cohen real over  $L[x]$  or*
- (2) *there is an inner model which satisfies generic  $\Sigma_3^1$  Cohen absoluteness.*

*Then any thin  $\Pi_2^1(x)$  equivalence relation has at most  $\aleph_1$  many equivalence classes.*

The conclusion from condition 1 is shown in in Harrington and Shelah [9]. Note that 2 implies 1 by Bartoszynski and Judah [2, theorems 9.2.12 and 9.2.1]. The previous facts generalize through the projective hierarchy:

**Theorem 4.20.** *Assume ZF and*

- (1)  $\Pi_{2n+1}^1$  *is scaled and*
- (2) *all  $\Delta_{2n+2}^1$  sets have the Baire property.*

*Then the equivalence classes of any thin  $\Pi_{2n+2}^1$  equivalence relation can be well-ordered with order type  $\leq \delta_{2n+1}^1$ . Moreover, there is a thin  $\Sigma_{2n+1}^1$  equivalence relation whose equivalence classes can be wellordered with order type  $\delta_{2n+1}^1$ .*

*Proof.* Let  $E$  be a thin  $\Pi_{2n+2}^1$  equivalence relation and fix some  $\Sigma_{2n+2}^1$ -scale  $(\leq_k: k < \omega)$  of length  $\delta_{2n+1}^1$  on  $\mathbb{R}^2 - E$ . In fact [30, 4C.14] and [21, theorem 38.4] imply that there is a scale of this length. We further have  $\Sigma_{2n+3}^1$  Cohen absoluteness by lemma 3.12. Then in every Cohen generic extension

- (1)  $E$  is an equivalence relation and
- (2)  $(\leq_k: k < \omega)$  is a scale on  $\mathbb{R}^2 - E$ ,

since both are  $\Pi_{2n+3}^1$  statements. Here  $E$  and  $(\leq_k: k < \omega)$  are understood as the corresponding sets in the generic extension with the same definition as in the ground model. Now Cohen forcing does not change the tree  $T$  from the scale since no new equivalence classes are added to the relevant prewellorders by lemma 3.18. Hence  $\mathbb{R}^2 - p[T]$  is an equivalence relation in any Cohen generic extension. Thus the equivalence classes of  $E$  can be wellordered with order type  $\leq \delta_{2n+1}^1$  by corollary 4.13.

Let  $\leq$  be a  $\Pi_{2n+1}^1$  norm on the complete  $\Pi_{2n+1}^1$  set  $A \subseteq \mathbb{R}$ , so  $\leq$  has length  $\delta_{2n+1}^1$ . The norm induces a  $\Sigma_{2n+1}^1$  equivalence relation  $E$  defined by  $(x, y) \in E$  if and only if  $(x \leq y \wedge y \leq x) \vee x, y \notin A$ . Moreover,  $E$  is thin by the argument in lemma 2.14. □

The previous theorem implies that there is no difference in the possible number of equivalence classes of thin  $\Sigma_{2n+1}^1$  equivalence relations and of thin  $\Pi_{2n+2}^1$  equivalence relations. On the other hand, we will see in theorem 5.26 that there are

inner models which have representatives in every equivalence class of every thin  $\Sigma_{2n+1}^1$  equivalence relation defined from a parameter in the inner model, but do not fulfill the same condition for thin  $\Pi_{2n+2}^1$  equivalence relations. For the odd levels we have

**Theorem 4.21.** *Assume ZF and*

- (1)  $\Pi_{2n+1}^1$  *is scaled and*
- (2) *all  $\Delta_{2n+2}^1$  sets have the Baire property.*

*Then the equivalence classes of any thin  $\Pi_{2n+1}^1$  equivalence relation can be well-ordered with order type  $< \delta_{2n+1}^1$ . Moreover, for every  $\alpha < \delta_{2n+1}^1$  there is a thin  $\Delta_{2n+1}^1$  equivalence relation whose equivalence classes can be wellordered with order type at least  $\alpha$ .*

*Proof.* Let  $E$  be a thin  $\Pi_{2n+1}^1$  equivalence relation. Let further  $A \subseteq \mathbb{R}^3$  be a  $\Pi_{2n}^1$  set with  $\mathbb{R}^2 - E = p[A]$  and fix a  $\Pi_{2n+1}^1$ -scale  $(\leq_k: k < \omega)$  on  $A$ . Then the prewellorders are actually  $\Delta_{2n+1}^1$ , since  $A$  is  $\Pi_{2n}^1$ . Since  $cf(\delta_{2n+1}^1) > \omega$ , this implies that the length  $\alpha$  of the scale is less than  $\delta_{2n+1}^1$ . Now  $\Sigma_{2n+3}^1$  Cohen absoluteness holds by lemma 3.12. So the  $\Pi_{2n+2}^1$  statements

- (1)  $E$  is an equivalence relation,
- (2)  $\mathbb{R}^2 - E = p[A]$ , and
- (3)  $(\leq_n: n < \omega)$  is a scale on  $A$ ,

hold in every Cohen generic extension. Cohen forcing does not change the tree  $T$  from the scale, since no new equivalence classes are added to the relevant prewellorders by lemma 3.18. Hence  $\mathbb{R}^2 - p[T]$  is an equivalence relation in any Cohen generic extension. Thus the equivalence classes of  $E$  can be wellordered with order type at most  $\alpha$  by corollary 4.13.

Clearly for every  $\alpha < \delta_{2n+1}^1$  there is a  $\Delta_{2n+1}^1$  prewellorder with order type at least  $\alpha$ , by the definition of  $\delta_{2n+1}^1$ . □

The extra assumptions in the previous theorems cannot be eliminated:

**Lemma 4.22.** *Let  $n < \omega$  and assume there are  $2n$  Woodin cardinals and a measurable above if  $n > 0$ . Then under  $\text{ZFC} + \text{Det}(\mathbf{\Pi}_{2n}^1)$  there is no upper bound for the number of equivalence classes of thin  $\Sigma_{2n+2}^1$  equivalence relations.*

*Proof.*  $M_{2n}$  denotes the inner class model defined by iterating the top extender of  $M_{2n}^\#$  out of the universe. In particular  $M_0 = L$ . Then  $M_{2n} \models \text{Det}(\mathbf{\Pi}_{2n}^1)$  by lemma 2.30.

Let  $\kappa$  be an uncountable cardinal in  $M_{2n}$  and  $G$  a generic filter over  $M_{2n}$  for the finite support product of  $(\kappa^+)^{M_{2n}}$  many Cohen forcings. Working in  $M_{2n}[G]$ , fix a set  $A \subseteq \mathbb{R}$  of size  $\kappa$ . We claim that there is a c.c.c. forcing  $\mathbb{P}$  in  $M_{2n}[G]$  such that  $A$  is  $\mathbf{\Pi}_{2n+2}^1$  in any  $\mathbb{P}$ -generic extension of  $M_{2n}[G]$ . For  $n = 0$  this is Harrington's forcing from [6, §1]. The forcing has to be adapted if  $n > 0$ ; in this case we have to find a sequence  $(d_{\alpha,n} : \alpha < \omega_1, n < \omega)$  of distinct reals in  $M_{2n}[G]$  which is  $\Delta_{2n+1}^{HC}$  over  $M_{2n}[G]$ .

As for the case  $n = 0$  we work with the sequence of all reals of  $M_{2n}$  in the order of constructibility. The canonical wellorder is shown to be  $\Delta_{2n+1}^{HC}$  over  $M_{2n}$  in [40, theorem 4.5] by comparing reals in  $\Pi_{2n}$ -iterable,  $2n$ -small,  $\omega$ -sound premice. The point is that  $\mathcal{M} \trianglelefteq M_{2n}^\#$  for such premice  $\mathcal{M}$  by [40, lemma 3.3]. Since  $\Pi_{2n}$ -iterability is  $\Pi_{2n+1}^1$  in the codes, it is absolute between  $M_{2n}$  and  $M_{2n}[G]$ . It follows that the sequence of reals of  $M_{2n}$  in the order of constructibility is  $\Delta_{2n+1}^{HC}$  over  $M_{2n}[G]$ .  $\square$

The number of equivalence classes of thin  $\mathbf{\Pi}_n^1$  and  $\Sigma_{2n+1}^1$  equivalence relations can be calculated under PD by the two theorems above. The next example defines a  $\Delta_3^1$  equivalence relation with exactly  $\text{Card}(\delta_2^1)$  many equivalence classes from a prewellorder. Hence it is consistent that there is a  $\Delta_3^1$  equivalence relation with  $\aleph_2$  many equivalence classes by lemma 2.18.

**Example 4.23.** Assume  $x^\#$  exists for every  $x \in \mathbb{R}$ . Let  $(\iota_\alpha^x : \alpha \in \text{Ord})$  enumerate the  $x$ -indiscernibles and define  $u_2^x := \iota_{\omega_1+1}^x$  for  $x \in \mathbb{R}$ . The prewellorder given by

$$x \leq y :\Leftrightarrow u_2^x \leq u_2^y$$

is  $\Delta_3^1$  and its length is  $\delta_2^1$ .

*Proof.* Note that the class of  $x$ -indiscernibles and the theory of  $L[x]$  are definable from  $x^\#$  since

$$x^\# = \{\ulcorner \varphi(v_0, \dots, v_n) \urcorner : L[x] \models \varphi(x, \omega_1^V, \dots, \omega_n^V)\}.$$

Thus  $u_2^x \leq u_2^y$  holds if and only if

$$L[x^\#, y^\#] \models \iota_{\omega_1^V+1}^x \leq \iota_{\omega_1^V+1}^y.$$

But this can be calculated from  $(x^\#, y^\#)^\#$  since  $\omega_1^V$  is an  $(x^\#, y^\#)$ -indiscernible. Now sharps for reals are defined by a  $\Pi_2^1$  formula. Hence  $u_2^x \leq u_2^y$  is  $\Delta_3^1$  in  $x, y$ .

The length of the prewellorder is  $\delta_2^1$  since  $\delta_2^1 = u_2 = \sup\{u_2^x : x \in \mathbb{R}\}$  in the presence of sharps for reals.  $\square$

A similar example for  $\Delta_5^1$  is not known. It could be possible to realize this by comparing the heights of the transitive direct limit of iterates of  $M_2^\#(x)$  and  $M_2^\#(y)$  via iteration trees living below the respective least Woodin cardinal.

4.2.2.  $\Sigma_{2n}^1$  equivalence relations. Hjorth [12, lemma 2.5] showed that every thin  $\Sigma_2^1(x)$  equivalence relation is  $\Pi_2^1$  in any real coding  $M_1^\#$ , assuming  $M_1^\#(x)$  exists and is  $\omega_1 + 1$ -iterable. This also works assuming  $M_1^\#(x)$  exists for every  $x \in \mathbb{R}$  via lemma 2.38. In this section this result and its proof are extended to the even levels of the projective hierarchy. The main ingredient is the next lemma, based on [12, lemma 2.2].

If  $\mathbb{P}$  is a forcing and  $\tau$  is a  $\mathbb{P}$ -name for a real, then in any  $\mathbb{P} \times \mathbb{P}$ -generic extension there are two corresponding reals from the  $\mathbb{P}$ -generic filters. We will denote the  $\mathbb{P} \times \mathbb{P}$ -names derived from  $\tau$  for these reals by  $\tau$  and  $\tau'$  throughout the rest of this paper.

**Lemma 4.24.** *Let  $n$  be even and  $k \geq n$ , Suppose  $M_k^\#(x)$  exists for every  $x \in \mathbb{R}$ . Let  $E \subseteq \mathbb{R}^2$  be a thin  $\Pi_{n+3}^1(x)$  equivalence relation where  $x \in \mathbb{R}$ . Let  $\mathcal{M}$  be a countable  $(k+1)$ -small  $X$ -premouse which is  $\omega_1$ -iterable above  $\delta$  and  $\omega$ -sound above  $\delta$  with  $\rho_\omega(\mathcal{M}) \leq \delta$ , where  $X$  is swo. Suppose there are  $n$  Woodin cardinals above  $\delta$  and an extender above them in  $\mathcal{M}$ . Let further  $\mathbb{P}$  be a forcing of size  $\leq \delta$  in  $\mathcal{M}$ . Then for every  $\mathbb{P}$ -name  $\tau \in \mathcal{M}$  for a real the set  $D$  of conditions  $p \in \mathbb{P}$  with*

$$(p, p) \Vdash_{\mathbb{P} \times \mathbb{P}}^{\mathcal{M}} \tau E \tau'$$

is dense in  $\mathbb{P}$ .

*Proof.* Suppose  $D$  is not dense. In this case let  $p_\emptyset \in \mathbb{P}$  be a condition such that for every  $q \leq p_\emptyset$  there are conditions  $r, u \leq q$  with

$$(r, u) \Vdash_{\mathbb{P} \times \mathbb{P}}^{\mathcal{M}} \neg \tau E \tau'.$$

Let  $(D_i : i < \omega)$  enumerate the dense open subsets of  $\mathbb{P} \times \mathbb{P}$  in  $\mathcal{M}$ . One can inductively define a family  $(p_s : s \in 2^{<\omega})$  of conditions in  $\mathbb{P}$  such that for all  $s, t \in 2^{<\omega}$

- (1)  $p_s \leq p_t$  if  $t \subseteq s$ ,
- (2)  $(p_{s \smallfrown 0}, p_{s \smallfrown 1}) \Vdash_{\mathbb{P} \times \mathbb{P}}^{\mathcal{M}} \neg \tau E \tau'$ ,
- (3)  $p_s$  decides  $\tau \upharpoonright lh(s)$ , and
- (4)  $(p_s, p_t) \in D_0 \cap \dots \cap D_i$  if  $s, t \in {}^i 2$  and  $s \neq t$ .

Moreover let

$$g_y := \{p \in \mathbb{P} : \exists n < \omega (p_{y \upharpoonright n} \leq p)\}$$

for each  $y \in \mathbb{R}$ . Then  $g_y \times g_z$  is  $\mathbb{P} \times \mathbb{P}$ -generic over  $\mathcal{M}$  for any  $y, z \in \mathbb{R}$  with  $y \neq z$  by condition 4. Then

$$\mathcal{M}[g_y, g_z] \models \neg \tau^{g_y} E \tau^{g_z}$$

by condition 2. We have  $\mathcal{M}[g_y, g_z] \prec_{\Sigma_{n+2}^1} V$  by lemma 2.45. Hence  $\neg \tau^{g_y} E \tau^{g_z}$  as  $\mathbb{R}^2 - E$  is  $\Sigma_{n+3}^1(x)$ . On the other hand the set  $P := \{\tau^{g_y} : y \in \mathbb{R}\}$  is perfect since  $\tau^{g_y}$  depends continuously on  $y$  by condition 2. This is a contradiction, since  $E$  is thin.  $\square$

We will need

**Lemma 4.25.** *Let  $n \leq k$  with  $k \geq 1$  and suppose  $M_k^\#(x)$  exists for every  $x \in \mathbb{R}$ . Let  $\mathcal{N}$  be a countable active  $\omega$ -sound  $\omega_1$ -iterable  $(k+1)$ -small  $X$ -premouse with  $\rho_\omega(\mathcal{N}) \leq \beta$ , where  $X$  is swo. Let  $\kappa$  be the critical point of the top extender of  $\mathcal{N}$  and  $\mathcal{M} := \mathcal{N}|\kappa$ . Let  $\delta > \beta$  be the least Woodin cardinal in  $\mathcal{N}$ . Let  $m < \omega$  be sufficiently large. Then there is a club  $C \subseteq \delta$  which is uniformly definable in  $\mathcal{M}$ , so that for every  $\gamma \in C$  we have for*

$$Y_\gamma := h_{\Sigma_m}^{\mathcal{M}}(V_\gamma^{\mathcal{M}})$$

and  $X_\gamma$  its transitive collapse that

- (1)  $X_\gamma \models$  " $\gamma$  is the least Woodin cardinal" and
- (2)  $X_\beta \triangleleft X_\gamma \triangleleft \mathcal{M}$  for all  $\beta \in C \cap \gamma$ .

*Proof.* We define a sequence  $(\gamma_\alpha : \alpha < \delta)$  by induction and then set

$$C := \{\gamma_\alpha : \alpha < \delta\}.$$

Note that  $\Sigma_1^{(m-1)}$  coincides with  $\Sigma_m$  over  $\mathcal{M}$  since  $\mathcal{M}$  is  $\omega$ -sound and is a model of ZF. To define  $\gamma_0$  let  $Y^0 := h_{\Sigma_1^{(m-1)}}^{\mathcal{M}}(\emptyset)$  via the canonical Skolem functions. Let

$$Y^{i+1} := h_{\Sigma_1^{(m-1)}}^{\mathcal{M}}(V_{\sup(Y^i \cap \delta)+1}^{\mathcal{M}})$$

for  $i < \omega$  and define

$$\gamma_0 := \sup\left(\bigcup_{i < \omega} Y^i \cap \delta\right).$$

We have  $\gamma_0 < \delta$  since  $\delta$  is inaccessible in  $\mathcal{M}$  and further

$$Y_{\gamma_0} = \bigcup_{i < \omega} Y^i = h_{\Sigma_1^{(m-1)}}^{\mathcal{M}}(V_{\gamma_0}^{\mathcal{M}}).$$

To define  $\gamma_{\alpha+1}$  in the successor step start with  $Y^0 := h_{\Sigma_1^{(m-1)}}^{\mathcal{M}}(V_{\sup(Y_{\gamma_\alpha} \cap \delta)+1}^{\mathcal{M}})$ . Let

$$Y^{i+1} := h_{\Sigma_1^{(m-1)}}^{\mathcal{M}}(V_{\sup(Y^i \cap \delta)+1}^{\mathcal{M}})$$

for  $i < \omega$  and let

$$\gamma_{\alpha+1} := \sup\left(\bigcup_{i < \omega} Y^i \cap \delta\right).$$

Again we have

$$Y_{\gamma_{\alpha+1}} = h_{\Sigma_1^{(m-1)}}^{\mathcal{M}}(V_{\gamma_{\alpha+1}}^{\mathcal{M}}).$$

For limits  $\mu < \delta$  define  $\gamma_\mu := \sup_{\alpha < \mu} \gamma_\alpha$ , so that  $Y_{\gamma_\mu} = h_{\Sigma_1^{(m-1)}}^{\mathcal{M}}(V_{\gamma_\mu}^{\mathcal{M}})$  as well. Since  $\delta$  is inaccessible in  $\mathcal{M}$ , we have  $\gamma_\alpha < \delta$  for each  $\alpha < \delta$ .

Let  $X_{\gamma_\alpha}$  be the transitive collapse of  $Y_{\gamma_\alpha}$  for  $\alpha < \delta$  and let  $\sigma_\alpha : X_{\gamma_\alpha} \rightarrow Y_{\gamma_\alpha}$  be the uncollapsing map. Then  $\gamma_\alpha$  is the least Woodin cardinal in  $X_{\gamma_\alpha}$ , since  $\pi_\alpha(\gamma_\alpha) = \delta$ . The construction ensures that  $J_{\gamma_\alpha}^{\vec{F}} = V_{\gamma_\alpha}^{\mathcal{M}}$  for each  $\alpha < \delta$  where  $\vec{F} = \vec{F}^{\mathcal{M}}$ .

Moreover  $X_{\gamma_\alpha}$  is  $m$ -sound above  $\gamma_\alpha$ , since  $Y_{\gamma_\alpha}$  is the  $\Sigma_1^{(m-1)}$ -hull of  $\gamma_\alpha$  in  $\mathcal{M}$ . It is clear that  $\text{crit}(\sigma_\alpha) = \gamma_\alpha$  and  $\rho_m(X_{\gamma_\alpha}) = \gamma_\alpha$ . Hence the condensation lemma can be applied, see [47, theorem 5.5.1] and [29, theorem 8.2]. We have  $F_{\gamma_\alpha}^{\mathcal{M}} = \emptyset$  since  $\gamma_\alpha$  is a cardinal. So the case that  $X_{\gamma_\alpha}$  is an ultrapower of an initial segment of  $\mathcal{M}$  by  $F_{\gamma_\alpha}$  can be ruled out. Hence  $X_{\gamma_\alpha} \trianglelefteq \mathcal{M}$ . One can now conclude that  $X_{\gamma_\alpha} \triangleleft X_{\gamma_\beta} \triangleleft \mathcal{M}$  for all  $\alpha < \beta < \delta$ .  $\square$

The previous lemma will also be used in the proof of the main lemma in section 4.1. For the next theorem we actually only need a single element of the club in the lemma. Hjorth [12, lemma 2.5] proved the next theorem for  $n = 1$ :

**Theorem 4.26.** *Let  $n \geq 1$  and suppose  $M_{2n-1}^\#(x)$  exists for every  $x \in \mathbb{R}$ . Then every thin  $\Sigma_{2n}^1(r)$  equivalence relation is  $\Pi_{2n}^1$  in any real coding  $M_{2n-1}^\#(r)$ , for  $r \in \mathbb{R}$ .*

*Proof.* Let  $E$  be a thin  $\Sigma_{2n}^1(r)$  equivalence relation. Define  $\mathcal{M} := M_{2n-1}^\#(r)$  and let  $\delta$  be the least Woodin cardinal in  $\mathcal{M}$ . Let  $\eta$  and  $\tau$  be  $\mathbb{W}_\delta$ -names in  $\mathcal{M}$  such that  $\Vdash_{\mathbb{W}_\delta}^{\mathcal{M}} \dot{x} = \eta \oplus \tau$ , where  $\dot{x}$  is a name for the  $\mathbb{W}_\delta$ -generic real. Then the set  $D$  of conditions  $p \in \mathbb{W}_\delta^{\mathcal{M}}$  with

$$(p, p) \Vdash_{\mathbb{W}_\delta \times \mathbb{W}_\delta}^{\mathcal{M}} \eta E \eta' \wedge \tau E \tau'$$

is dense in  $\mathbb{W}_\delta^{\mathcal{M}}$  by lemma 4.24. Let  $\kappa$  be the critical point of the top extender of  $\mathcal{M}$ . Let's choose some  $\gamma \in C$  where  $C \subseteq \delta$  is the club from the previous lemma. Let  $X_\gamma$  be the corresponding initial segment of  $\mathcal{M}$  with uncollapsing map  $\sigma : X_\gamma \rightarrow Y_\gamma$  and  $\sigma(\bar{D}) = D$ .

We claim that any two reals  $x$  and  $y$  are  $E$ -inequivalent if and only if there are

- (1) reals  $a$  and  $b$  and
- (2) an iteration tree on  $\mathcal{M}$  living on  $\mathcal{M}|_\gamma$  according to  $\Sigma$  with iteration map  $\pi : \mathcal{M} \rightarrow \mathcal{N}$

such that

- (1)  $a \oplus b$  is  $\mathbb{W}_{\pi(\gamma)}$ -generic over  $\mathcal{N}$ ,
- (2)  $\mathcal{N}[a, b] \models \neg a E b$ , and
- (3)  $a E x$  and  $b E y$ .

Condition 2 is equivalent to  $\neg aEb$  since  $\mathcal{N}[a, b] \prec_{\Sigma_{2n}^1} V$  by lemma 2.47. So these conditions imply that  $\neg xEy$ .

On the other hand, suppose  $\neg xEy$ . Let  $\mathcal{T}$  be a countable iteration tree on  $M_{2n-1}^\#(x)$  living below  $\gamma$  with iteration map  $\pi : \mathcal{M} \rightarrow \mathcal{N}$  such that  $x \oplus y$  is  $\mathbb{W}_{\pi(\gamma)}$ -generic over  $\pi(X)$  by lemma 2.44. Since  $\bar{D}$  is dense in  $\mathbb{W}_\gamma^{X_\gamma}$ , there is a condition  $p \in \pi(\bar{D})$  such that  $x \oplus y$  is  $\mathbb{W}_{\pi(\gamma)} \upharpoonright p$ -generic over  $\pi(X)$ . Now let  $a \oplus b$  be  $\mathbb{W}_{\pi(\gamma)} \upharpoonright p$ -generic over both  $\pi(X)[x, y]$  and over  $\mathcal{N}$ . We have

$$\pi(X)[a, x] \models aEx$$

$$\pi(X)[b, y] \models bEy,$$

since this is forced by  $(p, p)$ . Then lemma 2.45 shows that  $aEx$  and  $bEy$  hold, since there are  $2n - 2$  Woodin cardinals in  $\pi(X)[a, x]$  and in  $\pi(X)[b, y]$ . Thus  $\neg aEb$  and hence  $a, b, \pi$  satisfy the conditions.

It remains to show that the existence of  $a, b, \pi$  satisfying these conditions is a  $\Sigma_{2n}^1$  statement in any real coding  $\mathcal{M}$ . It suffices to check that the statement “ $\mathcal{T}$  is an iteration tree living on  $\mathcal{M} \upharpoonright \gamma$  according to  $\Sigma$ ” is  $\Sigma_{2n}^1$  in  $\mathcal{M}$ . Since  $X_\gamma$  is  $(2n - 1)$ -small, the  $\mathcal{Q}$ -structures for  $X_\gamma$  are  $(2n - 2)$ -small and hence  $\Pi_{2n-1}^1$  by lemma 2.48, so the statement is  $\Sigma_{2n}^1$  in  $\mathcal{M}$ .  $\square$

Note that Harrington and Sami [8, theorem 5] proved that every thin  $\Sigma_{2n}^1$  equivalence relation is  $\Delta_{2n}^1$  and every thin  $\Pi_{2n+1}^1$  equivalence relation is  $\Delta_{2n+1}^1$  from PD, without identifying the parameters.

## 5. INNER MODELS FOR THIN EQUIVALENCE RELATIONS

We consider transitive models  $M$  with the property that there are representatives in  $M$  for all equivalence classes of all thin  $\Pi_{2n}^1(z)$  equivalence relations defined from a real parameter  $z$  in  $M$ . We prove the main lemma in the first section and

use this to characterize the inner models with this property in the main theorem in the next section. The base theory is  $\text{ZF} + \text{DC}$ . In the last section, we will construct a transitive model satisfying the properties in the main theorem 5.15 with a similar method assuming the corresponding projective ordinal is below  $\omega_2$  in the base theory  $\text{ZFC}$ .

A first observation is that the set of reals of such a model has at least the size of  $\delta_{2n-1}^1$  if  $\Pi_{2n-1}^1$  determinacy holds by theorem 4.20 and computes  $\delta_{2n-1}^1$  correctly. Hjorth [10, theorem 3.1] characterized the inner models with this property for thin  $\Pi_2^1$  equivalence relations:

**Theorem 5.1.** (Hjorth [10]) *Assume  $x^\#$  exists for every  $x \in \mathbb{R}$ . The following are equivalent for an inner model  $M$ :*

- (1)  *$M$  has a representative in every equivalence class of every thin  $\Pi_2^1(x)$  equivalence relation for any  $x \in \mathbb{R} \cap M$ ,*
- (2) (a)  *$M \prec_{\Sigma_3^1} V$  and*  
 (b)  *$\omega_1^M = \omega_1^V$ .*

The suggestion how to extend this theorem to  $\Pi_{2n}^1$  is due to Greg Hjorth. We ask that the tree from the canonical  $\Pi_{2n+1}^1$  scale is computed correctly instead of  $\omega_1$ . However, the argument is more involved than the proof of theorem 5.1. The main lemma 5.14 shows that  $T_{2n+1}$  can be computed in an iterate of  $M_{2n}^\#$ . To prove this, we find a sequence of local Woodin cardinals below the least Woodin of  $M_{2n}^\#$  using lemma 4.25 and build an iteration tree to make reals generic at the local Woodins. A density argument will show that  $T_{2n+1}$  can be defined in the last model of the iteration tree. To apply the main lemma 5.14 in the proof of the main theorem, we assign an infinitary formula to a given real by the Harrington-Shelah theorem and express the existence of a real satisfying this formula in a projective way.

5.1. **The main lemma.** In this section we prove

**Main Lemma 5.2.** *Let  $n \geq 1$  and assume  $M_{2n}^\#(x)$  exists for every  $x \in \mathbb{R}$ . Suppose  $M$  is a transitive model of a sufficiently large finite fragment of ZF. Suppose  $\mathbb{R} \cap M$  is countable and  $M$  calculates  $M_{2n}^\#(x)$  correctly for each  $x \in \mathbb{R} \cap M$ . Let  $r \in \mathbb{R} \cap M$  and let  $\delta$  be the least Woodin cardinal in  $M_{2n}^\#(r)$ . There is a countable iteration tree on  $M_{2n}^\#(r)$  with iteration map  $\pi : M_{2n}^\#(r) \rightarrow \mathcal{N}$  so that  $T_{2n+1}^M$  is definable in  $\mathcal{N}$  uniformly in the parameter  $r \in \mathbb{R}$ .*

We will build an iteration tree on  $M_{2n}^\#(r)$  with last model  $\mathcal{N}$  and reconstruct  $T_{2n+1}^M$  in a  $Col(\omega, < \omega_1^M)$ -generic extension of  $\mathcal{N}$ . Let  $M$ ,  $r$ ,  $\delta$ , and  $n$  be as in the main lemma for the rest of this section.

**Claim 5.3.**  $M \prec_{\Sigma_{2n+1}^1} V$ .

*Proof.* Since every  $x \in \mathbb{R} \cap M$  is generic over some iterate of  $M_{2n}^\#(r)$  for the extender algebra at the least Woodin cardinal by lemma 2.44, one can construct  $L[\vec{E}, x]$  in this generic extension up to the critical point of the top extender. We then construct the  $M_{2n-1}^\#(x)$  of both  $M$  and  $V$  by attaching the top extender of the iterate of  $M_{2n}^\#(r)$  on top of this model. Now the claim follows from lemma 3.5.  $\square$

Let  $C \subseteq \delta$  be the club from lemma 4.25 applied to  $M_{2n}^\#(r)$ . Now the set

$$S := \{\gamma \in C : M_{2n}^\#(r) \models \gamma \text{ is inaccessible}\}$$

is stationary in  $\delta$ , since  $\delta$  is Mahlo in  $M_{2n}^\#(r)$ . Let  $\bar{S}$  be the set of limit points of  $S$  and

$$\lambda_r := \min(S \cap \bar{S}).$$

Let  $(\gamma_k : k < \omega)$  be a sequence in  $M$  of ordinals in  $S$  with supremum  $\lambda_r$ . We define  $X_k := X_{\gamma_k}$  for  $k < \omega$ . Note that each  $X_k$  is  $\omega_1$ -iterable via the  $\mathcal{Q}$ -structure iteration strategy  $\Sigma$ .

Now let  $(x_k : k < \omega)$  enumerate  $\mathbb{R} \cap M$  and set  $\mathcal{N}_0 := M_{2n}^\#(r)$ . We construct premice  $\mathcal{N}_k$  for  $k \geq 1$  and countable iteration trees  $\mathcal{T}_k$  on  $\mathcal{N}_k$  in  $M$  for  $k < \omega$  such that

- (1) the composition  $\mathcal{T}_0 \frown \dots \frown \mathcal{T}_k$  is an iteration tree according to  $\Sigma$  with map
 
$$\pi_{k+1} = \pi_{0,k+1} = \pi_{k,k+1} \circ \dots \circ \pi_{0,1} : \mathcal{N}_0 \rightarrow \mathcal{N}_{k+1},$$
- (2)  $x_k$  is  $\mathbb{W}_{\pi_{k+1}(\gamma_{k+1})}$ -generic over  $\pi_{k+1}(X_{k+1})$ , and
- (3)  $\mathcal{T}_k$  lives on  $\mathcal{N}_k \upharpoonright \pi_k(\gamma_{k+1})$  and all extenders in  $\mathcal{T}_k$  have critical points above  $\pi_k(\gamma_k)$ .

Suppose  $\mathcal{N}_k$  and  $\mathcal{T}_i$  have been defined for  $i < k$ . Note that  $\pi_k(X_{k+1}) \triangleleft \mathcal{N}_k$  and  $\pi_k(X_{k+1}) \upharpoonright \pi_k(\gamma_{k+1}) = \mathcal{N}_k \upharpoonright \pi_k(\gamma_{k+1})$ . There is a countable iteration tree  $\mathcal{T}_k$  on  $\mathcal{N}_k$  according to  $\Sigma$  with map  $\pi_{k,k+1}$  so that  $x_k$  is  $\mathbb{W}_{\pi_{k,k+1}(\pi_k(\gamma_{k+1}))}$ -generic over  $\pi_{k,k+1}(X_{k+1})$  by the genericity iteration. Here  $\mathcal{T}_k$  lives on  $\mathcal{N}_k \upharpoonright \pi_k(\gamma_{k+1})$  and all extenders have critical points above  $\pi_k(\gamma_k)$ . We define  $\mathcal{N}_{k+1}$  as the last model of  $\mathcal{T}_k$ .

One can easily check that the composition  $\mathcal{T}_0 \frown \dots \frown \mathcal{T}_k$  is an iteration tree on  $M_{2n}^\#(r)$ , since it follows from condition 3 and the rules of the iteration game that  $\mathcal{N}_k$  is the only model in  $\mathcal{T}_0 \frown \dots \frown \mathcal{T}_{k-1}$  to which an extender in  $\mathcal{T}_k$  can be applied. Since the composition  $\mathcal{T}$  of  $(\mathcal{T}_k : k < \omega)$  is according to  $\Sigma$ , the direct limit  $\mathcal{N}$  along the unique cofinal branch is wellfounded. Let  $\pi_{k,\omega} : \mathcal{N}_k \rightarrow \mathcal{N}$  denote the direct limit maps.

Note that it follows from condition 3 that  $\pi_k(\gamma_k) = \pi_{0,\omega}(\gamma_k)$  and

$$\pi_{k,\omega} \upharpoonright V_{\pi_k(\gamma_k)+\omega}^{\mathcal{N}_k} = id.$$

This implies that  $\mathbb{W}_{\pi_k(\gamma_k)}^{\mathcal{M}_k} = \mathbb{W}_{\pi_k(\gamma_k)}^{\pi_k(X_k)} = \mathbb{W}_{\pi_{0,\omega}(\gamma_k)}^{\pi_{0,\omega}(X_k)}$  and the forcing has the same subsets in  $\pi_k(X_k)$  and  $\pi_{0,\omega}(X_k)$ . Hence  $x_k$  is  $\mathbb{W}_{\pi_{0,\omega}(\gamma_{k+1})}^{\pi_{0,\omega}(X_{k+1})}$ -generic over  $\pi_{0,\omega}(X_{k+1})$ .

**Claim 5.4.**  $\sup_{k < \omega} \pi_k(\gamma_k) = \omega_1^M$ .

*Proof.* Since initial segments of  $\mathcal{T}$  are countable in  $M$ , we have  $\pi_k(\gamma_k) < \omega_1^M$  for every  $k < \omega$ . Thus  $\sup_{k < \omega} \pi_k(\gamma_k) \leq \omega_1^M$ .

To show that  $\sup_{k < \omega} \pi_k(\gamma_k) \geq \omega_1^M$ , suppose  $\alpha < \omega_1^M$  is given. Let  $x_k$  code  $\alpha$  where  $k < \omega$ . Then  $x_k$  is  $\mathbb{W}_{\pi_{k+1}(\gamma_{k+1})}$ -generic over  $\pi_{k+1}(X_{k+1})$ . Now  $\pi_{k+1}(\gamma_{k+2})$  is inaccessible in  $\pi_{k+1}(X_{k+2})$  and hence in  $\pi_{k+1}(X_{k+1}) \subseteq \pi_{k+1}(X_{k+2})$ . Thus it is still inaccessible in  $\pi_{k+1}(X_{k+1})[x_k]$ . This implies

$$\alpha < \omega_1^{\pi_{k+1}(X_{k+1})[x_k]} < \pi_{k+1}(\gamma_{k+2}) \leq \pi_{k+2}(\gamma_{k+2}).$$

□

If  $\mathbb{P}$  is a forcing and  $\tau$  is a  $\mathbb{P}$ -name for a real, then in any  $\mathbb{P} \times \mathbb{P}$ -generic extension  $\tau$  defines two reals via the  $\mathbb{P}$ -generic filters. Let  $\tau$  and  $\tau'$  be  $\mathbb{P} \times \mathbb{P}$ -names for these reals. Let  $\leq_i$  denote the  $\Pi_{2n+1}^1$  prewellorders from the canonical  $\Pi_{2n+1}^1$  scale on the complete  $\Pi_{2n+1}^1$  set  $A$  for  $i < \omega$ . We write  $\equiv_i$  for the induced thin  $\Sigma_{2n+1}^1$  equivalence relations, i.e.  $x \equiv_i y$  if and only if  $(x \leq_i y \wedge y \leq_i x) \vee x, y \notin A$ .

**Claim 5.5.** *Let  $\tau$  be a name for the  $\mathbb{W}_{\pi_k(\gamma_k)}$ -generic real. Then the set  $D_{j,k}$  of conditions  $p \in \mathbb{W}_{\pi_k(\gamma_k)}^{\pi_k(X_k)}$  with*

- (1)  $p$  decides  $\tau \upharpoonright j$  and
- (2)  $(p, p) \Vdash_{\mathbb{W}_{\pi_k(\gamma_k)}^{\pi_k(X_k)} \times \mathbb{W}_{\pi_k(\gamma_k)}} \tau \equiv_i \tau'$  for every  $i \leq j$

is dense for all  $j, k < \omega$ .

*Proof.* Let  $j, k < \omega$  and let  $\sigma$  be a name for the  $\mathbb{W}_\delta$ -generic real. Then the set of conditions  $p \in \mathbb{W}_\delta^{M_{2n}^\#(r)}$  with

$$(p, p) \Vdash_{\mathbb{W}_\delta \times \mathbb{W}_\delta}^{M_{2n}^\#(r)} \sigma \equiv_i \sigma'$$

is dense for every  $i < j$  by lemma 4.24. Since these sets are dense open and the set of conditions which decide  $\sigma \upharpoonright j$  is dense open, we have that their intersection is dense open. The claim follows by elementarity.  $\square$

We use the notation

$$[\alpha, \beta) := \{\gamma < \kappa : \alpha \leq \gamma < \beta\}$$

and

$$Col(\omega, [\alpha, \beta)) := \{f : \omega \times [\alpha, \beta) : \forall n < \omega \forall \gamma \in [\alpha, \beta) f(n, \gamma) < \gamma\}$$

ordered by reverse inclusion. Then

$$Col(\omega, < \beta) \cong Col(\omega, < \alpha) \times Col(\omega, [\alpha, \beta))$$

for all ordinals  $\alpha < \beta$ .

**Claim 5.6.** *There is a  $Col(\omega, < \omega_1^M)$ -generic filter  $g$  over  $\mathcal{N}$  in  $V$  with*

$$\mathbb{R}^{\mathcal{N}[g]} \subseteq M.$$

*Proof.* Let  $g_0$  be a  $Col(\omega, < \gamma_0)$ -generic filter over  $\mathcal{N}$  in  $M$ . Then  $\pi_2(\gamma_2)$  is inaccessible in  $\mathcal{N}[g_0]$  and hence  $\mathcal{P}(\pi_1(\gamma_1))^{\mathcal{N}[g_0]}$  is countable in  $M$ . Now let  $g_1$  be a  $Col(\omega, < \pi_1(\gamma_1))$ -generic filter over  $\mathcal{N}$  in  $M$  with

$$g_1 \cap Col(\omega, < \gamma_0) = g_0.$$

Similarly we choose  $Col(\omega, < \pi_k(\gamma_k))$ -generic filters  $g_k$  over  $\mathcal{N}$  with

$$g_{k+1} \cap Col(\omega, < \pi_k(\gamma_k))$$

for each  $k < \omega$ . Finally let  $g := \bigcup_{k < \omega} g_k$ .

To see that  $g$  is  $Col(\omega, < \omega_1^M)$ -generic over  $\mathcal{N}$ , note that the forcing  $Col(\omega, < \omega_1^M)$  has the  $\omega_1^M$ -c.c. in  $\mathcal{N}$  since  $\omega_1^M$  is regular in  $\mathcal{N}$ . So for any maximal antichain

$A \subseteq \text{Col}(\omega, < \omega_1^M)$  there is some  $k < \omega$  with  $A \subseteq \text{Col}(\omega, < \pi_k(\gamma_k))$ . Hence  $g_k \cap A \neq \emptyset$ .

By considering only the nice names for reals we get  $\mathbb{R}^{\mathcal{N}[g]} = \bigcup_{k < \omega} \mathbb{R} \cap \mathcal{N}[g_k]$ .  $\square$

We fix a  $\text{Col}(\omega, < \omega_1^M)$ -generic filter  $g$  over  $\mathcal{N}$  as in the previous claim and let  $\mathbb{R}^* := \mathbb{R}^{\mathcal{N}[g]}$ .

**Claim 5.7.** *For all  $x \in \mathbb{R} \cap M$  and  $j < \omega$  there is a real  $y \in \mathbb{R}^*$  such that  $x \upharpoonright j = y \upharpoonright j$  and  $M \models x \equiv_i y$  for every  $i \leq j$ .*

*Proof.* Let  $x \in \mathbb{R} \cap M$  and find  $k < \omega$  with  $x = x_k$ . Let

$$\mathbb{P} := \mathbb{W}_{\pi_{k+1}(\gamma_{k+1})}^{\pi_{k+1}(X_{k+1})}.$$

Then  $x$  is  $\mathbb{P}$ -generic over  $\pi_{k+1}(X_{k+1})$ . Let  $D_{j,k+1}$  be the dense set from claim 5.5 and choose a condition  $p \in D_{j,k+1}$  in the generic filter for  $x$ . Since the set  $\mathcal{P}(\pi_{k+1}(\gamma_{k+1}))^{\pi_{k+1}(X_{k+1})}$  is countable in  $\mathcal{N}[g]$ , there is a  $\mathbb{P} \upharpoonright p$ -generic real  $y$  over  $\pi_{k+1}(X_{k+1})$  in  $\mathcal{N}[g]$ . We directly get  $x \upharpoonright j = y \upharpoonright j$  by choice of  $p$ .

We claim that  $x \equiv_i y$  for every  $i \leq j$ . To see this, choose another real  $z \in \mathbb{R} \cap M$  which is  $\mathbb{P} \upharpoonright p$ -generic over both  $\pi_{k+1}(X_{k+1})[x]$  and  $\pi_{k+1}(X_{k+1})[y]$ . Since

$$(p, p) \Vdash_{\mathbb{P} \times \mathbb{P}}^{\pi_{k+1}(X_{k+1})} \tau \equiv_i \tau',$$

we have

$$\pi_{k+1}(X_{k+1})[x, z] \models x \equiv_i z$$

and

$$\pi_{k+1}(X_{k+1})[y, z] \models y \equiv_i z$$

for each  $i \leq j$ . Now  $\pi_{k+1}(X_{k+1})[x, z]$  and  $\pi_{k+1}(X_{k+1})[y, z]$  are  $2n$ -small boldface premice with  $2n - 1$  Woodin cardinals above  $\pi_{k+1}(\gamma_{k+1})$  which are  $\omega_1$ -iterable

above  $\pi_{k+1}(\gamma_{k+1})$  in  $M$  and project to  $\pi_{k+1}(\gamma_{k+1})$  or below. Hence both are  $\Sigma_{2n}^1$ -correct in  $M$  by lemma 2.45. We can conclude that

$$M \models x \equiv_i z \equiv_i y$$

by  $\Sigma_{2n+1}^1$  upwards absoluteness.  $\square$

**Claim 5.8.**  $T_{2n+1}^M$  is definable from  $r$  in  $\mathcal{N}[g]$ .

*Proof.* We have  $M \prec_{\Sigma_{2n+1}^1} V$  by claim 5.3. Now  $\mathcal{N}[g]$  is a countable  $\omega$ -sound  $2n$ -small boldface premouse with  $2n-1$  Woodin cardinals above  $\omega_1^M$  and  $\rho_\omega(\mathcal{N}[g]) \leq \omega_1^M$  which is  $\omega_1$ -iterable above  $\omega_1^M$  in  $V$ . Moreover  $\mathcal{N}[g]$  computes  $\Sigma_{2n+1}^1$  truth in  $V$  by lemma 2.45. So for any  $x, y \in \mathbb{R} \cap \mathcal{N}[g]$  and  $k < \omega$  we can calculate in  $\mathcal{N}[g]$  whether  $V \models x \leq_k y$  holds. Using the previous claim, we can define  $T_{2n+1}^M$  in  $\mathcal{N}[g]$  in the parameter  $\pi_{0,\omega}(\lambda_r)$ , which was defined from  $r$ .  $\square$

By homogeneity of  $Col(\omega, < \omega_1^M)$  we get that  $T_{2n+1}^M$  is definable from  $r$  in  $\mathcal{N}$  and hence an element of  $\mathcal{N}$ .

**Remark 5.9.**  $\mathcal{N}[g] \not\prec_{\Sigma_{2n+3}^1} V$ .

*Proof.*  $M_{2n}^\#(r)$  is a  $\Pi_{2n+2}^1(r)$  singleton by 2.49. Supposing  $\mathcal{N}[g] \prec_{\Sigma_{2n+3}^1} V$  we would have  $M_{2n}^\#(r) \in \mathcal{N}[g]$ . But this implies  $M_{2n}^\#(r) \in \mathcal{N}$  by homogeneity of  $Col(\omega, < \omega_1^M)$  and hence  $M_{2n}^\#(r) \in M_{2n}^\#(r)$ , a contradiction.  $\square$

**Remark 5.10.** If  $M_{2n}^\#(X)$  exists for every  $X \in H_{(2^{\aleph_0})^+}$ , then the iterability of  $M_{2n}^\#$  is not affected by forcing with  $Col(\omega, \mathbb{R})$  by lemma 3.4. In this case one can construct the iteration tree in the proof of the main lemma for  $M = V$  in  $V^{Col(\omega, \mathbb{R})}$ . The construction produces a forcing extension  $\mathcal{N}[g]$  of an iterate of  $M_{2n}^\#$  in  $V^{Col(\omega, \mathbb{R})}$ .

We get a simpler version of the main lemma for  $n = 0$  based on

**Lemma 5.11.** *let  $M$  be a transitive model of ZF. Then  $T_1^M = T_1^V$  if and only if  $\omega_1^M = \omega_1^V$ .*

*Proof.* Since  $ht(T_1) = \omega_1$  we know that  $T_1^M = T_1^V$  implies  $\omega_1^M = \omega_1^V$ . On the other hand the Shoenfield tree is absolute relative to  $\omega_1$ . Moreover, the tree  $T_1$  from the canonical  $\Pi_1^1$  scale is absolute relative to the Shoenfield tree by the proof of [21, theorem 36.12].  $\square$

**Lemma 5.12.** *Let  $M$  be a transitive model of ZF. Suppose  $r \in \mathbb{R} \cap M$  and  $r^\#$  exists in  $M$ . Let  $\kappa$  be the critical point of the top extender of  $M_0^\#(r)$  and  $\pi : M_0^\#(r) \rightarrow \mathcal{N}$  the map from iterating the top extender in  $\omega_1^M$  many steps. Then*

$$\pi(\kappa) = \omega_1^M$$

and

$$T_1^{\mathcal{N}[g]} = T_1^M$$

for every  $Col(\omega, < \omega_1^M)$ -generic filter  $g$  over  $\mathcal{N}$ .

*Proof.* Since  $\omega_1^{\mathcal{N}[g]} = \omega_1^M$  we have  $T_1^{\mathcal{N}[g]} = T_1^M$  by the previous lemma.  $\square$

One can derive a different version of the main lemma for  $M_{2n}^\dagger(r)$  with essentially the same proof, based on the following version of lemma 4.24:

**Lemma 5.13.** *Suppose  $m \leq k$  and  $M_k^\#(x)$  exists for every  $x \in \mathbb{R}$ . Let  $E \subseteq \mathbb{R}^2$  be a thin  $\Pi_{m+3}^1(x)$  equivalence relation with  $x \in \mathbb{R}$ . Let  $\mathcal{M}$  be a countable  $(k+1)$ -small  $X$ -premouse which is  $\omega_1$ -iterable above  $\delta$  and  $\omega$ -sound above  $\delta$  with  $\rho_\omega(\mathcal{M}) \leq \delta$ , where  $X$  is swo. Suppose there are  $m$  Woodin cardinals above  $\delta$  and at least two extenders above them in  $\mathcal{M}$ . Let  $\mathbb{P}$  be a forcing of size  $\leq \delta$  in  $\mathcal{M}$ . Then for every  $\mathbb{P}$ -name  $\tau$  in  $\mathcal{M}$  for a real the set  $D$  of conditions  $p \in \mathbb{P}$  with*

$$(p, p) \Vdash_{\mathbb{P} \times \mathbb{P}}^{\mathcal{M}} \tau E \tau'$$

is dense in  $\mathbb{P}$ .

*Proof.* The proof is the same as for lemma 4.24. We need the extra extender on top to make sure that  $\mathcal{M}[g_y, g_z] \prec_{\Sigma_{m+2}} V$  by lemma 2.47, where  $g_y \times g_z$  is  $\mathbb{P} \times \mathbb{P}$ -generic over  $\mathcal{M}$ .  $\square$

**Lemma 5.14.** *Let  $n \geq 1$  and assume  $M_{2n}^\#(x)$  exists for every  $x \in \mathbb{R}$ . Suppose  $M$  is a transitive model of ZF such that  $\mathbb{R} \cap M$  is countable and  $M$  calculates  $M_{2n}^\#(x)$  correctly for each  $x \in \mathbb{R} \cap M$ . Moreover suppose  $r \in \mathbb{R} \cap M$  and  $M_{2n}^\dagger(r)$  exists in  $M$  and is calculated correctly. Let  $\delta$  be the least Woodin cardinal in  $M_{2n}^\dagger(r)$ . There are*

- (1) *a countable iteration tree on  $M_{2n}^\dagger(r)$  with iteration map  $\pi : M_{2n}^\dagger(r) \rightarrow \mathcal{N}$  and*
- (2) *an ordinal  $\lambda_r < \delta$  definable in  $M_{2n}^\dagger(r)$  uniformly in the parameter  $r \in \mathbb{R}$*

such that

$$\pi(\lambda_r) = \omega_1^M$$

and

$$T_{2n+1}^{\mathcal{N}[g]} = T_{2n+1}^M,$$

where  $g$  is any  $\text{Col}(\omega, < \omega_1^M)$ -generic filter over  $\mathcal{N}$ .

**5.2. The main theorem.** In this section we show

**Main Theorem 5.15.** *Suppose  $n \geq 1$  and  $M_{2n-2}^\#(x)$  exists for every  $x \in \mathbb{R}$ . Let  $(\leq_k : k < \omega)$  denote the canonical scale on the complete  $\Pi_{2n-1}^1$  set and let  $\equiv_k$  denote the induced thin  $\Sigma_{2n-1}^1$  equivalence relations. The following are equivalent for any transitive model  $M$  of ZF:*

- (1) *every equivalence class of every thin  $\Pi_{2n}^1(r)$  equivalence relation has a representative in  $M$  for all  $r \in \mathbb{R} \cap M$*

- (2) (a)  $M \prec_{\Sigma_{2n+1}^1} V$  and  
 (b)  $T_{2n-1}^M = T_{2n-1}^V$
- (3) (a)  $M \prec_{\Sigma_{2n+1}^1} V$  and  
 (b) for some  $k < \omega$  every equivalence class of  $\equiv_k$  has a representative in  $M$ .

Note that theorem 5.1 is the special case of the equivalence of conditions 1 and 2 for  $n = 0$  since  $T_1^M = T_1^V$  if and only if  $\omega_1^M = \omega_1^V$  by lemma 5.11.

The first part of the proof of the main theorem is purely descriptive. Note that the assumptions of the main theorem imply  $\text{Det}(\mathbf{\Pi}_{2n-1}^1)$  by theorem 2.30.

**Claim 5.16.** *Under the assumptions of the main theorem, condition 1 implies conditions 2 and 3.*

*Proof.* 2 (a). It suffices to show that  $A \cap M \neq \emptyset$  for every nonempty  $\Pi_{2n}^1(r)$  set  $A$  with  $r \in \mathbb{R} \cap M$ . Let  $\leq$  be the  $\Pi_{2n}^1(r)$  prewellorder from a  $\Sigma_{2n}^1(r)$  norm on  $\mathbb{R} - A$ . We have  $[x]_{\leq} = A$  for every  $x \in A$ , where

$$[x]_{\leq} := \{y \in \mathbb{R} : x \leq y \wedge y \leq x\}.$$

Then  $A \cap M \neq \emptyset$ , since the induced  $\Pi_{2n}^1(r)$  equivalence relation is thin by lemma 2.15.

2 (b). Condition 1 implies

$$\text{rank}_k^M(x) = \text{rank}_k^V(x)$$

for all  $x \in \mathbb{R} \cap M$  and  $k < \omega$  since each norm in the  $\Pi_{2n-1}^1$ -scale induces a thin  $\Sigma_{2n-1}^1$  equivalence relation. Hence  $T_{2n-1}^M \subseteq T_{2n-1}$ . We have to show that  $T_{2n-1} \subseteq T_{2n-1}^M$ .

Suppose  $(s, f) \in T_{2n-1}$  and  $m = lh(s) = lh(f)$ . Let  $A$  be the canonical complete  $\Pi_{2n-1}^1$  set. Choose  $x_0 \in A$  with

$$(s, f) = (x_0 \upharpoonright m, (rank_0(x_0), \dots, rank_{m-1}(x_0))).$$

We inductively define reals  $x_k \in A \cap M$  for  $1 \leq k \leq m$  with

$$(s \upharpoonright k, f \upharpoonright k) = (x_k \upharpoonright k, (rank_0(x_k), \dots, rank_{k-1}(x_k)))$$

so  $x_m$  witnesses that  $(s, f) \in T_{\Pi_{2n-1}^1}^M$ . Let  $\leq_k$  denote the  $k^{th}$   $\Sigma_{2n-1}^1$  prewellorder from the canonical  $\Pi_{2n-1}^1$ -scale on  $A$  and  $\equiv_k$  the induced equivalence relation.

Moreover let

$$U_t := \{x \in \mathbb{R} : x \upharpoonright lh(t) = t\}$$

for  $t \in \omega^{<\omega}$ .

**Case 1.**  $k = 1$ . Define  $x E_1 y$  if and only if

$$x, y \notin U_{s \upharpoonright 1} \vee (x, y \in U_{s \upharpoonright 1} \wedge x \equiv_0 y).$$

Then  $E_1$  is a  $\Sigma_{2n-1}^1$  equivalence relation which is thin by lemma 2.13, since it is induced by a  $\Sigma_{2n-1}^1$  prewellorder. There is a real  $x_1 \in \mathbb{R} \cap M$  with  $x_1(0) = s(0)$  and  $x_1 \equiv_0 x_0$  by condition 1 applied to  $E_1$ .

**Case 2.**  $2 \leq k \leq m$ . Suppose  $x_i \in \mathbb{R} \cap M$  is defined for  $1 \leq i < k$ . Then the set

$$U := \{x \in \mathbb{R} : \forall i < k - 1 (x \equiv_i x_{k-1})\}$$

is  $\Delta_{2n-1}^1(x_{k-1})$  since  $x_{k-1} \in A$ . Now define  $x E_k y$  if and only if

$$x, y \notin U_{s \upharpoonright k} \cap U \vee (x, y \in U_{s \upharpoonright k} \cap U \wedge x \equiv_{k-1} y).$$

Then  $E_k$  is a  $\Sigma_{2n-1}^1(x_{k-1})$  equivalence relation. It is thin since it is induced by a  $\Sigma_{2n-1}^1(x_{k-1})$  prewellorder. Moreover we have  $x_0 \in U_{s \upharpoonright k} \cap U$ . Hence there is a real

$x_k \in \mathbb{R} \cap M$  with  $x_k \upharpoonright k = s \upharpoonright k$  and  $x_k \equiv_i x_0$  for all  $1 \leq i < k$  by condition 1 for  $E_k$ .

3 (b). Since  $\equiv_k$  is a thin  $\Sigma_{2n-1}^1$  equivalence relation for each  $k < \omega$ .  $\square$

**Remark 5.17.** In condition 1 one can equivalently replace thin  $\Pi_{2n}^1$  equivalence relations by  $\Pi_{2n}^1$  prewellorders, thin  $\Pi_{2n}^1$  linear preorders, or  $\Sigma_{2n}^1$  norms.

For the other implications will use

**Lemma 5.18.** Suppose  $T$  is the tree from a scale on a set containing  $x \in \mathbb{R}$  and  $\mathbb{A}$  is strongly admissible with  $x, T \in \mathbb{A}$ . Then  $\text{rank}_k(x)$  is definable from  $x$  and  $T$  in  $\mathbb{A}$  for every  $k < \omega$ .

*Proof.* Recall the definition 5.18 strongly admissible. Let  $\alpha < \delta_{2n-1}^1$  be least with

$$\mathbb{A} \models \exists f \in {}^\omega(\delta_{2n-1}^1) ((x, f) \in [T] \wedge f(k) = \alpha).$$

We show that  $\text{rank}_k(x) = \alpha$ .

For  $\text{rank}_k(x) \leq \alpha$  suppose  $f \in {}^\omega(\delta_{2n-1}^1)$  and  $(x, f) \in [T]$ . Let  $(x_i : i < \omega)$  be a sequence of reals with

$$(x_i \upharpoonright i, (\text{rank}_0(x_i), \dots, \text{rank}_{i-1}(x_i))) = (x \upharpoonright i, (f(0), \dots, f(i-1))) \in [T]$$

for every  $i < \omega$ . Hence  $x_i \rightarrow x$  and the sequence  $(\text{rank}_k(x_i) : i < \omega)$  is eventually constant with value  $f(k)$  for each  $k < \omega$ . The semicontinuity of the scale implies  $\text{rank}_k(x) \leq f(k)$ .

For  $\alpha \leq \text{rank}_k(x)$  we have to find a function  $f \in {}^\omega(\delta_{2n-1}^1)$  in  $\mathbb{A}$  with  $(x, f) \in [T]$  and  $f(k) = \alpha$ . Such functions are exactly the branches of the tree

$$S := \{s \in {}^{<\omega}(\delta_{2n-1}^1) : (x \upharpoonright \text{lh}(s), s) \in T \wedge s(k) = \alpha\}.$$

In  $V$  the sequence  $(\text{rank}_k(x) : k < \omega)$  is the pointwise minimal branch of  $S$  by the semicontinuity of the scale. Now the wellfoundedness is expressible by a  $\Delta_1$  predicate. Hence  $S$  has a branch  $f$  in  $\mathbb{A}$  as well by  $\Delta_1$  absoluteness.  $\square$

The previous lemma shows that condition 2 in the main theorem implies condition 3. We shall now prove condition 1 from condition 3. Let  $r \in \mathbb{R} \cap M$  and let  $E$  be a thin  $\Pi_{2n}^1(r)$  equivalence relation. We fix a real  $x \in \mathbb{R}$ . The goal is to find a real  $\bar{x} \in \mathbb{R} \cap M$  with  $x E \bar{x}$ .

The idea of the proof is as follows. Let  $\mathbb{A}$  be the least strongly admissible set containing  $T_{2n-1}$  as an element and choose a formula  $\varphi \in \mathcal{L}_{\infty,0,N} \cap \mathbb{A}$  with  $\varphi(x)$  as in theorem 4.8. One can find a real  $t \in \mathbb{R} \cap M$  so that  $\varphi$  can be defined from  $t$  and  $T_{2n-1}$ . One can now use the main lemma to reconstruct  $T_{2n-1}$  in an iterate of  $M_{2n-2}^\#(t \oplus u)$  for arbitrary reals  $u \in \mathbb{R}$ . This allows you to express the existence of a real  $\bar{x}$  with  $\varphi(\bar{x})$  by a  $\Sigma_{2n+1}^1(x, r)$  statement. Since  $M$  is sufficiently correct, there is such a real in  $M$ . Finally the choice of  $\varphi$  implies that  $x E \bar{x}$ .

Let  $T$  be a tree defined from  $T_{2n-1}$  and  $r$  as in lemma 2.11 with  $E = \mathbb{R}^2 - p[T]$ . In order to apply the theorem of Harrington and Shelah we need to know that

**Claim 5.19.**  $\mathbb{R}^2 - p[T]$  is an equivalence relation in any Cohen generic extension of  $V$ .

*Proof.* Let  $G$  be Cohen generic over  $V$ . We get  $\Sigma_{2n+1}^1$  Cohen forcing absoluteness from  $\text{Det}(\mathbf{\Pi}_{2n-1}^1)$  by lemma 3.12. Then  $T_{2n-1}^V = T_{2n-1}^{V[G]}$  since Cohen forcing does not add equivalence classes to the relevant prewellorders by lemma 3.18. So  $\mathbb{R}^2 - p[T]$  is defined by the same  $\Pi_{2n}^1$  formula in  $V[G]$ . Hence this is an equivalence relation in  $V[G]$  by  $\Sigma_{2n+1}^1$  Cohen absoluteness.  $\square$

Suppose  $x \in \mathbb{R}$  and  $\mathbb{A}$  is the least admissible set with  $T_{2n-1} \in \mathbb{A}$ . Let  $\varphi \in \mathcal{L}_{\infty,0,N} \cap \mathbb{A}$  be a formula with

- (1)  $\varphi(x)$  and

(2)  $\varphi(y) \Rightarrow xEy$  for every  $y \in \mathbb{R}$

by theorem 4.8.

**Claim 5.20.** *There is a real  $t \in \mathbb{R} \cap M$  and a formula  $\psi$  which defines  $\varphi$  from  $T_{2n-1}$  and  $t$  in every strongly admissible set  $A$  with  $T_{2n-1}, t \in A$ .*

*Proof.* Let  $A$  be a minimal strongly admissible set containing  $T_{2n-1}$  and  $t$  as elements. Then  $A$  is the  $\Sigma_2$  Skolem hull of  $\mathcal{Q}_{2n-1}^1 \cup \{T_{2n-1}\}$  in itself by minimality of  $A$ , since the defining axioms for strongly admissible sets (KP with Axiom Beta) are  $\Pi_2$ . Now there is a  $\Sigma_2$  Skolem function for  $A$  which is uniformly  $\Sigma_3$  over  $A$  by [37, theorem 1.15] and the following paragraph. So  $\varphi$  is  $\Sigma_3$ -definable in  $A$  from  $T_{2n-1}$  and some  $\vec{\alpha} = (\alpha_0, \dots, \alpha_j) \in (\mathcal{Q}_{2n-1}^1)^{<\omega}$ . Since the length of each  $\Pi_{2n-1}^1$  norm in the scale is  $\mathcal{Q}_{2n-1}^1$ , we can choose reals  $t_i \in \mathbb{R} \cap M$  with  $\text{rank}_k(t_i) = \alpha_i$  for  $i \leq j$  by condition 3 (b). Then the join  $t := t_0 \oplus \dots \oplus t_j$  works by the previous lemma.  $\square$

Let  $t \in \mathbb{R} \cap M$  and  $\psi$  be as in the previous claim. Let  $\varphi_{s,S}$  denote the formula defined by  $\psi$  from a real  $s$  and a tree  $S$ . Let  $T_u$  be the term from the main lemma 5.14 which defines  $T_{2n-1}$  from  $u \in \mathbb{R}$  in an iterate of  $M_{2n-2}^\#(u)$ . Note that  $T_u$  does not depend on the model  $M$  in the main lemma.

**Claim 5.21.** *For every real  $u \in \mathbb{R}$  we have that  $\varphi(u)$  holds if and only if  $M_{2n-2}^\#(t \oplus u) \models \varphi_{t, T_u}(u)$ .*

*Proof.* Let  $N \prec V_\eta$  be countable and contain all relevant parameters, where  $\eta$  is a large limit ordinal. Let  $\bar{N}$  be its transitive collapse with uncollapsing map  $\pi : \bar{N} \rightarrow N$  and  $\pi(\bar{\varphi}) = \varphi$ . There is an iteration  $M_{2n-2}^\#(t \oplus u) \rightarrow \mathcal{N}$  with

$$T_u^\mathcal{N} = T_{2n-1}^{\bar{N}}$$

by the main lemma 5.14 applied to  $\bar{N}$ . Since claim 5.20 is true in  $\bar{N}$  we have

$$\varphi_{t, T_u}^{\mathcal{N}} = \varphi_{t, T_{2n-1}^{\bar{N}}}^{\mathcal{N}} = \bar{\varphi}.$$

Hence

$$\varphi(u) \Leftrightarrow \bar{\varphi}(u) \Leftrightarrow \mathcal{N} \models \varphi_{t, T_u}(u) \Leftrightarrow M_{2n-2}^{\#}(t \oplus u) \models \varphi_{t, T_u}(u).$$

The last equivalence holds by elementarity of the iteration map.  $\square$

Now the previous claim expresses the existence of a real  $\bar{x}$  with  $\varphi(\bar{x})$  by a  $\Sigma_{2n+1}^1$  formula, since  $M_{2n-2}^{\#}(t \oplus u)$  is a  $\Pi_{2n}^1(t \oplus u)$  singleton uniformly in  $t$  and  $u$  by lemma 2.49. Since  $M \prec_{\Sigma_{2n+1}^1} V$ , there is a real  $\bar{x} \in \mathbb{R} \cap M$  with  $\varphi(\bar{x})$ .

Note that Harrington's proof [16, theorem 32.1] of Silver's theorem shows

**Lemma 5.22.** *(Harrington) For every equivalence class  $[x]$  of every thin  $\Pi_1^1$  equivalence relation, there is a  $\Delta_1^1$  set  $X \neq \emptyset$  with  $X \subseteq [x]$ .*

The technique from the proof of the main theorem can be used to show a similar fact. Let  $\text{rank}_k$  denote the  $k^{\text{th}}$  rank in the canonical  $\Pi_{2n-1}^1$ -scale for  $k < \omega$ .

**Lemma 5.23.** *Let  $n \geq 1$  and suppose  $\text{Det}(\Pi_{2n-1}^1)$  holds. Let  $A \subseteq \mathbb{R}$  be closed under finite join  $\oplus$ . Suppose that for every  $\alpha < \delta_{2n-1}^1$  there are  $r \in A$  and  $k < \omega$  with  $\text{rank}_k(r) = \alpha$ . Then for every thin  $\Pi_{2n}^1$  equivalence relation  $E$  and every real  $x$ , there exist a real  $r \in A$  and a nonempty  $\Delta_{2n+1}^1(r)$  set  $X$  such that  $x \in X \subseteq [x]_E$ .*

*Proof.* Given a formula  $\varphi$  with  $\varphi(x)$  as in the proof of the main theorem, we can choose a real  $t \in A$  satisfying claim 5.20. It follows from claim 5.21 that the set  $\{u \in \mathbb{R} : \varphi(u)\}$  is  $\Delta_{2n+1}^1(t)$ .  $\square$

We conclude this section with two remarks about the proof of the main theorem.

**Remark 5.24.** *It is unclear whether condition 2 (b) in the main theorem can be replaced by  $\delta_{2n-1}^1 = (\delta_{2n-1}^1)^M$ .*

Note that if there is model satisfying the assumptions of the main theorem for which forcing with  $Col(\omega, \omega_1)$  does not change  $\delta_3^1$  and at the same time  $V \prec_{\Sigma_5^1} V^{Col(\omega, \omega_1)}$  holds, then the condition  $T_3^M = T_3^V$  cannot be replaced by  $(\delta_3^1)^M = \delta_3^1$  in the main theorem for  $n = 2$ . However, it is not clear how one could obtain such a model.

**Remark 5.25.** *Claim 5.21 can be used to prove from PD that every thin projective equivalence relation is induced by a projective prewellorder.*

Note that this fact is proved from in [8, theorem 5] from a determinacy assumption which is locally weaker than in this case. Suppose  $E$  is a thin equivalence relation and  $T$  is a tree as in the main theorem. Let  $x$  and  $y$  be reals. A prewellorder which induces  $E$  can be defined by comparing infinitary formulas  $\varphi$  with  $\varphi(x)$  and  $\psi$  with  $\psi(y)$  as above in the constructibility order of the least admissible set containing  $T$ .

**5.3. An inner model.** In this section we construct an inner model which fulfills similar conditions as in the main theorem 5.15, assuming the corresponding projective ordinal is below  $\omega_2$ .

**Theorem 5.26.** *Suppose  $n \geq 2$  is even and  $\Pi_{n+1}^1$  determinacy holds. Suppose  $\delta_{n+1}^1 < \omega_2$ . There is a forcing extension  $\mathcal{N}[g]$  in  $V$  of an iterate  $\mathcal{N}$  of  $M_n^\#$  of size  $\omega_1$  such that  $\mathcal{N}[g]$  has representatives in all equivalence classes of all thin  $\Sigma_{n+1}^1(z)$  equivalence relations with  $z \in \mathbb{R} \cap \mathcal{N}[g]$ . Moreover,  $M \prec_{\Sigma_{n+2}^1} V$  and  $M \not\prec_{\Sigma_{n+3}^1} V$ .*

To prove the theorem, we build a stack of  $\omega_1$  many iteration trees on an iterate of  $M_n^\#$  with direct limit model  $\mathcal{N}$  such that every real is generic over an initial segment of  $\mathcal{N}$ . We then form a  $Col(\omega, < \omega_1^V)$ -generic extension of  $\mathcal{N}$  in  $V$ . The

difference to the proof of the main lemma 5.14 is that the initial segments of the generic filter have to be defined in the course of the construction to ensure the required property for  $\mathcal{N}[g]$ .

As the first model in the stack of iteration trees we construct an iterate of  $M_n^\#$  which contains a sequence of local Woodin cardinals of order type  $\omega_1$ . Let  $\delta$  be the least Woodin cardinal in  $M_n^\#$  and let  $\bar{C}$  be the set of limit points  $< \delta$  of the club  $C \subseteq \delta$  from lemma 4.25. Note that the set of critical points of extenders on the  $M_n^\#$ -sequence below  $\delta$  is stationary in  $\delta$ , since  $\delta$  is Woodin in  $M_n^\#$ . Choose such a critical point  $\gamma \in \bar{C}$ . We define  $\mathcal{N}_0$  as the iterate of  $M_n^\#$  obtained by iterating the extender with critical point  $\gamma$  on the  $M_n^\#$ -sequence with least index  $\omega_1$  many times.

Then  $\mathcal{N}_0$  is  $\omega_1$ -iterable with respect to iteration trees living on  $\mathcal{N}_0|_{\omega_1}$ , since the relevant iteration maps commute by the argument in the commutativity lemma [5, lemma 3.2]. Moreover the image  $D$  of  $C \cap \gamma$  is a club in  $\omega_1$ . We enumerate  $D$  by  $(\gamma_\alpha : \alpha < \omega_1)$  and let  $(X_{\gamma_\alpha} : \alpha < \omega_1)$  be the corresponding initial segments of  $\mathcal{N}_0$  from lemma 4.25 such that  $\gamma_\alpha$  is Woodin in  $X_{\gamma_\alpha}$  for all  $\alpha < \omega_1$ .

As a bookkeeping device we fix a bijective map  $f : \omega_1 \rightarrow \omega_1 \times \omega_1$  with  $\eta \leq \alpha$  if  $f(\alpha) = (\zeta, \eta)$ . Let's inductively construct

- (1) a premouse  $\mathcal{N}_\alpha$ ,
- (2) a countable iteration tree  $\mathcal{T}_\alpha$  on  $\mathcal{N}_\alpha$ ,
- (3) a filter  $g_\alpha$ , and
- (4) a set  $R_\alpha = \{x_{\eta,\alpha} : \eta < \omega_1\} \subseteq \mathbb{R}$

for each  $\alpha < \omega_1$ , such that for all  $\beta < \omega_1$

- (1) the composition of  $(\mathcal{T}_\alpha : \alpha < \beta)$  is an iteration tree on  $\mathcal{N}_0$  according to  $\Sigma$  with map  $\pi_\beta = \pi_{0,\beta} : \mathcal{N}_0 \rightarrow \mathcal{N}_\beta$ ,
- (2)  $g_\beta$  is  $Col(\omega, < \pi_\beta(\gamma_\beta))$ -generic over  $\mathcal{N}_\beta$ ,

- (3)  $g_\alpha = g_\beta \cap Col(\omega, < \pi_\alpha(\gamma_\alpha))$  for  $\alpha < \beta$ ,
- (4)  $x_{f(\beta)}$  is  $\mathbb{W}_{\pi_{\beta+1}(\gamma_{\beta+1})}$ -generic over  $\pi_{\beta+1}(X_{\gamma_{\beta+1}})$ ,
- (5)  $\mathcal{T}_\beta$  lives on  $\mathcal{N}_\beta | \pi_\beta(\gamma_{\beta+1})$  and all extenders in  $\mathcal{T}_\beta$  have critical points above  $\pi_\beta(\gamma_\beta)$ , and
- (6) there is a representative in  $R_\beta$  for every equivalence class of every thin  $\Sigma_{n+1}^1(z)$  equivalence relation with  $z \in \mathbb{R} \cap \mathcal{N}_\beta[g_\beta]$ .

In each successor step  $\beta + 1 < \omega_1$  there is a set  $R_\beta \subseteq \mathbb{R}$  which fulfills condition 6, since every thin  $\Sigma_{n+1}^1$  (even provably  $\Delta_{n+2}^1$ ) equivalence relation has at most  $Card(\delta_{n+1}^1) \leq \omega_1$  many equivalence classes by theorem 4.20. Let  $(x_{\alpha,\beta} : \alpha < \omega_1)$  enumerate  $R_\beta$ . There is a countable iteration tree  $\mathcal{T}_\beta$  on  $\mathcal{N}_\beta$  living on  $\mathcal{N}_\beta | \pi_\beta(\gamma_{\beta+1})$  with iteration map  $\pi_{\beta,\beta+1}$  so that  $x_{f(\beta)}$  is  $\mathbb{W}_{\pi_{\beta+1}(\gamma_{\beta+1})}$ -generic over  $\pi_{\beta,\beta+1}(\pi_\beta(X_{\gamma_{\beta+1}}))$  by lemma 2.44. Let  $\mathcal{N}_{\beta+1}$  be the last model of  $\mathcal{T}_\beta$ . We further choose a  $Col(\omega, < \pi_{\beta+1}(\gamma_{\beta+1}))$ -generic filter  $g_{\beta+1}$  over  $\mathcal{N}_{\beta+1}[g_\beta]$  in  $V$ .

In each limit step  $\beta \leq \omega_1$  let  $\mathcal{N}_\beta$  be the direct limit of the unique cofinal branch in the composition of  $(\mathcal{T}_\alpha : \alpha < \beta)$ . We define  $g_\beta := \bigcup_{\alpha < \beta} g_\alpha$ . Then  $g_\beta$  is  $Col(\omega, < \pi_\beta(\gamma_\beta))$ -generic over  $\mathcal{N}_\beta$ . The reason is that  $\pi_\beta(\gamma_\beta)$  is inaccessible in  $\mathcal{N}_\beta$ , so  $Col(\omega, \pi_\beta(\gamma_\beta))$  has the  $\pi_\beta(\gamma_\beta)$ -c.c. in  $\mathcal{N}_\beta$ . Finally let  $\mathcal{N} := \mathcal{N}_{\omega_1}$ ,  $g := g_{\omega_1}$ , and

$$\mathbb{R}^* := \mathbb{R} \cap \mathcal{N}[g].$$

**Claim 5.27.** *Let  $r \in \mathbb{R}^*$  and suppose  $E$  is a thin  $\Sigma_{n+1}^1(r)$  equivalence relation. Then for every  $x \in \mathbb{R}$  there is some  $y \in \mathbb{R}^*$  with  $xEy$ .*

*Proof.* Let  $\alpha < \omega_1$  be an ordinal with  $r \in \mathbb{R} \cap \mathcal{N}[g_\alpha]$ . We have

$$\mathbb{R} \cap \mathcal{N}[g_\alpha] = \mathbb{R} \cap \mathcal{N}_{\alpha+1}[g_\alpha],$$

since the set of nice  $Col(\omega, < \pi_\alpha(\gamma_\alpha))$ -names for reals in  $\mathcal{N}$  is contained in  $\mathcal{N}_{\alpha+1}$ . We can assume that  $x \in R_\alpha$ . Now let  $\beta, \eta < \omega_1$  be ordinals with  $x = x_{\eta,\alpha}$  and

$f(\beta) = (\eta, \alpha)$ . Note that this implies  $\beta \geq \alpha$ . Then  $x$  is  $\mathbb{P}$ -generic over  $\pi_{\beta+1}(X_{\beta+1})$  for

$$\mathbb{P} := \mathbb{W}_{\pi_{\beta+1}(\gamma_{\beta+1})}^{\pi_{\beta+1}(X_{\beta+1})}.$$

Let  $\tau$  be a name for the  $\mathbb{P}$ -generic real. Then there is a condition  $p$  in the generic filter for  $x$  with

$$(p, p) \Vdash_{\mathbb{P} \times \mathbb{P}}^{\pi_{\beta+1}(X_{\beta+1})} \tau E \tau'$$

by lemma 4.24.

Let  $y \in \mathbb{R}^*$  be  $\mathbb{P} \upharpoonright p$ -generic over  $\pi_{\beta+1}(X_{\gamma_{\beta+1}})$ . Let further  $z \in \mathbb{R}$  be  $\mathbb{P} \upharpoonright p$ -generic over both  $\pi_{\beta+1}(X_{\beta+1})[x]$  and  $\pi_{\beta+1}(X_{\beta+1})[y]$ . Then

$$\pi_{\beta+1}(X_{\beta+1})[x, z] \models x E z$$

and

$$\pi_{\beta+1}(X_{\beta+1})[y, z] \models y E z$$

hold, since this is forced by  $(p, p)$ . Now  $\pi_{\beta+1}(X_{\beta+1})[x, z]$  and  $\pi_{\beta+1}(X_{\beta+1})[y, z]$  are both  $\Sigma_n^1$ -correct in  $V$  by lemma 2.47. Hence  $x$ ,  $y$ , and  $z$  are  $E$ -equivalent in  $V$ .  $\square$

**Claim 5.28.**  $\mathcal{N}[g] \prec_{\Sigma_{n+2}^1} V$ .

*Proof.* Let's assume  $k \leq n+1$  and  $\mathcal{N}[g] \prec_{\Sigma_k^1} V$ . Suppose  $\exists x \varphi(x, a)$  holds, where  $\varphi$  is a  $\Pi_k^1$  formula and  $a \in \mathbb{R} \cap \mathcal{N}[g]$ . Let  $E$  denote the thin  $\Sigma_{n+1}^1$  equivalence relation induced by a  $\Pi_{n+1}^1$  norm on  $\{x : \varphi(x, a)\}$ . Then there is a real  $x \in \mathbb{R} \cap \mathcal{N}[g]$  with  $\mathcal{N}[g] \models \varphi(x, a)$  by the previous claim.  $\square$

**Claim 5.29.**  $\mathcal{N}[g] \not\prec_{\Sigma_{n+3}^1} V$ .

*Proof.* Suppose  $\mathcal{N}[g] \prec_{\Sigma_{n+3}^1} V$ . Since  $M_n^\#$  is a  $\Pi_{n+2}^1$  singleton by lemma 2.49, this implies  $M_n^\# \in \mathcal{N}[g]$ . Thus  $M_n^\# \in \mathcal{N}$  by homogeneity. But  $\mathcal{N}$  is an iterate of  $M_n^\#$ , so this is impossible.  $\square$

$\mathcal{N}[g]$  can be cut off at the critical point of its top extender to get a model of ZFC with the properties of theorem 5.26. With the same proof, but with an application of lemma 5.13 instead of lemma 4.24, we have

**Theorem 5.30.** *Let  $n \geq 1$  and suppose  $\Pi_{n+1}^1$  determinacy holds. Suppose  $M_n^{\dagger}$  exists and  $\delta_{n+1}^1 < \omega_2$ . There is a forcing extension  $\mathcal{N}[g]$  in  $V$  of an iterate  $\mathcal{N}$  of  $M_n^{\dagger}$  of size  $\omega_1$  such that  $\mathcal{N}[g]$  has representatives in all equivalence classes of all thin provably  $\Delta_{n+2}^1(z)$  equivalence relations with  $z \in \mathbb{R} \cap \mathcal{N}[g]$ . Moreover,  $M \prec_{\Sigma_{n+2}^1} V$  and  $M \not\prec_{\Sigma_{n+3}^1} V$ .*

Note that every model constructed in this way has exactly  $\aleph_1$  many reals, hence the method cannot work without a condition on the projective ordinals.

#### REFERENCES

- [1] Joan Bagaria and Roger Bosch. Projective forcing. *Ann. Pure Appl. Logic*, 86(3):237–266, 1997.
- [2] Tomek Bartoszyński and Haim Judah. *Set theory. On the structure of the real line*. A K Peters Ltd., Wellesley, MA, 1995.
- [3] Jon Barwise. *Admissible sets and structures*. Perspectives in Mathematical Logic. Springer, Berlin, 1975.
- [4] Matthew Foreman and Menachem Magidor. Large cardinals and definable counterexamples to the continuum hypothesis. *Ann. Pure Appl. Logic*, 76(1):47–97, 1995.
- [5] Gunter Fuchs, Itay Neeman, and Ralf Schindler. A criterion for coarse iterability. *Archive for Mathematical Logic*, to appear.
- [6] Leo Harrington. Long projective wellorderings. *Ann. Math. Logic*, 12(1):1–24, 1977.
- [7] Leo Harrington. Analytic determinacy and  $0^\sharp$ . *J. Symbolic Logic*, 43(4):685–693, 1978.
- [8] Leo Harrington and Ramez Sami. Equivalence relations, projective and beyond. In *Logic Colloquium '78 (Mons, 1978)*, volume 97 of *Stud. Logic Foundations Math.*, pages 247–264. North-Holland, Amsterdam, 1979.
- [9] Leo Harrington and Saharon Shelah. Counting equivalence classes for co- $\kappa$ -Souslin equivalence relations. In *Logic Colloquium '80 (Prague, 1980)*, volume 108 of *Stud. Logic Foundations Math.*, pages 147–152. North-Holland, Amsterdam, 1982.

- [10] Greg Hjorth. Thin equivalence relations and effective decompositions. *J. Symbolic Logic*, 58(4):1153–1164, 1993.
- [11] Greg Hjorth. A dichotomy for the definable universe. *J. Symbolic Logic*, 60(4):1199–1207, 1995.
- [12] Greg Hjorth. Some applications of coarse inner model theory. *J. Symbolic Logic*, 62(2):337–365, 1997.
- [13] Greg Hjorth. Glimm-Effros for coanalytic equivalence relations. *J. Symbolic Logic*, 74(2):402–422, 2009.
- [14] Greg Hjorth and Alexander S. Kechris. Analytic equivalence relations and Ulm-type classifications. *J. Symbolic Logic*, 60(4):1273–1300, 1995.
- [15] Steve Jackson. A computation of  $\delta_5^1$ . *Mem. Amer. Math. Soc.*, 140(670), 1999.
- [16] Thomas Jech. *Set theory. The third millennium edition, revised and expanded*. Springer Monographs in Mathematics. Springer, Berlin, 2003.
- [17] Akihiro Kanamori. *The higher infinite. Large cardinals in set theory from their beginnings*. Perspectives in Mathematical Logic. Springer, Berlin, 1994.
- [18] Alexander S. Kechris. Forcing in analysis. In *Higher set theory (Proc. Conf., Math. Forschungsinst., Oberwolfach, 1977)*, volume 669 of *Lecture Notes in Math.*, pages 277–302. Springer, Berlin, 1978.
- [19] Alexander S. Kechris. On transfinite sequences of projective sets with an application to  $\Sigma_2^1$  equivalence relations. In *Logic Colloquium '77 (Proc. Conf., Wrocław, 1977)*, volume 96 of *Stud. Logic Foundations Math.*, pages 155–160. North-Holland, Amsterdam, 1978.
- [20] Alexander S. Kechris. AD and projective ordinals. In *Cabal Seminar 76–77 (Proc. Caltech-UCLA Logic Sem., 1976–77)*, volume 689 of *Lecture Notes in Math.*, pages 91–132. Springer, Berlin, 1978.
- [21] Alexander S. Kechris. *Classical descriptive set theory*, volume 156 of *Graduate Texts in Mathematics*. Springer, New York, 1995.
- [22] Alexander S. Kechris and Robert M. Solovay. On the relative consistency strength of determinacy hypotheses. *Trans. Amer. Math. Soc.*, 290(1):179–211, 1985.
- [23] Peter Koellner and W. Hugh Woodin. Large cardinals from determinacy. In Matthew Foreman and Akihiro Kanamori, editors, *Handbook of Set Theory*. Springer, Berlin, to appear.
- [24] Alain Louveau and Christian Rosendal. Complete analytic equivalence relations. *Trans. Amer. Math. Soc.*, 357(12):4839–4866 (electronic), 2005.

- [25] Donald A. Martin. Measurable cardinals and analytic games. *Fund. Math.*, 66:287–291, 1969/1970.
- [26] Donald A. Martin and John R. Steel. The extent of scales in  $L(\mathbb{R})$ . In *Cabal seminar 79–81*, volume 1019 of *Lecture Notes in Math.*, pages 86–96. Springer, Berlin, 1983.
- [27] Donald A. Martin and John R. Steel. A proof of projective determinacy. *J. Amer. Math. Soc.*, 2(1):71–125, 1989.
- [28] Donald A. Martin and John R. Steel. Iteration trees. *J. Amer. Math. Soc.*, 7(1):1–73, 1994.
- [29] William J. Mitchell and John R. Steel. *Fine structure and iteration trees*, volume 3 of *Lecture Notes in Logic*. Springer-Verlag, Berlin, 1994.
- [30] Yiannis N. Moschovakis. *Descriptive set theory*, volume 100 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Co., Amsterdam, 1980.
- [31] Itay Neeman. Optimal proofs of determinacy. *Bull. Symbolic Logic*, 1(3):327–339, 1995.
- [32] Itay Neeman. Determinacy in  $L(\mathbb{R})$ . In Matthew Foreman and Akihiro Kanamori, editors, *Handbook of Set Theory*. Springer, Berlin, 2010.
- [33] Itay Neeman and Jindřich Zapletal. Proper forcing and  $L(\mathbb{R})$ . *J. Symbolic Logic*, 66(2):801–810, 2001.
- [34] Ernest Schimmerling. A core model toolbox and guide. In Matthew Foreman and Akihiro Kanamori, editors, *Handbook of Set Theory*. Springer, Berlin, 2010.
- [35] Ralf Schindler. The role of absoluteness and correctness. slides from the Barcelona conference on set theory, Sept 16-20, 03, 1996. available at <http://wwwmath.uni-muenster.de/logik/Personen/rds>.
- [36] Ralf Schindler and John R. Steel. The self-iterability of  $L[E]$ . *J. Symbolic Logic*, 74(3):751–779, 2009.
- [37] Ralf Schindler and Martin Zeman. Fine structure. In Matthew Foreman and Akihiro Kanamori, editors, *Handbook of Set Theory*. Springer, Berlin, 2010.
- [38] Saharon Shelah. On  $\text{co-}\kappa$ -Souslin relations. *Israel J. Math.*, 47(2-3):139–153, 1984.
- [39] Jack H. Silver. Counting the number of equivalence classes of Borel and coanalytic equivalence relations. *Ann. Math. Logic*, 18(1):1–28, 1980.
- [40] John R. Steel. Projectively well-ordered inner models. *Ann. Pure Appl. Logic*, 74(1):77–104, 1995.
- [41] John R. Steel. Scales in  $K(\mathbb{R})$ . In *Games, scales, and Suslin cardinals. The Cabal Seminar. Vol. I*, volume 31 of *Lecture Notes in Logic*. Association for Symbolic Logic, Chicago, 2008.

- [42] John R. Steel. An outline of inner model theory. In *Handbook of Set Theory*. Springer, Berlin, 2010.
- [43] John R. Steel and Robert Van Wesep. Two consequences of determinacy consistent with choice. *Trans. Amer. Math. Soc.*, 272(1):67–85, 1982.
- [44] John R. Steel and Philip D. Welch.  $\Sigma_3^1$  absoluteness and the second uniform indiscernible. *Israel J. Math.*, 104:157–190, 1998.
- [45] W. Hugh Woodin. On the consistency strength of projective uniformization. In *Proceedings of the Herbrand symposium (Marseilles, 1981)*, volume 107 of *Stud. Logic Found. Math.*, pages 365–384. North-Holland, Amsterdam, 1982.
- [46] W. Hugh Woodin. *The axiom of determinacy, forcing axioms, and the nonstationary ideal*, volume 1 of *de Gruyter Series in Logic and its Applications*. Walter de Gruyter & Co., Berlin, 1999.
- [47] Martin Zeman. *Inner models and large cardinals*, volume 5 of *de Gruyter Series in Logic and its Applications*. Walter de Gruyter & Co., Berlin, 2002.

MATHEMATICAL INSTITUTE, UNIVERSITY OF BONN, ENDENICHER ALLEE 60, 53115 BONN,  
GERMANY, PHONE +49 228 24003752

*E-mail address:* `schlicht@math.uni-bonn.de`