

# ÉTALE COHOMOLOGY OF DIAMONDS

PETER SCHOLZE

ABSTRACT. Motivated by problems on the étale cohomology of Rapoport–Zink spaces and their generalizations, as well as Fargues’s geometrization conjecture for the local Langlands correspondence, we develop a six functor formalism for the étale cohomology of diamonds, and more generally small  $v$ -stacks on the category of perfectoid spaces of characteristic  $p$ . Using a natural functor from analytic adic spaces over  $\mathbf{Z}_p$  to diamonds which identifies étale sites, this induces a similar formalism in that setting, which in the noetherian setting recovers the formalism from Huber’s book, [Hub96].

## CONTENTS

1. Introduction	2
2. Spectral Spaces	9
3. Perfectoid Spaces	13
4. Set-theoretic bounds	19
5. Morphisms of perfectoid spaces	20
6. Étale morphisms	25
7. Totally disconnected spaces	28
8. The pro-étale and $v$ -topology	37
9. Descent	41
10. Morphisms of $v$ -stacks	47
11. Diamonds	51
12. Small $v$ -stacks	67
13. Spatial morphisms	74
14. Comparison of étale, pro-étale and $v$ -cohomology	78
15. Analytic adic spaces as diamonds	86
16. General base change results	89
17. Four functors	94
18. Proper and partially proper morphisms	98
19. Proper base change	105
20. Constructible sheaves	110
21. Dimensions	114
22. Proper pushforward	120
23. Cohomologically smooth morphisms	132
24. Examples of smooth morphisms	143
References	149

## 1. INTRODUCTION

The aim of this manuscript is to lay foundations for the six operations in étale cohomology of adic spaces and diamonds, generalizing previous work of Huber, [Hub96].

In this manuscript, we will deal with analytic adic spaces  $X$  on which a fixed prime  $p$  is topologically nilpotent. Associated with any such  $X$ , we have an étale site  $X_{\text{ét}}$  defined (under noetherian hypotheses) by Huber, [Hub96]. For any ring  $\Lambda$  such that  $n\Lambda = 0$  for some  $n$  prime to  $p$ , we get a (left-completed) derived category  $D_{\text{ét}}(X, \Lambda)$  of étale sheaves of  $\Lambda$ -modules on  $X_{\text{ét}}$ . It comes equipped with  $- \otimes_{\Lambda} -$  and  $R\mathcal{H}om_{\Lambda}(-, -)$ , and for a map  $f : Y \rightarrow X$ , one gets adjoint functors

$$f^* : D_{\text{ét}}(X, \Lambda) \rightarrow D_{\text{ét}}(Y, \Lambda)$$

and

$$Rf_* : D_{\text{ét}}(Y, \Lambda) \rightarrow D_{\text{ét}}(X, \Lambda) .$$

Moreover, under certain conditions, one can define the pushforward with proper supports

$$Rf_! : D_{\text{ét}}(Y, \Lambda) \rightarrow D_{\text{ét}}(X, \Lambda) ,$$

with a right adjoint

$$Rf^! : D_{\text{ét}}(X, \Lambda) \rightarrow D_{\text{ét}}(Y, \Lambda) .$$

If  $f$  is smooth, then (a form of) Poincaré duality identifies  $Rf^!$  as a twist of  $f^*$ . Under suitable hypotheses, as for example for the adic spaces corresponding to rigid spaces over a fixed complete nonarchimedean base field, these results all appear in [Hub96].

Our aim here is to give a substantial generalization of this formalism. The motivation comes from some developments surrounding the theory of “local Shimura varieties” whose cohomology is expected to realize local Langlands correspondences for  $p$ -adic fields, and in particular Fargues’ conjecture geometrizing the local Langlands correspondence through sheaves on the stack of  $G$ -bundles on the Fargues–Fontaine curve, [RV14, Far16, FS].

In this context, it is necessary to work with certain objects that are not representable by adic spaces, but are merely quotients of adic spaces by “pro-étale” equivalence relations, akin to Artin’s algebraic spaces. If one is willing to work with stacks, this brings objects like the classifying stack of  $\text{GL}_n(\mathbb{Q}_p)$  into the picture.

In fact, as any analytic adic space over  $\mathbf{Z}_p$  is itself a quotient of a perfectoid space by a pro-étale equivalence relation, one can as well work with quotients of *perfectoid* spaces by pro-étale equivalence relations. By tilting, one can even assume that these perfectoid spaces are of characteristic  $p$ . This leads to the notion of a diamond. Let us quickly review the definition.

**Definition 1.1.** *A map  $f : Y = \text{Spa}(S, S^+) \rightarrow X = \text{Spa}(R, R^+)$  of affinoid perfectoid spaces is affinoid pro-étale if it can be written as a cofiltered limit of étale maps  $Y_i = \text{Spa}(S_i, S_i^+) \rightarrow X$  of affinoid perfectoid spaces. More generally, a map  $f : Y \rightarrow X$  of perfectoid spaces is pro-étale if it is locally on the source and target affinoid pro-étale.*

Let  $\text{Perf}$  denote the category of perfectoid spaces of characteristic  $p$ .<sup>1</sup> There are several different topologies one can consider on it, successively refining the previous.

(i) The analytic topology, where a cover  $\{f_i : X_i \rightarrow X\}$  consists of open immersions  $X_i \hookrightarrow X$  which jointly cover  $X$ .

---

<sup>1</sup>For set-theoretic reasons, we (carefully) choose some cut-off cardinal  $\kappa$ , and work with  $\kappa$ -small perfectoid spaces throughout. The formalism does not depend on the choice of  $\kappa$ .

- (ii) The étale topology, where a cover  $\{f_i : X_i \rightarrow X\}$  consists of étale maps  $X_i \rightarrow X$  with jointly cover  $X$ .
- (iii) The pro-étale topology, where a cover  $\{f_i : X_i \rightarrow X\}$  consists of pro-étale maps  $X_i \rightarrow X$  such that for any quasicompact open subset  $U \subset X$ , there are finitely many indices  $i$  and quasicompact open subsets  $U_i \subset X_i$  such that the  $U_i$  jointly cover  $U$ .
- (iv) The v-topology, where a cover  $\{f_i : X_i \rightarrow X\}$  consists of any maps  $X_i \rightarrow X$  such that for any quasicompact open subset  $U \subset X$ , there are finitely many indices  $i$  and quasicompact open subsets  $U_i \subset X_i$  such that the  $U_i$  jointly cover  $U$ .

We note that the last part of the definition in cases (iii) and (iv) is the same condition that appears in the definition of the fpqc topology for schemes, and is automatic in cases (i) and (ii) as étale maps are open.

It turns out that all topologies are well-behaved; the v-topology is the finest, so for brevity we only state it in this case:

**Theorem 1.2.** *The v-topology on  $\text{Perf}$  is subcanonical, and for any affinoid perfectoid space  $X = \text{Spa}(R, R^+)$ ,  $H_v^0(X, \mathcal{O}_X) = R$ ,  $H_v^0(X, \mathcal{O}_X^+) = R^+$  and for  $i > 0$ ,  $H_v^i(X, \mathcal{O}_X) = 0$  and  $H_v^i(X, \mathcal{O}_X^+)$  is almost zero.*

Analogous to Artin’s theory of algebraic spaces, it is for the intended applications of the formalism necessary to allow quotients of perfectoid spaces by pro-étale equivalence relations. These are called diamonds:

**Definition 1.3.** *Let  $Y$  be a pro-étale sheaf on  $\text{Perf}$ . Then  $Y$  is a diamond if  $Y$  can be written as a quotient  $Y = X/R$  of a perfectoid space  $X$  by a pro-étale equivalence relation  $R \subset X \times X$ .*

It turns out that all diamonds are v-sheaves. Now, for a diamond  $X$ , one can define sites  $X_v$  and  $X_{\text{ét}}$ . Actually, in the case of  $X_{\text{ét}}$ , one has to be careful, as in general there may not be enough étale maps into  $X$ . For this reason (and others), we often work with a restricted class of diamonds which turn out to be better-behaved. Here, we use the underlying topological space of a diamond  $Y$ , given as  $|Y| = |X|/|R|$  in case  $Y = X/R$  is a quotient of a perfectoid space  $X$  by a pro-étale equivalence relation  $R$ . There is a bijection between open subspaces of  $|Y|$  and open subfunctors of  $Y$ , for an obvious definition of the latter. If  $Y$  is quasicompact as a v-sheaf, i.e. any collection of collectively surjective maps  $Y_i \rightarrow Y$  of v-sheaves has a finite subcover, then  $|Y|$  is quasicompact, but the converse is not true in general.

**Definition 1.4.** *A diamond  $Y$  is spatial if it is qcqs, and  $|Y|$  admits a basis for the topology given by  $|U|$ , where  $U \subset Y$  ranges over quasicompact open subdiamonds. More generally,  $Y$  is locally spatial if it admits an open cover by spatial diamonds.*

In particular, any perfectoid space  $X$  defines a locally spatial diamond, which is spatial precisely when  $X$  is qcqs. If  $X$  is a (locally) spatial diamond, then  $|X|$  is a (locally) spectral topological space, and  $X$  is quasicompact (resp. quasiseparated) as a v-sheaf precisely when  $|X|$  is quasicompact (resp. quasiseparated) as a topological space. Now, if  $X$  is a locally spatial diamond, we define  $X_{\text{ét}}$  as consisting of (locally separated) étale maps  $Y \rightarrow X$  from diamonds  $Y$  (automatically locally spatial); it turns out that this is “big enough”.

The following theorem shows that this generalizes the classical notions.

**Theorem 1.5.** *There is a natural functor from analytic adic spaces over  $\mathbf{Z}_p$  to locally spatial diamonds, denoted  $X \mapsto X^\diamond$ , satisfying  $X_{\text{ét}} \cong X_{\text{ét}}^\diamond$ .*

Here, the functor  $X \mapsto X^\diamond$  is intuitively given as follows. Choose a pro-étale surjection  $\tilde{X} \rightarrow X$  from a perfectoid space  $\tilde{X}$ , and let  $R \subset \tilde{X} \times \tilde{X}$  be the induced equivalence relation. This is pro-étale over  $\tilde{X}$ , and thus perfectoid itself. Then

$$X^\diamond = \tilde{X}^b / R^b .$$

**Theorem 1.6.** *Let  $X$  be a locally spatial diamond. Then pullback induces a fully faithful functor*

$$\widehat{D}(X_{\text{ét}}, \Lambda) \rightarrow D(X_v, \Lambda) ,$$

where  $\widehat{D}$  denotes the left-completion of the derived category;  $D(X_v, \Lambda)$  is already left-complete.

Moreover, if  $A \in D(X_v, \Lambda)$  and  $f : Y \rightarrow X$  is a map of locally spatial diamonds that is surjective as a map of  $v$ -sheaves, then if  $f^*A$  lies in  $\widehat{D}(Y_{\text{ét}}, \Lambda)$ , then  $A$  lies in  $\widehat{D}(X_{\text{ét}}, \Lambda)$ .

Much of the formalism actually extends to all  $v$ -sheaves, or even  $v$ -stacks, subject to a small set-theoretic assumption of being “small”.

**Definition 1.7.** *Let  $X$  be a small  $v$ -stack, and consider the site  $X_v$  of all perfectoid spaces over  $X$ , with the  $v$ -topology. Define the full subcategory*

$$D_{\text{ét}}(X, \Lambda) \subset D(X_v, \Lambda)$$

as consisting of all  $A \in D(X_v, \Lambda)$  such that for all (equivalently, one surjective) map  $f : Y \rightarrow X$  from a locally spatial diamond  $Y$ ,  $f^*A$  lies in  $\widehat{D}(Y_{\text{ét}}, \Lambda)$ .

Now we can state that we have the following operations.

(i) A (derived) tensor product

$$- \otimes_{\Lambda}^{\mathbb{L}} - : D_{\text{ét}}(X, \Lambda) \times D_{\text{ét}}(X, \Lambda) \rightarrow D_{\text{ét}}(X, \Lambda) .$$

This is compatible with the inclusions into  $D(X_v, \Lambda)$ , and the usual derived tensor product on  $X_v$ .

(ii) An internal Hom

$$R\mathcal{H}om_{\Lambda}(-, -) : D_{\text{ét}}(X, \Lambda)^{\text{op}} \times D_{\text{ét}}(X, \Lambda) \rightarrow D_{\text{ét}}(X, \Lambda)$$

characterized by the adjunction

$$R\text{Hom}_{D_{\text{ét}}(X, \Lambda)}(A, R\mathcal{H}om_{\Lambda}(B, C)) = R\text{Hom}_{D_{\text{ét}}(X, \Lambda)}(A \otimes_{\Lambda}^{\mathbb{L}} B, C)$$

for all  $A, B, C \in D_{\text{ét}}(X, \Lambda)$ . In particular, for  $A = \Lambda$ ,

$$R\Gamma(X, R\mathcal{H}om_{\Lambda}(B, C)) = R\text{Hom}_{D_{\text{ét}}(X, \Lambda)}(B, C) .$$

In general, the formation of  $R\mathcal{H}om_{\Lambda}$  does not commute with the inclusion  $D_{\text{ét}}(X, \Lambda) \subset D(X_v, \Lambda)$ .

(iii) For any map  $f : Y \rightarrow X$  of small  $v$ -stacks, a pullback functor

$$f^* : D_{\text{ét}}(X, \Lambda) \rightarrow D_{\text{ét}}(Y, \Lambda) .$$

This is compatible with the inclusions into  $D(X_v, \Lambda)$  resp.  $D(Y_v, \Lambda)$ , and the pullback functor  $D(X_v, \Lambda) \rightarrow D(Y_v, \Lambda)$ .

(iv) For any map  $f : Y \rightarrow X$  of small  $v$ -stacks, a pushforward functor

$$Rf_* : D_{\text{ét}}(Y, \Lambda) \rightarrow D_{\text{ét}}(X, \Lambda)$$

which is right adjoint to  $f^*$ . In general, formation of  $Rf_*$  does not commute with the inclusions into  $D(X_v, \Lambda)$  resp.  $D(Y_v, \Lambda)$ , but this holds true if  $f$  is qcqs and one starts with an object of  $D^+$ .

(v) For any map  $f : Y \rightarrow X$  of small v-stacks that is compactifiable (cf. Definition 22.2), representable in locally spatial diamonds (cf. Definition 13.3) and with  $\dim. \operatorname{trg} f < \infty$  (cf. Definition 21.7), a functor

$$Rf_! : D_{\text{ét}}(Y, \Lambda) \rightarrow D_{\text{ét}}(X, \Lambda) .$$

(vi) For any map  $f : Y \rightarrow X$  of small v-stacks that is compactifiable, representable in locally spatial diamonds and with  $\dim. \operatorname{trg} f < \infty$ , a functor

$$Rf^! : D_{\text{ét}}(X, \Lambda) \rightarrow D_{\text{ét}}(Y, \Lambda)$$

that is right adjoint to  $Rf_!$ .

The first four operations are defined in Section 17, and then  $Rf_!$  is defined in Section 22, in particular Definition 22.18, and  $Rf^!$  is defined in Section 23.

These operations satisfy the following formulas.

**Theorem 1.8.** *Let  $f : Y \rightarrow X$  be a map of small v-stacks.*

(i) *For all  $A, B \in D_{\text{ét}}(X, \Lambda)$ , one has*

$$f^* A \otimes_{\Lambda}^{\mathbb{L}} f^* B \cong f^*(A \otimes_{\Lambda}^{\mathbb{L}} B) .$$

(ii) *For all  $A \in D_{\text{ét}}(X, \Lambda)$ ,  $B \in D_{\text{ét}}(Y, \Lambda)$ , one has*

$$Rf_* R\mathcal{H}om_{\Lambda}(f^* A, B) \cong R\mathcal{H}om_{\Lambda}(A, Rf_* B) .$$

(iii) *Assume that  $f$  is compactifiable and representable in locally spatial diamonds with  $\dim. \operatorname{trg} f < \infty$ . For all  $A \in D_{\text{ét}}(X, \Lambda)$ ,  $B \in D_{\text{ét}}(Y, \Lambda)$ , one has*

$$Rf_!(A \otimes_{\Lambda}^{\mathbb{L}} f^* B) \cong Rf_! A \otimes_{\Lambda}^{\mathbb{L}} B .$$

(iv) *Assume that  $f$  is compactifiable and representable in locally spatial diamonds with  $\dim. \operatorname{trg} f < \infty$ . For all  $A \in D_{\text{ét}}(Y, \Lambda)$ ,  $B \in D_{\text{ét}}(X, \Lambda)$ , one has*

$$R\mathcal{H}om_{\Lambda}(Rf_! A, B) \cong Rf_* R\mathcal{H}om_{\Lambda}(A, Rf^! B) .$$

(v) *Assume that  $f$  is compactifiable and representable in locally spatial diamonds with  $\dim. \operatorname{trg} f < \infty$ . For all  $A, B \in D_{\text{ét}}(X, \Lambda)$ , one has*

$$Rf^! R\mathcal{H}om_{\Lambda}(A, B) \cong R\mathcal{H}om_{\Lambda}(f^* A, Rf^! B) .$$

Parts (i) and (ii) are proved in Section 17. Part (iii) is Proposition 22.23, and parts (iv) and (v) are formal consequences of (iii) and adjunctions, cf. Proposition 23.3.

Moreover, there are the following base change results.

**Theorem 1.9.** *Let*

$$\begin{array}{ccc} Y' & \xrightarrow{\tilde{g}} & Y \\ \downarrow f' & & \downarrow f \\ X' & \xrightarrow{g} & X \end{array}$$

*be a cartesian diagram of small v-stacks.*

(i) *Assume that  $f$  is qcqs, and  $A \in D_{\text{ét}}^+(Y, \Lambda)$ . Then*

$$g^* Rf_* A \cong Rf'_* \tilde{g}^* A .$$

(ii) Assume that  $f$  is compactifiable and representable in locally spatial diamonds with  $\dim. \operatorname{trg} f < \infty$ , and  $A \in D_{\text{ét}}(Y, \Lambda)$ . Then

$$g^* Rf_! A \cong Rf'_! \tilde{g}^* A .$$

(iii) Assume that  $g$  is compactifiable and representable in locally spatial diamonds with  $\dim. \operatorname{trg} f < \infty$ , and  $A \in D_{\text{ét}}(Y, \Lambda)$ . Then

$$Rg^! Rf_* A \cong Rf'_* R\tilde{g}^! A .$$

Part (i) is Proposition 17.6, part (ii) is Proposition 22.19, and part (iii) is Proposition 23.16.

We also need a theory of smooth morphisms. From now on, fix a prime  $\ell \neq p$ , and assume that  $\Lambda$  is killed by a power of  $\ell$ . In Definition 23.8, we define a notion of  $\ell$ -cohomologically smooth morphisms for any prime  $\ell \neq p$  (which depends on the prime  $\ell$ ), which includes étale maps.

**Theorem 1.10.** *Let  $f : Y \rightarrow X$  be a map of small  $v$ -stacks that is separated and representable in locally spatial diamonds with  $\dim. \operatorname{trg} f < \infty$ . The property of  $f$  being  $\ell$ -cohomologically smooth is  $v$ -local on the target and  $\ell$ -cohomologically smooth local on the source; moreover, composites and base changes of  $\ell$ -cohomologically smooth maps are  $\ell$ -cohomologically smooth, and smooth maps of analytic adic spaces over  $\mathbf{Z}_p$  give rise to  $\ell$ -cohomologically smooth morphisms under  $X \mapsto X^\diamond$ . Assume that  $f$  is  $\ell$ -cohomologically smooth, and  $\Lambda$  is  $\ell$ -power torsion.*

(i) There is a natural equivalence of functors

$$Rf^! A \cong f^* A \otimes_{\Lambda}^{\mathbb{L}} Rf^! \Lambda : D_{\text{ét}}(X, \Lambda) \rightarrow D_{\text{ét}}(Y, \Lambda) ,$$

where  $Rf^! \Lambda$  is  $v$ -locally isomorphic to  $\Lambda[n]$  for some  $n \in \mathbb{Z}$ .

(ii) For  $A, B \in D_{\text{ét}}(X, \Lambda)$ , there is a natural equivalence

$$f^* R\mathcal{H}om_{\Lambda}(A, B) \cong R\mathcal{H}om_{\Lambda}(f^* A, f^* B) .$$

(iii) Let

$$\begin{array}{ccc} Y' & \xrightarrow{\tilde{g}} & Y \\ \downarrow f' & & \downarrow f \\ X' & \xrightarrow{g} & X \end{array}$$

be a cartesian diagram of small  $v$ -stacks, with  $f$  being  $\ell$ -cohomologically smooth as before. Then for all  $A \in D_{\text{ét}}(X', \Lambda)$ ,

$$f^* Rg_* A \cong R\tilde{g}_* f'^* A .$$

Moreover, for all  $A \in D_{\text{ét}}(X, \Lambda)$ ,

$$\tilde{g}^* Rf^! A \cong Rf'^! g^* A .$$

We refer to Sections 23 and 24 for these results.

Moreover, for  $\ell$ -cohomologically smooth morphisms, one can prove some finiteness results. For this, we restrict to the case  $\Lambda = \mathbb{F}_\ell$ . For any small  $v$ -stack  $X$ , one can define a full subcategory  $D_{\text{cons}}(X, \mathbb{F}_\ell) \subset D_{\text{ét}}(X, \mathbb{F}_\ell)$ ; containment can be checked  $v$ -locally, and on affinoid perfectoid spaces, those are bounded complexes that become locally constant and finite-dimensional in each degree over a *constructible* stratification. Note that constructible here refers to the standard notion for spectral spaces, and excludes the stratification of a closed unit disc into a point and its complement (as the open complement is not quasicompact). We note that this condition can be checked on cohomology sheaves. The following theorem is also proved in Section 23.

**Theorem 1.11.** *Let  $f : Y \rightarrow X$  be a quasicompact  $\ell$ -cohomologically smooth map (in particular, compactifiable, representable in locally spatial diamonds, and with  $\dim. \text{trg } f < \infty$ ) of small  $v$ -stacks. For all  $A \in D_{\text{cons}}(Y, \mathbb{F}_\ell)$ , one has  $Rf_! A \in D_{\text{cons}}(X, \mathbb{F}_\ell)$ .*

Finally, it is useful to have some “invariance under change of algebraically closed base field” statements. These come in different flavours, depending on whether the base field is discrete or nonarchimedean. The following is Theorem 19.5.

**Theorem 1.12.** *Let  $X$  be a small  $v$ -stack.*

(i) *Assume that  $X$  lives over  $k$ , where  $k$  is a discrete algebraically closed field of characteristic  $p$ , and  $k'/k$  is an extension of discrete algebraically closed base fields,  $X' = X \times_k k'$ . Then the pullback functor*

$$D_{\text{ét}}(X, \Lambda) \rightarrow D_{\text{ét}}(X', \Lambda)$$

*is fully faithful.*

(ii) *Assume that  $X$  lives over  $k$ , where  $k$  is an algebraically closed discrete field of characteristic  $p$ . Let  $C/k$  be an algebraically closed complete nonarchimedean field, and  $X' = X \times_k \text{Spa}(C, C^+)$  for some open and bounded valuation subring  $C^+ \subset C$  containing  $k$ . Then the pullback functor*

$$D_{\text{ét}}(X, \Lambda) \rightarrow D_{\text{ét}}(X', \Lambda)$$

*is fully faithful.*

(iii) *Assume that  $X$  lives over  $\text{Spa}(C, C^+)$ , where  $C$  is an algebraically closed complete nonarchimedean field with an open and bounded valuation subring  $C^+ \subset C$ ,  $C'/C$  is an extension of algebraically closed complete nonarchimedean fields, and  $C'^+ \subset C'$  an open and bounded valuation subring containing  $C^+$ , such that  $\text{Spa}(C', C'^+) \rightarrow \text{Spa}(C, C^+)$  is surjective. Then for  $X' = X \times_{\text{Spa}(C, C^+)} \text{Spa}(C', C'^+)$ , the pullback functor*

$$D_{\text{ét}}(X, \Lambda) \rightarrow D_{\text{ét}}(X', \Lambda)$$

*is fully faithful.*

Let us say some words about the proofs. We start with the proof of Theorem 1.2. The proof proceeds in steps, proving first the analytic and étale cases (already in [Sch12], [KL15]), deducing the pro-étale case by a limit argument, and then the  $v$ -case by making use of the notion of (strictly) totally disconnected spaces.

**Definition 1.13.** *Let  $X$  be a perfectoid space. Then  $X$  is totally disconnected (resp. strictly totally disconnected) if  $X$  is quasicompact, and every open cover of  $X$  splits (resp. and every étale cover of  $X$  splits).*

It turns out that totally disconnected spaces have a very simple form.

**Proposition 1.14.** *Let  $X$  be a perfectoid space. Then  $X$  is totally disconnected (resp. strictly totally disconnected) if and only if  $X$  is affinoid, and every connected component of  $X$  is of the form  $\text{Spa}(K, K^+)$ , where  $K$  is a perfectoid field with an open and bounded valuation subring  $K^+ \subset K$  (resp. and  $K$  is algebraically closed).*

Thus, (strictly) totally disconnected are essentially profinite sets of (geometric) points. Their use is justified by the following two lemmas.

**Lemma 1.15.** *Let  $X$  be any quasicompact perfectoid space. Then there is a pro-étale cover  $\tilde{X} \rightarrow X$  such that  $\tilde{X}$  is strictly totally disconnected.*

**Lemma 1.16.** *Let  $X = \mathrm{Spa}(R, R^+)$  be a totally disconnected perfectoid space, and  $f : Y = \mathrm{Spa}(S, S^+) \rightarrow X$  be any map of perfectoid spaces. Then  $R^+/\varpi \rightarrow S^+/\varpi$  is flat, for any pseudouniformizer  $\varpi \in R$ .*

Using the first lemma and pro-étale descent, general  $v$ -descent reduces to the case of strictly totally disconnected spaces. Then the second lemma reduces this to faithfully flat descent.

Another important use of strictly totally disconnected spaces is a classification result for pro-étale maps. This uses the notion of separated morphisms of perfectoid spaces, for which we refer to Definition 5.10.

**Lemma 1.17.** *Let  $X$  be a strictly totally disconnected perfectoid space. Then a quasicompact separated map  $f : Y \rightarrow X$  of perfectoid spaces is pro-étale if and only if for every rank-1-point  $x = \mathrm{Spa}(C(x), \mathcal{O}_{C(x)}) \in X$ , the fibre of  $f$  over  $x$  is isomorphic to  $\underline{S}_x \times \mathrm{Spa}(C(x), \mathcal{O}_{C(x)})$  for some profinite set  $S_x$ . In this case,  $f$  is affinoid pro-étale.*

Many arguments follow this pattern of reduction to (strictly) totally disconnected spaces, which are essentially just profinite sets. This is akin to covering compact Hausdorff spaces by profinite sets. Making these arguments work is often a question of delicate point-set topology, and we will need to use the properties of spectral spaces very intensely. One application of this is  $v$ -descent for separated étale maps:

**Theorem 1.18.** *The following prestacks are  $v$ -stacks.*

- (i) *The prestack of affinoid perfectoid spaces over the category of totally disconnected perfectoid spaces.*
- (ii) *The prestack of separated pro-étale perfectoid spaces over the category of strictly totally disconnected perfectoid spaces.*
- (iii) *The prestack of separated étale maps over the category of all perfectoid spaces.*

Using these descent theorems, the following definition becomes reasonable.

**Definition 1.19.** *A locally separated map  $f : Y \rightarrow X$  of  $v$ -stacks is étale (resp. quasi-pro-étale) if for any perfectoid space (resp. any strictly totally disconnected perfectoid space)  $Z$  mapping to  $X$ , the fibre product  $Y \times_X Z$  is representable by a perfectoid space, and étale (resp. pro-étale) over  $Z$ .*

This leads to the following characterization of diamonds.

**Proposition 1.20.** *A  $v$ -sheaf  $Y$  is a diamond if and only if there is a surjective quasi-pro-étale map  $X \rightarrow Y$  from a perfectoid space  $X$ .*

A positive side of the  $v$ -topology is that any small  $v$ -stack can be accessed through the following steps: from perfectoid spaces to diamonds by quotients under a pro-étale equivalence relation; from diamonds to general small  $v$ -sheaves by quotients under a diamond equivalence relation; and from small  $v$ -sheaves to small  $v$ -stacks by quotients under a small  $v$ -sheaf equivalence relation. In the key step, one uses that any sub- $v$ -sheaf of a diamond is automatically itself a diamond; this statement does not seem to have any classical analogue.

However, at some point, we have to prove some theorems, in particular invariance under change of algebraically closed base field, smooth and proper base change, and Poincaré duality. Invariance



under change of algebraically closed base field is reduced by approximation to the noetherian cases handled by Huber (which in turn are reduced to the case of schemes via nearby cycles). Proper base change is reduced through a series of reductions to a statement about Zariski–Riemann spaces of (algebraically closed) fields, which follows from proper base change for schemes. For smooth base change and Poincaré duality, things are reduced by a series of reductions to the case of a closed unit disc, where we can use Huber’s results. We note that our reductions reduce all appeals to Huber’s book to the case of smooth curves over  $\mathrm{Spa}(C, C^+)$ , where  $C$  is an algebraically closed complete nonarchimedean field, and  $C^+ \subset C$  is an open and bounded valuation subring; however, in that case, we need a solid theory of Poincaré duality.

We make a general warning to the reader that this manuscript contains essentially no examples, which probably makes it almost unreadable. We hope that forthcoming papers, including [FS], will give the reader the necessary examples and motivation to work through the long technical arguments of this paper.

**Acknowledgments.** These ideas were formed during a long period after my Berkeley course in 2014. I want to thank Laurent Fargues, David Hansen, Kiran Kedlaya and Jared Weinstein for very helpful discussions about various parts. In the winter term 2016/17, the ARGOS seminar in Bonn was going through the manuscript, and I want to thank all participants heartily for their detailed comments. Moreover, I want to thank the organizers of the Hadamard Lectures at the IHÉS for their invitation in March/April 2017, and in particular Ofer Gabber for his many helpful questions and suggestions. Part of this work was done while the author was a Clay Research Fellow.

## 2. SPECTRAL SPACES

We recall some basic properties of spectral spaces that are used throughout.

**Definition 2.1.** *A spectral space is a topological space  $X$  such that  $X$  is quasicompact,  $X$  has a basis of quasicompact open subsets stable under finite intersection, and every irreducible closed subset has a unique generic point. A locally spectral space is a topological space  $X$  that admits an open cover by spectral subspaces.*

*A morphism  $f : X \rightarrow Y$  between spectral spaces is spectral if it is continuous and the preimage of any quasicompact open subset is quasicompact open. A morphism  $f : X \rightarrow Y$  between locally spectral spaces is spectral if it is continuous and for any spectral open subspace  $U \subset X$  mapping into some spectral open  $V \subset Y$ , the map  $U \rightarrow V$  is spectral.*

**Theorem 2.2** (Hochster). *Let  $X$  be a topological space. The following conditions are equivalent.*

- (i) *The space  $X$  is spectral.*
- (ii) *There is a ring  $A$  such that  $X \cong \mathrm{Spec} A$ .*
- (iii) *The space  $X$  can be written as an inverse limit of finite  $T_0$ -spaces.*

*The category of spectral spaces with spectral maps is equivalent to the pro-category of finite  $T_0$ -spaces.*

Recall that a subset  $T \subset X$  of a spectral space  $X$  is constructible if it lies in the boolean algebra generated by quasicompact open subsets. The constructible topology on  $X$  is the topology generated by constructible subsets. Then  $X$  equipped with its constructible topology is a profinite set. If  $X$  is an inverse limit of finite  $T_0$ -spaces  $X_i$ , then  $X$  with the constructible topology is the inverse limit of the  $X_i$  equipped with the discrete topology.

A subset  $T \subset X$  is pro-constructible if it is an intersection of constructible subsets. More generally, a subset  $T \subset X$  of a locally spectral space  $X$  is constructible (resp. pro-constructible) if the intersection with any spectral open subspace is constructible (resp. pro-constructible).

**Lemma 2.3.** *Let  $X$  be a spectral space. A subset  $S \subset X$  is pro-constructible if and only if  $S$  is closed in the constructible topology. If  $f : X \rightarrow Y$  is a spectral map of spectral spaces, then the image of any pro-constructible subset of  $X$  is a pro-constructible subset of  $Y$ .*

*Proof.* Constructible subsets are open and closed in the constructible topology, so pro-constructible subsets are closed. Conversely, if a subset is closed in the constructible topology, then it is an intersection of open and closed subsets in the constructible topology. But open and closed subsets of the constructible topology are constructible, for example by writing  $X$  as a cofiltered inverse limit of finite  $T_0$ -space  $X_i$  (where any open and closed subset of  $X$  for the constructible topology is a preimage of a subset of some  $X_i$ ).

Now let  $f : X \rightarrow Y$  be spectral map of spectral spaces, and  $S \subset X$  a pro-constructible subset with image  $T \subset Y$ . If we equip  $X$  and  $Y$  with their constructible topology, then  $f$  is a continuous map of compact Hausdorff spaces, and  $S \subset X$  is a closed subset, which thus has closed image  $T \subset Y$ . Thus,  $T$  is pro-constructible.  $\square$

**Lemma 2.4.** *Let  $X$  be a spectral space, and  $S \subset X$  a pro-constructible subset. Then the closure  $\bar{S} \subset X$  of  $S$  is the set of specializations of points in  $S$ .*

*Proof.* Clearly, any specialization of a point in  $S$  lies in  $\bar{S}$ . Now let  $x \in X$  be a point such that no generalization of  $x$  lies in  $S$ . Let  $X_x \subset X$  be the set of generalizations of  $x$ , which is the intersection of all quasicompact open subsets  $U$  containing  $x$ ; in particular,  $X_x \subset X$  is a pro-constructible subset. Then

$$\emptyset = X_x \cap S = \bigcap_{U \ni x} (U \cap S) .$$

In other words, the  $(X \setminus U) \cap S$  cover  $S$ . As  $S$  is quasicompact in the constructible topology, and  $X \setminus U$  is constructible, it follows that there is some quasicompact open neighborhood  $U$  of  $x$  such that  $(X \setminus U) \cap S = S$ , i.e.  $U \cap S = \emptyset$ . Thus,  $x$  has an open neighborhood which does not meet  $S$ , so does not lie in the closure of  $S$ , as desired.  $\square$

All maps of analytic adic spaces are generalizing. This notably implies that surjective maps are quotient maps:

**Lemma 2.5.** *Let  $f : Y \rightarrow X$  be a surjective and generalizing spectral map of spectral spaces. Then  $f$  is a quotient map.*

*Proof.* We have to show a subset  $S \subset X$  is open if the preimage  $T = f^{-1}(S) \subset Y$  is open, so assume that  $T$  is open. Note that in particular  $Y \setminus T \subset Y$  is closed, and thus pro-constructible; therefore, its image  $X \setminus S \subset X$  is pro-constructible. But as  $f$  is generalizing, this subset is also closed under specializations, and thus closed. Thus,  $S$  is open, as desired.  $\square$

We also recall the following classical lemma.

**Lemma 2.6.** *Let  $f : S \rightarrow T$  be a continuous surjective map from a quasicompact space  $S$  to a compact Hausdorff space  $T$ . Then  $f$  is a quotient map.*

*Proof.* Note that for any  $x \in T$ , the open neighborhoods are cofinal with the closed neighborhoods. Assume that  $V \subset T$  is a subset whose preimage  $U = f^{-1}(V) \subset S$  is open with closed complement  $Z = S \setminus U$ , and let  $x \in V$ . Then

$$\emptyset = f^{-1}(x) \cap Z = \bigcap_{U_x \subset T} f^{-1}(\overline{U_x}) \cap Z.$$

Here,  $U_x \subset T$  runs over open neighborhoods of  $x$  in  $T$ . The intersection is over closed subsets of  $S$ ; by quasicompactness, there is some open neighborhood  $U_x$  of  $x$  such that  $f^{-1}(\overline{U_x}) \cap Z = \emptyset$ , i.e.  $f^{-1}(\overline{U_x}) \subset U$ . This implies that  $f^{-1}(U_x) \subset U$ , and thus  $U_x \subset V$ , as desired.  $\square$

Here is a lemma about equivalence relations on spectral spaces. Recall that a topological space is quasiseparated if the intersection of any two quasicompact open subsets is quasicompact.

**Lemma 2.7.** *Let  $X$  be a quasiseparated locally spectral space, and let  $R \subset X \times X$  be a pro-constructible equivalence relation such that the maps  $s, t : R \rightarrow X$  are quasicompact and generalizing. Then the quotient space  $X/R$  is  $T_0$ . Moreover, for any quasicompact open  $W \subset X$ , there exists an open  $R$ -invariant subset  $U \supset W$  such that  $U \subset E$ , where  $E$  is some  $R$ -invariant subset that is an intersection of a nonempty family of quasicompact open subsets.*

*Proof.* We start with the second part, for which we follow the arguments from [Sta, Tag 0APA, 0APB]. Let  $s, t : R \rightarrow X$  denote the two projections. Let  $W \subset X$  be a quasicompact open subset. As  $s$  is quasicompact,  $s^{-1}(W) \subset R$  is quasicompact open, and thus its image  $E = t(s^{-1}(W)) \subset X$  is quasicompact, pro-constructible, and generalizing. Note that by general nonsense about equivalence relations,  $E = t(s^{-1}(W))$  is  $R$ -invariant. By quasicompactness, there is some quasicompact open subset  $W' \subset X$  containing  $E$ . Now  $E$  is a pro-constructible and generalizing subset of the spectral space  $W'$ ; this implies that  $E$  is an intersection of quasicompact open subsets. Let  $Z = X \setminus W'$ , which is a closed subset of  $X$ . Then  $s^{-1}(Z) \subset R$  is closed, and in particular pro-constructible. As  $t$  is quasicompact, it follows that  $T = t(s^{-1}(Z)) \subset X$  is pro-constructible. Thus, its closure  $\bar{T} \subset X$  is the set of specializations of elements of  $T$ . We claim that  $\bar{T}$  is  $R$ -invariant: Indeed, if  $\xi \in \bar{T}$ , there is some point  $\xi' \in T$  specializing to  $\xi$ . If  $r \in R$  is a point mapping under  $s$  to  $\xi$ , we can find  $r' \in R$  mapping under  $s$  to  $\xi'$ , as  $s$  is generalizing. Then  $t(r)$  is a specialization of  $t(r') \in T$ , so  $t(r) \in \bar{T}$ . Finally,  $U = X \setminus \bar{T}$  is an open  $R$ -invariant subset of  $X$ , such that  $W \subset U \subset W'$ . The inclusion  $U \subset E$  does not hold with our definition of  $E$ , but we have  $U \subset E'$  where  $E' = t(s^{-1}(W'))$ .

To see that  $X/R$  is  $T_0$ , start with two distinct points  $\bar{x}, \bar{y} \in X/R$ , and lift them to  $x, y \in X$ , so that their  $R$ -orbits are  $R \cdot x = t(s^{-1}(x))$  and  $R \cdot y = t(s^{-1}(y))$ , respectively, and  $R \cdot x \cap R \cdot y = \emptyset$ . Note that  $R \cdot x$  and  $R \cdot y$  are both pro-constructible quasicompact subsets of  $X$ , as  $s$  is quasicompact and  $t$  is spectral; in particular,  $R \cdot x$ ,  $R \cdot y$  and  $R \cdot x \cup R \cdot y$  are spectral spaces. We claim that it cannot happen that there are  $x_1, x_2 \in R \cdot x$  and  $y_1, y_2 \in R \cdot y$  such that  $x_1$  generalizes to  $y_1$  and  $x_2$  specializes to  $y_2$ . Indeed, assume this was the case. As  $R \cdot x \cup R \cdot y \subset X$  is a pro-constructible subset, it is a spectral space, and thus admits a maximal point; possibly changing the roles of  $x$  and  $y$ , we may assume that  $y$  is a maximal point of  $R \cdot x \cup R \cdot y$ . As  $t : R \rightarrow X$  is generalizing,  $(y, y_2) \in R$ , and  $y_2$  generalizes to  $x_2$ , we can find some point  $(z, x_2) \in R$ , where  $z$  is a generalization of  $y$ . Then  $z \in R \cdot x \subset R \cdot x \cup R \cdot y$ . As  $y$  is maximal in that space, it follows that  $z = y$ , but then  $(y, x_2) \in R$ , so that  $R \cdot y = R \cdot x_2 = R \cdot x$ , which is a contradiction.

Thus, we may assume that no point of  $R \cdot x$  generalizes to a point of  $R \cdot y$ . In that case, we can find a quasicompact open subset  $W \subset X$  containing  $x$  such that  $W \cap R \cdot y = \emptyset$ : Indeed, the intersection of  $W \cap R \cdot y$  over all quasicompact open neighborhoods  $W$  of  $x$  is empty, as  $R \cdot y$

contains no generalization of  $x$ . But then by quasicompactness  $R \cdot y$  for the constructible topology,  $W \cap R \cdot y = \emptyset$  for some such  $W$ . Now we run the above argument to find some open  $R$ -invariant subset  $U \supset W$  of  $X$ . We claim that we can do this so that  $U \cap R \cdot y = \emptyset$ : This would finish the proof, as  $R \cdot x \subset U$ , while  $U \cap R \cdot y = \emptyset$ , so that the image of  $U$  in  $X/R$  separates  $x$  from  $y$ . To see that one can arrange  $U \cap R \cdot y = \emptyset$ , note that in the notation of the first paragraph,  $E \cap R \cdot y = \emptyset$ . As  $E$  is generalizing, one can for any point  $e \in E$  find a quasicompact open neighborhood  $W'_e$  of  $e$  such that  $W'_e \cap R \cdot y = \emptyset$ ; in other words, by quasicompactness of  $E$  we can choose  $W'$  so that  $W' \cap R \cdot y = \emptyset$ . In that case, also  $E' \cap R \cdot y = \emptyset$ , so as we have  $U \subset E'$  for the  $R$ -invariant open subset  $U \subset X$ , we see that  $U \cap R \cdot y = \emptyset$ , as desired.  $\square$

**Remark 2.8.** Even if  $X$  is spectral, it can happen under the hypothesis of the lemma that the quotient space  $X/R$  is not spectral. In fact, using just profinite sets  $X$  and  $R$ , one can get any compact Hausdorff space as the quotient  $X/R$ : Namely, given a compact Hausdorff space  $Y$ , let  $X$  be the Stone-Cech compactification of  $Y$  considered as a discrete set. Then  $X$  is a profinite set which comes with a natural continuous surjective map to  $Y$ , and the induced equivalence relation  $R \subset X \times X$  is closed, and thus profinite.

**Lemma 2.9.** *Let  $X$  be a spectral space, and let  $R \subset X \times X$  be a pro-constructible equivalence relation such that the maps  $s, t : R \rightarrow X$  are generalizing. Assume that  $X/R$  has a basis for the topology given by open subsets whose preimages in  $X$  are quasicompact. Then  $X/R$  is a spectral space, and  $X \rightarrow X/R$  is a spectral and generalizing map.*

*Proof.* The set of open subsets of  $X/R$  whose preimages in  $X$  are quasicompact is stable under finite intersections. To see that  $X/R$  is spectral, it remains to see that every irreducible closed subset  $Z \subset X/R$  has a unique generic point. Uniqueness holds because  $X/R$  is  $T_0$  by Lemma 2.7. For existence, look at the cofiltered intersection of all open subsets  $U \subset X/R$  with  $U \cap Z \neq \emptyset$  whose preimage in  $X$  is quasicompact. We need to see that this intersection is nonempty. But the preimages of such  $U$  are quasicompact open subsets, thus compact Hausdorff for the constructible topology. As they are all nonempty, their intersection is nonempty. Thus, the intersection of the  $U$ 's is nonempty as well, as desired.

As  $X/R$  has a basis of opens for which the preimage in  $X$  is quasicompact, it follows that  $X \rightarrow X/R$  is spectral. It remains to see that  $X \rightarrow X/R$  is generalizing. Assume that  $\bar{x}$  generalizes to  $\bar{x}' \in X/R$ . We claim first that there is some way to lift this to a generalization  $x$  to  $x'$  in  $X$ . If not, then the final paragraph of the proof of Lemma 2.7 would produce an open  $R$ -invariant neighborhood of  $x$  in  $X$  which does not contain  $x'$ , contradicting that  $\bar{x}$  generalizes to  $\bar{x}'$ . Now if  $x_0 \in X$  is any lift of  $\bar{x}$ , then  $(x_0, x) \in R$ . But  $t : R \rightarrow X$  is generalizing, and  $x'$  is a generalization of  $x$  in  $X$ , so we can find a generalization  $(x'_0, x')$  of  $(x_0, x)$ . But then  $x'_0 \in X$  is a generalization of  $x_0$  mapping to  $\bar{x}' \in X/R$ , as desired.  $\square$

**Lemma 2.10.** *In the situation of Lemma 2.7, assume that  $R \rightarrow X$  is open. Then the quotient space  $X/R$  is locally spectral and quasiseparated, and  $X \rightarrow X/R$  is an open spectral qcqs map.*

*Proof.* For any quasicompact open subset  $U \subset X$ , the set  $t(s^{-1}(U)) \subset X$  is a quasicompact  $R$ -invariant open subset under the assumptions. Note that  $t(s^{-1}(U)) \subset X$  is also the preimage of the image of  $U$  in  $X/R$ , so that  $X \rightarrow X/R$  is open. For the other assertions, replacing  $X$  by  $U$ , we can assume that  $X$  is quasicompact, thus spectral. We claim that the quasicompact open subsets  $V_0 \subset X/R$  whose preimages in  $X$  are quasicompact, form a basis for the topology. Thus, let  $W_0 \subset X/R$  be open, with preimage  $W \subset X$  an  $R$ -invariant open subset. Pick any  $x_0 \in W_0$ , and lift

it to  $x \in W$ . Then  $x$  has a quasicompact open neighborhood  $x \in V' \subset W$ , and  $V = t(s^{-1}(V')) \subset W$  is a quasicompact  $R$ -invariant open subset. Its image  $V_0 \subset W_0$  thus contains  $x_0$ , is quasicompact, with inverse image  $V \subset X$  still quasicompact.

It remains to see that every irreducible closed subset  $\bar{Z} \subset X/R$  has a generic point. Equivalently, the intersection of all quasicompact open subsets  $V_0 \subset X/R$  with  $V_0 \cap \bar{Z} \neq \emptyset$  is nonempty. But the corresponding intersection of the preimages in  $X$  is nonempty by compactness of the constructible topology, so the result follows.  $\square$

We will need the following result about inverse limits.

**Lemma 2.11.** *Let  $X_i, i \in I$ , be a cofiltered inverse system of spectral spaces along spectral maps, with inverse limit  $X = \varprojlim_i X_i$ . Then  $X$  is a spectral space, the maps  $X \rightarrow X_i$  are spectral, and a map  $Y \rightarrow X$  from a spectral space is spectral if and only if all composites  $Y \rightarrow X \rightarrow X_i$  are spectral. Moreover, if  $Y$  is a spectral space with a spectral map  $Y \rightarrow X$  such that the composite maps  $Y \rightarrow X_i$  are generalizing, then  $Y \rightarrow X$  is generalizing. In particular, if  $Y \rightarrow X$  is in addition surjective, then by Lemma 2.5 it is a quotient map.*

*Proof.* The first part follows for example from the interpretation of the category of spectral spaces with spectral maps as the pro-category of finite  $T_0$ -spaces. Now assume  $Y \rightarrow X$  is a spectral map of spectral spaces such that  $Y \rightarrow X_i$  is generalizing for all  $i \in I$ . Let  $y \in Y$  map to  $x \in X$ , and let  $\tilde{x} \in X$  be a generalization of  $x \in X$ . We look for a generalization  $\tilde{y}$  of  $y$  mapping to  $\tilde{x}$ . For this, we may replace  $Y$  by its localization  $Y_y$  at  $y$  (i.e. the set of generalizations of  $y$ ). Let  $x_i \in X_i$  be the image of  $x$ , and  $\tilde{x}_i \in X_i$  be the image of  $\tilde{x}$ . Then, for all  $i \in I$ ,  $\tilde{x}_i$  is a generalization of  $x_i$ . As  $Y \rightarrow X_i$  is generalizing, the preimage of  $\tilde{x}_i$  in  $Y$  is a non-empty pro-constructible subset of  $Y$ . For varying  $i$ , these preimages form a cofiltered inverse system of nonempty pro-constructible subsets of  $Y$ ; their intersection is thus nonempty (by Tychonoff). Any point in the intersection is a generalization of  $y$  mapping to  $\tilde{x}$ , as desired.  $\square$

### 3. PERFECTOID SPACES

Let  $p$  be a fixed prime throughout. Recall that a topological ring  $R$  is Tate if it contains an open and bounded subring  $R_0 \subset R$  and a topologically nilpotent unit  $\varpi \in R$ ; such elements are called pseudo-uniformizers. Let  $R^\circ \subset R$  denote the set of powerbounded elements. If  $\varpi \in R$  is a pseudo-uniformizer, then necessarily  $\varpi \in R^\circ$ . Furthermore, if  $R^+ \subset R$  is a ring of integral elements, i.e. an open and integrally closed subring of  $R^\circ$ , then  $\varpi \in R^+$ . Indeed, since  $\varpi^n \rightarrow 0$  as  $n \rightarrow \infty$  and since  $R^+$  is open, we have  $\varpi^n \in R^+$  for some  $n \geq 1$ . Since  $R^+$  is integrally closed,  $\varpi \in R^+$ .

The following definition is due to Fontaine, [Fon13].

**Definition 3.1.** *A Tate ring  $R$  is perfectoid if  $R$  is complete, uniform, i.e.  $R^\circ \subset R$  is bounded, and there exists a pseudo-uniformizer  $\varpi \in R$  such that  $\varpi^p | p$  in  $R^\circ$  and the Frobenius map*

$$\Phi: R^\circ/\varpi \rightarrow R^\circ/\varpi^p : x \mapsto x^p$$

*is an isomorphism.*

Hereafter we use the following notational convention. If  $R$  is a ring, and  $I, J \subset R$  are ideals containing  $p$  such that  $I^p \subset J$ , then  $\Phi: R/I \rightarrow R/J$  will refer to the ring homomorphism  $x \mapsto x^p$ .

**Remark 3.2.** The condition that  $\Phi : R^\circ/\varpi \rightarrow R^\circ/\varpi^p$  is an isomorphism is independent of the choice of a pseudo-uniformizer  $\varpi \in R$  such that  $\varpi^p|p$ . In fact, we claim that it is equivalent to the condition that  $\Phi : R^\circ/p \rightarrow R^\circ/p$  is surjective.

Indeed, for any complete Tate ring  $R$  and pseudo-uniformizer  $\varpi$  satisfying  $\varpi^p|p$  in  $R^\circ$ , the Frobenius map  $\Phi : R^\circ/\varpi \rightarrow R^\circ/\varpi^p$  is necessarily injective. Indeed, if  $x \in R^\circ$  satisfies  $x^p = \varpi^p y$  for some  $y \in R^\circ$  then the element  $x/\varpi \in R$  lies in  $R^\circ$  since its  $p$ -th power does. Thus, the isomorphism condition on  $\Phi$  in Definition 3.1 is equivalent to surjectivity of  $\Phi : R^\circ/\varpi \rightarrow R^\circ/\varpi^p$ . Thus, if  $\Phi : R^\circ/p \rightarrow R^\circ/p$  is surjective, then  $R$  is perfectoid. Conversely, assume that  $R$  is perfectoid. Let  $x \in R^\circ$  be any element. By using that  $\Phi : R^\circ/\varpi \rightarrow R^\circ/\varpi^p$  is surjective and successive  $\varpi^p$ -adic approximation, we can write

$$x = x_0^p + \varpi^p x_1^p + \varpi^{2p} x_2^p + \dots ,$$

where  $x_i \in R^\circ$ . But then

$$x \equiv (x_0 + \varpi x_1 + \varpi^2 x_2 + \dots)^p \pmod{p} ,$$

showing that  $\Phi : R^\circ/p \rightarrow R^\circ/p$  is surjective, as desired.

**Remark 3.3.** The field  $\mathbf{Q}_p$  is not perfectoid, even though the Frobenius map on  $\mathbf{F}_p$  is an isomorphism. The issue is that there is no element  $\varpi \in \mathbf{Z}_p$  whose  $p$ -th power divides  $p$ . More generally, a discretely valued non-archimedean field  $K$  cannot be perfectoid. Indeed, if  $\varpi$  is a pseudo-uniformizer as in Definition 3.1, then  $\varpi$  is a non-zero element of the maximal ideal, so the quotients  $K^\circ/\varpi$  and  $K^\circ/\varpi^p$  are Artin local rings of different lengths and hence cannot be isomorphic.

**Example 3.4.** (i) The cyclotomic field  $\mathbf{Q}_p^{\text{cycl}}$ , the completion of the cyclotomic extension  $\mathbf{Q}_p(\mu_{p^\infty})$ .

(ii) The  $t$ -adic completion of  $\mathbf{F}_p((t))(t^{1/p^\infty})$ , which we will write as  $\mathbf{F}_p((t^{1/p^\infty}))$ .

(iii) The algebra  $\mathbf{Q}_p^{\text{cycl}}\langle T^{1/p^\infty} \rangle$ . This is defined as  $A[1/p]$ , where  $A$  is the  $p$ -adic completion of  $\mathbf{Z}_p^{\text{cycl}}[T^{1/p^\infty}]$ .

(iv) As an example of a perfectoid Tate ring which does not live over a field, one may take

$$R = \mathbf{Z}_p^{\text{cycl}}[[T^{1/p^\infty}]]\langle (p/T)^{1/p^\infty} \rangle[1/T] .$$

Here we can take  $\varpi = T^{1/p}$ , because  $\varpi^p = T$  divides  $p$  in  $R^\circ$ .

**Proposition 3.5.** *Let  $R$  be a topological ring with  $pR = 0$ . The following are equivalent:*

- (1) *The topological ring  $R$  is a perfectoid Tate ring.*
- (2) *The topological ring  $R$  is a perfect complete Tate ring.*

Of course, perfect means that  $\Phi : R \rightarrow R$  is an isomorphism.

*Proof.* Let  $R$  be a complete Tate ring. If  $R$  is perfect, then take  $\varpi$  any pseudo-uniformizer. The condition  $\varpi^p|p = 0$  is vacuous. If  $x \in R$  is powerbounded, then so is  $x^p$ , and vice versa, which means that  $\Phi : R^\circ \rightarrow R^\circ$  is an isomorphism. This shows that  $\Phi : R^\circ/\varpi \rightarrow R^\circ/\varpi^p$  is surjective, and we had seen in Remark 3.2 that injectivity is automatic. It remains to see that  $R$  is automatically uniform. For this, take any open and bounded subring  $R_0 \subset R$ . As  $\Phi : R \rightarrow R$  is an isomorphism, which is automatically open by Banach's open mapping theorem, there is some  $n$  such that  $\varpi^n R_0 \subset \Phi(R_0)$ . Equivalently,  $\Phi^{-1}(R_0) \subset \varpi^{-n/p} R_0$ . This implies that

$$\Phi^{-2}(R_0) \subset \Phi^{-1}(\varpi^{-n/p} R_0) \subset \varpi^{-n/p - n/p^2} R_0$$

and inductively

$$\Phi^{-k}(R_0) \subset \varpi^{-n/p-n/p^2-\dots-n/p^k} R_0 .$$

In particular, for all  $k \geq 0$ ,

$$\Phi^{-k}(R_0) \subset \varpi^{-n} R_0 .$$

On the other hand,  $R^\circ \subset \bigcup_{k \geq 0} \Phi^{-k}(R_0)$ , and so  $R^\circ \subset \varpi^{-n} R_0$  is bounded, as desired.

Conversely, if  $R$  is perfectoid, then  $\Phi: R^\circ/\varpi \rightarrow R^\circ/\varpi^p$  is an isomorphism, and therefore so is  $R^\circ/\varpi^n \rightarrow R^\circ/\varpi^{np}$  by induction. Taking inverse limits and using completeness, we find that  $\Phi: R^\circ \rightarrow R^\circ$  is an isomorphism. Inverting  $\varpi$  shows that  $R$  is perfect.  $\square$

**Definition 3.6.** *A perfectoid field is a perfectoid Tate ring  $R$  which is a nonarchimedean field.*

**Remark 3.7.** It is not clear a priori that a perfectoid ring which is a field is a perfectoid field. However, this has recently been answered affirmatively by Kedlaya, [Ked16].

One easily checks the following equivalent characterization, which is the original definition, [Sch12], [KL15].

**Proposition 3.8.** *Let  $K$  be a nonarchimedean field. Then  $K$  is a perfectoid field if and only if the following conditions hold:*

- (i) *The nonarchimedean field  $K$  is not discretely valued,*
- (ii) *the absolute value  $|p| < 1$ , and*
- (iii) *the Frobenius  $\Phi: \mathcal{O}_K/p \rightarrow \mathcal{O}_K/p$  is surjective.*

$\square$

Next, we recall the process of tilting.

**Definition 3.9.** *Let  $R$  be a perfectoid Tate ring. The tilt of  $R$  is the topological ring*

$$R^{\flat} = \varprojlim_{x \mapsto x^p} R ,$$

*with the inverse limit topology, the pointwise multiplication, and the addition given by*

$$(x^{(0)}, x^{(1)}, \dots) + (y^{(0)}, y^{(1)}, \dots) = (z^{(0)}, z^{(1)}, \dots)$$

*where*

$$z^{(i)} = \lim_{n \rightarrow \infty} (x^{(i+n)} + y^{(i+n)})^{p^n} \in R .$$

**Lemma 3.10.** *The limit  $z^{(i)}$  above exists and defines a ring structure making  $R^{\flat}$  a perfectoid  $\mathbf{F}_p$ -algebra. The subset  $R^{\flat\circ}$  of power-bounded elements is given by the topological ring isomorphism*

$$R^{\flat\circ} = \varprojlim_{x \mapsto x^p} R^\circ \cong \varprojlim_{\Phi} R^\circ / \varpi ,$$

*where  $\varpi \in R$  is a pseudo-uniformizer which divides  $p$  in  $R^\circ$ .*

*Furthermore, there exists a pseudo-uniformizer  $\varpi \in R$  with  $\varpi^p | p$  in  $R^\circ$  which admits a sequence of  $p$ -power roots  $\varpi^{1/p^n}$ , giving rise to an element  $\varpi^{\flat} = (\varpi, \varpi^{1/p}, \dots) \in R^{\flat\circ}$ , which is a pseudo-uniformizer of  $R^{\flat}$ . Then  $R^{\flat} = R^{\flat\circ}[1/\varpi^{\flat}]$ .*

*Proof.* Let  $\varpi_0$  be a pseudo-uniformizer of  $R$  such that  $\varpi_0|p$  in  $R^\circ$ . Let us check that the map

$$\varprojlim_{x \mapsto x^p} R^\circ \rightarrow \varprojlim_{\Phi} R^\circ / \varpi_0$$

is an isomorphism. We have to see that any sequence  $(\bar{x}_0, \bar{x}_1, \dots) \in \varprojlim_{\Phi} R^\circ / \varpi_0$  lifts uniquely to a sequence  $(x_0, x_1, \dots) \in \varprojlim_{\Phi} R^\circ$ . This lift is given by  $x^{(i)} = \lim_{n \rightarrow \infty} x_{n+i}^{p^n}$ , where  $x_j \in R^\circ$  is any lift of  $\bar{x}_j$ . (For the convergence of that limit, note that if  $x \equiv y \pmod{\varpi_0^n}$ , then  $x^p \equiv y^p \pmod{\varpi_0^{n+1}}$ .) This shows that we get a well-defined ring

$$\varprojlim_{x \mapsto x^p} R^\circ \subset \varprojlim_{x \mapsto x^p} R = R^b,$$

which agrees with the powerbounded elements  $R^{b^\circ} \subset R^b$ .

Next, we construct the element  $\varpi^b$ . For this, we assume that  $\varpi_0 = \varpi_1^p$  for some pseudouniformizer  $\varpi_1 \in R$  such that  $\varpi_0 = \varpi_1^p|p$ . Any preimage of  $\varpi_0$  under  $R^{b^\circ} = \varprojlim_{\Phi} R^\circ / \varpi_0 \rightarrow R^\circ / \varpi_0$  is an element  $\varpi^b$  with the right properties. It is congruent to  $\varpi_1$  modulo  $\varpi_0$ , and therefore it is also a topologically nilpotent, and invertible in  $R^b$ . Then  $\varpi = \varpi^{b^\sharp}$  is the desired pseudo-uniformizer of  $R^\circ$ .

Now one sees that  $R^b = R^{b^\circ}[1/\varpi^b]$  as multiplicative monoids, and in fact as rings. This implies that the addition in  $R^b$  is always well-defined, and defines a ring of characteristic  $p$ , which is perfect by design. The rest of the lemma follows easily.  $\square$

We have a continuous, multiplicative (but not additive) map  $R^b = \varprojlim_{x \mapsto x^p} R \rightarrow R$  by projecting onto the zeroth coordinate; call this  $f \mapsto f^\sharp$ . This projection defines a ring isomorphism  $R^{b^\circ} / \varpi^b \cong R^\circ / \varpi$ . The open and integrally closed subrings of  $R^{b^\circ}$  and  $R^\circ$  correspond exactly to the integrally closed subrings of their common quotients modulo  $\varpi^b$  and  $\varpi$ . This defines an inclusion-preserving bijection between the sets of open and integrally closed subrings of  $R^{b^\circ}$  and  $R^\circ$ . This correspondence can be made more explicit:

**Lemma 3.11.** *The set of open and integrally closed subrings  $R^+ \subset R^\circ$  is in bijection with the set of open and integrally closed subrings  $R^{b^+} \subset R^{b^\circ}$ , via  $R^{b^+} = \varprojlim_{x \mapsto x^p} R^+$ . Also,  $R^{b^+} / \varpi^b = R^+ / \varpi$ .  $\square$*

The following two theorems belong to a pattern of “tilting equivalence”.

**Theorem 3.12** ([Sch12], [KL15]). *Let  $R$  be a perfectoid Tate ring with an open and integrally closed subring  $R^+ \subset R$ , with tilt  $(R^b, R^{b^+})$ .*

*The map sending  $x \in \text{Spa}(R, R^+)$  to  $x^b \in \text{Spa}(R^b, R^{b^+})$  defined by  $|f(x^b)| = |f^\sharp(x)|$  defines a homeomorphism  $\text{Spa}(R, R^+) \cong \text{Spa}(R^b, R^{b^+})$ . A subset  $U \subset \text{Spa}(R, R^+)$  is rational if and only if its image in  $\text{Spa}(R^b, R^{b^+})$  is rational.*

**Theorem 3.13** ([Sch12]). *Let  $R$  be a perfectoid Tate ring with tilt  $R^b$ . Then there is an equivalence of categories between perfectoid  $R$ -algebras and perfectoid  $R^b$ -algebras, via  $S \mapsto S^b$ .*

*Proof.* In [Sch12], these are only proved over a perfectoid field, but the proof works in general.  $\square$

Let us describe the inverse functor, along the lines of Fontaine’s Bourbaki talk, [Fon13]. In fact we will answer a more general question. Given a perfectoid Tate ring  $R$  in characteristic  $p$ , what are all the untilts  $R^\sharp$  of  $R$ ?



**Lemma 3.14.** *Let  $R$  be a perfectoid Tate ring with a ring of integral elements  $R^+ \subset R$ , and let  $(R^b, R^{b+})$  be its tilt.*

(i) *There is a canonical surjective ring homomorphism*

$$\begin{aligned} \theta : W(R^{b+})R^+ \\ \sum_{n \geq 0} [r_n]p^n \sum_{n \geq 0} r_n^\# p^n \end{aligned}$$

(ii) *The kernel of  $\theta$  is generated by a nonzero-divisor  $\xi$  of the form  $\xi = p + [\varpi]\alpha$ , where  $\varpi \in R^{b+}$  is a pseudo-uniformizer, and  $\alpha \in W(R^{b+})$ .*

Related results are discussed in [Fon13], [FF11] and [KL15, Theorem 3.6.5].

**Definition 3.15.** *An ideal  $I \subset W(R^{b+})$  is primitive of degree 1 if  $I$  is generated by an element of the form  $\xi = p + [\varpi^b]\alpha$ , with  $\varpi^b \in R^{b+}$  a pseudo-uniformizer and  $\alpha \in W(R^{b+})$ .*

An element  $\xi$  of this form is necessarily a non-zero-divisor:

**Lemma 3.16.** *Any  $\xi$  of the form  $\xi = p + [\varpi^b]\alpha$ , with  $\varpi^b \in R^{b+}$  a pseudo-uniformizer and  $\alpha \in W(R^+)$ , is a non-zero-divisor.*

*Proof.* Assume that  $\xi \sum_{n \geq 0} [c_n]p^n = 0$ . Modulo  $[\varpi^b]$ , this reads  $\sum_{n \geq 0} [c_n]p^{n+1} \equiv 0 \pmod{[\varpi^b]}$ , meaning that all  $c_n \equiv 0 \pmod{\varpi^b}$ . We can then divide all  $c_n$  by  $\varpi^b$ , and induct.  $\square$

*Proof of Lemma 3.14.* For part (i), we first check that  $\theta$  is a ring map. It is enough to check modulo  $\varpi^m$  for any  $m \geq 1$ . For this, we use that the  $m$ -th ghost map

$$W(R^+) \rightarrow R^+/\varpi^m : (x_0, x_1, \dots) \mapsto \sum_{n=0}^m x_n^{p^{m-n}} p^n$$

factors uniquely over  $W(R^+/\varpi)$ , by obvious congruences; the induced map  $W(R^+/\varpi) \rightarrow R^+/\varpi^m$  must be a ring homomorphism. Now the composite

$$W(R^{b+}) \rightarrow W(R^+/\varpi) \rightarrow R^+/\varpi^m,$$

where the first map is given by the  $m$ -th component map  $R^{b+} = \varprojlim_{x \mapsto x^p} R^+/\varpi \rightarrow R^+/\varpi$ , is a ring map, which we claim is equal to  $\theta$  modulo  $\varpi^m$ . This is a direct verification from the definitions.

For surjectivity of  $\theta$ , choose a pseudouniformizer  $\varpi^b$  of  $R^b$  such that the  $p$ -th power of  $\varpi = (\varpi^b)^\#$  divides  $p$ . We know that  $R^{b+} \rightarrow R^+/\varpi$  is surjective, which shows that  $\theta \pmod{[\varpi^b]}$  is surjective. As everything is  $[\varpi^b]$ -adically complete, this implies that  $\theta$  is surjective.

For part (ii), we claim first that there exists  $f \in \varpi^b R^{b+}$  such that  $f^\# \equiv p \pmod{p\varpi R^+}$ . Indeed, consider  $\alpha = p/\varpi \in R^+$ . There exists  $\beta \in R^{b+}$  such that  $\beta^\# \equiv \alpha \pmod{pR^+}$ . Then  $(\varpi^b \beta)^\# = \varpi \alpha \equiv p \pmod{p\varpi R^+}$ , and we can take  $f = \varpi^b \beta$ .

Thus we can write  $p = f^\# + p\varpi^\# \sum_{n \geq 0} r_n^\# p^n$ , with  $r_n \in R^{b+}$ . We can now define  $\xi = p - [f] - [\varpi^b] \sum_{n \geq 0} [r_n]p^{n+1}$ , which is of the desired form, and which lies in the kernel of  $\theta$ . Finally we need to show that  $\xi$  generates  $\ker(\theta)$ . For this, note that  $\theta$  induces a surjective map  $f : W(R^{b+})/\xi \rightarrow R^+$ . It is enough to show that  $f$  is an isomorphism modulo  $[\varpi^b]$ , because  $W(R^{b+})/\xi$  is  $[\varpi^b]$ -torsion free and  $[\varpi^b]$ -adically complete. But

$$W(R^{b+})/(\xi, [\varpi^b]) = W(R^{b+})/(p, [\varpi^b]) = R^{b+}/\varpi^b = R^+/\varpi,$$

as desired.  $\square$

It is now straightforward to check the following theorem.

**Theorem 3.17** ([KL15], [Fon13]). *There is an equivalence of categories between:*

- (i) *Pairs  $(S, S^+)$  of a perfectoid Tate ring  $S$  and an open and integrally closed subring  $S^+ \subset S^\circ$ , and*
- (ii) *Triples  $(R, R^+, \mathcal{J})$ , where  $R$  is a perfectoid Tate ring of characteristic  $p$ ,  $R^+ \subset R^\circ$  is an open and integrally closed subring, and  $\mathcal{J} \subset W(R^+)$  is an ideal which is primitive of degree 1.*

*The functors are given by  $(S, S^+) \mapsto (S^\flat, S^{\flat+}, \ker \theta)$  and  $(R, R^+, \mathcal{J}) \mapsto (W(R^+)[[\varpi]^{-1}]/\mathcal{J}, W(R^+)/\mathcal{J})$ , where  $\varpi \in R$  is any pseudouniformizer.  $\square$*

Now we can define perfectoid spaces.

**Theorem 3.18** ([Sch12, Thm. 6.3], [KL15, Thm. 3.6.14]). *Let  $R$  be a perfectoid Tate ring with an open and integrally closed subring  $R^+ \subset R^\circ$ , and let  $X = \mathrm{Spa}(R, R^+)$ . Then  $\mathcal{O}_X$  is a sheaf, and for all rational subsets  $U \subset X = \mathrm{Spa}(R, R^+)$ ,  $\mathcal{O}_X(U)$  is again perfectoid.*

*If  $(R^\flat, R^{\flat+})$  is the tilt of  $(R, R^+)$  and  $X^\flat = \mathrm{Spa}(R^\flat, R^{\flat+})$ , then recall that by Theorem 3.12,  $|X| \cong |X^\flat|$ , identifying rational subsets. Moreover, for any rational subset  $U \subset X$  with image  $U^\flat \subset X^\flat$ ,  $\mathcal{O}_X(U)$  is perfectoid with tilt  $\mathcal{O}_{X^\flat}(U^\flat)$ .*

**Definition 3.19.** *A perfectoid space is an adic space covered by open subspaces which are isomorphic to  $\mathrm{Spa}(R, R^+)$ , where  $R$  is a perfectoid Tate ring, and  $R^+ \subset R^\circ$  is an open and integrally closed subring.*

The tilting process glues to give a functor  $X \mapsto X^\flat$ . Theorem 3.13 globalizes to the following result.

**Corollary 3.20.** *Let  $X$  be a perfectoid space with tilt  $X^\flat$ . Then the functor  $X' \mapsto X'^\flat$  from perfectoid spaces over  $X$  to perfectoid spaces over  $X^\flat$  is an equivalence of categories.*

Finally, let us recall some almost mathematics. Let  $R$  be a perfectoid Tate ring.

**Definition 3.21.** *An  $R^\circ$ -module  $M$  is almost zero if  $\varpi M = 0$  for all pseudo-uniformizers  $\varpi$ . Equivalently, if  $\varpi$  is a fixed pseudo-uniformizer admitting  $p$ -power roots, then  $M$  is almost zero if and only if  $\varpi^{1/p^n} M = 0$  for all  $n$ .*

A similar definition applies to  $R^+$ -modules; in fact, the condition on  $M$  only uses the  $R^+$ -module structure.

**Example 3.22.** (i) If  $K$  is a perfectoid field, then the residue field  $k = \mathcal{O}_K/\mathfrak{m}_K$  is almost zero as  $\mathcal{O}_K$ -module, where  $\mathfrak{m}_K \subset \mathcal{O}_K$  is the maximal ideal. Conversely, any almost zero module over  $\mathcal{O}_K$  is a  $k$ -vector space, and thus a direct sum of copies of  $k$ .

(ii) If  $R$  is perfectoid and  $R^+ \subset R^\circ$  is any ring of integral elements, then  $R^\circ/R^+$  is almost zero. Indeed, if  $\varpi$  is a pseudo-uniformizer, and  $x \in R^\circ$ , then  $\varpi x$  is topologically nilpotent. Since  $R^+$  is open, there exists  $n$  with  $(\varpi x)^n \in R^+$ , so that  $\varpi x \in R^+$  by integral closedness.

Extensions of almost zero modules are almost zero. Thus the category of almost zero modules is a thick Serre subcategory of the category of all modules, and one can take the quotient.

**Definition 3.23.** *The category of almost  $R^\circ$ -modules, written  $R^{\circ a}\text{-mod}$ , is the quotient of the category of  $R^\circ$ -modules by the subcategory of almost zero modules.*

One can also define  $R^{+a}\text{-mod}$ , and the natural forgetful map  $R^{\circ a}\text{-mod} \rightarrow R^{+a}\text{-mod}$  is an equivalence.

**Theorem 3.24** ([Sch12], [KL15]). *Let  $R$  be a perfectoid Tate ring with an open and integrally closed subring  $R^+ \subset R^\circ$ , and let  $X = \text{Spa}(R, R^+)$ . Then the  $R^+$ -module  $H^i(X, \mathcal{O}_X^+)$  is almost zero for  $i > 0$ , and  $H^0(X, \mathcal{O}_X^+) = R^+$ .*

#### 4. SET-THEORETIC BOUNDS

In this paper, we will have to deal with many “large” constructions. However, we want to avoid set-theoretic issues, and in particular want to avoid the use of universes. The cleanest way to handle this seems to be to fix a certain cut-off cardinal  $\kappa$ .

**Lemma 4.1.** *There is an uncountable cardinal  $\kappa$  with the following properties.*

- (i) *For all cardinals  $\lambda < \kappa$ , one has  $2^\lambda < \kappa$ , i.e.  $\kappa$  is a strong limit cardinal.*
- (ii) *For all countable sequences  $\lambda_1, \lambda_2, \dots < \kappa$  of cardinals less than  $\kappa$ , the supremum of all  $\lambda_i$  is less than  $\kappa$ , i.e. the cofinality of  $\kappa$  is larger than  $\omega$ .*
- (iii) *For all cardinals  $\lambda < \kappa$ , there is a strong limit cardinal  $\kappa_\lambda < \kappa$  such that the cofinality of  $\kappa_\lambda$  is larger than  $\lambda$ .*

Note that in (iii), the set of such  $\kappa_\lambda$  is automatically cofinal (as for any  $\lambda \leq \lambda' < \kappa$ , one has  $\kappa_{\lambda'} \geq \lambda'$ , and  $\kappa_{\lambda'}$  has cofinality larger than  $\lambda$ ). Intuitively, cutting off at  $\kappa$  means the following. We are always allowed to take power sets, and countable unions. Moreover, we are allowed to do any construction that takes less than  $\kappa$  many steps, as long as the cardinality of the objects we are manipulating is uniformly bounded above by a cardinal strictly less than  $\kappa$ . Indeed, if for example we want to take a union of  $\lambda < \kappa$  many objects of size less than  $\kappa' < \kappa$ , then we can assume that  $\kappa' = \kappa_\lambda$  is of cofinality larger than  $\lambda$ , in which case the union is still of size less than  $\kappa_\lambda < \kappa$ . Moreover, if at each of the  $\lambda$  intermediate steps we are doing some auxiliary large construction that however respects sets that are less than  $\kappa_\lambda$ , then this is still admissible.

*Proof.* First, we show that for any cardinal  $\lambda$ , one can find a strong limit cardinal  $\kappa$  such that the cofinality of  $\kappa$  is larger than  $\lambda$ . For this, one starts with the countable cardinal  $\beth_0 = \aleph_0$  and then defines by transfinite induction  $\beth_\mu$  for any ordinal  $\mu$  as the power set of  $\beth_{\mu^-}$  if  $\mu$  is the successor of  $\mu^-$ , or the union of all previous  $\beth_{\mu'}$  if  $\mu$  is a limit ordinal. Then  $\beth_\mu$  is a strong limit cardinal with cofinality larger than  $\lambda$ , where  $\mu$  is any ordinal with cofinality larger than  $\lambda$  (for example, the smallest ordinal of cardinality larger than  $\lambda$ ).

Now build a sequence of cardinals  $\kappa(0), \kappa(1), \dots$  indexed by ordinals  $\mu$ , where  $\kappa(0)$  is a strong limit cardinal of cofinality  $> \omega$ , and for a successor ordinal  $\mu$  (of  $\mu^-$ ),  $\kappa(\mu)$  is a strong limit cardinal of cofinality  $> \kappa(\mu^-)$ . For limit ordinals,  $\kappa(\mu)$  is the union of all previous  $\kappa(\mu')$ . Let  $\omega_1$  be the first uncountable ordinal. We claim that  $\kappa(\omega_1)$  has the desired properties.

Indeed, any  $\lambda < \kappa(\omega_1)$  is also less than  $\kappa(\mu)$  for some  $\mu < \omega_1$ , which we may assume to be a successor ordinal, and so  $2^\lambda < \kappa(\mu) < \kappa(\omega_1)$ ; thus,  $\kappa(\omega_1)$  is a strong limit cardinal. Also,  $\omega_1$  has uncountable cofinality, so  $\kappa(\omega_1)$  has uncountable cofinality. Finally, for all  $\lambda < \kappa(\omega_1)$ , one has  $\lambda < \kappa(\mu)$  for some  $\mu < \omega_1$ , and then for all successor ordinals  $\mu' > \mu$  less than  $\omega_1$ , the cardinal  $\kappa(\mu') < \kappa(\omega_1)$  is by construction a strong limit cardinal with cofinality greater than  $\lambda$ .  $\square$

**Convention 4.2.** From now on, we fix a cardinal  $\kappa$  as in Lemma 4.1, and all of our perfectoid spaces will be assumed to be  $\kappa$ -small in the following sense.

**Definition 4.3.** A perfectoid space  $X$  is  $\kappa$ -small if the cardinality of  $|X|$  is less than  $\kappa$ , and for all open affinoid subspaces  $U = \mathrm{Spa}(A, A^+) \subset X$ , the cardinality of  $A$  is less than  $\kappa$ .

For the results of this section, it is actually enough to assume that  $\kappa$  is an uncountable strong limit cardinal.

**Remark 4.4.** If  $A$  has a dense subalgebra  $A_0 \subset A$  of cardinality less than  $\kappa$ , then  $A$  has cardinality less than  $\kappa$ . Indeed, identifying elements of  $A$  with convergent Cauchy sequences in  $A_0$ , one sees that if  $\lambda$  is the cardinality of  $A_0$ , then the cardinality of  $A$  is at most

$$\lambda^\omega \leq (2^\lambda)^\omega = 2^{\lambda \times \omega} = 2^\lambda < \kappa .$$

**Proposition 4.5.** A perfectoid space  $X$  is  $\kappa$ -small if and only if  $X$  admits a cover by less than  $\kappa$  many open affinoid subspaces  $U = \mathrm{Spa}(A, A^+)$  for which the cardinality of  $A$  is less than  $\kappa$ .

In particular, the category of  $\kappa$ -small perfectoid spaces is stable under finite products.

*Proof.* Assume that  $X$  is  $\kappa$ -small. In that case, the set of all subsets of  $X$  is of cardinality less than  $\kappa$ , and in particular  $X$  is covered by less than  $\kappa$  many open subspaces of the form  $U = \mathrm{Spa}(A, A^+)$ , where the cardinality of  $A$  is less than  $\kappa$ .

Conversely, for any open affinoid subspace  $U = \mathrm{Spa}(A, A^+) \subset X$ , the space  $|U|$  is a subset of the space of binary relations on  $A$  (where  $x \in U$  corresponds to the relation  $|a(x)| \leq |b(x)|$  for  $a, b \in A$ ). Thus,  $U$  embeds into the power set of  $A \times A$ , which is of cardinality  $2^{|A|^2} = 2^{|A|} < \kappa$ . If  $X$  is covered by less than  $\kappa$  many such  $U$ , then also the cardinality of  $|X|$  is less than  $\kappa$ . Also, for any open affinoid subspace  $U = \mathrm{Spa}(A, A^+)$  of  $X$ , one can (by quasicompactness) embed  $U$  into a union of finitely many rational subsets of  $V = \mathrm{Spa}(B, B^+)$  for which  $B$  has cardinality less than  $\kappa$ . Each of the rational subsets is of the form  $\mathrm{Spa}(B', B'^+)$ , where  $B'$  has a dense subalgebra which is finitely generated over  $B$ ; in particular, the cardinality of  $B'$  is also less than  $\kappa$ . As then  $A$  embeds into the finite product of the  $B'$ 's, one also sees that the cardinality of  $A$  is less than  $\kappa$ .  $\square$

## 5. MORPHISMS OF PERFECTOID SPACES

In this section, we define some basic classes of maps between perfectoid spaces.

**Definition 5.1.** A map  $f : Y \rightarrow X$  of perfectoid spaces is an injection if for all perfectoid spaces  $Z$ , the map  $f_* : \mathrm{Hom}(Z, Y) \rightarrow \mathrm{Hom}(Z, X)$  is injective.

We remind the reader that Convention 4.2 is in place; this means that implicitly, we assume that  $X$  and  $Y$  are  $\kappa$ -small, and the condition ranges over all  $\kappa$ -small  $Z$ . We note that this condition could therefore a priori depend on the choice of  $\kappa$ ; however, it follows from Proposition 5.3 (specifically, condition (iii)) that it does not. Similar remarks apply to all other definitions below.

It is enough to check the condition for affinoid perfectoid  $Z$ .

**Example 5.2.** Let  $X$  be a perfectoid space, and  $x \in X$  a point, giving rise to the map

$$i_x : \mathrm{Spa}(K(x), K(x)^+) \rightarrow X ,$$

where  $K(x)$  is the completed residue field at  $x$ , and  $K(x)^+ \subset K(x)$  the corresponding open and bounded valuation subring. Then the map  $i_x$  is an injection. To check this, we may assume that  $X$  is affinoid by replacing  $X$  by an affinoid neighborhood of  $x$ . Now

$$|\mathrm{Spa}(K(x), K(x)^+)| = \bigcap_{U \ni x} U \subset X ,$$

where the intersection runs over all rational neighborhoods  $U$  of  $x$  in  $X$ . In fact,  $\mathrm{Spa}(K(x), K(x)^+)$  is the inverse limit of  $U$  (considered as a perfectoid space) in the category of perfectoid spaces, as

$$K(x)^+/\varpi = \varinjlim_{U \ni x} \mathcal{O}_X^+(U)/\varpi ,$$

and so  $K(x)^+$  is the  $\varpi$ -adic completion of  $\varinjlim_{U \ni x} \mathcal{O}_X^+(U)$ . As each  $U \rightarrow X$  is an injection, it follows that  $i_x$  is an injection.

In particular, note that if  $X$  is qcqs and has a unique closed point  $x \in X$ , then  $X = \mathrm{Spa}(K(x), K(x)^+)$ , as in this case  $X$  is the only quasicompact open subset of  $X$  containing  $x$ .

We have the following characterizations of injections.

**Proposition 5.3.** *Let  $f : Y \rightarrow X$  be a map of perfectoid spaces. The following conditions are equivalent.*

- (i) *The map  $f$  is an injection.*
- (ii) *For all perfectoid fields  $K$  with an open and bounded valuation subring  $K^+ \subset K$ , the map  $f_* : Y(K, K^+) \rightarrow X(K, K^+)$  is injective.*
- (iii) *The map  $|f| : |Y| \rightarrow |X|$  is injective, and for all rank-1-points  $y \in Y$  with image  $x = f(y) \in X$ , the map of completed residue fields  $K(x) \rightarrow K(y)$  is an isomorphism.*
- (iv) *The map  $|f| : |Y| \rightarrow |X|$  is injective, and the map  $f : Y \rightarrow X$  is final in the category of maps  $g : Z \rightarrow X$  for which  $|g| : |Z| \rightarrow |X|$  factors over a continuous map  $|Z| \rightarrow |Y|$ .*

In particular, by (iv), given  $X$ , injections  $Y \hookrightarrow X$  of perfectoid spaces are determined by the injective map of topological spaces  $|Y| \rightarrow |X|$ . We warn the reader that in general  $|Y|$  may not have the subspace topology from  $|X|$ .

*Proof.* Clearly, (i) implies (ii). Next, we check that (ii) implies (iii). To see that  $|f|$  is injective, assume that two points  $y_1, y_2 \in Y$  map to  $x \in X$ . We may replace  $X$  by  $\mathrm{Spa}(K, K^+)$  via base change along  $i_x$  as in Example 5.2. There exists an extension  $(K, K^+) \subset (L, L^+)$  of affinoid perfectoid fields such that there are  $(L, L^+)$ -valued points  $\tilde{y}_1, \tilde{y}_2 \in Y(L, L^+)$  which send the closed point of  $\mathrm{Spa}(L, L^+)$  to  $y_1$  resp.  $y_2$ , and with same image in  $X(L, L^+)$  given by the map  $(K, K^+) \rightarrow (L, L^+)$ . As  $f$  satisfies (ii), this implies  $\tilde{y}_1 = \tilde{y}_2$ , and thus  $y_1 = y_2$ .

Now given a rank-1-point  $y \in Y$  with image  $x \in X$ , we get a commutative diagram

$$\begin{array}{ccc} \mathrm{Spa}(K(y), \mathcal{O}_{K(y)}) & \xrightarrow{i_y} & Y \\ \downarrow & & \downarrow f \\ \mathrm{Spa}(K(x), \mathcal{O}_{K(x)}) & \xrightarrow{i_x} & X . \end{array}$$

It follows that  $g : \mathrm{Spa}(K(y), \mathcal{O}_{K(y)}) \rightarrow \mathrm{Spa}(K(x), \mathcal{O}_{K(x)})$  is functorially injective on  $(K, K^+)$ -valued points for all  $(K, K^+)$ . We claim that the two maps

$$K(y) \rightarrow K(y) \widehat{\otimes}_{K(x)} K(y)$$

agree. Indeed, they agree after composition with  $K(y) \widehat{\otimes}_{K(x)} K(y) \rightarrow L$  for any perfectoid field  $L$  by assumption; but as  $K(y) \widehat{\otimes}_{K(x)} K(y)$  is perfectoid (in particular, uniform), no element lies in the kernel of all maps to perfectoid fields. This implies that the surjection  $K(y) \widehat{\otimes}_{K(x)} K(y) \rightarrow K(y)$  is an isomorphism. It follows that also the base change  $K(y) \rightarrow K(y) \widehat{\otimes}_{K(x)} K(y)$  of  $K(x) \rightarrow K(y)$  is an isomorphism. Let  $V = K(y)/K(x)$ , which is a  $K(x)$ -Banach space; then (by exactness of  $\widehat{\otimes}_{K(x)} K(y)$ ), we have  $V \widehat{\otimes}_{K(x)} K(y) = 0$ , which implies that  $V = 0$ . Thus,  $K(y) = K(x)$ , as desired, finishing the proof that (ii) implies (iii).

To check that (iii) implies (iv), we first check that for all perfectoid fields  $K$  with an open and bounded valuation subring  $K^+ \subset K$ , the map  $Y(K, K^+) \rightarrow X(K, K^+)$  is injective, with image those maps  $\mathrm{Spa}(K, K^+) \rightarrow X$  that factor over  $|Y|$ . Thus, assume given a map  $\mathrm{Spa}(K, K^+) \rightarrow X$ . The image of the closed point is a point  $x \in X$ , and  $\mathrm{Spa}(K, K^+)$  factors uniquely over the injection  $i_x : \mathrm{Spa}(K(x), K(x)^+) \rightarrow X$ . Replacing  $X$  by  $\mathrm{Spa}(K(x), K(x)^+)$  and  $Y$  by  $Y \times_X \mathrm{Spa}(K(x), K(x)^+)$ , we may assume that  $X = \mathrm{Spa}(K(x), K(x)^+)$ . Now  $|Y| \subset |X|$  is a subset stable under generalizations; in particular, any quasicompact open subset of  $Y$  has a unique closed point, so  $Y$  is an increasing union of open subfunctors of the form  $\mathrm{Spa}(K(x), (K(x)^+)' ) \subset \mathrm{Spa}(K(x), K(x)^+)$ , for varying  $(K(x)^+)'$ . But it is easy to see that for any such  $(K(x)^+)'$ ,

$$\mathrm{Spa}(K(x), (K(x)^+)' )(K, K^+) \rightarrow \mathrm{Spa}(K(x), K(x)^+)(K, K^+)$$

is injective, with image those maps that factor through  $|\mathrm{Spa}(K(x), (K(x)^+)' )| \subset |\mathrm{Spa}(K(x), K(x)^+)|$ ; thus, the same holds for  $Y$ .

Now take any map  $g : Z \rightarrow X$  such that  $|g|$  factors continuously over  $|Y| \subset |X|$ . We need to check that  $g$  factors uniquely over a map  $h : Z \rightarrow Y$ . This can be done locally on  $Z$ . Given a point  $z \in Z$ , we can replace  $Z$  by a small open affinoid neighborhood of  $z$  so that  $|Z| \rightarrow |X|$  factors over an affinoid open subset of  $X$ , and also the map  $|Z| \rightarrow |Y|$  factors over an affinoid open subset of  $Y$ . Thus, we can assume that  $X, Y$  and  $Z$  are all affinoid. Now consider the qcqs map  $Y \times_X Z \rightarrow Z$ . By Lemma 5.4 below, it is enough to check that  $(Y \times_X Z)(K, K^+) \rightarrow Z(K, K^+)$  is bijective for all perfectoid fields  $K$  with an open and bounded valuation subring  $K^+ \subset K$ . But this follows from the description of the map  $Y(K, K^+) \rightarrow X(K, K^+)$  above.

Finally, (iv) implies (i): Given any map  $Z \rightarrow X$ , we need to see that there is at most one way to factor it over  $Y$ . If  $|Z| \rightarrow |X|$  does not factor continuously over  $|Y|$ , there is no such factorization. However, if  $|Z| \rightarrow |X|$  factors continuously over  $|Y|$ , then there is a unique factorization over  $Y$ , by the universal property of  $Y \rightarrow X$  in (iv).  $\square$

**Lemma 5.4.** *Let  $f : Y \rightarrow X$  be a qcqs map of perfectoid spaces. The following conditions are equivalent.*

- (i) *The map  $f$  is an isomorphism.*
- (ii) *The map  $|f| : |Y| \rightarrow |X|$  is bijective, and for all rank-1-points  $y \in Y$  with image  $x \in X$ , the induced map of residue fields  $K(x) \rightarrow K(y)$  is an isomorphism.*
- (iii) *For all perfectoid fields  $K$  with open and bounded valuation subring  $K^+ \subset K$ , the map  $Y(K, K^+) \rightarrow X(K, K^+)$  is a bijection.*

(iv) For all algebraically closed perfectoid fields  $C$  with an open and bounded valuation subring  $C^+ \subset C$ , the map  $Y(C, C^+) \rightarrow X(C, C^+)$  is a bijection.

*Proof.* Clearly (i) implies (ii) and (iii). First, we check that (ii) and (iii) are equivalent. To see that (ii) implies (iii), take any map  $\mathrm{Spa}(K, K^+) \rightarrow X$ . Let  $x \in X$  be the image of the closed point; then the map factors over the immersion  $i_x : \mathrm{Spa}(K(x), K(x)^+) \rightarrow X$ . We can replace  $X$  by  $\mathrm{Spa}(K(x), K(x)^+)$ . Now  $Y$  has a unique closed point  $y \in Y$ , so that  $Y = \mathrm{Spa}(K(y), K(y)^+)$ . By the assumption in (ii), we have  $K(x) = K(y)$ , and then bijectivity forces  $K(y)^+ = K(x)^+$ . Now it is clear that (iii) holds.

For the converse (iii) implies (ii), assume that (iii) holds. To show that this implies (ii), it suffices to show that for any  $x \in X$ , the map  $Y \times_X \mathrm{Spa}(K(x), K(x)^+) \rightarrow \mathrm{Spa}(K(x), K(x)^+)$  is an isomorphism. We may assume that  $X = \mathrm{Spa}(K(x), K(x)^+)$ . The condition in (iii) gives us a unique section  $X \rightarrow Y$ . Let  $y \in Y$  be the image of the closed point  $x \in X$ . Note that the immersion  $i_y : Y' = \mathrm{Spa}(K(y), K(y)^+) \rightarrow Y$  has the property that  $Y' \rightarrow X$  still satisfies the conditions of (iii): Indeed,  $Y'(K, K^+) \subset Y(K, K^+) = X(K, K^+)$  is certainly injective, but there is a section  $X \rightarrow Y'$ , so that it is also surjective. In particular, any point of  $Y$  lies in  $Y'$ , so that in fact  $Y = Y'$ . As  $Y \rightarrow X$  has a section, we get maps  $K(x) \rightarrow K(y) \rightarrow K(x)$  of fields whose composite is the identity; as maps of fields are injective, this implies  $K(y) = K(x)$ , and then also  $K(y)^+ = K(x)^+$  as  $K(x)^+ \subset K(y)^+ \subset K(x)^+$ . Thus,  $Y \cong X$ , as desired.

Now we show that (ii) and (iii) together imply (i). We can assume that  $X$  is affinoid; then, by assumption on  $f$ ,  $Y$  is qcqs. Then  $|f| : |Y| \rightarrow |X|$  is a bijective, generalizing and spectral map of spectral spaces. By Lemma 2.5, it is a quotient map, and thus a homeomorphism. It remains to see that  $\mathcal{O}_X^+ \rightarrow \mathcal{O}_Y^+$  is an isomorphism of sheaves, for which it is enough to show that  $\mathcal{O}_X^+/\varpi \rightarrow \mathcal{O}_Y^+/\varpi$  is an isomorphism of sheaves, where  $\varpi$  is a pseudouniformizer on  $X$ . We can check this on stalks. But the stalk of  $\mathcal{O}_X^+/\varpi$  at  $x \in X$  is given by  $K(x)^+/\varpi$ , and similarly for  $\mathcal{O}_Y^+/\varpi$ . As  $K(x)^+/\varpi = K(y)^+/\varpi$  (by the proof that (iii) implies (ii)), the result follows.

Clearly, (iii) implies (iv), so it remains to prove that (iv) implies (iii). Note that injectivity of  $Y(K, K^+) \rightarrow X(K, K^+)$  is automatic by embedding  $(K, K^+)$  into an algebraic closure  $(C, C^+)$ . But in fact, given any map  $\mathrm{Spa}(K, K^+) \rightarrow X$ , we can lift to a map  $\mathrm{Spa}(C, C^+) \rightarrow Y$ , which factors over some affinoid  $\mathrm{Spa}(S, S^+) \subset Y$ , and is given by a map  $(S, S^+) \rightarrow (C, C^+)$ . But this map has to be Galois-equivariant, and thus factors over  $(K, K^+)$ . This gives the desired lift  $\mathrm{Spa}(K, K^+) \rightarrow Y$ .  $\square$

Moreover, injections have a simple behaviour on pullbacks.

**Corollary 5.5.** (i) Let  $f : Y \rightarrow X$  be an injection of perfectoid spaces, and  $X' \rightarrow X$  any map of perfectoid spaces. Then the pullback  $f' : Y' = X' \times_X Y \rightarrow X'$  is an injection, and the map

$$|Y'| \rightarrow |X'| \times_{|X|} |Y|$$

is a homeomorphism.

(ii) Let  $f : Y \rightarrow X$  be a map of perfectoid spaces. Then  $f$  is an injection if and only if  $f$  is universally injective, i.e. for all maps  $X' \rightarrow X$  with pullback  $f' : Y' = X' \times_X Y \rightarrow X'$ , the map  $|f'| : |Y'| \rightarrow |X'|$  is injective.

*Proof.* In part (i), it is clear from the definition that  $f'$  is an injection. Using that  $|f|$  and  $|f'|$  are injective by Proposition 5.3, it follows that  $|Y'| \rightarrow |X'| \times_{|X|} |Y|$  is injective. Moreover, all maps of perfectoid spaces are generalizing, so both  $|Y'|$  and  $|X'| \times_{|X|} |Y|$  are generalizing subsets of  $|Y|$ , and

thus the map  $|Y'| \rightarrow |X'| \times_{|X|} |Y|$  is generalizing. But this map is always surjective and spectral. By Lemma 2.5, it is a homeomorphism.

For part (ii), we have seen that if  $f$  is an injection, then it is universally injective. For the converse, assume that  $f$  is universally injective. Given any map  $g : Z \rightarrow X$ , we need to see that there is at most one way to factor it over  $f : Y \rightarrow X$ . Pulling back by  $g$ , we may assume that  $Z = X$ , and we have to show that there is at most one section of  $f$ . Assume that there is a section  $h : X \rightarrow Y$ . Being a section,  $h$  is an injection; but then  $|X| \rightarrow |Y| \rightarrow |X|$  is a factorization of the identity as a composition of injective maps. Thus,  $|Y| = |X|$ , and the map is a homeomorphism; then, by Proposition 5.3, it follows that  $Y = X$ . In particular,  $h$  is the only section, as desired.  $\square$

As injections are determined by their behaviour on topological spaces, the following definition is reasonable.

**Definition 5.6.** *A map  $f : Y \rightarrow X$  of perfectoid spaces is an immersion if  $f$  is an injection and  $|f| : |Y| \rightarrow |X|$  is a locally closed immersion. If  $|f|$  in addition is open (resp. closed), then  $f$  is an open (resp. closed) immersion.*

One cannot define closed immersions of schemes or adic spaces this way because of possible non-reduced structures. This is not a concern for perfectoid spaces, which makes this general definition possible.

Zariski closed embeddings as defined in [Sch15, Section II.2] are a special class of closed embeddings.

**Definition 5.7** ([Sch15, Definition II.2.1, Definition II.2.6]). *Let  $f : Z \rightarrow X$  be a map of perfectoid spaces, where  $X = \mathrm{Spa}(R, R^+)$  is affinoid.*

- (i) *The map  $f$  is Zariski closed if  $f$  is a closed immersion and there is an ideal  $I \subset R$  such that  $|Z| \subset |X|$  is the locus where  $|f| = 0$  for all  $f \in I$ .*
- (ii) *The map  $f$  is strongly Zariski closed if  $Z = \mathrm{Spa}(S, S^+)$  is affinoid,  $R \rightarrow S$  is surjective, and  $S^+$  is the integral closure of  $R^+$  in  $S$ .*

By [KL15, Proposition 3.6.9 (c)], if  $f$  is strongly Zariski closed, then  $R^+ \rightarrow S^+$  is almost surjective, so that the definition agrees with [Sch15, Definition II.2.6].

**Proposition 5.8.** *Let  $f : Z \rightarrow X$  be a map of perfectoid spaces, where  $X$  is affinoid.*

- (i) *If  $f$  is strongly Zariski closed, then  $f$  is Zariski closed, and in particular a closed immersion.*
- (ii) *If  $f$  is Zariski closed, then  $Z$  is affinoid.*
- (iii) *If  $X$  is of characteristic  $p$  and  $f$  is Zariski closed, then  $f$  is strongly Zariski closed.*

*Proof.* Cf. [Sch15, Section II.2].  $\square$

A general class of immersions is given by diagonal morphisms.

**Proposition 5.9.** *Let  $f : Y \rightarrow X$  be any map of perfectoid spaces. Then the diagonal morphism*

$$\Delta_f : Y \rightarrow Y \times_X Y$$

*is an immersion.*

*Proof.* Clearly,  $\Delta_f$  is an injection. Thus, it is enough to show that  $|\Delta_f|$  identifies  $|Y|$  with a locally closed subspace of  $|Y \times_X Y|$ . This statement can be checked locally on  $X$  and  $Y$ ; thus, we can



assume that  $X = \mathrm{Spa}(R, R^+)$  and  $Y = \mathrm{Spa}(S, S^+)$  are affinoid. But then  $\Delta_f$  is strongly Zariski closed, as  $S \widehat{\otimes}_R S \rightarrow S$  is surjective. By Proposition 5.8, it follows that  $\Delta_f$  is a closed immersion.  $\square$

**Definition 5.10.** *A map  $f : Y \rightarrow X$  of perfectoid spaces is separated if  $\Delta_f$  is a closed immersion.*

**Proposition 5.11.** *Let  $f : Y \rightarrow X$  be a map of perfectoid spaces. The following conditions are equivalent.*

- (i) *The map  $f$  is separated.*
- (ii) *The map  $|\Delta_f| : |Y| \rightarrow |Y \times_X Y|$  is a closed immersion.*
- (iii) *The map  $f$  is quasiseparated (i.e.,  $|f| : |Y| \rightarrow |X|$  is quasiseparated), and for all perfectoid fields  $K$  with ring of integers  $\mathcal{O}_K$  and an open and bounded valuation subring  $K^+ \subset K$  and any commutative diagram*

$$\begin{array}{ccc} \mathrm{Spa}(K, \mathcal{O}_K) & \longrightarrow & Y \\ \downarrow & \dashrightarrow & \downarrow f \\ \mathrm{Spa}(K, K^+) & \longrightarrow & X \end{array},$$

*there exists at most one dotted arrow making the diagram commute.*

*Proof.* The equivalence of (i) and (ii) follows from Proposition 5.9. Next, we check that (ii) implies (iii). If  $|\Delta_f|$  is a closed immersion, then it is in particular quasicompact; this implies that  $f$  is quasiseparated. Now assume that there are two dotted arrows making the diagram commute. These define a point  $z \in (Y \times_X Y)(K, K^+)$  such that  $z|_{(K, \mathcal{O}_K)} \in \Delta_f(Y)(K, \mathcal{O}_K)$ . But if  $|\Delta_f| : |Y| \rightarrow |Y \times_X Y|$  is a closed immersion, then the map  $z : \mathrm{Spa}(K, K^+) \rightarrow Y \times_X Y$  maps into  $|\Delta_f|(|Y|)$  if this is true for  $z|_{(K, \mathcal{O}_K)}$ , as  $\mathrm{Spa}(K, \mathcal{O}_K) \subset \mathrm{Spa}(K, K^+)$  is dense. As  $\Delta_f$  is an immersion, this implies that  $z$  maps  $\mathrm{Spa}(K, K^+)$  into  $\Delta_f(Y)$ , so that the dotted arrows agree.

Finally, (iii) implies (ii): The conditions in (iii) imply that the map  $|\Delta_f| : |Y| \rightarrow |Y \times_X Y|$  is a quasicompact locally closed immersion of locally spectral spaces, which is moreover specializing. We claim that this implies that  $|\Delta_f|$  is a closed immersion. By quasicompactness, the image of  $|\Delta_f|$  is pro-constructible, thus the closure is given by the set of specializations; but the image is already closed under specializations.  $\square$

## 6. ÉTALE MORPHISMS

In this section, we recall briefly some facts about étale morphisms of perfectoid spaces. We start with the almost purity theorem.

**Theorem 6.1** ([Fal02], [KL15, Theorem 3.6.21, Theorem 5.5.9], [Sch12, Theorem 5.25, Theorem 7.9 (iii)]). *Let  $R$  be a perfectoid Tate ring with tilt  $R^\flat$ .*

- (i) *For any finite étale  $R$ -algebra  $S$ ,  $S$  is perfectoid.*
- (ii) *Tilting induces an equivalence*

$$\begin{aligned} \{ \text{Finite étale } R\text{-algebras} \} &\rightarrow \{ \text{Finite étale } R^\flat\text{-algebras} \} \\ S &\mapsto S^\flat \end{aligned}$$

- (iii) *For any finite étale  $R$ -algebras  $S$ ,  $S^\circ$  is almost finite étale over  $R^\circ$ .*

**Definition 6.2.** *Let  $f : Y \rightarrow X$  be a morphism of perfectoid spaces.*

- (i) The morphism  $f$  is finite étale if for all open affinoid perfectoid  $\mathrm{Spa}(R, R^+) = U \subset X$ , the preimage  $V = f^{-1}(U) = \mathrm{Spa}(S, S^+) \subset Y$  is affinoid perfectoid,  $R \rightarrow S$  is a finite étale morphism of rings, and  $S^+$  is the integral closure of  $R^+$  in  $S$ .
- (ii) The morphism  $f$  is étale if for all  $y \in Y$ , there are open subsets  $V \subset Y$ ,  $U \subset X$  such that  $y \in V$ ,  $f(V) \subset U$ , and there is a factorization

$$\begin{array}{ccc} V & \hookrightarrow & W \\ & \searrow f|_V & \downarrow \\ & & U, \end{array}$$

where  $V \hookrightarrow W$  is an open immersion, and  $W \rightarrow U$  is finite étale.

We refer to [Sch12, Section 7] for an extensive discussion. In particular, it is proved there that a morphism is finite étale if and only if there is a cover by open affinoid perfectoid  $U = \mathrm{Spa}(R, R^+) \subset X$  such that their preimages  $V = \mathrm{Spa}(S, S^+) \subset Y$  are affinoid perfectoid,  $R \rightarrow S$  is finite étale, and  $S^+$  is the integral closure of  $R^+$  in  $S$ . Moreover, composites, base changes, and maps between (finite) étale maps are (finite) étale.

By  $X_{\mathrm{fét}}$ , resp.  $X_{\mathrm{ét}}$ , we denote the categories of finite étale, resp. étale, perfectoid spaces over  $X$ . If  $X = \mathrm{Spa}(R, R^+)$  is affinoid, then  $X_{\mathrm{fét}} \cong R_{\mathrm{fét}}^{\mathrm{op}}$ .

**Theorem 6.3** ([Sch12], [KL15]). *Let  $X$  be a perfectoid space with tilt  $X^{\flat}$ . Then tilting induces an equivalence  $X_{\mathrm{ét}} \cong X_{\mathrm{ét}}^{\flat}$ .*

*If  $X = \mathrm{Spa}(R, R^+)$  is affinoid perfectoid, then the  $R^+$ -module  $H^i(X_{\mathrm{ét}}, \mathcal{O}_X^+)$  is almost zero for  $i > 0$ , and  $H^0(X_{\mathrm{ét}}, \mathcal{O}_X^+) = R^+$ .*

We will need a result about étale morphisms to an inverse limit of spaces.

**Proposition 6.4.** *Let  $X_i = \mathrm{Spa}(R_i, R_i^+)$ ,  $i \in I$ , be a cofiltered inverse system of affinoid perfectoid spaces for some  $\kappa$ -small index category  $I$ , where  $\lambda < \kappa$ . Assume that all  $X_i$  are  $\kappa'$ -small for some strong limit cardinal  $\kappa' < \kappa$ . Then the inverse limit  $X = \mathrm{Spa}(R, R^+)$  is still  $\kappa$ -small, where  $R^+$  is the  $\varpi$ -adic completion of  $\varinjlim_i R_i^+$ , and  $R = R^+[\frac{1}{\varpi}]$ . Here,  $\varpi \in R_i^+$  denotes any compatible choice of pseudouniformizers for large  $i$ . Moreover, one has the following results.*

- (o) The map

$$|X| \rightarrow \varprojlim_i |X_i|$$

is a homeomorphism of spectral spaces.

- (i) The base change functors  $(X_i)_{\mathrm{fét}} \rightarrow X_{\mathrm{fét}}$  induce an equivalence of categories

$$\varprojlim_i (X_i)_{\mathrm{fét}} \rightarrow X_{\mathrm{fét}}.$$

- (ii) Let  $X_{\mathrm{ét}, \mathrm{qcqs}} \subset X_{\mathrm{ét}}$  denote the full subcategory of qcqs étale perfectoid spaces over  $X$  (and similarly for  $X_i$ ). Then the base change functors  $(X_i)_{\mathrm{ét}, \mathrm{qcqs}} \rightarrow X_{\mathrm{ét}, \mathrm{qcqs}}$  induce an equivalence of categories

$$\varprojlim_i (X_i)_{\mathrm{ét}, \mathrm{qcqs}} \rightarrow X_{\mathrm{ét}, \mathrm{qcqs}}.$$

(iii) Let  $X_{\text{ét, qc, sep}} \subset X_{\text{ét, qcqs}}$  be the full subcategory of quasicompact separated étale perfectoid spaces over  $X$  (and similarly for  $X_i$ ). Then the base change functors  $(X_i)_{\text{ét, qc, sep}} \rightarrow X_{\text{ét, qc, sep}}$  induce an equivalence of categories

$$2\text{-}\varinjlim_i (X_i)_{\text{ét, qc, sep}} \rightarrow X_{\text{ét, qc, sep}} .$$

(iv) Let  $X_{\text{ét, aff}} \subset X_{\text{ét, qc, sep}}$  be the full subcategory of affinoid perfectoid spaces which are étale over  $X$  (and similarly for  $X_i$ ). Then the base change functors  $(X_i)_{\text{ét, aff}} \rightarrow X_{\text{ét, aff}}$  induce an equivalence of categories

$$2\text{-}\varinjlim_i (X_i)_{\text{ét, aff}} \rightarrow X_{\text{ét, aff}} .$$

*Proof.* For the set-theoretic part, we can by Lemma 4.1 assume that  $\kappa' = \kappa_\lambda$  has cofinality larger than  $\lambda$ , in which case the cardinality of the direct limit of the  $R_i^+$  is still less than  $\kappa_\lambda$ , and so  $R$  is  $\kappa$ -small.

Part (o) follows from [Hub93a, Proposition 3.9], cf. [Sch12, Lemma 6.13 (ii)]. The statement in part (i) is equivalent to

$$2\text{-}\varinjlim_i (R_i)_{\text{fét}} \rightarrow R_{\text{fét}}$$

being an equivalence of categories. This is a consequence [Sch12, Lemma 7.5 (i)] of a theorem of Elkik, [Elk73], (in the noetherian case) and Gabber–Ramero, [GR03, Proposition 5.4.53], in general.

For part (ii), we first check fully faithfulness. Thus, let  $Y_i, Y'_i \rightarrow X_i$  be two qcqs étale perfectoid spaces over  $X_i$ , and denote by  $Y_j, Y'_j \rightarrow X_j$  their pullbacks to  $X_j$  for  $j \geq i$ , and  $Y, Y' \rightarrow X$  their pullbacks to  $X$ . Assume that two morphisms  $f_i, g_i : Y_i \rightarrow Y'_i$  become equal after pullback to  $X$ ,  $f = g : Y \rightarrow Y'$ . Let  $\Gamma_{f_i}, \Gamma_{g_i} : Y_i \hookrightarrow Y_i \times_X Y'_i =: Z_i$  be the graphs of  $f_i$  and  $g_i$ . Then  $\Gamma_{f_i}$  and  $\Gamma_{g_i}$  are injections, and thus are determined by the quasicompact open image of  $|Y_i|$  in  $|Z_i|$ . We know that after pullback along  $|Z| \rightarrow |Z_i|$ ,  $Z = Z_i \times_{X_i} X$ , the two quasicompact open images  $\Gamma_{f_i}(|Y_i|)$ ,  $\Gamma_{g_i}(|Y_i|)$  agree. As  $|Z| = \varprojlim_j |Z_j|$  is an inverse limit of spectral spaces along spectral maps, these two quasicompact open images agree after pullback to  $|Z_j|$  for large  $j$ , which means that  $\Gamma_{f_j} = \Gamma_{g_j}$ , as desired. This proves faithfulness.

As an intermediate step to fullness, we check that if  $f_i : Y_i \rightarrow Y'_i$  is a morphism over  $X$  whose pullback  $f : Y \rightarrow Y'$  to  $X$  is an isomorphism, then the pullback  $f_j : Y_j \rightarrow Y'_j$  to  $X_j$  is an isomorphism for large enough  $j$ . Note that the image of  $f_i$  is a quasicompact open subset, over which  $Y' \rightarrow Y'_i$  factors; a standard quasicompactness argument implies that  $Y'_j \rightarrow Y'_i$  factors over this open subset for large enough  $j$ , so that  $f_j$  is surjective for large enough  $j$ . Applying the same reasoning to  $\Delta_{f_i} : Y_i \rightarrow Y_i \times_{Y'_i} Y_i$  shows that  $\Delta_{f_j}$  is surjective for large enough  $j$ . As  $\Delta_{f_j}$  is always an injection, this implies that  $\Delta_{f_j}$  is an isomorphism, which then implies that  $f_j$  is an isomorphism.

Now, let  $f : Y \rightarrow Y'$  be a morphism over  $X$ ; we need to see that this is a base change of a morphism  $f_j : Y_j \rightarrow Y'_j$  over  $X_j$  for large enough  $j$ . The graph  $\Gamma_f : Y \hookrightarrow Y \times_X Y'$  is a quasicompact open embedding. By approximation of quasicompact open subsets, we can find a quasicompact open immersion  $V_j \subset Y_j \times_{X_j} Y'_j$  whose pullback agrees with the image of  $\Gamma_f$ . In particular, the projection  $V_j \rightarrow Y_j$  of qcqs étale perfectoid spaces over  $X_j$  becomes an isomorphism after pullback to  $X$ . By the intermediate step, we see that  $V_j \rightarrow Y_j$  becomes an isomorphism for large enough  $j$ ; composing the inverse with the map  $V_j \hookrightarrow Y_j \times_{X_j} Y'_j \rightarrow Y'_j$  gives the desired map  $Y_j \rightarrow Y'_j$ .

Finally, for essential surjectivity, take any qcqs étale  $f : Y \rightarrow X$ . It is enough to cover  $Y$  by quasicompact open subsets which descend to  $X_i$  for large  $i$ , as by fully faithfulness (and the assumption that  $Y$  is qcqs), the pieces will automatically glue for large enough  $i$ . Thus, we can assume that  $f$  is a composite of a rational open subset, a finite étale map, and a rational open subset. Each of these maps descends to  $X_i$  for large enough  $i$  (using (i)), finishing the proof of (ii).

For (iii), fully faithfulness follows from (ii). It remains to see that if  $f_i : Y_i \rightarrow X_i$  is a qcqs étale map whose base change  $f : Y \rightarrow X$  to  $X$  is separated, then the base change  $f_j : Y_j \rightarrow X_j$  to  $X_j$  is separated for large enough  $j$ . The map  $f_j : Y_j \rightarrow X_j$  is separated if and only if the quasicompact open immersion  $\Delta_{f_j} : Y_j \rightarrow Y_j \times_{X_j} Y_j$  is also a closed immersion. But in general, if  $S_i$ ,  $i \in I$ , is a cofiltered inverse system of spectral spaces (with spectral maps) and inverse limit  $S$ , then if  $U_i \subset S_i$  is a quasicompact open subset whose inverse image  $U \subset S$  is open and closed, then the inverse image  $U_j \subset S_j$  is open and closed for large  $j$ . Indeed, the open and closed decomposition given by  $U$  descends to  $S_j$  for large enough  $j$ , and for large enough  $j$ , one of the two quasicompact open subsets in this decomposition is equal to  $U_j$ . Applied with  $S_i = |Y_i \times_{X_i} Y_i|$ ,  $S = |Y \times_X Y|$ , we see that if  $\Delta_f$  is an open and closed immersion, then  $\Delta_{f_j}$  is an open and closed immersion for large enough  $j$ .

In (iv), fully faithfulness again follows from (ii). Essential surjectivity follows from [Hub96, Proposition 1.7.1], which holds for affinoid perfectoid spaces by using the theory of pseudocoherent sheaves of Kedlaya–Liu, [KL16].  $\square$

## 7. TOTALLY DISCONNECTED SPACES

In this section, we introduce a notion of (strictly) totally disconnected spaces. These are analogues of profinite sets in our setting.

**Definition 7.1.** *A perfectoid space  $X$  is totally disconnected if  $X$  is qcqs and every open cover of  $X$  splits, i.e. if  $\{U_i \subset X\}$  is an open cover of  $X$ , then  $\bigsqcup U_i \rightarrow X$  has a splitting.*

The following characterization of spectral spaces for which every open cover splits is due to L. Fargues:

**Lemma 7.2.** *Let  $X$  be a spectral space. The following conditions are equivalent.*

- (i) *Every open cover of  $X$  splits.*
- (ii) *Every connected component of  $X$  has a unique closed point.*
- (iii) *The functor  $\mathcal{F} \mapsto \Gamma(X, \mathcal{F})$  on sheaves on  $X$  is exact, i.e. commutes with all finite colimits.*
- (iv) *The functor  $\mathcal{F} \mapsto \Gamma(X, \mathcal{F})$  on sheaves of abelian groups on  $X$  is exact.*
- (v) *For all sheaves of abelian groups  $\mathcal{F}$  on  $X$ , one has  $H^i(X, \mathcal{F}) = 0$  for all  $i > 0$ .*
- (vi) *For all sheaves of abelian groups  $\mathcal{F}$  on  $X$ , one has  $H^1(X, \mathcal{F}) = 0$ .*

*Proof.* For (i) implies (ii), note that the condition that any open cover splits passes to closed subsets (by adding the open complement to the open cover). Thus, we may assume that  $X$  is connected. But then if  $x, y \in X$  are two distinct closed points, then  $(X \setminus \{x\}) \sqcup (X \setminus \{y\}) \rightarrow X$  does not have a splitting. For the converse, let  $\{U_i \subset X\}$  be an open cover of  $X$ . Let  $c \in \pi_0 X$ , corresponding to a connected component  $X_c \subset X$ . Take some  $U_i$  which contains the unique closed point of  $X_c$ ; then  $X_c \subset U_i$ . It follows that there is some open and closed neighborhood  $U_c \subset X$  such that  $U_c \subset U_i$ . Thus, we can find a cover of  $\pi_0 X$  by open and closed subsets such that  $\bigsqcup U_i \rightarrow X$  splits

on the preimage. As  $\pi_0 X$  is profinite, we can find a disjoint cover of  $\pi_0 X$  by such subsets, and then assemble the local splittings into a splitting of  $\bigsqcup U_i \rightarrow X$ .

Moreover, (i) implies (iii): We have to see that  $\Gamma(X, -)$  commutes with coequalizers. Let  $\mathcal{H} = \text{coeq}(\mathcal{F} \rightrightarrows \mathcal{G})$  be a coequalizer diagram. First, we check that

$$\text{coeq}(\Gamma(X, \mathcal{F}) \rightrightarrows \Gamma(X, \mathcal{G})) \rightarrow \Gamma(X, \mathcal{H})$$

is injective. If  $t, t' \in \Gamma(X, \mathcal{G})$  are sections whose images in  $\Gamma(X, \mathcal{H})$  agree, then after a cover  $\{U_i \rightarrow X\}$ , one can find  $s_i \in \mathcal{F}(U_i)$  mapping to  $t|_{U_i}$  and  $t'|_{U_i}$ . But under (i), we can split  $\bigsqcup U_i \rightarrow X$ , giving us such a section  $s \in \Gamma(X, \mathcal{F})$ . Now assume  $s \in \Gamma(X, \mathcal{H})$  is a section. Then after a cover  $\{U_i \rightarrow X\}$ , we can lift it to sections  $s_i \in \mathcal{G}(U_i)$ . But under (i), we can split  $\bigsqcup U_i \rightarrow X$ , giving us a lift of  $s$  to  $\Gamma(X, \mathcal{G})$ , proving the desired surjectivity.

It is clear that (iii) $\Rightarrow$ (iv) $\Rightarrow$ (v) $\Rightarrow$ (vi) $\Rightarrow$ (iv). To see that (iv) implies (i), let  $\{U_1, \dots, U_n\}$  be a finite quasicompact open cover of  $X$ , let  $j_i : U_i \rightarrow X$  be the open immersions, and consider the surjection

$$\mathcal{F} = \bigoplus_i j_{i!} \mathbb{Z} \rightarrow \mathbb{Z}$$

of sheaves on  $X$ . Under condition (iv), this is surjective on global sections. This implies that there are locally constant functions  $f_i : X \rightarrow \mathbb{Z}$  with  $V_i = \text{supp } f_i \subset U_i$  such that  $\sum_i f_i = 1$ . In particular, all  $V_i$  are open and closed, and

$$X = \bigsqcup_i (V_i \setminus \bigcup_{j < i} V_j) \rightarrow \bigsqcup_i U_i \rightarrow X$$

gives a section, as desired.  $\square$

For perfectoid spaces, one can give a finer structural analysis.

**Lemma 7.3.** *Let  $X$  be a totally disconnected perfectoid space.*

- (i) *There is a continuous projection  $\pi : X \rightarrow \pi_0(X)$  to the profinite set  $\pi_0(X)$  of connected components.*
- (ii) *All fibres of  $\pi$  are of the form  $\text{Spa}(K, K^+)$  for some perfectoid field  $K$  with an open and bounded valuation subring  $K^+ \subset K$ .*

*Conversely, if  $X$  is a qcqs perfectoid space all of whose connected components are of the form  $\text{Spa}(K, K^+)$  for some perfectoid field  $K$  with an open and bounded valuation subring  $K^+ \subset K$ , then  $X$  is totally disconnected.*

*Proof.* Part (i) is a consequence of the assumption that the underlying topological space of  $X$  is a spectral space. For part (ii), note that in a spectral space, any connected component is an intersection of open and closed subsets; this implies that connected components of qcqs perfectoid spaces are again perfectoid spaces. Thus, for (ii), we have to classify the connected totally disconnected perfectoid spaces. By Lemma 7.2,  $X$  has a unique closed point. By Example 5.2, it follows that if  $x \in X$  is the unique closed point of a perfectoid space  $X$ , then  $X = \text{Spa}(K(x), K(x)^+)$ , as desired.

The converse follows from Lemma 7.2, as then all connected components have a unique closed point.  $\square$

In [BS15], we worked with a stronger notion of w-local spaces; this includes the extra condition that the subset of closed points is closed. This stronger notion will have little relevance for us.

**Definition 7.4.** A perfectoid space  $X$  is *w-local* if its underlying topological space is a *w-local spectral space* in the sense of [BS15, Definition 2.1.1]. Equivalently,  $X$  is *qcqs*, for any open cover  $X = \bigcup_{i \in I} U_i$ , the map  $\bigsqcup_{i \in I} U_i \rightarrow X$  splits, and the subset  $X^c \subset X$  of closed points is closed.

In particular, w-local spaces are totally disconnected.

**Lemma 7.5.** Let  $X$  be a totally disconnected perfectoid space. Then  $X$  is affinoid.

*Proof.* Let  $c \in \pi_0(X)$  be any point, and let  $x_c \in X$  be the unique closed point mapping to  $c$ . There is an open affinoid neighborhood  $U \subset X$  of  $x_c$ ; in particular,  $U$  contains the connected component  $X_c \subset X$ , as all points of  $X_c$  specialize to  $x_c$ . We claim that there is an open compact neighborhood  $V$  of  $c$  in  $\pi_0(X)$  such that  $\pi^{-1}(V) \subset U$ . Indeed, the intersection

$$\bigcap_{c \in V} (X \setminus U) \cap \pi^{-1}(V) = \emptyset ,$$

where the intersection runs over open and closed subsets  $V \subset \pi_0(X)$  containing  $c$ . By quasicompactness of the constructible topology, it follows that there is some  $V$  such that  $(X \setminus U) \cap \pi^{-1}(V) = \emptyset$ , i.e.  $\pi^{-1}(V) \subset U$ . In this case,  $U$  has a closed and open decomposition  $U = \pi^{-1}(V) \sqcup \pi^{-1}(\pi_0(X) \setminus V)$ , so that  $\pi^{-1}(V)$  is also affinoid. Thus,  $\pi_0(X)$  is covered by open and closed subsets  $V \subset \pi_0(X)$  such that  $\pi^{-1}(V) \subset X$  is affinoid. By quasicompactness, finitely many such  $V$  cover  $\pi_0(X)$ , and we may moreover assume that they form a disjoint cover  $\pi_0(X) = V_1 \sqcup \dots \sqcup V_n$ . Then  $X$  is the disjoint union  $\pi^{-1}(V_1) \sqcup \dots \sqcup \pi^{-1}(V_n)$ , and thus affinoid.  $\square$

**Lemma 7.6.** Let  $X$  be a totally disconnected perfectoid space. Let  $U \subset X$  be a pro-constructible generalizing subset. Then  $U$  is an intersection of subsets of the form  $\{|f| \leq 1\}$  for varying  $f \in H^0(X, \mathcal{O}_X)$ . In particular,  $U$  is an affinoid perfectoid space, which is moreover totally disconnected.

**Remark 7.7.** Note that the lemma implies in particular that any quasicompact open subset of  $X$  is affinoid.

*Proof.* Let  $x \in X \setminus U$  be a point, lying in a connected component  $c = \pi(x) \in \pi_0(X)$ . As the fibre  $X_c$  is of the form  $\mathrm{Spa}(K_c, K_c^+)$  for some perfectoid field  $K_c$  and open and bounded valuation subring  $K_c^+ \subset K_c$ , one can find some  $f_c \in K_c$  such that  $|f_c| \leq 1$  on  $U_c = U \cap X_c$ , but  $|f_c(x)| > 1$ . Modulo  $K_c^+$ , we can lift  $f_c$  to a function  $f_V$  on  $\pi^{-1}(V) \subset X$  for some compact open neighborhood  $V \subset \pi_0(X)$  of  $c$ . Then by assumption

$$\bigcap_{c \in V' \subset V} (U \cap \{|f_V| > 1\}) \cap \pi^{-1}(V') = \emptyset .$$

Here, all terms are pro-constructible subsets of  $X$ , and thus compact in the constructible topology; by Tychonoff, it follows that for some open compact neighborhood  $V'$  of  $v$ ,  $(U \cap \{|f_V| > 1\}) \cap \pi^{-1}(V') = \emptyset$ ; in other words, after replacing  $V$  by  $V'$ , we can assume that  $U \cap \pi^{-1}(V) \subset \{|f_V| \leq 1\}$ . Extending  $f_V$  by 0 on  $\pi^{-1}(\pi_0(X) \setminus V)$ , we get a function  $f \in H^0(X, \mathcal{O}_X)$  with  $|f| \leq 1$  on  $U$ , but  $|f(x)| > 1$ . Taking the intersection over all such  $f$  for varying  $x \in X \setminus U$  gives  $U$ .

Thus,  $U$  is affinoid. Every connected component of  $U$  is an intersection of a connected component  $\mathrm{Spa}(K_c, K_c^+)$  of  $X$  with  $U$ , which is necessarily of the form  $\mathrm{Spa}(K_c, (K_c^+)')$  for another open and bounded valuation subring  $(K_c^+)'$  of  $K_c$ .  $\square$

The notion of w-local and totally disconnected spaces is useful as on the hand, they are rather simple, but on the other hand, there are many examples. More precisely, any qcqs perfectoid space

$X$  admits a universal map  $X^{\text{wl}} \rightarrow X$  from a w-local space  $X^{\text{wl}}$ , and the map  $X^{\text{wl}} \rightarrow X$  is pro-étale. Before we formulate this result, we recall the definition of pro-étale morphisms. We recall that Convention 4.2 is in place throughout the whole text, but we stress it again in the following definition.

**Definition 7.8.** *Let  $f : Y \rightarrow X$  be a map of  $\kappa$ -small perfectoid spaces.*

(i) *The map  $f$  is affinoid pro-étale if  $Y = \text{Spa}(S, S^+)$  and  $X = \text{Spa}(R, R^+)$  are affinoid, and one can write  $Y = \varprojlim Y_i \rightarrow X$  as a cofiltered inverse limit of étale maps  $Y_i \rightarrow X$  from affinoid perfectoid spaces  $Y_i = \text{Spa}(S_i, S_i^+)$  along a  $\kappa$ -small index category  $I$ . Any such presentation  $Y = \varprojlim Y_i \rightarrow X$  is called a pro-étale presentation.*

(ii) *The map  $f$  is pro-étale if for all  $y \in Y$ , there is an open neighborhood  $V \subset Y$  of  $y$  and an open subset  $U \subset X$  such that  $f(V) \subset U$ , and the restriction  $f|_V : V \rightarrow U$  is affinoid pro-étale.*

For example, if  $X = \text{Spa}(C, \mathcal{O}_C)$  is a geometric point, then any affinoid pro-étale map  $Y \rightarrow X$  is of the form  $Y = X \times \underline{S}$  for some profinite set  $S$ , where if  $S = \varprojlim_i S_i$  is a limit of finite sets, we set

$$X \times \underline{S} = \varprojlim_i X \times S_i .$$

**Remark 7.9.** A slightly curious example of pro-étale morphisms are Zariski closed immersions: If  $f : Z \hookrightarrow X$  is a Zariski closed immersion of affinoid perfectoid spaces, then  $f$  is affinoid pro-étale. Indeed, if  $Z = \{\forall f \in I : f = 0\} \subset X$  for an ideal  $I \subset R = \mathcal{O}_X(X)$ , then  $Z$  can be written as the intersection of the rational subsets

$$U_{f_1, \dots, f_n} = \{|f_1|, \dots, |f_n| \leq 1\}$$

over varying  $n$  and  $f_1, \dots, f_n \in I$ . Thus,  $Z = \varprojlim U_{f_1, \dots, f_n} \rightarrow X$ , which gives a pro-étale presentation of  $Z$ .

In particular, if  $f : Y \rightarrow X$  is any morphism of perfectoid spaces, then  $\Delta_f : Y \rightarrow Y \times_X Y$  is pro-étale. Indeed, this can be checked locally, so we can assume that  $X$  and  $Y$  are affinoid; then  $\Delta_f$  is a Zariski closed immersion, and thus affinoid pro-étale.

**Proposition 7.10.** *Fix an affinoid perfectoid space  $X$ , and let  $X_{\text{ét}}^{\text{aff}}$  be the category of étale maps  $f : Y \rightarrow X$  from affinoid perfectoid spaces  $Y$ , and  $X_{\text{proét}}^{\text{aff}}$  the category of affinoid pro-étale maps  $f : Y \rightarrow X$ . Then the functor*

$$\text{Pro}_{\kappa}(X_{\text{ét}}^{\text{aff}}) \rightarrow X_{\text{proét}}^{\text{aff}} : \text{“}\varprojlim\text{” } Y_i \mapsto \varprojlim_i Y_i$$

*is an equivalence of categories.*

Here,  $\text{Pro}_{\kappa}$  denotes the category of pro-systems whose index category is bounded by  $\kappa$ .

*Proof.* By definition, the functor is essentially surjective, so we need to see that it is fully faithful. For this, write  $X = \text{Spa}(R, R^+)$ , and take  $Y = \text{Spa}(S, S^+)$ ,  $Z = \text{Spa}(T, T^+)$  two objects of  $X_{\text{proét}}^{\text{aff}}$  that are written as limits of  $Y_i = \text{Spa}(S_i, S_i^+) \in X_{\text{ét}}^{\text{aff}}$ ,  $Z_j = \text{Spa}(T_j, T_j^+) \in X_{\text{ét}}^{\text{aff}}$ , for cofiltered index categories  $I, J$ . We need to show that

$$\text{Hom}_X(Y, Z) = \varprojlim_j \varinjlim_i \text{Hom}_X(Y_i, Z_j) .$$

Without loss of generality  $J$  is a singleton, so  $Z \rightarrow X$  is étale. Now have to check that

$$\mathrm{Hom}_X(Y, Z) = \varinjlim_i \mathrm{Hom}_X(Y_i, Z) .$$

Recall that  $Y_{\acute{e}t, \mathrm{qcqs}} = 2\text{-}\varinjlim_i (Y_i)_{\acute{e}t, \mathrm{qcqs}}$  by Proposition 6.4 (ii). Thus,

$$\mathrm{Hom}_X(Y, Z) = \mathrm{Hom}_Y(Y, Y \times_X Z) = \varinjlim_i \mathrm{Hom}_{Y_i}(Y_i, Y_i \times_X Z) = \varinjlim_i \mathrm{Hom}_X(Y_i, Z) ,$$

as desired.  $\square$

We need some basic properties of pro-étale maps.

**Lemma 7.11.**

- (i) *Let  $g : Z \rightarrow Y$  and  $f : Y \rightarrow X$  be pro-étale (resp. affinoid pro-étale) morphisms. Then  $f \circ g : Z \rightarrow X$  is pro-étale (resp. affinoid pro-étale).*
- (ii) *Let  $f : Y \rightarrow X$  be pro-étale (resp. affinoid pro-étale), and let  $g : X' \rightarrow X$  be a map from a perfectoid space (resp. affinoid perfectoid space). Then  $f' : Y' = X' \times_X Y \rightarrow X'$  is pro-étale (resp. affinoid pro-étale).*
- (iii) *Let  $f : Y \rightarrow X$  and  $f' : Y' \rightarrow X$  be pro-étale (resp. affinoid pro-étale). Then any map  $g : Y \rightarrow Y'$  over  $X$  is pro-étale (resp. affinoid pro-étale).*
- (iv) *For any affinoid perfectoid space  $X$ , the category  $X_{\mathrm{pro\acute{e}t}}^{\mathrm{aff}}$  has all  $\kappa$ -small limits.*

*Proof.* It is enough to prove the assertions in the affinoid pro-étale case. For part (i), we use Proposition 7.10, Proposition 6.4 (iv) and the natural functor  $\mathrm{Pro}_\kappa(\mathrm{Pro}_\kappa(\mathcal{C})) \rightarrow \mathrm{Pro}_\kappa(\mathcal{C})$ , which exists for any category  $\mathcal{C}$ . Part (ii) follows directly from the definition. For part (iii), we can factor  $g$  as the composite of a section of the affinoid pro-étale map  $Y \times_X Y' \rightarrow Y$  (given by the graph), and the affinoid pro-étale map  $Y \times_X Y' \rightarrow Y'$ ; thus, it is enough to show that sections of affinoid pro-étale maps are affinoid pro-étale. But if  $Y = \varinjlim_i Y_i \rightarrow X$  is a pro-étale presentation, and  $s : X \rightarrow Y$  is a section, then  $s$  is given by a compatible system of sections  $s_i : X \rightarrow Y_i$ , and  $X = \varinjlim_i (X \times_{Y_i} Y) \rightarrow Y$  is a pro-étale presentation.

Finally, for part (iv), as  $X_{\mathrm{pro\acute{e}t}}^{\mathrm{aff}} = \mathrm{Pro}_\kappa(X_{\acute{e}t}^{\mathrm{aff}})$  by Proposition 7.10, it is enough to prove that  $X_{\acute{e}t}^{\mathrm{aff}}$  has all finite limits. For this, it is enough to check that it has a final object and fibre products, which is clear.  $\square$

**Proposition 7.12.** *The inclusion from the category of  $w$ -local perfectoid spaces with  $w$ -local maps, to the category of qcqs perfectoid spaces, admits a left adjoint  $X \mapsto X^{\mathrm{wl}}$ . The adjunction map  $X^{\mathrm{wl}} \rightarrow X$  is pro-étale. If  $X$  is affinoid, the map  $X^{\mathrm{wl}} \rightarrow X$  is an inverse limit of surjective maps of the form  $\bigsqcup_{i \in I} U_i \rightarrow X$ , where  $I$  is a finite set, and  $U_i \subset X$  are rational subsets; in particular, it is affinoid pro-étale.*

*Proof.* By [BS15, Lemma 2.1.10], there is a similar  $w$ -localization  $|X|^{\mathrm{wl}}$  of the underlying topological space  $|X|$ . Consider the topological space  $|X|^{\mathrm{wl}}$  endowed with the presheaves  $\mathcal{O}_{X^{\mathrm{wl}}}^+$ ,  $\mathcal{O}_{X^{\mathrm{wl}}}$  of complete topological rings given by  $\mathcal{O}_{X^{\mathrm{wl}}}^+ = (p^* \mathcal{O}_X^+)_{\varpi}^{\wedge}$ ,  $\mathcal{O}_{X^{\mathrm{wl}}} = \mathcal{O}_{X^{\mathrm{wl}}}^+[\varpi^{-1}]$ , where  $p : |X|^{\mathrm{wl}} \rightarrow |X|$  denotes the projection, and  $\varpi$  is a (local choice of) pseudouniformizer. We claim that this defines a perfectoid space  $X^{\mathrm{wl}}$ ; it is then clear that it is final in the category of  $w$ -local perfectoid spaces over  $X$  (with  $w$ -local maps).

We handle the affinoid case first. We use the following general presentation of the  $w$ -localization.



**Lemma 7.13.** *Let  $X$  be a spectral topological space, and let  $B$  be a basis for the topology consisting of quasicompact open subsets, such that for  $U, V \in B$ , also  $U \cap V \in B$ . Let  $X^{\text{wl}} \rightarrow X$  be the  $w$ -localization of  $X$ , and let  $C$  be the category of factorizations  $X^{\text{wl}} \rightarrow T \rightarrow X$ , where  $T$  is a finite disjoint union of elements of  $B$ . Then  $C$  is cofiltered, and the natural map*

$$f : X^{\text{wl}} \rightarrow \varprojlim_C T$$

*is a homeomorphism.*

*Proof.* Note that the category of  $C$  is cofiltered: Given any two factorizations  $X^{\text{wl}} \rightarrow T_1 \rightarrow X$  and  $X^{\text{wl}} \rightarrow T_2 \rightarrow X$ , one gets another one  $X^{\text{wl}} \rightarrow T_1 \times_X T_2 \rightarrow X$ , and for any two given maps  $f, g : T_1 \rightarrow T_2$  between two such factorizations, the locus  $T \subset T_1$  where  $f = g$  is open and closed (more precisely, if  $T_1 = \bigsqcup_{i \in I} U_i$ , then  $T = \bigsqcup_{i \in J} U_i$  for some subset  $J \subset I$ ), and  $X^{\text{wl}} \rightarrow T_1$  factors over  $T$ . In particular, we can assume that  $X \in B$ , as the two resulting categories are cofinal.

By Lemma 2.5, it is enough to see that  $f : X^{\text{wl}} \rightarrow \varprojlim_C T$  is bijective and generalizing. Recall that  $\pi_0(X^{\text{wl}}) = X^{\text{cons}}$ , where  $X^{\text{cons}}$  is  $X$  endowed with the constructible topology, and the connected component corresponding to  $x \in X$  is given by the localization  $X_x$ , i.e. the set of points specializing to  $x$ . In particular, points of  $X^{\text{wl}}$  are in bijection with pairs  $(x, y)$  of points in  $X$ , where  $y \in X_x$ . The map  $X^{\text{wl}} \rightarrow X$  is given by  $(x, y) \mapsto y$ , which is generalizing; more precisely, for any  $(x, y) \in X^{\text{wl}}$ , the localization  $X_{(x,y)}^{\text{wl}}$  is identified with the localization  $X_y$ . As  $T \rightarrow X$  is a finite disjoint union of open subsets, it also identifies localizations, and thus the map  $f : X^{\text{wl}} \rightarrow \varprojlim_C T$  identifies localizations, and in particular is generalizing.

For injectivity of  $f : X^{\text{wl}} \rightarrow \varprojlim_C T$ , it is enough to show that given points  $(x, y), (x', y') \in X^{\text{wl}}$  with  $x \neq x'$ ,  $f(x, y) \neq f(x', y')$  (as the  $y$ -coordinate is seen by the projection to  $X$ ). For this, choose some  $U \in B$  such that  $U$  contains exactly one of  $x$  and  $x'$ , say  $x$ , and look at  $T = U \sqcup X \rightarrow X$ . We have a factorization

$$X^{\text{wl}} = X^{\text{wl}} \times_{X^{\text{cons}}} U^{\text{cons}} \sqcup X^{\text{wl}} \times_{X^{\text{cons}}} (X \setminus U)^{\text{cons}} \rightarrow U \sqcup X = T \rightarrow X$$

which sends  $(x, y)$  into  $U \subset T$ , and  $(x', y')$  into  $X \subset T$ . Thus, we get injectivity.

For surjectivity, we claim that it is enough to show that if  $X^{\text{wl}} \rightarrow T \rightarrow X$  is a factorization, and  $t \in T$  is a point that is not in the image of  $X^{\text{wl}} \rightarrow T$ , then one can find another factorization  $X^{\text{wl}} \rightarrow \tilde{T} \rightarrow T \rightarrow X$  such that  $t$  is not in the image of  $\tilde{T} \rightarrow T$ . Indeed, this implies that the image of  $X^{\text{wl}} \rightarrow T$  agrees with the image  $T_0 \subset T$  of  $\varprojlim_C T \rightarrow T$ . But then  $\varprojlim_C T_0 = \varprojlim_C T$ , and we have a family of surjective maps  $X^{\text{wl}} \rightarrow T_0$ ; by a usual quasicompactness argument, this implies that  $X^{\text{wl}} \rightarrow \varprojlim_C T_0 = \varprojlim_C T$  is surjective (the fibre over any point is a cofiltered limit of nonempty spectral spaces along spectral maps, thus nonempty by Tychonoff applied to the constructible topology).

Thus, assume  $t \in T$  is not in the image of  $X^{\text{wl}} \rightarrow T = \bigsqcup_{i \in I} U_i$ . Assume  $t \in U_{i_0}$ , and choose finitely many  $V_j \subset U_{i_0}$ ,  $j \in J$ ,  $V_j \in B$  with  $t \notin V_j$ , such that  $X^{\text{wl}} \rightarrow T$  factors over

$$\bigcup_{j \in J} V_j \sqcup \bigsqcup_{i \in I, i \neq i_0} U_i .$$

This is possible, as the map  $X^{\text{wl}} \rightarrow T$  is generalizing, so if  $t$  is not in the image, the closure of  $t$  is not in the image as well. We have an open cover

$$\tilde{T} := \bigsqcup_{j \in J} V_j \sqcup \bigsqcup_{i \in I, i \neq i_0} U_i \rightarrow \bigcup_{j \in J} V_j \sqcup \bigsqcup_{i \in I, i \neq i_0} U_i .$$

After pullback to  $X^{\text{wl}}$ , this open cover splits by the definition of w-local spaces. In particular, the map

$$X^{\text{wl}} \rightarrow \bigcup_{j \in J} V_j \sqcup \bigsqcup_{i \in I, i \neq i_0} U_i$$

lifts to a map  $X^{\text{wl}} \rightarrow \tilde{T}$ , giving the desired factorization  $X^{\text{wl}} \rightarrow \tilde{T} \rightarrow T \rightarrow X$ , with  $t$  not in the image of  $\tilde{T} \rightarrow T$ .  $\square$

We apply this in the case where  $B$  is the basis of rational subsets of  $X$ . Clearly, any  $T$  as in the lemma admits a canonical structure as an affinoid perfectoid space, for which  $T \rightarrow X$  is étale. Then  $\varprojlim_C T = X^{\text{wl}}$ , on topological spaces by Lemma 7.13, and on  $\mathcal{O}$  and  $\mathcal{O}^+$  by a direct verification; this finishes the argument.

In general, we can reduce to the affinoid case by the following lemma.

**Lemma 7.14.** *Let  $X$  be a spectral topological space, and  $U \subset X$  a quasicompact open subset. Then  $U^{\text{wl}} \rightarrow X^{\text{wl}}$  is a quasicompact open embedding. If  $X = \cup_{i \in I} U_i$  is a covering of  $X$  by quasicompact open subsets  $U_i \subset X$ , then the induced map  $\sqcup U_i^{\text{wl}} \rightarrow X^{\text{wl}}$  is an open covering by quasicompact open subsets.*

*Proof.* In general  $X^{\text{wl}}$  comes with functorial maps to  $X$  (by adjunction) and  $X^{\text{cons}}$ , which is  $X$  endowed with the constructible topology, via the identification  $\pi_0(X^{\text{wl}}) = X^{\text{cons}}$  from [BS15, Lemma 2.1.10]. In particular, there is a natural map

$$U^{\text{wl}} \rightarrow X^{\text{wl}} \times_{X^{\text{cons}}} U^{\text{cons}} \rightarrow X^{\text{wl}} .$$

Here,  $U^{\text{cons}} \subset X^{\text{cons}}$  is a quasicompact open subset; thus, it suffices to see that the map  $U^{\text{wl}} \rightarrow X^{\text{wl}} \times_{X^{\text{cons}}} U^{\text{cons}}$  is a homeomorphism. By Lemma 2.5, it is enough to show that it is bijective and generalizing. Recall that  $\pi_0(X^{\text{wl}}) = X$ , and for  $x \in X$ , the corresponding connected component of  $X^{\text{wl}}$  is the set  $X_x \subset X$  of all generalizations of  $x$ , i.e. the localization of  $X$  at  $x$ . If  $x \in U$ , it follows that  $U_x = X_x$ ; this shows bijectivity. Moreover, it shows that the map is generalizing, as desired.

The final statement follows from  $X^{\text{cons}} = \cup_{i \in I} U_i^{\text{cons}}$ .  $\square$

$\square$

Moreover, we have the following variant.

**Definition 7.15.** *A perfectoid space  $X$  is strictly totally disconnected if it is qcqs and every étale cover splits.*

**Proposition 7.16.** *Let  $X$  be a qcqs perfectoid space. Then  $X$  is strictly totally disconnected if and only if every connected component is of the form  $\text{Spa}(C, C^+)$ , where  $C$  is algebraically closed.*

*Proof.* Assume first that  $X$  is strictly totally disconnected. This condition passes to closed subsets, so we can assume that  $X$  is connected. In that case, as  $X$  is in particular totally disconnected,  $X = \text{Spa}(K, K^+)$  for some perfectoid field  $K$  with an open and bounded valuation subring  $K^+ \subset K$ . If  $K$  is not algebraically closed, then for any finite extension  $L$  of  $K$  with integral closure  $L^+ \subset L$

of  $K^+$ , the map  $\mathrm{Spa}(L, L^+) \rightarrow \mathrm{Spa}(K, K^+)$  is a nonsplit finite étale cover. Thus, necessarily  $K$  is algebraically closed.

Conversely, assume that all connected components of  $X$  are of the form  $\mathrm{Spa}(C, C^+)$ . By Lemma 7.3,  $X$  is totally disconnected. To see that it is strictly totally disconnected, it is enough to see that any étale cover is locally split (as then the étale cover admits a refinement by an open cover, which is split as  $X$  is totally disconnected). We may assume that the étale cover is qcqs. But any étale cover of any connected component  $\mathrm{Spa}(C, C^+)$  splits, and then the splitting extends to a small neighborhood by Proposition 6.4.  $\square$

Again, one could consider the following strengthening.

**Definition 7.17.** *A  $w$ -strictly local perfectoid space is a  $w$ -local perfectoid space  $X$  such that for all  $x \in X$ , the completed residue field  $K(x)$  is algebraically closed.*

In particular, a  $w$ -strictly local space is strictly totally disconnected.

For any  $X$ , one can find strictly totally disconnected spaces over  $X$ , as the following lemma shows. The construction of this lemma is slightly different than the  $w$ -localization functor, and in particular also ensures that  $\tilde{X} \rightarrow X$  is universally open, which is not true for  $X^{\mathrm{wl}} \rightarrow X$ .

**Lemma 7.18.** *Let  $X$  be an affinoid perfectoid space. Then there is an affinoid perfectoid space  $\tilde{X}$  with an affinoid pro-étale surjective and universally open map  $\tilde{X} \rightarrow X$ , such that  $\tilde{X}$  is strictly totally disconnected.*

As usual, per Convention 4.2,  $X$  is assumed to be  $\kappa$ -small, and we require  $\tilde{X}$  (and the implicit pro-system) to be  $\kappa$ -small as well.

*Proof.* We want to take the cofiltered limit of “all” affinoid étale covers over  $X$ . For this, fix a set of representatives  $\{X_i \rightarrow X\}_{i \in I}$  of all affinoid étale and surjective spaces over  $X$ . We note that this is of cardinality less than  $\kappa$ : The cardinality of  $\mathcal{O}_{X_i}(X_i)$  is bounded independently of  $i$ , say by  $\lambda$  (which is essentially the cardinality of  $\mathcal{O}_X(X)$ ). The set of all affinoid perfectoid spaces  $Y$  for which  $\mathcal{O}_Y(Y)$  has cardinality bounded by  $\lambda$  is also of cardinality less than  $\kappa$  (by thinking about all possible ways of endowing a set of cardinality  $\leq \lambda$  with the structure of a perfectoid algebra, and possible rings of integral elements). For each such  $Y$ , the set of maps from  $Y$  to  $X$  has cardinality less than  $\kappa$ . In total, one sees that there are less than  $\kappa$  such étale maps  $Y \rightarrow X$ . For each finite subset  $J \subset I$ , let  $X_J$  be the product of all  $X_i$ ,  $i \in J$ , over  $X$ , which is still affinoid. Then  $X_J \rightarrow X$  is étale and surjective, and the transition maps  $X_J \rightarrow X_{J'}$  for  $J' \subset J$  are also étale and surjective. In particular, if we set  $X_\infty = \varprojlim_J X_J$ , then  $X_\infty \rightarrow X$  is affinoid pro-étale, and universally open. Indeed, it is open as any quasicompact open subset  $U_\infty \subset X_\infty$  comes as the preimage of some quasicompact open  $U_J \subset X_J$ , and then  $U_J$  is also equal to the image of  $U_\infty$  in  $X_J$ , as  $X_\infty \rightarrow X_J$  is surjective. Thus, the image of  $U_\infty$  in  $X$  agrees with the image of  $U_J$  in  $X$ , which is open, as étale maps are open. The same argument applies after any base change, so  $X_\infty \rightarrow X$  is universally open. Repeating the construction  $X \mapsto X_\infty$  countably often (here, we use our assumption that countable unions of cardinals less than  $\kappa$  are less than  $\kappa$ ), we can find an affinoid pro-étale  $\tilde{X} \rightarrow X$  such that all étale covers of  $\tilde{X}$  split. Indeed, any étale cover can be refined by an affinoid étale cover. Such a cover comes from some finite level by Proposition 6.4 (iv), and becomes split at the next.  $\square$

In the following, we will analyze perfectoid spaces which are pro-étale over a strictly totally disconnected space.

**Lemma 7.19.** *Let  $X$  be a strictly totally disconnected perfectoid space. Let  $f : Y \rightarrow X$  be a quasicompact separated map of perfectoid spaces. Then  $f$  is pro-étale if and only if for all rank-1-points  $x = \mathrm{Spa}(C, \mathcal{O}_C) \in X$ , the fibre  $Y_x = Y \times_X x$  is isomorphic to  $x \times \underline{S}_x$  for some profinite set  $S_x$ . In this case,  $Y$  is strictly totally disconnected, and  $Y \rightarrow X$  is affinoid pro-étale.*

Note that the lemma applies in particular if  $Y$  is affinoid, as any map of affinoid perfectoid spaces is separated.

*Proof.* The condition is clearly necessary. Thus assume that  $f : Y \rightarrow X$  has the property that for all rank-1-points  $x = \mathrm{Spa}(C, \mathcal{O}_C) \in X$ , the fibre  $Y_x = Y \times_X x$  is isomorphic to  $x \times \underline{S}_x$  for some profinite set  $S_x$ . We want to show that  $Y$  can be written as an inverse limit  $Y = \varprojlim_i Y_i \rightarrow X$  of étale maps  $Y_i \rightarrow X$ , where all  $Y_i$  are affinoid.

First, note that  $f : Y \rightarrow X$  factors as  $Y \rightarrow X \times_{\pi_0(X)} \pi_0(Y) \rightarrow X$ , where  $X \times_{\pi_0(X)} \pi_0(Y) \rightarrow X$  is affinoid pro-étale, and  $X \times_{\pi_0(X)} \pi_0(Y)$  is strictly totally disconnected. Thus, replacing  $X$  by  $X \times_{\pi_0(X)} \pi_0(Y)$ , we can reduce to the case that  $\pi_0(f) : \pi_0(Y) \rightarrow \pi_0(X)$  is a homeomorphism.

We claim that in this situation,  $f$  is an injection. To check this, we can check on connected components, so we may assume that  $X = \mathrm{Spa}(C, C^+)$ , and then by assumption also  $Y$  is connected. To check that  $f$  is injective, it is by Proposition 5.3 enough to check injectivity on  $(K, K^+)$ -valued points, and then as  $f$  is separated, it is enough to check on  $(K, \mathcal{O}_K)$ -valued points. Thus, assume two rank-1-points  $y_1, y_2 \in Y$  map to the same point  $x \in X$ . Now  $x$  is the unique rank-1-point of  $X$ , and the set of preimages of  $x$  forms a profinite set  $Y_x \subset Y$  by assumption. By our assumption,  $Y_x$  contains at least two points, so we can find a closed and open decomposition  $Y_x = U_{1,x} \sqcup U_{2,x}$ , for some quasicompact open subsets  $U_1, U_2 \subset Y$ . Let  $V_1, V_2 \subset Y$  be the closures of  $U_{1,x}$  and  $U_{2,x}$ . As  $U_{1,x}$  and  $U_{2,x}$  are pro-constructible subsets, their closures are precisely the subsets of specializations of points in  $U_{1,x}$  resp.  $U_{2,x}$ . As any point of  $Y$  generalizes to a unique point of  $Y_x$ ,  $Y = V_1 \sqcup V_2$ . As  $V_1$  and  $V_2$  are both closed, this gives a contradiction to our assumption that  $Y$  is connected, finishing the proof that  $f$  is an injection.

Now, since  $|f| : |Y| \rightarrow |X|$  is injective, the image of  $|f|$  is a pro-constructible generalizing subset of  $X$ , and thus by Lemma 7.6 an intersection of subsets of the form  $\{|g| \leq 1\}$  for varying  $g \in H^0(X, \mathcal{O}_X)$ . As such, the image is affinoid pro-étale in  $X$ , and itself (strictly) totally disconnected. Thus, replacing  $X$  by the image of  $f$ , we can assume that  $|f|$  is a bijection. But then  $f$  is an isomorphism by Lemma 5.4.  $\square$

Lemma 5.4 allows us to give a purely topological description of perfectoid spaces which are pro-étale over a strictly totally disconnected space.

**Definition 7.20.** *Let  $T$  be a spectral space such that every connected component is a totally ordered chain of specializations. A spectral map  $S \rightarrow T$  of spectral spaces is called affinoid pro-étale if the induced map  $S \rightarrow T \times_{\pi_0(T)} \pi_0(S)$  is a pro-constructible and generalizing embedding. A spectral map  $S \rightarrow T$  from a locally spectral space  $S$  is called pro-étale if  $S$  is covered by spectral subsets  $S' \subset S$  for which the restriction  $S' \rightarrow T$  is affinoid pro-étale.*

**Remark 7.21.** If  $S \rightarrow T$  is pro-étale and  $S$  is spectral, it is not automatic that  $S \rightarrow T$  is affinoid pro-étale; the analogue of the separatedness condition can fail.

**Corollary 7.22.** *Let  $X$  be a strictly totally disconnected perfectoid space. Then the functor sending a map  $f : Y \rightarrow X$  to  $|f| : |Y| \rightarrow |X|$  induces equivalences between:*

- (i) the category of ( $\kappa$ -small) affinoid pro-étale perfectoid spaces  $Y \rightarrow X$ , and the category of affinoid pro-étale  $S \rightarrow |X|$  (of cardinality less than  $\kappa$ );
- (ii) the category of ( $\kappa$ -small) pro-étale perfectoid spaces  $Y \rightarrow X$ , and the category of pro-étale  $S \rightarrow |X|$  (of cardinality less than  $\kappa$ ).

*Proof.* Lemma 7.19 implies that if  $f : Y \rightarrow X$  is (affinoid) pro-étale, then  $|f| : |Y| \rightarrow |X|$  is (affinoid) pro-étale. We construct an inverse functor. For this, given a pro-étale map  $g : S \rightarrow |X|$ , set  $\mathcal{O}_S^+ := (g^* \mathcal{O}_X^+)_{\varpi}^{\wedge}$  and  $\mathcal{O}_S = \mathcal{O}_S^+[\varpi^{-1}]$ , where  $\varpi$  is a pseudouniformizer on the affinoid space  $X$ . For any point of  $S$ , one gets an induced valuation on  $\mathcal{O}_S$  by pullback from  $X$ . One checks using Lemma 7.6 that if  $g : S \rightarrow |X|$  is affinoid pro-étale, then this defines an affinoid perfectoid space  $Y$ ; Lemma 7.19 implies that the corresponding map  $f : Y \rightarrow X$  is affinoid pro-étale. In general,  $S \rightarrow |X|$  is locally affinoid pro-étale, so again one gets a perfectoid space  $Y$ , pro-étale over  $X$ .  $\square$

Another important property of totally disconnected spaces is an automatic flatness assertion.

**Proposition 7.23.** *Let  $X = \mathrm{Spa}(R, R^+)$  be a totally disconnected perfectoid space, and  $f : Y = \mathrm{Spa}(S, S^+) \rightarrow X$  any map from an affinoid perfectoid space  $Y$ . Then  $S^+/\varpi$  is flat over  $R^+/\varpi$ , for any pseudouniformizer  $\varpi \in R$ .*

*Moreover, if  $|f| : |Y| \rightarrow |X|$  is surjective, then the map is faithfully flat.*

*Proof.* Let  $g : |X| \rightarrow \pi_0(X)$  be the projection. First, we check that  $((g \circ f)_* \mathcal{O}_Y^+)/\varpi$  is flat over  $(g_* \mathcal{O}_X^+)/\varpi$ , as sheaves on  $\pi_0(X)$ . This can be checked on stalks. But the stalk of  $(g_* \mathcal{O}_X^+)/\varpi$  at  $c \in \pi_0(X)$  is given by  $K_c^+/\varpi$ , where  $\mathrm{Spa}(K_c, K_c^+) \subset X$  is the connected component corresponding to  $c$ . Similarly, the stalk of  $((g \circ f)_* \mathcal{O}_Y^+)/\varpi$  is given by  $S_c^+/\varpi$ , where  $\mathrm{Spa}(S_c, S_c^+) = Y \times_X \mathrm{Spa}(K_c, K_c^+)$ . But  $S_c^+$  is  $\varpi$ -torsionfree, which implies that it is flat over  $K_c^+$ . By base change,  $S_c^+/\varpi$  is flat over  $K_c^+/\varpi$ , as desired.

As flatness can be checked locally, it follows that  $S^+/\varpi$  is flat over  $R^+/\varpi$ . More precisely, flatness can be checked on connected components of  $\mathrm{Spec}(R^+/\varpi)$ , but  $\pi_0(\mathrm{Spec}(R^+/\varpi)) = \pi_0(X)$ , and on each connected component, we have verified the desired flatness.

To check faithful flatness in case  $|f|$  is surjective, we have to see that  $\mathrm{Spec}(S^+/\varpi) \rightarrow \mathrm{Spec}(R^+/\varpi)$  is surjective. This can be checked on connected components, so we can assume that  $X = \mathrm{Spa}(K, K^+)$  for some perfectoid field  $K$  with an open and bounded valuation subring  $K^+ \subset K$ . In that case, as  $|f|$  is surjective, we can find a map  $\mathrm{Spa}(L, L^+) \rightarrow Y$  with  $L$  a perfectoid field with an open and bounded valuation subring  $L^+ \subset L$ , such that  $\mathrm{Spa}(L, L^+) \rightarrow \mathrm{Spa}(K, K^+)$  is surjective. But one can identify  $\mathrm{Spa}(L, L^+) = \mathrm{Spec}(L^+/\varpi)$  and  $\mathrm{Spa}(K, K^+) = \mathrm{Spec}(K^+/\varpi)$ , so  $\mathrm{Spec}(L^+/\varpi) \rightarrow \mathrm{Spec}(S^+/\varpi) \rightarrow \mathrm{Spec}(K^+/\varpi)$  is surjective, as desired.  $\square$

## 8. THE PRO-ÉTALE AND V-TOPOLOGY

In this section, we define two Grothendieck topologies on the category of perfectoid spaces.

**Definition 8.1.** *Let  $\mathrm{Perfd}$  be the category of  $\kappa$ -small perfectoid spaces.*

- (i) *The big pro-étale site is the Grothendieck topology on  $\mathrm{Perfd}$  for which a collection  $\{f_i : Y_i \rightarrow X\}_{i \in I}$  of morphisms is a covering if all  $f_i$  are pro-étale, and for each quasicompact open subset  $U \subset X$ , there exists a finite subset  $J \subset I$  and quasicompact open subsets  $V_i \subset Y_i$  for  $i \in J$ , such that  $U = \bigcup_{i \in J} f_i(V_i)$ .*

- (ii) Let  $X$  be a perfectoid space. The small pro-étale site of  $X$  is the Grothendieck topology on the category of perfectoid spaces  $f : Y \rightarrow X$  pro-étale over  $X$ , with covers the same as in the big pro-étale site.
- (iii) The  $v$ -site is the Grothendieck topology on  $\text{Perfd}$  for which a collection  $\{f_i : Y_i \rightarrow X\}_{i \in I}$  of morphisms is a covering if for each quasicompact open subset  $U \subset X$ , there exists a finite subset  $J \subset I$  and quasicompact open subsets  $V_i \subset Y_i$  for  $i \in J$ , such that  $U = \bigcup_{i \in J} f_i(V_i)$ .

We note that the slightly technical condition on quasicompact subsets is the same condition that appears in the definition of the fpqc topology for schemes. Also, there is no “small”  $v$ -site of a perfectoid space  $X$ , as this would by design include all perfectoid spaces over  $X$ .

Our goal in this section is prove that both the pro-étale and the  $v$ -site are well-behaved. We start with some general remarks about quasicompact and quasiseparated objects. Let us quickly recall their definition.

**Definition 8.2.** *Let  $T$  be a topos.*

- (i) An object  $X \in T$  is quasicompact if for any collection of objects  $X_i \in T$ ,  $i \in I$ , with maps  $f_i : X_i \rightarrow X$  such that  $\bigsqcup f_i : \bigsqcup X_i \rightarrow X$  is surjective, there is a finite subset  $J \subset I$  such that  $\bigsqcup_{i \in J} X_i \rightarrow X$  is surjective, cf. SGA 4 VI Définition 1.1.
- (ii) An object  $X \in T$  is quasiseparated if for all quasicompact objects  $Y, Z \in T$ , the fibre product  $Y \times_X Z \in T$  is quasicompact, cf. SGA 4 VI Définition 1.13.
- (iii) A map  $f : Y \rightarrow X$  is quasicompact if for all quasicompact  $Z \in T$ , the fibre product  $Z \times_X Y$  is quasicompact, cf. SGA 4 VI Définition 1.7.
- (iv) A map  $f : Y \rightarrow X$  is quasiseparated if  $\Delta_f : Y \rightarrow Y \times_X Y$  is quasicompact, cf. SGA 4 VI Définition 1.7.

The combination of quasicompact and quasiseparated is abbreviated to “qcqs”.

In the following, we assume that  $T$  is an *algebraic* topos, cf. SGA 4 VI Définition 2.3. This means that there is a generating full subcategory  $C \subset T$  consisting of qcqs objects that is stable under fibre products, and moreover for  $X \in C$ , the map  $X \rightarrow *$  is quasiseparated. This assumption will be satisfied in all our examples. Under this assumption, it suffices to check these various conditions after pullback to objects of  $C$ ; also, it suffices to check after one cover by objects of  $C$ . Thus, for example  $X \in T$  is quasiseparated if and only if for one cover of  $X$  by objects  $X_i \in C$ , all  $X_i \times_X X_j$  are quasicompact, cf. SGA 4 VI Corollaire 1.17. Also,  $f : Y \rightarrow X$  is quasiseparated if and only if for all  $Z \in C$  with a map  $Z \rightarrow X$ , the object  $Y \times_X Z$  is quasiseparated, cf. SGA 4 VI Corollaire 2.6. Another equivalent characterization is that for all quasiseparated  $Z \in T$  with a map  $Z \rightarrow X$ , the object  $Y \times_X Z$  is quasiseparated, cf. SGA 4 VI Corollaire 2.8.

We warn the reader that if the final object of  $T$  is not quasiseparated (as in the case for (sheaves on)  $\text{Perfd}$ ), it is not equivalent to say that  $X$  is quasicompact (or quasiseparated) and to say that the map  $X \rightarrow *$  is quasicompact (or quasiseparated). It is true that  $X$  quasiseparated implies  $X \rightarrow *$  is quasiseparated, and that  $X \rightarrow *$  being quasicompact implies that  $X$  is quasicompact. Indeed, the former follows from our assumption that  $T$  is algebraic by SGA 4 VI Proposition 2.2, and the latter follows from SGA 4 VI Proposition 1.3, by choosing any quasicompact object  $Y \in T$ , and the corresponding cover  $X \times Y \rightarrow X$ , where  $X \times Y$  is quasicompact if  $X \rightarrow *$  is.

**Proposition 8.3.** *The topoi of sheaves on  $\text{Perfd}$  for either the (big) pro-étale or  $v$ -topology, and the topos of sheaves on  $X_{\text{proét}}$  for a perfectoid space  $X$ , are algebraic. A basis of qcqs objects stable*

under fibre products is in all cases given by affinoid perfectoid spaces. Moreover, a perfectoid space  $X$  is quasicompact resp. quasiseparated in any of these settings if and only if  $|X|$  is quasicompact resp. quasiseparated.

*Proof.* Left to the reader.  $\square$

**Convention 8.4.** Further below, we will sometimes ask that a map  $f : Y' \rightarrow Y$  of stacks on some topos  $T$  is quasiseparated. Following the convention of [Sta, Tag 04YW], we define this to mean that  $\Delta_f : Y' \rightarrow Y' \times_Y Y'$  is quasicompact and quasiseparated, where the latter condition is not automatic for stacks (as the diagonal need not be injective).

**Proposition 8.5.** *Let  $X$  be a perfectoid space, and  $\nu : X_{\text{proét}} \rightarrow X_{\text{ét}}$  the natural map of sites.*

- (i) *For any sheaf  $\mathcal{F}$  on  $X_{\text{ét}}$ , the natural adjunction map  $\mathcal{F} \rightarrow \nu_* \nu^* \mathcal{F}$  is an equivalence. If  $\mathcal{F}$  is a sheaf of abelian groups, then  $R^i \nu_* \nu^* \mathcal{F} = 0$  for all  $i \geq 1$ .*
- (ii) *Let  $Y = \varprojlim_i Y_i \rightarrow X_0 \subset X$  be a pro-étale presentation of an affinoid pro-étale map  $Y \rightarrow X_0$  to an affinoid open subset  $X_0 \subset X$ , and let  $\mathcal{F}$  be a sheaf on  $X_{\text{ét}}$ . The natural map*

$$\varinjlim_i \mathcal{F}(Y_i) \rightarrow (\nu^* \mathcal{F})(Y)$$

*is an isomorphism.*

- (iii) *The presheaves  $\mathcal{O}, \mathcal{O}^+$  on  $X_{\text{proét}}$  sending  $Y \in X_{\text{proét}}$  to  $\mathcal{O}_Y(Y)$  resp.  $\mathcal{O}_Y^+(Y)$ , are sheaves. If  $X$  is affinoid perfectoid, then  $H^i(X_{\text{proét}}, \mathcal{O}) = 0$  for  $i > 0$ , and  $H^i(X_{\text{proét}}, \mathcal{O}^+)$  is almost zero for  $i > 0$ .*

*Proof.* All assertions reduce to the case that  $X$  is affinoid. In this case, the sites  $X_{\text{proét}}^{\text{aff}}$  and  $X_{\text{proét}}$  define equivalent topoi, so we can work with  $X_{\text{proét}}^{\text{aff}}$  instead. Let  $\mathcal{F}$  be a sheaf on  $X_{\text{ét}}$ . It follows from Proposition 7.10 that  $\nu^* \mathcal{F}$  is the sheafification of the presheaf  $\nu^+ \mathcal{F}$  sending a pro-étale presentation  $Y = \varprojlim_i Y_i \rightarrow X$  to

$$(\nu^+ \mathcal{F})(Y) = \varinjlim_i \mathcal{F}(Y_i) ;$$

in fact, this presheaf is the pullback on the level of presheaves. To prove (ii), one argues as in [BS15, Lemma 5.1.1]. More precisely, we want to show that  $\nu^+ \mathcal{F}$  is a sheaf. Thus, for any surjective map  $\tilde{Y} \rightarrow Y$  in  $X_{\text{proét}}^{\text{aff}}$ , we need to see that

$$(\nu^+ \mathcal{F})(Y) \rightarrow \text{eq}((\nu^+ \mathcal{F})(\tilde{Y}) \rightrightarrows (\nu^+ \mathcal{F})(\tilde{Y} \times_Y \tilde{Y}))$$

is an isomorphism. But we can write  $\tilde{Y} = \varprojlim_i \tilde{Y}_i \rightarrow Y$  as a cofiltered limit of affinoid étale surjective  $\tilde{Y}_i \rightarrow Y$ , and

$$(\nu^+ \mathcal{F})(\tilde{Y}) = \varinjlim_i (\nu^+ \mathcal{F})(\tilde{Y}_i) ,$$

and similarly for  $(\nu^+ \mathcal{F})(\tilde{Y} \times_Y \tilde{Y})$ , which reduces us to the case that  $\tilde{Y} \rightarrow Y$  is étale. In that case, writing  $Y = \varprojlim_j Y_j$  as a cofiltered limit of affinoid étale  $Y_j \rightarrow X$ , the map  $\tilde{Y} \rightarrow Y$  comes as the pullback of some affinoid étale  $\tilde{Y}_j \rightarrow Y_j$  for  $j$  large enough by Proposition 6.4 (iv). Then

$$\text{eq}((\nu^+ \mathcal{F})(\tilde{Y}') \rightrightarrows (\nu^+ \mathcal{F})(\tilde{Y}' \times_Y \tilde{Y}')) = \varinjlim_j \text{eq}(\mathcal{F}(\tilde{Y}'_j) \rightrightarrows \mathcal{F}(\tilde{Y}'_j \times_{Y_j} \tilde{Y}'_j)) = \varinjlim_j \mathcal{F}(Y_j) = \mathcal{F}(Y) ,$$

as desired.

Part (i) is proved similarly, following [BS15, Corollary 5.1.6]. Finally, for part (iii), choose a pseudouniformizer  $\varpi \in \mathcal{O}_X^+(X)$ . Then by [Sch12, Proposition 7.13], we have a sheaf of almost  $\mathcal{O}_X^+(X)/\varpi$ -modules on  $X_{\text{ét}}^{\text{aff}}$ , sending  $Y \in X_{\text{ét}}^{\text{aff}}$  to  $(\mathcal{O}_Y^+(Y)/\varpi)^a$ . By part (ii), the pullback of this sheaf to the pro-étale site sends  $Y = \varprojlim_i Y_i \rightarrow X$  to

$$(\varinjlim_i \mathcal{O}_{Y_i}^+(Y_i)/\varpi)^a = (\mathcal{O}_Y^+(Y)/\varpi)^a ,$$

so sending  $Y \in X_{\text{proét}}^{\text{aff}}$  to  $(\mathcal{O}_Y^+(Y)/\varpi)^a$  is a sheaf of almost  $\mathcal{O}_X^+(X)/\varpi$ -modules. By induction, one gets sheaves  $(\mathcal{O}_Y^+(Y)/\varpi^n)^a$  for all  $n \geq 1$ , and then passing to the inverse limit, a sheaf  $Y \mapsto \mathcal{O}_Y^+(Y)^a$ ; inverting  $\varpi$  gives a sheaf  $\mathcal{O} : Y \mapsto \mathcal{O}_Y(Y)$ . All of these sheaves have vanishing higher cohomology groups on  $X$  by part (i) and [Sch12, Proposition 7.13]. Finally,  $\mathcal{O}^+ \subset \mathcal{O}$  is the subsheaf of those functions which are everywhere of absolute value  $\leq 1$ , and so is a sheaf as  $\mathcal{O}$  is.  $\square$

**Corollary 8.6.** *The presheaves  $\mathcal{O} : X \mapsto \mathcal{O}_X(X)$  and  $\mathcal{O}^+ : X \mapsto \mathcal{O}_X^+(X)$  on the big pro-étale site are sheaves. Moreover, the big pro-étale site is subcanonical, i.e. for every perfectoid space  $X$ , the functor  $Y \mapsto \text{Hom}(Y, X)$  is a sheaf for the big pro-étale site.*

*Proof.* As any cover  $\{f_i : Y_i \rightarrow X\}$  in the big pro-étale site can also be regarded as a cover in the small pro-étale site of  $X$ , these questions reduce to the small pro-étale site of  $X$ . Then the first part follows from Proposition 8.5 (iii). For the second part, assume that  $\{f_i : Y_i \rightarrow Y\}$  is a pro-étale cover of  $Y$ , and one has maps  $g_i : Y_i \rightarrow X$  which agree on overlaps  $Y_i \times_Y Y_j$ . Note first that by Lemma 2.5, the maps  $|Y_i| \rightarrow |X|$  glue to a continuous map  $|Y| \rightarrow |X|$ ; in particular, the problem can be considered locally on  $X$ , so we can assume that  $X = \text{Spa}(R, R^+)$  is affinoid. Then we are given maps  $(R, R^+) \rightarrow (\mathcal{O}(Y_i), \mathcal{O}^+(Y_i))$  which agree on overlaps; as  $\mathcal{O}$  and  $\mathcal{O}^+$  are sheaves for the pro-étale site, these maps glue to a map  $(R, R^+) \rightarrow (\mathcal{O}(Y), \mathcal{O}^+(Y))$ , which gives a map  $Y \rightarrow X$ , as desired.  $\square$

Next, we want to extend these results to the v-site. Here, the strategy is to combine pro-étale localization with the automatic faithful flatness results over totally disconnected bases, Proposition 7.23.

**Theorem 8.7.** *The presheaves  $\mathcal{O} : X \mapsto \mathcal{O}_X(X)$  and  $\mathcal{O}^+ : X \mapsto \mathcal{O}_X^+(X)$  on the v-site are sheaves. Moreover, the v-site is subcanonical.*

*Proof.* As in Corollary 8.6, it is enough to show that  $\mathcal{O}$  and  $\mathcal{O}^+$  are sheaves. As  $\mathcal{O}^+ \subset \mathcal{O}$  is the subsheaf of functions which are everywhere of absolute value  $\leq 1$  (which can be checked after v-covers), it is enough to prove that  $\mathcal{O}$  is a sheaf. As for every perfectoid space  $X$ ,  $\mathcal{O}(X)$  injects into  $\prod_{x \in X} K(x)$ , it is clear that  $\mathcal{O}$  is separated. It remains to see that given a v-cover  $\{f_i : Y_i \rightarrow X\}$  of a perfectoid space and functions  $g_i \in \mathcal{O}(Y_i)$  which agree on overlaps  $Y_i \times_X Y_j$ , they glue uniquely to a function  $g \in \mathcal{O}(X)$ . This can be checked locally on  $X$ , so we can assume that  $X$  is affinoid. In that case, the quasicompactness assumptions on a v-cover ensure that we can refine the given cover into a single  $f : Y \rightarrow X$ , where  $Y$  is affinoid. Let  $\tilde{X} \rightarrow X$  be a totally disconnected cover of  $X$  along an affinoid pro-étale map (for example the w-localization). We can also replace  $f$  by the composite  $Y \times_X \tilde{X} \rightarrow \tilde{X} \rightarrow X$ . By Corollary 8.6, we have descent for  $\tilde{X} \rightarrow X$ , so it is enough to handle the map  $Y \times_X \tilde{X} \rightarrow \tilde{X}$ ; in other words, we can assume that  $X$  is totally disconnected, so let  $f : Y = \text{Spa}(S, S^+) \rightarrow X = \text{Spa}(R, R^+)$  be a map of affinoid perfectoid spaces, where  $X$  is totally disconnected.



But now  $S^+/\varpi$  is flat over  $R^+/\varpi$  by Proposition 7.23. In fact, as  $f$  is surjective, it is faithfully flat, so  $R^+/\varpi$  is the equalizer of the two maps from  $S^+/\varpi$  to  $S^+/\varpi \otimes_{R^+/\varpi} S^+/\varpi$ . But if  $Y \times_X Y = \mathrm{Spa}(T, T^+)$ , then the map  $S^+/\varpi \otimes_{R^+/\varpi} S^+/\varpi \rightarrow T^+/\varpi$  is almost an isomorphism by [Sch12, proof of Proposition 6.18]. Thus,  $R^+/\varpi$  is almost the equalizer of the two maps from  $S^+/\varpi$  to  $T^+/\varpi$ ; passing to the limit over the similar statement for  $\varpi^n$  and inverting  $\varpi$  shows that  $R$  is the equalizer of the two maps from  $S$  to  $T$ , as desired.  $\square$

In fact, the vanishing of cohomology also extends to the v-site.

**Proposition 8.8.** *Let  $X$  be an affinoid perfectoid space. Then  $H_v^i(X, \mathcal{O}) = 0$  for  $i > 0$ , and  $H_v^i(X, \mathcal{O}^+)$  is almost zero for  $i > 0$ .*

*Proof.* Assume that  $X$  is totally disconnected. First, we check vanishing of Čech cohomology, i.e. for any v-cover  $f : Y = \mathrm{Spa}(S, S^+) \rightarrow X = \mathrm{Spa}(R, R^+)$ , we show that the complex

$$0 \rightarrow R^+ \rightarrow S^+ \rightarrow \mathcal{O}^+(Y \times_X Y) \rightarrow \dots$$

is almost exact. As everything is  $\varpi$ -adically complete and  $\varpi$ -torsionfree, where  $\varpi \in R^+$  is a pseudouniformizer, we can check this modulo  $\varpi$ . But then the complex is almost the same as

$$0 \rightarrow R^+/\varpi \rightarrow S^+/\varpi \rightarrow S^+/\varpi \otimes_{R^+/\varpi} S^+/\varpi \rightarrow \dots,$$

which is exact by Proposition 7.23. Similarly, for general affinoid  $X$ , the Čech complex for the affinoid pro-étale cover  $X^{\mathrm{wl}} \rightarrow X$  is almost exact by Proposition 8.5 (iii).

We claim that this implies that  $H_v^i(X, \mathcal{O}^+)$  is almost zero for  $i > 0$  for affinoid  $X$ . Indeed, choose  $i$  minimal for which  $H_v^i(X, \mathcal{O}^+)$  is not almost zero for all affinoid  $X$ , and choose  $X$  and a class  $\alpha \in H_v^i(X, \mathcal{O}^+)$  that is not almost zero. Let  $f : \tilde{X} \rightarrow X$  be a totally disconnected cover; if  $f^*(\alpha)$  is almost zero in  $H_v^i(\tilde{X}, \mathcal{O}^+)$ , a Čech-to-sheaf cohomology spectral sequence gives that  $\check{H}^i(\tilde{X}/X, \mathcal{O}^+)$  is not almost zero, which is a contradiction. Thus, we can replace  $X$  by  $\tilde{X}$  and assume that  $X$  is totally disconnected. But there is some v-cover, without loss of generality by an affinoid  $f : Y \rightarrow X$ , such that  $f^*(\alpha) = 0$ . A Čech-to-sheaf cohomology spectral sequence would then give that  $\check{H}^i(Y/X, \mathcal{O}^+)$  is not almost zero, which contradicts the first paragraph.  $\square$

## 9. DESCENT

In this section, we establish some descent results for perfectoid spaces. As a piece of general notation, we define the category of descent data as follows.

**Definition 9.1.** *Let  $\mathcal{C}$  be a site with fibre products, and let  $F$  be a prestack on  $\mathcal{C}$ , i.e. a functor from  $\mathcal{C}^{\mathrm{op}}$  to groupoids. Let  $f : Y \rightarrow X$  be a map in  $\mathcal{C}$ ,  $p_1, p_2 : Y \times_X Y \rightarrow Y$  the two projections, and  $p_{12}, p_{13}, p_{23} : Y \times_X Y \times_X Y \rightarrow Y \times_X Y$  the three projections. Then*

$$F(Y/X) := \{(s, \alpha) \mid s \in F(Y), \alpha : p_1^*(s) \cong p_2^*(s) \in F(Y \times_X Y), p_{23}^*\alpha \circ p_{12}^*\alpha = p_{13}^*\alpha\},$$

which comes with a natural map  $F(X) \rightarrow F(Y/X)$ , sending  $t \in F(X)$  to  $(s, \alpha)$  with  $s = f^*(t)$  and  $\alpha$  the natural identification  $p_1^*(s) = (p_1 \circ f)^*(t) = (p_2 \circ f)^*(t) = p_2^*(s)$ .

This is the category of objects in  $F(Y)$  equipped with a descent datum to  $X$ . Note that if  $f$  is a covering and  $F$  is a stack, then the natural map  $F(X) \rightarrow F(Y/X)$  is an equivalence.

In this section, we will be interested in several examples of prestacks  $F$  on the pro-étale or v-site of perfectoid spaces.

**Lemma 9.2.** *Let  $F$  be the prestack on the category of perfectoid spaces sending  $X$  to the groupoid of perfectoid spaces over  $X$ . Let  $X$  be a perfectoid space, and  $Y \rightarrow X$  a  $v$ -cover. The functor  $F(X) \rightarrow F(Y/X)$  is fully faithful.*

*Proof.* Given two perfectoid spaces  $X_1, X_2 \rightarrow X$ , we have to see that we can glue morphisms over  $v$ -covers. This follows formally from the fact that the  $v$ -site is subcanonical.  $\square$

In the following, we want to prove that in some instances, the map is an equivalence.

**Proposition 9.3.** *Let  $F$  be the prestack on the category of affinoid perfectoid spaces, sending  $X$  to the groupoid of affinoid perfectoid spaces over  $X$ . Let  $X$  be a totally disconnected perfectoid space, and  $Y \rightarrow X$  a  $v$ -cover, where  $Y$  is an affinoid perfectoid space. The functor  $F(X) \rightarrow F(Y/X)$  is an equivalence of categories.*

*Proof.* Let  $X = \mathrm{Spa}(R, R^+)$ ,  $Y = \mathrm{Spa}(S, S^+)$ , and  $\tilde{Y} = \mathrm{Spa}(\tilde{S}, \tilde{S}^+)$  an affinoid perfectoid space over  $Y$  with a descent datum to  $X$ . Choose a pseudouniformizer  $\varpi \in R^+$  dividing  $p$ . By Proposition 7.23,  $S^+/\varpi$  is faithfully flat over  $R^+/\varpi$ ; in particular,  $(S^+/\varpi)^a$  is faithfully flat over  $(R^+/\varpi)^a$ . One can then use faithfully flat descent in the almost world, cf. [GR03, Section 3.4.1], to see that the  $(S^+/\varpi)^a$ -algebra  $(\tilde{S}^+/\varpi)^a$  descends to an  $(R^+/\varpi)^a$ -algebra, which is perfectoid in the sense of [Sch12, Definition 5.1 (iii)], and thus, by [Sch12, Theorem 5.2], is of the form  $(\tilde{R}^\circ/\varpi)^a$ , where  $\tilde{R}$  is a perfectoid  $R$ -algebra. In particular,  $\tilde{R} \otimes_R S = \tilde{S}$ .

It remains to see that one can find the correct  $\tilde{R}^+ \subset \tilde{R}$ . Let  $\tilde{R}_{\min}^+ \subset \tilde{R}$  be the integral closure of  $R^+ + \tilde{R}^{\circ\circ} \subset \tilde{R}^\circ$ , and  $\tilde{X}' = \mathrm{Spa}(\tilde{R}, \tilde{R}_{\min}^+)$ . We get a map  $\tilde{Y} \rightarrow \tilde{X}'$  such that the induced map  $\tilde{Y} \rightarrow \tilde{Y}' := \tilde{X}' \times_X Y$  is a pro-(open immersion): It is cut out by the conditions  $|f| \leq 1$  for all  $f \in \tilde{S}^+$ . Let  $|\tilde{X}| \subset \tilde{X}'$  denote the image of  $\tilde{Y} \rightarrow \tilde{X}'$ ; then  $\tilde{Y} \subset \tilde{Y}'$  is precisely the preimage of  $|\tilde{X}| \subset \tilde{X}'$  (as  $|\tilde{X}'|$  is the quotient of  $|\tilde{Y}'|$  by the equivalence relation which is the image of

$$|\tilde{Y}' \times_{\tilde{X}'} \tilde{Y}'| \rightarrow |\tilde{Y}'| \times |\tilde{Y}'|).$$

We need to see that  $|\tilde{X}| \subset \tilde{X}'$  is cut out by conditions  $|f| \leq 1$  for elements  $f \in \tilde{R}$ . This is the content of the following lemma.  $\square$

**Lemma 9.4.** *Let  $X = \mathrm{Spa}(R, R^+)$  be a totally disconnected affinoid perfectoid space, let  $\tilde{X} = \mathrm{Spa}(\tilde{R}, \tilde{R}^+)$  be an affinoid perfectoid space over  $X$ , and let  $A \subset |\tilde{X}|$  be a subset. Assume that there is a surjective map  $Y = \mathrm{Spa}(S, S^+) \rightarrow X$  such that the preimage  $B$  of  $A$  in  $|\tilde{Y}|$ , where  $\tilde{Y} = \tilde{X} \times_X Y = \mathrm{Spa}(\tilde{S}, \tilde{S}^+)$ , is an intersection of subsets of the form  $|g| \leq 1$  for elements  $g \in \tilde{S}$ . Then  $A$  is an intersection of subsets of the form  $|f| \leq 1$  for elements  $f \in \tilde{R}$ .*

*Proof.* We may assume that  $X$  is of characteristic  $p$ , as the subsets of the form  $|f| \leq 1$  are unchanged under tilting. Note that  $B$  is pro-constructible, and thus the image  $A$  of  $B$  is also pro-constructible.

As a first reduction step, we reduce to the case that  $X$  is connected. Let  $\pi : X \rightarrow \pi_0(X)$  be the projection. Let  $\tilde{x} \in \tilde{X} \setminus A$  be any point, and let  $c \in \pi_0(X)$  be the connected component  $X_c = \pi^{-1}(c) \subset X$  whose preimage contains  $\tilde{x}$ . If the result is true in the connected case, it holds true for the intersection  $A_c$  of  $A$  with  $\tilde{X}_c := \tilde{X} \times_X X_c$ . Thus, there is a function  $f_c \in H^0(\tilde{X}_c, \mathcal{O}_{\tilde{X}_c})$  such that  $|f_c(\tilde{x})| > 1$ , but  $|f_c| \leq 1$  on  $A_c$ . Modulo functions in  $H^0(\tilde{X}_c, \mathcal{O}_{\tilde{X}_c}^+)$ ,  $f_c$  can be lifted to a function  $f_U \in H^0(\tilde{X}_U, \mathcal{O}_{\tilde{X}_U})$  for some compact open neighborhood  $U$  of  $c$  in  $\pi_0(X)$ , where  $\tilde{X}_U \subset \tilde{X}$

is the preimage of  $U \subset \pi_0(X)$ . Then  $f_U$  still satisfies  $|f_U(\tilde{x})| > 1$ , and the pro-constructible subset  $\{|f_U| > 1\} \cap A_U \subset \tilde{X}_U$  does not meet the fibre over  $c$ . As  $\{|f_U| > 1\} \cap A_U \subset \tilde{X}_U$  is a pro-constructible subset, it is a spectral space, and hence a quasicompactness argument implies that after shrinking  $U$  to a smaller compact open neighborhood of  $c$ ,  $\{|f_U| > 1\} \cap A_U = \emptyset$ , so that  $A_U \subset \{|f_U| \leq 1\}$ . Now, extending  $f_U$  to a function  $f$  on all of  $\tilde{X}$  by setting it to 0 on the preimage of  $\pi_0(X) \setminus U$ , we get a function  $f$  on  $\tilde{X}$  with  $|f(\tilde{x})| > 1$ , but  $|f| \leq 1$  on  $A$ . The intersection of  $\{|f| \leq 1\}$  over all such functions thus gives  $A$ .

Thus, we can assume that  $R = K$  is a perfectoid field, and  $R^+ = K^+$  is an open and bounded valuation subring; fix a pseudouniformizer  $\varpi \in K$ . As a second reduction step, we reduce to the case that  $K$  is algebraically closed. Indeed, let  $C$  be a completed algebraic closure of  $K$ , and  $C^+ \subset C$  the completion of the integral closure of  $K^+$  in  $C$ . If the result is known in the algebraically closed case, we can assume that  $Y = \text{Spa}(C, C^+)$ . Fix a point  $\tilde{x} \in \tilde{X} \setminus A$ , and a lift  $\tilde{y} \in \tilde{Y} \setminus B$  of  $\tilde{x}$ . By assumption, there is some function  $g_{\tilde{y}} \in \tilde{S}$  with  $|g_{\tilde{y}}(\tilde{y})| > 1$ , but  $|g_{\tilde{y}}| \leq 1$  on  $B$ . Now  $Y$  is an inverse limit of  $\text{Spa}(L, L^+)$  over all finite extensions  $L \subset C$  of  $K$  with integral closure  $L^+ \subset L$  of  $K^+$  in  $L$ . Approximating the function  $g_{\tilde{y}}$  modulo  $\tilde{S}^+$ , we can assume that  $g_{\tilde{y}}$  is a function on  $\tilde{X} \times_X \text{Spa}(L, L^+) =: \text{Spa}(\tilde{R}_L, \tilde{R}_L^+)$ , for some big enough finite extension  $L$  of  $K$ . In this case,  $\tilde{R}_L = \tilde{R} \otimes_K L$  is a finite free  $\tilde{R}$ -algebra; consider the characteristic polynomial  $P_{g_{\tilde{y}}}(X) = X^d + f_1 X^{d-1} + \dots + f_d$  of  $g_{\tilde{y}}$  acting on  $\tilde{R}_L$ , so all  $f_i \in \tilde{R}$ . As  $|g_{\tilde{y}}| \leq 1$  on the preimage of  $A$ , it follows that  $|f_i| \leq 1$  on  $A$  for all  $i = 1, \dots, d$ . On the other hand, if we had  $|f_i(\tilde{x})| \leq 1$  for all  $i = 1, \dots, d$ , then the equation

$$0 = P_{g_{\tilde{y}}}(g_{\tilde{y}}) = g_{\tilde{y}}^d + f_1 g_{\tilde{y}}^{d-1} + \dots + f_d$$

would imply that  $|g_{\tilde{y}}(\tilde{y})| \leq 1$ , which is a contradiction. Thus, at least one  $f_i$  satisfies  $|f_i| \leq 1$  on  $A$  while  $|f_i(\tilde{x})| > 1$ . Intersecting all such  $f_i$  for varying  $\tilde{x}$  thus writes  $A$  as an intersection of sets of the form  $\{|f| \leq 1\}$  for varying  $f \in \tilde{R}$ .

Finally, assume that  $R = C$  is a complete algebraically closed nonarchimedean field of characteristic  $p$ , and  $R^+ = C^+$  is an open and bounded valuation subring. We may assume that similarly  $S = L$  is a perfectoid field, and  $S^+ = L^+$  is an open and bounded valuation subring. Fix a point  $\tilde{x} \in \tilde{X} \setminus A$ , with preimage  $\tilde{Y}_{\tilde{x}} \subset \tilde{Y}$ , a pro-constructible subset. Let  $x \in X$  be the image of  $\tilde{x}$ . By assumption, for any  $\tilde{y} \in \tilde{Y}_{\tilde{x}}$ , there is some function  $g_{\tilde{y}} \in \tilde{S}$  with  $|g_{\tilde{y}}(\tilde{y})| > 1$ , but  $|g_{\tilde{y}}| \leq 1$  on  $B$ . By quasicompactness, we can find finitely many  $g_1, \dots, g_n$  such that  $|g_i| \leq 1$  on  $B$ , but

$$\tilde{Y}_{\tilde{x}} \subset \bigcup_{i=1}^n \{|g_i| > 1\}.$$

Up to an element of  $\tilde{S}^+$ , we can approximate  $g_i \in \tilde{S} = S \widehat{\otimes}_R \tilde{R}$  as a sum  $\varpi^{-n_i} \sum_{j=1}^{m_i} s_{ij} \otimes \tilde{r}_{ij}$  with  $s_{ij} \in S$ ,  $\tilde{r}_{ij} \in \tilde{R}$ . Let  $Y' = \text{Spa}(S', S'^+)$ , with  $S' = C \langle (T_{ij}^{1/p^\infty})_{i=1, \dots, n, j=1, \dots, m_i} \rangle$ ,  $S'^+ = C^+ \langle (T_{ij}^{1/p^\infty})_{i=1, \dots, n, j=1, \dots, m_i} \rangle$ , and consider the map  $Y \rightarrow Y'$  given by  $T_{ij} \mapsto s_{ij}$ . Then there are functions  $g'_i = \varpi^{-n_i} \sum_{j=1}^{m_i} T_{ij} \otimes \tilde{r}_{ij} \in \tilde{S}'$ , where  $\tilde{Y}' = \tilde{X} \times_X Y' = \text{Spa}(\tilde{S}', \tilde{S}'^+)$ . These functions pull back to  $g_i$ , and in particular they satisfy  $|g'_i| \leq 1$  on  $B$ . Let  $B' \subset |\tilde{Y}'|$  be the preimage of  $A$ . Let  $f : \tilde{Y}' \rightarrow Y'$  be the projection. Let  $W \subset Y'$  be the image of  $Y \rightarrow Y'$ ; thus  $W = \text{Spa}(L', L'^+)$  is the set of generalizations of a point  $y' \in Y'$  (given by the image of the closed point of  $Y$ ), and  $W$  can thus be written as the intersection of all rational subsets  $V \subset Y'$  containing  $W$ . Then

$B' \cap f^{-1}(W) \subset \{|g'_i| \leq 1\}$ , as  $B$  surjects onto  $B' \cap f^{-1}(W)$ . By a quasicompactness argument, this implies that there is some rational neighborhood  $V \subset Y'$  of  $W$  such that  $B' \cap f^{-1}(V) \subset \{|g'_i| \leq 1\}$ . Similarly, the fibre of  $\tilde{Y}'_{\tilde{x}}$  over  $W$  is contained in  $\bigcup_{i=1}^n \{|g'_i| > 1\}$ ; said differently, the spectral space

$$M = \{\tilde{y} \in \tilde{Y}'_{\tilde{x}} \mid \forall i = 1, \dots, n : |g'_i(\tilde{y})| \leq 1\}$$

has empty fiber over  $W \subset Y'$ . By a quasicompactness argument, this implies that replacing  $V$  by a smaller rational subset of  $Y'$  containing  $W$ , the fiber of  $M$  over  $V$  is empty. Thus, we can choose  $V$  with the properties

$$(9.1) \quad B' \cap f^{-1}(V) \subset \bigcap_{i=1}^n \{|g'_i| \leq 1\} ,$$

$$(9.2) \quad \tilde{Y}'_{\tilde{x}} \cap f^{-1}(V) \subset \bigcup_{i=1}^n \{|g'_i| > 1\} .$$

Now  $V$  is a rational subset of

$$\mathrm{Spa}(C\langle\langle(T_{ij}^{1/p^\infty})_{i=1, \dots, n, j=1, \dots, m_i}\rangle\rangle, C^+\langle\langle(T_{ij}^{1/p^\infty})_{i=1, \dots, n, j=1, \dots, m_i}\rangle\rangle)$$

mapping surjectively to  $\mathrm{Spa}(C, C^+)$ . By Lemma 9.5 this implies that there is a  $(C, C^+)$ -valued point  $z : \mathrm{Spa}(C, C^+) \rightarrow V$ . Let  $f_i \in \tilde{R}$  be the evaluation of  $g'_i$  at  $z$ . Then  $|f_i| \leq 1$  on  $A$  by (9.1), but  $\tilde{x} \in \bigcup_{i=1}^n \{|f_i| > 1\}$  by (9.2). This finishes the proof, as now some  $f = f_i$  satisfies  $|f| \leq 1$  on  $A$ , but  $|f(\tilde{x})| > 1$ , and intersecting  $\{|f| \leq 1\}$  over all such  $f$  for varying  $\tilde{x}$  gives  $A$ .  $\square$

**Lemma 9.5.** *Let  $C$  be an algebraically closed nonarchimedean field with open and bounded valuation subring  $C^+ \subset C$ . Let  $V$  be an open subset of*

$$\mathrm{Spa}(C\langle T_1, \dots, T_d \rangle, C^+\langle T_1, \dots, T_d \rangle)$$

*mapping surjectively to  $\mathrm{Spa}(C, C^+)$ . Then there is a section  $\mathrm{Spa}(C, C^+) \rightarrow V$ .*

*Proof.* We may assume that  $V$  is rational, and in particular affinoid. Then  $\mathfrak{X} = \mathrm{Spf} \mathcal{O}^+(V)$  is a formal scheme which is flat and topologically of finite type (thus, of finite presentation) over  $\mathrm{Spf} C^+$ ; moreover,  $\mathfrak{X} \rightarrow \mathrm{Spf} C^+$  is surjective. We need to see that  $\mathfrak{X} \rightarrow \mathrm{Spf} C^+$  has a section. First,  $R = C^+/C^\circ$  is a valuation ring with algebraically closed fraction field  $K$ , and  $X = \mathfrak{X} \times_{\mathrm{Spf} C^+} \mathrm{Spec} R$  is a scheme of finite presentation and faithfully flat over  $R$ . This implies that  $X(R)$  is nonempty: Any finitely presented faithfully flat cover of a strictly henselian local ring is refined by a finite flat locally free cover, cf. [Gro68, after Lemma 11.2], and finite flat covers of  $\mathrm{Spec} R$  split. But  $\mathfrak{X}(C^+) = \mathfrak{X}(\mathcal{O}_C) \times_{X(K)} X(R)$ , and the reduction map  $\mathfrak{X}(\mathcal{O}_C) \rightarrow X(K)$  is surjective.  $\square$

We need another result for morphisms which are not necessarily affinoid. We will however need to make a separatedness assumption.

**Proposition 9.6.** *Let  $F$  be the prestack on the category of perfectoid spaces sending a perfectoid space  $X$  to the groupoid of separated pro-étale perfectoid spaces over  $X$ . Let  $X$  be a strictly totally disconnected perfectoid space, and  $Y \rightarrow X$  a  $v$ -cover. Then  $F(X) \rightarrow F(Y/X)$  is an equivalence of categories.*

*Proof.* By Lemma 9.2, we only have to prove that all descent data are effective; for this, we may assume that  $Y$  is strictly totally disconnected as well.

Let  $R = \tilde{Y} \times_X Y \subset \tilde{Y} \times \tilde{Y}$  be the equivalence relation on  $\tilde{Y}$  corresponding to the descent datum. We claim that the image  $R' \subset |\tilde{Y}| \times |\tilde{Y}|$  satisfies the hypothesis of Lemma 2.7. First,  $|\tilde{Y}|$  is a quasiseparated locally spectral space. As all maps of perfectoid spaces are generalizing, it follows that  $R' \rightarrow \tilde{Y}$  is generalizing; also, it is quasicompact (as it is a quotient of the map  $|R| = |\tilde{Y} \times_X Y| \rightarrow |\tilde{Y}|$ , and  $Y \rightarrow X$  is quasicompact). Finally,  $R'$  is pro-constructible: For any quasicompact open subset  $W \subset \tilde{Y}$ , the intersection of  $R'$  with  $|W| \times |W|$  is the image of the spectral space  $W \times_X Y \cap Y \times_X W$  under the spectral map to  $|W| \times |W|$ .

Thus, by Lemma 2.7, for any quasicompact open  $W \subset \tilde{Y}$  we can find  $R$ -invariant subsets  $U \subset E \subset \tilde{Y}$  such that  $U$  is open in  $\tilde{Y}$  and contains  $W$ , and  $E$  is an intersection of a nonempty family of quasicompact open subsets. Thus,  $E \subset \tilde{Y}$  corresponds by Lemma 7.19 to an affinoid pro-étale perfectoid space, which we will still denote by  $E$ . As  $E$  is  $R$ -invariant, it comes with an induced descent datum, which is effective by Proposition 9.3. We find some affinoid perfectoid space  $E_X \rightarrow X$  whose pullback to  $Y$  is  $E$ ; it follows (from Lemma 7.19) that  $E_X \rightarrow X$  is affinoid pro-étale. Now the open  $R$ -invariant subset  $U \subset E$  descends to some open subset  $U_X \subset E_X$ , which is thus pro-étale over  $X$ . For varying  $W$ , the  $U$  will cover  $\tilde{Y}$ , and thus the  $U_X$  will glue to give the desired space  $\tilde{X} \rightarrow X$  pro-étale over  $X$ .  $\square$

Finally, we need some stronger descent results for étale morphisms.

**Proposition 9.7.** *The prestack on the category of perfectoid spaces sending any perfectoid space  $X$  to the groupoid of separated étale perfectoid spaces over  $X$  is a stack in the  $v$ -topology.*

*Similarly, the prestack on the category of perfectoid spaces sending any perfectoid space  $X$  to the groupoid of finite étale perfectoid spaces over  $X$  is a stack in the  $v$ -topology.*

*Proof.* We give the proof in the more difficult case of separated étale maps; it is easy to see that the property of being finite étale passes through all the arguments.

Let  $F$  be the prestack on the category of perfectoid spaces sending any perfectoid space  $X$  to the groupoid of separated étale perfectoid spaces over  $X$ . Let  $f : Y \rightarrow X$  be any  $v$ -cover; we need to prove that  $F(X) \rightarrow F(Y/X)$  is an equivalence. From Lemma 9.2, we know that the functor is fully faithful, so we need to show effectivity of descent. Assume first that  $X$  is strictly totally disconnected; we may also assume that  $Y$  is strictly totally disconnected. Then Proposition 9.6 implies that any separated étale  $\tilde{Y} \rightarrow Y$  with descent data descends to a separated pro-étale  $\tilde{X} \rightarrow X$ . It remains to see that  $\tilde{X} \rightarrow X$  is étale. For this, we may assume that  $\tilde{X}$  is affinoid. Under the equivalence of Corollary 7.22, we have to see that  $|\tilde{X}| \rightarrow |X|$  is a local isomorphism. But  $|\tilde{X} \times_X Y| \rightarrow |\tilde{X}| \times_{|X|} |Y|$  is a homeomorphism, and so we know that  $|\tilde{X}| \rightarrow |X|$  becomes a local isomorphism after pullback along the surjection  $|Y| \rightarrow |X|$ . By Lemma 9.8 below, we see that  $|\tilde{X}| \rightarrow |X|$  is a local isomorphism, as desired.

This finishes the case that  $X$  is strictly totally disconnected. In general, we can assume that  $X$  is affinoid, and by Lemma 7.18 find an affinoid pro-étale surjective and universally open map  $Y' \rightarrow X$  with  $Y'$  strictly totally disconnected. Then we have a  $v$ -cover  $Y \times_X Y' \rightarrow X$  refining  $Y \rightarrow X$ , and we already know effectivity of descent for  $Y \times_X Y' \rightarrow Y'$ . In other words, we may replace  $Y$  by  $Y'$ , and assume that  $Y \rightarrow X$  is affinoid pro-étale and universally open. Write  $Y = \varprojlim_i Y_i \rightarrow X$  as a cofiltered inverse limit of étale maps  $Y_i \rightarrow X$ , where all  $Y_i$  are affinoid.

Now let  $\tilde{Y} \rightarrow Y$  be a separated étale map equipped with a descent datum to  $X$ , which induces an equivalence relation  $\tilde{R} \subset \tilde{Y} \times \tilde{Y}$ . The map  $\tilde{R} \rightarrow \tilde{Y}$  can be identified with the projection  $\tilde{Y} \times_X Y \rightarrow \tilde{Y}$ , which is open (as  $Y \rightarrow X$  is universally open). From Lemma 2.10, we see that  $|\tilde{Y}|/|\tilde{R}|$  is a quasiseparated locally spectral space. Covering it by quasicompact open subsets, we can reduce to the case that  $\tilde{Y}$  is quasicompact.

In this situation, Proposition 6.4 (iii) says that

$$Y_{\text{ét, qc, sep}} = 2\text{-}\varinjlim_i (Y_i)_{\text{ét, qc, sep}},$$

so any quasicompact separated étale  $Y$ -space  $\tilde{Y}$  comes via pullback from some quasicompact separated étale  $Y_i$ -space  $\tilde{Y}_i$  for  $i$  large enough. Similarly, any descent datum is defined at some finite level, so we can replace  $Y$  by  $Y_i$  for  $i$  sufficiently large, and reduce to the case that  $Y \rightarrow X$  is étale. By the variant [KL15, Proposition 8.2.20] of an argument of de Jong and van der Put, [dJvdP96, Proposition 3.2.2], one further reduces to the case that  $Y \rightarrow X$  is finite étale. We can also assume that it is Galois, with Galois group  $G$ .

Finally, take any point  $x \in X$ . As  $(X_x)_{\text{ét, qc, sep}} = 2\text{-}\varinjlim_{U \ni x} U_{\text{ét, qc, sep}}$  (and similarly for pullbacks to  $Y$  and  $Y \times_X Y$ ), it is enough to show effectivity of descent after base change to  $X_x$ ; thus we may assume that  $X = \text{Spa}(K, K^+)$  for some perfectoid field  $K$  with open and bounded valuation subring  $K^+ \subset K$ . It remains to see that in this case, if  $\tilde{Y} \in X_{\text{ét, qc, sep}}$  has a free  $G$ -action for some finite group  $G$ , then the quotient  $\tilde{Y}/G$  exists (with  $\tilde{Y} \times_{\tilde{Y}/G} \tilde{Y} = \tilde{Y} \times G$ ). For this, we use Lemma 9.9 below to reduce to the case that  $\tilde{Y} \rightarrow X$  is finite étale, which reduces us to the algebraic case of finite étale  $K$ -algebras.  $\square$

The following two lemmas were used in the proof.

**Lemma 9.8.** *Let  $f : S' \rightarrow S$  be a map of spectral spaces, and let  $p : \tilde{S} \rightarrow S$  be a surjective, spectral and generalizing map of spectral spaces, with pullback  $\tilde{f} : \tilde{S}' = S' \times_S \tilde{S} \rightarrow \tilde{S}$ . If  $\tilde{f}$  is a local isomorphism, then  $f$  is a local isomorphism.*

*Proof.* Let  $p' : \tilde{S}' \rightarrow S'$  denote the projection. Let  $s' \in S'$  be any point, with image  $s \in S$ . We claim that there is a quasicompact open neighborhood  $U' \subset S'$  of  $s'$  such that  $f|_{U'} : U' \rightarrow S$  is injective. For this, note that  $\Delta_{\tilde{f}} : \tilde{S}' \rightarrow \tilde{S}' \times_{\tilde{S}} \tilde{S}'$  is an open immersion (as  $\tilde{f}$  is a local isomorphism); this implies by Lemma 2.5 that also  $\Delta_f : S' \rightarrow S' \times_S S'$  is an open immersion. In particular, there is a quasicompact open neighborhood  $U'$  of  $s'$  in  $S'$  such that  $U' \times_S U' \subset \Delta_f(S')$ , i.e.  $U' \rightarrow S$  is injective.

Replacing  $S'$  by  $U'$ , we may assume that  $f$  is injective. In that case  $\tilde{f}$  is injective and a local isomorphism. This implies that  $\tilde{f}$  is an open immersion (as the image of  $\tilde{f}$  is open, and the map to the image is a continuous open bijective map). By Lemma 2.5, it follows that  $f$  is an open immersion.  $\square$

**Lemma 9.9.** *Let  $X = \text{Spa}(K, K^+)$ , where  $K$  is a perfectoid field, and  $K^+ \subset K$  is an open and bounded integrally closed subring (not necessarily a valuation subring). Let  $Y \rightarrow X$  be a quasicompact separated étale map. Then there is a functorial factorization  $Y \hookrightarrow \bar{Y} \rightarrow X$ , where  $Y \hookrightarrow \bar{Y}$  is a quasicompact open immersion, and  $\bar{Y} \rightarrow X$  is finite étale.*

**Remark 9.10.** This compactification is a special case of the canonical compactifications introduced in Proposition 18.6 below.

*Proof.* Let  $X^\circ = \mathrm{Spa}(K, \mathcal{O}_K)$ , and  $Y^\circ = Y \times_X X^\circ$ . Then  $Y^\circ$  is finite étale over  $X^\circ$ ; in particular,  $|Y^\circ|$  is a finite set of maximal points. Each of them is a pro-constructible subset, and their closures (which are given by their sets of specializations) do not intersect, as any point generalizes to a unique maximal point. Thus, replacing  $Y$  by an open and closed subset, we may assume that  $Y^\circ = \mathrm{Spa}(L, \mathcal{O}_L)$  is a point, where  $L$  is a finite extension of  $K$ . Then  $Y^\circ \subset Y$  is the unique maximal point, so  $Y$  is connected, and  $\mathcal{O}_Y$  is the constant sheaf  $L$ . Let  $\bar{Y} = \mathrm{Spa}(L, L^+)$ , where  $L^+$  is the integral closure of  $K^+$  in  $L$ . Then  $\bar{Y} \rightarrow X$  is finite étale, and there is a natural map  $Y \rightarrow \bar{Y}$ , given by the map  $(L, L^+) \rightarrow (\mathcal{O}_Y(Y), \mathcal{O}_Y^+(Y))$  of algebras. Then  $Y \rightarrow \bar{Y}$  is a separated étale map of perfectoid spaces, which after pullback along  $X^\circ \subset X$  is an isomorphism. In particular, it is an isomorphism on rank-1-valued points, so by the valuative criterion for separatedness (Proposition 5.11), it is an injection (cf. Proposition 5.3 (ii)). Thus,  $Y$  is determined by the image of the injective map  $|Y| \rightarrow |\bar{Y}|$ , which is a quasicompact open subspace of  $\bar{Y}$ , as  $Y \rightarrow \bar{Y}$  is étale. Therefore,  $Y \rightarrow \bar{Y}$  is a quasicompact open immersion, as desired.  $\square$

**Corollary 9.11.** *Let  $f : Y \rightarrow X$  be a map of perfectoid spaces, and let  $\tilde{X} \rightarrow X$  be a v-cover, with pullback  $f : \tilde{Y} = Y \times_X \tilde{X} \rightarrow \tilde{X}$ .*

- (i) *Assume that  $\tilde{f}$  is pro-étale, and  $X$  is strictly totally disconnected. Then  $f$  is pro-étale.*
- (ii) *Assume that  $\tilde{f}$  is étale. Then  $f$  is étale.*
- (iii) *Assume that  $\tilde{f}$  is finite étale. Then  $f$  is finite étale.*

*Proof.* We can assume that  $X$  is affinoid. In cases (i) and (ii), the conclusion is local on  $Y$ , so we can in these cases assume that  $Y$  is affinoid; in particular,  $f$  is separated. Now (i) follows from Proposition 9.6, and parts (ii) and (iii) from Proposition 9.7.  $\square$

## 10. MORPHISMS OF V-STACKS

The descent results of the previous section make it possible to define good notions of étale and (quasi-)pro-étale morphisms between pro-étale stacks on  $\mathrm{Perfd}$ . In the following, we will often confuse a perfectoid space and the sheaf that it represents.

**Definition 10.1.** *Let  $f : Y' \rightarrow Y$  be a map of pro-étale stacks on the category of ( $\kappa$ -small) perfectoid spaces  $\mathrm{Perfd}$ .*

- (i) *Assume that  $f$  is locally separated, i.e. there is an open cover of  $Y'$  over which  $f$  becomes separated. The map  $f$  is quasi-pro-étale if for any strictly totally disconnected  $X$  with a map  $X \rightarrow Y$  (i.e., a section in  $Y(X)$ ), the pullback  $Y' \times_Y X$  is representable and  $Y' \times_Y X \rightarrow X$  is pro-étale.*
- (ii) *Assume that  $f$  is locally separated. The map  $f$  is étale if for any perfectoid space  $X$  with a map  $X \rightarrow Y$ , the pullback  $Y' \times_Y X$  is representable, and  $Y' \times_Y X \rightarrow X$  is étale.*
- (iii) *The map  $f$  is finite étale if for any perfectoid space  $X$  with a map  $X \rightarrow Y$ , the pullback  $Y' \times_Y X \rightarrow X$  is representable, and  $Y' \times_Y X \rightarrow X$  is finite étale.*

In parts (i) and (ii), we restrict the definition to the case that  $f$  is locally separated, as otherwise one may want to give a slightly different definition. Note that  $f$  is always locally separated if  $Y'$  is a perfectoid space, and  $Y$  is a v-sheaf. We make the following important convention.

**Convention 10.2.** In the remainder of this text, we take “locally separated” as part of the definition of étale or quasi-pro-étale maps of pro-étale stacks.

Any pro-étale morphism of perfectoid spaces is quasi-pro-étale, but the converse need not hold. This is similar to the difference between pro-étale and weakly étale morphisms of schemes, cf. [BS15].

The following proposition is immediate from the definitions.

**Proposition 10.3.** *Let  $f : X' \rightarrow X$  be a map of perfectoid spaces. Then  $f$  is étale, resp. finite étale, if and only if the corresponding map of pro-étale sheaves on  $\text{Perfd}$  is étale, resp. finite étale. If  $X$  is strictly totally disconnected, then  $f$  is pro-étale if and only if the corresponding map of pro-étale sheaves on  $\text{Perfd}$  is quasi-pro-étale.  $\square$*

**Proposition 10.4.** *Let  $f : Y_1 \rightarrow Y_2$ ,  $g : Y_2 \rightarrow Y_3$  be morphisms of pro-étale stacks on  $\text{Perfd}$ , with composite  $h = g \circ f : Y_1 \rightarrow Y_3$ .*

- (i) *If  $f$  and  $g$  are quasi-pro-étale (resp. étale, finite étale), then so is  $h$ .*
- (ii) *If  $g$  and  $h$  are quasi-pro-étale (resp. étale, finite étale), then so is  $f$ .*
- (iii) *Any pullback of a quasi-pro-étale (resp. étale, finite étale) map is quasi-pro-étale (resp. étale, finite étale).*

In some situations, one can also deduce that  $g$  is quasi-pro-étale from knowing that  $f$  and  $h$  are quasi-pro-étale, cf. Proposition 11.30 below.

*Proof.* These are all straightforward. Let us check the most subtle part, which is (i) in the quasi-pro-étale case. We may assume that  $X_3 = Y_3$  is a strictly totally disconnected perfectoid space. In this case,  $Y_2$  is representable and pro-étale over  $X_3$ . As such,  $Y_2$  admits an open cover by strictly totally disconnected perfectoid spaces  $X_2 \subset Y_2$ . The preimage of any such  $X_2 \subset Y_2$  in  $Y_1$  is representable and pro-étale over  $X_2$ . It follows that  $Y_1$  has an open cover by representable functors, so  $Y_1$  is itself representable. Also  $Y_1 \rightarrow Y_2$  is pro-étale locally on  $Y_2$ , thus pro-étale, and then  $Y_1 \rightarrow Y_3$  is pro-étale as a composite of pro-étale maps.  $\square$

We will also need to extend other notions like injections, open and closed immersions, and separated maps to morphisms of sheaves. These notions work better for v-sheaves, because of the following result.

**Proposition 10.5.** *Let  $X$  be a totally disconnected perfectoid space, and let  $Y \subset X$  be a sub-v-sheaf. Then  $Y$  is ind-representable. More precisely, one can write  $Y$  as a ( $\kappa$ -small) filtered colimit of  $Y_i \subset Y \subset X$  which are pro-constructible and generalizing subsets of  $X$ , in particular affinoid pro-étale over  $X$ .*

*Proof.* Let  $Z$  be any affinoid perfectoid space with a map  $Z \rightarrow Y$ . Then the image of the composition  $Z \rightarrow Y \rightarrow X$  on topological spaces is a pro-constructible and generalizing subset  $\bar{Z} \subset X$ . By Lemma 7.6, the subset  $\bar{Z}$  is naturally an affinoid perfectoid space, and  $Z \rightarrow \bar{Z}$  is a v-cover. As  $Y$  is sub-v-sheaf of  $X$ , it follows that one gets a map  $\bar{Z} \rightarrow Y$ . Note that the category of pro-constructible generalizing subsets  $Y_i \subset X$  for which the inclusion  $Y_i \rightarrow X$  factors over  $Y$  is filtered: This follows from the sheaf property of  $Y$ . This implies that  $Y$  is the filtered colimit of such  $Y_i$ , as desired.  $\square$

**Corollary 10.6.** *Let  $f : Y' \rightarrow Y$  be a quasicompact injection of v-stacks. Then  $f$  is quasi-pro-étale, and for every totally disconnected perfectoid space  $X$  mapping to  $Y$ , the fibre product  $Y' \times_Y X$  is representable by a pro-constructible and generalizing subset of  $X$ .*

*Proof.* We may assume that  $Y = X$  is a totally disconnected perfectoid space. The previous proposition shows that  $Y' \subset X$  is ind-representable. By quasicompactness of  $Y'$ , this implies that



$Y'$  is representable; applying the proposition again, we see that  $Y' \subset X$  is affinoid pro-étale, as desired.  $\square$

**Definition 10.7.** *Let  $f : Y' \rightarrow Y$  be a map of pro-étale stacks.*

- (i) *The map  $f$  is an open immersion if for every perfectoid space  $X$  mapping to  $Y$ , the pullback  $Y' \times_Y X \rightarrow X$  is representable by an open immersion.*
- (ii) *The map  $f$  is a closed immersion if for every totally disconnected perfectoid space  $X$  mapping to  $Y$ , the pullback  $Y' \times_Y X \subset X$  is representable by a closed immersion.*
- (iii) *The map  $f$  is separated if  $\Delta_f : Y' \rightarrow Y' \times_Y Y'$  is a closed immersion.*
- (iv) *The map  $f$  is 0-truncated if for all  $X \in \text{Perfd}$ , the map of groupoids  $f(X) : Y'(X) \rightarrow Y(X)$  is faithful.*

*Moreover, the pro-étale stack  $Y$  is separated if the map  $Y \rightarrow *$  is separated.*

**Remark 10.8.** We will only occasionally use the last definition, with which one has to be a bit careful. Namely, it can happen that  $Y$  is separated, but  $Y$  is not quasiseparated: The issue is that the notions of  $Y$  being quasiseparated and of  $Y \rightarrow *$  being quasiseparated do not coincide. (An example is given by  $Y = X/\varphi^{\mathbb{Z}}$ , where  $X$  is some perfectoid space of characteristic  $p$ , and  $\varphi$  is its absolute Frobenius: This defines a diamond  $Y$  that is separated, but not quasiseparated.)

In part (iv), it is equivalent to ask that for all affinoid perfectoid spaces (or more generally, all pro-étale sheaves)  $X$  mapping to  $Y$ , the pro-étale stack  $Y' \times_Y X$  is discrete, i.e. a pro-étale sheaf. It is also equivalent to ask that  $\Delta_f : Y' \rightarrow Y' \times_Y Y'$  is an injection; in particular, separated maps are automatically 0-truncated. (This differs from the convention in [Sta, Tag 04YW].)

For a map of perfectoid spaces  $f : X' \rightarrow X$ , these notions are equivalent to the previously defined notions. Moreover, there is again a valuative criterion for separatedness.

**Proposition 10.9.** *Let  $f : Y' \rightarrow Y$  be a map of  $v$ -stacks. Then  $f$  is separated if and only if  $f$  is 0-truncated, quasiseparated, and for every perfectoid field  $K$  with ring of integers  $\mathcal{O}_K$  and an open and bounded valuation subring  $K^+$ , and any diagram*

$$\begin{array}{ccc} \text{Spa}(K, \mathcal{O}_K) & \longrightarrow & Y' \\ \downarrow & \nearrow \text{dotted} & \downarrow f \\ \text{Spa}(K, K^+) & \longrightarrow & Y \end{array},$$

*there exists at most one dotted arrow making the diagram commute.*

Note that if  $f$  is 0-truncated, the groupoid of dotted arrows is automatically discrete, i.e. there are no automorphisms.

*Proof.* If  $f$  is separated, then  $\Delta_f$  is a closed immersion, which implies that  $\Delta_f$  is quasicompact, so that  $f$  is quasiseparated. Also, closed immersions are specializing, which shows that there is at most one such dotted arrow. Conversely, if  $f$  is quasiseparated, then  $\Delta_f$  is quasicompact. Thus, by Corollary 10.6, the diagonal  $\Delta_f$  is quasi-pro-étale. It remains to check that for any totally disconnected perfectoid space  $X$  mapping to  $Y' \times_Y Y'$ , the pro-constructible generalizing subset

$$Y' \times_{Y' \times_Y Y'} X \subset X$$

is closed. For this, it is enough to check that it is closed under specializations, which is exactly the valuative criterion.  $\square$

Sometimes it is useful to know that a version of the valuative criterion holds true for general pairs  $(R, R^+)$ .

**Proposition 10.10.** *Let  $f : Y' \rightarrow Y$  be a separated map of v-stacks, and let  $R$  be a perfectoid Tate ring with an open and integrally closed subring  $R^+ \subset R^\circ$ . Then, for any diagram*

$$\begin{array}{ccc} \mathrm{Spa}(R, R^\circ) & \longrightarrow & Y' \\ \downarrow & \nearrow & \downarrow f \\ \mathrm{Spa}(R, R^+) & \longrightarrow & Y \end{array},$$

*there exists at most one dotted arrow making the diagram commute.*

*Proof.* Assume that  $a, b : \mathrm{Spa}(R, R^+) \rightarrow Y'$  are two maps projecting to the same map  $\bar{a} = \bar{b} : \mathrm{Spa}(R, R^+) \rightarrow Y$ , and with same restriction  $a^\circ = b^\circ : \mathrm{Spa}(R, R^\circ) \rightarrow Y'$ . Consider the sub-v-sheaf

$$Z = \mathrm{Spa}(R, R^+) \times_{Y' \times_Y Y'} Y' \subset \mathrm{Spa}(R, R^+),$$

where the two projections  $\mathrm{Spa}(R, R^+) \rightarrow Y'$  are given by  $a$  and  $b$ . As  $f$  is separated,  $\Delta_f$  is a closed immersion, so  $Z \subset \mathrm{Spa}(R, R^+)$  is a closed immersion. To check that it is an equality, we can assume that  $R$  is totally disconnected; then  $Z$  is a totally disconnected perfectoid space as well. Now for any perfectoid field  $(K, K^+)$ , we know that  $Z(K, K^+) = \mathrm{Spa}(R, R^+)(K, K^+)$  by the valuative criterion, Proposition 10.9. Thus,  $|Z| = |\mathrm{Spa}(R, R^+)|$ , and then  $Z = \mathrm{Spa}(R, R^+)$ .  $\square$

The next proposition shows that many of these conditions can be checked v-locally on the target.

**Proposition 10.11.** *Let  $f : Y' \rightarrow Y$  be a map of v-stacks, and let  $g : \tilde{Y} \rightarrow Y$  be a surjective map of v-stacks. Let  $\tilde{f} : \tilde{Y}' = \tilde{Y} \times_Y Y' \rightarrow \tilde{Y}$  be the pullback of  $f$ .*

- (o) *If  $\tilde{f}$  is quasicompact (resp. quasiseparated), then  $f$  is quasicompact (resp. quasiseparated).*
- (i) *If  $\tilde{f}$  is an open (resp. closed) immersion, then  $f$  is an open (resp. closed) immersion.*
- (ii) *If  $\tilde{f}$  is separated, then  $f$  is separated.*
- (iii) *If  $\tilde{f}$  is finite étale, then  $f$  is finite étale.*
- (iv) *If  $\tilde{f}$  is separated and étale, then  $f$  is separated and étale.*
- (v) *If  $\tilde{f}$  is separated and quasi-pro-étale, then  $f$  is separated and quasi-pro-étale.*

*Proof.* For (o), the quasiseparated case reduces to the quasicompact case by passing to the diagonal. One can also replace  $Y$  by an affinoid perfectoid space  $X$ , and  $\tilde{Y}$  by a v-cover  $\tilde{X} \rightarrow X$ , where  $\tilde{X}$  may be assumed to be an affinoid perfectoid space. Now let  $Y'_i, i \in I$ , be a set of v-sheaves with a surjection  $\bigsqcup_{i \in I} Y'_i \rightarrow Y'$ . The pullback  $\bigsqcup_{i \in I} \tilde{Y}'_i \rightarrow \tilde{Y}'$  has a finite subcover by quasicompactness of  $\tilde{Y}'$ , so assume  $J \subset I$  is a finite subset such that  $\bigsqcup_{i \in J} \tilde{Y}'_i \rightarrow \tilde{Y}'$  is surjective. In particular, the composite  $\bigsqcup_{i \in J} \tilde{Y}'_i \rightarrow \tilde{Y}' \rightarrow Y'$  is surjective, which implies that  $\bigsqcup_{i \in J} Y'_i \rightarrow Y'$  is surjective.

Part (ii) reduces to (i) by passing to the diagonal. Now for (i), (iii), (iv) and (v), we need to check that something holds after any pullback  $X \rightarrow Y$  for an affinoid perfectoid space  $X$ . As  $\tilde{Y} \rightarrow Y$  is surjective, we can find a v-cover  $\tilde{X} \rightarrow X$  such that  $X \rightarrow Y$  lifts to  $\tilde{X} \rightarrow \tilde{Y}$ . We may then replace  $Y$  by  $X$  and  $\tilde{Y}$  by  $\tilde{X}$ .

Now (i) for open immersions follows because  $|\tilde{X}| \rightarrow |X|$  is a quotient map by Lemma 2.5. For closed immersions, one uses in addition that if  $X$  is totally disconnected, then  $f$  is representable by Corollary 10.6, as  $\tilde{f}$  is a quasicompact injection, and thus  $f$  is a quasicompact injection.

Finally, parts (iii) and (iv) follow from Proposition 9.7, and part (v) (in which case one assumes that  $X$  is strictly totally disconnected) from Proposition 9.6.  $\square$

As an application, let us discuss  $\underline{G}$ -torsors for locally profinite groups  $G$ . Here, for any topological space  $T$ , we denote by  $\underline{T}$  the associated v-sheaf given by

$$\underline{T}(X) = C^0(|X|, T) .$$

**Definition 10.12.** *Let  $G$  be a locally profinite group. A  $\underline{G}$ -torsor is map  $f : \tilde{X} \rightarrow X$  of v-stacks with an action of  $\underline{G}$  on  $\tilde{X}$  over  $X$ , such that v-locally on  $X$ , there is a  $\underline{G}$ -equivariant isomorphism  $\tilde{X} \cong \underline{G} \times X$ .*

**Lemma 10.13.** *Let  $G$  be a locally profinite group,  $X$  a perfectoid space, and  $f : \tilde{X} \rightarrow X$  a  $\underline{G}$ -torsor. Then the v-sheaf  $\tilde{X}$  is representable by a perfectoid space, and  $\tilde{X} \rightarrow X$  is pro-étale, universally open and a v-cover. More precisely, for any open subgroup  $K \subset G$ , pushout along the discrete  $G$ -set  $G/K$  defines a separated étale map*

$$\tilde{X}_K := \tilde{X} \times_{\underline{G}} \underline{G/K} \rightarrow X ,$$

the transition map  $\tilde{X}_{K'} \rightarrow \tilde{X}_K$  is finite étale if  $K' \subset K$  is of finite index, and

$$\tilde{X} = \varprojlim_K \tilde{X}_K \rightarrow X .$$

Related results had been obtained by Hansen, [Han16].

*Proof.* In case  $\tilde{X} = \underline{G} \times X$ , the results are clear, as then  $\tilde{X}_K = \underline{G/K} \times X$  is a disjoint union of copies of  $X$ . Thus, Proposition 10.11 (iv) implies that  $\tilde{X}_K \rightarrow X$  is representable by a separated étale map in general, and Proposition 10.11 (iii) implies that  $\tilde{X}_{K'} \rightarrow \tilde{X}_K$  is finite étale in case  $K' \subset K$  is of finite index. Also,  $\tilde{X} = \varprojlim_K \tilde{X}_K$ , as this can be checked v-locally. This presentation shows that  $\tilde{X} \rightarrow X$  is pro-étale, as desired. It is also a v-cover, as this can be checked v-locally, where it admits a section. Finally, the map  $\tilde{X} \rightarrow X$  is universally open, as any open subspace comes via pullback from  $\tilde{X}_K$  for sufficiently small  $K$ , the map  $\tilde{X} \rightarrow \tilde{X}_K$  is surjective, and  $\tilde{X}_K \rightarrow X$  is open (as it is étale).  $\square$

## 11. DIAMONDS

From now on, we change the setup, and work with the full subcategory  $\text{Perf} \subset \text{Perfd}$  of ( $\kappa$ -small) perfectoid spaces of characteristic  $p$ .

**Definition 11.1.** *A diamond is a sheaf  $Y$  for the pro-étale topology of  $\text{Perf}$  such that  $Y$  can be written as a quotient  $X/R$ , where  $X$  is representable by a perfectoid space, and  $R \subset X \times X$  is a representable equivalence relation for which the projections  $s, t : R \rightarrow X$  are pro-étale.*

For convenience, we introduce the following name for the equivalence relations.

**Definition 11.2.** *Let  $X$  be a perfectoid space. A pro-étale equivalence relation on  $X$  is a representable equivalence relation  $R \subset X \times X$  such that the projections  $s, t : R \rightarrow X$  are pro-étale.*

Note that we do not make any assumptions like representability of the diagonal; the issue is that there is no good notion of morphisms that are relatively representable (in perfectoid spaces).

**Proposition 11.3.** *Let  $X \in \text{Perf}$ , and  $R \subset X \times X$  a pro-étale equivalence relation.*

- (i) *The quotient sheaf  $Y = X/R$  is a diamond.*
- (ii) *The natural map  $R \rightarrow X \times_Y X$  of sheaves on  $\text{Perf}$  is an isomorphism.*
- (iii) *Let  $\tilde{X} \rightarrow X$  be a pro-étale cover by a perfectoid space  $\tilde{X}$ , and*

$$\tilde{R} = R \times_{X \times X} (\tilde{X} \times \tilde{X}) \subset \tilde{X} \times \tilde{X}$$

*the induced equivalence relation. Then  $\tilde{R}$  is a pro-étale equivalence relation on  $\tilde{X}$ , and the natural map  $\tilde{Y} := \tilde{X}/\tilde{R} \rightarrow Y = X/R$  is an isomorphism.*

- (iv) *The map  $X \rightarrow Y$  is quasi-pro-étale.*

Recall that quasi-pro-étale morphisms are defined in Definition 10.1 (i).

*Proof.* Part (i) is clear by definition. For part (ii), note that both are subsheaves of  $X \times X$ , so  $R \rightarrow X \times_Y X$  is injective. On the other hand, let  $Z$  be any perfectoid space with a map  $Z \rightarrow X \times_Y X$ . By definition, this means that we have two maps  $a, b : Z \rightarrow X$  such that the two induced maps  $Z \rightarrow Y$  agree. This condition means that after some pro-étale cover  $\tilde{Z} \rightarrow Z$ , the composite map  $\tilde{Z} \rightarrow Z \rightarrow X \times X$  factors over the equivalence relation  $R$ . We get a map  $\tilde{Z} \rightarrow R$ , which descends to a map  $Z \rightarrow R$  as the two maps  $\tilde{Z} \times_Z \tilde{Z} \rightarrow R$  coming from projection on either factor agree: Namely, their composites with  $R \hookrightarrow X \times X$  agree. We have produced a map  $Z \rightarrow R$ , which factors the map  $Z \rightarrow X \times X$ , as this holds true after composition with the pro-étale cover  $\tilde{Z} \rightarrow Z$ . Therefore,  $R \rightarrow X \times_Y X$  is surjective, as desired.

In part (iii), first note that  $\tilde{R}$  is representable as a fibre product of representable objects, and the two projections  $\tilde{s}, \tilde{t} : \tilde{R} \rightarrow \tilde{X}$  are pro-étale: For example, one can write  $\tilde{s}$  as the composite of

$$(R \times_X \tilde{X}) \times_X (\tilde{X} \rightarrow X) : \tilde{R} = R \times_{X \times X} (\tilde{X} \times \tilde{X}) \rightarrow R \times_{X \times X} (\tilde{X} \times X) = R \times_X \tilde{X}$$

and

$$s \times_X \tilde{X} : R \times_X \tilde{X} \rightarrow \tilde{X},$$

both of which are a base change of a pro-étale map, and thus pro-étale.

Now the map  $\tilde{Y} \rightarrow Y$  of pro-étale sheaves on  $\text{Perf}$  is surjective, as the composite  $\tilde{X} \rightarrow \tilde{Y} \rightarrow Y$  is, as this can also be written as the composite  $\tilde{X} \rightarrow X \rightarrow Y$ . To check whether  $\tilde{Y} \rightarrow Y$  is injective, let  $Z$  be a perfectoid space with two maps  $a, b : Z \rightarrow \tilde{Y}$  which agree after composing with  $\tilde{Y} \rightarrow Y$ . Replacing  $Z$  by a pro-étale cover, we may assume that  $a, b$  factor over  $\tilde{a}, \tilde{b} : Z \rightarrow \tilde{X}$ . The associated map  $Z \rightarrow \tilde{X} \times \tilde{X} \rightarrow X \times X$  factors over  $R$  by assumption and part (ii), so we get a map  $Z \rightarrow R \times_{X \times X} (\tilde{X} \times \tilde{X}) = \tilde{R}$ . This means that  $\tilde{a}, \tilde{b} : Z \rightarrow \tilde{X}$  induce the same map  $Z \rightarrow \tilde{Y}$ , proving injectivity of  $\tilde{Y} \rightarrow Y$ , thereby finishing the proof of part (iii).

For part (iv), we can replace  $X$  by  $\tilde{X} = \bigsqcup_i U_i \rightarrow X$ , where the  $U_i$  form an affinoid cover of  $X$  and  $R$  by the induced equivalence relation  $\tilde{R} \subset \tilde{X} \times \tilde{X}$ : Indeed, these spaces still satisfy the same assumptions by part (iii), and to check whether  $X' \times_Y X$  is representable and pro-étale over  $X'$ , it is enough to check for the open subspaces  $X' \times_Y U_i$ . In particular, after this replacement,  $s, t : R \rightarrow X$  are separated: Indeed, both maps  $R \subset X \times X$  and  $X \times X \rightarrow X$  are separated.

As by definition  $X \rightarrow Y$  is surjective in the pro-étale topology, we can find a pro-étale cover  $\widetilde{X}' \rightarrow X'$  and a map  $\widetilde{X}' \rightarrow X$  lying over  $X' \rightarrow Y$ . We can assume that  $\widetilde{X}'$  is affinoid. Let  $W = X' \times_Y X \rightarrow X'$  be the fibre product. Then

$$\widetilde{X}' \times_{X'} W = \widetilde{X}' \times_X (X \times_Y X) = \widetilde{X}' \times_X R,$$

which is representable and pro-étale over  $\widetilde{X}'$ . Moreover, as  $s, t : R \rightarrow X$  are separated, also the base change  $\widetilde{X}' \times_{X'} W = \widetilde{X}' \times_X R \rightarrow \widetilde{X}'$  is separated. Now applying Proposition 9.6, we see that  $W \rightarrow X'$  is representable, pro-étale and separated over  $X'$ .  $\square$

Another basic fact is the following.

**Proposition 11.4.** *Fibre products exist in the category of diamonds.*

*Proof.* Let  $Y_1 \rightarrow Y_3 \leftarrow Y_2$  be a diagram of diamonds. Choose presentations  $Y_i = X_i/R_i$  as in the definition of a diamond, for  $i = 1, 2, 3$ . After replacing  $X_i$  for  $i = 1, 2$  by a pro-étale cover, we can assume that there are maps  $X_i \rightarrow X_3$  lying over  $Y_i \rightarrow Y_3$ . We can then further replace  $X_i$  by  $X_i \times_{Y_3} X_3 = X_i \times_{X_3} R_3$  to assume that the maps  $X_i \rightarrow Y_1 \times_{Y_3} X_3$  are surjective in the pro-étale topology.

In this case, the map  $X_4 := X_1 \times_{X_3} X_2 \rightarrow Y_1 \times_{Y_3} Y_2 =: Y_4$  is surjective in the pro-étale topology. The induced equivalence relation  $R_4 = X_4 \times_{Y_4} X_4$  can be calculated as  $R_4 = R_1 \times_{R_3} R_2$ , which is representable. It remains to see that  $R_4 \rightarrow X_4$  is pro-étale. But  $R_1 \rightarrow X_1$  and  $R_2 \rightarrow X_2$  are pro-étale, so  $R_1 \times R_2 \rightarrow X_1 \times X_2$  and its base change  $R_1 \times_{X_3} R_2 \rightarrow X_1 \times_{X_3} X_2 = X_4$  are pro-étale. It remains to see that  $R_4 = R_1 \times_{R_3} R_2 \rightarrow R_1 \times_{X_3} R_2$  is pro-étale; this is a base change of  $R_3 \rightarrow R_3 \times_{X_3} R_3$ . But the diagonal of any representable map is pro-étale by Remark 7.9.  $\square$

Another characterization of diamonds is given as follows.

**Proposition 11.5.** *Let  $Y$  be a pro-étale sheaf on  $\text{Perf}$ . Then  $Y$  is a diamond if and only if there is a surjective quasi-pro-étale morphism  $X \rightarrow Y$  from a perfectoid space  $X$ . If  $X$  is a disjoint union of strictly totally disconnected spaces, then  $R = X \times_Y X \subset X \times X$  is a pro-étale equivalence relation with  $Y = X/R$ .*

*Proof.* We may assume that  $X$  is a disjoint union of strictly totally disconnected spaces. In this case, by definition of quasi-pro-étale morphisms (Definition 10.1 (i)),  $R = X \times_Y X \rightarrow X$  is pro-étale. As in the proof of Proposition 11.3 (iii), one finds that  $Y = X/R$ , so  $Y$  is a diamond.  $\square$

One can use this to obtain some stability properties for diamonds.

**Proposition 11.6.** *Let  $Y$  be a pro-étale sheaf on  $\text{Perf}$ , and assume that there is a surjective quasi-pro-étale map  $Y' \rightarrow Y$ , where  $Y'$  is a diamond. Then  $Y$  is a diamond.*

*Proof.* Choose a surjective quasi-pro-étale map  $X \rightarrow Y'$ . Then  $X \rightarrow Y' \rightarrow Y$  is a surjective quasi-pro-étale map, so  $Y$  is a diamond by Proposition 11.5.  $\square$

**Proposition 11.7.** *Let  $f : Y' \rightarrow Y$  be a quasi-pro-étale map of pro-étale sheaves on  $\text{Perf}$ , and assume that  $Y$  is a diamond. Then  $Y'$  is a diamond.*

*Proof.* Choose a surjective quasi-pro-étale map  $X \rightarrow Y$  where  $X$  is a disjoint union of strictly totally disconnected spaces. Then  $X' = Y' \times_Y X$  is representable (and pro-étale over  $X$ ), and  $X' \rightarrow Y'$  is a surjective quasi-pro-étale map. Now Proposition 11.5 implies that  $Y'$  is a diamond.  $\square$

**Proposition 11.8.** *Let  $X$  be a diamond, and let  $R \subset X \times X$  be an equivalence relation such that  $s, t : R \rightarrow X$  are quasi-pro-étale. Then the quotient  $Y = X/R$  is a diamond.*

In the statement,  $R$  is just assumed to be a pro-étale sheaf, but note that as  $s : R \rightarrow X$  is quasi-pro-étale, it follows from Proposition 11.7 that  $R$  is automatically a diamond.

*Proof.* We may assume that  $X$  is a separated perfectoid space. In that case, the map  $X \rightarrow Y$  is a surjective separated map which is quasi-pro-étale: Indeed, its base change  $R = X \times_Y X \rightarrow X$  is quasi-pro-étale, and  $X \rightarrow Y$  is a v-cover, so we can apply Proposition 10.11 (v). Thus, the result follows from Proposition 11.6.  $\square$

A nice property of diamonds is that they are always sheaves for the v-topology. This resembles a result of Gabber, [Sta, Tag 0APL], that all algebraic spaces are sheaves for the fpqc topology, and our proof will follow his arguments.

**Proposition 11.9.** *Let  $Y$  be a diamond. Then  $Y$  is a sheaf for the v-topology.*

*Proof.* Choose a presentation  $Y = X/R$  of  $Y$  as a quotient of a perfectoid space  $X$  by a pro-étale equivalence relation  $R \subset X \times X$ . By Proposition 11.3 (iii), we may assume that  $X$  is a disjoint union of totally disconnected (in particular, affinoid) perfectoid spaces.

Let  $Z$  be a perfectoid space with a v-cover  $\tilde{Z} \rightarrow Z$ . First, we prove injectivity of  $Y(Z) \rightarrow Y(\tilde{Z})$ . Thus, assume that  $a, b : Z \rightarrow Y$  are two maps such that the composite maps  $\tilde{a}, \tilde{b} : \tilde{Z} \rightarrow Z \rightarrow Y$  agree. After replacing  $Z$  by a pro-étale cover (and using that  $Y$  is a pro-étale sheaf), we can assume that  $a, b$  lift to maps  $a_X, b_X : Z \rightarrow X$ ; then the composite maps  $\tilde{a}_X, \tilde{b}_X : \tilde{Z} \rightarrow Z \rightarrow X$  have the property that  $(\tilde{a}_X, \tilde{b}_X) : \tilde{Z} \rightarrow X \times X$  factors over  $R \subset X \times X$ . In other words, the map  $(a_X, b_X) : Z \rightarrow X \times X$  factors over  $R$  after precomposition with  $\tilde{Z} \rightarrow Z$ . As  $R$  is a v-sheaf, this implies that  $(a_X, b_X) : Z \rightarrow X \times X$  factors over  $R$ , showing that  $a = b$ , as desired.

It remains to prove surjectivity of

$$Y(Z) \rightarrow \text{eq}(Y(\tilde{Z}) \rightrightarrows Y(\tilde{Z} \times_Z \tilde{Z})) .$$

By standard reductions (using that  $F$  is a pro-étale sheaf and separated for the v-topology), we may assume that  $Z$  and  $\tilde{Z}$  are strictly totally disconnected. Let  $\tilde{a} : \tilde{Z} \rightarrow Y$  be a map such that the two induced maps  $\tilde{Z} \times_Z \tilde{Z} \rightarrow Y$  agree. By Proposition 11.3 (iv), the fibre product  $\tilde{Z} \times_Y X \rightarrow \tilde{Z}$  is representable and pro-étale; moreover, by our assumption on  $X$ , it is also separated over  $\tilde{Z}$ . Note that  $\tilde{W} = \tilde{Z} \times_Y X \rightarrow \tilde{Z}$  comes with a descent datum relative to  $\tilde{Z}/Z$ : Indeed, the fibre product

$$\tilde{W} \times_Z \tilde{Z} = (\tilde{Z} \times_Z \tilde{Z}) \times_Y X$$

is independent of which of the two induced maps  $\tilde{Z} \times_Z \tilde{Z} \rightarrow Y$  is chosen, as they agree by assumption. By Proposition 9.6, it follows that  $\tilde{W} \rightarrow \tilde{Z}$  descends to a surjective separated pro-étale map  $W \rightarrow Z$ . In particular,  $\tilde{W} = W \times_Z \tilde{Z} \rightarrow W$  is a v-cover, and the map  $\tilde{W} \rightarrow X$  descends to  $W \rightarrow X$  by construction of the descent datum, and using that  $X$  is a v-sheaf. We have constructed a pro-étale cover  $W \rightarrow Z$  with a map  $W \rightarrow X$ . The induced map  $W \times_Z W \rightarrow X \times X$  factors over  $R \subset X \times X$ , as this is true for the v-cover

$$(W \times_Z W) \times_Z \tilde{Z} = \tilde{W} \times_{\tilde{Z}} \tilde{W} = \tilde{Z} \times_Y X$$

of  $W \times_Z W$ . Thus, the map composite map  $W \rightarrow X \rightarrow Y$  factors over a map  $Z \rightarrow Y$ , which pulls back to the given map  $\tilde{Z} \rightarrow Y$ .  $\square$

Another interesting stability property is the following.

**Proposition 11.10.** *Let  $f : Y' \rightarrow Y$  be an injection of  $v$ -sheaves, where  $Y$  is a diamond. Then  $Y'$  is a diamond.*

*Proof.* We have to find a surjective quasi-pro-étale morphism  $X' \rightarrow Y'$ . Pulling back via a surjective quasi-pro-étale morphism  $X \rightarrow Y$ , we may assume that  $Y$  is a disjoint union of totally disconnected spaces; we may replace  $Y$  by a totally disconnected space  $X$ . In that case, Proposition 10.5 says that  $Y'$  is a filtered colimit of pro-constructible generalizing subsets  $X_i \subset X$ . Let  $X' = \bigsqcup_i X_i$ , with its natural morphism to  $Y'$ . Then  $X' \rightarrow Y'$  is surjective, and quasi-pro-étale (as each map  $X_i \rightarrow Y'$  is quasi-pro-étale).  $\square$

A convenient property of diamonds is that it is easy to check whether a map is an isomorphism. For an even more general statement, see Lemma 12.5 below.

**Lemma 11.11.** *Let  $f : Y \rightarrow X$  be a qcqs map of diamonds. Then  $f$  is an isomorphism if and only if for every algebraically closed perfectoid field  $K$  with open and bounded valuation subring  $K^+ \subset K$ , the map  $f(K, K^+) : Y(K, K^+) \rightarrow X(K, K^+)$  is a bijection.*

*Proof.* Pulling back by some presentation of  $X$ , we may replace  $X$  by a perfectoid space, which we can then assume to be affinoid. In that case,  $Y$  is a qcqs diamond. Write  $Y = \tilde{Y}/R$  as the quotient of a qcqs perfectoid space  $\tilde{Y}$  by a pro-étale equivalence relation  $R$ , which is still qcqs. The map  $\tilde{Y} \rightarrow X$  is surjective in the  $v$ -topology, as it is a map of qcqs perfectoid spaces which by assumption is surjective on points, i.e. a  $v$ -cover. As both  $Y$  and  $X$  are  $v$ -sheaves by Proposition 11.9, it suffices to see that the map  $R \rightarrow \tilde{Y} \times_X \tilde{Y}$  is an isomorphism. Now both  $R$  and  $\tilde{Y} \times_X \tilde{Y}$  are qcqs perfectoid spaces. Moreover, the map  $R(K, K^+) \rightarrow (\tilde{Y} \times_X \tilde{Y})(K, K^+)$  is a bijection for all  $(K, K^+)$  as in the statement of the lemma. Thus, the result follows from Lemma 5.4.  $\square$

Let us give a slightly weird example of diamonds.

**Example 11.12.** Fix a perfectoid field  $K$  of characteristic  $p$ . We claim that there is a fully faithful functor from the category of ( $\kappa$ -small) compact Hausdorff spaces to the category of diamonds over  $\mathrm{Spa}(K, \mathcal{O}_K)$ . Indeed, for any compact Hausdorff space  $T$ , one has the functor  $\underline{T}$  on  $\mathrm{Perf}$  which takes a perfectoid space  $X$  to  $C^0(|X|, T)$ , where  $C^0$  denotes the continuous functions. We claim that the functor

$$T \mapsto \underline{T} \times \mathrm{Spa}(K, \mathcal{O}_K)$$

is a fully faithful functor from the category of compact Hausdorff spaces to the category of diamonds over  $\mathrm{Spa}(K, \mathcal{O}_K)$ .

Let us check first that it takes values in diamonds. If  $S$  is a profinite set, then we can write  $S$  as an inverse limit of finite sets  $S_i$ , and if we also denote by  $S_i$  the corresponding constant sheaf on  $\mathrm{Perf}$ , then  $\underline{S} = \varprojlim S_i$ , and  $\underline{S} \times \mathrm{Spa}(K, \mathcal{O}_K) = \varprojlim (S_i \times \mathrm{Spa}(K, \mathcal{O}_K))$  is an affinoid perfectoid space, given by

$$\underline{S} \times \mathrm{Spa}(K, \mathcal{O}_K) = \mathrm{Spa}(C^0(S, K), C^0(S, \mathcal{O}_K)) .$$

Thus, profinite sets are mapped to affinoid perfectoid spaces. Now recall that any compact Hausdorff space  $T$  admits a surjection  $S \rightarrow T$  from a profinite set  $S$ : One can take for  $S$  the Stone-Cech compactification of  $T$  considered as a discrete set. The induced equivalence relation

$R = S \times_T S \subset S \times S$  is a closed subspace of the profinite set  $S \times S$ , and thus a profinite set itself. Now one can form the equivalence relation

$$\underline{R} \times \mathrm{Spa}(K, \mathcal{O}_K) \subset (\underline{S} \times \mathrm{Spa}(K, \mathcal{O}_K)) \times_{\mathrm{Spa}(K, \mathcal{O}_K)} (\underline{S} \times \mathrm{Spa}(K, \mathcal{O}_K)) ,$$

which is representable and pro-étale over  $\underline{S} \times \mathrm{Spa}(K, \mathcal{O}_K)$ . Let  $Y$  be the quotient  $\underline{S} \times \mathrm{Spa}(K, \mathcal{O}_K) / \underline{R} \times \mathrm{Spa}(K, \mathcal{O}_K)$ . We claim that  $Y = \underline{T} \times \mathrm{Spa}(K, \mathcal{O}_K)$ . Certainly, there is a natural map  $Y \rightarrow \underline{T} \times \mathrm{Spa}(K, \mathcal{O}_K)$ . It is injective: If we have two maps  $a, b : Z \rightarrow \underline{S} \times \mathrm{Spa}(K, \mathcal{O}_K)$  whose composites to  $\underline{T} \times \mathrm{Spa}(K, \mathcal{O}_K)$  agree, then we have two continuous maps  $|Z| \rightarrow S$  whose composite to  $T$  agree, i.e. a continuous map  $|Z| \rightarrow R = S \times_T S$ , or in other words a map  $Z \rightarrow \underline{R} \times \mathrm{Spa}(K, \mathcal{O}_K)$ . For surjectivity, let  $Z$  be any perfectoid space with a map  $Z \rightarrow \underline{T} \times \mathrm{Spa}(K, \mathcal{O}_K)$ . We want to find a pro-étale cover  $\tilde{Z} \rightarrow Z$  and a lift  $\tilde{Z} \rightarrow \underline{S} \times \mathrm{Spa}(K, \mathcal{O}_K)$ . We may assume that  $Z$  is affinoid. In that case any continuous map  $|Z| \rightarrow T$  to a compact Hausdorff space  $T$  factors through the profinite set  $\pi_0(Z)$  of connected components. We get an induced map  $\pi_0(Z) \rightarrow T$ , and a cover by the profinite set  $\pi_0(Z) \times_T S \subset \pi_0(Z) \times S \rightarrow \pi_0(Z)$ . It follows that

$$\tilde{Z} = Z \times_{\pi_0(Z)} (\pi_0(Z) \times_T S)$$

is affinoid perfectoid, pro-étale over  $Z$ , and admits a lift  $|\tilde{Z}| \rightarrow S$  of  $|Z| \rightarrow T$ , as desired.

Finally, to check that  $T \mapsto \underline{T} \times \mathrm{Spa}(K, \mathcal{O}_K)$  is fully faithful, we need to prove that the map

$$C^0(T_1, T_2) \rightarrow \mathrm{Hom}_{\mathrm{Spa}(K, \mathcal{O}_K)}(\underline{T}_1 \times \mathrm{Spa}(K, \mathcal{O}_K), \underline{T}_2 \times \mathrm{Spa}(K, \mathcal{O}_K))$$

is bijective for all compact Hausdorff sets  $T_1, T_2$ . For injectivity, we can assume that  $T_1$  is a point. Now any map  $\mathrm{Spa}(K, \mathcal{O}_K) \rightarrow \underline{T}_2 \times \mathrm{Spa}(K, \mathcal{O}_K)$  is given by a map  $|\mathrm{Spa}(K, \mathcal{O}_K)| \rightarrow T_2$  whose image recovers the given point of  $T_2$ .

For surjectivity, assume first that  $T_1$  is profinite. In that case,  $\underline{T}_1 \times \mathrm{Spa}(K, \mathcal{O}_K)$  is representable, and so

$$\mathrm{Hom}_{\mathrm{Spa}(K, \mathcal{O}_K)}(\underline{T}_1 \times \mathrm{Spa}(K, \mathcal{O}_K), \underline{T}_2 \times \mathrm{Spa}(K, \mathcal{O}_K)) = \underline{T}_2(\underline{T}_1 \times \mathrm{Spa}(K, \mathcal{O}_K)) = C^0(T_1, T_2)$$

by definition, using that  $|\underline{T}_1 \times \mathrm{Spa}(K, \mathcal{O}_K)| = T_1$ .

Now choose a surjection  $S_1 \rightarrow T_1$  from a profinite set. Given any map  $\underline{T}_1 \times \mathrm{Spa}(K, \mathcal{O}_K) \rightarrow \underline{T}_2 \times \mathrm{Spa}(K, \mathcal{O}_K)$ , we get a composite map  $\underline{S}_1 \times \mathrm{Spa}(K, \mathcal{O}_K) \rightarrow \underline{T}_2 \times \mathrm{Spa}(K, \mathcal{O}_K)$ , which comes from a map  $S_1 \rightarrow T_2$  by what we have just shown. The two induced maps  $S_1 \times_{T_1} S_1 \rightarrow T_2$  agree by faithfulness. Thus, we get a map from the quotient  $T_1 \rightarrow T_2$  (recalling that any surjective map of compact Hausdorff spaces is a quotient map), which induces the given map  $\underline{T}_1 \times \mathrm{Spa}(K, \mathcal{O}_K) \rightarrow \underline{T}_2 \times \mathrm{Spa}(K, \mathcal{O}_K)$  (as this is true after precomposing with the pro-étale cover  $\underline{S}_1 \times \mathrm{Spa}(K, \mathcal{O}_K) \rightarrow \underline{T}_1 \times \mathrm{Spa}(K, \mathcal{O}_K)$ ).

Although compact Hausdorff spaces may have some appeal, we will generally work with spaces whose behaviour is closer to that of schemes. They are essentially characterized by having a spectral underlying topological space.

**Proposition 11.13.** *Let  $Y$  be a diamond, and let  $Y = X/R$  be a presentation as a quotient of a perfectoid space  $X$  by a pro-étale equivalence relation  $R \subset X \times X$ . There is a canonical bijection between  $|X|/|R|$  and the set*

$$|Y| = \{\mathrm{Spa}(K, K^+) \rightarrow Y\} / \sim ,$$



where  $(K, K^+)$  runs over all pairs of a perfectoid field  $K$  with an open and bounded valuation subring  $K^+ \subset K$ , and two such maps  $\mathrm{Spa}(K_i, K_i^+) \rightarrow Y$ ,  $i = 1, 2$ , are equivalent, if there is a third pair  $(K_3, K_3^+)$ , and a commutative diagram

$$\begin{array}{ccc} & \mathrm{Spa}(K_1, K_1^+) & \\ & \nearrow & \searrow \\ \mathrm{Spa}(K_3, K_3^+) & \longrightarrow & Y, \\ & \searrow & \nearrow \\ & \mathrm{Spa}(K_2, K_2^+) & \end{array}$$

where the maps  $\mathrm{Spa}(K_3, K_3^+) \rightarrow \mathrm{Spa}(K_i, K_i^+)$ ,  $i = 1, 2$ , are surjective.

Moreover, the quotient topology on  $|Y|$  induced by the surjection  $|X| \rightarrow |Y|$  is independent of the choice of presentation  $Y = X/R$ .

*Proof.* First, note that the given relation  $\sim$  on  $\{\mathrm{Spa}(K, K^+) \rightarrow Y\}$  is an equivalence relation. This follows from the observation that for a diagram  $(K_1, K_1^+) \leftarrow (K_0, K_0^+) \rightarrow (K_2, K_2^+)$  of perfectoid fields with open and bounded valuation subrings for which  $\mathrm{Spa}(K_i, K_i^+) \rightarrow \mathrm{Spa}(K_0, K_0^+)$  is surjective for  $i = 1, 2$ , one can find another such pair  $(K_3, K_3^+)$  sitting in a diagram

$$\begin{array}{ccc} & \mathrm{Spa}(K_1, K_1^+) & \\ & \nearrow & \searrow \\ \mathrm{Spa}(K_3, K_3^+) & \longrightarrow & \mathrm{Spa}(K_0, K_0^+), \\ & \searrow & \nearrow \\ & \mathrm{Spa}(K_2, K_2^+) & \end{array}$$

with all transition maps surjective. Indeed, this follows from surjectivity of

$$|\mathrm{Spa}(K_1, K_1^+) \times_{\mathrm{Spa}(K_0, K_0^+)} \mathrm{Spa}(K_2, K_2^+)| \rightarrow |\mathrm{Spa}(K_1, K_1^+)| \times_{|\mathrm{Spa}(K_0, K_0^+)|} |\mathrm{Spa}(K_2, K_2^+)|.$$

There is a canonical map  $|X| \rightarrow |Y|$  by first interpreting  $|X|$  as the similar set of equivalence classes of maps  $\mathrm{Spa}(K, K^+) \rightarrow X$ , which is a standard fact. This map is surjective. Indeed, given a map  $\mathrm{Spa}(K, K^+) \rightarrow Y$ , one can lift it to  $X$  after a pro-étale cover. Thus, after increasing  $K$ , we can lift it to a map  $\mathrm{Spa}(K, K^+) \rightarrow X$ , as desired. One checks directly that this map  $|X| \rightarrow |Y|$  factors over  $|X|/|R|$ .

Now assume that two maps  $\mathrm{Spa}(K_i, K_i^+) \rightarrow X$ ,  $i = 1, 2$ , project to the same point of  $|Y|$ . By definition of  $|Y|$ , this means that up to enlarging  $K_1$  and  $K_2$ , we can assume that  $K = K_1 = K_2$ ,  $K^+ = K_1^+ = K_2^+$ , and the composite maps  $\mathrm{Spa}(K, K^+) \rightrightarrows X \rightarrow Y$  agree. In other words, we get a map  $\mathrm{Spa}(K, K^+) \rightarrow R = X \times_Y X$ , proving that  $|X|/|R| \rightarrow |Y|$  is injective, finishing the proof that  $|X|/|R|$  maps bijectively to  $|Y|$ .

Finally, to see that the quotient topology on  $|Y|$  is independent of the presentation, choose another presentation  $Y = X'/R'$ . Then  $X \times_Y X'$  is a diamond, so we can find a pro-étale surjection  $X'' \rightarrow X \times_Y X'$  from a perfectoid space  $X''$ . Now  $|X''| \rightarrow |X|$  and  $|X''| \rightarrow |X'|$  are quotient maps

by Lemma 2.5, so the quotient topologies on  $|Y|$  induced by the surjection from  $|X|$ ,  $|X''|$  and  $|X'|$  are equivalent.  $\square$

In particular, the following definition makes sense.

**Definition 11.14.** *Let  $Y$  be a diamond, and write  $Y = X/R$  as the quotient of a perfectoid space  $X$  by a pro-étale equivalence relation  $R \subset X \times X$ . The underlying topological space of  $|Y|$  is the quotient space  $|X|/|R|$ .*

It is easy to see that this construction is functorial in  $Y$ . One can read off the open subspaces of a diamond  $Y$  on the topological space  $|Y|$ .

**Proposition 11.15.** *Let  $Y$  be a diamond with underlying topological space  $|Y|$ . Any open subfunctor of  $Y$  is a diamond. Moreover, there is bijective correspondence between open immersions  $U \subset Y$  and open subsets  $V \subset |Y|$ , given by sending  $U$  to  $V = |U|$ , and  $V$  to the subfunctor  $U \subset Y$  of those maps  $X \rightarrow Y$  for which  $|X| \rightarrow |Y|$  factors over  $V \subset |Y|$ .*

*Moreover, if  $f : Y' \rightarrow Y$  is a map of diamonds which is a surjection of v-sheaves, then  $|f| : |Y'| \rightarrow |Y|$  is a quotient map.*

*Proof.* Write  $Y = X/R$  as usual. Given an open immersion  $U \subset Y$ , one gets an open subspace  $U_X \subset X$  which is stable under the equivalence relation  $R$ ; then  $U = U_X/(R \cap (U_X \times U_X))$ , where  $R \cap (U_X \times U_X) \subset U_X \times U_X$  is a pro-étale equivalence relation, so that  $U$  is a diamond.

In particular, in this situation  $|U_X| \subset |X|$  is an open  $|R|$ -invariant subspace of  $|X|$ , giving rise to the open subspace  $V = |U| \subset |X|$ . Conversely, if  $V \subset |Y|$  is open, then the preimage  $V_X \subset |X|$  is an open  $|R|$ -invariant subset, giving rise to an open  $R$ -invariant subspace  $U_X \subset X$ , whose quotient  $U_X/R \subset Y$  defines an open immersion. One checks easily that the two processes are inverse.

The final statement follows from the fact that the prestack sending a v-sheaf  $Y$  to its open subsheaves is a v-stack, which follows from Proposition 10.11 (i).  $\square$

**Example 11.16.** If  $Y$  is a perfectoid space, then  $|Y|$  is the usual underlying topological space of  $Y$ . In the context of Example 11.12, if  $T$  is a compact Hausdorff space and  $K$  is a perfectoid field, then

$$|\underline{T} \times \mathrm{Spa}(K, \mathcal{O}_K)| = T .$$

In particular, the example shows that in general  $|Y|$  can be quite far from a spectral space.

**Definition 11.17.** *Let  $Y$  be a diamond. Then  $Y$  is spatial if  $Y$  is quasicompact and quasiseparated, and  $|Y|$  admits a basis of open subsets given by  $|U|$  for quasicompact open immersions  $U \subset Y$ . More generally,  $Y$  is locally spatial if  $Y$  admits an open cover by spatial diamonds.*

We note that any perfectoid space is locally spatial, and it is spatial precisely when it is qcqs.

**Proposition 11.18.** *Let  $Y$  be a spatial diamond.*

- (i) *The underlying topological space  $|Y|$  is spectral.*
- (ii) *Any quasicompact open subfunctor  $U \subset Y$  is spatial.*
- (iii) *For any perfectoid space  $X'$  with a map  $X' \rightarrow Y$ , the induced map of locally spectral spaces  $|X'| \rightarrow |Y|$  is spectral and generalizing.*

*Proof.* For part (i), first note that  $|Y|$  is quasicompact, as by Proposition 11.15, any open cover of  $|Y|$  determines an open cover of  $Y$ . Note that the set of quasicompact open immersions  $U \subset Y$  is stable under finite intersections (as  $Y$  is quasiseparated); it follows that the set of  $|U| \subset |Y|$  for  $U$  running over quasicompact open immersions forms a basis of quasicompact open subsets stable under finite intersections. It remains to check that  $|Y|$  is sober, i.e. every irreducible closed subset has a unique generic point. Uniqueness holds because  $|Y|$  is  $T_0$  by Lemma 2.7. For existence of generic points, write  $Y = X/R$ , where  $X$  is a qcqs perfectoid space and  $R$  is a pro-étale equivalence relation, so that  $|X| \rightarrow |Y|$  is a surjective spectral map of spectral spaces. Let  $Z \subset |Y|$  be an irreducible closed subset, with preimage  $W \subset |X|$ . It is enough to see that the intersection of all nonempty quasicompact open subsets of  $Z$  is nonempty, as any point in the intersection is a generic point for  $Z$ . But the preimages of quasicompact open subsets of  $Z$  are quasicompact open subsets of  $W$ , and by quasicompactness of the constructible topology on  $W$ , their intersection is nonempty; thus, the intersection is nonempty in  $Z$ , as desired.

In (ii), any quasicompact open subfunctor  $U \subset Y$  is given by a quasicompact open subset  $|U| \subset |Y|$ . As  $Y$  is quasiseparated,  $U$  is quasiseparated. Moreover, if  $V \subset Y$  is a quasicompact open subfunctor, then  $V \times_Y U \subset U$  is a quasicompact open subfunctor. Now the  $|V \times_Y U| \subset |U|$  form a basis of open subsets, verifying that  $U$  is spatial.

For part (iii), we may assume that  $X'$  is qcqs, in which case  $|X'|$  and  $|Y|$  are spectral. We need to check that the preimage of a quasicompact open is quasicompact. If  $V \subset |Y|$  is a quasicompact open, then it is covered by finitely many  $|U_i| \subset |Y|$  for quasicompact open  $U_i \subset Y$ , so it remains to check that  $X' \times_Y U_i \subset X'$  is quasicompact. But this follows from the assumption that  $Y$  is quasiseparated. To see that  $|X'| \rightarrow |Y|$  is generalizing, write  $Y = X/R$  as usual, with  $X$  strictly totally disconnected. Then we can replace  $X'$  by  $X' \times_Y X$ , and assume that  $X' \rightarrow Y$  lifts to  $X' \rightarrow X$ . Now  $|X'| \rightarrow |X|$  is generalizing, so it remains to see that  $|X| \rightarrow |Y|$  is generalizing. But this follows from Lemma 2.9.  $\square$

Let us note that some of these properties extend to locally spatial diamonds.

**Proposition 11.19.** *Let  $Y$  be a locally spatial diamond.*

- (i) *The underlying topological space  $|Y|$  is locally spectral.*
- (ii) *Any open subfunctor  $U \subset Y$  is locally spatial.*
- (iii) *The functor  $Y$  is quasicompact (resp. quasiseparated) if and only if  $|Y|$  is quasicompact (resp. quasiseparated).*
- (iv) *For any locally spatial diamond  $Y'$  with a map  $Y' \rightarrow Y$ , the induced map of locally spectral spaces  $|Y'| \rightarrow |Y|$  is spectral and generalizing.*

*Proof.* Left as an exercise to the reader.  $\square$

We need some permanence properties of the class of (locally) spatial diamonds.

**Proposition 11.20.** *Let  $f : Y' \rightarrow Y$  be a quasicompact injection of  $v$ -sheaves, where  $Y$  is a locally spatial diamond. Then  $Y'$  is a locally spatial diamond,  $|Y'| \subset |Y|$  is a pro-constructible and generalizing subset (with the subspace topology), and the map*

$$Y' \rightarrow Y \times_{|Y|} |Y'|$$

*of  $v$ -sheaves is an isomorphism.*

*Proof.* We may assume that  $Y$  is spatial. Let  $X \rightarrow Y$  be a surjective quasi-pro-étale map where  $X$  is a strictly totally disconnected perfectoid space. By Corollary 10.6,  $X' = X \times_Y Y'$  is a pro-constructible and generalizing subspace of  $X$ ; in particular  $X' = X \times_{|X|} |X'|$ . The spectral map  $|X| \rightarrow |Y|$  of spectral spaces is generalizing, so the image  $V \subset |Y|$  of  $|X'|$  in  $|Y|$  is a pro-constructible and generalizing subset, with  $|X'| = |X| \times_{|Y|} V$ . Also, if  $R = X \times_Y X$  is the equivalence relation, which is affinoid pro-étale over  $X$ , then  $R' = R \times_Y Y'$  is a pro-constructible and generalizing subset of  $R$ . One has a homeomorphism  $|X'|/|R'| = V$ : Indeed, the map  $|X'|/|R'| \rightarrow V$  is a continuous bijective map, and  $|X'| \rightarrow V$  is a quotient map by Lemma 2.5. Thus,  $|Y'| = V$ , and any quasicompact open subset of  $V$  comes via pullback from a quasicompact open subset of  $|Y|$ , which corresponds to a quasicompact open subspace of  $Y$ , and pulls back to a quasicompact open subspace of  $Y'$ : Thus,  $Y'$  is spatial. The natural map  $Y' \rightarrow Y \times_{|Y|} |Y'|$  becomes an isomorphism after pullback along  $X \rightarrow Y$ . As it is a map of v-sheaves, we see that  $Y' = Y \times_{|Y|} |Y'|$ , as desired.  $\square$

**Lemma 11.21.** *Let  $Y$  be a (locally) spatial diamond, and let  $Y' \rightarrow Y$  be a finite étale map of pro-étale sheaves. Then  $Y'$  is a (locally) spatial diamond.*

*Proof.* We may assume that  $Y$  is spatial. By Proposition 11.7,  $Y'$  is a diamond. Choose a presentation  $Y = X/R$  as a quotient of an affinoid perfectoid space  $X$ , and let  $X' = X \times_Y Y'$ , which is finite étale over  $X$ . Let  $R$  and  $R'$  be the respective equivalence relations. We need to find enough quasicompact open subspaces of  $Y'$ . Pick any point  $y' \in |Y'|$  with image  $y \in |Y|$ . Let  $Y_y \subset Y$  be the localization of  $Y$  at  $y$ , i.e. the intersection of all quasicompact open subspaces containing  $y$ . For any qcqs perfectoid space  $Z$  over  $Y$ , let  $Z_y = Z \times_Y Y_y$ . Then quasicompact open subspaces of  $Z_y$  come via pullback from quasicompact open subspaces of  $Z \times_Y U$  for some quasicompact open subspace  $U$  of  $Y$ , and any two such extensions agree after shrinking  $U$ . By applying this to  $Z = X'$  and  $Z = R'$ , we find that any quasicompact open subspace of  $Y'_y$  extends to a quasicompact open subspace of  $Y'_U$  for some  $U$ . This reduces us to the case  $Y = Y_y$ .

Thus, we may assume that  $|Y|$  has a unique closed point. In this case, we can take  $X = \text{Spa}(C, C^+)$  for some algebraically closed field  $C$  with an open and bounded valuation subring  $C^+ \subset C$ . Then  $X'$  is a finite disjoint union of copies of  $X$ . This implies that any pro-constructible generalizing subset of  $X'$  is open. In particular, for any quasicompact open subset  $V \subset X'$ , the  $R'$ -invariant subset  $R' \cdot V \subset X'$  is open (as it is pro-constructible and generalizing). This descends to a quasicompact open subspace of  $Y'$ , and these give a basis, as desired.  $\square$

**Lemma 11.22.** *Let  $Y_i$ ,  $i \in I$ , be a cofiltered inverse system of diamonds with qcqs transition maps, where  $I$  has cardinality less than  $\kappa$ . Assume that there is some strong limit cardinal  $\kappa' < \kappa$  and  $\kappa'$ -small perfectoid spaces  $X_i$  surjecting onto  $Y_i$ . Then  $Y = \varprojlim_i Y_i$  is a diamond with a bijective continuous map  $|Y| \rightarrow \varprojlim_i |Y_i|$ , and the maps  $Y \rightarrow Y_i$  are qcqs. If all  $Y_i$  are (locally) spatial, then  $Y$  is (locally) spatial, and  $|Y| \rightarrow \varprojlim_i |Y_i|$  is a homeomorphism.*

*Proof.* For the first part, it is enough to see that for any ordinal  $\lambda$  (less than  $\kappa$ ), if  $I$  is the category of ordinals  $\mu < \lambda$ , then the cofiltered limit along  $I$  exists. We can assume that  $Y_0$  (and then all  $Y_i$ ) are qcqs. Given  $\lambda$ , we fix  $\kappa_\lambda < \kappa$  a strong limit cardinal with cofinality larger than  $\lambda$ . We can also assume that  $\kappa_\lambda \geq \kappa'$ . In this situation, we will lift the diagram  $Y_\mu$ ,  $\mu < \lambda$ , to a diagram of  $\kappa_\lambda$ -small strictly totally disconnected spaces  $X_\mu$ ,  $\mu < \lambda$ , with compatible quasi-pro-étale surjections

$X_\mu \rightarrow Y_\mu$ ; in fact, with quasi-pro-étale surjections

$$X_\mu \rightarrow Y_\mu \times_{\varprojlim_{\mu' < \mu} Y_{\mu'}} \varprojlim_{\mu' < \mu} X_{\mu'} .$$

That this can be done follows easily by transfinite induction (noting that  $\varprojlim_{\mu' < \mu} X_{\mu'}$  is still  $\kappa_\lambda$ -small, as the cofinality of  $\kappa_\lambda$  is larger than  $\lambda$ ), and we note that one obtains a quasi-pro-étale surjection

$$\varprojlim_{\mu < \lambda} X_\mu \rightarrow \varprojlim_{\mu < \lambda} Y_\mu$$

in the limit. Indeed, after pullback to any strictly totally disconnected  $Z$ , this morphism is a transfinite composition of surjective affinoid pro-étale morphisms (using Lemma 7.19).

In this construction, one has

$$|Y| = \left| \varprojlim_{\mu < \lambda} X_\mu \right| / \left| \varprojlim_{\mu < \lambda} R_\mu \right| = \varprojlim_{\mu < \lambda} |X_\mu| / |R_\mu| = \varprojlim_{\mu < \lambda} |Y_\mu| ,$$

as sets, where  $R_\mu = X_\mu \times_{Y_\mu} X_\mu$ . If all  $Y_i$  are spatial, then Lemma 2.11 applied to the inverse system  $|Y_\mu|$  and the map from  $\left| \varprojlim_{\mu < \lambda} X_\mu \right|$  to the inverse limit shows that

$$\left| \varprojlim_{\mu < \lambda} X_\mu \right| \rightarrow \varprojlim_{\mu < \lambda} |Y_\mu|$$

is a quotient map, and thus  $|Y| \rightarrow \varprojlim_{\mu < \lambda} |Y_\mu|$  is a homeomorphism. Moreover, any quasicompact open subspace of  $|Y|$  comes via pullback from some quasicompact open subspace of  $|Y_i|$ , which corresponds to a quasicompact open subspace of  $Y_i$ , which in turn pulls back to a quasicompact open subspace of  $Y$ ; thus,  $Y$  is spatial. If the  $Y_i$  are just locally spatial, one can fix a spatial open subset of some  $Y_{i_0}$ , and taking the preimage gives a spatial open subset of  $Y$ ; also  $|Y| = \varprojlim_i |Y_i|$  follows.  $\square$

Moreover, in this situation, an analogue of Proposition 6.4 holds true.

**Proposition 11.23.** *Let  $Y_i$ ,  $i \in I$ , be a cofiltered inverse system of qcqs diamonds with set-theoretic bounds as in Lemma 11.22, with inverse limit  $Y = \varprojlim_i Y_i$ .*

(i) *Let  $Y_{\text{fét}}$  denote the category of finite étale diamonds over  $Y$ , and similarly for  $Y_i$ . The base change functors  $(Y_i)_{\text{fét}} \rightarrow Y_{\text{fét}}$  induce an equivalence of categories*

$$\mathcal{2}\text{-}\varprojlim_i (Y_i)_{\text{fét}} \rightarrow Y_{\text{fét}} .$$

(ii) *Let  $Y_{\text{ét, qcqs}}$  denote the category of (locally separated) étale qcqs diamonds over  $Y$ , and similarly for  $Y_i$ . The base change functors  $(Y_i)_{\text{ét, qcqs}} \rightarrow Y_{\text{ét, qcqs}}$  induce an equivalence of categories*

$$\mathcal{2}\text{-}\varprojlim_i (Y_i)_{\text{ét, qcqs}} \rightarrow Y_{\text{ét, qcqs}} .$$

(iii) *Let  $Y_{\text{ét, qc, sep}} \subset Y_{\text{ét, qcqs}}$  be the full subcategory of quasicompact separated étale diamonds over  $Y$  (and similarly for  $Y_i$ ). Then the base change functors  $(Y_i)_{\text{ét, qc, sep}} \rightarrow Y_{\text{ét, qc, sep}}$  induce an equivalence of categories*

$$\mathcal{2}\text{-}\varprojlim_i (Y_i)_{\text{ét, qc, sep}} \rightarrow Y_{\text{ét, qc, sep}} .$$

*Proof.* As in Lemma 11.22, we can assume that  $I$  is the set of ordinals  $\mu < \lambda$  for some fixed ordinal  $\lambda$ . We pick strictly totally disconnected perfectoid spaces  $X_\mu$  mapping compatibly to  $Y_\mu$  as in the proof of Lemma 11.22. Let  $R_\mu = X_\mu \times_{Y_\mu} X_\mu$  be the induced equivalence relations, and let  $X = \varprojlim_{\mu} X_\mu$ ,  $R = \varprojlim_{\mu} R_\mu$ .

Let  $F$  be the prestack on the category of diamonds of finite étale, resp. quasicompact separated étale morphisms. Then  $F$  is a stack for the  $v$ -topology by Proposition 10.11. Note that  $F(X/Y)$  involves the value of  $F$  on  $X$ ,  $R$ , and  $R \times_X R$ , all of which are affinoid perfectoid spaces, and similarly for  $F(X_\mu/Y_\mu)$ . By Proposition 6.4, we see that

$$F(Y) = F(X/Y) = 2\text{-}\varinjlim_{\mu < \lambda} F(X_\mu/Y_\mu) = 2\text{-}\varinjlim_{\mu < \lambda} F(Y_\mu) ,$$

proving parts (i) and (iii). For part (ii), essentially the same argument applies, except that a priori the functor  $F(Y) \rightarrow F(X/Y)$  is only fully faithful for the prestack  $F$  of qcqs étale morphisms. To check essential surjectivity in (ii), pick any étale qcqs map  $f : \tilde{Y} \rightarrow Y$ , where it is understood that  $f$  is locally separated. In particular, one can cover  $\tilde{Y}$  by quasicompact open subsets  $\tilde{Y}_j \subset \tilde{Y}$  such that  $f|_{\tilde{Y}_j}$  is separated. By (iii), this comes via pullback from some finite level. Moreover, the gluing data between the different  $\tilde{Y}_j$  will also be defined at some finite level, and the cocycle condition satisfied. This produces a (locally separated) étale qcqs map to some finite stage with pullback  $\tilde{Y}$ , as desired.  $\square$

To prove further permanence properties of spatial diamonds, we need the following result.

**Proposition 11.24.** *Let  $Y$  be a spatial diamond. Then one can find a strictly totally disconnected perfectoid space  $X$  with a surjective and universally open quasi-pro-étale map  $f : X \rightarrow Y$  that can be written as a cofiltered inverse limit of étale maps which are composites of quasicompact open immersions and finite étale maps.*

*Conversely, assume that  $Y$  is a qcqs diamond such that  $Y$  admits a surjective and universally open quasi-pro-étale map  $X \rightarrow Y$ , where  $X$  is a perfectoid space. Then  $Y$  is a spatial diamond.*

**Remark 11.25.** Of course, the proposition implies the more general converse that if  $Y$  is a qcqs diamond such that  $Y$  admits a surjective and universally open quasi-pro-étale map  $Y' \rightarrow Y$ , where  $Y'$  is a (locally) spatial diamond, then  $Y$  is a spatial diamond.

*Proof.* The converse is easy: If  $X \rightarrow Y$  is surjective and universally open quasi-pro-étale, and without loss of generality  $X$  is affinoid, and in fact strictly totally disconnected (cf. Lemma 7.18), then the induced equivalence relation  $R \subset X \times X$  is a qcqs perfectoid space, for which the maps  $s, t : R \rightarrow X$  are open; thus,  $|Y| = |X|/|R|$  is spectral and  $|X| \rightarrow |Y|$  is spectral by Lemma 2.10, which implies that  $Y$  is spatial.

Now assume that  $Y$  is spatial and consider the set  $I$  of isomorphism classes of surjective étale maps  $Y' \rightarrow Y$  that can be written as a composite of quasicompact open embeddings and finite étale maps. (To see that this is of cardinality less than  $\kappa$ , realize these by descent data along some fixed surjective map from a  $\kappa$ -small affinoid perfectoid space. Moreover, each  $Y'$  admits a surjection from a  $\kappa'$ -small affinoid perfectoid space for some fixed strong limit cardinal  $\kappa' < \kappa$ .) By Lemma 11.21, all such  $Y'$  are spatial diamonds. Choose a representative  $Y_i \rightarrow Y$  for each  $i \in I$ . For any subset  $J \subset I$ , let  $Y_J$  be the product of all  $Y_i$ ,  $i \in J$ , over  $Y$ , which is still a composite of quasicompact open embeddings and finite étale maps over  $Y$ , and thus a spatial diamond. By Lemma 11.22, the

cofiltered limit  $Y_\infty$  of all  $Y_J$  over finite subsets  $J \subset I$  is again a spatial diamond, and  $Y_\infty \rightarrow Y$  is a surjective and universally open quasi-pro-étale map that can be written as a cofiltered inverse limit of étale maps which are composites of quasicompact open immersions and finite étale maps. Repeating the construction  $Y \mapsto Y_\infty$  countably often, we can find a surjective and universally open quasi-pro-étale map  $X \rightarrow Y$  of spatial diamonds, which can be written as a cofiltered inverse limit of étale maps which are composites of quasicompact open immersions and finite étale maps. Now any étale cover  $\tilde{X} \rightarrow X$  that can be written as a composite of quasicompact open immersions and finite étale maps splits: Indeed, by Proposition 11.23, any such map comes from some finite level, and becomes split at the next. Now the result follows from the next proposition.  $\square$

**Proposition 11.26.** *Let  $Y$  be a spatial diamond. Assume that any surjective étale map  $\tilde{Y} \rightarrow Y$  that can be written as a composite of quasicompact open immersions and finite étale maps splits. Then  $Y$  is a strictly totally disconnected perfectoid space.*

*Proof.* First note that any quasicompact open cover of  $|Y|$  splits by assumption; thus, every connected component of  $|Y|$  has a unique closed point. Thus, any connected component  $Y_0 \subset Y$  admits a pro-étale surjection  $\mathrm{Spa}(C, C^+) \rightarrow Y_0$  for some algebraically closed field  $C$  with an open and bounded valuation subring  $C^+ \subset C$ . First, we claim this map  $\mathrm{Spa}(C, C^+) \rightarrow Y_0$  is an isomorphism. Let  $R_0 = \mathrm{Spa}(C, C^+) \times_{Y_0} \mathrm{Spa}(C, C^+)$  be the equivalence relation, which is affinoid pro-étale over  $\mathrm{Spa}(C, C^+)$ . To check that  $\mathrm{Spa}(C, C^+) = Y_0$ , we have to check that  $R_0 = \mathrm{Spa}(C, C^+)$ . If not, then  $R_0$  has another maximal point. We see that it is enough to show that  $\mathrm{Spa}(C, \mathcal{O}_C) \rightarrow Y_0^\circ$  is an isomorphism, where  $Y_0^\circ \subset Y_0$  is the open subfunctor corresponding to the maximal point (which is open). But now  $R_0^\circ = R_0 \times_{Y_0} Y_0^\circ$  is isomorphic to  $\underline{S} \times \mathrm{Spa}(C, \mathcal{O}_C)$  for some profinite set  $S$ , where in fact  $S = G$  is a profinite group (by the equivalence relation structure). Assume  $G$  is nontrivial, and let  $H \subset G$  be a proper open subgroup. Then  $\underline{H} \times \mathrm{Spa}(C, \mathcal{O}_C) \subset \underline{G} \times \mathrm{Spa}(C, \mathcal{O}_C) = R_0^\circ$  is another equivalence relation, and the corresponding quotient of  $\mathrm{Spa}(C, \mathcal{O}_C)$  is a nontrivial finite étale cover of  $Y_0^\circ$ . Note that finite étale covers of  $Y_0^\circ$  agree with finite étale covers of  $Y_0$  (as this is true for the pair  $\mathrm{Spa}(C, \mathcal{O}_C)$  and  $\mathrm{Spa}(C, C^+)$ , and the pair  $R_0$  and  $R_0^\circ$ ). Now any finite étale cover of  $Y_0$  extends to a finite étale cover of an open and closed subset of  $Y$ , which together with the complementary open and closed subset of  $Y$  forms an étale cover as in the statement of the proposition. As this is assumed to be split, the original finite étale cover of  $Y_0$  has to be split, which is a contradiction. Thus,  $Y_0 = \mathrm{Spa}(C, C^+)$ .

In other words, we have seen that any connected component of  $Y_0$  is given by  $\mathrm{Spa}(C, C^+)$  for some algebraically closed field  $C$  with an open and bounded valuation subring  $C^+ \subset C$ . The result now follows from the next lemma.  $\square$

**Lemma 11.27.** *Let  $Y$  be a spatial diamond. Assume that every connected component of  $Y$  is representable by an affinoid perfectoid space. Then  $Y$  is representable by an affinoid perfectoid space.*

*Proof.* Write  $Y = X/R$  as a quotient of a strictly totally disconnected perfectoid space  $X$  by an affinoid pro-étale equivalence relation  $R \subset X \times X$ . Let  $X = \mathrm{Spa}(B, B^+)$  and  $R = \mathrm{Spa}(C, C^+)$ , and define

$$(A, A^+) = \mathrm{eq}((B, B^+) \rightrightarrows (C, C^+)) .$$

The goal is to prove that  $A$  is perfectoid,  $A^+ \subset A$  is open and integrally closed, and  $Y = \mathrm{Spa}(A, A^+)$ .

As a first step, we find a pseudouniformizer  $\varpi_B \in B$  such that  $p_1^*(\varpi_B) = p_2^*(\varpi_B)u$  for some unit  $u \in 1 + C^{\circ\circ}$ ; here,  $p_1, p_2 : R \rightarrow X$  denote the two projections. We can find such a pseudouniformizer

over each connected component  $c \in \pi_0(Y)$  (by taking one coming via pullback from the affinoid connected component of  $Y$ ), and arbitrarily lifting it to  $B$ , this relation will be satisfied in the preimage of an open and closed neighborhood of  $c \in \pi_0(Y)$ . As the set of connected components is profinite, we can then find such a  $\varpi_B \in B$  globally.

Now we check that if an element  $t \in p_1^*(\varpi_B)^n C^{\circ\circ} = p_2^*(\varpi_B)^n C^{\circ\circ}$  (for some  $n \geq 0$ ) satisfies the cocycle relation

$$p_{23}^*(t) + p_{12}^*(t) = p_{13}^*(t)$$

(corresponding to the three maps  $p_{12}, p_{13}, p_{23} : R \times_X R \rightarrow R$ ), then there is an element  $s \in \varpi_B^n B^{\circ\circ}$  with  $p_1^*(s) - p_2^*(s) = t$ . It suffices to check that we can find  $s \in \varpi_B^n B^{\circ\circ}$  such that

$$p_1^*(s) - p_2^*(s) - t \in p_1^*(\varpi_B)^{n+1} C^{\circ\circ} ,$$

as then the result follows by induction. But in each connected component  $Y_c \subset Y$ , we can find such an  $s$  (by almost vanishing of  $H_v^1(Y_c, \mathcal{O}^+)$ ), and lifting it arbitrarily to  $B$ , it will satisfy the desired congruence in a neighborhood; again, as  $\pi_0(Y)$  is profinite, we can find some such  $s$  globally.

Applying this to  $t = p_1^*(\varpi_B) - p_2^*(\varpi_B) \in p_1^*(\varpi_B) C^{\circ\circ}$ , we find some  $s \in \varpi_B B^{\circ\circ}$  such that  $p_1^*(s) - p_2^*(s) = t$ . This implies that  $\varpi := \varpi_B - s \in A$ , and is a topologically nilpotent unit in  $B$ , and thus in  $A$ . Now we replace  $\varpi_B$  by  $\varpi$ , so  $\varpi$  gives a compatible choice of pseudouniformizer in all rings.

From the definition of  $A$ , it is clear that  $A$  is uniform and perfect, so  $A$  is perfectoid. Also,  $A^+ \subset A$  is open and integrally closed. Thus,  $Y' := \mathrm{Spa}(A, A^+)$  is an affinoid perfectoid space, and we get a natural map  $Y \rightarrow Y'$ . To see that this is an isomorphism, it suffices by Lemma 11.11 to check on connected components. It remains to see that the construction of  $(A, A^+)$  commutes with passage to connected components. But the arguments above show that

$$(A^+/\varpi)^a = \mathrm{eq}((B^+/\varpi)^a \rightrightarrows (C^+/\varpi)^a) ,$$

and this statement passes to open and closed subsets (i.e., direct summands on the level of rings), and filtered colimits. By the equivalence of perfectoid  $A$ -algebras with perfectoid  $(A^+/\varpi)^a$ -algebras, this gives the result on the level of the perfectoid Tate ring. For the open and integrally closed subalgebra, it suffices to check that if  $Y_c = \mathrm{Spa}(A_c, A_c^+)$  is a connected component of  $Y$ , then  $A^+ \rightarrow A_c^+$  is surjective. But any  $f \in A_c^+$  can be lifted to some function in  $g \in B^+$  with  $p_1^*(g) - p_2^*(g) \in \varpi C^{\circ\circ}$ , and then using the claim above we can correct it by some function  $s \in \varpi B^{\circ\circ}$  such that  $g - s \in A^+$ . Thus,  $A^+ \rightarrow A_c^+/\varpi$  is surjective, which implies that  $A^+ \rightarrow A_c^+$  is surjective, as  $A^+$  and  $A_c^+$  are  $\varpi$ -adically complete.  $\square$

Now we get the following two permanence properties.

**Corollary 11.28.** *Let  $Y$  be a locally spatial diamond, and  $Y' \rightarrow Y$  a quasi-pro-étale map of pro-étale sheaves. Then  $Y'$  is a locally spatial diamond.*

Recall that by Convention 10.2, the map  $Y' \rightarrow Y$  is automatically required to be locally separated.

*Proof.* We may assume that  $Y$  is spatial, and that  $Y' \rightarrow Y$  is separated. By Proposition 11.24, we can find a surjective and universally open quasi-pro-étale map  $X \rightarrow Y$ , where  $X$  is strictly totally disconnected. Then  $X' = X \times_Y Y'$  is representable and pro-étale over  $X$ , and  $R' = X' \times_{Y'} X'$  is representable and qcqs over  $X'$ . Moreover,  $X'$  is quasiseparated, so  $|Y'| = |X'|/|R'|$  is a locally



spectral space and  $|X'| \rightarrow |Y'|$  is a spectral map, by Lemma 2.10. This proves that  $Y'$  is locally spatial.  $\square$

**Corollary 11.29.** *A fibre product of (locally) spatial diamonds is (locally) spatial.*

*Proof.* Let  $Y_1 \rightarrow Y_3: Y_2$  be a diagram of locally spatial diamonds. We may assume that  $Y_3, Y_1$ , and  $Y_2$  are spatial. Assume first that  $Y_1 \rightarrow Y_3$  is a quasi-separated quasi-pro-étale map. Then the fibre product  $Y_1 \times_{Y_3} Y_2$  is quasi-separated quasi-pro-étale over  $Y_2$ , so the result follows from Corollary 11.28. In general, let  $X_3 \rightarrow Y_3$  be a surjective and universally open quasi-pro-étale map from a qcqs perfectoid space  $X_3$ , as guaranteed by Proposition 11.24. By Remark 11.25, it is enough to prove that

$$(Y_1 \times_{Y_3} Y_2) \times_{Y_3} X_3 = (Y_1 \times_{Y_3} X_3) \times_{X_3} (Y_2 \times_{Y_3} X_3)$$

is a spatial diamond. As the fibre products  $Y_1 \times_{Y_3} X_3$  and  $Y_2 \times_{Y_3} X_3$  are known to exist by the beginning of the proof (as  $X_3 \rightarrow Y_3$  is quasi-separated quasi-pro-étale), we may assume that  $Y_3 = X_3$  is a qcqs perfectoid space.

Now choose  $X_1 \rightarrow Y_1$  and  $X_2 \rightarrow Y_2$  surjective and universally open quasi-pro-étale maps. Then  $X_1 \times_{Y_3} X_2 \rightarrow Y_1 \times_{Y_3} Y_2$  is again surjective and universally open quasi-pro-étale, so the result follows from Proposition 11.24.  $\square$

Finally, there is the following 2-out-of-3 property for quasi-pro-étale maps.

**Proposition 11.30.** *Let  $f : Y_1 \rightarrow Y_2$  and  $g : Y_2 \rightarrow Y_3$  be maps of locally spatial diamonds, with composite  $h = g \circ f : Y_1 \rightarrow Y_3$ . Assume that  $f$  is quasi-pro-étale and surjective,  $h$  is quasi-pro-étale, and  $g$  is separated. Then  $g$  is quasi-pro-étale.*

*Proof.* By the previous corollary, we may assume that  $X_3 = Y_3$  is a strictly totally disconnected perfectoid space. Moreover, we can assume that  $Y_2$  is spatial. Now  $Y_1 = X_1$  is representable and pro-étale over  $X_3$ . Replacing  $X_1$  by an open cover, we can assume that  $X_1$  is a disjoint union of strictly totally disconnected perfectoid spaces. A finite union already covers  $Y_2$ , so we can assume that  $X_1$  is strictly totally disconnected, and thus affinoid pro-étale over  $X_3$ . Finally, we can replace  $X_3$  by  $X_3 \times_{\pi_0(X_3)} \pi_0(Y_2)$ , so that  $\pi_0(Y_2) = \pi_0(X_3)$ .

We claim that in this situation,  $|f| : |Y_2| \rightarrow |X_3|$  is automatically injective. The argument is the same as in the proof of Lemma 7.19, but we repeat it for convenience. We can check on connected components, so we may assume that  $X_3 = \text{Spa}(C, C^+)$  is a connected strictly totally disconnected space, and then by assumption also  $Y_2$  is connected. Now assume that two distinct points  $y_1, y_2 \in |Y_2|$  map to the same point  $x \in X_3$ . If  $x$  corresponds to a valuation ring  $(C^+)' \subset C$ , then  $y_1, y_2$  give rise to points still denoted  $y_1, y_2 \in Y_2(C, (C^+)')$  (by lifting further to  $X_1$ ). By the valuative criterion of separatedness, if  $y_1 \neq y_2$ , then the corresponding  $(C, \mathcal{O}_C)$ -points are still distinct, so we can assume that  $(C^+)' = \mathcal{O}_C$ . Then  $x$  is the unique rank 1 point of  $X$ , and the set of preimages of  $x$  forms a profinite set  $Y_{2,x} \subset Y_2$ , as they form a spectral set without specializations. By our assumption,  $Y_{2,x}$  contains at least two points, so we can find a closed and open decomposition  $Y_{2,x} = U_{1,x} \sqcup U_{2,x}$ , for some quasicompact open subsets  $U_1, U_2 \subset Y_2$ . Let  $V_1, V_2 \subset Y_2$  be the closures of  $U_{1,x}$  and  $U_{2,x}$ . As  $U_{1,x}$  and  $U_{2,x}$  are pro-constructible subsets, their closures are precisely the subsets of specializations of points in  $U_{1,x}$  resp.  $U_{2,x}$ . As any point of  $Y_2$  generalizes to a unique point of  $Y_{2,x}$ , we have  $Y_2 = V_1 \sqcup V_2$ . As  $V_1$  and  $V_2$  are both closed, this gives a contradiction to our assumption that  $Y_2$  is connected, finishing the proof that  $|f| : |Y_2| \rightarrow |X_3|$  is injective.

Moreover, the image of  $|f|$  is a pro-constructible and generalizing subset of  $|X_3|$ , and thus by Lemma 7.6 there is an affinoid pro-étale  $X_2 \subset X_3$  whose image is precisely  $|Y_2|$ ; thus  $Y_2 \rightarrow X_3$

factors over  $Y_2 \rightarrow X_2$ ; we can thus also replace  $X_3$  by  $X_2$ . We claim that the map  $Y_2 \rightarrow X_2$  is an isomorphism. For this, we will use Lemma 11.11. This reduces us to the case that  $X_3 = \mathrm{Spa}(C, C^+)$  where  $C$  is algebraically closed, and  $C^+ \subset C$  is an open and bounded valuation subring. But now as  $|Y_2| = |X_2| = |X_3| = |\mathrm{Spa}(C, C^+)|$ , also  $|Y_2|$  has a unique closed point, and so we can replace  $X_1 = \mathrm{Spa}(C', C'^+)$  by its localization at one point. But then  $(C', C'^+) = (C, C^+)$  (as  $X_1 \rightarrow X_2$  is affinoid pro-étale and surjective), and the composite  $X_1 \rightarrow Y_1 \rightarrow X_2$  is an isomorphism, where the first map is a surjection. Thus,  $X_1 = Y_1 = X_2$ , as desired.  $\square$

Another useful lemma is a structure result for the local nature of étale morphisms of locally spatial diamonds, generalizing Definition 6.2 (ii).

**Lemma 11.31.** *Let  $f : Y' \rightarrow Y$  be an étale map of locally spatial diamonds. Then for every  $y' \in |Y'|$  with image  $y \in |Y|$ , one can find open neighborhoods  $V' \subset Y'$  of  $y'$  and  $V \supset f(V')$  of  $y$  such that  $f|_{V'} : V' \rightarrow V$  factors as the composite of a quasicompact open immersion  $V' \hookrightarrow W$  and a finite étale map  $W \rightarrow V$ .*

We note again that by Convention 10.2, the map  $f$  is required to be locally separated. (This is necessary, as  $f|_{V'}$  is separated.)

*Proof.* We can assume that  $Y$  is spatial, and then also that  $Y'$  is spatial; thus  $f : Y' \rightarrow Y$  is a qcqs étale map of spatial diamonds, which we may moreover assume to be separated. Let  $Y'_{y'}$  resp.  $Y_y$  be the localization of  $Y'$  at  $y'$ , resp.  $Y$  at  $y$ , i.e. the intersection of all open subfunctors containing  $y'$  resp.  $y$ . Then  $|Y'_{y'}|$  resp.  $|Y_y|$  have a unique closed point given by  $y'$  resp.  $y$ . We claim that it suffices to prove that  $Y'_{y'}$  factors as the composite of a quasicompact open immersion  $Y'_{y'} \rightarrow W_y$  and a finite étale map  $W_y \rightarrow Y_y$ . Indeed, both the finite étale map  $W_y \rightarrow Y_y$  and the quasicompact open immersion arise via base change from  $V$  for some quasicompact open subfunctor  $V \subset Y$ , by Proposition 11.23 (i) and Lemma 11.22, respectively. Up to replacing  $Y$  by  $V$ , we get some map  $V' \rightarrow Y$  which is a composite of a quasicompact open immersion and a finite étale morphism, such that  $V' \times_Y Y_y \cong Y'_{y'} \subset Y' \times_Y Y_y$  over  $Y_y$ . Shrinking  $Y$  again, we may assume by Proposition 11.23 (ii) that the map  $V' \times_Y Y_y \rightarrow Y' \times_Y Y_y$  is defined over  $Y$  already, as a quasicompact open immersion  $V' \rightarrow Y'$ , with  $y'$  in the image. This gives the desired result.

Thus, we can assume that  $y$  is the unique closed point of  $Y$ . In that case,  $|Y|$  is a totally ordered chain of specializations (being a quotient of  $|\mathrm{Spa}(K, K^+)|$  for a perfectoid field  $K$  with an open and bounded valuation subring  $K^+ \subset K$ ). Let  $Y^\circ \subset Y$  correspond to the unique open point, and  $Y^\circ = Y' \times_Y Y^\circ$ . Then  $Y'^\circ \rightarrow Y^\circ$  is étale, and thus finite étale: Indeed,  $Y^\circ$  is covered by  $\mathrm{Spa}(K, \mathcal{O}_K)$ , and  $\mathrm{Spa}(K, \mathcal{O}_K)_{\text{ét}} = \mathrm{Spa}(K, \mathcal{O}_K)_{\text{fét}}$ . Moreover,  $Y'^\circ_{\text{fét}} \cong Y_{\text{fét}}$ , as finite étale spaces are insensitive to  $\mathcal{O}^+$ . We see that there is some finite étale space  $W \rightarrow Y$  with  $W \times_Y Y^\circ \cong Y'^\circ$ . The composite map  $Y'^\circ \cong W \times_Y Y^\circ \subset W$  extends uniquely to a map  $Y' \rightarrow W$  over  $Y$ , by checking the similar result after pullback to  $\mathrm{Spa}(K, K^+)$  (and uniqueness). Moreover, by Lemma 9.9, the map  $Y' \rightarrow W$  is an injection. In particular, if we let  $w \in |W|$  be the image of  $y' \in |Y'|$ , then  $Y'_{y'}$  maps isomorphically to  $W_w$ . This finishes the proof, as  $W_w \subset W$  is a quasicompact open subset, as  $|W|$  is a quotient of finitely many totally ordered chains of specializations, and so any pro-constructible generalizing subset is open.  $\square$

## 12. SMALL v-STACKS

The goal of this section is two-fold. First, we generalize many basic results to the setting of general (small) v-stacks. Secondly, we give a criterion for when a v-sheaf is a diamond, Theorem 12.18, without exhibiting an explicit quasi-pro-étale surjection. In this way, it is similar to Artin's theorem on algebraic spaces reducing smooth (or even flat) groupoids to étale groupoids.

For several results, we will need a set-theoretic smallness assumption.

**Definition 12.1.** *A small v-sheaf is a v-sheaf  $Y$  on  $\text{Perf}$  such that there is a surjective map of v-sheaves  $X \rightarrow Y$  for some ( $\kappa$ -small) perfectoid space  $X$ .*

**Remark 12.2.** Clearly, if  $Y$  is a diamond, then  $Y$  is a small v-sheaf. Also, if  $Y$  is a v-sheaf such that there is a surjective map of v-sheaves  $X \rightarrow Y$  for some diamond  $X$ , then  $Y$  is a small v-sheaf. Finally, if  $Y$  is a quasicompact v-sheaf, then it is small (as one can always cover  $Y$  by the disjoint union of all maps  $X \rightarrow Y$  from perfectoid spaces  $X$ ).

Perhaps surprisingly, the following proposition shows that any small v-sheaf has a reasonable geometric structure.

**Proposition 12.3.** *Let  $Y$  be a small v-sheaf, and let  $X \rightarrow Y$  be a surjective map of v-sheaves, where  $X$  is a diamond. Then  $R = X \times_Y X$  is a diamond, and  $Y = X/R$  as v-sheaves.*

*If  $Y$  is a quasiseparated small v-sheaf, and  $X \rightarrow Y$  is a surjective map of v-sheaves, where  $X$  is a locally spatial diamond, then  $R = X \times_Y X$  is a locally spatial diamond. In particular, if  $Y$  is a qcqs v-sheaf, and  $X$  is a spatial diamond, then  $R$  is a spatial diamond.*

*Proof.* As  $R \subset X \times X$  is a sub-v-sheaf, and  $X \times X$  is a diamond, the first part follows from Proposition 11.10. If  $Y$  is quasiseparated, then  $R = X \times_Y X \subset X \times X$  is a quasicompact sub-v-sheaf, so if  $X$  is locally spatial, then Proposition 11.20 shows that  $R$  is a locally spatial diamond. For the final sentence, note that if  $Y$  is quasicompact, then it is small. Now if  $X$  and  $Y$  are qcqs, then so is  $R = X \times_Y X$ , so  $R$  is spatial.  $\square$

This allows us to go one step further and pass to v-stacks.

**Definition 12.4.** *A small v-stack is a v-stack  $Y$  on  $\text{Perf}$  such that there is a surjective map of v-stacks  $X \rightarrow Y$  from a perfectoid space  $X$ , for which  $R = X \times_Y X$  is a small v-sheaf.*

In particular, all qcqs v-stacks are small. Lemma 11.11 extends to v-sheaves and more generally v-stacks.

**Lemma 12.5.** *Let  $f : Y' \rightarrow Y$  be a qcqs map of v-stacks. Then  $f$  is an isomorphism if and only if for all algebraically closed perfectoid fields  $K$  with an open and bounded valuation subring  $K^+ \subset K$ , the map  $f(K, K^+) : Y'(K, K^+) \rightarrow Y(K, K^+)$  is an equivalence of groupoids.*

**Remark 12.6.** We remind the reader of Convention 8.4 concerning quasiseparated maps of stacks.

*Proof.* It suffices to check the result after pullback to any affinoid perfectoid space  $X$  mapping to  $Y$ . In that case,  $Y'$  is a qcqs v-stack. Write  $Y' = X'/R'$  as the quotient of an affinoid perfectoid space  $X'$ ; then  $R'$  is a qcqs v-sheaf. The map  $X' \rightarrow X$  is a v-cover, as it is a map of qcqs perfectoid spaces which by assumption is surjective on points. As both  $X'$  and  $X$  are v-sheaves, it suffices to see that the map of v-sheaves  $R' \rightarrow X' \times_X X'$  is an isomorphism.

After this reduction, we have a new map  $Y' \rightarrow X$  from a qcqs v-sheaf  $Y'$  (the  $R'$  from above) to an affinoid perfectoid space  $X$  (the  $X' \times_X X'$  above). Again, write  $Y' = X'/R'$  as the quotient of an

affinoid perfectoid space  $X'$  by the equivalence relation  $R'$ , which by Proposition 12.3 is a spatial diamond. Repeating the arguments, it is enough to see that  $R' \rightarrow X' \times_X X'$  is an isomorphism of spatial diamonds. But the map  $R(K, K^+) \rightarrow (X' \times_X X')(K, K^+)$  is a bijection for all  $(K, K^+)$  as in the statement of the lemma. Thus, the result follows from Lemma 11.11.  $\square$

Moreover, one can define an underlying topological space for small v-stacks.

**Proposition 12.7.** *Let  $Y$  be a small v-stack, and let  $Y = X/R$  be a presentation as a quotient of a diamond  $X$  by a small v-sheaf  $R \rightarrow X \times X$ ; let  $\tilde{R} \rightarrow R$  be a surjection from a diamond. There is a canonical bijection between  $|X|/|\tilde{R}|$  and the set*

$$|Y| = \{\mathrm{Spa}(K, K^+) \rightarrow Y\} / \sim,$$

where  $(K, K^+)$  runs over all pairs of a perfectoid field  $K$  with an open and bounded valuation subring  $K^+ \subset K$ , and two such maps  $\mathrm{Spa}(K_i, K_i^+) \rightarrow Y$ ,  $i = 1, 2$ , are equivalent, if there is a third pair  $(K_3, K_3^+)$ , and a commutative diagram

$$\begin{array}{ccc} & \mathrm{Spa}(K_1, K_1^+) & \\ & \nearrow & \searrow \\ \mathrm{Spa}(K_3, K_3^+) & \longrightarrow & Y, \\ & \searrow & \nearrow \\ & \mathrm{Spa}(K_2, K_2^+) & \end{array}$$

where the maps  $\mathrm{Spa}(K_3, K_3^+) \rightarrow \mathrm{Spa}(K_i, K_i^+)$ ,  $i = 1, 2$ , are surjective.

Moreover, the quotient topology on  $|Y|$  induced by the surjection  $|X| \rightarrow |Y|$  is independent of the choice of presentation  $Y = X/R$ .

*Proof.* The proof of Proposition 11.13 carries over.  $\square$

Thus, the following definition is independent of the choice made.

**Definition 12.8.** *Let  $Y$  be a small v-stack with a presentation  $Y = X/R$  as a quotient of a diamond  $X$  by a small v-sheaf  $R \rightarrow X \times X$ , and  $\tilde{R} \rightarrow R$  a surjection from a diamond. The underlying topological space of  $Y$  is  $|Y| := |X|/|\tilde{R}|$ .*

Note that we are doing two steps at once here, really; one could also define this first for small v-sheaves, and then for small v-stacks as  $|Y| = |X|/|R|$ . But as  $|\tilde{R}| \rightarrow |R|$  is surjective, this recovers the same answer.

The following version of Proposition 11.15 holds for small v-stacks.

**Proposition 12.9.** *Let  $Y$  be a small v-stack with underlying topological space  $|Y|$ . Any open sub-v-stack of  $Y$  is a small v-stack. Moreover, there is bijective correspondence between open sub-v-stacks  $U \subset Y$  and open subsets  $V \subset |Y|$ , given by sending  $U$  to  $V = |U|$ , and  $V$  to the subfunctor  $U \subset Y$  of those maps  $X \rightarrow Y$  for which  $|X| \rightarrow |Y|$  factors over  $V \subset |Y|$ .*

Moreover, if  $f : Y' \rightarrow Y$  is a surjective map of small v-stacks, then  $|f| : |Y'| \rightarrow |Y|$  is a quotient map.

*Proof.* The same arguments as in the proof of Proposition 11.15 apply.  $\square$

The following proposition follows from the similar result for perfectoid spaces.

**Proposition 12.10.** *Let  $Y_1 \rightarrow Y_3 \leftarrow Y_2$  be a diagram of small v-stacks. Then  $Y = Y_1 \times_{Y_3} Y_2$  is a small v-stack, and*

$$|Y| \rightarrow |Y_1| \times_{|Y_3|} |Y_2|$$

*is surjective.* □

In the following, when we say that map  $f : Y' \rightarrow Y$  of (small) v-stacks is surjective, we mean that it is surjective as a map of v-stacks. There is a related notion that  $|f| : |Y'| \rightarrow |Y|$  is surjective; if we mean this, we say it explicitly. The relation is as follows.<sup>2</sup>

**Lemma 12.11.** *Let  $f : Y' \rightarrow Y$  be a map of small v-stacks. If  $f$  is a surjective map of v-stacks, then  $|f| : |Y'| \rightarrow |Y|$  is surjective. Conversely, if  $f$  is quasicompact and  $|f| : |Y'| \rightarrow |Y|$  is surjective, then  $f$  is a surjective map of v-stacks.*

*Proof.* The first part is clear. For the converse, assume that  $f$  is quasicompact and that  $|f| : |Y'| \rightarrow |Y|$  is surjective. If  $X$  is an affinoid perfectoid space with a map  $X \rightarrow Y$ , then the pullback  $Y' \times_Y X$  is a quasicompact v-stack, and thus admits a surjection  $X' \rightarrow Y' \times_Y X$  from an affinoid perfectoid space. Thus,  $|X'| \rightarrow |Y' \times_Y X|$  is surjective, and also  $|Y' \times_Y X| \rightarrow |Y'| \times_{|Y|} |X| \rightarrow |X|$  is a composition of surjections (using Proposition 12.10). Therefore,  $X' \rightarrow X$  is a map of affinoid perfectoid spaces for which  $|X'| \rightarrow |X|$  is surjective; thus, by definition, it is a v-cover. Thus, the map  $X \rightarrow Y$  lifts to  $X' \rightarrow Y'$  after the v-cover  $X' \rightarrow X$ , as desired. □

We can now introduce a notion of spatial v-sheaves.

**Definition 12.12.** *A v-sheaf  $Y$  is spatial if it is qcqs (in particular, small), and  $|Y|$  has a basis for the topology given by  $|U|$  for quasicompact open subfunctors  $U \subset Y$ . More generally,  $Y$  is locally spatial if it is small, and has an open cover by spatial v-sheaves.*

Proposition 11.18 extends to spatial v-sheaves.

**Proposition 12.13.** *Let  $Y$  be a spatial v-sheaf.*

- (i) *The underlying topological space  $|Y|$  is spectral.*
- (ii) *Any quasicompact open subfunctor  $U \subset Y$  is spatial.*
- (iii) *For any perfectoid space  $X'$  with a map  $X' \rightarrow Y$ , the induced map of locally spectral spaces  $|X'| \rightarrow |Y|$  is spectral and generalizing.*

*Proof.* The proof is identical to the proof of Proposition 11.18, noting that any qcqs v-sheaf  $Y$  can be written in the form  $Y = X/R$  for spatial diamonds  $X$  and  $R$ . □

**Proposition 12.14.** *Let  $Y$  be a locally spatial v-sheaf.*

- (i) *The underlying topological space  $|Y|$  is locally spectral.*
- (ii) *Any open subfunctor  $U \subset Y$  is locally spatial.*
- (iii) *The functor  $Y$  is quasicompact (resp. quasiseparated) if and only if  $|Y|$  is quasicompact (resp. quasiseparated).*
- (iv) *For any locally spatial v-sheaf  $Y'$  with a map  $Y' \rightarrow Y$ , the induced map of locally spectral spaces  $|Y'| \rightarrow |Y|$  is spectral and generalizing.*

<sup>2</sup>We thank David Hansen for related discussions.

□

There is the following analogue of Proposition 5.3 for locally spatial  $v$ -sheaves.

**Proposition 12.15.** *Let  $f : Y' \rightarrow Y$  be a map of small  $v$ -sheaves, and assume that  $f$  is qcqs, or  $Y'$  and  $Y$  are locally spatial. The following conditions are equivalent.*

- (i) *The map  $f$  is an injective map of  $v$ -sheaves.*
- (ii) *For all perfectoid fields  $K$  with an open and bounded valuation subring  $K^+ \subset K$ , the map  $f(K, K^+) : Y'(K, K^+) \rightarrow Y(K, K^+)$  is injective.*
- (iii) *The map  $|f| : |Y'| \rightarrow |Y|$  is injective, and the map  $f : Y' \rightarrow Y$  is final in the category of maps  $g : Z \rightarrow Y$  from small  $v$ -sheaves  $Z$  for which  $|g| : |Z| \rightarrow |Y|$  factors over a continuous map  $|Z| \rightarrow |Y'|$ .*

Note that (iii) says equivalently that

$$Y' = Y \times_{|Y|} \underline{|Y'|}.$$

*Proof.* Clearly, (iii) implies (i) implies (ii), so we have to see that (ii) implies (iii). Injectivity of  $|f|$  follows from the description of the set  $|Y|$  in Proposition 12.7. Now assume that  $g : Z \rightarrow Y$  has the property that  $|Z| \rightarrow |Y|$  lifts continuously to  $|Z| \rightarrow |Y'|$ . We claim that there is a unique map  $Z \rightarrow Y'$ . This can be done  $v$ -locally on  $Z$ , so we can assume that  $Z$  is a perfectoid space, or even a strictly totally disconnected perfectoid space. In case  $Y$  and  $Y'$  are locally spatial, we can assume that  $|Z|$  maps into qcqs open subsets of  $|Y|$  and  $|Y'|$ , we can work locally on  $Z$ , which allows us to assume that  $Y$  and  $Y'$  are spatial, and in particular  $f$  is qcqs. Now  $Z \times_Y Y' \rightarrow Z$  is a qcqs map of  $v$ -sheaves, and we need to see that it is an isomorphism. By Lemma 12.5, this can be checked on  $(K, K^+)$ -valued points, so we can further assume that  $Z = \mathrm{Spa}(K, K^+)$ . In that case, we know that  $Z \times_Y Y' \rightarrow Z$  is an injection. But quasicompact injections into  $\mathrm{Spa}(K, K^+)$  are of the form  $\mathrm{Spa}(K, (K^+)')$  by Corollary 10.6. Looking at topological spaces shows that  $(K^+) = K^+$ , so that indeed  $Z \times_Y Y' = Z$ , as desired. □

Moreover, we will need an analogue of Lemma 11.21.

**Lemma 12.16.** *Let  $Y$  be a spatial  $v$ -sheaf, and let  $Y' \rightarrow Y$  be a finite étale map of  $v$ -sheaves. Then  $Y'$  is a spatial  $v$ -sheaf.*

*Proof.* The proof is identical to Lemma 11.21, but we repeat it for convenience. Choose a presentation  $Y = X/R$  as a quotient of an affinoid perfectoid space  $X$ , and let  $X' = X \times_Y Y'$ , which is finite étale over  $X$ . Let  $R$  and  $R'$  be the respective equivalence relations. We need to find enough quasicompact open subspaces of  $Y'$ . Pick any point  $y' \in |Y'|$  with image  $y \in |Y|$ . Let  $Y_y \subset Y$  be the localization of  $Y$  at  $y$ , i.e. the intersection of all quasicompact open subspaces containing  $y$ . For any spatial diamond  $Z$  over  $Y$ , let  $Z_y = Z \times_Y Y_y$ . Then quasicompact open subspaces of  $Z_y$  come via pullback from quasicompact open subspaces of  $Z \times_Y U$  for some quasicompact open subspace  $U$  of  $Y$ , and any two such extensions agree after shrinking  $U$ . By applying this to  $Z = X'$  and  $Z = R'$ , we find that any quasicompact open subspace of  $Y'_y$  extends to a quasicompact open subspace of  $Y'_U$  for some  $U$ . This reduces us to the case  $Y = Y_y$ .

Thus, we may assume that  $|Y|$  has a unique closed point. In this case, we can take  $X = \mathrm{Spa}(C, C^+)$  for some algebraically closed field  $C$  with an open and bounded valuation subring  $C^+ \subset C$ . Then  $X'$  is a finite disjoint union of copies of  $X$ . This implies that any pro-constructible

generalizing subset of  $X'$  is open. In particular, for any quasicompact open subset  $V \subset X'$ , the  $R'$ -invariant subset  $R' \cdot V \subset X'$  is open (as it is pro-constructible and generalizing). This descends to a quasicompact open subspace of  $Y'$ , and these give a basis, as desired.  $\square$

Finally, we will need an analogue of Lemma 11.22 and Proposition 11.23.

**Lemma 12.17.** *Let  $Y_i$ ,  $i \in I$ , be a cofiltered inverse system of small  $v$ -sheaves with qcqs transition maps, where the index category  $I$  is of cardinality less than  $\kappa$ . Assume that all  $Y_i$  admit surjections from  $\kappa'$ -small perfectoid spaces for some fixed strong limit cardinal  $\kappa' < \kappa$ . Then  $Y = \varprojlim_i Y_i$  is a small  $v$ -sheaf, the maps  $Y \rightarrow Y_i$  are qcqs, and the continuous map  $|Y| \rightarrow \varprojlim_i |Y_i|$  is bijective. If all  $Y_i$  are locally spatial, then  $Y$  is locally spatial, and  $|Y| \rightarrow \varprojlim_i |Y_i|$  is a homeomorphism.*

*Moreover, if all  $Y_i$  are qcqs, then  $Y$  is qcqs, and the base change functors  $(Y_i)_{\text{fét}} \rightarrow Y_{\text{fét}}$ , resp.  $(Y_i)_{\text{ét, qcqs}} \rightarrow Y_{\text{ét, qcqs}}$ , resp.  $(Y_i)_{\text{ét, qc, sep}} \rightarrow Y_{\text{ét, qc, sep}}$ , induce an equivalence of categories*

$$\mathcal{2}\text{-}\varinjlim_i (Y_i)_{\text{fét}} \rightarrow Y_{\text{fét}} ,$$

respectively

$$\mathcal{2}\text{-}\varinjlim_i (Y_i)_{\text{ét, qcqs}} \rightarrow Y_{\text{ét, qcqs}} ,$$

respectively

$$\mathcal{2}\text{-}\varinjlim_i (Y_i)_{\text{ét, qc, sep}} \rightarrow Y_{\text{ét, qc, sep}} .$$

We remark again that by our Convention 10.2, all étale maps are locally separated.

*Proof.* It is easy to reduce to the case that all  $Y_i$  are qcqs, by fixing some index  $i = i_0$  and a surjection  $Y'_{i_0} \rightarrow Y_{i_0}$ , where  $Y'_{i_0}$  is a disjoint union of qcqs  $v$ -sheaves, and replacing the whole diagram by the diagram of  $Y_i \times_{Y_{i_0}} Y'_{i_0}$ .

We can assume that  $I$  is the category of ordinals  $\mu$  less than some fixed ordinal  $\lambda$  (which is less than  $\kappa$ ). Using Lemma 4.1, we can also assume that  $\kappa' = \kappa_\lambda$  is of cofinality larger than  $\lambda$ ; one then verifies that all constructions below happen with  $\kappa_\lambda$ -small sheaves.

By transfinite induction on  $\mu$ , we can find a diagram  $X_\mu$  of  $\kappa_\lambda$ -small spatial diamonds with compatible surjections  $X_\mu \rightarrow Y_\mu$ ; in fact, such that

$$X_\mu \rightarrow Y_\mu \times \varprojlim_{\mu' < \mu} Y_{\mu'} \varprojlim_{\mu' < \mu} X_{\mu'}$$

is surjective. To see that this can be done, assume it has been done for  $\mu' < \mu$ . Then

$$\varprojlim_{\mu' < \mu} X_{\mu'} \rightarrow \varprojlim_{\mu' < \mu} Y_{\mu'}$$

is a surjection of qcqs  $v$ -sheaves: Indeed, given a map  $Z \rightarrow \varprojlim_{\mu' < \mu} Y_{\mu'}$  from a spatial diamond, the diagram

$$Z_{\mu'} := Z \times \varprojlim_{\mu' < \mu} Y_{\mu'} X_{\mu'} \subset Z \times X_{\mu'}$$

over varying  $\mu' < \mu$  is a diagram of spatial diamonds (using Proposition 11.20), where all maps  $Z_{\mu'} \rightarrow Z$  are surjective. Then  $Z_\mu := \varprojlim_{\mu' < \mu} Z_{\mu'} \rightarrow Z$  is also a surjective map of spatial diamonds, using Lemma 11.22. Therefore,

$$\varprojlim_{\mu' < \mu} X_{\mu'} \rightarrow \varprojlim_{\mu' < \mu} Y_{\mu'}$$

is a qcqs surjection of v-sheaves, and the same holds for the base change

$$Y_\mu \times_{\varprojlim_{\mu' < \mu} Y_{\mu'}} \varprojlim_{\mu' < \mu} X_{\mu'} \rightarrow Y_\mu .$$

In particular, the source is a qcqs v-sheaf, and we can find a surjection  $X_\mu \rightarrow Y_\mu$  from a spatial diamond  $X_\mu$ .

Rereading the above arguments in the case  $\mu = \lambda$ , we see that  $Y = \varprojlim_{\mu < \lambda} Y_\mu$  admits a qcqs surjection from the spatial diamond  $X = \varprojlim_{\mu < \lambda} X_\mu$ , so  $Y$  is a qcqs v-sheaf.

Let  $R_\mu = X_\mu \times_{Y_\mu} X_\mu$  and  $R = X \times_Y X = \varprojlim_{\mu < \lambda} R_\mu$  be the equivalence relations, which are spatial diamonds. Then

$$|Y| = |X|/|R| = \varprojlim |X_\mu|/|R_\mu| = \varprojlim |Y_\mu|$$

as sets. If all  $Y_\mu$  are spatial, then the same arguments as in the proof of Lemma 11.22 ensure that  $|Y| = \varprojlim |Y_\mu|$  is a homeomorphism, and  $Y$  is spatial.

For the final statement about finite étale maps, let  $F$  be the prestack on the category of small v-sheaves of finite étale, resp. quasicompact separated étale, morphisms. Then  $F$  is a stack for the v-topology by Proposition 10.11. Note that  $F(X/Y)$  involves the value of  $F$  on  $X$ ,  $R$ , and  $R \times_X R$ , all of which are spatial diamonds, and similarly for  $F(X_\mu/Y_\mu)$ . By Proposition 11.23, we see that

$$F(Y) = F(X/Y) = 2\text{-}\varinjlim F(X_\mu/Y_\mu) = 2\text{-}\varinjlim F(Y_\mu) ,$$

as desired.

This leaves the case of  $Y_{\text{ét, qcqs}}$ . In that case, one still gets a fully faithful functor

$$2\text{-}\varinjlim_i (Y_i)_{\text{ét, qcqs}} \rightarrow Y_{\text{ét, qcqs}} .$$

For essential surjectivity, note that any  $\tilde{Y} \rightarrow Y$  in  $Y_{\text{ét, qcqs}}$  is per convention locally separated, and thus locally lies in  $Y_{\text{ét, qc, sep}}$ . Using that case, we can descend to some finite level, and then we can also descend the gluing data.  $\square$

The following theorem will be used heavily to show that certain functors are (spatial) diamonds.

**Theorem 12.18.** *Let  $Y$  be a spatial v-sheaf such that there exists a perfectoid space  $X$  with a quasi-pro-étale map  $f : X \rightarrow Y$  for which  $|f| : |X| \rightarrow |Y|$  is surjective. Then  $Y$  is a spatial diamond.*

**Remark 12.19.** The hypothesis here is much weaker than asking that  $f$  is surjective. An equivalent condition is to ask that for any  $y \in |Y|$ , there exists a quasi-pro-étale map  $\text{Spa}(C, C^+) \rightarrow Y$  having  $y$  in its image, where  $C$  is some algebraically closed nonarchimedean field with an open and bounded valuation subring  $C^+ \subset C$ . Indeed, the disjoint union of all such  $\text{Spa}(C, C^+) \rightarrow Y$  over varying  $y \in |Y|$  will then give the desired map  $f : X \rightarrow Y$  from a perfectoid space  $X$  (which will be highly non-quasicompact, so that Lemma 12.11 does not apply).

In particular, the condition in Theorem 12.18 is only a condition about the points of  $Y$ .

*Proof.* We need to see that  $Y$  is a diamond, as the notion of being spatial is compatible. To see that  $Y$  is a diamond, we need to find a surjective quasi-pro-étale map  $X \rightarrow Y$ , where  $X$  is a perfectoid space. For this, we follow the argument in the proof of Proposition 11.24. Consider the set  $I$  of isomorphism classes of surjective étale maps  $Y' \rightarrow Y$  that can be written as a composite of quasicompact open embeddings and finite étale maps. (To see that this is of cardinality less than



$\kappa$ , realize these by descent data along some fixed surjective map from a  $\kappa$ -small affinoid perfectoid space. Moreover, each  $Y'$  is  $\kappa'$ -small for some fixed strong limit cardinal  $\kappa' < \kappa$ .) By Lemma 12.16, all such  $Y'$  are spatial  $v$ -sheaves. Choose a representative  $Y_i \rightarrow Y$  for each  $i \in I$ . For any subset  $J \subset I$ , let  $Y_J$  be the product of all  $Y_i$ ,  $i \in J$ , over  $Y$ , which is still a composite of quasicompact open embeddings and finite étale maps over  $Y$ , and thus a spatial  $v$ -sheaf. Thus, by Lemma 12.17, the cofiltered limit  $Y_\infty$  of all  $Y_J$  over finite subsets  $J \subset I$  is again a spatial  $v$ -sheaf, and  $Y_\infty \rightarrow Y$  is a surjective separated quasi-pro-étale map. Repeating the construction  $Y \mapsto Y_\infty$  countably often, we find a surjective separated quasi-pro-étale map  $X \rightarrow Y$  of spatial  $v$ -sheaves. Now any étale cover  $\tilde{X} \rightarrow X$  that can be written as a composite of quasicompact open immersions and finite étale maps splits: Indeed, by Lemma 12.17, any such map comes from some finite level, and becomes split at the next. Now the result follows from the following generalization of Proposition 11.26, noting that the condition on points lifts to  $X$  (as  $X \rightarrow Y$  is quasi-pro-étale).  $\square$

**Proposition 12.20.** *Let  $Y$  be a spatial  $v$ -sheaf such that there exists a quasi-pro-étale map  $f : X \rightarrow Y$  from a perfectoid space  $X$  for which  $|f| : |X| \rightarrow |Y|$  is surjective. Assume that any surjective étale map  $\tilde{Y} \rightarrow Y$  that can be written as a composite of quasicompact open immersions and finite étale maps splits. Then  $Y$  is a strictly totally disconnected perfectoid space.*

*Proof.* The proof is identical to the proof of Proposition 11.26, but we repeat it for convenience. First note that any quasicompact open cover of  $|Y|$  splits by assumption; thus, every connected component of  $|Y|$  has a unique closed point. Thus, any connected component  $Y_0 \subset Y$  admits a quasi-pro-étale surjection  $\mathrm{Spa}(C, C^+) \rightarrow Y_0$  for some algebraically closed field  $C$  with an open and bounded valuation subring  $C^+ \subset C$ . First, we claim this map  $\mathrm{Spa}(C, C^+) \rightarrow Y_0$  is an isomorphism. Let  $R_0 = \mathrm{Spa}(C, C^+) \times_{Y_0} \mathrm{Spa}(C, C^+)$  be the equivalence relation. To check that  $\mathrm{Spa}(C, C^+) = Y_0$ , we have to check that  $R_0 = \mathrm{Spa}(C, C^+)$ . If not, then  $R_0$  has another maximal point. We see that it is enough to show that  $\mathrm{Spa}(C, \mathcal{O}_C) \rightarrow Y_0^\circ$  is an isomorphism, where  $Y_0^\circ \subset Y_0$  is the open subfunctor corresponding to the maximal point (which is open). But now  $R_0^\circ = R_0 \times_{Y_0} Y_0^\circ$  is isomorphic to  $\underline{S} \times \mathrm{Spa}(C, \mathcal{O}_C)$  for some profinite set  $S$  as  $R_0^\circ \rightarrow \mathrm{Spa}(C, \mathcal{O}_C)$  is a qcqs pro-étale map, where in fact  $\underline{S} = G$  is a profinite group (by the equivalence relation structure). Assume  $G$  is nontrivial, and let  $H \subset G$  be a proper open subgroup. Then  $\underline{H} \times \mathrm{Spa}(C, \mathcal{O}_C) \subset \underline{G} \times \mathrm{Spa}(C, \mathcal{O}_C) = R_0^\circ$  is another equivalence relation, and the corresponding quotient of  $\mathrm{Spa}(C, \mathcal{O}_C)$  is a nontrivial finite étale cover of  $Y_0^\circ$ . Note that finite étale covers of  $Y_0^\circ$  agree with finite étale covers of  $Y_0$  (as this is true for the pair  $\mathrm{Spa}(C, \mathcal{O}_C)$  and  $\mathrm{Spa}(C, C^+)$ , and the pair  $R_0$  and  $R_0^\circ$ ). Now any finite étale cover of  $Y_0$  extends to a finite étale cover of an open and closed subset of  $Y$  by Lemma 12.17, which together with the complementary open and closed subset of  $Y$  forms an étale cover as in the statement of the proposition. As this is assumed to be split, the original finite étale cover of  $Y_0$  has to be split, which is a contradiction. Thus,  $Y_0 = \mathrm{Spa}(C, C^+)$ .

In other words, we have seen that any connected component of  $Y_0$  is given by  $\mathrm{Spa}(C, C^+)$  for some algebraically closed field  $C$  with an open and bounded valuation subring  $C^+ \subset C$ . The result now follows from the next lemma, which generalizes Lemma 11.27.  $\square$

**Lemma 12.21.** *Let  $Y$  be a spatial  $v$ -sheaf. Assume that every connected component of  $Y$  is representable by an affinoid perfectoid space. Then  $Y$  is representable by an affinoid perfectoid space.*

*Proof.* Write  $Y = X/R$  as the quotient of an affinoid perfectoid space  $X$  by a spatial diamond equivalence relation  $R \subset X \times X$ . By Lemma 11.27, the spatial diamond  $R$  is an affinoid perfectoid space. Now the same argument as in the proof of Lemma 11.27 applies.  $\square$

### 13. SPATIAL MORPHISMS

In this section, we define a notion of (locally) spatial morphisms, and prove that it behaves well. All notions we will consider will be examples of 0-truncated maps.

**Definition 13.1.** *A map  $f : Y' \rightarrow Y$  of  $v$ -stacks is representable in diamonds if for all diamonds  $X$  with a map  $X \rightarrow Y$ , the fibre product  $Y' \times_Y X$  is a diamond.*

The following proposition ensures that this notion is well-behaved.

**Proposition 13.2.** *Let  $f : Y' \rightarrow Y$  and  $\tilde{Y} \rightarrow Y$  be maps of  $v$ -stacks, with pullback  $\tilde{f} : \tilde{Y}' = Y' \times_Y \tilde{Y} \rightarrow \tilde{Y}$ .*

- (i) *If  $Y$  is a diamond, then  $f$  is representable in diamonds if and only if  $Y'$  is a diamond.*
- (ii) *If  $f$  is representable in diamonds, then  $\tilde{f}$  is representable in diamonds.*
- (iii) *If  $\tilde{Y} \rightarrow Y$  is surjective as a map of pro-étale stacks and  $\tilde{f}$  is representable in diamonds, then  $f$  is representable in diamonds.*

*Proof.* Part (i) follows from Proposition 11.4, and part (ii) is clear by definition. For part (iii), we may assume that  $Y$  is a diamond. We can find a surjective quasi-pro-étale  $X \rightarrow Y$ , where  $X$  is a disjoint union of strictly totally disconnected perfectoid spaces, and it suffices to see that  $X \times_Y Y'$  is a diamond by Proposition 11.6. This can be checked locally on  $X$ , so we can reduce to the case that  $Y = X$  is a strictly totally disconnected perfectoid space. As  $\tilde{Y} \rightarrow Y = X$  is surjective as a map of pro-étale sheaves, we can find a surjective pro-étale map  $\tilde{X} \rightarrow X$  with a lift  $\tilde{X} \rightarrow \tilde{Y}$ . Then  $\tilde{X} \times_X Y'$  is a diamond, and thus  $Y'$  by Proposition 11.6.  $\square$

Now we can define locally spatial morphisms.

**Definition 13.3.** *A map  $f : Y' \rightarrow Y$  of  $v$ -stacks is representable in (locally) spatial diamonds if for all (locally) spatial diamonds  $X$  with a map  $X \rightarrow Y$ , the fibre product  $Y' \times_Y X$  is a (locally) spatial diamond.*

If  $f : Y' \rightarrow Y$  is a map of diamonds which is representable in (locally) spatial diamonds, we will sometimes simply say that  $f : Y' \rightarrow Y$  is a (locally) spatial map of diamonds.

Clearly, if a map of  $v$ -stacks  $f : Y' \rightarrow Y$  is representable in spatial diamonds, then it is representable in locally spatial diamonds, and the converse holds precisely when  $f$  is qcqs.

**Proposition 13.4.** *Let  $f : Y' \rightarrow Y$  and  $\tilde{Y} \rightarrow Y$  be maps of  $v$ -stacks, with pullback  $\tilde{f} : \tilde{Y}' = Y' \times_Y \tilde{Y} \rightarrow \tilde{Y}$ .*

- (i) *If  $f$  is representable in locally spatial diamonds, then  $\tilde{f}$  is representable in diamonds.*
- (ii) *If  $Y$  is a locally spatial diamond, then  $f$  is representable in locally spatial diamonds if and only if  $Y'$  is a locally spatial diamond.*
- (iii) *If  $f$  is representable in locally spatial diamonds, then  $\tilde{f}$  is representable in locally spatial diamonds.*

- (iv) If  $\tilde{Y} \rightarrow Y$  is surjective as a map of pro-étale sheaves,  $f$  is quasiseparated and  $\tilde{f}$  is representable in locally spatial diamonds, then  $f$  is representable in locally spatial diamonds.
- (v) If  $f$  is representable in diamonds,  $\tilde{Y} \rightarrow Y$  is a surjective map of  $v$ -stacks,  $f$  is quasiseparated and  $\tilde{f}$  is representable in locally spatial diamonds, then  $f$  is representable in locally spatial diamonds.

*Proof.* Part (i) follows from Proposition 13.2 (i) and (iii). Part (ii) follows from Corollary 11.29, and part (iii) is clear by definition.

For part (iv), we may assume that  $Y$  is a locally spatial diamond; in fact, we can assume that  $Y$  is spatial. By Proposition 13.2 (iii), we know that  $Y'$  is a diamond. By Proposition 11.24 and Lemma 2.10, we can further reduce to the case that  $Y$  is strictly totally disconnected: By Proposition 11.18, we can find a universally open map  $X \rightarrow Y$  where  $X$  is strictly totally disconnected, and if  $X \times_Y Y'$  is a locally spatial diamond, then so is  $Y'$  by Lemma 2.10.

Now assume  $Y = X$  is strictly totally disconnected. We can find an affinoid pro-étale map  $\tilde{X} \rightarrow X$  which lifts to  $\tilde{Y}$ , and may assume that  $\tilde{Y} = \tilde{X}$ . Now  $\tilde{Y}' := \tilde{X} \times_X Y'$  is a locally spatial diamond. We claim that this implies that  $Y'$  is a locally spatial diamond.

Let  $V \subset Y'$  be an open subfunctor, and let  $y \in |V| \subset |Y'|$  be a point, lying in a connected component  $c \in \pi_0 X$ . By a subscript  $c$ , we denote the fiber of all objects over  $c$ . Note that  $\tilde{X}_c \rightarrow X_c$  splits as  $X_c$  is a connected strictly totally disconnected space; this implies that  $Y'_c$  is locally spatial. Fix a quasicompact open subset  $U_c \subset Y'_c$  containing  $y$ . We can find a quasicompact open subset  $U \subset \tilde{Y}'$  contained in the preimage of  $V$ , whose intersection with  $\tilde{Y}'_c$  is given by the preimage of  $U_c$ . The two preimages of  $U$  in  $\tilde{Y}' \times_{Y'} \tilde{Y}'$  are two quasicompact open subsets  $W_1, W_2 \subset \tilde{Y}' \times_{Y'} \tilde{Y}'$  whose fibers over  $c$  agree. By standard properties of spectral spaces, it follows that there is some open and closed neighborhood  $U_c$  of  $c$  in  $\pi_0 X$  such that the intersection of  $W_1$  and  $W_2$  with the preimage of  $U_c$  agree. Thus, the intersection of  $U$  with the preimage of  $U_c$  descends to  $Y'$ , and defines a quasicompact open subfunctor of  $Y'$  containing  $y$ . This proves that  $Y'$  is locally spatial, as desired.

Finally, for part (v), we can use part (iv) to reduce to the case that  $Y = X$  is a strictly totally disconnected perfectoid space, in which case  $Y'$  is a qcqs diamond. By Lemma 13.5 below, we can assume that  $X = \mathrm{Spa}(C, C^+)$  is a connected strictly totally disconnected perfectoid space. Then we can assume that similarly  $\tilde{X} = \mathrm{Spa}(\tilde{C}, \tilde{C}^+)$  is a connected strictly totally disconnected perfectoid space. Write  $\tilde{X}$  as an inverse limit of rational subspaces  $\tilde{X}_i$  of perfected closed balls over  $X$ . Let  $V \subset Y'$  be an open subspace, with preimage  $\tilde{V} \subset \tilde{Y}'$ , and let  $y \in |V|$ . Then there is some quasicompact open subspace  $\tilde{U} \subset \tilde{V}$  containing the preimage of  $y$ . This quasicompact open subspace spreads to a quasicompact open subspace  $U_i \subset Y'_i = Y' \times_X \tilde{X}_i$  for  $i$  sufficiently large by arguing as in Lemma 13.5. Moreover,  $\tilde{U}_i \subset \tilde{V}_i$  for  $i$  sufficiently large by a standard quasicompactness argument. Now the map  $\tilde{X}_i \rightarrow X$  has a section by Lemma 9.5. Pulling back  $\tilde{U}_i$  under this section gives a quasicompact open subspace of  $Y'$  contained in  $V$ , and containing  $y$ , as desired.  $\square$

**Lemma 13.5.** *Let  $Y$  be a quasiseparated small  $v$ -sheaf with a map  $Y \rightarrow \underline{S}$  for some profinite set  $S$ . Assume that for all  $s \in S$ , the fiber  $Y_s$  is locally spatial. Moreover, assume that there is a surjective qcqs map  $X \rightarrow Y$  from a locally spatial diamond  $X$ . Then  $Y$  is locally spatial.*

We note that the existence of the surjective qcqs map  $X \rightarrow Y$  from a locally spatial diamond  $X$  is automatic if  $Y$  is quasicompact.

*Proof.* Choose a surjection  $X \rightarrow Y$  as in the statement of the lemma, and let  $R = X \times_Y X$  be the induced equivalence relation, which is a locally spatial diamond by Proposition 12.3. Let  $V \subset Y$  be an open subspace with preimage  $U \subset X$ , and let  $y \in |U|$  be a point, with image  $s \in S$ . Then we can find a quasicompact open subspace  $V'_s \subset Y_s \cap V$ . Its preimage  $U'_s \subset X_s$  extends to a quasicompact open subspace  $U'_T \subset X_T \cap U$  for any sufficiently small compact open neighborhood  $T$  of  $s$  in  $S$ . Moreover,  $U'_T$  is invariant under the equivalence relation for  $T$  sufficiently small (as the two preimages under  $s, t : R_T \rightarrow X_T$  are quasicompact open subspace whose fibers over  $s$  agree, so a quasicompactness argument applies), and thus descends to a quasicompact open subspace  $V'_T \subset Y_T \cap U$  for  $T$  sufficiently small. But then in particular  $V'_T \subset U$  is a quasicompact open subspace of  $Y$  containing  $y$ , which proves that  $Y$  is locally spatial.  $\square$

The following characterization of quasi-pro-étale maps was suggested by L. Fargues.

**Proposition 13.6.** *Let  $f : Y' \rightarrow Y$  be a separated map of  $v$ -stacks. Then  $f$  is quasi-pro-étale if and only if it is representable in locally spatial diamonds and for all complete algebraically closed fields  $C$  with a map  $\mathrm{Spa}(C, \mathcal{O}_C) \rightarrow Y$ , the pullback  $Y' \times_Y \mathrm{Spa}(C, \mathcal{O}_C) \rightarrow \mathrm{Spa}(C, \mathcal{O}_C)$  is pro-étale.*

*Proof.* If  $f$  is quasi-pro-étale, then Proposition 13.4 (iv) implies that  $f$  is representable in locally spatial diamonds (noting that to check this, one can assume that  $Y$  is a spatial diamond, which admits a quasi-pro-étale cover by a strictly totally disconnected space).

Conversely, we may assume that  $Y = X$  is a strictly totally disconnected perfectoid space. Moreover, we can assume that  $Y'$  is spatial. In this case, we can find a surjective separated quasicompact quasi-pro-étale map  $X' \rightarrow Y'$  as in Proposition 11.24. By Lemma 7.19, the map  $X' \rightarrow X$  is affinoid pro-étale. By Proposition 11.30, the map  $Y' \rightarrow X$  is quasi-pro-étale.  $\square$

Moreover, we will need to know that for any quasicompact separated diamond  $Y$ , there is a compact Hausdorff space  $T$  and a map  $Y \rightarrow \underline{T}$  which is representable in locally spatial diamonds. We will deduce this from a general discussion of “Berkovich spaces”.

**Definition 13.7.** *Let  $X = \mathrm{Spa}(R, R^+)$  be an affinoid perfectoid space, and fix some topologically nilpotent unit  $\varpi \in R$ . The Berkovich space  $|X|^B$  associated with  $X$  is the space of all multiplicative bounded nonarchimedean seminorms  $|\cdot| : R \rightarrow \mathbb{R}_{\geq 0}$  with  $|\varpi| = \frac{1}{2}$ , equipped with the weakest topology making the functions  $f \mapsto |f|$  continuous for all  $f \in R$ .*

**Remark 13.8.** The space  $|X|^B$  is canonically independent of the choice of  $\varpi$ , for example by Proposition 13.9 below.

Note that there is a natural map  $|X|^B \rightarrow |X|$ , as any multiplicative bounded nonarchimedean seminorm is in particular a valuation. There is also a map  $|X| \rightarrow |X|^B$ : Given any  $x \in X$ , we get a corresponding map  $\mathrm{Spa}(K(x), K(x)^+) \rightarrow \mathrm{Spa}(R, R^+)$ , where  $K(x)$  is a complete nonarchimedean field with an open and bounded valuation subring  $K(x)^+ \subset K(x)$ . In particular,  $K(x)$  comes with a unique up to scaling nonarchimedean norm  $|\cdot| : K(x) \rightarrow \mathbb{R}_{\geq 0}$ ; it can be normalized by  $|\varpi| = \frac{1}{2}$ . Thus, the composition  $R \rightarrow K(x) \rightarrow \mathbb{R}_{\geq 0}$  defines a multiplicative bounded nonarchimedean seminorm, and thus a point of  $|X|^B$ . One checks easily that the composition  $|X|^B \rightarrow |X| \rightarrow |X|^B$  is the identity.

**Proposition 13.9.** *Let  $X$  be an affinoid perfectoid space. The topological space  $|X|^B$  is compact Hausdorff, and the map  $|X| \rightarrow |X|^B$  is a continuous map identifying  $|X|^B$  as the maximal Hausdorff quotient of  $|X|$ .*

We warn the reader that the map  $|X|^B \rightarrow |X|$  is not continuous.

*Proof.* The space  $|X|^B$  is Hausdorff, as if  $|\cdot|, |\cdot|'$  are two distinct points, then there is some  $f \in R$  such that  $|f| \neq |f|'$ , so there is some real number  $r \in \mathbb{R}_{>0}$  lying strictly between  $|f|$  and  $|f|'$ ; then the subsets of points giving  $f$  absolute value strictly less than  $r$  resp. strictly bigger than  $r$  are open subsets of  $|X|^B$  which contain exactly one of  $|\cdot|$  and  $|\cdot|'$ .

Moreover, the map  $|X| \rightarrow |X|^B$  is surjective (as it has a set-theoretic section) and continuous, as follows easily from the definition. Thus,  $|X|^B$  is quasicompact, as  $|X|$  is quasicompact. Also, the fibers of  $|X| \rightarrow |X|^B$  have a unique maximal point by construction: They all share the same completed residue field  $K(x)$ , and thus generalize to the point given by  $\text{Spa}(K(x), \mathcal{O}_{K(x)})$ . Thus, any map from  $|X|$  to a Hausdorff space factors set-theoretically over  $|X|^B$ . It remains to see that  $|X| \rightarrow |X|^B$  is a quotient map, which follows from Lemma 2.6.  $\square$

We can now extend to general small v-sheaves, much as in the case of  $|Y|$ .

**Proposition 13.10.** *There is a unique colimit-preserving functor  $Y \mapsto |Y|^B$  from small v-sheaves to topological spaces extending  $X \mapsto |X|^B$  on affinoid perfectoid spaces. It comes with natural transformations  $|Y|^B \rightarrow |Y| \rightarrow |Y|^B$  whose composite is the identity, and the map  $|Y| \rightarrow |Y|^B$  is continuous and a quotient map.*

*Proof.* For uniqueness, note that the first is first uniquely defined for disjoint unions of affinoid perfectoid spaces, and then (by taking equivalence relations) for separated perfectoid spaces, for diamonds, and for small v-sheaves. One explicit construction of  $|Y|^B$  as a set is as the subset of  $|Y|$  consisting of those maps  $\text{Spa}(K, K^+) \rightarrow Y$  that can be represented by a map  $\text{Spa}(K, \mathcal{O}_K) \rightarrow Y$ . One endows  $|Y|^B$  with the quotient topology from  $|X|^B$ , for any surjection  $X \rightarrow Y$  from a disjoint union of affinoid perfectoid spaces  $X$ , and checks that this is independent of the choice of  $X$ . From the explicit description of  $|Y|^B$ , we also get the natural transformations  $|Y|^B \rightarrow |Y| \rightarrow |Y|^B$  (noting that any map  $\text{Spa}(K, K^+) \rightarrow Y$  in particular gives a map  $\text{Spa}(K, \mathcal{O}_K) \rightarrow Y$  for the construction of  $|Y| \rightarrow |Y|^B$ ).

Finally, to see that  $|Y| \rightarrow |Y|^B$  is a quotient map, take a cover by a disjoint union of affinoid perfectoid spaces  $X \rightarrow Y$ . Then  $|X|^B \rightarrow |Y|^B$  is a quotient map by definition, and  $|X| \rightarrow |X|^B$  is a quotient map by Proposition 13.9, which implies that  $|Y| \rightarrow |Y|^B$  is a quotient map.  $\square$

**Proposition 13.11.** *Let  $Y$  be a qcqs v-sheaf. Then  $|Y|^B$  is a compact Hausdorff space, and it is the maximal Hausdorff quotient of  $|Y|$ .*

*Proof.* First, we prove this if  $Y$  is a spatial diamond. Then  $Y = X/R$ , where  $X$  and  $R$  are affinoid perfectoid spaces, and then  $|Y|^B = |X|^B/|R|^B$  is a quotient of compact Hausdorff spaces, and thus compact Hausdorff itself. Now, if  $Y$  is any qcqs v-sheaf, we can write  $Y = X/R$  as a quotient, where  $X$  and  $R$  are spatial diamonds. Repeating the argument, we see that  $|Y|^B$  is compact Hausdorff.

As all points in the fiber of  $|Y| \rightarrow |Y|^B$  over  $y \in |Y|^B$  generalize to the image of  $y$  under  $|Y|^B \rightarrow |Y|$ , it follows that any map from  $|Y|$  to a Hausdorff space factors set-theoretically over  $|Y|^B$ . By Lemma 2.6,  $|Y| \rightarrow |Y|^B$  is a quotient map, so the result follows.  $\square$

As promised, we can now show that a general quasicompact separated diamond differs from a (locally) spatial diamond only through a map to a compact Hausdorff space.

**Proposition 13.12.** *Let  $Y$  be a quasicompact separated diamond. Then the map  $Y \rightarrow \underline{|Y|^B}$  is representable in locally spatial diamonds.*

*Proof.* By Proposition 13.11, the space  $|Y|^B$  is compact Hausdorff. Choose a profinite set  $S$  with a surjection  $S \rightarrow |Y|^B$ . By Proposition 13.4 (iv), it is enough to show that  $Y \times_{|Y|^B} \underline{S}$  is a spatial diamond. By Lemma 13.5, it is enough to check that for any  $s \in |Y|^B$ , the fiber product  $Y_s = Y \times_{|Y|^B} \underline{s}$  is a spatial diamond. It then follows from the definitions that  $|Y_s|^B = \{s\}$  is a point.

In other words, we have to prove that if  $Y$  is a quasicompact separated diamond such that  $|Y|^B$  is a point, then  $Y$  is spatial. Let  $y \in |Y|$  be the image of  $|Y|^B \rightarrow |Y|$ . Then one can find a quasi-pro-étale map  $f : \mathrm{Spa}(C, \mathcal{O}_C) \rightarrow Y$  with image  $y$ , for some algebraically closed nonarchimedean field  $C$ . Let  $Y_y \subset Y$  be the image of  $f$ , so that  $Y_y \rightarrow Y$  is a quasicompact injection, and  $|Y_y| = \{y\}$ . One finds that  $\mathrm{Spa}(C, \mathcal{O}_C) \times_{Y_y} \mathrm{Spa}(C, \mathcal{O}_C) = \mathrm{Spa}(C, \mathcal{O}_C) \times \underline{G}$  for some profinite group  $G$  acting continuously and faithfully on  $C$ , as in the proof of Proposition 11.26. Consider  $\bar{Y} = \mathrm{Spa}(C, C^+)/\underline{G}$ , where  $C^+$  is the integral closure of  $\mathbb{F}_p + C^{\circ\circ}$  in  $C$  (cf. Proposition 18.7 for a generalization of this construction). Then  $\mathrm{Spa}(C, C^+) \rightarrow \bar{Y}$  is a  $\underline{G}$ -torsor, and thus universally open by Lemma 10.13. Thus,  $\bar{Y}$  is a spatial diamond by Proposition 11.24.

To finish the proof, it suffices (by Proposition 11.20) to show that there is a (necessarily quasicompact) injection  $Y \rightarrow \bar{Y}$ . We construct the map  $Y \rightarrow \bar{Y}$  as a natural transformation on totally disconnected perfectoid spaces  $X = \mathrm{Spa}(R, R^+)$ . Given a map  $X \rightarrow Y$ , we note that the induced map  $\mathrm{Spa}(R, R^\circ) \rightarrow Y$  factors over  $Y_y$ ; indeed,  $\mathrm{Spa}(R, R^+) \times_Y Y_y \subset \mathrm{Spa}(R, R^+)$  is a quasicompact injection which contains all maximal points; it then follows from Lemma 7.6 that it contains  $\mathrm{Spa}(R, R^\circ)$  (as all occurring functions will necessarily lie in  $R^\circ$ ). The map  $\mathrm{Spa}(R, R^\circ) \rightarrow Y_y$  gives a map  $\mathrm{Spa}(R, R^+) \rightarrow \bar{Y}$  (cf. Proposition 18.7 (iv)). This defines the desired map  $Y \rightarrow \bar{Y}$ . To check that it is injective, it suffices by Proposition 12.15 to check injectivity on  $(K, K^+)$ -valued points, where  $K$  is a perfectoid field with an open and bounded valuation subring  $K^+ \subset K$ . As  $Y$  and thus  $Y \rightarrow \bar{Y}$  are separated, the valuative criterion of separatedness shows that it is enough to check injectivity on  $(K, \mathcal{O}_K)$ -valued points. But on  $(K, \mathcal{O}_K)$ -valued points, one has  $Y(K, \mathcal{O}_K) = Y_y(K, \mathcal{O}_K) = \bar{Y}(K, \mathcal{O}_K)$ .  $\square$

#### 14. COMPARISON OF ÉTALE, PRO-ÉTALE AND V-COHOMOLOGY

We consider the following sites.

**Definition 14.1.** *Let  $Y$  be a small v-stack on  $\mathrm{Perf}$ .*

- (i) *Assume that  $Y$  is a locally spatial diamond. The étale site  $Y_{\acute{e}t}$  is the site whose objects are (locally separated) étale maps  $Y' \rightarrow Y$ , with coverings given by families of jointly surjective maps.*
- (ii) *Assume that  $Y$  is a diamond. The quasi-pro-étale site  $Y_{\mathrm{qpro}\acute{e}t}$  is the site whose objects are (locally separated) quasi-pro-étale maps  $Y' \rightarrow Y$ , with coverings given by families of jointly surjective maps.*
- (iii) *The v-site  $Y_v$  is the site whose objects are all maps  $Y' \rightarrow Y$  from small v-sheaves  $Y'$ , with coverings given by families of jointly surjective maps.*

We note that here, in all cases, surjectivity refers to surjectivity as v-stacks on  $\mathrm{Perf}$ . Thus, if  $X$  and  $\{X_i \rightarrow X\}$  are perfectoid spaces, then surjectivity means that  $\{X_i \rightarrow X\}$  is a cover in the v-topology. Again, the topoi are algebraic.

**Proposition 14.2.** *The topoi  $Y_{\acute{e}t}$  resp.  $Y_{\mathrm{qpro}\acute{e}t}$  resp.  $Y_v$  for a locally spatial diamond resp. diamond resp. small v-stack  $Y$  are algebraic. If  $Y$  is 0-truncated (i.e., if  $Y$  is a small v-sheaf), then an object is quasicompact resp. quasiseparated if and only if it is quasicompact resp. quasiseparated as a small v-stack on  $\mathrm{Perf}$ .*

*Proof.* Left to the reader.  $\square$

**Proposition 14.3.** *Let  $Y$  be a locally spatial diamond. Then the étale site  $Y_{\text{ét}}$  has enough points. More precisely, for any  $y \in |Y|$ , choose a quasi-pro-étale map  $\bar{y} : \text{Spa}(C(y), C(y)^+) \rightarrow Y$  where  $C(y)$  is algebraically closed and  $C(y)^+ \subset C(y)$  is an open and bounded valuation subring, such that  $f_{\bar{y}}$  maps the closed point to  $y$ . Then*

$$\mathcal{F} \mapsto \mathcal{F}_{\bar{y}} = \varinjlim_{\bar{y} \rightarrow U \in Y_{\text{ét}}} \mathcal{F}(U)$$

defines a point of the topos  $Y_{\text{ét}}^{\sim}$ , and a section  $s \in \mathcal{F}(Y)$  of a sheaf  $\mathcal{F}$  on  $Y_{\text{ét}}$  is zero if and only if

$$s_{\bar{y}} = 0$$

for all  $y \in |Y|$ .

*Proof.* Note that equalizers exist in  $Y_{\text{ét}}$ , so the colimit defining  $\mathcal{F}_{\bar{y}}$  is filtered.

The functor  $\mathcal{F} \mapsto \mathcal{F}_{\bar{y}}$  is the composition of pullback along  $\bar{y}_{\text{ét}} : \text{Spa}(C(y), C(y)^+)_{\text{ét}} \rightarrow Y_{\text{ét}}$ , with global sections on  $\text{Spa}(C(y), C(y)^+)_{\text{ét}}$ . But note that any étale map to  $\text{Spa}(C(y), C(y)^+)$  splits and  $\text{Spa}(C(y), C(y)^+)$  is connected, so the functor of global sections is exact and commutes with all colimits, i.e. defines a point.

Now if  $s \in \mathcal{F}(Y)$  has the property that  $s_{\bar{y}} = 0$  for all  $y \in |Y|$ , then for all  $y \in |Y|$  we can find some  $\bar{y} \rightarrow U_{\bar{y}} \in Y_{\text{ét}}$  such that  $s|_{U_{\bar{y}}} = 0$ . As étale maps are open, it follows that the disjoint union of all  $U_{\bar{y}}$  is an étale cover of  $Y$ , over which  $s$  becomes 0, so that  $s = 0$ , as desired.  $\square$

There are obvious functors of sites

$$\lambda_Y : Y_v \rightarrow Y_{\text{qproét}}$$

if  $Y$  is a diamond, and

$$\nu_Y : Y_{\text{qproét}} \rightarrow Y_{\text{ét}}$$

if  $Y$  is a locally spatial diamond. Recall that this means that the underlying functor of categories goes the other way, and is given by observing that any étale map is quasi-étale, and any quasi-pro-étale map has source given by some small v-sheaf. These functors commute with all finite limits (which exist in all cases), so they define maps of topoi. In particular, pullback gives functors

$$Y_{\text{ét}}^{\sim} \xrightarrow{\nu_X^*} Y_{\text{qproét}}^{\sim} \xrightarrow{\lambda_X^*} Y_v^{\sim}$$

on the corresponding categories of sheaves.

A surprising feature of the situation is the following observation.

**Lemma 14.4.** *Let  $Y$  be a diamond. Then the functor*

$$\lambda_X^* : Y_{\text{qproét}}^{\sim} \rightarrow Y_v^{\sim}$$

*commutes with all limits.*

*Proof.* We have to show that for any category  $I$  and functor  $I \rightarrow Y_{\text{qproét}}^{\sim} : i \mapsto \mathcal{F}_i$ , the natural map

$$\lambda_X^* \left( \varprojlim_i \mathcal{F}_i \right) \rightarrow \varprojlim_i (\lambda_X^* \mathcal{F}_i)$$

is an isomorphism. This assertion is local in  $Y_{\text{qproét}}$ , so we can reduce to the case that  $Y = X$  is representable. One can then further assume that  $X$  is affinoid, and strictly totally disconnected.

In this situation,  $X_{\text{qproét}}^{\sim}$  is equivalent to  $X_{\text{qproét, qc, sep}}^{\sim}$ , where  $X_{\text{qproét, qc, sep}} \subset X_{\text{qproét}}$  denotes the site of quasicompact separated pro-étale morphisms  $X' \rightarrow X$ . Similarly,  $X_v^{\sim}$  is equivalent to  $X_{v, \text{qcqs}}^{\sim}$ , where  $X_{v, \text{qcqs}} \subset X_v$  denotes the site of qcqs and representable  $X' \rightarrow X$ . In particular,  $X_{\text{qproét, qc, sep}}^{\sim}$  and  $X_{v, \text{qcqs}}^{\sim}$  consist of qcqs objects (in the site-theoretic sense), stable under fibre products; it follows that  $X_{\text{qproét, qc, sep}}^{\sim}$  and  $X_{v, \text{qcqs}}^{\sim}$  are coherent. Now we need the following lemma.

**Lemma 14.5.** *Let  $X$  be a strictly totally disconnected perfectoid space, and  $X' \rightarrow X$  a map from a qcqs perfectoid space  $X'$ . Then there is a quasicompact separated pro-étale morphism  $\lambda_{X^\circ}(X') \rightarrow X$  with a factorization*

$$X' \rightarrow \lambda_{X^\circ}(X') \rightarrow X$$

*such that any map from  $X'$  to a separated pro-étale perfectoid space over  $X$  factors uniquely over  $\lambda_{X^\circ}(X')$ . Moreover:*

- (i) *The map  $X' \rightarrow \lambda_{X^\circ}(X')$  is surjective. In particular, if  $X'_2 \rightarrow X'_1$  is a surjection of quasicompact separated perfectoid spaces over  $X$ , then  $\lambda_{X^\circ}(X'_2) \rightarrow \lambda_{X^\circ}(X'_1)$  is surjective as well.*
- (ii) *If  $X'_1 \rightarrow X'_3 \leftarrow X'_2$  is a diagram of strictly totally disconnected perfectoid spaces over  $X$ , then*

$$\lambda_{X^\circ}(X'_1 \times_{X'_3} X'_2) \rightarrow \lambda_{X^\circ}(X'_1) \times_{\lambda_{X^\circ}(X'_3)} \lambda_{X^\circ}(X'_2)$$

*is an isomorphism.*

*Proof.* Recall that by Corollary 7.22, the category of quasicompact separated pro-étale perfectoid spaces over  $X$  is equivalent to the category of spectral maps of spectral spaces  $T \rightarrow |X|$  for which  $T \rightarrow |X| \times_{\pi_0 X} \pi_0 T$  is a pro-constructible generalizing embedding. Now for any quasicompact separated map  $X' \rightarrow X$ , we can define  $T$  as the image of  $|X'| \rightarrow |X| \times_{\pi_0 X} \pi_0 X'$ , which is of this form. The corresponding quasicompact separated pro-étale map  $\lambda_{X^\circ}(X') \rightarrow X$  has the desired universal property, and  $X' \rightarrow \lambda_{X^\circ}(X')$  is surjective by construction.

For part (ii), we first check surjectivity, for which we need to see that

$$X'_1 \times_{X'_3} X'_2 \rightarrow \lambda_{X^\circ}(X'_1) \times_{\lambda_{X^\circ}(X'_3)} \lambda_{X^\circ}(X'_2)$$

is surjective. We may assume that all spaces are connected, so  $X = \text{Spa}(C, C^+)$ , and  $X_i = \text{Spa}(C_i, C_i^+)$  for  $i = 1, 2, 3$ , where  $C$  and  $C_i$  are algebraically closed nonarchimedean fields with open and bounded valuation subrings  $C^+$  resp.  $C_i^+$ . The fibre product  $\lambda_{X^\circ}(X'_1) \times_{\lambda_{X^\circ}(X'_3)} \lambda_{X^\circ}(X'_2)$  is given by  $\text{Spa}(C, (C^+)^{\prime})$  for some other open and bounded valuation subring. We may pullback everything under the open immersion  $\text{Spa}(C, (C^+)^{\prime}) \rightarrow \text{Spa}(C, C^+)$ . After this replacement,  $\lambda_{X^\circ}(X'_i) = X$  for all  $i = 1, 2, 3$ , so  $\text{Spa}(C_i, C_i^+) \rightarrow \text{Spa}(C, C^+)$  is surjective. As  $\text{Spa}(C_3, C_3^+)$  is a totally ordered chain of specializations and the images of  $\text{Spa}(C_i, C_i^+) \rightarrow \text{Spa}(C_3, C_3^+)$  for  $i = 1, 2$  are generalizing, one of them is contained in the other; we may replace  $\text{Spa}(C_3, C_3^+)$  by the image of the smaller one (and  $X'_1, X'_2$  by the corresponding preimages). This preserves the condition that the maps  $X'_i \rightarrow X$  are surjective as now they will have the same image given by the minimal image that occurred previously (which was all of  $X$ ). After this further replacement,  $X'_i \rightarrow X'_3$  is surjective for  $i = 1, 2$ . Thus,

$$|X'_1 \times_{X'_3} X'_2| \rightarrow |X'_1| \times_{|X'_3|} |X'_2| \rightarrow |X'_3| \rightarrow |X|$$

is a series of surjections, as desired.

Finally, we need to show that the map is an isomorphism. Again, we can assume that all spaces are connected, and that  $X'_i \rightarrow X'_3$  are surjective for  $i = 1, 2$ , and  $X'_3 \rightarrow X$  is surjective. We have to see that  $X'_1 \times_{X'_3} X'_2$  is connected. This follows from the following general result.



**Lemma 14.6.** *Let  $X = \mathrm{Spa}(C, C^+)$ , where  $C$  is an algebraically closed nonarchimedean field with an open and bounded valuation subring  $C^+ \subset C$ , and let  $Z \rightarrow X$  be a connected affinoid perfectoid space over  $X$ . Let  $X' = \mathrm{Spa}(C', C'^+) \rightarrow X$ , where  $C'$  is another algebraically closed nonarchimedean field with an open and bounded valuation subring  $C'^+ \subset C'$ . Then  $Z' := Z \times_X X'$  is connected.*

*Proof.* Assume first that  $C' = C$  and  $C'^+ = \mathcal{O}_C$ . Then any disconnection of  $Z'$  extends to a disconnection of  $Z$  by taking closures, noting that any point of  $Z$  has a unique maximal generalization, which is a point of  $Z'$ . Thus, in general, we can assume that  $C^+ = \mathcal{O}_C$  and  $C'^+ = \mathcal{O}_{C'}$ . We can write  $Z$  as a cofiltered inverse limit of p-finite perfectoid spaces  $Z_i$  over  $(C, \mathcal{O}_C)$ , cf. [Sch12, Lemma 6.13]. We can without loss of generality assume that all  $Z_i$  are connected. By pullback,  $Z' = \varprojlim_i Z'_i$ , with  $Z'_i = Z_i \times_X X'$ . Any disconnection of  $Z'$  comes from a disconnection of  $Z'_i$  for  $i$  sufficiently large. But for rigid spaces, being geometrically connected passes to extensions of algebraically closed fields, so  $Z_i$  is disconnected for some  $i$ , which is a contradiction.  $\square$

$\square$

The first part of Lemma 14.5 implies that for any sheaf  $\mathcal{F}$  on  $X_{\mathrm{qproét}, \mathrm{ét}, \mathrm{sep}}$ , the pullback  $\lambda_X^* \mathcal{F}$  on  $X_{v, \mathrm{qcqs}}$  is the sheafification of the presheaf sending  $X' \in X_{v, \mathrm{qcqs}}$  to  $\mathcal{F}(\lambda_{X^\circ}(X'))$ . Now Lemma 14.5 (i) implies that this defines a separated presheaf, and Lemma 14.5 (ii) implies that for strictly totally disconnected  $X' \in X_{v, \mathrm{qcqs}}$ , the sheafification does not change anything (using that any cover can be refined by a cover by a strictly totally disconnected space); thus, for strictly totally disconnected  $X' \in X_{v, \mathrm{qcqs}}$ , we have

$$(\lambda_X^* \mathcal{F})(X') = \mathcal{F}(\lambda_{X^\circ}(X')) .$$

As evaluation commutes with limits, this equation commutes with limits of sheaves. As strictly totally disconnected spaces form a basis of  $X_{v, \mathrm{qcqs}}$ , we get the desired result.  $\square$

**Proposition 14.7.** *Let  $Y$  be a diamond. The functor*

$$\lambda_Y^* : Y_{\mathrm{qproét}}^\sim \rightarrow Y_v^\sim$$

*is fully faithful. Moreover, for any sheaf  $\mathcal{F}$  on  $Y_{\mathrm{qproét}}$ , the adjunction map*

$$\mathcal{F} \rightarrow \lambda_{Y*} \lambda_Y^* \mathcal{F}$$

*is an isomorphism, and if  $\mathcal{F}$  is a sheaf of abelian groups (resp. groups), then*

$$R^i \lambda_{Y*} \lambda_Y^* \mathcal{F} = 0$$

*for all  $i > 0$  (resp. for  $i = 1$ ).*

*Proof.* To prove fully faithfulness, it is enough to show that the adjunction map  $\mathcal{F} \rightarrow \lambda_{Y*} \lambda_Y^* \mathcal{F}$  is an isomorphism for all sheaves  $\mathcal{F}$  on  $Y_{\mathrm{qproét}}$ . This statement, as well as the similar statements about higher direct images, is local in  $Y_{\mathrm{qproét}}$ . Thus, we can assume that  $Y = X$  is a strictly totally disconnected perfectoid space.

In this case, we know by the proof of Lemma 14.4 that  $(\lambda_X^* \mathcal{F})(X') = \mathcal{F}(\lambda_{X^\circ}(X'))$  for all strictly totally disconnected  $X'$  over  $X$ . In particular, if  $X' \in X_{\mathrm{qproét}, \mathrm{qc}, \mathrm{sep}}$ , this implies that

$$(\lambda_{X*} \lambda_X^* \mathcal{F})(X') = (\lambda_X^* \mathcal{F})(X') = \mathcal{F}(\lambda_{X^\circ}(X')) = \mathcal{F}(X') .$$

As such  $X'$  form a base for  $X_{\mathrm{qproét}} = X_{\mathrm{qproét}}$ , it follows that  $\mathcal{F} \rightarrow \lambda_{Y*} \lambda_Y^* \mathcal{F}$  is an isomorphism.

For the statement about cohomology, note that by Lemma 14.5, the functor  $X' \mapsto \lambda_{X^\circ}(X')$  from strictly totally disconnected perfectoid spaces  $X' \rightarrow X$  to affinoid pro-étale perfectoid spaces over

$X$  takes surjections to surjections, and commutes with all finite limits. This implies that if  $X'_\bullet \rightarrow X$  is any v-hypercover of  $X$  by strictly totally disconnected spaces, then  $\overline{X'_\bullet} \rightarrow X$  is a hypercover of  $X$  in  $X_{\text{qproét}}$ . But the simplicial objects

$$(\lambda_X^* \mathcal{F})(X'_\bullet) = \mathcal{F}(\lambda_{X \circ}(X'_\bullet))$$

agree, so by writing cohomology as the filtered colimit over all hypercovers, we get

$$R\Gamma(X_v, \lambda_X^* \mathcal{F}) = R\Gamma(X_{\text{qproét}}, \mathcal{F})$$

for all sheaves of abelian groups  $\mathcal{F}$  on  $X_{\text{qproét}}$  (and a similar statement on  $H^1$  for sheaves of groups). Using the same result for  $X$  replaced by a space affinoid pro-étale over  $X$  and sheafifying gives the result.  $\square$

Next, we will need a similar result for  $\nu_Y : Y_{\text{qproét}} \rightarrow Y_{\text{ét}}$  in case  $Y$  is a locally spatial diamond.

**Proposition 14.8.** *Let  $Y$  be a locally spatial diamond. The functor*

$$\nu_Y^* : Y_{\text{ét}}^\sim \rightarrow Y_{\text{qproét}}^\sim$$

*is fully faithful. Moreover, for any sheaf  $\mathcal{F}$  on  $Y_{\text{ét}}$ , the adjunction map*

$$\mathcal{F} \rightarrow \nu_{Y*} \nu_Y^* \mathcal{F}$$

*is an isomorphism, and if  $\mathcal{F}$  is a sheaf of abelian groups (resp. groups), then*

$$R^i \nu_{Y*} \nu_Y^* \mathcal{F} = 0$$

*for all  $i > 0$  (resp. for  $i = 1$ ).*

*Proof.* We may assume that  $Y$  is spatial. Let  $Y_{\text{ét,qc,sep}} \subset Y_{\text{ét}}$  be the full subcategory of quasicompact separated étale maps. As all étale maps are by Convention 10.2 locally separated, and using Corollary 11.28, this is a basis for the topology, so  $Y_{\text{ét,qc,sep}}^\sim \cong Y_{\text{ét}}^\sim$ . Moreover, there is a functor

$$\text{Pro}_\kappa(Y_{\text{ét,qc,sep}}) \rightarrow Y_{\text{qproét}} .$$

As in the proof of Proposition 7.10, Proposition 11.23 (iii) implies that this functor is fully faithful. Moreover, the objects in  $\text{Pro}_\kappa(Y_{\text{ét,qc,sep}}) \subset Y_{\text{qproét}}$  form a basis for the topology. Indeed, choose a map  $X \rightarrow Y$  from a strictly totally disconnected space  $X$  as in Proposition 11.24, so that  $X \in \text{Pro}_\kappa(Y_{\text{ét,qc,sep}})$ . If  $Y' \rightarrow Y$  is any quasi-pro-étale map, then  $X' := X \times_Y Y'$  is a cover of  $Y'$  in  $Y_{\text{qproét}}$ , and is representable and pro-étale over  $X$ , so can be covered by perfectoid spaces which are affinoid pro-étale over  $X$ . All such affinoid pro-étale maps to  $X$  are inverse limits of affinoid étale, and in particular quasicompact separated étale maps, to  $X$ . Using Proposition 11.23 (iii), any such map lies in  $\text{Pro}_\kappa(Y_{\text{ét,qc,sep}})$  again, giving the claim.

Thus, we can replace the map  $\nu_Y : Y_{\text{qproét}} \rightarrow Y_{\text{ét}}$  by  $\text{Pro}_\kappa(Y_{\text{ét,qc,sep}}) \rightarrow Y_{\text{ét,qc,sep}}$ . Now the same arguments as in Proposition 8.5 apply.  $\square$

Moreover, we have the following commutation of étale cohomology with inverse limits.

**Proposition 14.9.** *Let  $Y_i$ ,  $i \in I$ , be a cofiltered inverse system of spatial diamonds with inverse limit  $Y$ . Assume that  $I$  has a final object  $0 \in I$ , and let  $\mathcal{F}_0$  be an étale sheaf on  $Y_0$ , with pullbacks  $\mathcal{F}_i$  to  $Y_i$ , and  $\mathcal{F}$  to  $Y$ . Then the natural map*

$$\varinjlim_{i \in I} H^j(Y_i, \mathcal{F}_i) \rightarrow H^j(Y, \mathcal{F})$$

is an isomorphism for  $j = 0$ , resp.  $j = 0, 1$ , resp. all  $j \geq 0$ , if  $\mathcal{F}_0$  is sheaf of sets, resp. sheaf of groups, resp. sheaf of abelian groups.

*Proof.* As in the previous proof, we can replace  $Y_{\acute{e}t}$  (and  $(Y_i)_{\acute{e}t}$ ) by  $Y_{\acute{e}t, \text{qc, sep}}$  (and  $(Y_i)_{\acute{e}t, \text{qc, sep}}$ ). Then Proposition 11.23 (iii) implies that  $Y_{\acute{e}t, \text{qc, sep}}^{\sim} = Y_{\acute{e}t}^{\sim}$  is a limit of the fibres topos  $(Y_i)_{\acute{e}t, \text{qc, sep}}^{\sim} = (Y_i)_{\acute{e}t}^{\sim}$ . As all intervening topoi are coherent, the result follows from SGA 4 VI.8.7.7 (in the case of sheaves of abelian groups, the other cases being similar).  $\square$

Using the previous results, we can compare some derived categories. Fix a ring  $\Lambda$ . If  $T$  is a topos and  $A$  is a ring on  $T$ , we denote by  $D(T, A)$  the derived category of  $A$ -modules on  $T$ , and by  $D^+(T, A), D^-(T, A), D^b(T, A) \subset D(T, A)$  the subcategories of complexes which are (cohomologically) bounded below, resp. above, resp. below and above.

**Proposition 14.10.** *Let  $Y$  be a diamond. Then the pullback functor*

$$\lambda_Y^* : D(Y_{\text{qproét}}, \Lambda) \rightarrow D(Y_v, \Lambda)$$

*is fully faithful. If  $Y$  is locally spatial, then the pullback functor*

$$\nu_Y^* : D^+(Y_{\acute{e}t}, \Lambda) \rightarrow D^+(Y_{\text{qproét}}, \Lambda)$$

*is fully faithful. If  $Y$  is a strictly totally disconnected perfectoid space, then the pullback functor*

$$\nu_Y^* : D(Y_{\acute{e}t}, \Lambda) \rightarrow D(Y_{\text{qproét}}, \Lambda)$$

*is fully faithful.*

*Proof.* The statements for  $D^+$  follow easily from Proposition 14.7 and Proposition 14.8. For the unbounded statements, we need some convergence results. For this, note that  $Y_{\text{qproét}}$  and  $Y_{\acute{e}t}$ , in case  $Y$  is strictly totally disconnected, have a basis for the topology consisting of  $U \in Y_{\text{qproét}}$  resp.  $U \in Y_{\acute{e}t}$  such that for all sheaves  $\mathcal{F}$ ,  $H^i(U, \mathcal{F}) = 0$  for  $i > 0$ . Indeed, in the case of  $Y_{\text{qproét}}$ , one can take strictly w-local perfectoid spaces  $U$  for which  $\pi_0 U$  is extremally disconnected (as then every pro-étale cover splits), and in the case of  $Y_{\acute{e}t}$ , one can take strictly totally disconnected  $U$ . This implies that for any  $A \in D(Y_{\text{qproét}}, \Lambda)$  resp.  $A \in D(Y_{\acute{e}t}, \Lambda)$ , we have  $A = R\varprojlim_n \tau^{\geq -n} A$  by testing the values on such  $U$ . In fact, the similar result holds true also in  $D(Y_v, \Lambda)$  as  $Y_v$  is replete, using [BS15, Proposition 3.3.3].

Now, in the case of  $Y_{\text{qproét}}$ , we have to see that for all  $A \in D(Y_{\text{qproét}}, \Lambda)$ , the adjunction map

$$A \rightarrow R\lambda_{Y*} \lambda_Y^* A$$

is an isomorphism. But the result holds true for all  $\tau^{\geq -n} A$ , so writing  $A = R\varprojlim_n \tau^{\geq -n} A$ , the result follows from the convergence results of the first paragraph. The same discussion applies to the case of  $Y_{\acute{e}t}$ .  $\square$

Note that in the first paragraph of the preceding proof, we have proved the following.

**Proposition 14.11.** *Let  $Y$  be a small  $v$ -stack.*

- (i) *The derived category  $D(Y_v, \Lambda)$  is left-complete.*
- (ii) *If  $Y$  is a diamond, then  $D(Y_{\text{qproét}}, \Lambda)$  is left-complete.*
- (iii) *If  $Y$  is a strictly totally disconnected perfectoid space, then  $D(Y_{\acute{e}t}, \Lambda)$  is left-complete.*  $\square$

Thus, if  $Y$  is a locally spatial diamond, we have full subcategories

$$D^+(Y_{\text{ét}}, \Lambda) \subset D(Y_{\text{qproét}}, \Lambda) \subset D(Y_v, \Lambda) .$$

Interestingly, containment in these subcategories can be checked  $v$ -locally.

**Theorem 14.12.** *Let  $Y$  be a diamond,  $A \in D(Y_v, \Lambda)$ , and  $f : Y' \rightarrow Y$  a  $v$ -cover by a diamond  $Y'$ .*

- (i) *If  $f^*A \in D(Y'_{\text{qproét}}, \Lambda)$ , then  $A \in D(Y_{\text{qproét}}, \Lambda)$ .*
- (ii) *If  $Y$  and  $Y'$  are locally spatial and  $f^*A \in D^+(Y'_{\text{ét}}, \Lambda)$ , then  $A \in D^+(Y_{\text{ét}}, \Lambda)$ .*
- (iii) *If  $Y$  and  $Y'$  are strictly totally disconnected and  $f^*A \in D(Y'_{\text{ét}}, \Lambda)$ , then  $A \in D(Y_{\text{ét}}, \Lambda)$ .*

*Proof.* As pullbacks commute with canonical truncations, we can assume that  $A \in D^+(Y_v, \Lambda)$ , using the convergence results from the proof of Proposition 14.10. Using Proposition 14.10, we can then further reduce to the case that  $C = \mathcal{F}$  is a sheaf of  $\Lambda$ -modules.

In part (i), we have to see that the adjunction map  $\lambda_Y^* \lambda_{Y*} A \rightarrow A$  is an isomorphism. This question is pro-étale local on  $Y$ , so we may assume that  $Y = X$  is a strictly totally disconnected perfectoid space. We can also replace  $Y'$  by a strictly totally disconnected perfectoid space  $X'$ . We have to see that for any strictly totally disconnected  $\tilde{X} \in X_v$ , the natural map

$$\Gamma(\lambda_{X \circ}(\tilde{X})_v, A) \rightarrow \Gamma(\tilde{X}_v, A)$$

is an isomorphism. It is enough to do this locally, so we may assume that  $\tilde{X} \rightarrow X$  lifts to a map  $\tilde{X} \rightarrow X'$ . For the proof of the displayed identity, we can also replace  $X$  by  $\lambda_{X \circ}(\tilde{X})$ , and then assume that  $X' = \tilde{X}$ . Thus,  $X$  and  $X'$  are strictly totally disconnected spaces,  $f : X' \rightarrow X$  is a  $v$ -cover such that  $\lambda_{X \circ}(X') = X$ , and  $\mathcal{F}$  is a small  $v$ -sheaf on  $X$  such that  $f^*X'$  is the pullback of a pro-étale sheaf  $\mathcal{G}'$  on  $X'$ , and we need to show that  $\mathcal{F}(X) = \mathcal{F}(X')$ . Applying Theorem 16.1 below to  $f : Y' = X' \rightarrow Y = X$ , the sheaf  $\mathcal{G}'$  on  $X'_{\text{qproét}}$  and  $\tilde{X} = X'$ , we see that the pullback map

$$p_1^* : \mathcal{F}(X') \rightarrow \mathcal{F}(X' \times_X X')$$

is an isomorphism. In particular, all maps in the commutative diagram

$$\begin{array}{ccc} \mathcal{F}(X') & & \\ p_1^* \downarrow & \searrow & \\ \mathcal{F}(X' \times_X X') & \xrightarrow{\Delta^*} & \mathcal{F}(X') \\ p_2^* \uparrow & \swarrow & \\ \mathcal{F}(X') & & \end{array}$$

are isomorphisms, and thus

$$\mathcal{F}(X) = \text{eq}(\mathcal{F}(X') \rightrightarrows \mathcal{F}(X' \times_X X')) = \text{eq}(\mathcal{F}(X') \rightrightarrows \mathcal{F}(X')) = \mathcal{F}(X') ,$$

as desired.

It remains to handle part (ii). For this, we have to see that for any  $\tilde{Y} = \varprojlim_j \tilde{Y}_j \in \text{Pro}(Y_{\text{ét}, \text{qc}, \text{sep}})$ , one has

$$\mathcal{F}(\tilde{Y}) = \varinjlim_j \mathcal{F}(\tilde{Y}_j) .$$

But if one lets  $\tilde{Y}' = \tilde{Y} \times_Y Y'$ ,  $\tilde{Y}'_j = \tilde{Y}_j \times_Y Y'$ , then

$$\mathcal{F}(\tilde{Y}') = \varinjlim_j \mathcal{F}(\tilde{Y}'_j)$$

by assumption, and a similar equation over the fibre product  $Y' \times_Y Y'$ . As equalizers commute with filtered colimits, the result follows.  $\square$

Theorem 14.12 shows that the following definition is reasonable.

**Definition 14.13.** *Let  $Y$  be a small  $v$ -stack. Define full subcategories*

$$D_{\text{ét}}(Y, \Lambda) \subset D_{\text{qproét}}(Y, \Lambda) \subset D(Y, \Lambda) = D(Y_v, \Lambda)$$

as consisting of those  $A \in D(Y, \Lambda)$  such that for all strictly totally disconnected perfectoid spaces  $f : X \rightarrow Y$ , the pullback  $f^* A \in D(X_{\text{ét}}, \Lambda)$ , resp.  $f^* A \in D(X_{\text{qproét}}, \Lambda)$ .

**Remark 14.14.** By Theorem 14.12, it suffices to check the condition for one  $v$ -cover of  $Y$ . If  $Y$  is a diamond, then  $D_{\text{qproét}}(Y, \Lambda) = D(Y_{\text{qproét}}, \Lambda)$ , if  $Y$  is a locally spatial diamond,  $D_{\text{ét}}^+(Y, \Lambda) = D^+(Y_{\text{ét}}, \Lambda)$ , and if  $Y$  is a strictly totally disconnected perfectoid space,  $D_{\text{ét}}(Y, \Lambda) = D(Y_{\text{ét}}, \Lambda)$ . If  $Y$  is a locally spatial diamond, then in general  $D_{\text{ét}}(Y, \Lambda) \neq D(Y_{\text{ét}}, \Lambda)$ , but one has the following result.

**Proposition 14.15.** *Let  $Y$  be a small  $v$ -stack. Then  $D_{\text{ét}}(Y, \Lambda)$  and  $D_{\text{qproét}}(Y, \Lambda)$  are left-complete. If  $Y$  is a locally spatial diamond, then  $D_{\text{ét}}(Y, \Lambda)$  is the left-completion of  $D(Y_{\text{ét}}, \Lambda)$ .*

*Proof.* The first part is clear. For the final statement, the proof is the same as in [BS15, Proposition 5.3.2].  $\square$

Containment in these subcategories can be checked on cohomology sheaves.

**Proposition 14.16.** *Let  $Y$  be a small  $v$ -stack and  $A \in D(Y, \Lambda)$ . Then  $A \in D_{\text{ét}}(Y, \Lambda)$  (resp.  $A \in D_{\text{qproét}}(Y, \Lambda)$ ) if and only if for all  $i \in \mathbb{Z}$ , the  $v$ -sheaf  $\mathcal{H}^i(C)[0] \in D_{\text{ét}}(Y, \Lambda)$  (resp.  $\mathcal{H}^i(C)[0] \in D_{\text{qproét}}(Y, \Lambda)$ ).*

*Proof.* As all relevant derived categories are left-complete, one can assume that  $A$  is bounded below. Then  $A = \varinjlim \tau^{\leq n} A$ , so one can assume that  $A$  is bounded. As the inclusions  $D_{\text{ét}}(Y, \Lambda) \subset D_{\text{qproét}}(Y, \Lambda) \subset D(Y, \Lambda)$  are fully faithful, one can further reduce to the case where  $A$  is concentrated in one degree, as desired.  $\square$

Moreover, the inclusion  $D_{\text{qproét}}(Y, \Lambda) \subset D(Y, \Lambda)$  can be described as those sheaves that are invariant under change of algebraically base field, in the following sense.

**Proposition 14.17.** *A complex  $A \in D(Y, \Lambda)$  lies in  $D_{\text{qproét}}(Y, \Lambda)$  if and only if for all maps of strictly totally disconnected perfectoid spaces  $\tilde{X} \rightarrow X$  in  $Y_v$  which induce a homeomorphism  $|\tilde{X}| \rightarrow |X|$ , the map*

$$R\Gamma(X, A) \rightarrow R\Gamma(\tilde{X}, A)$$

*is a quasi-isomorphism.*

*In particular, if  $I$  is any category and  $F \in D(Y^I, \Lambda)$  is in the derived category of functors from  $I$  to small sheaves of abelian groups on  $Y_v$ , such that for all  $i \in I$ , the complex  $F(i) \in D_{\text{qproét}}(Y, \Lambda)$ , then the (derived) limit  $\lim_I F \in D(Y, \Lambda)$  also satisfies  $\lim_I F \in D_{\text{qproét}}(Y, \Lambda)$ .*

*Proof.* For the first part, we can assume that  $Y$  is strictly totally disconnected. In that case, we first claim that for all  $E \in D(Y_{\text{qproét}}, \Lambda)$  and all strictly totally disconnected  $\tilde{Y} \in Y_v$ , one has

$$R\Gamma(\tilde{Y}_v, \lambda_Y^* E) = R\Gamma(\tilde{Y}_{\text{qproét}}, E|_{\tilde{Y}}) = R\Gamma(\lambda_{Y \circ}(\tilde{Y})_{\text{qproét}}, E),$$

where in the middle  $E|_{\tilde{Y}}$  denotes the pullback of  $E$  to  $D(\tilde{Y}_{\text{qproét}}, \Lambda)$ . Here, the first equation follows from Proposition 14.10, and the second equation from the equivalence  $\tilde{Y}_{\text{qproét}} \cong \lambda_{Y \circ}(\tilde{Y})_{\text{qproét}}$ , which results from the identification  $|\tilde{Y}| = |\lambda_{Y \circ}(\tilde{Y})|$  and Corollary 7.22.

In particular, we see that if  $A \in D_{\text{qproét}}(Y, \Lambda)$ , then indeed for all maps of strictly totally disconnected perfectoid spaces  $\tilde{X} \rightarrow X$  in  $Y_v$  which induce a homeomorphism  $|\tilde{X}| = |X|$  (equivalently,  $\lambda_{X \circ}(\tilde{X}) = X$ ), one has

$$R\Gamma(X, A) = R\Gamma(\tilde{X}, A).$$

For the converse, one needs to check that the map  $A \rightarrow \lambda_Y^* R\lambda_{Y*} A$  is an isomorphism, or equivalently that for all strictly totally disconnected  $\tilde{Y} \in Y_v$ , one has

$$R\Gamma(\tilde{Y}_v, A) = R\Gamma(\tilde{Y}, \lambda_Y^* R\lambda_{Y*} A) = R\Gamma(\lambda_{Y \circ}(\tilde{Y})_v, A),$$

using the first paragraph in the last equality. But this follows from the assumption.

The final statement is a formal consequence, noting that global sections commute with homotopy limits.  $\square$

## 15. ANALYTIC ADIC SPACES AS DIAMONDS

There is a natural functor from rigid-analytic varieties over  $\mathbb{Q}_p$  to locally spatial diamonds. In fact, more generally, there is a functor from analytic adic spaces over  $\mathbb{Z}_p$  to locally spatial diamonds. Our goal in this section is to construct this functor, and show that it preserves the étale site.

We start with the affinoid case.

**Lemma 15.1.** (i) *The functor  $\text{Spd } \mathbb{Z}_p$  taking  $X \in \text{Perf}$  to the set of pairs  $(X^\sharp, \iota)$  up to isomorphism, where  $X^\sharp$  is a perfectoid space and  $\iota : (X^\sharp)^\flat \cong X$  is an identification of  $X^\sharp$  as an untilt of  $X$ , is a v-sheaf.*

(ii) *Let  $A$  be a Tate  $\mathbb{Z}_p$ -algebra, and let  $A^+ \subset A$  be an open and integrally closed subring. The functor  $\text{Spd}(A, A^+)$  taking  $X \in \text{Perf}$  to the set of pairs*

$$((X^\sharp, \iota), f : (A, A^+) \rightarrow (\mathcal{O}_{X^\sharp}, \mathcal{O}_{X^\sharp}^+))$$

*up to isomorphism, where  $(X^\sharp, \iota)$  is as in (i), and  $f$  is a continuous map of pairs of topological algebras, is a v-sheaf.*

*Proof.* Part (ii) follows from part (i) and Theorem 8.7. For part (i), we first have to see that this in fact defines a presheaf: It is not a priori clear what the maps are. But if  $f : X' \rightarrow X$  is a map of perfectoid spaces and  $X^\sharp$  is an untilt of  $X$ , then by Corollary 3.20, perfectoid spaces over  $X$  are equivalent to perfectoid spaces over  $X^\sharp$ ; then we define  $(X')^\sharp$  to be the perfectoid space over  $X^\sharp$  corresponding to  $X' \rightarrow X$ . Also, note that pairs  $(X^\sharp, \iota)$  have no automorphisms, by the same argument.

To see that this is a v-sheaf, we have to see that if  $X = \text{Spa}(R, R^+)$  is an affinoid perfectoid space of characteristic  $p$  with a v-cover  $Y = \text{Spa}(S, S^+) \rightarrow X$ , and  $Y^\sharp = \text{Spa}(S^\sharp, S^{\sharp+})$  is an untilt

of  $Y$  such that the two corresponding untits of  $Z := Y \times_X Y = \mathrm{Spa}(T, T^+)$  agree, then there is a unique untit  $X^\sharp = \mathrm{Spa}(R^\sharp, R^{\sharp+})$  whose pullback to  $Y$  is  $Y^\sharp$ .

Fix a pseudouniformizer  $\varpi \in R$ , and consider the map  $f : W(R^+) \rightarrow W(S^+) \rightarrow S^{\sharp+}$ , where the second map is Fontaine's map  $\theta : W(S^+) \rightarrow S^{\sharp+}$ . Then  $f([\varpi]) \in S^{\sharp+}$  is a pseudouniformizer. In particular, since  $p$  is topologically nilpotent in  $S^\sharp$ , there is some  $N$  such that  $f([\varpi])$  divides  $p^N$ . Replacing  $\varpi$  by a  $p$ -power root, we may assume that  $N = 0$ . In that case,  $S^+/\varpi \cong S^{\sharp+}/f([\varpi])$ , and similarly for  $T^+$ . It follows that

$$(R^+/\varpi)^a = \mathrm{eq}((S^+/\varpi)^a \rightrightarrows (T^+/\varpi)^a) = \mathrm{eq}((S^{\sharp+}/f([\varpi]))^a \rightrightarrows (T^{\sharp+}/f([\varpi]))^a).$$

Moreover, extending to a long exact sequence, the higher cohomology groups vanish by Theorem 8.7. We see that

$$(W(R^+)/[\varpi])^a \rightarrow \mathrm{eq}((S^{\sharp+}/f([\varpi]))^a \rightrightarrows (T^{\sharp+}/f([\varpi]))^a)$$

is surjective, with kernel generated by  $p$ . Now the vanishing of higher cohomology groups and the five lemma imply inductively that one gets surjective maps

$$(W(R^+)/[\varpi]^n)^a \rightarrow \mathrm{eq}((S^{\sharp+}/f([\varpi])^n)^a \rightrightarrows (T^{\sharp+}/f([\varpi])^n)^a)$$

for all  $n \geq 1$ , whose kernel is generated by an element  $\xi \in W(R^+)$  with  $\xi \equiv p \pmod{[\varpi]}$ . In the limit, one gets an isomorphism

$$(W(R^+)/\xi)^a = \mathrm{eq}(S^{\sharp+a} \rightrightarrows T^{\sharp+a}).$$

In particular,  $R^\sharp := (W(R^+)/\xi)[[\varpi]^{-1}]$  is a perfectoid Tate ring which satisfies

$$R^\sharp = \mathrm{eq}(S^\sharp \rightrightarrows T^\sharp),$$

which is an untit of  $R$ , and then comes with a canonical  $R^{\sharp+} \subset R^\sharp$  determined by  $R^+ \subset R$ . It follows that  $X^\sharp = \mathrm{Spa}(R^\sharp, R^{\sharp+})$  is the unique untit.  $\square$

In case  $A$  is a perfectoid Tate ring, it is easy to identify  $\mathrm{Spd}(A, A^+)$ .

**Lemma 15.2.** *Let  $A$  be a perfectoid Tate ring with an open and integrally closed subring  $A^+ \subset A$ . Then  $\mathrm{Spd}(A, A^+)$  is representable by the affinoid perfectoid space  $\mathrm{Spa}(A^b, A^{b+})$ .*

*Proof.* This is a direct consequence of Corollary 3.20.  $\square$

To understand  $\mathrm{Spd}(A, A^+)$  in general, we use the following general lemma.

**Lemma 15.3.** *Let  $A$  be any ring. Then one can find a cofiltered inverse system of finite groups  $G_i$  with surjective transition maps, and a compatible filtered direct system of finite étale  $G_i$ -torsors  $A \rightarrow A_i$  such that  $A_\infty = \varinjlim_i A_i$  has no nonsplit finite étale covers.*

*If  $A$  is a Tate  $\mathbb{Z}_p$ -algebra, then the uniform completion  $\widehat{A}_\infty$  of  $A_\infty$  is perfectoid.*

Results of this type were proved for example by Colmez, [Col02], and Faltings, [Fal02].

*Proof.* The first part is standard. To see that  $R = \widehat{A}_\infty$  is perfectoid we find first a pseudo-uniformizer  $\varpi \in R$  such that  $\varpi^p | p$  in  $R^\circ$ . To do this, let  $\varpi_0 \in R$  be any pseudo-uniformizer. Let  $n$  be large enough so that  $\varpi_0 | p^n$ . Now look at the equation  $x^{p^n} - \varpi_0 x = \varpi_0$ . This determines a finite étale  $R$ -algebra, and so it admits a solution  $x = \varpi \in R$ . Note that  $\varpi^{p^n} | \varpi_0$  in  $R^\circ$ , and  $\varpi_1$  is a unit in  $R$ . As  $p^{n-1} \geq n$ , we see that  $\varpi^p | p$ , as desired.

Now we must check that  $\Phi: R^\circ/\varpi \rightarrow R^\circ/\varpi^p$  is surjective. Let  $f \in R^\circ$ , and consider the equation  $x^p - \varpi^p x - f$ . This determines a finite étale  $R$ -algebra, which consequently has a section, i.e. there is some  $x \in R^\circ$  with  $x^p - \varpi^p x = f$ . But then  $x^p \equiv f \pmod{\varpi^p R^\circ}$ , as desired.  $\square$

Now we can prove that  $\mathrm{Spd}(A, A^+)$  is always a spatial diamond.

**Proposition 15.4.** *Let  $A$  be a Tate  $\mathbb{Z}_p$ -algebra with an open and integrally closed subring  $A^+ \subset A$ , and choose a cofiltered inverse system of finite groups  $G_i$  with surjective transition maps, and a compatible filtered direct system of finite étale  $G_i$ -torsors  $A \rightarrow A_i$  as in Lemma 15.3. Let  $A_i^+ \subset A_i$  be the integral closure of  $A^+$ . Also, let  $\widehat{A_\infty^+} \subset \widehat{A_\infty}$  be the closure of  $\varprojlim_i A_i^+$  in  $\widehat{A_\infty}$ .*

*Then  $\mathrm{Spd}(A_i, A_i^+) \rightarrow \mathrm{Spd}(A, A^+)$  is a  $G_i$ -torsor of  $v$ -sheaves, and*

$$\mathrm{Spd}(\widehat{A_\infty}, \widehat{A_\infty^+}) = \varprojlim_i \mathrm{Spd}(A_i, A_i^+) \rightarrow \mathrm{Spd}(A, A^+)$$

*is a  $G$ -torsor, where  $G = \varprojlim_i G_i$ . In particular,*

$$\mathrm{Spa}(\widehat{A_\infty}^b, \widehat{A_\infty^+}^b) = \mathrm{Spd}(\widehat{A_\infty}, \widehat{A_\infty^+}) \rightarrow \mathrm{Spd}(A, A^+)$$

*is a universally open qcqs quasi-pro-étale map from an affinoid perfectoid space, so that  $\mathrm{Spd}(A, A^+)$  is a spatial diamond, with*

$$|\mathrm{Spd}(A, A^+)| = |\mathrm{Spa}(\widehat{A_\infty}^b, \widehat{A_\infty^+}^b)|/G = |\mathrm{Spa}(\widehat{A_\infty}, \widehat{A_\infty^+})|/G = |\mathrm{Spa}(A, A^+)|.$$

*Proof.* It follows from Theorem 6.1 that  $\mathrm{Spd}(A_i, A_i^+) \rightarrow \mathrm{Spd}(A, A^+)$  is a  $G_i$ -torsor. The rest, except for the identification of the topological space, follows formally from Lemma 15.3, Lemma 15.2, Lemma 10.13 and Proposition 11.24.

For the identification of the topological space, only the last identification needs justification. But

$$|\mathrm{Spa}(\widehat{A_\infty}, \widehat{A_\infty^+})| = \varprojlim_i |\mathrm{Spa}(A_i, A_i^+)|,$$

and for each  $i$ ,  $|\mathrm{Spa}(A_i, A_i^+)|/G_i = |\mathrm{Spa}(A, A^+)|$ . Passing to the limit gives the result.  $\square$

One can now glue the functor  $(A, A^+) \mapsto \mathrm{Spd}(A, A^+)$ , and pass to adic spaces. Indeed, if  $U \subset \mathrm{Spa}(A, A^+)$  is a rational open subset, then

$$\mathrm{Spd}(\mathcal{O}(U), \mathcal{O}^+(U)) \rightarrow \mathrm{Spd}(A, A^+)$$

is the open subfunctor of  $\mathrm{Spd}(A, A^+)$  corresponding to the open subset  $U \subset |\mathrm{Spa}(A, A^+)| = |\mathrm{Spd}(A, A^+)|$ .

**Definition 15.5.** *Let  $Y$  be an analytic adic space over  $\mathbb{Z}_p$ . The diamond associated with  $Y$  is the  $v$ -sheaf defined by*

$$Y^\diamond : X \mapsto \{((X^\sharp, \iota), f : X^\sharp \rightarrow Y)\} / \cong,$$

*where  $X^\sharp$  is a perfectoid space with an isomorphism  $\iota : (X^\sharp)^b \cong X$ .*

We note that this definition (and the next lemma) work in the generality of adic spaces as defined in [SW13, Section 2.1].

**Lemma 15.6.** *Let  $Y$  be an analytic adic space over  $\mathbb{Z}_p$ . Then  $Y^\diamond$  is a locally spatial diamond, with  $|Y^\diamond| = |Y|$ . Moreover, with  $Y_{\text{ét}}$  and  $Y_{\text{fét}}$  defined as in [KL15, Definition 8.2.19], one has equivalences of sites  $Y_{\text{ét}}^\diamond \cong Y_{\text{ét}}$  and  $Y_{\text{fét}}^\diamond \cong Y_{\text{fét}}$ .*



*Proof.* All statements reduce readily to the affinoid case  $Y = \mathrm{Spa}(A, A^+)$ , where the first part follows from Proposition 15.4. Now, for the equivalence of finite étale sites, note that by [KL15, Lemma 8.2.17], we need to see that the category of finite étale  $A$ -algebras is equivalent to (the opposite of)  $Y_{\mathrm{fét}}^{\diamond}$ . Choosing a  $\underline{G}$ -torsor as in Proposition 15.4, this follows from the equivalence

$$(\widehat{A}_\infty)_{\mathrm{fét}} = 2\text{-}\varinjlim_i (A_i)_{\mathrm{fét}}$$

from [Sch12, Lemma 7.5 (i)] (and similar results

$$(C^0(G, \widehat{A}_\infty))_{\mathrm{fét}} = 2\text{-}\varinjlim_i (C^0(G_i, A_i))_{\mathrm{fét}},$$

$$(C^0(G \times G, \widehat{A}_\infty))_{\mathrm{fét}} = 2\text{-}\varinjlim_i (C^0(G_i \times G_i, A_i))_{\mathrm{fét}}$$

for the algebras corresponding to the fibre products) and usual descent along the finite étale  $G_i$ -torsors  $A \rightarrow A_i$ .

Finally, the case of the étale site follows by combining the description of the topological space with the finite étale case, as by Lemma 11.31 resp. [KL15, Definition 8.2.16], in both cases an étale map is locally given by a composite of a quasicompact open immersion and a finite étale map.  $\square$

## 16. GENERAL BASE CHANGE RESULTS

In this section, we establish some general base change results. We will first prove results comparing étale, pro-étale and v-pushforwards, and then deduce more classical base change results.

**Theorem 16.1.** *Let  $f : Y' \rightarrow Y$  be a map of diamonds, and consider the diagram of sites*

$$\begin{array}{ccc} Y'_v & \xrightarrow{\lambda_{Y'}} & Y'_{\mathrm{qproét}} \\ f_v \downarrow & & \downarrow f_{\mathrm{qproét}} \\ Y_v & \xrightarrow{\lambda_Y} & Y_{\mathrm{qproét}}. \end{array}$$

*Let  $\mathcal{F}$  be a small sheaf of abelian groups on  $Y'_{\mathrm{qproét}}$ . Then the base change morphism*

$$\lambda_Y^* R^i f_{\mathrm{qproét}*} \mathcal{F} \rightarrow R^i f_{v*} \lambda_{Y'}^* \mathcal{F}$$

*is an isomorphism in the following cases:*

- (i) *if  $i = 0$ , or*
- (ii) *for all  $i \geq 0$  if  $f$  is quasi-pro-étale, or*
- (iii) *for all  $i \geq 0$  if there is some integer  $n$  prime to  $p$  such that  $n\mathcal{F} = 0$ .*

*Moreover, under the same conditions, if  $Y = X$  and  $\widetilde{X} \in X_v$  are strictly totally disconnected and  $\widetilde{X} \rightarrow \lambda_{X^\circ}(\widetilde{X}) \rightarrow X$  is the factorization from Lemma 14.5, then the natural map*

$$H^i((\lambda_{X^\circ}(\widetilde{X}) \times_X Y')_{\mathrm{qproét}}, \mathcal{F}) \rightarrow H^i((\widetilde{X} \times_X Y')_v, \lambda_{Y'}^* \mathcal{F})$$

*is an isomorphism.*

**Remark 16.2.** The final statement combines an “invariance under change of algebraically closed base field” statement with a “pro-étale cohomology = v-cohomology”-statement. Indeed, if  $X$  and  $\tilde{X}$  are geometric points, then  $\lambda_{X^\circ}(\tilde{X}) = X$ , and the statement becomes

$$H^i(Y'_{\text{qproét}}, \mathcal{F}) = H^i(Y'_v, \lambda_{Y'}^* \mathcal{F}) = H^i((\tilde{X} \times_X Y')_v, \lambda_{Y'}^* \mathcal{F}) .$$

*Proof.* The claim is pro-étale local on  $Y$ , so we can assume that  $Y = X$  is strictly totally disconnected, and it suffices to prove the final statement, as this gives the statement about sheaves by sheafification.

The final statement is again pro-étale local on  $Y'$ , so we may also assume that  $Y' = X'$  is strictly totally disconnected. In case (ii), note that  $\tilde{X} \times_X X'$  is again strictly totally disconnected, and so

$$\begin{aligned} H^i((\lambda_{X^\circ}(\tilde{X}) \times_X X')_{\text{qproét}}, \mathcal{F}) &= H^i(\lambda_{X'^\circ}(\tilde{X} \times_X X')_{\text{qproét}}, \mathcal{F}) \\ &= H^i((\tilde{X} \times_X X')_{\text{qproét}}, \mathcal{F}) \\ &= H^i((\tilde{X} \times_X X')_v, \mathcal{F}) , \end{aligned}$$

where we have written  $\mathcal{F}$  also for any of its pullbacks. Here, the first equation follows from the definitions, the second equation follows from Corollary 7.22 and

$$|\lambda_{X'^\circ}(\tilde{X} \times_X X')| = |\tilde{X} \times_X X'| ,$$

and the final equation from Proposition 14.7. It remains to handle parts (i) and (iii).

We know from Lemma 14.4 that  $\lambda_{Y'}$  commutes with all limits. Moreover, limits are exact in all relevant categories. As  $\mathcal{F}$  is small, we can find a set of affinoid pro-étale perfectoid spaces  $f_j : Y'_j \rightarrow Y'$  and injective abelian groups  $M_j$  (annihilated by  $n$  in case (iii)) such that  $\mathcal{F}$  injects into  $\prod_j f_{j\text{qproét}*} M_j$ . An easy dévissage then reduces us to the case  $\mathcal{F} = Rf_{j\text{qproét}*} M_j$ . Replacing  $Y'$  by  $Y'_j$  and  $f$  by the composite  $f \circ f_j$  (and using case (ii)) reduces us further to the case that  $\mathcal{F} = M$  is the constant sheaf associated with some abelian group  $M$  (killed by  $n$  in case (iii)).

It remains to see that if  $M$  is an abelian group and  $\tilde{X} \rightarrow X \leftarrow X'$  is a diagram of strictly totally disconnected perfectoid spaces, then the natural map

$$H^i((\lambda_{X^\circ}(\tilde{X}) \times_X X')_{\text{qproét}}, M) \rightarrow H^i((\tilde{X} \times_X X')_v, M)$$

is an isomorphism for  $i = 0$ , and for all  $i \geq 0$  if  $nM = 0$  for some  $n$  prime to  $p$ . Here, we may replace pro-étale and v-cohomology by étale cohomology by Proposition 14.7 and Proposition 14.8. Let  $g : \tilde{X} \times_X X' \rightarrow \lambda_{X^\circ}(\tilde{X}) \times_X X'$  be the natural map. It suffices to see that  $g_{\text{ét}*} M = M$  and  $R^i g_{\text{ét}*} M = 0$  for  $i > 0$  if  $nM = 0$  for some  $n$  prime to  $p$ . This can be checked on stalks. This reduces us to the next lemma.  $\square$

**Lemma 16.3.** *Let  $X_i = \text{Spa}(C_i, C_i^+)$  for  $i = 1, 2, 3$ , where  $C_i$  is an algebraically closed nonarchimedean field with an open and bounded valuation subring  $C_i^+ \subset C_i$ , and let  $X_1 \rightarrow X_3 \leftarrow X_2$  be a diagram. Then, for any abelian group  $M$ ,*

$$H^0((X_1 \times_{X_3} X_2)_{\text{ét}}, M) = M$$

and if  $nM = 0$  for some  $n$  prime to  $p$ , then

$$H^i((X_1 \times_{X_3} X_2)_{\text{ét}}, M) = 0$$

for all  $i > 0$ .

*Proof.* The statement for  $H^0$  follows from Lemma 14.6. For the statement about higher cohomology, we prove more generally that if  $X_2$  is any perfectoid space over  $X_3$ , then

$$H^i((X_1 \times_{X_3} X_2)_{\text{ét}}, M) = H^i(X_{2,\text{ét}}, M)$$

for all  $i \geq 0$ . We can write  $X_2 = \varprojlim_j \text{Spa}((R_j)^{\text{perf}}, (R_j^+)^{\text{perf}})$  as a cofiltered inverse limit, where each  $R_j$  is topologically of finite type over  $C_3$ . Applying Proposition 14.9, it is enough to prove the result for all  $\text{Spa}((R_j)^{\text{perf}}, (R_j^+)^{\text{perf}})$ , so we may assume  $X_2 = \text{Spa}(R^{\text{perf}}, (R^+)^{\text{perf}})$ , where  $R$  is topologically of finite type over  $C$ . Now the result follows from [Hub96, Theorem 4.1.1 (c)], using the equivalence of sites  $X_{2,\text{ét}} = \text{Spa}(R, R^+)_{\text{ét}}$ , which follows for example from Lemma 15.6 and the observation  $\text{Spa}(R, R^+)^\diamond = \text{Spa}(R^{\text{perf}}, (R^+)^{\text{perf}})$ .  $\square$

Let us note the following consequence for derived categories.

**Corollary 16.4.** *Let  $f : Y' \rightarrow Y$  be a map of diamonds. Assume that  $f$  is quasi-pro-étale, or  $n\Lambda = 0$  for some  $n$  prime to  $p$ . Then for any  $A \in D(Y'_{\text{qproét}}, \Lambda)$ , the base change morphism*

$$\lambda_Y^* Rf_{\text{qproét}*} A \rightarrow Rf_{v*} \lambda_{Y'}^* A$$

*in  $D(Y_v, \Lambda)$  is an isomorphism.*

*Proof.* The statement is local in  $Y_{\text{qproét}}$ , so we may assume  $Y = X$  is a strictly totally disconnected perfectoid space. We prove the finer statement that for all  $\tilde{X} \in X_v$  strictly totally disconnected,

$$R\Gamma((\lambda_{X_c}(\tilde{X}) \times_X Y')_{\text{qproét}}, A) = R\Gamma((\tilde{X} \times_X Y')_v, \lambda_{Y'}^* A).$$

For this, note that both sides commute with the Postnikov limit  $A = R\varprojlim_n \tau^{\geq -n} A$  by Proposition 14.11.  $\square$

**Corollary 16.5.** *Let  $f : Y' \rightarrow Y$  be a map of small  $v$ -stacks. Assume that  $f$  is quasi-pro-étale, or  $n\Lambda = 0$  for some  $n$  prime to  $p$ . Then for any  $A \in D_{\text{qproét}}(Y', \Lambda)$ , the pushforward  $Rf_{v*} C$  lies in  $D_{\text{qproét}}(Y, \Lambda)$ . If  $Y$  and  $Y'$  are diamonds, then under the identifications  $D_{\text{qproét}}(Y', \Lambda) = D(Y'_{\text{qproét}}, \Lambda)$  and  $D_{\text{qproét}}(Y, \Lambda) = D(Y_{\text{qproét}}, \Lambda)$ , one has  $Rf_{v*} = Rf_{\text{qproét}*}$ .*

*Proof.* We may assume that  $Y$  is a diamond. If  $f$  is quasi-pro-étale, then  $Y'$  is a diamond, and  $D_{\text{qproét}}(Y', \Lambda) = D(Y'_{\text{qproét}}, \Lambda)$ , so the result follows directly from Corollary 16.4.

If  $n\Lambda = 0$  for some  $n$  prime to  $p$ , choose a perfectoid space  $X'$  with a surjection  $X' \rightarrow Y'$ . Let  $X'_\bullet \rightarrow Y'$  be the corresponding simplicial nerve, which is a simplicial diamond. Let  $g_\bullet : X'_\bullet \rightarrow Y$  be the resulting map. Then  $Rf_{v*} C$  is the derived limit of the simplicial object  $Rg_{\bullet*} A|_{X'_\bullet}$  (more precisely, of the object in the derived category  $D(X'_{\bullet,v}, \Lambda)$  of the simplicial topos  $X'_{\bullet,v}$ ). By Corollary 16.4, all  $Rg_{i*} A|_{X'_i} \in D(Y_{\text{qproét}}, \Lambda)$ . But  $D(Y_{\text{qproét}}, \Lambda) \subset D(Y_v, \Lambda)$  is closed under derived limits by Proposition 14.17, so the result follows.  $\square$

We need a similar result relating the étale and pro-étale pushforward.

**Proposition 16.6.** *Let  $f : Y' \rightarrow Y$  be a qcqs morphism of locally spatial diamonds. Let  $\mathcal{F}$  be a sheaf of abelian groups on  $Y'_{\text{ét}}$ . Then the base change morphism*

$$\nu_Y^* R^i f_{\text{ét}*} \mathcal{F} \rightarrow R^i f_{\text{qproét}*} \nu_{Y'}^* \mathcal{F}$$

*of sheaves of abelian groups on  $Y_{\text{qproét}}$  is an isomorphism for all  $i \geq 0$ .*

*Proof.* The statement is étale local on  $Y$ , so we can assume that  $Y$  is spatial. By the proof of Proposition 14.8, it suffices to check that for all  $\tilde{Y} = \varprojlim_j \tilde{Y}_j \rightarrow Y$  in  $\text{Pro}(Y_{\text{ét}, \text{qc}, \text{sep}})$ , one has

$$H^i((\tilde{Y} \times_Y Y')_{\text{qproét}}, \mathcal{F}) = \varinjlim_j H^i((\tilde{Y}_j \times_Y Y')_{\text{ét}}, \mathcal{F}),$$

where we denote by  $\mathcal{F}$  also any of its base changes. But by Proposition 14.8, the left-hand side is given by  $H^i((\tilde{Y} \times_Y Y')_{\text{ét}}, \mathcal{F})$ , and then Proposition 14.9 gives the result.  $\square$

Again, we can give derived consequences.

**Corollary 16.7.** *Let  $f : Y' \rightarrow Y$  be a qcqs map of locally spatial diamonds. Then for any  $A \in D^+(Y'_{\text{ét}}, \Lambda)$ , the base change morphism*

$$\nu_Y^* Rf_{\text{ét}*} A \rightarrow Rf_{\text{qproét}*} \nu_{Y'}^* A$$

*in  $D^+(Y_{\text{qproét}}, \Lambda)$  is an isomorphism. If  $Y$  and  $Y'$  are strictly totally disconnected perfectoid spaces, then the same result holds true more generally for any  $A \in D(Y'_{\text{ét}}, \Lambda)$ .*

*Proof.* The first part is a direct consequence of Proposition 16.6. If  $Y$  and  $Y'$  are strictly totally disconnected perfectoid spaces, then as in Corollary 16.4, the statement follows from the convergence results in the proof of Proposition 14.10.  $\square$

**Corollary 16.8.** *Let  $f : Y' \rightarrow Y$  be a qcqs map of small v-stacks.*

- (i) *If  $f$  is quasi-pro-étale, then for any  $A \in D_{\text{ét}}(Y', \Lambda)$ , the pushforward  $Rf_{v*} A$  lies in  $D_{\text{ét}}(Y, \Lambda)$ .*
- (ii) *If  $n\Lambda = 0$  for some  $n$  prime to  $p$ , then for any  $A \in D_{\text{ét}}^+(Y', \Lambda)$ , the pushforward  $Rf_{v*} A$  lies in  $D_{\text{ét}}^+(Y, \Lambda)$ . If  $Y$  and  $Y'$  are locally spatial, then under the identifications  $D_{\text{ét}}^+(Y', \Lambda) = D^+(Y'_{\text{ét}}, \Lambda)$  and  $D_{\text{ét}}^+(Y, \Lambda) = D^+(Y_{\text{ét}}, \Lambda)$ , one has  $Rf_{v*} = Rf_{\text{ét}*}$ .*

*Proof.* We may assume that  $Y$  is a strictly totally disconnected perfectoid space. In case (i),  $Y'$  is a qcqs perfectoid space. If  $Y'$  is separated, then it is strictly totally disconnected by Lemma 7.19, and the result follows from Corollary 16.7 and Corollary 16.5. In general, there is a finite quasicompact open cover of  $Y'$  by affinoid perfectoid spaces, where all intersections are separated. Using the corresponding Čech cover, we get the result in general.

For part (ii), choose an affinoid perfectoid space  $X'$  with a surjection  $X' \rightarrow Y'$ . Let  $X'_\bullet \rightarrow Y'$  be the corresponding simplicial nerve, which is a simplicial qcqs v-sheaf. If  $Y'$  was already a v-sheaf, then in fact it is a simplicial spatial diamond; assume that we are in this case for the moment. Let  $g_\bullet : X'_\bullet \rightarrow Y$  be the resulting map. Then  $Rf_{v*} C$  is the limit of the simplicial object  $Rg_{\bullet, v*} C|_{X'_\bullet}$ . By Corollary 16.7 and Corollary 16.4, all  $Rg_{i, v*} C|_{X'_i} \in D^+(Y_{\text{ét}}, \Lambda)$ , and there is some  $n$  such that all of them lie in  $D^{\geq -n}(Y_{\text{ét}}, \Lambda)$ . This implies that the derived limit  $Rf_{v*} C$  also lies in  $D^+(Y_{\text{ét}}, \Lambda)$ .

In general (if  $Y'$  is any qcqs v-stack), you repeat the same argument, using that the result is now known for qcqs v-sheaves.  $\square$

Moreover, we get the following classical base change results.

**Corollary 16.9.** *Let*

$$\begin{array}{ccc} Y' & \xrightarrow{f'} & Y \\ g' \downarrow & & \downarrow g \\ X' & \xrightarrow{f} & X \end{array}$$

be a cartesian diagram of diamonds, and let  $\mathcal{F}$  be a small sheaf of abelian groups on  $Y_{\text{qproét}}$ . Then the base change morphism

$$f_{\text{qproét}}^* R^i g_{\text{qproét}*} \mathcal{F} \rightarrow R^i g'_{\text{qproét}*} f'_{\text{qproét}*} \mathcal{F}$$

is an isomorphism in the following cases:

- (i) if  $i = 0$ , or
- (ii) for all  $i \geq 0$  if  $f$  is quasi-pro-étale, or
- (iii) for all  $i \geq 0$  if  $n\mathcal{F} = 0$  for some  $n$  prime to  $p$ .

Moreover, if  $f$  is quasi-pro-étale or  $n\Lambda = 0$  for some  $n$  prime to  $p$ , then for any  $A \in D(Y_{\text{qproét}}, \Lambda)$ , the base change morphism

$$f_{\text{qproét}}^* Rg_{\text{qproét}*} A \rightarrow Rg'_{\text{qproét}*} f'_{\text{qproét}*} A$$

is an isomorphism.

*Proof.* The similar result is formal if one replaces the pro-étale site by the v-site everywhere, as then  $X'_v$  is the slice  $X_v/X'$  (and similarly for  $Y'_v$ ), and one always has base change for slice topoi. However, one can reduce from the pro-étale site to the v-site by Theorem 16.1 (and Proposition 14.7). The final statement follows similarly from Proposition 14.10 and Corollary 16.4.  $\square$

**Corollary 16.10.** *Let*

$$\begin{array}{ccc} Y' & \xrightarrow{f'} & Y \\ g' \downarrow & & \downarrow g \\ X' & \xrightarrow{f} & X \end{array}$$

be a cartesian diagram of locally spatial diamonds where  $g$  is qcqs, and let  $\mathcal{F}$  be a sheaf of abelian groups on  $Y_{\text{ét}}$ . Then the base change morphism

$$f_{\text{ét}}^* R^i g_{\text{ét}*} \mathcal{F} \rightarrow R^i g'_{\text{ét}*} f'_{\text{ét}*} \mathcal{F}$$

is an isomorphism in the following cases:

- (i) if  $i = 0$ , or
- (ii) for all  $i \geq 0$  if  $f$  is quasi-pro-étale, or
- (iii) for all  $i \geq 0$  if  $n\mathcal{F} = 0$  for some  $n$  prime to  $p$ .

Moreover, if  $f$  is quasi-pro-étale or  $n\Lambda = 0$  for some  $n$  prime to  $p$ , then for any  $A \in D^+(Y'_{\text{ét}}, \Lambda)$ , the base change morphism

$$f_{\text{ét}}^* Rg_{\text{ét}*} A \rightarrow Rg'_{\text{ét}*} f'_{\text{ét}*} A$$

is an isomorphism.

*Proof.* This follows from Corollary 16.9 and Proposition 16.6. Note that here the final statement is a formal consequence of the statement about sheaves.  $\square$

## 17. FOUR FUNCTORS

We will use the formalism of the previous section to set up the first parts of our six functor formalism. This will notably introduce a pushforward functor  $Rf_*$  that is in general different from any of the functors  $Rf_{v*}$ ,  $Rf_{\text{qproét}*}$  and  $Rf_{\text{ét}*}$ , and we will clarify their relation.

First, recall that for any small  $v$ -stack  $Y$ , we have defined the full subcategory  $D_{\text{ét}}(Y, \Lambda) \subset D(Y_v, \Lambda)$ . We will at certain points invoke Lurie's  $\infty$ -categorical adjoint functor theorem. For this reason, we need to upgrade our constructions to functors of  $\infty$ -categories at certain points.

**Lemma 17.1.** *There is a (natural) presentable stable  $\infty$ -category  $\mathcal{D}_{\text{ét}}(Y, \Lambda)$  whose homotopy category is  $D_{\text{ét}}(Y, \Lambda)$ . More precisely, the  $\infty$ -derived category  $\mathcal{D}(Y_v, \Lambda)$  of  $\Lambda$ -modules on  $Y_v$  is a presentable stable  $\infty$ -category, and  $\mathcal{D}_{\text{ét}}(Y, \Lambda)$  is a full presentable stable  $\infty$ -subcategory closed under all colimits.*

*Proof.* First,  $\mathcal{D}(Y_v, \Lambda)$  is a presentable stable  $\infty$ -category, as this is true for any ringed topos. Next, we check that the full  $\infty$ -subcategory  $\mathcal{D}_{\text{ét}}(Y, \Lambda)$ , with objects those of  $D_{\text{ét}}(Y, \Lambda)$ , is closed under all colimits in  $\mathcal{D}(Y_v, \Lambda)$ . This is clear for cones, so we are reduced to filtered colimits. Those commute with canonical truncations, and filtered colimits of étale sheaves are still étale sheaves, as desired.

By [Lur09, Proposition 5.5.3.12], it is enough to prove the claim if  $Y$  is a disjoint union of strictly totally disconnected perfectoid spaces. In that case,  $\mathcal{D}_{\text{ét}}(Y, \Lambda) = \mathcal{D}(Y_{\text{ét}}, \Lambda)$  (as the functor of stable  $\infty$ -categories  $\mathcal{D}(Y_{\text{ét}}, \Lambda) \rightarrow \mathcal{D}(Y_v, \Lambda)$  is fully faithful (as it is on homotopy categories), and has the same objects as  $\mathcal{D}_{\text{ét}}(Y, \Lambda)$ ), which is a presentable  $\infty$ -category.  $\square$

One consequence of this is the following corollary.

**Corollary 17.2.** *For any small  $v$ -stack  $Y$ , the inclusion  $D_{\text{ét}}(Y, \Lambda) \subset D(Y_v, \Lambda)$  has a right adjoint*

$$R_{Y_{\text{ét}}} : D(Y_v, \Lambda) \rightarrow D_{\text{ét}}(Y, \Lambda) .$$

In general, it is hard to compute  $R_{Y_{\text{ét}}}$ . If  $Y$  is a locally spatial diamond, then on  $D^+(Y_v, \Lambda)$ , it is given by  $R(\nu \circ \lambda)_*$ , where  $\nu \circ \lambda : Y_v \rightarrow Y_{\text{ét}}$  is the map of sites; for strictly totally disconnected spaces, this formula even holds true on all of  $D(Y_v, \Lambda)$ .

*Proof.* The inclusion can be lifted to a colimit-preserving functor  $\mathcal{D}_{\text{ét}}(Y, \Lambda) \rightarrow \mathcal{D}(Y_v, \Lambda)$  of presentable  $\infty$ -categories. As such, it admits a right adjoint by Lurie's  $\infty$ -categorical adjoint functor theorem, [Lur09, Corollary 5.5.2.9].  $\square$

We now deal first with the pullback functor. We claim that if  $f : Y' \rightarrow Y$  is a map of small  $v$ -stacks, there is a pullback functor

$$f^* : D(Y_v, \Lambda) \rightarrow D(Y'_v, \Lambda) ,$$

inducing by restriction

$$f^* : D_{\text{ét}}(Y, \Lambda) \rightarrow D_{\text{ét}}(Y', \Lambda) .$$

This functor is easy to construct for 0-truncated maps, as then  $f$  induces a map of sites  $f_v : Y'_v \rightarrow Y_v$ . In that case, one even gets naturally a functor of  $\infty$ -categories

$$f_v^* : \mathcal{D}(Y_v, \Lambda) \rightarrow \mathcal{D}(Y'_v, \Lambda) ,$$

which induces by restriction

$$f^* : \mathcal{D}_{\text{ét}}(Y, \Lambda) \rightarrow \mathcal{D}_{\text{ét}}(Y', \Lambda) .$$

In general, we need simplicial techniques, so let us recall this.

**Proposition 17.3.** *Let  $Y$  be a small  $v$ -stack, and let  $Y_\bullet \rightarrow Y$  be a simplicial  $v$ -hypercover of  $Y$  by small  $v$ -stacks  $Y_\bullet$  with 0-truncated maps  $Y_i \rightarrow Y$ . Consider the simplicial site  $Y_{\bullet,v}$ , and let  $\mathcal{D}(Y_{\bullet,v}, \Lambda)$  be the  $\infty$ -derived category of sheaves of  $\Lambda$ -modules on  $Y_{\bullet,v}$ , with homotopy category  $D(Y_{\bullet,v}, \Lambda)$ .*

*Pullback along  $Y_\bullet \rightarrow Y$  induces a fully faithful functor*

$$\mathcal{D}(Y_v, \Lambda) \rightarrow \mathcal{D}(Y_{\bullet,v}, \Lambda)$$

*whose essential image is given by the full  $\infty$ -subcategory of cartesian objects. In particular, on homotopy categories,*

$$D(Y_v, \Lambda) \rightarrow D(Y_{\bullet,v}, \Lambda)$$

*is fully faithful with essential image given by the full subcategory of cartesian objects.*

*Proof.* This follows from [Sta, Tag 0DC7] (on homotopy categories, thus on stable  $\infty$ -categories), cf. also [BS15, Proposition 3.3.6].  $\square$

**Remark 17.4.** One may worry that  $D_{\text{ét}}(Y, \Lambda) \subset D(Y_v, \Lambda)$  depends on the choice of  $\kappa$ , as  $Y_v$  does. However, by choosing a simplicial  $v$ -hypercover  $Y_\bullet \rightarrow Y$  by disjoint unions of strictly totally disconnected spaces, and using that  $D_{\text{ét}}(Y_i, \Lambda) = D(Y_{i,\text{ét}}, \Lambda)$  is independent of the choice of  $\kappa$ , one finds that  $D_{\text{ét}}(Y, \Lambda) = D_{\text{ét, cart}}(Y_\bullet, \Lambda)$  is also independent of the choice of  $\kappa$ . Similarly, one checks that the whole 6-functor formalism does not depend on the choice of  $\kappa$ .

Now, if  $f : Y' \rightarrow Y$  is any map of small  $v$ -stacks, choose a perfectoid space  $Y'_0$  with a surjective map of  $v$ -stacks  $Y'_0 \rightarrow Y'$ , and let  $Y'_\bullet$  be the corresponding Čech nerve, all of whose terms are small  $v$ -sheaves. Then in particular all the maps  $Y'_i \rightarrow Y'$  and  $Y'_i \rightarrow Y$  are 0-truncated, and we get a well-defined pullback functor

$$\mathcal{D}(Y_v, \Lambda) \rightarrow \mathcal{D}(Y'_{\bullet,v}, \Lambda) ,$$

taking values in cartesian objects (as composites of pullbacks are pullbacks). Thus, we get a functor

$$f_v^* : \mathcal{D}(Y_v, \Lambda) \rightarrow \mathcal{D}_{\text{cart}}(Y'_{\bullet,v}, \Lambda) \simeq \mathcal{D}(Y'_v, \Lambda) .$$

This carries  $\mathcal{D}_{\text{ét}}(Y_v, \Lambda)$  into  $\mathcal{D}_{\text{ét}}(Y'_v, \Lambda)$  (as it is, at least up to homotopy, compatible with further pullbacks). Thus, we get a functor

$$f^* : \mathcal{D}_{\text{ét}}(Y, \Lambda) \rightarrow \mathcal{D}_{\text{ét}}(Y', \Lambda) ,$$

and in particular

$$f^* : D_{\text{ét}}(Y, \Lambda) \rightarrow D_{\text{ét}}(Y', \Lambda) .$$

It is easy to see that this is canonically independent of the choices made, and compatible with composition, i.e. for another map  $g : Y'' \rightarrow Y'$  of small  $v$ -stacks with composite  $f \circ g : Y'' \rightarrow Y$ , one has a natural equivalence  $(f \circ g)^* \simeq f^* \circ g^*$ , satisfying the usual coherences.

**Lemma 17.5.** *For any map of small  $v$ -stacks  $f : Y' \rightarrow Y$ , the functor*

$$f^* : D_{\text{ét}}(Y, \Lambda) \rightarrow D_{\text{ét}}(Y', \Lambda)$$

*has a right adjoint*

$$Rf_* : D_{\text{ét}}(Y', \Lambda) \rightarrow D_{\text{ét}}(Y, \Lambda) .$$

*Proof.* The functor  $f_v^* : \mathcal{D}(Y_v, \Lambda) \rightarrow \mathcal{D}(Y'_v, \Lambda)$  commutes with all colimits by construction, and thus so does  $f^* : \mathcal{D}_{\text{ét}}(Y_v, \Lambda) \rightarrow \mathcal{D}_{\text{ét}}(Y'_v, \Lambda)$ . Thus, the result follows from the  $\infty$ -categorical adjoint functor theorem, [Lur09, Corollary 5.5.2.9], and Lemma 17.1.  $\square$

Again, it is in general hard to compute  $Rf_*$ . If  $Y'$  and  $Y$  are locally spatial diamonds, then it agrees with (the left-completed)  $Rf_{\text{ét}*}$ . In general, it is clear that  $Rf_* = R_{Y\text{ét}} Rf_{v*}$ ; however, this involves the complicated functor  $R_{Y\text{ét}}$ . However, in the following situation, it is not necessary to apply  $R_{Y\text{ét}}$ ; this makes use of the results of the previous section.

**Proposition 17.6.** *Assume that  $n\Lambda = 0$  for some  $n$  prime to  $p$ . Let  $f : Y' \rightarrow Y$  be a qcqs map of small  $v$ -stacks. Then for any  $A \in D_{\text{ét}}^+(Y', \Lambda)$ , one has  $Rf_{v*}A \in D_{\text{ét}}^+(Y, \Lambda)$ , and thus*

$$Rf_*A = Rf_{v*}A .$$

Moreover, in this case, the formation of  $Rf_*$  commutes with any base change, i.e. for all maps  $g : \tilde{Y} \rightarrow Y$  of small  $v$ -stacks with pullback  $\tilde{f} : \tilde{Y}' = Y' \times_Y \tilde{Y} \rightarrow \tilde{Y}$ ,  $g' : \tilde{Y}' \rightarrow Y'$ , the natural transformation

$$g^*Rf_*A \rightarrow R\tilde{f}_*g'^*A$$

is an equivalence for all  $A \in D_{\text{ét}}^+(Y', \Lambda)$ .

If  $Rf_*$  has finite cohomological dimension, i.e. there is some integer  $N$  such that for all  $A \in D_{\text{ét}}(Y', \Lambda)$  concentrated in degree 0, one has  $R^i f_*A = 0$  for  $i > N$ , then for all  $A \in D_{\text{ét}}(U', \Lambda)$ , one has

$$Rf_*A = Rf_{v*}A \in D_{\text{ét}}(Y, \Lambda) ,$$

and moreover the base change morphism

$$g^*Rf_*A \rightarrow R\tilde{f}_*g'^*A$$

is an isomorphism.

We note that to get base change, we only need to assume that  $Rf_*$  has finite cohomological dimension (and not the same about  $R\tilde{f}_{ast}$ ).

*Proof.* Assume first that  $A \in D_{\text{ét}}^+(Y', \Lambda)$ . The statement  $Rf_{v*}A \in D_{\text{ét}}^+(Y, \Lambda)$  is given by Corollary 16.8 (ii). The base change result follows formally as  $v$ -pushforward commutes with  $v$ -slices.

Now, if  $Rf_*$  has finite cohomological dimension, then by left-completeness, it follows that in general

$$Rf_{v*}A = R\varprojlim_n Rf_*\tau^{\geq -n}A ,$$

where the limit becomes eventually constant in each degree. In a replete topos, the cohomological dimension of  $R\varprojlim_n$  is bounded by 1, cf. [BS15, Proposition 3.1.11], so we see that each cohomology sheaf of  $Rf_{v*}A$  agrees with the cohomology sheaf of  $Rf_*\tau^{\geq -n}A$  for  $n$  sufficiently large, which shows that  $Rf_{v*}A \in D_{\text{ét}}(Y, \Lambda)$  by Proposition 14.16, and thus  $Rf_*A = Rf_{v*}A$ .

Moreover, concerning base change, one has, by commuting limits,

$$\begin{aligned} R\tilde{f}_*g'^*A &= R\varprojlim_n R\tilde{f}_*g'^*\tau^{\geq -n}A \\ &= R\varprojlim_n g^*Rf_*\tau^{\geq -n}A . \end{aligned}$$

This admits a canonical map from  $g^*Rf_*A$ . The cone of the map

$$g^*Rf_*A \rightarrow g^*Rf_*\tau^{\geq -n}A$$

lies in degrees  $\leq -n + N$ . This implies that the  $R\varprojlim_n$  is equal to zero, as we work in a replete topos, and so the derived category is left-complete.  $\square$



Next, we define the tensor product functor. Note that there is a natural functor

$$- \otimes_{\Lambda}^{\mathbb{L}} - : D(Y_v, \Lambda) \times D(Y_v, \Lambda) \rightarrow D(Y_v, \Lambda) ,$$

like for any ringed topoi. In fact, this functor is defined as a functor

$$- \otimes_{\Lambda}^{\mathbb{L}} - : \mathcal{D}(Y_v, \Lambda) \times \mathcal{D}(Y_v, \Lambda) \rightarrow \mathcal{D}(Y_v, \Lambda)$$

which preserves all colimits separately in each variable.<sup>3</sup> If  $f : Y' \rightarrow Y$  is any map of small v-stacks, then there is a natural equivalence

$$f^*(A \otimes_{\Lambda}^{\mathbb{L}} B) \simeq f^* A \otimes_{\Lambda}^{\mathbb{L}} f^* B$$

for all  $A, B \in D(Y_v, \Lambda)$ : This is clear for 0-truncated maps by the general formalism of ringed topoi, and then follows in general by following the construction of  $f^*$ .

**Lemma 17.7.** *For any small v-stack  $Y$ , the functor*

$$- \otimes_{\Lambda}^{\mathbb{L}} - : D(Y_v, \Lambda) \times D(Y_v, \Lambda) \rightarrow D(Y_v, \Lambda)$$

*induces by restriction a functor*

$$- \otimes_{\Lambda}^{\mathbb{L}} - : D_{\text{ét}}(Y, \Lambda) \times D_{\text{ét}}(Y, \Lambda) \rightarrow D_{\text{ét}}(Y, \Lambda) .$$

One can use this lemma (together with the previous discussion) to deduce that  $\mathcal{D}_{\text{ét}}(Y, \Lambda)$  is a presentably symmetric monoidal  $\infty$ -category (in a unique way compatible with the standard symmetric monoidal structure on  $\mathcal{D}(Y_v, \Lambda)$ ).

*Proof.* This can be checked v-locally on  $Y$ , so we can assume that  $Y$  is a disjoint union of strictly totally disconnected perfectoid spaces. In this case,  $D_{\text{ét}}(Y, \Lambda) = D(Y_{\text{ét}}, \Lambda)$ , and the result follows from the existence of the natural tensor product on  $D(Y_{\text{ét}}, \Lambda)$  (compatible with pullback along  $Y_v \rightarrow Y_{\text{ét}}$ ).  $\square$

**Lemma 17.8.** *For any small v-stack  $Y$  and  $A \in D_{\text{ét}}(Y, \Lambda)$ , the functor*

$$D_{\text{ét}}(Y, \Lambda) \rightarrow D_{\text{ét}}(Y, \Lambda) : B \mapsto B \otimes_{\Lambda}^{\mathbb{L}} A$$

*admits a right adjoint*

$$D_{\text{ét}}(Y, \Lambda) \rightarrow D_{\text{ét}}(Y, \Lambda) : C \mapsto R\mathcal{H}om_{\Lambda}(A, C) ,$$

*i.e.*

$$\text{Hom}_{D_{\text{ét}}(Y, \Lambda)}(B \otimes_{\Lambda}^{\mathbb{L}} A, C) = \text{Hom}_{D_{\text{ét}}(Y, \Lambda)}(B, R\mathcal{H}om_{\Lambda}(A, C)) .$$

*For varying  $A$ , these assemble into a functor*

$$R\mathcal{H}om_{\Lambda}(-, -) : D_{\text{ét}}(Y, \Lambda)^{\text{op}} \times D_{\text{ét}}(Y, \Lambda) \rightarrow D_{\text{ét}}(Y, \Lambda) .$$

Again,  $R\mathcal{H}om_{\Lambda}(-, -)$  is in general hard to compute, and not compatible with the inner Hom in  $D(Y_v, \Lambda)$ ; rather, one has to apply  $R_{Y_{\text{ét}}}$  to the inner Hom in  $D(Y_v, \Lambda)$ .

<sup>3</sup>Even better,  $\mathcal{D}(Y_v, \Lambda)$  is a presentably symmetric monoidal  $\infty$ -category.

*Proof.* We have a functor from  $D_{\text{ét}}(Y, \Lambda)^{\text{op}} \times D_{\text{ét}}(Y, \Lambda)$  to the presheaf category on  $D_{\text{ét}}(Y, \Lambda)$ , given by sending a pair  $(A, C)$  to the functor from  $D_{\text{ét}}(Y, \Lambda)^{\text{op}}$  to sets,

$$B \mapsto \text{Hom}_{D_{\text{ét}}(Y, \Lambda)}(B \otimes_{\Lambda}^{\mathbb{L}} A, C) .$$

The claim is that this functor factors over the full subcategory  $D_{\text{ét}}(Y, \Lambda)$ , i.e. is representable for any pair  $(A, C)$ . But for any fixed  $(A, C)$ , the functor

$$B \mapsto \text{Map}_{D_{\text{ét}}(Y, \Lambda)}(B \otimes_{\Lambda}^{\mathbb{L}} A, C)$$

from  $D_{\text{ét}}(Y, \Lambda)^{\text{op}}$  to the  $\infty$ -category of spaces takes all limits (in  $D_{\text{ét}}(Y, \Lambda)^{\text{op}}$ , which are colimits in  $D_{\text{ét}}(Y, \Lambda)$ ) to limits (as  $-\otimes_{\Lambda}^{\mathbb{L}} A$  preserves colimits). Thus, by [Lur09, Proposition 5.5.2.2], it is representable, by what is denoted  $R\mathcal{H}om_{\Lambda}(A, C)$ .  $\square$

**Corollary 17.9.** *Let  $f : Y' \rightarrow Y$  be a map of small v-stacks. There is a natural equivalence*

$$Rf_* R\mathcal{H}om_{\Lambda}(f^* A, B) \cong R\mathcal{H}om_{\Lambda}(A, Rf_* B)$$

of functors

$$D_{\text{ét}}(Y, \Lambda)^{\text{op}} \times D_{\text{ét}}(Y', \Lambda) \rightarrow D_{\text{ét}}(Y, \Lambda) .$$

*Proof.* For any  $C \in D_{\text{ét}}(Y, \Lambda)$ , we have a series of natural equivalences

$$\begin{aligned} \text{Hom}_{D_{\text{ét}}(Y, \Lambda)}(C, Rf_* R\mathcal{H}om_{\Lambda}(f^* A, B)) &\cong \text{Hom}_{D_{\text{ét}}(Y', \Lambda)}(f^* C, R\mathcal{H}om_{\Lambda}(f^* A, B)) \\ &\cong \text{Hom}_{D_{\text{ét}}(Y', \Lambda)}(f^* C \otimes_{\Lambda}^{\mathbb{L}} f^* A, B) \\ &\cong \text{Hom}_{D_{\text{ét}}(Y', \Lambda)}(f^*(C \otimes_{\Lambda}^{\mathbb{L}} A), B) \\ &\cong \text{Hom}_{D_{\text{ét}}(Y, \Lambda)}(C \otimes_{\Lambda}^{\mathbb{L}} A, Rf_* B) \\ &\cong \text{Hom}_{D_{\text{ét}}(Y, \Lambda)}(C, R\mathcal{H}om_{\Lambda}(A, Rf_* B)) , \end{aligned}$$

giving the desired result.  $\square$

## 18. PROPER AND PARTIALLY PROPER MORPHISMS

As a preparation for the proper base change theorem in the next section, we define proper and partially proper morphisms of v-sheaves.

**Definition 18.1.** *Let  $f : Y' \rightarrow Y$  be a map of v-stacks. The map  $f$  is proper if it is quasicompact, separated, and universally closed, i.e. for all small v-sheaves  $X$  with a map  $X \rightarrow Y$ , the map  $|Y' \times_Y X| \rightarrow |X|$  is closed.*

Note that as  $f$  is quasicompact, it follows in particular that for any small v-sheaf  $X$  with a map  $X \rightarrow Y$ , the pullback  $Y' \times_Y X$  is a small v-sheaf. By the definition of the topology on  $|X|$  for a small v-sheaf, it suffices to check this condition for perfectoid spaces  $X$ , or even just for strictly totally disconnected perfectoid space  $X$ .

**Remark 18.2.** Any closed immersion of v-sheaves is a proper map. As a somewhat amusing exercise, we leave it to the reader to check that if  $T' \rightarrow T$  is a map of locally compact Hausdorff spaces, then the map of v-sheaves  $\underline{T}' \rightarrow \underline{T}$  is proper if and only if  $T' \rightarrow T$  is proper in the usual sense.

Again, there is a valuative criterion.

**Proposition 18.3.** *Let  $f : Y' \rightarrow Y$  be a map of  $v$ -stacks. Then  $f$  is proper if and only if it is 0-truncated, qcqs, and for every perfectoid field  $K$  with an open and bounded valuation subring  $K^+ \subset K$ , and any diagram*

$$\begin{array}{ccc} \mathrm{Spa}(K, \mathcal{O}_K) & \longrightarrow & Y' \\ \downarrow & \nearrow \text{dotted} & \downarrow f \\ \mathrm{Spa}(K, K^+) & \longrightarrow & Y, \end{array}$$

there exists a unique dotted arrow making the diagram commute.

Moreover, if  $f$  is proper, then for every perfectoid Tate ring  $R$  with an open and integrally closed subring  $R^+ \subset R$ , and every diagram

$$\begin{array}{ccc} \mathrm{Spa}(R, R^\circ) & \longrightarrow & Y' \\ \downarrow & \nearrow \text{dotted} & \downarrow f \\ \mathrm{Spa}(R, R^+) & \longrightarrow & Y, \end{array}$$

there exists a unique dotted arrow making the diagram commute.

*Proof.* Assume first that the valuative criterion holds. By Proposition 10.9, we know that  $f$  is separated. Replacing  $f$  by a pullback, it is enough to see that  $|f| : |Y'| \rightarrow |Y|$  is closed if  $Y$  is a small  $v$ -sheaf. Choosing a  $v$ -cover  $X \rightarrow Y$ , where  $X$  is a perfectoid space, we can further reduce to the case that  $Y = X$  is a perfectoid space. As we can moreover work locally on  $X$ , we can assume that  $X$  is an affinoid perfectoid space. Then  $Y'$  is a qcqs  $v$ -sheaf; choose an affinoid perfectoid space  $X'$  with a  $v$ -cover  $X' \rightarrow Y'$ . Let  $Z \subset |Y'|$  be a closed subset. Then the image  $W \subset |X|$  of  $Z$  in  $|Y'|$  is a pro-constructible subset, as it is also the image of the preimage of  $Z$  in  $|X'|$ , under the spectral map of spectral spaces  $|X'| \rightarrow |X|$ . To see that  $W$  is closed, it suffices to see that  $W$  is specializing. For this, let  $w, w' \in X$  be two points such that  $w$  specializes to  $w'$ , and  $w \in W$ . This corresponds to maps  $\mathrm{Spa}(K, K^+) \rightarrow \mathrm{Spa}(K, (K^+)') \rightarrow X$  for a perfectoid field  $K$  with open and bounded valuation subrings  $(K^+) \subset K^+ \subset K$ , such that  $w$  is the image of the closed point of  $\mathrm{Spa}(K, K^+)$ , and  $w'$  is the image of the closed point of  $\mathrm{Spa}(K, (K^+)')$ . As  $w$  lies in the image of  $Z$ , we can assume that, after enlarging  $K$ , there is a commutative diagram

$$\begin{array}{ccc} \mathrm{Spa}(K, K^+) & \longrightarrow & Y' \\ \downarrow & \nearrow \text{dotted} & \downarrow f \\ \mathrm{Spa}(K, (K^+)') & \longrightarrow & X, \end{array}$$

such that the image of the closed point of  $\mathrm{Spa}(K, K^+)$  in  $|Y'|$  lies in  $Z$ . Applying the valuative criterion for the two pairs  $(K, K^+)$  and  $(K, (K^+)')$ , we see that there exists a unique dotted arrow in the diagram. As  $Z$  is closed and  $\mathrm{Spa}(K, K^+)$  is dense in  $\mathrm{Spa}(K, (K^+)')$ , the closed point of  $\mathrm{Spa}(K, (K^+)')$  maps to  $Z$ . Thus,  $w'$  lies in the image  $W$  of  $Z$  in  $|X|$ , as desired.

Conversely, assume that  $f$  is proper. We will prove that for all pairs  $(R, R^+)$  of a perfectoid Tate ring with an open and integrally closed subring  $R^+ \subset R$ , and every diagram

$$\begin{array}{ccc} \mathrm{Spa}(R, R^\circ) & \longrightarrow & Y' \\ \downarrow & \nearrow & \downarrow f \\ \mathrm{Spa}(R, R^+) & \longrightarrow & Y \end{array},$$

there exists a unique dotted arrow making the diagram commute. By Proposition 10.10, we know that the map is injective. Thus, assume given a map  $\mathrm{Spa}(R, R^+) \rightarrow Y$  and a lift of  $\mathrm{Spa}(R, R^\circ) \rightarrow Y$  to a map  $\mathrm{Spa}(R, R^\circ) \rightarrow Y'$ . We may replace  $Y$  by  $\mathrm{Spa}(R, R^+)$  and  $Y'$  by the corresponding pullback. We get a quasicompact injection

$$\mathrm{Spa}(R, R^\circ) \rightarrow Y'.$$

By Proposition 12.15, it is characterized by the subspace  $|\mathrm{Spa}(R, R^\circ)| \subset |Y'|$ . Let  $X' \rightarrow Y'$  be a surjection from a totally disconnected perfectoid space  $X'$ . Then  $\mathrm{Spa}(R, R^\circ) \times_{Y'} X' \subset X'$  is a quasicompact injection, which by Corollary 10.6 corresponds to a pro-constructible and generalizing subset  $W \subset |X'|$ , given by  $W = |\mathrm{Spa}(R, R^\circ)| \times_{|Y'|} |X'|$ . Consider the closure  $Z$  of  $|\mathrm{Spa}(R, R^\circ)|$  in  $|Y'|$ . Its preimage  $Z' \subset |X'|$  is a closed subset containing  $W$ . In fact, it is precisely the closure of  $W$ , as the closure of  $W$  can be checked to be invariant under the equivalence relation  $|X'| \times_{|Y'|} |X'|$ , and  $|Y'|$  has the quotient topology from  $|X'|$ . Thus,  $Z' \subset |X'|$  is a closed and generalizing subset, and therefore corresponds to a totally disconnected perfectoid space; passing back to  $Z$ , we see that  $Z$  corresponds to a qcqs  $v$ -sheaf, still denoted by  $Z$ . The map  $Z \rightarrow \mathrm{Spa}(R, R^+)$  is an injection: Indeed, to check this, it suffices by Proposition 12.15 to check that  $Z(K, K^+) \rightarrow \mathrm{Spa}(R, R^+)(K, K^+)$  is injective for all perfectoid fields  $K$  with an open and bounded valuation subring  $K^+ \subset K$ . This is true if  $K^+ = \mathcal{O}_K$  as  $Z(K, \mathcal{O}_K) = \mathrm{Spa}(R, R^\circ)(K, \mathcal{O}_K)$ ; in general, it follows as  $Z \rightarrow \mathrm{Spa}(R, R^+)$  is separated (as a sub- $v$ -sheaf of the separated map  $Y' \rightarrow \mathrm{Spa}(R, R^+)$ ).

On the other hand, as  $f$  is by assumption universally closed, the image of  $|Z|$  in  $|\mathrm{Spa}(R, R^+)|$  is closed, and thus all of  $|\mathrm{Spa}(R, R^+)|$ , as it contains the dense subspace  $|\mathrm{Spa}(R, R^\circ)|$ . Thus,  $|Z| \rightarrow |\mathrm{Spa}(R, R^+)|$  is surjective, thus a quotient map by Lemma 2.5 (applied to a cover of  $Z$  by an affinoid perfectoid space), and therefore (using Proposition 12.15 again)  $Z$  maps isomorphically to  $\mathrm{Spa}(R, R^+)$ . Composing the inverse with the natural map  $Z \rightarrow Y'$  gives the desired map  $\mathrm{Spa}(R, R^+) \rightarrow Y'$ .  $\square$

The following generalization of proper maps is useful.

**Definition 18.4.** *Let  $f : Y' \rightarrow Y$  be a map of  $v$ -stacks. Then  $f$  is partially proper if  $f$  is separated, and for every perfectoid Tate ring  $R$  with an open and integrally closed subring  $R^+ \subset R$ , and every diagram*

$$\begin{array}{ccc} \mathrm{Spa}(R, R^\circ) & \longrightarrow & Y' \\ \downarrow & \nearrow & \downarrow f \\ \mathrm{Spa}(R, R^+) & \longrightarrow & Y \end{array},$$

*there exists a (necessarily unique) dotted arrow making the diagram commute.*

*Moreover, the  $v$ -stack  $Y$  is partially proper if  $Y \rightarrow *$  is partially proper.*

**Remark 18.5.** We refrain from making the definition that  $Y$  is proper if  $Y \rightarrow *$  is proper. The problem here is that proper should be equivalent to being partially proper and quasicompact; however, the notions of  $Y$  being quasicompact and of  $Y \rightarrow *$  being quasicompact are different. In the partially proper case, this issue does not come up.

A useful feature of the world of analytic adic spaces is the presence of canonical compactifications, cf. [Hub96, Theorem 5.1.5]. In our context, this is the following construction.

**Proposition 18.6.** *Let  $f : Y' \rightarrow Y$  be a separated map of  $v$ -stacks. The functor sending any totally disconnected perfectoid space  $X = \mathrm{Spa}(R, R^+)$  to*

$$Y'(R, R^\circ) \times_{Y(R, R^\circ)} Y(R, R^+)$$

*extends to a  $v$ -stack  $\overline{Y'}^{/Y}$  with a map  $\overline{f}^{/Y} : \overline{Y'}^{/Y} \rightarrow Y$ , and a natural map  $Y' \rightarrow \overline{Y'}^{/Y}$  over  $Y$ . The map  $\overline{f}^{/Y}$  is partially proper, and for every partially proper map  $g : Z \rightarrow Y$  of  $v$ -stacks, composition with  $Y' \rightarrow \overline{Y'}^{/Y}$  induces a bijection*

$$\mathrm{Hom}_Y(\overline{Y'}^{/Y}, Z) \rightarrow \mathrm{Hom}_Y(Y', Z).$$

We will refer to  $\overline{f}^{/Y} : \overline{Y'}^{/Y} \rightarrow Y$  as the canonical compactification of  $f : Y' \rightarrow Y$ . The formation of  $\overline{f}^{/Y}$  is clearly functorial in  $f$ .

*Proof.* To see that  $\overline{Y'}^{/Y}$  is a  $v$ -stack, observe first that if  $\mathrm{Spa}(S, S^+) \rightarrow \mathrm{Spa}(R, R^+)$  is a  $v$ -cover of totally disconnected perfectoid spaces, then also  $\mathrm{Spa}(S, S^\circ) \rightarrow \mathrm{Spa}(R, R^\circ)$  is a  $v$ -cover. This is true as  $\mathrm{Spa}(R, R^\circ) \subset \mathrm{Spa}(R, R^+)$  is the minimal pro-constructible generalizing subset containing all rank-1-points by Lemma 7.6, and the image of  $\mathrm{Spa}(S, S^\circ) \rightarrow \mathrm{Spa}(R, R^\circ)$  is a pro-constructible and generalizing subset containing all rank-1-points. Also, to check that  $\overline{Y'}^{/Y}$  is a  $v$ -stack, we may work locally on  $Y$ , and in particular assume that  $Y$  is representable, in which case  $Y'$  is a  $v$ -sheaf, and  $\overline{Y'}^{/Y}$  is a presheaf. Choose a surjection  $\mathrm{Spa}(T, T^+) \rightarrow \mathrm{Spa}(S, S^+) \times_{\mathrm{Spa}(R, R^+)} \mathrm{Spa}(S, S^+)$ , where  $\mathrm{Spa}(T, T^+)$  is totally disconnected. We need to see that

$$\overline{Y'}^{/Y}(R, R^+) = \mathrm{eq}(\overline{Y'}^{/Y}(S, S^+) \rightrightarrows \overline{Y'}^{/Y}(T, T^+)).$$

As  $\mathrm{Spa}(S, S^\circ) \rightarrow \mathrm{Spa}(R, R^\circ)$  is a  $v$ -cover, we see that the map

$$\overline{Y'}^{/Y}(R, R^+) = Y'(R, R^\circ) \times_{Y(R, R^\circ)} Y(R, R^+) \rightarrow \overline{Y'}^{/Y}(S, S^+) = Y'(S, S^\circ) \times_{Y(S, S^\circ)} Y(S, S^+)$$

is injective. Now let  $a \in \overline{Y'}^{/Y}(S, S^+) = Y'(S, S^\circ) \times_{Y(S, S^\circ)} Y(S, S^+)$  be a section whose two pullbacks to  $\overline{Y'}^{/Y}(T, T^+)$  agree. In particular, we get a section  $b \in Y(R, R^+)$  as  $Y$  is a  $v$ -sheaf. On  $Y'$ , we want to see that the section in  $Y'(S, S^\circ)$  descends to  $Y'(R, R^\circ)$ . For this, we need to see that the two induced sections in  $Y'(T, (T^+)')$  agree, where  $\mathrm{Spa}(T, (T^+)') = \mathrm{Spa}(S, S^\circ) \times_{\mathrm{Spa}(R, R^\circ)} \mathrm{Spa}(S, S^\circ)$ . But we know that they agree in  $Y'(T, T^\circ)$  and  $Y(T, (T^+)')$ , and  $Y' \rightarrow Y$  is separated, so the result follows from Proposition 10.10.

It remains to see that  $\overline{f}^{/Y}$  is partially proper, for then the definition of partially proper maps ensures that any map  $Y' \rightarrow Z$  to a partially proper  $Z \rightarrow Y$  factors uniquely over  $\overline{Y'}^{/Y}$ . This question is  $v$ -local on  $Y$ , so we may assume that  $Y$  is an affinoid perfectoid space (and in particular separated). Now the result follows from Proposition 18.7 (i) below.  $\square$

In fact, for any separated v-sheaf  $Y$ , one can define a new v-sheaf  $\bar{Y}$  by setting

$$\bar{Y}(R, R^+) = Y(R, R^\circ)$$

for any totally disconnected space  $\mathrm{Spa}(R, R^+)$ , which comes with an injective map  $Y \rightarrow \bar{Y}$ . The verification that  $\bar{Y}$  is a v-sheaf is identical to the verification for  $\bar{Y}'/Y$  above (and uses that  $Y$  is separated). With this notation, we have

$$\bar{Y}'/Y = \bar{Y}' \times_{\bar{Y}} Y .$$

We will need the following properties.

**Proposition 18.7.** *Let  $Y$  be a separated v-sheaf.*

- (i) *The v-sheaf  $\bar{Y}$  is partially proper.*
- (ii) *If  $Y$  is a small v-sheaf, then  $\bar{Y}$  is a small v-sheaf.*
- (iii) *If  $Y$  is a diamond, then  $\bar{Y}$  is a diamond.*
- (iv) *If  $Y = \mathrm{Spa}(R, R^+)$  is an affinoid perfectoid space, then  $\bar{Y} = \mathrm{Spa}(R, (R^+)')$ , where  $(R^+) \subset R^+$  is the smallest open and integrally closed subring (which is the integral closure of  $\mathbb{F}_p + R^{\circ\circ}$ ).*
- (v) *The functor  $Y \mapsto \bar{Y}$  commutes with all limits.*
- (vi) *If  $f : Y' \rightarrow Y$  is a surjective map of separated v-sheaves, then  $\bar{f} : \bar{Y}' \rightarrow \bar{Y}$  is a surjective map of v-sheaves.*
- (vii) *If  $f : Y' \rightarrow Y$  is a quasicompact map of separated v-sheaves, then  $\bar{f} : \bar{Y}' \rightarrow \bar{Y}$  is a proper map of v-sheaves.*
- (viii) *If  $f : Y' \rightarrow Y$  is a quasi-pro-étale map of separated v-sheaves, then  $\bar{f} : \bar{Y}' \rightarrow \bar{Y}$  is quasi-pro-étale.*

*Proof.* Part (iv) follows from the equations

$$\mathrm{Hom}((R, (R^+)'), (S, S^+)) = \mathrm{Hom}(R, S) = \mathrm{Hom}((R, R^+), (S, S^\circ)) ,$$

which hold for any pair  $(S, S^+)$ . Part (v) is clear from the definition. For part (vi), consider a totally disconnected space  $X = \mathrm{Spa}(R, R^+)$  with a map  $X \rightarrow \bar{Y}$ , which corresponds to a map  $\mathrm{Spa}(R, R^\circ) \rightarrow Y$ . Then, after replacing  $X$  by a v-cover, we can lift this to a map  $\mathrm{Spa}(R, R^\circ) \rightarrow Y'$ , which corresponds to a map  $X \rightarrow \bar{Y}'$  lifting  $X \rightarrow \bar{Y}$ .

For part (i), we check first that  $\bar{Y}$  is quasiseparated. If  $X_1, X_2 \rightarrow \bar{Y}$  are two affinoid perfectoid spaces mapping to  $\bar{Y}$ , we have to show that  $X_1 \times_{\bar{Y}} X_2$  is quasicompact. But using (v), we have

$$X_1 \times_{\bar{Y}} X_2 = X_1 \times_{\bar{X}_1} \overline{X_1 \times_Y X_2} \times_{\bar{X}_2} X_2 ,$$

where the maps  $X_i \rightarrow \bar{X}_i$  are quasicompact by (iv), and  $\overline{X_1 \times_Y X_2}$  is quasicompact, using (vi) and the assumption that  $X_1 \times_Y X_2$  is quasicompact (as  $Y$  is quasiseparated). Now, since  $\bar{Y}$  is quasiseparated, it follows that  $\bar{Y}$  is partially proper, as it satisfies the desired valuative criterion by construction (for totally disconnected  $\mathrm{Spa}(R, R^+)$ , which gives the general case by v-descent); this finishes the proof of (i).

Part (ii) follows directly from (iv) and (vi). For part (vii), note that  $\bar{f}$  is always partially proper by (i); to show that  $\bar{f}$  is proper, it remains to see that  $\bar{f}$  is quasicompact. But this follows from (iv), (v) and (vi) (similarly to the argument that  $\bar{Y}$  is quasicompact).

For part (viii), we need to see that for any strictly totally disconnected space  $X \rightarrow \overline{Y}$ ,  $\overline{Y'} \times_{\overline{Y}} X \rightarrow X$  is quasi-pro-étale. Let  $X = \mathrm{Spa}(R, R^+)$  and  $X^\circ = \mathrm{Spa}(R, R^\circ)$ . Then the map  $X \rightarrow \overline{Y}$  corresponds to a map  $X^\circ \rightarrow Y$ . We get separated quasi-pro-étale maps  $Y' \times_Y X^\circ \rightarrow X^\circ \rightarrow X$ ; let  $g : Z := Y' \times_Y X^\circ \rightarrow X$  denote their composite. Then  $\overline{Y'} \times_{\overline{Y}} X \rightarrow X$  can be identified with  $\overline{Z}^{/X} \rightarrow X$ . But recall that by Corollary 7.22,  $Z \rightarrow X$  has a canonical factorization  $Z \rightarrow X \times_{\pi_0 X} \pi_0 Z \rightarrow X$ . It is then easy to see that  $\overline{Z}^{/X} = X \times_{\pi_0 X} \pi_0 Z$ , which is quasi-pro-étale over  $X$ .

Finally, part (iii) follows from (iv), (vi) and (viii).  $\square$

Coming back to the setting of Proposition 18.6, there is the following relative version.

**Corollary 18.8.** *Let  $Y' \rightarrow Y$  be a separated map of v-stacks.*

- (i) *The map of v-stacks  $\overline{Y'}^{/Y} \rightarrow Y$  is partially proper.*
- (ii) *If  $Y$  and  $Y'$  are small v-stacks, then  $\overline{Y'}^{/Y}$  is a small v-stack.*
- (iii) *If  $Y$  and  $Y'$  are diamonds, then  $\overline{Y'}^{/Y}$  is a diamond.*
- (iv) *The functor  $Y' \mapsto \overline{Y'}^{/Y}$  from v-sheaves over  $Y$  to v-sheaves over  $Y$  commutes with all limits.*
- (v) *If  $f : Y'_2 \rightarrow Y'_1$  is a surjective map of separated v-stacks over  $Y$ , then  $\overline{f} : \overline{Y'_2}^{/Y} \rightarrow \overline{Y'_1}^{/Y}$  is a surjective map of v-stacks.*
- (vi) *If  $f : Y'_2 \rightarrow Y'_1$  is a quasicompact map of separated v-stacks over  $Y$ , then  $\overline{f} : \overline{Y'_2}^{/Y} \rightarrow \overline{Y'_1}^{/Y}$  is a proper map of v-stacks.*
- (vii) *If  $f : Y'_2 \rightarrow Y'_1$  is a quasi-pro-étale map of separated v-stacks over  $Y$ , then  $\overline{f} : \overline{Y'_2}^{/Y} \rightarrow \overline{Y'_1}^{/Y}$  is quasi-pro-étale.*

*Proof.* If  $Y$  is separated (and thus a sheaf), then also  $Y'$  is separated (and a sheaf), and the results follow from Proposition 18.7 and the formula  $\overline{Y'}^{/Y} = \overline{Y'} \times_{\overline{Y}} Y$ . In general, the formation of  $\overline{Y'}^{/Y}$  commutes with base change in  $Y$ , so all properties reduce to the case where  $Y$  is affinoid perfectoid (and in particular separated).  $\square$

In the case of sheaves, another characterization of partially proper maps is given by the following proposition.

**Proposition 18.9.** *Let  $f : Y' \rightarrow Y$  be a map of v-sheaves. Then  $f$  is partially proper if and only if  $f$  can be written as a (possibly large) filtered colimit of proper maps  $f_i : Y'_i \rightarrow Y$  along closed immersions  $Y'_i \rightarrow Y'_j$ . If  $Y'$  is small, then the filtered colimit is also small.*

*Proof.* It follows from Proposition 18.3 that if  $f$  can be written as such a filtered colimit of proper maps, then  $f$  is partially proper. Conversely, assume that  $f$  is partially proper, and consider the category of all closed immersions  $Y'_i \subset Y'$  for which the composite  $Y'_i \rightarrow Y' \rightarrow Y$  is proper. If  $Y'$  is small, this category is small. Any map in this category is a closed immersion. To finish the proof, we need to see that it is filtered, and that  $Y'$  is the colimit of all those  $Y'_i$ .

First, we claim that if  $Z \rightarrow Y$  is any proper map of v-sheaves and  $Z \rightarrow Y'$  is a map over  $Y$ , then the (sheaf-theoretic) image  $Z'$  of  $Z$  in  $Y'$  is proper over  $Y$ , and closed in  $Y'$ . (In this step, we only use that  $f$  is separated.) Indeed,  $Z'$  is separated as it is a subsheaf of  $Y'$ , and quasicompact as a quotient of  $Z$ , and the valuative criterion for  $Z'$  follows from that for  $Z$ . That the image is closed

follows from the definition of properness, as it can be written as the image of the graph of  $Z \rightarrow Y'$ , which is a closed subset of  $Z \times_Y Y'$ , under the projection  $Z \times_Y Y' \rightarrow Y'$ .

In particular, if  $Y'_i \subset Y'$ ,  $i = 1, 2$ , are two closed subsets which are proper over  $Y$ , then the image of  $Y'_1 \sqcup Y'_2$  is another such subset, showing that the category is filtered. On the other hand, if  $Z = \mathrm{Spa}(R, R^+) \rightarrow Y'$  is any map from an affinoid perfectoid space, then the map is separated (as  $Z$  is separated), and so we have the canonical compactification  $\overline{Z}^{/Y}$ , which is proper over  $Y$  by Corollary 18.8 (vi). Now the image of  $\overline{Z}^{/Y} \rightarrow Y'$  is closed in  $Y'$  and proper over  $Y$ , and the map  $Z \rightarrow Y'$  factors over it, finishing the proof.  $\square$

In the case of locally spatial v-sheaves, there is another description of partially proper maps.

**Proposition 18.10.** *Let  $f : Y' \rightarrow Y$  be a map from a locally spatial v-sheaf  $Y'$  to a spatial v-sheaf  $Y$ . Then  $f$  is partially proper if and only if  $|Y'|$  is taut, and for every perfectoid field  $K$  with an open and bounded valuation subring  $K^+ \subset K$ , and any diagram*

$$\begin{array}{ccc} \mathrm{Spa}(K, \mathcal{O}_K) & \longrightarrow & Y' \\ \downarrow & \nearrow \text{dotted} & \downarrow f \\ \mathrm{Spa}(K, K^+) & \longrightarrow & Y \end{array}$$

there exists a unique dotted arrow making the diagram commute.

Recall that a locally spectral topological space  $T$  is *taut* if it is quasiseparated, and for every quasicompact open subset  $U \subset T$ , the closure  $\bar{U} \subset T$  of  $U$  is quasicompact, cf. [Hub96, Definition 5.1.2].

*Proof.* In one direction, we have to see that if  $f$  is partially proper, then  $|Y'|$  is taut. Fix a surjection of v-sheaves  $\mathrm{Spa}(S, S^+) \rightarrow Y$  from an affinoid perfectoid space. Let  $|U| \subset |Y'|$  be a quasicompact open subspace, which necessarily comes from some quasicompact open sub-v-sheaf  $U \subset Y'$ . Choose an affinoid perfectoid space  $\mathrm{Spa}(R, R^+)$  with a surjective map of v-sheaves  $\mathrm{Spa}(R, R^+) \rightarrow U \times_Y \mathrm{Spa}(S, S^+)$ . Let  $(R^+)' \subset R$  be the smallest open and integrally closed subring containing the image of  $S^+ \rightarrow R^+$ . As  $f$  is partially proper, we get a unique map  $\mathrm{Spa}(R, (R^+)') \rightarrow Y' \times_Y \mathrm{Spa}(S, S^+)$  extending  $\mathrm{Spa}(R, R^+) \rightarrow U \times_Y \mathrm{Spa}(S, S^+)$ . We claim that the image of  $|\mathrm{Spa}(R, (R^+)')| \rightarrow |Y'|$  is given by  $\bar{U}$ , which is thus quasicompact. Note that as  $\mathrm{Spa}(R, R^+)$  is dense in  $\mathrm{Spa}(R, (R^+)')$ , the image is contained in  $\bar{U}$ . On the other hand, if  $x \in \bar{U}$  is any point, then it corresponds to a map  $\mathrm{Spa}(K, K^+) \rightarrow Y'$  such that  $\mathrm{Spa}(K, \mathcal{O}_K) \subset \mathrm{Spa}(K, K^+) \rightarrow Y'$  factors over  $U$  (as  $\bar{U}$  is the set of specializations of points of  $U$ ). One can lift the image of  $x$  in  $Y$  to  $\mathrm{Spa}(S, S^+)$ . After enlarging  $K$ , we can find a map  $(R, R^+) \rightarrow (K, \mathcal{O}_K)$  inducing the given  $(K, \mathcal{O}_K)$ -point of  $U \times_Y \mathrm{Spa}(S, S^+)$ . Then we get a map  $(R, (R^+)') \rightarrow (K, K^+)$ , so that  $x$  lies in the image of  $|\mathrm{Spa}(R^+, (R^+)')| \rightarrow |Y'|$ , as desired.

For the converse, note that if  $|Y'|$  is taut, we can write it as an increasing union of quasicompact closed generalizing subsets (along closed immersions): Indeed, write  $|Y'| = \bigcup_{i \in I} U_i$  as an increasing union of quasicompact open subsets  $U_i$ ; then their closures  $\bar{U}_i \subset |Y'|$  form an increasing union of quasicompact closed generalizing subsets. Now each  $\bar{U}_i \subset |Y'|$  corresponds to a spatial sub-v-sheaf  $Y'_i \subset Y'$ , which still satisfies the valuative criterion for pairs  $(K, K^+)$ . It follows from Proposition 18.3 that  $Y'_i \rightarrow Y'$  is proper, so that  $Y' \rightarrow Y$  satisfies the criterion of Proposition 18.9.  $\square$



## 19. PROPER BASE CHANGE

In this section, we prove the proper base change theorem for diamonds. It states a certain commutation between pushforward under proper maps, and the functor  $j_!$  for open morphisms  $j$ . Let us briefly recall the latter functor.

**Definition/Proposition 19.1.** *Let  $f : Y' \rightarrow Y$  be an étale morphism of small  $v$ -stacks. Then  $f^* : D_{\text{ét}}(Y, \Lambda) \rightarrow D_{\text{ét}}(Y', \Lambda)$  has a left adjoint  $Rf_! : D_{\text{ét}}(Y', \Lambda) \rightarrow D_{\text{ét}}(Y, \Lambda)$ . Moreover, for any map  $g : \tilde{Y} \rightarrow Y$  with pullback  $\tilde{f} : \tilde{Y}' = Y' \times_Y \tilde{Y} \rightarrow \tilde{Y}$ ,  $g' : \tilde{Y}' \rightarrow Y'$ , the natural transformation*

$$R\tilde{f}_! g'^* \rightarrow g^* Rf_!$$

of functors  $D_{\text{ét}}(Y', \Lambda) \rightarrow D_{\text{ét}}(\tilde{Y}, \Lambda)$ , adjoint to

$$g'^* \rightarrow g'^* f^* Rf_! = \tilde{f}^* g^* Rf_! ,$$

is an equivalence.

As  $Rf_!$  commutes with canonical truncations, we will often denote it simply by  $f_! : D_{\text{ét}}(Y', \Lambda) \rightarrow D_{\text{ét}}(Y, \Lambda)$ .

*Proof.* We start with the case that  $Y$  is a perfectoid space. Then  $Y'$  is also a perfectoid space. In that case,  $Y'_{\text{ét}}$  is a slice of  $Y_{\text{ét}}$ , and thus the functor  $f^*$  on étale sheaves has an exact left adjoint  $f_!$ . This induces a corresponding functor on (non-left-completed) derived categories, which is left adjoint to  $f^*$  as a functor on non-left-completed derived categories; moreover, both  $f^*$  and  $Rf_!$  commute with canonical truncations. This implies that one gets a similar adjunction on left-completions. To check that  $Rf_!$  commutes with base change along maps of perfectoid spaces, use that all functors commute with canonical truncations, so it is enough to prove the assertion if  $\mathcal{F}$  is an étale sheaf on  $Y'$ . Moreover, all functors are defined on all sheaves (not just sheaves of abelian groups), and the adjunction exists in that setting; as all functors commute with all colimits, one reduces to the case of the sheaf represented by some étale map  $Z \rightarrow Y'$ . In that case,  $f_!$  is given by the sheaf represented by the composite étale map  $Z \rightarrow Y' \rightarrow Y$ . It is clear from this description that  $f_!$  commutes with base change.

In general, as  $Y$  is a small  $v$ -stack, we can find a simplicial  $v$ -hypercover of  $Y$  by a simplicial perfectoid space  $Y_\bullet$ ; for simplicity, we assume that all  $Y_i$  are strictly totally disconnected. Let  $f_\bullet : Y'_\bullet \rightarrow Y_\bullet$  be the pullback of  $Y' \rightarrow Y$ , which is again a simplicial perfectoid space. We have the simplicial topos of étale sheaves on  $Y_\bullet$  (and  $Y'_\bullet$ ), roughly given as systems of sheaves  $\mathcal{F}_i$  on  $Y_i$  together with maps  $g^* \mathcal{F}_i \rightarrow \mathcal{F}_j$  for all maps  $g : \Delta^j \rightarrow \Delta^i$ . Correspondingly, we have the derived category  $D(Y_{\bullet, \text{ét}}, \Lambda)$  of sheaves of  $\Lambda$ -modules on the simplicial site  $Y_{\bullet, \text{ét}}$ , and its full subcategory  $D_{\text{cart}}(Y_{\bullet, \text{ét}}, \Lambda)$  consisting of those objects for which the maps  $g^* A_i \rightarrow A_j$  are all equivalences. Then pullback along  $Y_\bullet \rightarrow Y$  induces an equivalence

$$D_{\text{ét}}(Y, \Lambda) \cong D_{\text{cart}}(Y_{\bullet, \text{ét}}, \Lambda) .$$

Similarly,

$$D_{\text{ét}}(Y', \Lambda) \cong D_{\text{cart}}(Y'_{\bullet, \text{ét}}, \Lambda) .$$

The functor  $f_\bullet^* : D(Y_{\bullet, \text{ét}}, \Lambda) \rightarrow D(Y'_{\bullet, \text{ét}}, \Lambda)$  has a left adjoint given by  $Rf_\bullet!$  (which in every degree is given by  $Rf_{i!}$ ). By the base change result, this carries  $D_{\text{cart}}(Y_{\bullet, \text{ét}}, \Lambda)$  into  $D_{\text{cart}}(Y'_{\bullet, \text{ét}}, \Lambda)$ , and thus gives the desired left adjoint  $Rf_!$ . Compatibility with base change follows for pullback along  $Y_0 \rightarrow Y$  by construction, and this implies the general case.  $\square$

Now we can state the proper base change theorem.

**Theorem 19.2.** *Let  $f : Y' \rightarrow Y$  be a proper morphism of small  $v$ -stacks, and let  $j : V \subset Y$  be an open immersion, with pullback  $g : V' = V \times_Y Y' \rightarrow U$ , and  $j' : V' \subset Y'$ . There is a natural transformation of functors*

$$j_! Rg_* \rightarrow Rf_* j'_! : D_{\text{ét}}(U', \Lambda) \rightarrow D_{\text{ét}}(Y, \Lambda) .$$

*If  $f$  is quasi-pro-étale or  $n\Lambda = 0$  for some  $n$  prime to  $p$ , then for any  $A \in D_{\text{ét}}^+(U', \Lambda)$ , the map*

$$j_! Rg_* A \rightarrow Rf_* j'_! A$$

*is an isomorphism.*

*If  $Rf_*$  has finite cohomological dimension, i.e. there is some integer  $N$  such that for all  $A \in D_{\text{ét}}(Y', \Lambda)$  concentrated in degree 0, one has  $R^i f_* A = 0$  for  $i > N$ , then for all  $A \in D_{\text{ét}}^+(U', \Lambda)$ , the map*

$$j_! Rg_* A \rightarrow Rf_* j'_! A$$

*is an isomorphism.*

**Remark 19.3.** This does not quite look like a base change theorem. However, morally it is equivalent to the statement  $Rf_!$  commutes with pullback to the complementary closed subset  $Y \setminus U$ . However, in general  $Y \setminus U \subset Y$  is not generalizing, and so does not correspond to a closed sub- $v$ -sheaf. In [Hub96], Huber defines a notion of pseudo-adic spaces which allows one to treat such subsets as spaces in their own right. In this language, the theorem becomes a base change theorem. We have decided not to introduce an analogue of pseudo-adic spaces in our setup, and so leave the statement of Theorem 19.2 as it is.

*Proof.* The natural transformation  $j_! Rg_* \rightarrow Rf_* j'_!$  is adjoint to

$$f^* j_! Rg_* = j'_! g^* Rg_* \rightarrow j'_! ,$$

using base change for  $j_!$  in Proposition 19.1.

First, we note that if  $Rf_*$  has finite cohomological dimension, then the general case reduces to the case  $A \in D_{\text{ét}}^+(U', \Lambda)$ . Indeed, in that case also  $Rg_* = j^* Rf_* j'_!$  has finite cohomological dimension. Thus, any cohomology sheaf of both sides of

$$j_! Rg_* A \rightarrow Rf_* j'_! A$$

agrees with those of

$$j_! Rg_* \tau^{\geq -n} A \rightarrow Rf_* j'_! \tau^{\geq -n} A$$

for  $n$  sufficiently large.

Now let  $A \in D_{\text{ét}}^+(U', \Lambda)$ ; we want to show that the map

$$j_! Rg_* A \rightarrow Rf_* j'_! A$$

is an isomorphism. By Proposition 17.6 and Proposition 19.1, we can assume that  $Y = X$  is a strictly totally disconnected perfectoid space. The map

$$j_! Rg_* A \rightarrow Rf_* j'_! A$$

becomes a map in  $D_{\text{ét}}^+(X, \Lambda) = D^+(X_{\text{ét}}, \Lambda)$ . To check that it is an isomorphism, we can check on stalks, i.e. after pullback to maps  $\text{Spa}(C, C^+) \rightarrow X$ , where  $C$  is algebraically closed and  $C^+ \subset C$  is an open and bounded valuation subring. Thus, we can assume that  $X = \text{Spa}(C, C^+)$ , and we need

only check the statement on global sections. If  $U = X$ , the result is clear. Otherwise,  $j_! Rg_* A$  has trivial global sections, so we need to see that

$$R\Gamma(X, Rf_* j_! A) = 0 .$$

But this is given by

$$R\Gamma(X, Rf_* j_! A) = R\Gamma(Y', j_! A) .$$

If  $f$  is quasi-pro-étale, then  $Y'$  is a qcqs perfectoid space proper and pro-étale over  $X = \mathrm{Spa}(C, C^+)$ , i.e.  $Y' = \mathrm{Spa}(C, C^+) \times \underline{S}$  for some profinite set  $S$ . In that case, the result is clear. Thus, from now on assume that  $n\Lambda = 0$  for some  $n$  prime to  $p$ .

Let  $X'$  be a strictly totally disconnected space with a surjection  $X' \rightarrow Y'$ . By Corollary 18.8 (vi), the canonical compactification  $\overline{X}'/X$  is proper over  $X$ , and  $\overline{X}'/X = \overline{X}' \times_{\overline{X}} X$  is an affinoid perfectoid space by Proposition 18.7 (iv). There is a unique extension of  $X' \rightarrow Y'$  to  $\overline{X}'/X \rightarrow Y'$  by Proposition 18.6. The fibre product  $\overline{X}'/X \times_{Y'} \overline{X}'/X$  is again proper over  $X$ ; continuing, we can produce a v-hypercover  $Y'_\bullet \rightarrow Y'$  where each  $Y'_i = \overline{X}'_i/X$  is the canonical compactification of some strictly totally disconnected  $X'_i \rightarrow X$ . Then, by unbounded cohomological descent in a replete topos, cf. [BS15, Proposition 3.3.6],

$$R\Gamma(Y', j_! A)$$

is the (derived) limit of the simplicial object

$$R\Gamma(\overline{X}'_i/X, j_! A) .$$

Thus, it is enough to handle the case that  $Y' = \overline{X}'/X$  is the canonical compactification of some strictly totally disconnected  $X' \rightarrow X$ . In particular, in this case  $Y'$  is an affinoid perfectoid space, and  $A \in D_{\text{ét}}^+(Y', \Lambda) = D^+(Y'_{\text{ét}}, \Lambda)$ . Let  $X' \rightarrow \pi_0 X'$  be the projection; this extends to a projection  $Y' \rightarrow \pi_0 Y' = \pi_0 Y'$ . Computing  $R\Gamma(Y', -)$  via a Leray spectral sequence along  $Y' \rightarrow \pi_0 Y'$ , it is enough to check that the fibres of the pushforward vanish; but these are given by the cohomology of the fibers. Thus, we can assume that  $Y'$  is connected. In other words,  $X' = \mathrm{Spa}(C', C'^+)$  for some complete algebraically closed field  $C'$  (over  $C$ ), and some open and bounded valuation subring  $C'^+ \subset C'$ . Then  $Y' = \overline{X}'/X$  is given by

$$Y' = \mathrm{Spa}(C', C'^{\circ\circ} + C'^+) .$$

Now  $Y'$  has only one rank-1-point  $\mathrm{Spa}(C', \mathcal{O}_{C'})$ , where  $C'$  is algebraically closed; it follows that any map in  $Y'_{\text{ét}}$  is a local isomorphism, and thus the topos  $(Y'_{\text{ét}})^\sim$  is equivalent to the topos  $|Y'|^\sim$ . Let  $K'$  resp.  $K$  be the (algebraically closed) residue field of  $C'$  resp.  $C$ , and let  $V \subset K$  correspond to  $C^+ \subset \mathcal{O}_C$ , i.e.  $V = C^+/C^{\circ\circ} \subset K = \mathcal{O}_C/C^{\circ\circ}$ . Then  $|Y'| = |\mathrm{Spa}(K', V)|$ . Thus, we are reduced to the following lemma about the Zariski–Riemann spaces of algebraically closed fields.  $\square$

The following lemma is related to results of Huber, [Hub93b].

**Lemma 19.4.** *Let  $K \subset K'$  be algebraically closed fields, and let  $V \subset K$  be a valuation ring of  $K$ . Let  $T' = \mathrm{Spa}(K', V)$  be the spectral space of all valuation rings  $V' \subset K'$  that contain  $V$ , and let  $T = \mathrm{Spa}(K, V)$ , with the natural projection map  $f : T' \rightarrow T$  sending  $V' \subset K'$  to  $V' \cap K \subset K$ . Let  $s \in T$  denote the unique closed point, corresponding to the valuation ring  $V \subset K$ . Let  $\mathcal{F}$  be a sheaf of torsion abelian groups on  $T'$  such that for all  $x' \in f^{-1}(s)$ , the stalk  $\mathcal{F}_{x'} = 0$ . Then*

$$R\Gamma(T', \mathcal{F}) = 0 .$$

*Proof.* Consider the category  $C$  of all proper  $V$ -schemes  $X$  with a point  $x \in X(K')$ . This category is cofiltered: Given such  $(X, x)$  and  $(X', x')$ , the product  $X \times_V X'$  with product point  $(x, x')$  dominates both, and if  $f, g : X \rightarrow X'$  are two morphisms mapping  $x$  to  $x'$ , then the equalizer of  $f$  and  $g$  is also proper over  $V$  and contains  $x$ . Then  $T' = \varprojlim_C |X|$ ; in fact, one could replace  $C$  by the subcategory of integral and projective  $X$  for which the point  $x$  is dominant, cf. [Hub93b, Lemma 2.1]. Moreover, in the limit, the topoi  $\varprojlim_C |X|^\sim$  and  $\varprojlim_C X_{\text{ét}}^\sim$  are equivalent (as any étale map becomes a local isomorphism after some pullback), cf. [Hub93b, Lemma 2.4].

We may assume that  $\mathcal{F}$  is constructible. Then it comes via pullback from some  $\mathcal{F}_X$  on  $X_{\text{ét}}^\sim$  (with trivial restriction to the special fiber) for some  $X \in C$ , and the result follows from the proper base change theorem in étale cohomology applied to all  $X' \in C_{/X}$ , noting that

$$R\Gamma(T', \mathcal{F}) = \varinjlim_{X' \in C_{/X}} R\Gamma(X'_{\text{ét}}, \mathcal{F}_{X'}) ,$$

where  $\mathcal{F}_{X'}$  denotes the pullback of  $\mathcal{F}_X$  to  $X'$ . □

As a first application of the proper base change theorem, we can complete the proof of invariance under change of algebraically closed base field. Let us recall the statement from the introduction.

**Theorem 19.5.** *Let  $Y$  be a small  $v$ -stack, and assume that  $n\Lambda = 0$  for some  $n$  prime to  $p$ .*

(i) *Assume that  $Y$  lives over  $k$ , where  $k$  is a discrete algebraically closed field of characteristic  $p$ , and  $k'/k$  is an extension of discrete algebraically closed base fields,  $Y' = Y \times_k k'$ . Then the pullback functor*

$$D_{\text{ét}}(Y, \Lambda) \rightarrow D_{\text{ét}}(Y', \Lambda)$$

*is fully faithful.*

(ii) *Assume that  $Y$  lives over  $k$ , where  $k$  is an algebraically closed discrete field of characteristic  $p$ . Let  $C/k$  be an algebraically closed complete nonarchimedean field, and  $Y' = Y \times_k \text{Spa}(C, C^+)$  for some open and bounded valuation subring  $C^+ \subset C$  containing  $k$ . Then the pullback functor*

$$D_{\text{ét}}(Y, \Lambda) \rightarrow D_{\text{ét}}(Y', \Lambda)$$

*is fully faithful.*

(iii) *Assume that  $Y$  lives over  $\text{Spa}(C, C^+)$ , where  $C$  is an algebraically closed complete nonarchimedean field with an open and bounded valuation subring  $C^+ \subset C$ ,  $C'/C$  is an extension of algebraically closed complete nonarchimedean fields, and  $C'^+ \subset C'$  an open and bounded valuation subring containing  $C^+$ , such that  $\text{Spa}(C', C'^+) \rightarrow \text{Spa}(C, C^+)$  is surjective. Then for  $Y' = Y \times_{\text{Spa}(C, C^+)} \text{Spa}(C', C'^+)$ , the pullback functor*

$$D_{\text{ét}}(Y, \Lambda) \rightarrow D_{\text{ét}}(Y', \Lambda)$$

*is fully faithful.*

*Proof.* We note that (i) follows from (ii) and (iii). Moreover, (ii) follows from (iii) and the restricted version of (ii) where we demand that  $C$  is the completed algebraic closure of  $k((t))$ .

Let us start by proving (iii). Let  $f : Y' \rightarrow Y$  be the map, which is qcqs. We need to see that the adjunction map

$$A \rightarrow Rf_* f^* A$$

is an isomorphism for all  $A \in D_{\text{ét}}(X, \Lambda)$ . First, note that by writing  $A$  as the homotopy limit of  $\tau^{\geq -n} A$ , it is enough to prove this for  $A \in D_{\text{ét}}^+(X, \Lambda)$ . In that case,  $Rf_* = Rf_{v*}$  commutes with any

base change by Proposition 17.6. Thus, we may assume that  $Y = X$  is a strictly totally disconnected perfectoid space. Now the statement follows from (the final paragraph of) Theorem 16.1.

It remains to prove (ii) in the case that  $C$  is the completed algebraic closure of  $k((t))$ . Let  $Y_\bullet \rightarrow Y$  be a simplicial  $v$ -hypercover of  $Y$  by disjoint unions of strictly totally disconnected spaces  $Y_i$ ; then  $Y'_i = Y' \times_Y Y_i$  is a perfectoid space. Under the identifications

$$D_{\text{ét}}(Y, \Lambda) \simeq D_{\text{ét, cart}}(Y_\bullet, \Lambda) , \quad D_{\text{ét}}(Y', \Lambda) \simeq D_{\text{ét, cart}}(Y'_\bullet, \Lambda) ,$$

it is enough to prove that

$$D_{\text{ét}}(Y_i, \Lambda) \rightarrow D_{\text{ét}}(Y'_i, \Lambda)$$

is fully faithful for all  $i$ . In other words, we can assume that  $Y = \text{Spa}(A, A^+)$  is a strictly totally disconnected space.

Now fix a pseudouniformizer  $\varpi \in A$ . In that case,  $Y'$  is the increasing union of the affinoid perfectoid spaces

$$Y'_n = \{|t|^n \leq |[\varpi]| \leq |t|^{1/n}\} \subset Y' .$$

As  $R\text{Hom}$ 's in  $D_{\text{ét}}(Y', \Lambda)$  are the derived limit of  $R\text{Hom}$ 's in  $D_{\text{ét}}(Y'_n, \Lambda)$ , it is enough to prove that for all  $n \geq 1$ , the functor

$$D_{\text{ét}}(Y, \Lambda) \rightarrow D_{\text{ét}}(Y'_n, \Lambda)$$

is fully faithful. Let  $f_n : Y'_n \rightarrow Y$  be the map of affinoid perfectoid spaces. We need to see that the adjunction map

$$A \rightarrow Rf_{n*} f_n^* A$$

is an equivalence for all  $A \in D_{\text{ét}}(Y, \Lambda)$ . Again, this reduces to the case  $A \in D_{\text{ét}}^+(Y, \Lambda)$ . The desired statement can be checked on stalks, so we can assume that  $Y = \text{Spa}(C', C'^+)$ , where  $C'$  is an algebraically closed nonarchimedean field with an open and bounded valuation subring  $C'^+ \subset C'$ , and we only need to verify the statement on global sections. Thus, for any  $A \in D_{\text{ét}}(Y, \Lambda)$ , we need to see that

$$R\Gamma(Y, A) = R\Gamma(Y'_n, f_n^* A) .$$

By Theorem 19.2, both sides vanish if  $A = j_! A_0$  for some  $A_0 \in D_{\text{ét}}^+(U, \Lambda)$ , where  $U = Y \setminus \{s\} \subset Y$  is the complement of the closed point  $s \in Y$ . Thus, replacing  $A$  by the cone of  $j_! j^* A \rightarrow A$ , we can assume that  $A$  is concentrated at the closed point  $s$ . Repeating the argument in the other direction, we can assume that  $A$  is constant. We can also assume that  $\Lambda = \mathbb{Z}/\ell^n \mathbb{Z}$ , and then by triangles that  $\Lambda = \mathbb{F}_\ell$ . In that case,  $A$  is a direct sum of shifted copies of  $\mathbb{F}_\ell$  (but bounded below). This finally reduces us to the case  $A = \mathbb{F}_\ell$ .

Thus, it remains to see that

$$R\Gamma(Y'_n, \mathbb{F}_\ell) = \mathbb{F}_\ell .$$

But note that  $Y'_n$  is the inverse limit over all finite extensions  $L \subset C$  of  $k((t^{1/p^\infty}))$  of the system of affinoid perfectoid spaces given by

$$Y'_{n,L} = \{|t|^n \leq |[\varpi]| \leq |t|^{1/n}\} \subset \text{Spa}(L, \mathcal{O}_L) \times_k \text{Spa}(C', C'^+) ,$$

which implies

$$H^i(Y'_n, \mathbb{F}_\ell) = \varinjlim H^i(Y'_{n,L}, \mathbb{F}_\ell)$$

for all  $i \geq 0$  by Proposition 14.9. On the other hand, all  $L$  are isomorphic to  $k((t_L^{1/p^\infty}))$  for some pseudo-uniformizer  $t_L$ , as one can take for  $t_L$  a uniformizer of the finite separable extension of

$k((t))$  corresponding to  $L$  (noting that finite separable extensions of  $k((t))$  are equivalent to finite extensions of  $k((t^{1/p^\infty}))$ ). Thus,  $Y'_{n,L}$  is an annulus over  $\mathrm{Spa}(C, C^+)$ , which implies that

$$H^i(Y'_{n,L}, \mathbb{F}_\ell) = \begin{cases} \mathbb{F}_\ell & i = 0, 1 \\ 0 & \text{else.} \end{cases}$$

Taking the direct limit over all  $L$  kills the class in degree 1 by extracting  $\ell$ -power roots of the uniformizer; this finishes the proof.  $\square$

Before going on and using the proper base change theorem to define the functor  $Rf_!$ , we pause to obtain certain criteria guaranteeing finite cohomological dimension for  $Rf_*$ .

## 20. CONSTRUCTIBLE SHEAVES

It will be convenient to have a general notion of constructible sheaves. This notion is slightly subtle as locally closed subsets of adic spaces are not adic spaces themselves. This is one reason that Huber considers pseudo-adic spaces in [Hub96]. We will work around this issue.

**Definition 20.1.** *Let  $\Lambda$  be a noetherian ring.*

- (i) *Let  $X$  be a strictly totally disconnected perfectoid space, and identify the topos of étale sheaves on  $X$  with the topos of sheaves on  $|X|$ . A sheaf  $\mathcal{F}$  of  $\Lambda$ -modules on  $X_{\text{ét}}$  (equivalently, on  $|X|$ ) is constructible if there is stratification of  $X$  into constructible locally closed subsets  $S_i \subset |X|$  such that  $\mathcal{F}|_{S_i}$  is the constant sheaf on  $S_i$  associated with some finitely generated  $\Lambda$ -module.*
- (ii) *Let  $Y$  be a small  $v$ -sheaf, and  $\mathcal{F}$  a small sheaf of  $\Lambda$ -modules on  $Y_v$ . Then  $\mathcal{F}$  is constructible if  $\mathcal{F} \in D_{\text{ét}}(Y, \Lambda) \subset D(Y, \Lambda)$ , and for every strictly totally disconnected space  $f : X \rightarrow Y$ , the pullback  $f^*\mathcal{F}$  is constructible.*

**Remark 20.2.** Note that in part (i), we made a switch from  $X$  as a perfectoid space to  $|X|$  as a mere topological space, which allowed us to restrict to the topological space  $S_i$ .

**Remark 20.3.** It is clear from the definition that the class of constructible sheaves is stable under kernels, cokernels, images, and extensions; in particular, constructible sheaves form an abelian category. Indeed, this reduces immediately to the case where  $X$  is strictly totally disconnected, in which case it reduces further (by passing to suitably refined stratifications) to the case of a map of constant sheaves (associated with finitely generated  $\Lambda$ -modules) on a spectral space, where finally it reduces (on open and closed subsets) to the category of finitely generated  $\Lambda$ -modules.

We will show that this property has good descent properties, and will give a better definition in the case of spatial diamonds. First, we analyze constructible sheaves on strictly totally disconnected perfectoid spaces  $X$ . In fact, this reduces to sheaves on the spectral space  $|X|$ , where we have the following general lemma.

**Lemma 20.4.** *Let  $\Lambda$  be a noetherian ring, let  $X$  be a spectral space, and let  $\mathcal{F}$  be a sheaf of  $\Lambda$ -modules on  $X$ . Then  $\mathcal{F}$  is constructible if and only if  $\mathcal{F}$  is compact in the category of sheaves of  $\Lambda$ -modules on  $X$ , i.e.  $\mathrm{Hom}(\mathcal{F}, -)$  commutes with filtered colimits. Moreover, any sheaf of  $\Lambda$ -modules on  $X$  can be written as a filtered colimit of constructible sheaves.*

*Proof.* First, we check that constructible sheaves are compact. For this, note that both  $j_!$  for constructible open immersions  $j$  and  $i_*$  for constructible closed immersions  $i$  preserve compact objects (as their right adjoints  $j^*$  resp.  $i^!$  preserve filtered colimits). Passing to a filtration, this

reduces the problem to the case that  $\mathcal{F}$  is the constant sheaf associated with some finitely generated  $\Lambda$ -module, where the result follows by taking a finite free 2-term resolution.

Next, we show that every sheaf of  $\Lambda$ -modules on  $X$  can be written as a filtered colimit of constructible sheaves. Note that any sheaf  $\mathcal{F}$  admits a surjection from a direct sum  $\mathcal{G}$  of sheaves of the form  $j_!\Lambda$ , where  $j : U \hookrightarrow X$  ranges over quasicompact open immersions into  $X$ . Applying the same to the kernel of  $\mathcal{G} \rightarrow \mathcal{F}$ , we get a 2-term resolution

$$\mathcal{H} \rightarrow \mathcal{G} \rightarrow \mathcal{F} \rightarrow 0 ,$$

where  $\mathcal{G}$  and  $\mathcal{H}$  are direct sums of sheaves of the form  $j_!\Lambda$ . The map  $\mathcal{H} \rightarrow \mathcal{G}$  is then a filtered colimit of similar maps  $\mathcal{H}_i \rightarrow \mathcal{G}_i$ , where  $\mathcal{H}_i$  and  $\mathcal{G}_i$  are finite sums of sheaves of the form  $j_!\Lambda$  (as such sheaves are compact). The cokernels  $\mathcal{F}_i$  of the maps  $\mathcal{H}_i \rightarrow \mathcal{G}_i$  are constructible, and  $\mathcal{F}$  is their filtered colimit, as desired.

Now, if  $\mathcal{F}$  is compact, then write  $\mathcal{F} = \varinjlim_i \mathcal{F}_i$  as a filtered colimit of constructible sheaves. By compactness, one can factor the map  $\mathcal{F} \rightarrow \varinjlim_i \mathcal{F}_i = \mathcal{F}$  over  $\mathcal{F}_i$  for  $i$  sufficiently large. This shows that  $\mathcal{F}$  is a direct summand of  $\mathcal{F}_i$ , and in particular constructible itself.  $\square$

**Proposition 20.5.** *Let  $\Lambda$  be a noetherian ring, let  $f : \tilde{Y} \rightarrow Y$  be a surjective map of small  $v$ -sheaves, and let  $\mathcal{F}$  be a small sheaf of  $\Lambda$ -modules. If  $f^*\mathcal{F}$  is constructible, then  $\mathcal{F}$  is constructible.*

*Proof.* We may assume that both  $Y = X$  and  $\tilde{Y} = \tilde{X}$  are strictly totally disconnected spaces. In that case, it follows from Lemma 20.4 as  $\mathrm{Hom}_X(\mathcal{F}, -)$  can be written as the equalizer of

$$\mathrm{Hom}_{\tilde{X}}(f^*\mathcal{F}, f^*- ) \rightrightarrows \mathrm{Hom}_{\tilde{X}}(g^*\mathcal{F}, g^*- ) ,$$

where  $\tilde{X} \rightarrow \tilde{X} \times_X \tilde{X}$  (with composite  $g : \tilde{X} \rightarrow X$ ) is some strictly totally disconnected cover, and these functors commute with filtered colimits by assumption.  $\square$

**Proposition 20.6.** *Let  $\Lambda$  be a noetherian ring, and let  $Y$  be a spatial diamond. Let  $\mathcal{F}$  be an étale sheaf of  $\Lambda$ -modules on  $Y$ . The following conditions are equivalent.*

- (i) *The sheaf  $\mathcal{F}$  is constructible.*
- (ii) *The sheaf  $\mathcal{F}$  is compact in the category of étale sheaves of  $\Lambda$ -modules on  $Y$ .*
- (iii) *There is a stratification of  $|Y|$  into constructible locally closed subsets  $S_i \subset |Y|$  such that the restriction of  $\mathcal{F}$  to  $S_i$  satisfies the following condition: For any strictly totally disconnected perfectoid space  $f : X \rightarrow Y$ , the pullback  $f^*\mathcal{F}|_{f^{-1}(S_i)}$  is the constant sheaf associated with some finitely generated  $\Lambda$ -module.*

*Moreover, any étale sheaf of  $\Lambda$ -modules on  $Y$  can be written as a filtered colimit of constructible sheaves.*

*Proof.* First, (i) implies (ii), as compactness descends over a  $v$ -cover  $X \rightarrow Y$  by a strictly totally disconnected perfectoid space, as in the proof of Proposition 20.5. Moreover, by definition (iii) implies (i).

It remains to see that (ii) implies (iii). Let us for the moment call a sheaf  $\mathcal{F}$  satisfying the hypothesis of (iii) strongly constructible. Then strongly constructible sheaves are constructible. It now suffices to show that any étale sheaf of  $\Lambda$ -modules on  $Y$  can be written as a filtered colimit of strongly constructible sheaves. Indeed, it will then follow that any compact  $\mathcal{F}$  is a direct summand of a strongly constructible sheaf, and therefore strongly constructible itself.

Thus, it remains to see that any étale sheaf of  $\Lambda$ -modules on  $Y$  can be written as a filtered colimit of strongly constructible sheaves. As in Remark 20.3, the category of strongly constructible sheaves is closed under kernels, cokernels, images, and extensions. By Lemma 11.31, any sheaf of  $\Lambda$ -modules on  $Y_{\text{ét}}$  can be written as a quotient of a direct sum of sheaves of the form  $j_!\Lambda$  where  $j : U \rightarrow Y$  ranges over étale maps which can be written as a composite  $U \hookrightarrow V \rightarrow W \hookrightarrow Y$ , where  $U \hookrightarrow V$  and  $W \hookrightarrow Y$  are quasicompact open immersions, and  $V \rightarrow W$  is finite étale. Arguing as in Lemma 20.4, it is enough to prove that  $j_!\Lambda$  is strongly constructible. Consider the function  $a : |Y| \rightarrow \mathbb{Z}_{\geq 0}$  which assigns to any point  $y \in Y$  the number of geometric points of  $U$  above a fixed geometric point  $\bar{y}$  over  $y$ . We claim that the level sets  $a^{-1}(n)$  are constructible locally closed subsets of  $|Y|$  for all  $n \geq 0$ . As  $a$  is bounded, it suffices to show that  $a^{-1}(\mathbb{Z}_{\geq n}) \subset |Y|$  is a quasicompact open subset for all  $n \geq 0$ . This statement can be checked  $v$ -locally, so we can assume that  $Y = X$  is a strictly disconnected perfectoid space, where it is a direct verification. Let  $S_i = a^{-1}(i)$  for  $i \geq 0$ . It remains to see that for all strictly totally disconnected perfectoid spaces  $f : X \rightarrow Y$ , the pullback  $f^*(j_!\Lambda)|_{f^{-1}(S_i)}$  is the constant sheaf on  $f^{-1}(S_i)$  associated with  $\Lambda^i$ . Again, this is a direct verification.  $\square$

**Proposition 20.7.** *Let  $Y_i$ ,  $i \in I$ , be a cofiltered inverse system of spatial diamonds with inverse limit  $Y = \varprojlim_i Y_i$ , which is again a spatial diamond. Let  $\Lambda$  be a noetherian ring, and denote by  $\text{Cons}(Y, \Lambda)$  (resp.  $\text{Cons}(Y_i, \Lambda)$ ) the category of constructible étale sheaves of  $\Lambda$ -modules on  $Y$  (resp.  $Y_i$ ). Then the natural functor*

$$2\text{-}\varinjlim_i \text{Cons}(Y_i, \Lambda) \rightarrow \text{Cons}(Y, \Lambda)$$

*is an equivalence of categories.*

*Proof.* First, we check fully faithfulness. For this, let  $\mathcal{F}_{i_0}, \mathcal{G}_{i_0} \in \text{Cons}(Y_{i_0}, \Lambda)$  for some  $i_0$  with pullbacks  $\mathcal{F}_i, \mathcal{G}_i \in \text{Cons}(Y_i, \Lambda)$  and  $\mathcal{F}, \mathcal{G} \in \text{Cons}(Y, \Lambda)$ . Let  $f_j : Y \rightarrow Y_j$  and  $f_{ij} : Y_i \rightarrow Y_j$  be the natural maps. Then

$$\begin{aligned} \text{Hom}_Y(\mathcal{F}, \mathcal{G}) &= \text{Hom}_{Y_{i_0}}(\mathcal{F}_{i_0}, f_{i_0*}\mathcal{G}) \\ &= \text{Hom}_{Y_{i_0}}(\mathcal{F}_{i_0}, \varinjlim_i f_{i,i_0*}\mathcal{G}_i) \\ &= \varinjlim_i \text{Hom}_{Y_{i_0}}(\mathcal{F}_{i_0}, f_{i,i_0*}\mathcal{G}_i) \\ &= \varinjlim_i \text{Hom}_{Y_i}(\mathcal{F}_i, \mathcal{G}_i), \end{aligned}$$

using obvious adjunctions, (the relative version of) Proposition 14.9 to write  $f_{i_0*}\mathcal{G} = \varinjlim_i f_{i,i_0*}\mathcal{G}_i$ , and compactness of  $\mathcal{F}_{i_0}$ .

Now, for essential surjectivity, note that any constructible sheaf  $\mathcal{F}$  on  $Y$  is a quotient of a map of finite direct sums of  $j_!\Lambda$ , where  $j : U \rightarrow Y$  runs through quasicompact separated étale maps, as the proof of Proposition 20.6 shows. Using Proposition 11.23 and fully faithfulness, we see that this data is defined over some  $Y_i$ , as desired.  $\square$

Now we can prove a slightly more explicit characterization of constructibility, which will be necessary for the passage to derived categories.

**Proposition 20.8.** *Let  $\Lambda$  be a noetherian ring, and let  $Y$  be a spatial diamond. Then an étale sheaf  $\mathcal{F}$  of  $\Lambda$ -modules on  $Y$  is constructible if and only if  $\mathcal{F}$  has a filtration whose graded pieces*



$\mathcal{F}_i$  are of the form  $j_!(\mathcal{L}|_Z)$ , where  $j : U \rightarrow Y$  is a quasicompact separated étale map,  $Z \subset U$  is a constructible closed subset, and  $\mathcal{L}$  is a sheaf of  $\Lambda$ -modules on  $U$  that is locally on  $U_{\text{ét}}$  isomorphic to the constant sheaf associated with some finitely generated  $\Lambda$ -module.

Here,  $\mathcal{L}|_Z$  is the cokernel of the injective map  $j'_!\mathcal{L}|_{U \setminus Z} \rightarrow \mathcal{L}$ , where  $j' : U \setminus Z \rightarrow U$  is the open embedding.

*Proof.* It is clear that any sheaf admitting such a filtration is constructible. Conversely, we can find a stratification  $S_1, \dots, S_m$  of  $Y$  as in Proposition 20.6 (iii), with the extra property that for all  $i = 1, \dots, m$ , the union  $U_i = S_1 \sqcup \dots \sqcup S_i$  is (quasicompact and) open. One can filter  $\mathcal{F}$  by the extensions by 0 of  $\mathcal{F}|_{U_i}$ ; passing to associated graded, we can assume that there is some constructible locally closed subset  $S \subset |Y|$  such that the stalks of  $\mathcal{F}$  at all points outside of  $S$  vanish, and  $\mathcal{F}|_S$  satisfies the final condition of Proposition 20.6 (iii): For all strictly totally disconnected perfectoid spaces  $f : X \rightarrow Y$ , the restriction  $f^*\mathcal{F}|_{f^{-1}(S)}$  is constant.

Moreover, the claim can be checked locally: Indeed, assume it is true for the restrictions  $\mathcal{F}|_{V_i}$  for an open covering of  $\{V_i\}$  of  $Y$ . By induction, we may assume that there are only two subsets  $V_1, V_2 \subset Y$ ; let  $j_i : V_i \rightarrow Y$  be the open subsets for  $i = 1, 2$ , as well as  $j_{12} : V_{12} = V_1 \cap V_2 \subset Y$ . Then  $\mathcal{F}$  has a subsheaf  $j_{1!}\mathcal{F}|_{V_1}$  of the desired form, and the quotient  $\mathcal{F}/j_{1!}\mathcal{F}|_{V_1}$  can be identified with  $j_{2!}(\mathcal{F}|_{V_2}|_{V_1 \setminus V_{12}})$ , which is again of the desired form.

Now we claim that locally,  $\mathcal{F}$  is indeed of the form  $j_!(\mathcal{L}|_Z)$  for a quasicompact separated étale map  $j : U \rightarrow Y$ , some  $\mathcal{L}$  on  $U$  as in the statement, and a constructible closed subset  $Z \subset U$ . This can be checked after pullback to the localization  $Y_y$  at varying points  $y \in |Y|$ , by Proposition 20.7.

Thus, we can assume that  $Y$  is local, so let  $\text{Spa}(C, C^+) \rightarrow Y$  be a quasi-pro-étale surjection, where  $C$  is algebraically closed and  $C^+ \subset C$  an open and bounded valuation subring. Then  $|Y| = |\text{Spa}(C, C^+)|$  is a totally ordered chain of points with a unique closed point  $s \in |Y|$ . Moreover, by the first paragraph, there is some constructible locally closed subset  $S \subset |Y|$  such that  $\mathcal{F}|_{\text{Spa}(C, C^+) \setminus S} = 0$  and  $\mathcal{F}|_{S \subset \text{Spa}(C, C^+)}$  is constant, with value some finitely generated  $\Lambda$ -module  $M$ . We can assume that  $s \in S$ , so that  $S$  is actually closed. Let  $\eta_S \in S$  be the generic point of  $S$ , and let  $G_S = G_{\eta_S}$  be the profinite group given as the fibre of  $|R| \rightarrow |Y|$ ,  $R = \text{Spa}(C, C^+) \times_Y \text{Spa}(C, C^+)$ , over  $\eta_S$  (where the group structure comes from the equivalence relation structure). Then  $\mathcal{F}$  is given by a continuous action of  $G_S$  on  $M$ . Similarly, let  $\eta \in |Y|$  be the generic point, and  $G_\eta$  the profinite group that is the fibre of  $|\text{Spa}(C, C^+) \times_Y \text{Spa}(C, C^+)| \rightarrow |Y|$  over  $\eta$  (so that the open subspace  $Y^\circ$  of  $Y$  with underlying space  $\{\eta\}$  is given by  $\text{Spa}(C, \mathcal{O}_C)/\overline{G_\eta}$ ). There is natural closed immersion  $G_S \hookrightarrow G_\eta$  of groups, given by generalization. One can find an open subgroup  $H \subset G_\eta$  containing  $G_S$  such that the action of  $G_S$  on  $M$  extends to  $H$ . Let  $R_H \subset R$  be the open and closed subspace given as the closure of  $H \subset G_\eta = R \times_Y \{\eta\}$ . Then  $U = \text{Spa}(C, C^+)/\underline{H}$  is separated and étale over  $Y$ , and the continuous  $H$ -module  $M$  defines a local system  $\mathcal{L}$  on  $U$ , which is easily checked to have the right property.  $\square$

We will also need a compactness result in the derived category. We start with an easy result.

**Proposition 20.9.** *Let  $Y$  be a spatial diamond, and assume that  $\mathcal{F}$  is a constructible étale sheaf of  $\mathbb{F}_\ell$ -vector spaces on  $Y$ . Then for all  $C_j \in D^{\geq -n}(Y_{\text{ét}}, \mathbb{F}_\ell)$ ,  $j \in J$ , of complexes uniformly bounded to the left, the natural map*

$$\bigoplus_{j \in J} \text{Hom}_{D(Y_{\text{ét}}, \mathbb{F}_\ell)}(\mathcal{F}[0], C_j) \rightarrow \text{Hom}_{D(Y_{\text{ét}}, \mathbb{F}_\ell)}(\mathcal{F}[0], \bigoplus_{j \in J} C_j)$$

is an isomorphism.

*Proof.* By descent (and using boundedness), this can be reduced to the case that  $Y = X$  is strictly totally disconnected. Decomposing  $\mathcal{F}$  into finitely many triangles, we can assume that  $\mathcal{F} = j_! \mathbb{F}_\ell$  for some quasicompact open immersion  $j : U \hookrightarrow X$ . Then the statement becomes

$$\bigoplus_{j \in J} R\Gamma(U_{\text{ét}}, C_j) \xrightarrow{\cong} R\Gamma(U_{\text{ét}}, \bigoplus_{j \in J} C_j),$$

which holds true as  $U_{\text{ét}}$  is coherent and the  $C_j$  are uniformly bounded to the left.  $\square$

One gets a stronger result if one assumes that  $Y$  is locally of finite  $\ell$ -cohomological dimension.

**Proposition 20.10.** *Let  $Y$  be a spatial diamond, and assume that there is an integer  $N$  such that for all quasicompact separated étale maps  $U \rightarrow Y$  the  $\ell$ -cohomological dimension of  $U_{\text{ét}}$  is  $\leq N$ , i.e. for all  $\ell$ -torsion sheaves  $\mathcal{F}$  on  $U_{\text{ét}}$ , one has  $H^i(U_{\text{ét}}, \mathcal{F}) = 0$  for  $i > N$ .*

*Then  $D(Y_{\text{ét}}, \mathbb{F}_\ell)$  is left-complete (thus  $D(Y_{\text{ét}}, \mathbb{F}_\ell) = D_{\text{ét}}(Y, \mathbb{F}_\ell)$ ), compactly generated, and a complex  $C \in D_{\text{ét}}(Y, \mathbb{F}_\ell)$  is compact if and only if it is bounded and all cohomology sheaves are constructible.*

*Proof.* Left-completeness of  $D(Y_{\text{ét}}, \mathbb{F}_\ell)$  follows from [Sta, Tag 0719]. This implies  $D(Y_{\text{ét}}, \mathbb{F}_\ell) = D_{\text{ét}}(Y, \mathbb{F}_\ell)$  by Proposition 14.15.

First, we check that a complex is compact if it is bounded with all cohomology sheaves constructible. This reduces immediately to the case that  $C = \mathcal{F}[0]$  for some constructible sheaf  $\mathcal{F}$ . Using Proposition 20.8, this reduces further to the case  $\mathcal{F} = j_! \mathcal{L}$  for some quasicompact separated étale map  $j : U \rightarrow Y$  and  $\mathbb{F}_\ell$ -local system  $\mathcal{L}$  on  $U$ . But then

$$\text{Hom}_{D(Y_{\text{ét}}, \mathbb{F}_\ell)}(j_! \mathcal{L}, -) = \text{Hom}_{D(U_{\text{ét}}, \mathbb{F}_\ell)}(\mathcal{L}, -) = R\Gamma(U_{\text{ét}}, \mathcal{L}^\vee \otimes_{\mathbb{F}_\ell} -),$$

so we have to prove that  $R\Gamma(U_{\text{ét}}, -)$  commutes with direct sums in  $D(U_{\text{ét}}, \mathbb{F}_\ell)$ . This follows easily from left-completeness of  $D_{\text{ét}}(U, \mathbb{F}_\ell)$ , the assumption that  $U_{\text{ét}}$  has finite  $\ell$ -cohomological dimension, and Proposition 20.9.

As the derived category is generated by  $j_! \mathbb{F}_\ell$  for varying quasicompact separated étale maps  $j : U \rightarrow Y$ , this shows that  $D(Y_{\text{ét}}, \mathbb{F}_\ell)$  is compactly generated. By abstract nonsense, any compact object is a direct summand of a finite complex whose terms are finite direct sums of sheaves of the form  $j_! \mathbb{F}_\ell$  for varying quasicompact separated étale maps  $j : U \rightarrow Y$ . All of those are bounded with constructible cohomology sheaves, as desired.  $\square$

## 21. DIMENSIONS

To discuss duality, it will be important to impose a “finite-dimensionality” assumption on morphisms  $f$ . For this, we restrict to locally spatial morphisms, and we will put a condition on transcendence degrees of residue fields.

For a morphism  $f : X' \rightarrow X$  of perfectoid spaces, one has the definition  $\dim f$  as given in [Hub96, Definition 1.8.4], and of  $\dim. \text{tr } f$  as given in [Hub96, Definition 1.8.4]. Let us briefly recall those.

### Definition 21.1.

- (i) *Let  $X$  be a locally spectral space. The dimension  $\dim X \in \mathbb{Z}_{\geq 0} \cup \{-\infty, \infty\}$  of  $X$  is the supremum of all integers  $n$  for which there exists a chain  $x_0, \dots, x_n \in X$  of distinct points in  $X$  such that  $x_i$  is a specialization of  $x_{i+1}$  for  $i = 0, \dots, n-1$ .*

(ii) Let  $f : X' \rightarrow X$  be a spectral map of locally spectral spaces. Then

$$\dim f = \sup_{x \in X} \dim f^{-1}(x) \in \mathbb{Z}_{\geq 0} \cup \{-\infty, \infty\} .$$

**Definition 21.2.** Let  $K \subset K'$  be an extension of complete algebraically closed nonarchimedean fields. Then the topological transcendence degree  $\text{tr. c}(K'/K) \in \mathbb{Z}_{\geq 0} \cup \{\infty\}$  of  $K'$  over  $K$  is the minimum over all integers  $n$  such that there exists a dense subfield  $L \subset K'$  containing  $K$  with transcendence degree  $n$  over  $K$ .

The definition in [Hub96, Definition 1.8.2] is slightly different, but in fact they agree in the case of algebraically closed fields, as one easily checks using Krasner's lemma.

**Lemma 21.3.**

- (i) Let  $K \subset K' \subset K''$  be extensions of algebraically closed complete nonarchimedean fields. Then  $\text{tr. c}(K''/K) \leq \text{tr. c}(K''/K') + \text{tr. c}(K'/K)$ .
- (ii) Let  $K \subset K'$  and  $L \subset L'$  be extensions of algebraically closed complete nonarchimedean fields such that there is an embedding  $K' \hookrightarrow L'$  sending  $K$  into  $L$ , and such that the algebraic closure of  $L \cdot K'$  in  $L'$  is dense. Then  $\text{tr. c}(L'/L) \leq \text{tr. c}(K'/K)$ .

*Proof.* The first part is [Hub96, Remark 1.8.3 (i)]. More precisely, assume that  $\text{tr. c}(K'/K) = n$  and  $\text{tr. c}(K''/K') = m$ . Then there exist  $n$  elements  $x_1, \dots, x_n \in K'$  such that  $K'$  is the minimal complete and algebraically closed subfield of  $K'$  containing  $K$  and  $x_1, \dots, x_n$ , and there exist  $m$  elements  $x_{n+1}, \dots, x_{n+m} \in K''$  such that  $K''$  is the minimal complete and algebraically closed subfield of  $K''$  containing  $K'$  and  $x_{n+1}, \dots, x_{n+m}$ . Then the minimal complete and algebraically closed subfield of  $K''$  containing  $K$  and  $x_1, \dots, x_{n+m}$  is  $K''$ : Indeed, it contains  $K'$ , and then is all of  $K''$ . This shows that  $\text{tr. c}(K''/K) \leq n + m$ , as desired.

Similarly, one proves the second part, which is also [Hub96, Remark 1.8.3 (ii)].  $\square$

Unfortunately, we were not able to resolve the following question.

**Question 21.4.** Let  $K \subset K' \subset K''$  be extensions of algebraically closed complete nonarchimedean fields. Is it true that  $\text{tr. c}(K'/K) \leq \text{tr. c}(K''/K)$ ?

As we cannot answer this, we consider instead the variant  $\widetilde{\text{tr. c}}(K'/K)$ , which is defined as the minimum over  $\text{tr. c}(K''/K)$  over all complete algebraically closed extensions  $K''$  of  $K$ . Clearly,  $\widetilde{\text{tr. c}}(K'/K) \leq \text{tr. c}(K'/K)$ . Moreover, Lemma 21.3 holds true with  $\widetilde{\text{tr. c}}$  in place of  $\text{tr. c}$ .

**Definition 21.5.** Let  $f : X' \rightarrow X$  be a map of analytic adic spaces. Then

$$\dim. \text{trg } f \in \mathbb{Z}_{\geq 0} \cup \{-\infty, \infty\}$$

is the supremum over all  $x' \in X'$  of  $\widetilde{\text{tr. c}}(C(x')/C(x))$ , where  $x = f(x') \in X$ , and  $C(x')$  resp.  $C(x)$  denote completed algebraic closures of the completed residue fields.

**Lemma 21.6.** Let  $f : X' \rightarrow X$  be a map of analytic adic spaces. Then

$$\dim f \leq \dim. \text{trg } f .$$

*Proof.* As the rationalized value group does not change under passage to a completed algebraic closure (and only gets larger under passage to extensions), the proof of [Hub96, Lemma 1.8.5 (i)] applies to show that  $\dim f \leq \dim. \text{trg } f$ .  $\square$

The definition of  $\dim. \operatorname{trg}$  extends to maps of diamonds.

**Definition 21.7.** *Let  $f : Y' \rightarrow Y$  be a map of diamonds. For each  $y' \in |Y'|$  with image  $y \in |Y|$ , choose quasi-pro-étale maps  $\operatorname{Spa}(C(y), C(y)^+) \rightarrow Y$  with  $y$  in the image, and  $\operatorname{Spa}(C(y'), C(y')^+) \rightarrow \operatorname{Spa}(C(y), C(y)^+) \times_Y Y'$  with  $y'$  in the image. Then*

$$\dim. \operatorname{trg} f \in \mathbb{Z}_{\geq 0} \cup \{-\infty, \infty\}$$

*is defined as the supremum of  $\widetilde{\operatorname{tr}.c}(C(y')/C(y))$  over all  $y' \in |Y'|$ .*

*More generally, if  $f : Y' \rightarrow Y$  is a map of  $v$ -stacks that is representable in diamonds, then  $\dim. \operatorname{trg} f$  is the supremum of  $\dim. \operatorname{trg}(f \times_Y X)$  over all maps  $X \rightarrow Y$  from diamonds  $X$ .*

**Remark 21.8.** Note that a pullback will not increase  $\dim. \operatorname{trg}$  by Lemma 21.3 (ii), so the definition of  $\dim. \operatorname{trg} f$  for morphisms of  $v$ -sheaves agrees with the previous definition if  $Y$  and  $Y'$  are diamonds.

Note that if  $Y$  is a locally spatial diamond or if  $f : Y' \rightarrow Y$  is map of  $v$ -stacks that is representable in locally spatial diamonds, we also have a definition  $\dim Y$  and of  $\dim f$ , by Definition 21.1. In the second case, we take the supremum over all locally spatial diamonds  $X$  with a map  $X \rightarrow Y$  of  $\dim(f \times_Y X)$ . To evaluate this, it is enough to range over  $X$  of the form  $\operatorname{Spa}(C, C^+)$ .

Next, we want to prove a bound on the cohomological dimension of spatial diamonds. For this, we need to briefly discuss points of diamonds.

**Proposition 21.9.** *Let  $Y$  be a quasiseparated diamond such that  $|Y|$  consists of only one point. Then  $Y = \operatorname{Spa}(C, \mathcal{O}_C)/\underline{G}$  for some algebraically closed nonarchimedean field  $C$  and some profinite group  $G$  acting continuously and faithfully on  $C$ . Moreover, the pair  $(C, G)$  is unique up to (non-unique) isomorphism.*

*Proof.* As  $|Y|$  consists of only one point, we can find a quasi-pro-étale surjection  $\operatorname{Spa}(C, \mathcal{O}_C) \rightarrow Y$ , which is necessarily separated (as  $\operatorname{Spa}(C, \mathcal{O}_C)$  is separated). Then the fibre product  $R = \operatorname{Spa}(C, \mathcal{O}_C) \times_Y \operatorname{Spa}(C, \mathcal{O}_C)$  is quasicompact separated pro-étale over  $\operatorname{Spa}(C, \mathcal{O}_C)$ , and in particular affinoid pro-étale by Lemma 7.19. Thus,  $R = \operatorname{Spa}(C, \mathcal{O}_C) \times \underline{S}$  for some profinite set  $S$ . As  $R$  is an equivalence relation,  $S$  becomes a profinite group, which we denote by  $G$ . The equivalence relation structure  $R = \operatorname{Spa}(C, \mathcal{O}_C) \times \underline{G} \subset \operatorname{Spa}(C, \mathcal{O}_C) \times \operatorname{Spa}(C, \mathcal{O}_C)$  is then equivalent to a continuous and faithful action of  $G$  on  $C$ , and  $Y = \operatorname{Spa}(C, \mathcal{O}_C)/\underline{G}$ .

For uniqueness, note that if  $(C', G')$  is another such pair, then  $\operatorname{Spa}(C', \mathcal{O}_{C'}) \times_Y \operatorname{Spa}(C, \mathcal{O}_C)$  is affinoid pro-étale over both  $\operatorname{Spa}(C, \mathcal{O}_C)$  and  $\operatorname{Spa}(C', \mathcal{O}_{C'})$ ; the choice of a point will then give an isomorphism  $C \cong C'$  commuting with the maps to  $Y$ . In this case, the group  $G$  is also the same, as it is the group of automorphisms of the map  $\operatorname{Spa}(C, \mathcal{O}_C) \rightarrow Y$ .  $\square$

In particular, it follows that in the situation of Proposition 21.9, étale sheaves on  $Y$  are equivalent to continuous discrete  $G$ -modules, and étale cohomology is equivalent to continuous  $G$ -cohomology.

**Definition 21.10.** *Let  $Y$  be a quasiseparated diamond, and let  $\ell$  be a prime. For each maximal point  $y \in |Y|$ , let  $Y_y \subset Y$  be the corresponding subdiamond with  $|Y_y| = \{y\} \subset |Y|$ , and write  $Y_y = \operatorname{Spa}(C_y, \mathcal{O}_{C_y})/\underline{G}_y$ . Then the cohomological dimension of  $Y$  at  $y$  is*

$$\operatorname{cd}_\ell y = \operatorname{cd}_\ell G_y .$$

We get the following bound on the cohomological dimension of spatial diamonds, which is an analogue of [Hub96, Corollary 2.8.3].

**Proposition 21.11.** *Let  $Y$  be a spatial diamond, let  $\ell$  be a prime, and let  $\mathcal{F}$  be an  $\ell$ -power-torsion étale sheaf on  $Y$  such that  $\mathcal{F}|_U = 0$  for some open subset  $U \subset Y$ . Then*

$$H^i(Y, \mathcal{F}) = 0$$

for  $i > \dim(Y \setminus U) + \sup_y \text{cd}_\ell y$ , where  $y \in Y$  runs through maximal points of  $Y$ .

**Remark 21.12.** Even if we are only interested in the statement for  $U = \emptyset$ , the proof runs by an induction that involves the statement for general  $U$ . However, we will actually need the statement for general  $U$  below.

*Proof.* We argue by induction on  $\dim(Y \setminus U)$  (noting that if it is infinite, there is nothing to prove). Also, we may assume that  $\mathcal{F}$  is  $\ell$ -torsion.

We may assume that there is a constructible sheaf  $\mathcal{F}_0$  such that

$$\mathcal{F} = \mathcal{F}_0 / j_{U!} \mathcal{F}_0|_U ,$$

where  $j_U : U \rightarrow Y$  denotes the open inclusion. Filtering  $\mathcal{F}_0$ , we may assume that there is a constructible locally closed subset  $S \subset |Y|$  such that the restriction of  $\mathcal{F}_0$  to  $|Y| \setminus S$  is trivial (i.e., the stalk at all geometric points above  $|Y| \setminus S$  vanishes), and for any strictly totally disconnected  $f : X \rightarrow Y$ , the pullback  $f^* \mathcal{F}_0|_{f^{-1}(S)}$  is the constant sheaf associated with some finite abelian group  $M$  with  $\ell M = 0$ . In particular,  $S = V \cap Z_0$  for a quasicompact open subset  $V \subset |Y|$  and a constructible closed subset  $Z_0 \subset |Y|$ . In this case,

$$\mathcal{F} = \mathcal{F}_0 / j_{U!} \mathcal{F}_0|_U$$

is concentrated (and constant after pullback to a strictly totally disconnected space) on the locally closed subset  $S \setminus U = V \cap Z$ , where  $Z = Z_0 \cap (Y \setminus U)$ .

We also denote by  $V \subset Y$  the corresponding open subdiamond, and let  $j : V \hookrightarrow Y$  be the quasicompact open immersion. Then  $\mathcal{F} = j_! \mathcal{F}$  for some sheaf  $\mathcal{F}_V$  on  $V$ . Let  $\tilde{\mathcal{F}} = j_* \mathcal{F}$ . Note that  $R^i j_* \mathcal{F}_V = 0$  for  $i > 0$  by Lemma 21.13 below. Thus,  $H^i(Y, \tilde{\mathcal{F}}) = H^i(V, \mathcal{F}_V)$ . Moreover, we have an injection  $\mathcal{F} \hookrightarrow \tilde{\mathcal{F}}$ ; let  $\mathcal{G}$  be its cokernel. Then  $\mathcal{G}$  is concentrated on  $\overline{V \setminus U} \setminus V$ . Note that  $\dim(\overline{V \setminus U} \setminus V) < \dim(Y \setminus U)$ : Indeed,  $\overline{V \setminus U} \setminus V \subset Y \setminus U$ , and any chain of specializations inside  $\overline{V \setminus U} \setminus V$  can be prolonged by including a point of  $V \setminus U$  as a proper generalization.

By induction, it follows that  $H^i(Y, \mathcal{G}) = 0$  for

$$i > \dim(\overline{V \setminus U} \setminus V) + \sup_y \text{cd}_\ell y ,$$

and in particular for  $i > \dim(Y \setminus U) + \sup_y \text{cd}_\ell y - 1$ . Thus, the long exact sequence

$$\dots \rightarrow H^{i-1}(Y, \mathcal{G}) \rightarrow H^i(Y, \mathcal{F}) \rightarrow H^i(Y, \tilde{\mathcal{F}}) \rightarrow \dots$$

reduces us to proving the desired vanishing for  $\tilde{\mathcal{F}}$ . As  $H^i(Y, \tilde{\mathcal{F}}) = H^i(V, \mathcal{F}_V)$ , we can replace  $Y$  by  $V$  (and  $\mathcal{F}$  by  $\mathcal{F}_V$ ) and assume that there is a closed subset  $Z \subset |Y|$  such that  $\mathcal{F}$  is concentrated on  $Z$ , and for any strictly totally disconnected perfectoid space  $f : X \rightarrow Y$ , the pullback  $f^* \mathcal{F}|_{f^{-1}(Z)}$  is the constant sheaf associated with some finite abelian group  $M$  with  $\ell M = 0$ .

Now we use the Leray spectral sequence for  $g : Y_{\text{ét}} \rightarrow |Y|$ . Note that  $Rg_* \mathcal{F}|_{|Y| \setminus Z} = 0$ , so the derived pushforward is concentrated on  $Z$ , which is a spectral space contained in  $Y \setminus U$ , and thus is of dimension  $\leq \dim(Y \setminus U)$ . By [Sch92, Corollary 4.6], the cohomological dimension of  $Z$  is bounded by  $\dim(Y \setminus U)$ .

Thus, it suffices to prove that  $R^i g_* \mathcal{F} = 0$  for  $i > \sup_y \text{cd}_\ell y$ . This can be checked on stalks, so we can assume that  $|Y|$  is local. In that case,  $|Y|$  is a totally ordered chain of specializations, and  $Z \subset |Y|$  is a closed subset of finite dimension. It follows that  $Z$  is a finite set of points totally ordered under specialization; let  $\eta \in Z$  be the generic point. Then the generalizations of  $\eta$  in  $|Y|$  form a quasicompact open subspace corresponding to a quasicompact open subdiamond  $j : V \hookrightarrow Y$ . Now the adjunction map  $\mathcal{F} \rightarrow j_* j^* \mathcal{F}$  is an isomorphism (as can be checked after pullback to a strictly totally disconnected cover) by our assumption on  $\mathcal{F}|_Z$ . Using Lemma 21.13 again, we can assume replace  $Y$  by  $V$ , and so assume that  $Z$  is the closed point of  $|Y|$ . Finally, we are reduced to Proposition 21.15 below.  $\square$

**Lemma 21.13.** *Let  $j : U \rightarrow Y$  be a quasicompact injection of locally spatial diamonds. Then for any sheaf of abelian groups  $\mathcal{F}$  on  $U_{\text{ét}}$ , one has  $R^i j_{\text{ét}*} \mathcal{F} = 0$  for  $i > 0$ .*

**Remark 21.14.** In fact, this holds true more generally for quasicompact separated quasi-pro-étale morphisms  $j : U \rightarrow X$ . The proof reduces to the case where  $X$  is strictly totally disconnected, in which case  $U$  is also strictly totally disconnected by Lemma 7.19. Then it follows from vanishing of étale cohomology on strictly totally disconnected spaces.

*Proof.* By Corollary 16.10, and as the result can be checked on stalks, we can assume that  $Y = \text{Spa}(C, C^+)$  for an algebraically closed nonarchimedean field  $C$  with an open and bounded valuation subring  $C^+ \subset C$ , and we need to check that  $H^i(U_{\text{ét}}, \mathcal{F}) = 0$  for  $i > 0$ . But  $U_{\text{ét}}$  and  $|U|$  define equivalent topoi, and  $|U|$  has a unique closed point, so  $H^i(U_{\text{ét}}, \mathcal{F}) = H^i(|U|, \mathcal{F}) = 0$  for  $i > 0$ , as desired.  $\square$

**Proposition 21.15.** *Let  $Y$  be a spatial diamond such that  $|Y|$  is local with closed point  $s \in |Y|$  and complement  $U = Y \setminus \{s\} \subset Y$ . Then for any  $\ell$ -torsion étale sheaf  $\mathcal{F}$  on  $Y_{\text{ét}}$  such that  $\mathcal{F}|_U = 0$ ,*

$$H^i(Y_{\text{ét}}, \mathcal{F}) = 0$$

*for any  $i > \text{cd}_\ell \eta$ , where  $\eta \in |Y|$  is the unique generic point.*

*Proof.* By assumption, there is a surjective quasi-pro-étale map  $f : \text{Spa}(C, C^+) \rightarrow Y$ , where  $C$  is an algebraically closed nonarchimedean field, and  $C^+ \subset C$  is an open and bounded valuation subring.

As  $\text{Spa}(C, C^+)$  is separated, the map  $f$  is separated, and thus  $R = \text{Spa}(C, C^+) \times_Y \text{Spa}(C, C^+)$  is quasicompact separated and pro-étale over  $\text{Spa}(C, C^+)$ , i.e.  $R$  is affinoid pro-étale over  $\text{Spa}(C, C^+)$  by Lemma 7.19.

Note that the map  $|\text{Spa}(C, C^+)| \rightarrow |Y|$  is bijective. For any point  $y \in |Y|$ , let  $\tilde{y} \in |\text{Spa}(C, C^+)|$  be the unique lift, which corresponds to a valuation subring  $C_y^+ \subset \mathcal{O}_C$  containing  $C^+$ . Then the fiber of  $|R|_{\tilde{y}}$  of  $|R|$  over  $\tilde{y}$  identifies with the sections of  $f$  over  $\text{Spa}(C, C_y^+)$ . It is a profinite set, which in fact is a profinite group  $G_y$  by the equivalence relation structure. If  $y'$  is a generalization of  $y$ , there is a natural generalization map  $|R|_{\tilde{y}} \rightarrow |R|_{\tilde{y}'}$  as all maps are uniquely generalizing, so one gets an inclusion of profinite groups  $G_y \subset G_{y'}$ . In particular,  $G_s$  is a closed subgroup of  $G_\eta$ , and so  $\text{cd}_\ell G_s \leq \text{cd}_\ell G_\eta = \text{cd}_\ell \eta$ .

It remains to see that the sheaf  $\mathcal{F}$  can be identified with a discrete continuous  $G_s$ -module, and its cohomology with continuous  $G_s$ -cohomology. But giving  $\mathcal{F}$  is equivalent to giving a sheaf on  $|\text{Spa}(C, C^+)|$  concentrated at the closed point, i.e. an abelian group, and the descent data over  $R$  amount to a continuous  $G_s$ -action. Computing cohomology via the Cartan–Leray spectral sequence for the covering  $g : \text{Spa}(C, C^+) \rightarrow Y$  gives the complex of continuous cochains, as desired.  $\square$

Moreover, we need the following result bounding the cohomological dimension of points.

**Proposition 21.16.** *Let  $f : Y \rightarrow \mathrm{Spa}(C, C^+)$  be a map of locally spatial diamonds, and let  $y \in |Y|$  be a maximal point. Then, for all  $\ell \neq p$ , one has*

$$\mathrm{cd}_\ell y \leq \dim. \mathrm{trg} f .$$

*Proof.* We may replace  $Y$  by  $Y_y$  and  $\mathrm{Spa}(C, C^+)$  by  $\mathrm{Spa}(C, \mathcal{O}_C)$ . Then  $Y = \mathrm{Spa}(C', \mathcal{O}_{C'})/\underline{G}$  for a profinite group  $G$  acting continuously and faithfully on  $C'$ , fixing  $C$  pointwise. In particular,  $G$  acts continuously on the quotient

$$C'^{\times}/(1 + C'^{\circ\circ})$$

of  $C'^{\times}$ . Let  $P \subset G$  be the normal closed subgroup which acts trivially; this is an analogue of the wild inertia subgroup. First, we note that  $P$  is a pro- $p$ -group. This follows from Lemma 21.17 below. Thus,  $\mathrm{cd}_\ell G = \mathrm{cd}_\ell G/P$  for  $\ell \neq p$ .

Let  $k'/k$  be the extension of residue fields of  $C'/C$ , and let  $\Gamma \subset \Gamma' \subset \mathbb{R}_{>0}$  be the value groups. Then the group  $C'^{\times}/(1 + C''^{\times})$  is an extension

$$1 \rightarrow k'^{\times} \rightarrow C'^{\times}/(1 + C''^{\times}) \rightarrow \Gamma' \rightarrow 1 .$$

As  $\Gamma' \subset \mathbb{R}_{>0}$ , the action of  $G/P$  on  $\Gamma'$  is trivial. Considering the action of  $G/P$  on  $k'$ , let  $I$  be the pointwise stabilizer of  $k'$ ; this is an analogue of the inertia group. Then  $\overline{G} = G/I$  acts faithfully and continuously on the discrete field  $k'$ , fixing  $k$  pointwise. It follows that  $k'_0 = k'^{\overline{G}}$  is a perfect field with algebraic closure  $k'$ , and  $\mathrm{Gal}(k'/k'_0) = \overline{G}$ . Thus,

$$\mathrm{cd}_\ell \overline{G} \leq \mathrm{tr. deg}(k'_0/k) = \mathrm{tr. deg}(k'/k) .$$

Finally, we have to understand the ‘‘tame inertia’’ group  $I/P$ . This acts through maps  $\Gamma' \rightarrow k^\times$  which are trivial on  $\Gamma$ , i.e. through maps  $\Gamma'/\Gamma \rightarrow k^\times$ . Moreover, as the elements are topologically nilpotent, the image of  $\Gamma'/\Gamma \rightarrow k^\times$  lands in the roots of unity; assuming that  $\Gamma'/\Gamma$  is finitely-dimensional over  $\mathbb{Q}$ , this gives an embedding

$$I/P \hookrightarrow \mathrm{Hom}(\Gamma'/\Gamma, \mu_\infty(k)) \cong (\mathbb{A}_f^p)^{\dim_{\mathbb{Q}} \Gamma'/\Gamma} .$$

This implies that

$$\mathrm{cd}_\ell I/P \leq \dim_{\mathbb{Q}} \Gamma'/\Gamma .$$

Therefore we get the desired inequality

$$\mathrm{cd}_\ell G = \mathrm{cd}_\ell G/P \leq \mathrm{cd}_\ell \overline{G} + \mathrm{cd}_\ell I/P \leq \mathrm{tr. deg}(k'/k) + \dim_{\mathbb{Q}} \Gamma'/\Gamma \leq \widetilde{\mathrm{tr. c}}(C'/C) ,$$

using [Bou98, VI.10.3 Corollary 1] in the final inequality (which also shows that indeed  $\Gamma'/\Gamma$  is finite-dimensional in case  $\mathrm{tr. c}(C'/C) < \infty$ ); here, we use that rationalized value groups and transcendence degrees of residue fields do not change under completion, and only increase after passage to further extensions.  $\square$

**Lemma 21.17.** *Let  $C$  be an algebraically closed complete nonarchimedean field of characteristic  $p$ , and let  $\gamma$  be a continuous automorphism of  $C$  such that  $\gamma^{n!}$  converges (pointwise) to the identity for  $n \rightarrow \infty$ . Assume that  $\gamma$  acts trivially on  $C^\times/(1 + C^{\circ\circ})$ . Then  $\gamma^{p^n}$  converges (pointwise) to the identity for  $n \rightarrow \infty$ .*

*Proof.* Let  $G$  be the subgroup of the continuous automorphisms of  $C$  generated by  $\gamma$ ; then  $G$  is a cyclic profinite group, and we have to see that  $G$  is pro- $p$ . If  $G$  is not pro- $p$ , then take some prime  $\ell \neq p$  dividing the pro-order of  $G$ ; replacing  $G$  by a pro- $\ell$ -Sylow subgroup (and  $\gamma$  by a generator of it), we can assume that  $G$  is pro- $\ell$ . Then  $\gamma^{\ell^n}$  converges pointwise to the identity for  $n \rightarrow \infty$ , and we have to see that  $\gamma$  is the identity. For any  $x \in C^\times$ , the quotient  $\frac{\gamma(x)}{x} \in C^\times$  lies in  $1 + C^{\circ\circ}$  by assumption. If it is not equal to 1, pick some nonzero  $y \in C^{\circ\circ}$  such that

$$\frac{\frac{\gamma(x)}{x} - 1}{y} \in \mathcal{O}_C^\times.$$

Then the association

$$g \in G \mapsto \frac{\frac{g(x)}{x} - 1}{y} \in \mathcal{O}_C \rightarrow k = \mathcal{O}_C / C^{\circ\circ}$$

defines a continuous group homomorphism  $G \rightarrow k$ . As  $k$  is  $p$ -torsion and  $G$  is pro- $\ell$ , this implies that the map is 0, which contradicts the assumption on  $y$ . Thus, for all  $x \in C^\times$ , one has  $\gamma(x) = x$ , as desired.  $\square$

## 22. PROPER PUSHFORWARD

In this section, we finally define the functor  $Rf_!$ .

**Convention 22.1.** In this section, we will always work with unbounded derived categories and therefore assume that  $f$  is representable in locally spatial diamonds with  $\dim.\mathrm{trg} f < \infty$ , which will guarantee the assumptions on bounded cohomological dimension in Theorem 19.2 and Proposition 17.6. Everything below works as well for  $D^+$  without the assumption  $\dim.\mathrm{trg} f < \infty$ . Moreover, we assume throughout that  $n\Lambda = 0$  for some  $n$  prime to  $p$ .

As usual, one needs to restrict attention to morphisms admitting a compactification.

**Definition 22.2.** A morphism  $f : Y' \rightarrow Y$  of  $v$ -stacks is compactifiable if it can be written as a composite of an open immersion and a partially proper morphism.

A key difference to the world of schemes is the presence of a canonical compactification.

**Proposition 22.3.** Let  $f : Y' \rightarrow Y$  and  $\tilde{Y} \rightarrow Y$  be morphisms of  $v$ -stacks, with pullback  $\tilde{f} : \tilde{Y}' = Y' \times_Y \tilde{Y} \rightarrow \tilde{Y}$ .

- (i) The morphism  $f$  is compactifiable if and only if it is separated and the natural map  $Y' \rightarrow \overline{Y'}^Y$  is an open immersion.
- (ii) If  $f$  is compactifiable, then  $\tilde{f}$  is compactifiable.
- (iii) If  $\tilde{f}$  is compactifiable and  $\tilde{Y} \rightarrow Y$  is a surjective map of  $v$ -stacks, then  $f$  is compactifiable.
- (iv) If  $Y_1 \rightarrow Y_2$  and  $Y_2 \rightarrow Y_3$  are compactifiable morphisms of  $v$ -stacks, then the composite  $Y_1 \rightarrow Y_3$  is compactifiable.
- (v) If  $f$  is separated and representable in locally spatial diamonds and  $Y'$  admits a cover by open subfunctors  $V \subset Y'$  such that  $f|_V : V \rightarrow Y$  is compactifiable, then  $f$  is compactifiable. Moreover, for any such open subfunctor  $V \subset Y'$ , the composite  $V \rightarrow Y' \rightarrow \overline{Y'}^Y$  is an open immersion.
- (vi) If  $f$  is separated and étale, then  $f$  is compactifiable.



(vii) *If  $f$  is separated and representable in locally spatial diamonds and there is a separated surjective map  $g : Z \rightarrow Y'$  of  $v$ -stacks such that  $f \circ g$  is compactifiable, then  $f$  is compactifiable.*

(viii) *If  $g : Y \rightarrow Z$  is a separated map of  $v$ -stacks such that  $g \circ f$  is compactifiable, then  $f$  is compactifiable.*

In particular, the property of being compactifiable is  $v$ -local on the target by (iii), and for separated maps which are representable in locally spatial diamonds, it is also “compactifiable-local” on the source by (iv) and (vii). (Under the hypothesis of (vii), it is necessarily the case that  $g$  is compactifiable, by (viii).)

*Proof.* For part (i), one direction is clear as  $\overline{Y'}^/Y$  is partially proper. For the converse, assume that  $f$  is compactifiable, and let  $Y' \hookrightarrow Z$  be an open immersion into a partially proper  $Z \rightarrow Y$ . As open immersions and partially proper morphisms are separated, we see that  $f$  is separated. Moreover, we get a map  $\overline{Y'}^/Y \rightarrow Z$ , which is still an injection. Thus  $Y' \rightarrow \overline{Y'}^/Y$  is a pullback of the open immersion  $Y' \rightarrow Z$ , and so an open immersion itself.

Now parts (ii) and (iii) follow easily from (i). In Part (iv), we may now assume that  $Y_3$ , and thus all  $Y_i$ , are separated. Then  $Y_2 \rightarrow \overline{Y_2}^/Y_3$  and  $Y_1 \rightarrow \overline{Y_1}^/Y_2$  are open immersions, thus so is

$$Y_1 \rightarrow \overline{Y_1}^/Y_2 = \overline{Y_1} \times_{\overline{Y_2}} Y_2 \rightarrow \overline{Y_1} \times_{\overline{Y_2}} \overline{Y_2}^/Y_3 = \overline{Y_1} \times_{\overline{Y_3}} Y_3 = \overline{Y_1}^/Y_3 ,$$

as desired.

In part (v), it is enough to prove that for any open subspace  $V \subset Y'$ , the composite  $V \rightarrow \overline{Y'}^/Y$  is an open immersion. For this, we can assume  $Y = \overline{Y'}^/Y$ , and we can work  $v$ -locally on  $Y$ , so we can assume  $Y = \text{Spa}(A, A^+)$  is strictly totally disconnected. In that case, by Proposition 10.5,  $Y'$  is a filtered union of open subspaces (of  $Y'$ ; not a priori of  $Y$ !) of the form  $\text{Spa}(A, (A^+)')$  for certain varying rings of integral elements  $(A^+) \subset A$  containing  $A^+$ . One can assume that the subspaces  $V$  are quasicompact, in which case  $V \subset \text{Spa}(A, (A^+)') \subset Y'$  for some such  $(A^+)'$ , and for any  $x \in V$  there are  $f_1, \dots, f_n, g \in A$  generating the unit ideal such that the rational subset  $U = \{|f_i| \leq |g|\} \subset Y$  satisfies  $x \in \text{Spa}(A, (A^+)') \cap U \subset V$ . But then  $U \subset \overline{V} = \overline{V}^/Y$ . As  $V \subset \overline{V}^/Y$  is an open immersion by assumption, this implies that  $U \cap V \subset U \cap \overline{V} = U$  is an open immersion. In other words,  $x \in U \cap V \subset U \subset Y$  is an open neighborhood of  $x$  in  $Y$  which is contained in  $Y'$ . These cover  $Y'$ , so that  $Y'$  is open in  $Y$ , as desired.

For part (vi), we may assume (by (iii)) that  $Y = X$  is a strictly totally disconnected perfectoid space. Then  $Y' = X'$  is a perfectoid space which is separated and étale over  $X$ . By (v), we may assume that  $Y'$  is quasicompact. In that case,  $Y'$  decomposes as a disjoint union of quasicompact open subspaces of  $Y$ , so the result is clear.

For part (vii), we can assume that  $Y = \overline{Y'}^/Y$ , and we need to see that  $f : Y' \hookrightarrow Y$  is an open immersion. This can be checked  $v$ -locally on  $Y$ , so we can assume that  $Y = \text{Spa}(A, A^+)$  is a strictly totally disconnected space, so that  $Y'$  is a locally spatial diamond. By part (vi), we can assume that  $Y'$  is quasicompact. Then by Proposition 10.5, one has  $Y' = \text{Spa}(A, (A^+)')$  for some ring of integral elements  $(A^+) \subset A$  containing  $A^+$ . Moreover, after a  $v$ -cover  $Z' = \text{Spa}(B, B^+) \rightarrow Y'$ , there is a map  $Z' \rightarrow Z$ , and there is an open subspace  $W = Z \times_{\overline{Z}^/Y} \overline{Z'}^/Y \subset \overline{Z'}^/Y$  containing  $Z'$  that maps into  $Y' \subset Y$ . For any finite collection  $\{f_i\}$  of elements of  $(A^+)'$ , consider the subspace  $\{|f_i| \leq 1\} \subset Y$ , which contains  $Y'$ . Their cofiltered intersection is  $Y'$ , and the preimage of this cofiltered intersection in  $\overline{Z'}^/Y$  is contained in  $W$ . By quasicompactness of  $\overline{Z'}^/Y \setminus W$  for the constructible topology, some such

subset  $\{|f_i| \leq 1\} \subset Y$  has the property that the preimage in  $\overline{Z}'/Y$  is contained in  $W$ . But the image of  $W$  is contained in  $Y'$  (and the map  $|\overline{Z}'/Y| \rightarrow |Y|$  surjective), so it follows that  $\{|f_i| \leq 1\} \subset Y'$ . As the reverse inclusion was also true, we see that  $Y' = \{|f_i| \leq 1\} \subset Y$  is open.

In part (viii), we have injections

$$Y' \hookrightarrow \overline{Y}'/Y \hookrightarrow \overline{Y}'/Z .$$

If the composite is an open immersion, then so is the first map (as it is a pullback of the composite).  $\square$

Now if  $f : Y' \rightarrow Y$  is a quasicompact compactifiable map of small  $v$ -stacks, we can write  $f$  as a composite of an open immersion  $j : Y' \hookrightarrow \overline{Y}'/Y$  and the proper map  $\overline{f}'/Y : \overline{Y}'/Y \rightarrow Y$ . In that case, we will momentarily define  $Rf_!$  as the composite  $R\overline{f}'_* \circ j_!$ .

However, we will be interested in the case where  $f$  is not necessarily quasicompact. In that case  $\overline{f}'/Y : \overline{Y}'/Y \rightarrow Y$  is only partially proper, and one should restrict to sections with proper support. In a fully derived setting, this is somewhat nontrivial to do. But we are already restricting to maps which are representable in locally spatial diamonds, so we can get around this difficulty. Namely, if  $Y$  and  $Y'$  are locally spatial diamonds, and say that  $Y'$  is even spatial, then one can write  $Y'$  as an increasing union of quasicompact open subspaces  $V \subset Y'$ , and  $Rf_!$  should be defined as the filtered colimit of  $R(f|_V)_!$ , which are defined as before. One can then descend this construction to the case where  $f$  is merely representable in locally spatial diamonds, as  $Rf_!$  satisfies base change.

We will now execute this line of thoughts, one step at a time.

**Definition 22.4.** *Let  $f : Y' \rightarrow Y$  be a compactifiable map of small  $v$ -stacks which is representable in spatial diamonds with  $\dim. \operatorname{trg} f < \infty$ , and assume  $n\Lambda = 0$  for some  $n$  prime to  $p$ . Write  $f$  as the composite of the open immersion  $j : Y' \hookrightarrow \overline{Y}'/Y$  and the proper map  $\overline{f}'/Y : \overline{Y}'/Y \rightarrow Y$ . Then we define*

$$Rf_! = R\overline{f}'_* \circ j_! : D_{\text{ét}}(Y', \Lambda) \rightarrow D_{\text{ét}}(Y, \Lambda) .$$

Our first aim is to prove that this definition is well-behaved on unbounded derived categories. For this reason, we need to know that  $R\overline{f}'_*$  has bounded cohomological dimension.

**Theorem 22.5.** *Let  $f : Y' \rightarrow Y$  be a compactifiable map of small  $v$ -stacks which is representable in spatial diamonds with  $\dim. \operatorname{trg} f < \infty$ , and assume  $n\Lambda = 0$  for some  $n$  prime to  $p$ . Write  $f$  as the composite of the open immersion  $j : Y' \hookrightarrow \overline{Y}'/Y$  and the proper map  $\overline{f}'/Y : \overline{Y}'/Y \rightarrow Y$ . Then  $R\overline{f}'_*$  has bounded cohomological dimension; more precisely, for all  $A \in D_{\text{ét}}(\overline{Y}'/Y, \Lambda)$  concentrated in degree 0, one has*

$$R^i \overline{f}'_* A = 0$$

for  $i > 3 \dim. \operatorname{trg} f$ .

The constant 3 may well be an artifact of the proof; if  $\overline{f}'/Y$  is still representable in spatial diamonds, it can be replaced by 2, as expected.

*Proof.* By Proposition 17.6, we can assume that  $Y$  is a strictly totally disconnected space. In fact, it is enough to check a statement about stalks, so we can reduce to the case  $Y = \operatorname{Spa}(C, C^+)$ , where  $C$  is an algebraically closed nonarchimedean field and  $C^+ \subset C$  an open and bounded valuation

subring, and we only need to check the statement on global sections. In that case, we need to prove that

$$H^i(\overline{Y'}^/Y, A) = 0$$

for  $i \geq 3 \dim. \operatorname{trg} f$ . Let  $s \in Y$  be the closed point and  $U = X \setminus \{s\}$ ,  $V = f^{-1}(U) \subset \overline{Y'}^/Y$ , with inclusion  $j : V \hookrightarrow \overline{Y'}^/Y$ . By Theorem 19.2, we know that

$$R\Gamma(\overline{Y'}^/Y, j_! j^* A) = 0.$$

Thus, replacing  $A$  by the cone of  $j_! j^* A \rightarrow A$ , we can assume that  $j^* A = 0$ .

Now, if  $\overline{Y'}^/Y$  is a spatial diamond, note that  $\dim(\overline{Y'}^/Y \setminus V) = \dim(\overline{f}^/Y)^{-1}(s) \leq \dim \overline{f}^/Y \leq \dim. \operatorname{trg} \overline{f}^/Y = \dim. \operatorname{trg} f$  and  $\operatorname{cd}_\ell y' \leq \dim. \operatorname{trg} f$  for all maximal points  $y' \in \overline{Y'}^/Y$  (which all lie in  $Y'$ ) and  $\ell \neq p$  by Proposition 21.16, so the result follows from Proposition 21.11.

Unfortunately, for a compactifiable map  $f : Y' \rightarrow Y$  of spatial diamonds, we do not know whether the canonical compactification  $\overline{Y'}^/Y$  is spatial. However, by Proposition 13.12 and Proposition 13.9, there is a compact Hausdorff space  $T$  (given as the maximal Hausdorff quotient of  $|\overline{Y'}^/Y|$ , which is also the maximal Hausdorff quotient of the spectral space  $|Y' \times_{\operatorname{Spa}(C, C^+)} \operatorname{Spa}(C, \mathcal{O}_C)|$ , as they share the same maximal points) and a map  $\overline{Y'}^/Y \rightarrow \underline{T}$  which is representable in locally spatial diamonds. In particular, one gets a map

$$g : \overline{Y'}^/Y \rightarrow \underline{T} \times Y$$

of proper diamonds over  $Y$ , which is thus necessarily proper; moreover, it is representable in spatial diamonds. The previous arguments imply that

$$R^i g_* A = 0$$

for  $i > 2 \dim. \operatorname{trg} g = 2 \dim. \operatorname{trg} f$ . Let  $h : \underline{T} \times Y \rightarrow Y$  be the projection. It remains to see that for any  $B \in D_{\text{ét}}(\underline{T} \times Y, \Lambda)$  concentrated in degree 0 and with trivial restriction to  $\underline{T} \times U$ , we have  $H^i(\underline{T} \times Y, B) = 0$  for  $i > \dim. \operatorname{trg} f$ .

In fact, we claim that  $H^i(\underline{T} \times Y, B) = 0$  for  $i > \dim f$ . This follows from Proposition 22.7 below, noting that if  $B$  is trivial on  $\underline{T} \times U$ , one gets

$$H^i(\underline{T} \times Y, B) = H^i(T, B|_{T \times \{s\}}),$$

and the observation that  $H^i(T, -) = H^i(|Y' \times_{\operatorname{Spa}(C, C^+)} \operatorname{Spa}(C, \mathcal{O}_C)|, -)$ , the latter of which is a spectral space of dimension  $\leq \dim f$ , so we conclude using [Sch92, Corollary 4.6].  $\square$

In the final step, we used a result on proper diamonds over strictly totally disconnected spaces of  $\dim. \operatorname{trg} = 0$ .

**Proposition 22.6.** *Let  $X$  be a strictly totally disconnected perfectoid space. Then the category of proper diamonds  $f : Y \rightarrow X$  with  $\dim. \operatorname{trg} f = 0$  is equivalent to the category of compact Hausdorff spaces over  $\pi_0 X$ , via sending a compact Hausdorff space  $T \rightarrow \pi_0 X$  to  $X \times_{\pi_0 X} \underline{T}$ .*

As a special case, if  $X = \operatorname{Spa}(C, C^+)$ , then the category of proper diamonds  $f : Y \rightarrow X$  of  $\dim. \operatorname{trg} f = 0$  is equivalent to the category of compact Hausdorff spaces. This result was suggested to the author by M. Rapoport.

*Proof.* Let  $f : Y \rightarrow X$  be any proper map of diamonds with  $\dim. \operatorname{trg} f = 0$ . Let  $\tilde{X} \rightarrow Y$  be a quasi-pro-étale surjection from a strictly totally disconnected space. Then  $\tilde{X}^{\overline{}/X} \rightarrow X$  is a proper map of strictly totally disconnected perfectoid spaces which induces an isomorphism on completed residue fields (by the assumption  $\dim. \operatorname{trg} f = 0$ ); this implies that  $\tilde{X}^{\overline{}/X} = X \times_{\pi_0 X} \underline{S}$  for some profinite set  $S \rightarrow \pi_0 X$ . Write  $X_S = X \times_{\pi_0 X} \underline{S}$ . We get a surjective map  $X_S \rightarrow \tilde{Y}$  over  $X$ . The equivalence relation  $R = X_S \times_Y X_S$  is again proper and pro-étale over  $X$ , so  $R = X \times_{\pi_0 X} \underline{S}'$  for some profinite  $S' \subset S \times S$ . Moreover, the maps  $s, t : R \rightarrow X_S$  over  $X$  are given by maps  $S' \rightarrow S$ . Now  $T = S/S'$  is a compact Hausdorff space, and  $Y = X \times_{\pi_0 X} \underline{T}$ . Using Example 11.12, we see that this gives the desired equivalence of categories.  $\square$

In this situation, we also need a characterization of the category  $D_{\text{ét}}^+(Y, \Lambda)$ .

**Proposition 22.7.** *Let  $X$  be a strictly totally disconnected perfectoid space, and let  $f : Y \rightarrow X$  be a proper map of diamonds of  $\dim. \operatorname{trg} f = 0$ , so that by Proposition 22.6, one has  $Y = X \times_{\pi_0 X} \underline{T}$  for some compact Hausdorff space  $T \rightarrow \pi_0 X$ . Then pullback under  $Y \rightarrow |Y|$  defines an equivalence*

$$D^+(|Y|, \Lambda) \xrightarrow{\sim} D_{\text{ét}}^+(Y, \Lambda) ,$$

where  $D^+(|Y|, \Lambda)$  is the derived category of sheaves of  $\Lambda$ -modules on the topological space  $|Y|$ .

*Proof.* There is a natural map of topoi  $t : Y_v \rightarrow |Y|$ . It follows from the definitions that if  $\mathcal{F}$  is a sheaf of  $\Lambda$ -modules on  $|Y|$ , then  $t^* \mathcal{F}$  lies in  $D_{\text{ét}}^+(Y_v, \Lambda)$ . Thus, it is enough to prove the following assertions.

- (i) For any sheaf of abelian groups  $\mathcal{F}$  on  $|Y|$ , the adjunction map  $\mathcal{F} \rightarrow R t_* t^* \mathcal{F}$  is an isomorphism.
- (ii) If  $\mathcal{G}$  is a small sheaf of abelian groups on  $Y_v$  such that for some cover  $g : \tilde{Y} \rightarrow Y$  by some strictly totally disconnected perfectoid space  $\tilde{Y}$ , the restriction  $g^* \mathcal{G}$  lies in  $\tilde{Y}_{\text{ét}}^{\sim} \subset \tilde{Y}_v^{\sim}$ , then the adjunction map  $\mathcal{G} \rightarrow t^* R t_* \mathcal{G}$  is an isomorphism.

First, note that for any small v-sheaf  $Y$ , if  $C \in D_{\text{ét}}^+(Y_v, \Lambda)$ , then for any cofiltered inverse system of qcqs diamonds  $Z_i \rightarrow Y$  with inverse limit  $Z$ , one has

$$R\Gamma(Z, C) = \varinjlim_i R\Gamma(Z_i, C) .$$

This follows from Proposition 14.9 if all  $Z_i$  are spatial; in general, writing all  $Z_i$  compatibly as quotients of affinoid perfectoid spaces  $X_i$  as in the proof of Lemma 11.22, the equivalence relations  $R_i = X_i \times_{Z_i} X_i$  are spatial diamonds by Proposition 12.3. The result then follows by descent from the case of the limits of the  $X_i$ , the  $R_i$ , the  $R_i \times_{X_i} R_i$ , etc. .

To check claim (i) above, we have to check an equality of stalks. But note that if  $y \in |Y|$ , then the open neighborhoods of  $y$  are cofinal with the neighborhoods of the form  $U \times_{\pi_0 X} \underline{S}$ , where  $U \subset X$  is a quasicompact open neighborhood of the image of  $y$  in  $X$ , and  $S \subset T$  is the closure of an open neighborhood of the image of  $y$  in  $T$ ; indeed, this follows from the similar observation for compact Hausdorff spaces. Note that such  $U \times_{\pi_0 X} \underline{S}$  are qcqs diamonds which form a cofiltered inverse system with inverse limit the set of generalizations of  $Y_y$  in  $|Y|$ , which is of the form  $\operatorname{Spa}(C(y), C(y)^+)$ , where  $C(y)$  is algebraically closed, and  $C(y)^+ \subset C(y)$  an open and bounded valuation subring. Thus, the stalk of  $R t_* t^* \mathcal{F}$  can also be computed as a direct limit of the cohomologies of  $t^* \mathcal{F}$  over these neighborhoods  $U \times_{\pi_0 X} \underline{S}$ , which by the remark above reduces to

$$R\Gamma(Y_y, t^* \mathcal{F}) .$$

But this is given by the stalk of  $\mathcal{F}$  at  $y$ , as desired.

Now for claim (ii), we have to similarly check a claim on stalks, which again follows from the cofinality of open and qcqs neighborhoods.  $\square$

Now we come back to the analysis of the functor  $Rf_!$ .

**Proposition 22.8.** *Let  $f : Y' \rightarrow Y$  be a compactifiable map of small  $v$ -stacks which is representable in spatial diamonds with  $\dim. \operatorname{trg} f < \infty$ , and assume  $n\Lambda = 0$  for some  $n$  prime to  $p$ . Let  $g : \tilde{Y} \rightarrow Y$  be any map of small  $v$ -stacks, with base change  $\tilde{f} : \tilde{Y}' = Y' \times_Y \tilde{Y} \rightarrow \tilde{Y}$  and  $g' : \tilde{Y}' \rightarrow Y'$ .*

*There is a natural base change equivalence*

$$g^* Rf_! \simeq R\tilde{f}_! g'^*$$

of functors  $D_{\text{ét}}(Y', \Lambda) \rightarrow D_{\text{ét}}(\tilde{Y}, \Lambda)$ .

*Proof.* This follows by combining Theorem 22.5, Proposition 17.6 and Proposition 19.1.  $\square$

**Proposition 22.9.** *Let  $g : Y'' \rightarrow Y'$  and  $f : Y' \rightarrow Y$  be compactifiable maps of small  $v$ -stacks which are representable in spatial diamonds with  $\dim. \operatorname{trg} f, \dim. \operatorname{trg} g < \infty$ , and assume  $n\Lambda = 0$  for some  $n$  prime to  $p$ . Then there is a natural equivalence*

$$Rf_! \circ Rg_! \simeq R(f \circ g)_!$$

of functors  $D_{\text{ét}}(Y'', \Lambda) \rightarrow D_{\text{ét}}(Y, \Lambda)$ .

*Proof.* Consider the following diagram

$$\begin{array}{ccccc} Y'' & \xrightarrow{j_1} & \overline{Y''}/Y' & \xrightarrow{j_2} & \overline{Y''}/Y \\ & \searrow g & \downarrow g_1 & & \downarrow g_2 \\ & & Y' & \xrightarrow{j_3} & \overline{Y'}/Y \\ & & \searrow f & & \downarrow f_1 \\ & & & & Y \end{array}$$

Then

$$Rf_! Rg_! = Rf_{1*} j_{3!} Rg_{1*} j_{1!}$$

and

$$R(f \circ g)_! = Rf_{1*} Rg_{2*} j_{2!} j_{1!}.$$

However, by Theorem 19.2 and Theorem 22.5 (applied to  $Y'' \rightarrow \overline{Y''}/Y$ ), we have  $j_{3!} Rg_{1*} = Rg_{2*} j_{2!}$ , as desired.  $\square$

In case of overlap, the current definition of  $Rf_!$  agrees with Definition 19.1.

**Proposition 22.10.** *Let  $f : Y' \rightarrow Y$  be a quasicompact separated étale map of small  $v$ -stacks, and assume  $n\Lambda = 0$  for some  $n$  prime to  $p$ . Then  $Rf_!$  as defined in Definition 19.1 agrees with  $Rf_!$  as defined in Definition 22.4.*

*Proof.* Let  $Rf_!^{\text{ét}}$  denote the one from Definition 19.1 temporarily, which is the left adjoint of  $f^*$ . There is a natural transformation

$$Rf_!^{\text{ét}} \rightarrow Rf_!$$

adjoint to the map  $\text{id} \rightarrow f^*Rf_!$  coming from the base change identity  $f^*Rf_! = R\pi_{2!}\pi_1^*$ , where  $\pi_1, \pi_2 : Y' \times_Y Y' \rightarrow Y'$  are the two projections, and the map

$$\text{id} = R\pi_{2!}R\Delta_!\Delta^*\pi_1^* \rightarrow R\pi_{2!}\pi_1^*,$$

using that  $\Delta : Y' \hookrightarrow Y' \times_Y Y'$  is an open and closed immersion.

To check that this is an equivalence, we use that both functors commute with any base change to reduce to the case that  $Y$  is strictly totally disconnected. In that case,  $Y'$  decomposes into a disjoint union of quasicompact open subspaces of  $Y$ , so we can reduce to that case, in which the claim is clear.  $\square$

We also get the projection formula.

**Proposition 22.11.** *Let  $f : Y' \rightarrow Y$  be a compactifiable map of small  $v$ -sheaves which is representable in spatial diamonds with  $\dim. \text{trg } f < \infty$ , and let  $\Lambda$  be a ring with  $n\Lambda = 0$  for some  $n$  prime to  $p$ . Then there is a functorial isomorphism*

$$Rf_!B \otimes_{\Lambda}^{\mathbb{L}} A \simeq Rf_!(B \otimes_{\Lambda}^{\mathbb{L}} f^*A)$$

in  $B \in D_{\text{ét}}(Y', \Lambda)$  and  $A \in D_{\text{ét}}(Y, \Lambda)$ .

*Proof.* If  $f = j$  is an open immersion, there is a natural map

$$j_!(B \otimes_{\Lambda}^{\mathbb{L}} j^*A) \rightarrow j_!B \otimes_{\Lambda}^{\mathbb{L}} A$$

adjoint to

$$B \otimes_{\Lambda}^{\mathbb{L}} \Lambda j^*A = j^*j_!B \otimes_{\Lambda}^{\mathbb{L}} A = j^*(j_!B \otimes_{\Lambda}^{\mathbb{L}} A).$$

To check whether this is an equivalence, we can assume that  $Y$  is a strictly totally disconnected space (as all operations commute with pullback). In that case  $D_{\text{ét}}(Y, \Lambda) = D_{\text{ét}}(|Y|, \Lambda)$ ; let  $i : Z \hookrightarrow |Y|$  be the closed complement. As the map is clearly an isomorphism away from  $Z$  and the left-hand side vanishes on  $Z$ , the statement follows from

$$i^*(j_!B \otimes_{\Lambda}^{\mathbb{L}} A) = i^*j_!B \otimes_{\Lambda}^{\mathbb{L}} i^*A = 0$$

as  $i^*j_!B = 0$ .

In general, we now get a map

$$Rf_!B \otimes_{\Lambda}^{\mathbb{L}} A = R\bar{f}_*^Y j_!B \otimes_{\Lambda}^{\mathbb{L}} A \rightarrow R\bar{f}_*^Y (j_!B \otimes_{\Lambda}^{\mathbb{L}} R\bar{f}^*A) \simeq R\bar{f}_*^Y j_!(B \otimes_{\Lambda}^{\mathbb{L}} j^*R\bar{f}_*^Y A) = Rf_!(B \otimes_{\Lambda}^{\mathbb{L}} f^*A),$$

where the middle map comes from a standard adjunction. To check whether this is an isomorphism, we may by Proposition 17.6 assume that  $Y$  is a strictly totally disconnected space. In that case, we have to prove a statement on stalks, so we can assume that  $Y = \text{Spa}(C, C^+)$  is connected, and it is enough to check the statement on global sections. Let  $s \in Y$  be the closed point,  $j_U : U = Y \setminus \{s\} \hookrightarrow Y$  the open immersion. If  $A = j_{U!}A_U$  for some  $A_U \in D_{\text{ét}}(U, \Lambda)$ , then

$$R\Gamma(Y, Rf_!B \otimes_{\Lambda}^{\mathbb{L}} j_{U!}A_U) = 0 = R\Gamma(\bar{Y}^{\prime/Y}, j_!(B \otimes_{\Lambda}^{\mathbb{L}} f^*j_{U!}A_U)) = R\Gamma(Y, Rf_!(B \otimes_{\Lambda}^{\mathbb{L}} f^*j_{U!}A_U)),$$

using Theorem 19.2 in the middle equality. Thus, we may replace  $A$  by the cone of  $j_{U!}j_U^*A \rightarrow A$ , and assume that  $A$  is a complex of  $\Lambda$ -modules concentrated on  $\{s\} \subset Y$ . On the other hand, we can also replace  $A$  by the constant complex of  $\Lambda$ -modules on  $Y$ . Repeating this argument in the other

direction, we may assume that  $A$  is a constant complex of  $\Lambda$ -modules. As such, it is a (derived) filtered colimit of perfect complexes of  $\Lambda$ -modules. The case of a perfect complex is clear, and so the general case follows from the next proposition.  $\square$

**Proposition 22.12.** *Let  $f : Y' \rightarrow Y$  be a compactifiable map of small  $v$ -sheaves which is representable in spatial diamonds with  $\dim. \operatorname{trg} f < \infty$ , and let  $\Lambda$  be a ring with  $n\Lambda = 0$  for some  $n$  prime to  $p$ . Then for any collection of objects  $A_i \in D_{\text{ét}}(Y', \Lambda)$ ,  $i \in I$  for some set  $I$ , the natural map*

$$\bigoplus_i Rf_! A_i \rightarrow Rf_! \left( \bigoplus_i A_i \right)$$

is an isomorphism in  $D_{\text{ét}}(Y, \Lambda)$ .

*Proof.* By Proposition 22.8, we can assume that  $Y = \operatorname{Spa}(C, C^+)$ , and it suffices to check on global sections. In any given degree  $d$ , the groups  $H^d(Y, \bigoplus_i Rf_! A_i)$  and  $H^d(Y, Rf_! (\bigoplus_i A_i))$  depend only on the truncations  $\tau_{\geq d-3} \dim. \operatorname{trg} f A_i$ ; thus, we may assume that all  $A_i$  are uniformly bounded below. We need to see that

$$R\Gamma(Y, \bigoplus_i R\bar{f}_*^{/Y} j_! A_i) = R\Gamma(Y, R\bar{f}_*^{/Y} j_! (\bigoplus_i A_i)) .$$

Here, the left-hand side is given by  $\bigoplus_i R\Gamma(Y, R\bar{f}_*^{/Y} j_! A_i) = \bigoplus_i R\Gamma(\bar{Y}'^{/Y}, j_! A_i)$  as  $Y$  is strictly local, and the right-hand side is given by  $R\Gamma(\bar{Y}'^{/Y}, \bigoplus_i j_! A_i)$ . In other words, we have to check that  $R\Gamma(\bar{Y}'^{/Y}, -)$  commutes with arbitrary direct sums in  $D_{\text{ét}}^{\geq -n}(\bar{Y}'^{/Y}, \Lambda)$ . But cohomology is given by  $v$ -cohomology, and the  $v$ -topos of a qcqs  $v$ -sheaf is coherent, so we get the result by SGA 4 VI Corollaire 5.2.  $\square$

This finishes the case that  $f$  is quasicompact. It remains to consider the non-quasicompact case, and obtain all previous results in this generality. The idea here is simple: If  $f : Y' \rightarrow Y$  is a compactifiable map of small  $v$ -stacks that is representable in locally spatial diamonds with  $\dim. \operatorname{trg} f < \infty$ , then, at least  $v$ -locally on  $Y$ , we can write  $Y'$  as an increasing union of quasicompact open subspaces, to which we can apply the preceding discussion; now we take the filtered direct limit. Unfortunately, it is somewhat tricky to resolve all homotopy coherence issues in this approach. For this reason, we make rather heavy use of Lurie's book, [Lur09], in the construction of  $Rf_!$ . Specifically, we will use the theory of left Kan extensions (along full embeddings), cf. [Lur09, Section 4.3.2].

First, we consider the case where  $Y$  is itself a locally spatial diamond.

**Definition 22.13.** *Let  $f : Y' \rightarrow Y$  be a compactifiable map of locally spatial diamonds with  $\dim. \operatorname{trg} f < \infty$ , and assume  $n\Lambda = 0$  for some  $n$  prime to  $p$ . Let  $j : Y' \hookrightarrow \bar{Y}'^{/Y}$  and  $\bar{f}^{/Y} : \bar{Y}'^{/Y} \rightarrow Y$  be the usual maps, where  $\bar{f}^{/Y}$  is partially proper. Let  $\mathcal{D}_{\text{ét}, \operatorname{prop}/Y}(Y', \Lambda) \subset \mathcal{D}_{\text{ét}}(Y', \Lambda)$  be the full  $\infty$ -subcategory of  $A \in \mathcal{D}_{\text{ét}}(Y', \Lambda)$  such that  $A \simeq j_{V!} j_V^* A$  for some open subspace  $j_V : V \hookrightarrow Y'$  that is quasicompact over  $Y$ .*

The functor

$$Rf_! : \mathcal{D}_{\text{ét}}(Y', \Lambda) \rightarrow \mathcal{D}_{\text{ét}}(Y, \Lambda)$$

is defined as the left Kan extension of

$$R\bar{f}_*^{/Y} j_! : \mathcal{D}_{\text{ét}, \operatorname{prop}/Y}(Y', \Lambda) \rightarrow \mathcal{D}_{\text{ét}}(Y, \Lambda)$$

along the full inclusion  $\mathcal{D}_{\text{ét,prop}/Y}(Y', \Lambda) \subset \mathcal{D}_{\text{ét}}(Y', \Lambda)$ .

In other words, on the full  $\infty$ -subcategory  $\mathcal{D}_{\text{ét,prop}/Y}(Y', \Lambda) \subset \mathcal{D}_{\text{ét}}(Y', \Lambda)$  of sheaves “with proper support over  $Y$ ”, the functor  $Rf_!$  is given by  $R\bar{f}_*^{/Y} j_!$ . For a general object  $A \in \mathcal{D}_{\text{ét}}(Y', \Lambda)$ , one can write  $A$  as a filtered colimit of objects  $A_V = j_{V!} j_V^* A \in \mathcal{D}_{\text{ét,prop}/Y}(Y', \Lambda)$ , and then by definition (cf. [Lur09, Definition 4.3.2.2])

$$Rf_! A = \varinjlim_V Rf_! A_V .$$

Note that here,  $Rf_! A_V = R(f|_V)_!(j_V^* A)$ , as

$$R\bar{f}_*^{/Y} j_! A_V = R\bar{f}_*^{/Y} (j \circ j_V)_!(j_V^* A) = R\bar{f}_*^{/Y} g_* j'_V!(j_V^* A) = R(f|_V)_!(j_V^* A) ,$$

where  $j'_V : V \hookrightarrow \bar{V}^{/Y}$  is the open immersion, and  $g : \bar{V}^{/Y} \hookrightarrow \bar{Y}^{/Y}$  the closed immersion.

Before going on with the definition of  $Rf_!$  in the general case, we need to establish base change in this situation. For this, it is helpful to first establish that  $Rf_!$  commutes with all colimits.

**Proposition 22.14.** *Let  $f : Y' \rightarrow Y$  be a compactifiable map of locally spatial diamonds with  $\dim. \text{trg } f < \infty$ , and assume  $n\Lambda = 0$  for some  $n$  prime to  $p$ . The functor*

$$Rf_! : \mathcal{D}_{\text{ét}}(Y', \Lambda) \rightarrow \mathcal{D}_{\text{ét}}(Y, \Lambda)$$

*commutes with all direct sums; equivalently (cf. [Lur16, Proposition 1.4.4.1 (2)]), with all colimits.*

*Proof.* Let  $A_i, i \in I$ , be any set of objects of  $\mathcal{D}_{\text{ét}}(Y', \Lambda)$ . Write each  $A_i$  as the colimit of  $A_{i,V} = j_{V!} j_V^* A_i$  over all open immersions  $j_V : V \hookrightarrow Y'$  which are quasicompact over  $Y$ . Then  $Rf_! A_{i,V} = R(f|_V)_!(j_V^* A_i)$  by construction, which commutes with arbitrary direct sums by Proposition 22.12. In general, we see that  $Rf_! \bigoplus_i A_i$  is the colimit of

$$R(f|_V)_! j_V^* \bigoplus_i A_i = \bigoplus_i R(f|_V)_! j_V^* A_i ,$$

which gives the desired result.  $\square$

**Proposition 22.15.** *Let  $f : Y' \rightarrow Y$  be a compactifiable map of locally spatial diamonds with  $\dim. \text{trg } f < \infty$ , and assume  $n\Lambda = 0$  for some  $n$  prime to  $p$ . Let  $g : \tilde{Y} \rightarrow Y$  be any map of locally spatial diamonds, with base change  $\tilde{f} : \tilde{Y}' = Y' \times_Y \tilde{Y} \rightarrow \tilde{Y}$  and  $g' : \tilde{Y}' \rightarrow Y'$ .*

*There is a natural base change equivalence*

$$g^* Rf_! \simeq R\tilde{f}_! g'^*$$

*of functors  $\mathcal{D}_{\text{ét}}(Y', \Lambda) \rightarrow \mathcal{D}_{\text{ét}}(\tilde{Y}, \Lambda)$ .*

*Proof.* To get a natural transformation from  $g^* Rf_!$  to  $R\tilde{f}_! g'^*$ , note that in the definition of  $Rf_!$  as the left Kan extension of  $R\bar{f}_*^{/Y} j_!$ , the functor  $j_!$  commutes with any base change, and there is a general base change adjunction for  $R\bar{f}_*^{/Y}$ . On  $\mathcal{D}_{\text{ét,prop}/Y}(Y', \Lambda)$ , the base change map is an equivalence by Proposition 22.8. In general, write any  $A \in \mathcal{D}_{\text{ét}}(Y', \Lambda)$  as the filtered colimit of  $A_V = j_{V!} j_V^* A$ . The functor  $g^* Rf_!$  commutes with this by definition; but so does  $R\tilde{f}_! g'^*$ , by Proposition 22.14.  $\square$

Let us now define  $Rf_!$  in general. Thus, let  $f : Y' \rightarrow Y$  be a compactifiable map of small  $v$ -stacks that is representable in locally spatial diamonds with  $\dim. \text{trg } f < \infty$ . In that case, we pick a simplicial  $v$ -hypercover  $Y_\bullet \rightarrow Y$  such that all  $Y_i$  are locally spatial diamonds, and let



$Y'_\bullet = Y' \times_Y Y_\bullet$ , which is a simplicial v-hypercover of  $Y'$ . The association  $i \mapsto \mathcal{D}_{\acute{e}t}(Y'_i, \Lambda)$  defines a functor  $\Delta \rightarrow \mathcal{C}at_\infty$  (with functors given by pullback); cf. [LZ14, Section 2] for a construction even as presentable symmetric monoidal  $\infty$ -categories for any diagram of ringed topoi (the extra condition  $\acute{e}t$  only amounts to passage to full  $\infty$ -subcategories, and poses no homotopy coherence issues). This is encoded in a coCartesian fibration  $\mathcal{D}_{\acute{e}t}(Y'_\bullet, \Lambda)^0 \rightarrow \Delta$ , whose  $\infty$ -category of sections is the  $\infty$ -derived category  $\mathcal{D}_{\acute{e}t}(Y'_\bullet, \Lambda)$  of the simplicial space  $Y'_\bullet$ .

Let

$$\mathcal{D}_{\acute{e}t, \text{prop}/Y_\bullet}(Y'_\bullet, \Lambda)^0 \subset \mathcal{D}_{\acute{e}t}(Y'_\bullet, \Lambda)^0$$

be the full  $\infty$ -subcategory whose fibre over any  $i \in \Delta$  is given by

$$\mathcal{D}_{\acute{e}t, \text{prop}/Y_i}(Y'_i, \Lambda) \subset \mathcal{D}_{\acute{e}t}(Y'_i, \Lambda) .$$

As pullbacks preserve this condition, this is still a coCartesian fibration over  $\Delta$ . Similarly, we have coCartesian fibrations  $\mathcal{D}_{\acute{e}t}(\overline{Y}'/Y_\bullet, \Lambda)^0$  and  $\mathcal{D}_{\acute{e}t}(Y_\bullet, \Lambda)^0$  over  $\Delta$ . The restriction functor

$$j_\bullet^* : \mathcal{D}_{\acute{e}t}(\overline{Y}'/Y_\bullet, \Lambda)^0 \rightarrow \mathcal{D}_{\acute{e}t}(Y'_\bullet, \Lambda)^0$$

has a fully faithful left adjoint  $j_{\bullet!}$ , which is in every fibre over  $i \in \Delta$  given by  $j_{i!}$ , where  $j_i : Y'_i \hookrightarrow \overline{Y}'/Y_i$  is the open immersion. Indeed, this follows from the fact that  $j_{i!}$  commutes with base change. Moreover, one has the pushforward functor

$$R\overline{f}_{\bullet*}^{Y_\bullet} : \mathcal{D}_{\acute{e}t}(\overline{Y}'/Y_\bullet, \Lambda)^0 \rightarrow \mathcal{D}_{\acute{e}t}(Y_\bullet, \Lambda)^0 ,$$

right adjoint to  $\overline{f}_{\bullet*}^{Y_\bullet, *}$ , which in every fibre over  $i \in \Delta$  is given by  $R\overline{f}_{i*}^{Y_i}$ . In particular, we get the functor

$$R\overline{f}_{\bullet*}^{Y_\bullet} j_{\bullet!} : \mathcal{D}_{\acute{e}t, \text{prop}/Y_\bullet}(Y'_\bullet, \Lambda)^0 \rightarrow \mathcal{D}_{\acute{e}t}(Y_\bullet, \Lambda)^0 .$$

We let

$$Rf_{\bullet!}^0 : \mathcal{D}_{\acute{e}t}(Y'_\bullet, \Lambda)^0 \rightarrow \mathcal{D}_{\acute{e}t}(Y_\bullet, \Lambda)^0$$

be its left Kan extension, which exists by [Lur09, Corollary 4.3.2.14].

**Lemma 22.16.** *The functor  $Rf_{\bullet!}^0$  is given by*

$$Rf_{i!} : \mathcal{D}_{\acute{e}t}(Y'_i, \Lambda) \rightarrow \mathcal{D}_{\acute{e}t}(Y_i, \Lambda)$$

in the fibre over  $i \in \Delta$ .

*Proof.* As  $\mathcal{D}_{\acute{e}t, \text{prop}/Y_\bullet}(Y'_\bullet, \Lambda)^0$  is itself a coCartesian fibration over  $\Delta$ , one sees that for all  $A \in \mathcal{D}_{\acute{e}t}(Y'_i, \Lambda)$ , the functor

$$\mathcal{D}_{\acute{e}t, \text{prop}/Y_i}(Y'_i, \Lambda)/_A \rightarrow \mathcal{D}_{\acute{e}t, \text{prop}/Y_\bullet}(Y'_\bullet, \Lambda)^0/_A$$

is cofinal. Moreover, the index category is filtered. Taken together, these imply that the relevant colimits agree (using [Lur09, Proposition 4.3.1.7] to see that one may pass to a cofinal subcategory).  $\square$

**Lemma 22.17.** *The functor*

$$Rf_{\bullet!}^0 : \mathcal{D}_{\acute{e}t}(Y'_\bullet, \Lambda)^0 \rightarrow \mathcal{D}_{\acute{e}t}(Y_\bullet, \Lambda)^0$$

sends coCartesian edges to coCartesian edges.

*Proof.* This follows from the previous lemma and Proposition 22.15.  $\square$

Passing to coCartesian sections over  $\Delta$ , we get a functor

$$Rf_! : \mathcal{D}_{\text{ét}}(Y', \Lambda) \simeq \mathcal{D}_{\text{ét, cart}}(Y'_\bullet, \Lambda) \rightarrow \mathcal{D}_{\text{ét, cart}}(Y_\bullet, \Lambda) \simeq \mathcal{D}_{\text{ét}}(Y, \Lambda),$$

as desired.

**Definition 22.18.** *Let  $f : Y' \rightarrow Y$  be a compactifiable map of small  $v$ -stacks that is representable in locally spatial diamonds with  $\dim. \text{trg } f < \infty$ , and assume  $n\Lambda = 0$  for some  $n$  prime to  $p$ . The functor*

$$Rf_! : D_{\text{ét}}(Y', \Lambda) \rightarrow D_{\text{ét}}(Y, \Lambda)$$

*is the functor obtained from the previous discussion by passage to homotopy categories.*

It is easy to see that this is independent of the choice of the simplicial  $v$ -hypercover, by passing to common refinements. Again, we need to check that this still satisfies all desired properties, starting with base change.

**Proposition 22.19.** *Let  $f : Y' \rightarrow Y$  be a compactifiable map of small  $v$ -stacks which is representable in locally spatial diamonds with  $\dim. \text{trg } f < \infty$ , and assume  $n\Lambda = 0$  for some  $n$  prime to  $p$ . Let  $g : \tilde{Y} \rightarrow Y$  be any map of small  $v$ -stacks, with base change  $\tilde{f} : \tilde{Y}' = Y' \times_Y \tilde{Y} \rightarrow \tilde{Y}$  and  $g' : \tilde{Y}' \rightarrow Y'$ .*

*There is a natural base change equivalence*

$$g^* Rf_! \simeq R\tilde{f}_! g'^*$$

*of functors  $D_{\text{ét}}(Y', \Lambda) \rightarrow D_{\text{ét}}(\tilde{Y}, \Lambda)$ .*

*Proof.* Let  $Y_\bullet \rightarrow Y$  be a simplicial  $v$ -hypercover by locally spatial diamonds  $Y_i$ , and similarly let  $\tilde{Y}_\bullet \rightarrow \tilde{Y}$  be a simplicial  $v$ -hypercover by locally spatial diamonds  $\tilde{Y}_i$ , such that  $g : \tilde{Y} \rightarrow Y$  extends to a map of simplicial spaces  $g_\bullet : \tilde{Y}_\bullet \rightarrow Y_\bullet$ . Repeating the previous discussion over  $\Delta$  again over  $\Delta \times \Delta^1$  gives the result.  $\square$

**Proposition 22.20.** *Let  $f : Y' \rightarrow Y$  be a compactifiable map of small  $v$ -stacks that is representable in locally spatial diamonds with  $\dim. \text{trg } f < \infty$ , and assume  $n\Lambda = 0$  for some  $n$  prime to  $p$ . The functor*

$$Rf_! : \mathcal{D}_{\text{ét}}(Y', \Lambda) \rightarrow \mathcal{D}_{\text{ét}}(Y, \Lambda)$$

*commutes with all direct sums; equivalently (cf. [Lur16, Proposition 1.4.4.1 (2)]), with all colimits.*

*Proof.* This follows from Proposition 22.14 and Proposition 22.19.  $\square$

**Proposition 22.21.** *Let  $g : Y'' \rightarrow Y'$  and  $f : Y' \rightarrow Y$  be compactifiable maps of small  $v$ -stacks which are representable in locally spatial diamonds with  $\dim. \text{trg } f, \dim. \text{trg } g < \infty$ , and assume  $n\Lambda = 0$  for some  $n$  prime to  $p$ . Then there is a natural equivalence*

$$Rf_! \circ Rg_! \simeq R(f \circ g)_!$$

*of functors  $D_{\text{ét}}(Y'', \Lambda) \rightarrow D_{\text{ét}}(Y, \Lambda)$ .*

*Proof.* Let  $Y_\bullet \rightarrow Y$  be a simplicial  $v$ -hypercover by locally spatial diamonds, as usual, and let  $Y'_\bullet \rightarrow Y'$ ,  $Y''_\bullet \rightarrow Y''$  be the pullbacks, which are again simplicial  $v$ -hypercovers by locally spatial diamonds. We seek to find an equivalence of the composite of the functors

$$Rg_{\bullet!}^0 : \mathcal{D}_{\text{ét}}(Y''_\bullet, \Lambda)^0 \rightarrow \mathcal{D}_{\text{ét}}(Y'_\bullet, \Lambda)^0$$

and

$$Rf_{\bullet!}^0 : \mathcal{D}_{\text{ét}}(Y', \Lambda)^0 \rightarrow \mathcal{D}_{\text{ét}}(Y_{\bullet}, \Lambda)^0$$

with

$$R(f \circ g)_{\bullet!}^0 : \mathcal{D}_{\text{ét}}(Y'', \Lambda)^0 \rightarrow \mathcal{D}_{\text{ét}}(Y_{\bullet}, \Lambda)^0 .$$

This gives the result by passage to coCartesian sections. But one can directly identify the functors on the full  $\infty$ -subcategory  $\mathcal{D}_{\text{ét,prop}/Y_{\bullet}}(Y'', \Lambda)^0$  (as there is a natural transformation, which is an equivalence, by Proposition 22.9 and its proof). On the other hand, all functors commute with all colimits by Proposition 22.20, so the full functors can be recovered by left Kan extension (as above).  $\square$

In case of overlap, the current definition of  $Rf_{\bullet}$  agrees with Definition 19.1.

**Proposition 22.22.** *Let  $f : Y' \rightarrow Y$  be a separated étale map of small  $v$ -stacks, and assume  $n\Lambda = 0$  for some  $n$  prime to  $p$ . Then  $Rf_{\bullet}$  as defined in Definition 19.1 agrees with  $Rf_{\bullet}$  as defined in Definition 22.18.*

*Proof.* As in the proof of Proposition 22.10, one has a natural transformation. To check whether it is an equivalence, one can reduce to the case where  $Y$  is strictly totally disconnected. As moreover both functors commute with all direct sums, one can reduce to the case where  $Y'$  is quasicompact, where it follows from Proposition 22.10.  $\square$

Finally, we also get the projection formula in general.

**Proposition 22.23.** *Let  $f : Y' \rightarrow Y$  be a compactifiable map of small  $v$ -sheaves which is representable in locally spatial diamonds with  $\dim. \text{trg } f < \infty$ , and let  $\Lambda$  be a ring with  $n\Lambda = 0$  for some  $n$  prime to  $p$ . Then there is a functorial isomorphism*

$$Rf_{\bullet} B \otimes_{\Lambda}^{\mathbb{L}} A \simeq Rf_{\bullet}(B \otimes_{\Lambda}^{\mathbb{L}} f^* A)$$

in  $B \in D_{\text{ét}}(Y', \Lambda)$  and  $A \in D_{\text{ét}}(Y, \Lambda)$ .

*Proof.* We fix  $A \in D_{\text{ét}}(Y, \Lambda)$ , and define an isomorphism

$$Rf_{\bullet} B \otimes_{\Lambda}^{\mathbb{L}} A \simeq Rf_{\bullet}(B \otimes_{\Lambda}^{\mathbb{L}} f^* A)$$

functorial in  $B \in D_{\text{ét}}(Y', \Lambda)$ . One then checks that varying  $A$ , the relevant diagrams commute in the derived category.

Choose a simplicial  $v$ -hypercover  $Y_{\bullet} \rightarrow Y$  by locally spatial diamonds as in the definition of  $Rf_{\bullet}$ , with pullback  $Y'_{\bullet} \rightarrow Y'$ . Consider the functor

$$Rf_{\bullet!}^0 : \mathcal{D}_{\text{ét}}(Y'_{\bullet}, \Lambda)^0 \rightarrow \mathcal{D}_{\text{ét}}(Y_{\bullet}, \Lambda)^0 .$$

Both  $\infty$ -categories have an endofunctor given by tensoring with the pullback of  $A$ ; we denote this operation by  $- \otimes_{\Lambda}^{\mathbb{L}} A|_{Y'_{\bullet}}$  respectively  $- \otimes_{\Lambda}^{\mathbb{L}} A|_{Y_{\bullet}}$ . We claim that there is a natural equivalence of functors

$$Rf_{\bullet!}^0(B \otimes_{\Lambda}^{\mathbb{L}} A|_{Y'_{\bullet}}) \simeq Rf_{\bullet!}^0(B \otimes_{\Lambda}^{\mathbb{L}} A|_{Y_{\bullet}})$$

from  $B \in \mathcal{D}_{\text{ét}}(Y'_{\bullet}, \Lambda)^0$  to  $\mathcal{D}_{\text{ét}}(Y_{\bullet}, \Lambda)^0$ . This implies the desired result by passage to coCartesian sections (noting that both functors map coCartesian edges to coCartesian edges, as the second functor does).

To construct the natural equivalence, note that both functors are left Kan extensions from  $\mathcal{D}_{\text{ét,prop}/Y_\bullet}(Y'_\bullet, \Lambda)^0$ . On this full  $\infty$ -subcategory, the first functor is the composite

$$\mathcal{D}_{\text{ét,prop}/Y_\bullet}(Y'_\bullet, \Lambda)^0 \xrightarrow{\otimes_{\Lambda}^{\mathbb{L}} A|_{Y'_\bullet}} \mathcal{D}_{\text{ét,prop}/Y_\bullet}(Y'_\bullet, \Lambda)^0 \xrightarrow{j_{\bullet!}} \mathcal{D}_{\text{ét}}(\overline{Y'_\bullet}/Y_\bullet, \Lambda)^0 \xrightarrow{R\overline{f}_{\bullet*}^{/Y_\bullet}} \mathcal{D}_{\text{ét}}(Y_\bullet, \Lambda)^0 .$$

This admits a natural map to the composite

$$\mathcal{D}_{\text{ét,prop}/Y_\bullet}(Y'_\bullet, \Lambda)^0 \xrightarrow{j_{\bullet!}} \mathcal{D}_{\text{ét}}(\overline{Y'_\bullet}/Y_\bullet, \Lambda)^0 \xrightarrow{\otimes_{\Lambda}^{\mathbb{L}} A|_{\overline{Y'_\bullet}/Y_\bullet}} \mathcal{D}_{\text{ét}}(\overline{Y'_\bullet}/Y_\bullet, \Lambda)^0 \xrightarrow{R\overline{f}_{\bullet*}^{/Y_\bullet}} \mathcal{D}_{\text{ét}}(Y_\bullet, \Lambda)^0 ,$$

by using that  $j_{\bullet!}$  is left adjoint to  $j_{\bullet*}$ , and that pullback commutes with  $- \otimes_{\Lambda}^{\mathbb{L}} -$ . This natural transformation is an equivalence (even without postcomposition with  $R\overline{f}_{\bullet*}^{/Y_\bullet}$ ), as can be checked in each fibre, where it reduces to the projection formula for an open embedding.

On the other hand, the second functor is the composite

$$\mathcal{D}_{\text{ét,prop}/Y_\bullet}(Y'_\bullet, \Lambda)^0 \xrightarrow{j_{\bullet!}} \mathcal{D}_{\text{ét}}(\overline{Y'_\bullet}/Y_\bullet, \Lambda)^0 \xrightarrow{R\overline{f}_{\bullet*}^{/Y_\bullet}} \mathcal{D}_{\text{ét}}(Y_\bullet, \Lambda)^0 \xrightarrow{\otimes_{\Lambda}^{\mathbb{L}} A|_{Y_\bullet}} \mathcal{D}_{\text{ét}}(Y_\bullet, \Lambda)^0 .$$

This also admits a natural map to the composition

$$\mathcal{D}_{\text{ét,prop}/Y_\bullet}(Y'_\bullet, \Lambda)^0 \xrightarrow{j_{\bullet!}} \mathcal{D}_{\text{ét}}(\overline{Y'_\bullet}/Y_\bullet, \Lambda)^0 \xrightarrow{\otimes_{\Lambda}^{\mathbb{L}} A|_{\overline{Y'_\bullet}/Y_\bullet}} \mathcal{D}_{\text{ét}}(\overline{Y'_\bullet}/Y_\bullet, \Lambda)^0 \xrightarrow{R\overline{f}_{\bullet*}^{/Y_\bullet}} \mathcal{D}_{\text{ét}}(Y_\bullet, \Lambda)^0$$

considered above, by using that  $R\overline{f}_{\bullet*}^{/Y_\bullet}$  is right adjoint to  $\overline{f}_{\bullet*}^{/Y_\bullet,*}$ , and that pullback commutes with  $- \otimes_{\Lambda}^{\mathbb{L}} -$ . Again, one can check that it is a natural equivalence by checking it on fibres, where it reduces to the projection formula in the quasicompact case, Proposition 22.11. Combining these observations finishes the proof.  $\square$

### 23. COHOMOLOGICALLY SMOOTH MORPHISMS

We start with the following theorem, which is essentially a corollary of Proposition 22.20.

**Theorem 23.1.** *Let  $f : Y' \rightarrow Y$  be a compactifiable map of small  $v$ -stacks which is representable in locally spatial diamonds with  $\dim. \text{trg } f < \infty$ , and let  $\Lambda$  be a ring with  $n\Lambda = 0$  for some  $n$  prime to  $p$ . Then the functor*

$$Rf_! : D_{\text{ét}}(Y', \Lambda) \rightarrow D_{\text{ét}}(Y, \Lambda)$$

*admits a right adjoint*

$$Rf^! : D_{\text{ét}}(Y, \Lambda) \rightarrow D_{\text{ét}}(Y', \Lambda) .$$

*Proof.* This follows from Lurie's  $\infty$ -categorical adjoint functor theorem, [Lur09, Corollary 5.5.2.9], and Proposition 22.20.  $\square$

**Remark 23.2.** The functor  $Rf^!$  is compatible with change of rings  $g : \Lambda' \rightarrow \Lambda$  in the following sense. The map  $g$  induces a map  $D_{\text{ét}}(Y', \Lambda) \rightarrow D_{\text{ét}}(Y', \Lambda')$  by restricting the action along  $g$ . Then the diagram

$$\begin{array}{ccc} D_{\text{ét}}(Y, \Lambda) & \xrightarrow{Rf^!} & D_{\text{ét}}(Y', \Lambda) \\ \downarrow & & \downarrow \\ D_{\text{ét}}(Y, \Lambda') & \xrightarrow{Rf^!} & D_{\text{ét}}(Y', \Lambda') \end{array}$$

commutes. Indeed, this identity of functors is the right adjoint of the identity of functors

$$Rf_!(- \otimes_{\Lambda'} \Lambda) = Rf_! \otimes_{\Lambda'} \Lambda ,$$

which follows from the projection formula, Proposition 22.23.

Before going on, we note that the following results follow from the definitions and Proposition 22.23.

**Proposition 23.3.** *Let  $f : Y' \rightarrow Y$  be a compactifiable map of small  $v$ -stacks that is representable in locally spatial diamonds with  $\dim. \operatorname{trg} f < \infty$ , and let  $\Lambda$  be a ring with  $n\Lambda = 0$  for some  $n$  prime to  $p$ .*

(i) *For all  $A \in D_{\acute{e}t}(Y', \Lambda)$ ,  $B \in D_{\acute{e}t}(Y, \Lambda)$ , one has*

$$R\mathcal{H}om_{\Lambda}(Rf_!A, B) \cong Rf_*R\mathcal{H}om_{\Lambda}(A, Rf^!B) .$$

(ii) *For all  $A, B \in D_{\acute{e}t}(Y, \Lambda)$ , one has*

$$Rf^!R\mathcal{H}om_{\Lambda}(A, B) \cong R\mathcal{H}om_{\Lambda}(f^*A, Rf^!B) .$$

*Proof.* For part (i), note that for all  $C \in D_{\acute{e}t}(Y, \Lambda)$ , one has

$$\begin{aligned} \operatorname{Hom}_{D_{\acute{e}t}(Y, \Lambda)}(C, R\mathcal{H}om_{\Lambda}(Rf_!A, B)) &= \operatorname{Hom}_{D_{\acute{e}t}(Y, \Lambda)}(C \otimes_{\Lambda}^{\mathbb{L}} Rf_!A, B) \\ &= \operatorname{Hom}_{D_{\acute{e}t}(Y, \Lambda)}(Rf_!(f^*C \otimes_{\Lambda}^{\mathbb{L}} A), B) \\ &= \operatorname{Hom}_{D_{\acute{e}t}(Y', \Lambda)}(f^*C \otimes_{\Lambda}^{\mathbb{L}} A, Rf^!B) \\ &= \operatorname{Hom}_{D_{\acute{e}t}(Y', \Lambda)}(f^*C, R\mathcal{H}om_{\Lambda}(A, Rf^!B)) \\ &= \operatorname{Hom}_{D_{\acute{e}t}(Y, \Lambda)}(C, Rf_*R\mathcal{H}om_{\Lambda}(A, Rf^!B)) , \end{aligned}$$

so the result follows from the Yoneda lemma. Similarly, for part (ii), note that for all  $C \in D_{\acute{e}t}(Y', \Lambda)$ , one has

$$\begin{aligned} \operatorname{Hom}_{D_{\acute{e}t}(Y', \Lambda)}(C, Rf^!R\mathcal{H}om_{\Lambda}(A, B)) &= \operatorname{Hom}_{D_{\acute{e}t}(Y, \Lambda)}(Rf_!C, R\mathcal{H}om_{\Lambda}(A, B)) \\ &= \operatorname{Hom}_{D_{\acute{e}t}(Y, \Lambda)}(Rf_!C \otimes_{\Lambda}^{\mathbb{L}} A, B) \\ &= \operatorname{Hom}_{D_{\acute{e}t}(Y, \Lambda)}(Rf_!(C \otimes_{\Lambda}^{\mathbb{L}} f^*A), B) \\ &= \operatorname{Hom}_{D_{\acute{e}t}(Y', \Lambda)}(C \otimes_{\Lambda}^{\mathbb{L}} f^*A, Rf^!B) \\ &= \operatorname{Hom}_{D_{\acute{e}t}(Y', \Lambda)}(C, R\mathcal{H}om_{\Lambda}(f^*A, Rf^!B)) . \end{aligned}$$

□

As a preparation for the definition of smooth morphisms, we prove the following proposition.

**Proposition 23.4.** *Let  $X$  be a strictly totally disconnected perfectoid space, let  $f : Y \rightarrow X$  be a compactifiable map from a locally spatial diamond  $Y$  of  $\dim. \operatorname{trg} f < \infty$ , and fix a prime  $\ell \neq p$ . The following conditions are equivalent.*

(i) *The natural transformation*

$$Rf^! \mathbb{F}_{\ell} \otimes_{\mathbb{F}_{\ell}} f^* \rightarrow Rf^! : D_{\acute{e}t}(X, \mathbb{F}_{\ell}) \rightarrow D_{\acute{e}t}(Y, \mathbb{F}_{\ell})$$

*adjoint to*

$$Rf_!(Rf^! \mathbb{F}_{\ell} \otimes_{\mathbb{F}_{\ell}} f^* -) = Rf_!Rf^! \mathbb{F}_{\ell} \otimes_{\mathbb{F}_{\ell}} \operatorname{id} \rightarrow \operatorname{id}$$

*is an equivalence (using the projection formula, Proposition 22.23, in the equality).*

- (ii) The functor  $Rf^! : D_{\text{ét}}(X, \mathbb{F}_\ell) \rightarrow D_{\text{ét}}(Y, \mathbb{F}_\ell)$  is equivalent to a functor of the form  $A \otimes_{\mathbb{F}_\ell} f^*$  for some  $A \in D_{\text{ét}}(Y, \mathbb{F}_\ell)$ .
- (iii) The functor  $Rf^! : D_{\text{ét}}(X, \mathbb{F}_\ell) \rightarrow D_{\text{ét}}(Y, \mathbb{F}_\ell)$  commutes with arbitrary direct sums, and for all connected components  $X_0 = \text{Spa}(C, C^+) \subset X$  with an open subset  $j : U \subset X_0$  and pullbacks

$$\begin{array}{ccccc} V & \xrightarrow{j'} & Y_0 & \longrightarrow & Y \\ \downarrow f_U & & \downarrow f_0 & & \downarrow f \\ U & \xrightarrow{j} & X_0 & \longrightarrow & X \end{array},$$

the map

$$j'_! Rf_U^! \mathbb{F}_\ell \rightarrow Rf_0^! j_! \mathbb{F}_\ell$$

adjoint to

$$Rf_0^! j'_! Rf_U^! \mathbb{F}_\ell = j_! Rf_U^! Rf_0^! \mathbb{F}_\ell \rightarrow j_! \mathbb{F}_\ell$$

is an equivalence.

- (iv) For all affinoid pro-étale maps  $g : X' \rightarrow X$  with pullback

$$\begin{array}{ccc} Y' & \xrightarrow{h} & Y \\ \downarrow f' & & \downarrow f \\ X' & \xrightarrow{g} & X \end{array},$$

the natural transformation of functors

$$h^* Rf^! \rightarrow Rf'^! g^* : D_{\text{ét}}(X, \mathbb{F}_\ell) \rightarrow D_{\text{ét}}(Y', \mathbb{F}_\ell)$$

adjoint to

$$Rf'_! h^* Rf^! = g^* Rf_! Rf^! \rightarrow g^*$$

is an equivalence, and for all connected components  $X_0 = \text{Spa}(C, C^+) \subset X$  with an open subset  $j : U \subset X_0$  and pullbacks

$$\begin{array}{ccccc} V & \xrightarrow{j'} & Y_0 & \longrightarrow & Y \\ \downarrow f_U & & \downarrow f_0 & & \downarrow f \\ U & \xrightarrow{j} & X_0 & \longrightarrow & X \end{array},$$

the natural transformation of functors

$$j'_! Rf_U^! \rightarrow Rf_0^! j_! : D_{\text{ét}}(U, \mathbb{F}_\ell) \rightarrow D_{\text{ét}}(Y_0, \mathbb{F}_\ell)$$

adjoint to

$$Rf_0^! j'_! Rf_U^! = j_! Rf_U^! Rf_0^! \rightarrow j_!$$

is an equivalence.

Moreover, under these conditions, for any  $\ell$ -power-torsion ring  $\Lambda$ , the natural transformation

$$Rf^! \Lambda \otimes_{\Lambda} f^* \rightarrow Rf^! : D_{\text{ét}}(X, \Lambda) \rightarrow D_{\text{ét}}(Y, \Lambda)$$

is an equivalence.

**Remark 23.5.** As in the discussion around Theorem 19.2, the second part of condition (iv) can be regarded as a version of the first part for pullback to closed but non-generalizing subsets. Thus, condition (iv) is expressing a version of the idea that “ $Rf^!$  commutes with arbitrary pro-étale base change”. Of course, if  $Rf^!$  is essentially given by  $f^*$ , this should be true. What is maybe surprising is that the converse also holds.

*Proof.* It is clear that (i) implies (ii), and (ii) implies (iii). To see that (iii) implies (iv), we first check that if  $Rf^!$  commutes with arbitrary direct sums, then the first part of condition (iv) is satisfied. For this, let  $g : X' = \varprojlim X'_i \rightarrow X$  be an inverse limit of affinoid étale maps  $g_i : X'_i \rightarrow X$ , and consider the cartesian diagrams

$$\begin{array}{ccc} Y' & \xrightarrow{h} & Y \\ \downarrow f' & & \downarrow f \\ X' & \xrightarrow{g} & X \end{array} \quad \begin{array}{ccc} Y'_i & \xrightarrow{h_i} & Y \\ \downarrow f'_i & & \downarrow f \\ X'_i & \xrightarrow{g_i} & X . \end{array}$$

We need to check that the natural transformation

$$h^* Rf^! \rightarrow Rf'^! g^*$$

is an equivalence. Note that by Remark 21.14, the functor  $h_* = Rh_*$  is exact, and one checks easily that it is also conservative (like pushforward along any quasicompact separated quasi-pro-étale map). Thus, it suffices to show that the natural transformation

$$h_* h^* Rf^! \rightarrow h_* Rf'^! g^* = Rf^! g_* g^*$$

is an equivalence. But note that  $g_* g^*$  is the filtered colimit of  $g_{i*} g_i^*$ , and correspondingly  $h_* h^*$  is the filtered colimit of  $h_{i*} h_i^*$ . If  $Rf^!$  commutes with arbitrary direct sums,  $Rf^!$  commutes with arbitrary colimits, and thus the preceding argument reduces us to the case of  $g_i : X'_i \rightarrow X$ . But if  $g = g_i$  is étale, then  $g^* = Rg^!$  and  $h^* = Rh^!$ , so  $h^* Rf^! = Rh^! Rf^! = Rf'^! Rg^! = Rf'^! g^*$ , as desired.

Now, for the second part of condition (iv), one checks that the full subcategory of all  $K \in D_{\text{ét}}(U, \mathbb{F}_\ell)$  which satisfy the conclusion is triangulated and stable under arbitrary direct sums, and contains  $j'_! \mathbb{F}_\ell$  for all open immersions  $j' : U' \hookrightarrow U$ . This implies that it is all of  $D_{\text{ét}}(U, \mathbb{F}_\ell)$ .

For (iv) implies (i), we have to check that

$$Rf^! \mathbb{F}_\ell \otimes f^* \rightarrow Rf^! : D_{\text{ét}}(X, \mathbb{F}_\ell) \rightarrow D_{\text{ét}}(Y, \mathbb{F}_\ell)$$

is an equivalence. This can be checked in fibers over points of  $X$ . As both sides commute with base change by the first part of condition (iv), we can thus assume that  $X = \text{Spa}(C, C^+)$  is strictly local. Fix a geometric point  $y$  of  $Y$ ; we want to prove that the map  $(Rf^! \mathbb{F}_\ell \otimes f^*)_y \rightarrow (Rf^!)_y$  of stalks at  $y$  is an isomorphism. Let  $x \in X$  be the image of  $y$ , let  $X_1 = \text{Spa}(C, (C^+)' ) \subset X$  be the set of generalizations of  $x$ , and let  $U = X_1 \setminus \{x\}$ . Then we can pullback further under the open immersion  $X_1 \hookrightarrow X$ , and assume that  $X = X_1$ . For any  $K \in D_{\text{ét}}(X, \mathbb{F}_\ell)$ , we have a triangle  $j_! j^* K \rightarrow K \rightarrow K'$ , where  $j : U \hookrightarrow X_1 = X$  denotes the open immersion. By the second condition of (iv), we see that the stalk at  $y$  of the target of the map

$$Rf^! \mathbb{F}_\ell \otimes f^* j_! K|_U \rightarrow Rf^! j_! K|_U$$

vanishes; it also vanishes on the source, as the second factor  $f^* j_! K|_U$  does. Thus, we can replace  $K$  by  $K'$ , and assume that  $K|_U = 0$ . In that case,  $K = i_* K_0$  for some complex  $K_0 \in D(\mathbb{F}_\ell)$ , where

$i : \{x\} \rightarrow |X|$  denotes the closed inclusion. Repeating the argument with the displayed triangle, we can also replace  $K$  by the constant sheaf  $K_0$ .

As a further reduction step, we reduce to the case that  $K_0$  is concentrated in degree 0. Assume for the moment that the result holds true in this case. By triangles, it holds if  $K_0 \in D^b(\mathbb{F}_\ell)$  is bounded. We claim that it also holds if  $K_0 \in D^+(\mathbb{F}_\ell)$ . For this, it is enough to show that if  $K_0 \in D^{\geq n}(\mathbb{F}_\ell)$ , then for some constant  $c$ , both sides lie in  $D^{\geq n-c}(Y, \Lambda)$ . This in turn reduces to proving a similar result for  $Rf^!$ , which follows formally from the fact that  $Rf_!$  has finite cohomological dimension. It remains to handle the case of  $K_0 \in D^-(\mathbb{F}_\ell)$ . Writing this as a limit of its Postnikov truncations and using that  $Rf^!$  commutes with derived limits, it is enough to prove that  $Rf^!\mathbb{F}_\ell \in D_{\text{ét}}^{\leq 0}(Y, \Lambda)$ . This can be checked on stalks, which are given by colimits of  $R\text{Hom}_X(Rf_!\mathbb{F}_\ell, \mathbb{F}_\ell)$  for  $f' : Y' \rightarrow Y \rightarrow X$  the composition of an étale map  $Y' \rightarrow Y$  with  $f : Y \rightarrow X$ . But if  $j_\eta : \{\eta\} \rightarrow X$  is the inclusion of the maximal point, then using Lemma 21.13

$$R\text{Hom}_X(-, \mathbb{F}_\ell) = R\text{Hom}_X(-, Rj_{\eta*}\mathbb{F}_\ell) = R\text{Hom}_\eta(j^*- , \mathbb{F}_\ell) = \text{Hom}_\eta(j^*- , \mathbb{F}_\ell) ,$$

as on a geometric point,  $\text{Hom}(-, \mathbb{F}_\ell)$  is exact. As  $Rf_!\mathbb{F}_\ell \in D^{\geq 0}(X, \mathbb{F}_\ell)$ , this implies that

$$R\text{Hom}_X(Rf_!\mathbb{F}_\ell, \mathbb{F}_\ell) \in D^{\leq 0}(\mathbb{F}_\ell) ,$$

as desired.

Thus, it remains to handle the case that  $K_0$  is concentrated in degree 0. In this case,  $K_0 = V[0]$  for some  $\mathbb{F}_\ell$ -vector space  $V$ . We can assume that  $V = C^0(S, \mathbb{F}_\ell)$  is the space of continuous functions on some profinite set  $S$ . Let  $X' = X \times \underline{S}$ , so that  $h : X' \rightarrow X$  is quasicompact separated pro-étale. Let  $f' : Y' = X' \times_X Y \rightarrow X'$  be the pullback, which is given by  $Y' = Y \times \underline{S}$ . Using the first part of condition (iv), we get

$$Rf'^!h^*\mathbb{F}_\ell = h'^*Rf^!\mathbb{F}_\ell .$$

Applying  $Rh'_*$  and using that  $Rh'_*Rf'^! = Rf^!Rh_*$  by Proposition 23.16 (i), we get

$$Rf^!Rh_*h^*\mathbb{F}_\ell = Rh'_*h'^*Rf^!\mathbb{F}_\ell .$$

By Lemma 23.6 below, this translates into

$$Rf^!V = V \otimes_{\mathbb{F}_\ell} Rf^!\mathbb{F}_\ell ,$$

as desired.

This shows the equivalence of conditions (i) through (iv). To handle the case of a general  $\ell$ -power-torsion ring  $\Lambda$ , choose  $m$  such that  $\ell^m = 0$  in  $\Lambda$ . Then  $\Lambda$  is a  $\mathbb{Z}/\ell^m\mathbb{Z}$ -algebra, and using Remark 23.2, one reduces to the case  $\Lambda = \mathbb{Z}/\ell^m\mathbb{Z}$ . Any  $K \in D_{\text{ét}}(X, \Lambda)$  is filtered by  $m$  copies of  $K \otimes_{\mathbb{Z}/\ell^m\mathbb{Z}} \mathbb{F}_\ell$ , so one can assume that  $K$  comes from  $D_{\text{ét}}(X, \mathbb{F}_\ell)$  via restriction of coefficients. Using the result for  $\mathbb{F}_\ell$ , this reduces to showing that

$$Rf^!\mathbb{Z}/\ell^m\mathbb{Z} \otimes_{\mathbb{Z}/\ell^m\mathbb{Z}} \mathbb{F}_\ell \rightarrow Rf^!\mathbb{F}_\ell$$

is an isomorphism. For this, note that condition (iv) also holds for  $\mathbb{Z}/\ell^m\mathbb{Z}$  in place of  $\mathbb{F}_\ell$ , by filtering by  $m$  copies of  $- \otimes_{\mathbb{Z}/\ell^m\mathbb{Z}} \mathbb{F}_\ell$ . Thus, as above, we can reduce to the case that  $X = \text{Spa}(C, C^+)$  is connected. Now one uses the standard infinite resolution of  $\mathbb{F}_\ell$  as a  $\mathbb{Z}/\ell^m\mathbb{Z}$ -module, and the observation  $Rf^!\mathbb{Z}/\ell^m\mathbb{Z} \in D_{\text{ét}}^{\leq 0}(Y, \mathbb{Z}/\ell^m\mathbb{Z})$ . The latter is proved in the same way as  $Rf^!\mathbb{F}_\ell \in D_{\text{ét}}^{\leq 0}(Y, \mathbb{F}_\ell)$  above, using that  $\mathbb{Z}/\ell^m\mathbb{Z}$  is self-injective.  $\square$



**Lemma 23.6.** *Let  $S$  be a profinite set, let  $Y$  be a small  $v$ -sheaf, and consider  $h : Y \times \underline{S} \rightarrow Y$ . Then for any ring  $\Lambda$  and any  $C \in D_{\text{ét}}(Y, \Lambda)$ , one has a natural isomorphism*

$$Rh_*h^*C \simeq C^0(S, \Lambda) \otimes_{\Lambda} C,$$

where  $C^0(S, \Lambda)$  is the  $\Lambda$ -module of continuous maps  $S \rightarrow \Lambda$ .

*Proof.* We may assume that  $Y$  is a strictly totally disconnected perfectoid space. In this case, write  $S = \varprojlim_i S_i$  as an inverse limit of finite sets, and compute both sides as a filtered colimit, using Proposition 14.9 if  $C \in D_{\text{ét}}^+(Y, \Lambda)$ ; in general pass to a Postnikov limit.  $\square$

**Proposition 23.7.** *Let  $X$  be a strictly totally disconnected perfectoid space,  $f : Y \rightarrow X$  a compactifiable map from a spatial diamond  $Y$  with  $\dim. \text{trg } f < \infty$ , and fix a prime  $\ell \neq p$ .*

*The functor*

$$Rf^! : D_{\text{ét}}(X, \mathbb{F}_{\ell}) \rightarrow D_{\text{ét}}(Y, \mathbb{F}_{\ell})$$

*commutes with arbitrary direct sums if and only if for all constructible sheaves  $\mathcal{F}$  of  $\mathbb{F}_{\ell}$ -vector spaces on  $Y_{\text{ét}}$  and all  $i \geq 0$ , the  $!$ -pushforward  $R^i f_! \mathcal{F}$  is constructible on  $X_{\text{ét}}$ .*

*Proof.* First, we claim that  $Y$  satisfies the hypothesis of Proposition 20.10. For this, we have to see that for any quasicompact separated étale map  $j : U \rightarrow Y$ , the  $\ell$ -cohomological dimension of  $U_{\text{ét}}$  is bounded by  $N$ , for some fixed integer  $N$ . Here, we claim that we can take  $N = 2 \dim. \text{trg } f$ . Indeed, let  $\bar{U}^{/X} \rightarrow X$  be the canonical compactification of  $U \rightarrow X$ ; then, using Lemma 21.13, it suffices to show that the  $\ell$ -cohomological dimension of  $\bar{U}^{/X}$  is bounded by  $2 \dim. \text{trg } f$ . As  $\pi_0 X$  is profinite, this can be checked on connected components of  $\pi_0 X$ , so we can assume that  $X = \text{Spa}(C, C^+)$  is connected. As  $\bar{U}^{/X} \rightarrow X$  is proper, Theorem 19.2 allows us to restrict to sheaves concentrated on the fiber of  $\bar{U}^{/X}$  over the closed point of  $|X|$ . But then Proposition 21.11 combined with  $\dim f \leq \dim. \text{trg } f$  and  $\text{cd}_{\ell} G_y \leq \dim. \text{trg } f$  finishes the argument.

Now it follows from Proposition 20.10 that both  $D_{\text{ét}}(Y, \mathbb{F}_{\ell})$  and  $D_{\text{ét}}(X, \mathbb{F}_{\ell})$  are compactly generated, with compact objects given by bounded complexes whose cohomology sheaves are constructible. It is easy to see that the second condition is equivalent to the condition that  $Rf_! : D_{\text{ét}}(Y, \mathbb{F}_{\ell}) \rightarrow D_{\text{ét}}(X, \mathbb{F}_{\ell})$  preserves compact objects. It follows from adjunction that  $Rf^!$  commutes with arbitrary direct sums if and only if  $Rf_!$  preserves compact objects, so we get the result.  $\square$

Finally, we can give the definition of cohomological smoothness.

**Definition 23.8.** *Let  $f : Y' \rightarrow Y$  be a separated map of small  $v$ -stacks that is representable in locally spatial diamonds, and let  $\ell \neq p$  be a prime. Then  $f$  is  $\ell$ -cohomologically smooth if  $f$  is compactifiable,  $\dim. \text{trg } f < \infty$ , and for any strictly totally disconnected perfectoid space  $X$  with a map  $X \rightarrow Y$  with pullback  $f_X : Y' \times_Y X \rightarrow X$ , the functor*

$$Rf_X^! : D_{\text{ét}}(X, \mathbb{F}_{\ell}) \rightarrow D_{\text{ét}}(Y' \times_Y X, \mathbb{F}_{\ell})$$

*is equivalent to a functor of the form  $Rf_X^! = D_{f_X} \otimes_{\mathbb{F}_{\ell}} f^*$  for some invertible object  $D_{f_X} \in D_{\text{ét}}(Y' \times_Y X, \mathbb{F}_{\ell})$ .*

Here, for a locally spatial diamond  $Y$ , an object  $D \in D_{\text{ét}}(Y, \mathbb{F}_{\ell})$  is *invertible* if it is locally (equivalently, in the  $v$ -, quasi-pro-étale, or étale topology of  $Y$ ) isomorphic to  $\mathbb{F}_{\ell}[n]$  for some integer  $n \in \mathbb{Z}$ . We note that we are not a priori asking that the equivalence between  $Rf_X^!$  and  $D_{f_X} \otimes_{\mathbb{F}_{\ell}} f^*$  is natural, that  $D_{f_X}$  commutes with base change, or similar results. These are however automatic consequences, cf. Proposition 23.12.

**Remark 23.9.** The definition is phrased in a way that leaves room for a more general definition of  $\ell$ -cohomological smoothness for morphisms which are not separated, or not representable in locally spatial diamonds. For example, the map  $\underline{M} \rightarrow *$  for a topological manifold  $M$  should be considered smooth, but it is not representable in locally spatial diamonds. Moreover, smoothness should be a local condition, while being separated is not.

Before investigating cohomologically smooth morphisms in detail, we give a different characterization that is easier to check in practice.

**Proposition 23.10.** *Let  $f : Y' \rightarrow Y$  be a compactifiable map of small  $v$ -stacks which is representable in spatial diamonds, and let  $\ell \neq p$  be a prime. Then  $f$  is  $\ell$ -cohomologically smooth if and only if the following conditions are satisfied.*

- (i) *The dimension  $\dim. \text{trg } f < \infty$  is finite.*
- (ii) *For any strictly totally disconnected perfectoid space  $X$  with a map  $g : X \rightarrow Y$  and pullback  $f_X : Y' \times_Y X \rightarrow X$ , and any constructible étale sheaf  $\mathcal{F}$  of  $\mathbb{F}_\ell$ -vector spaces on the spatial diamond  $Y' \times_Y X$ , the  $!$ -pushforward  $R^i f_{X!} \mathcal{F}$  is constructible on  $X$ , for all  $i \geq 0$ .*
- (iii) *For any  $X = \text{Spa}(C, C^+)$  with  $C$  an algebraically closed field and  $C^+ \subset C$  an open and bounded valuation subring, with an open subset  $j : U \rightarrow X$ , and pullbacks*

$$\begin{array}{ccccc} Y' \times_Y U & \xrightarrow{j'} & Y' \times_Y X & \longrightarrow & Y' \\ \downarrow f_U & & \downarrow f_X & & \downarrow f \\ U & \xrightarrow{j} & X & \longrightarrow & Y, \end{array}$$

the natural map

$$j'_! Rf_U^! \mathbb{F}_\ell \rightarrow Rf_X^! j_! \mathbb{F}_\ell$$

adjoint to

$$Rf_{X!} j'_! Rf_U^! \mathbb{F}_\ell = j_! Rf_{U!} Rf_U^! \mathbb{F}_\ell \rightarrow j_! \mathbb{F}_\ell$$

is an equivalence.

- (iv) *For any strictly totally disconnected perfectoid space  $X$  with a map  $g : X \rightarrow Y$  and pullback  $f_X : Y' \times_Y X \rightarrow X$ , the  $!$ -pullback  $Rf_X^! \mathbb{F}_\ell \in D_{\text{ét}}(Y' \times_Y X, \mathbb{F}_\ell)$  is invertible.*

*Proof.* This follows from the equivalence of (ii) and (iii) in Proposition 23.4 and Proposition 23.7.  $\square$

One can deduce that cohomologically smooth maps are universally open.

**Proposition 23.11.** *Let  $f : Y' \rightarrow Y$  be separated map of small  $v$ -stacks that is representable in locally spatial diamonds and  $\ell$ -cohomologically smooth for some  $\ell \neq p$ . Then  $f$  is universally open, i.e. for all  $X \rightarrow Y$ , the map  $|Y' \times_Y X| \rightarrow |X|$  is open.*

*Proof.* We can assume that  $X$  is strictly totally disconnected, and  $Y'$  is spatial; it is enough to see that the image of  $|Y'| \rightarrow |X|$  is open. As the image is generalizing, it is enough to see that the image is constructible. Consider  $Rf_! Rf^! \mathbb{F}_\ell \in D_{\text{ét}}(X, \mathbb{F}_\ell)$ . As  $f$  is  $\ell$ -cohomologically smooth,  $Rf^! \mathbb{F}_\ell$  is invertible, and in particular constructible, and thus by Proposition 23.10 (ii), one sees that  $Rf_! Rf^! \mathbb{F}_\ell$  is constructible. In particular, its support is a constructible subset of  $X$ . We claim that the support of  $Rf_! Rf^! \mathbb{F}_\ell$  agrees with the image of  $|Y'| \rightarrow |X|$ . For this, we can assume  $X = \text{Spa}(C, C^+)$  is strictly

local (using that by Proposition 23.4 and Proposition 23.10,  $Rf^!\mathbb{F}_\ell$  commutes with quasi-pro-étale base change). If the closed point  $s \in |X|$  is not in the image of  $f$ , then  $Rf^!\mathbb{F}_\ell$  is concentrated on the preimage on  $X \setminus \{s\}$ , which implies the same for  $Rf_!Rf^!\mathbb{F}_\ell$  (by commuting  $Rf_!$  with  $j_!$ ). Now assume  $s$  lies in the image of  $f$ , so  $f$  is surjective. Let  $j : U = X \setminus \{s\} \hookrightarrow X$  and  $\tilde{j} : V = f^{-1}(U) \hookrightarrow Y'$  be the open immersions, and  $f_U : V \rightarrow U$  the pullback of  $f$ .

We have

$$Rf_!Rf^!j_!\mathbb{F}_\ell = Rf_!\tilde{j}_!Rf_U^!\mathbb{F}_\ell = j_!Rf_{U!}Rf_U^!\mathbb{F}_\ell = j_!j^*(Rf_!Rf^!\mathbb{F}_\ell) ,$$

using Proposition 23.10 (iii) in the first equality. In particular, if we define  $\mathbb{F}_\ell|_s$  as the cone of  $j_!\mathbb{F}_\ell \rightarrow \mathbb{F}_\ell$ , one finds that  $Rf_!Rf^!\mathbb{F}_\ell|_s$  is concentrated at  $s$ , with stalk equal to the stalk of  $Rf_!Rf^!\mathbb{F}_\ell$ . Now note that

$$\mathrm{Hom}_{D_{\text{ét}}(X, \mathbb{F}_\ell)}(Rf_!Rf^!\mathbb{F}_\ell|_s, \mathbb{F}_\ell) = \mathrm{Hom}_{D_{\text{ét}}(Y', \mathbb{F}_\ell)}(Rf^!\mathbb{F}_\ell|_s, Rf^!\mathbb{F}_\ell) ,$$

and  $Rf^!\mathbb{F}_\ell|_s$  is the restriction  $Rf^!\mathbb{F}_\ell|_{f^{-1}(s)}$  of the invertible sheaf  $Rf^!\mathbb{F}_\ell$  to  $f^{-1}(s)$ . In particular, there is a natural map  $Rf^!\mathbb{F}_\ell|_s \rightarrow Rf^!\mathbb{F}_\ell$ , which is nonzero as soon as the fibre  $f^{-1}(s)$  is nonempty. Thus, one has  $Rf_!Rf^!\mathbb{F}_\ell|_s \neq 0$ , and therefore  $s$  lies in the support of  $Rf_!Rf^!\mathbb{F}_\ell$ .  $\square$

For cohomologically smooth morphisms, the functor  $Rf^!$  commutes with any base change, and has a very simple form.

**Proposition 23.12.** *Let  $f : Y' \rightarrow Y$  be a separated map of small  $v$ -stacks that is representable in locally spatial diamonds such that  $f$  is  $\ell$ -cohomologically smooth, and let  $\Lambda$  be an  $\ell$ -power-torsion ring. Then the natural transformation*

$$Rf^!\Lambda \otimes_\Lambda f^* \rightarrow Rf^! : D_{\text{ét}}(Y, \Lambda) \rightarrow D_{\text{ét}}(Y', \Lambda)$$

is an equivalence, and the dualizing complex  $D_f := Rf^!\Lambda \in D_{\text{ét}}(Y', \Lambda)$  is invertible, in fact  $v$ -locally isomorphic to  $\Lambda[n]$  for some integer  $n \in \mathbb{Z}$ .

Moreover, if  $g : \tilde{Y} \rightarrow Y$  is a map of small  $v$ -sheaves with base change

$$\begin{array}{ccc} \tilde{Y}' & \xrightarrow{g'} & Y' \\ \downarrow \tilde{f} & & \downarrow f \\ \tilde{Y} & \xrightarrow{g} & Y \end{array} ,$$

then the natural transformation of functors

$$g'^*Rf^! \rightarrow R\tilde{f}^!g^* : D_{\text{ét}}(Y, \Lambda) \rightarrow D_{\text{ét}}(\tilde{Y}', \Lambda)$$

adjoint to

$$R\tilde{f}_!g'^*Rf^! = g^*Rf_!Rf^! \rightarrow g^*$$

is an equivalence. In particular, the dualizing complex  $D_f$  commutes with base change, i.e.

$$g'^*D_f = D_{\tilde{f}} .$$

*Proof.* By Proposition 23.4, we know that if  $Y = X$  is strictly totally disconnected, then

$$Rf^!\Lambda \otimes_\Lambda f^* \rightarrow Rf^!$$

is an equivalence. Moreover, we see that  $Rf^!\Lambda \otimes_{\Lambda} \mathbb{F}_{\ell} = Rf^!\mathbb{F}_{\ell}$  is invertible, which implies formally that  $Rf^!\Lambda$  is invertible (as it is an extension of  $m$  copies of  $\mathbb{F}_{\ell}[n]$ ). Thus, the first part holds true in case  $Y$  is a strictly totally disconnected perfectoid space.

Next, we check that the base change property holds true in case  $Y = X$  and  $\tilde{Y} = \tilde{X}$  are both strictly totally disconnected perfectoid spaces. By the first part, it suffices to check that  $g'^*D_f = D_{\tilde{f}}$ . Using part (iv) of Proposition 23.4, we can assume that  $X$  and  $\tilde{X}$  are connected, and in fact we can assume that  $X = \mathrm{Spa}(C, \mathcal{O}_C)$  and  $\tilde{X} = \mathrm{Spa}(\tilde{C}, \mathcal{O}_{\tilde{C}})$ , using that  $Rf^!$  and  $g'^*$  commute with  $j_*$  for quasicompact open immersions. Also, the equality  $g'^*D_f = D_{\tilde{f}}$  can be checked locally on  $Y'$ , so we can assume that  $Y'$  is spatial.

Thus, assume that  $X = \mathrm{Spa}(C, \mathcal{O}_C)$  and  $\tilde{X} = \mathrm{Spa}(\tilde{C}, \mathcal{O}_{\tilde{C}})$ , and that  $Y'$  is spatial. First, note that by assumption  $Rf^!\mathbb{F}_{\ell}$  is invertible and in particular constructible, so by Proposition 23.7,  $Rf_!Rf^!\mathbb{F}_{\ell} \in D_{\acute{e}t}(X, \mathbb{F}_{\ell}) = D(\mathbb{F}_{\ell})$  is constructible, i.e. a bounded complex of finite-dimensional vector spaces. Thus,  $\mathrm{Hom}_X(Rf_!Rf^!\mathbb{F}_{\ell}, \mathbb{F}_{\ell}) = \mathrm{Hom}_{Y'}(Rf^!\mathbb{F}_{\ell}, Rf^!\mathbb{F}_{\ell}) = \mathrm{Hom}_{Y'}(\mathbb{F}_{\ell}, \mathbb{F}_{\ell})$  is finite-dimensional, which implies that  $\pi_0 Y'$  is finite. Passing to a connected component, we can thus assume that  $Y'$  is connected. In that case,  $\tilde{Y}'$  is still connected by Lemma 14.6, and so both  $D_f$  and  $D_{\tilde{f}}$  are isomorphic to  $\mathbb{L}[n]$  resp.  $\tilde{\mathbb{L}}[\tilde{n}]$  for some  $\mathbb{F}_{\ell}$ -local systems  $\mathbb{L}$  on  $Y'$  resp.  $\tilde{\mathbb{L}}$  on  $\tilde{Y}'$ , and integers  $n$  resp.  $\tilde{n}$ . The map  $g'^*D_f \rightarrow D_{\tilde{f}}$  is given by a map

$$g'^*\mathbb{L}[n] \rightarrow \tilde{\mathbb{L}}[\tilde{n}] ,$$

such that for any  $K \in D_{\acute{e}t}(Y', \mathbb{F}_{\ell})$ , one has a commutative diagram

$$\begin{array}{ccccc} \mathrm{Hom}_{D_{\acute{e}t}(Y', \mathbb{F}_{\ell})}(K, \mathbb{L}[n]) & \longrightarrow & \mathrm{Hom}_{D_{\acute{e}t}(\tilde{Y}', \mathbb{F}_{\ell})}(g'^*K, g'^*\mathbb{L}[n]) & \longrightarrow & \mathrm{Hom}_{D_{\acute{e}t}(\tilde{Y}', \mathbb{F}_{\ell})}(g'^*K, \tilde{\mathbb{L}}[\tilde{n}]) \\ \downarrow \cong & & & & \downarrow \cong \\ \mathrm{Hom}_{D_{\acute{e}t}(X, \mathbb{F}_{\ell})}(Rf_!K, \mathbb{F}_{\ell}) & \xrightarrow{\cong} & \mathrm{Hom}_{D_{\acute{e}t}(\tilde{X}, \mathbb{F}_{\ell})}(g^*Rf_!K, \mathbb{F}_{\ell}) & \xrightarrow{\cong} & \mathrm{Hom}_{D_{\acute{e}t}(\tilde{X}, \mathbb{F}_{\ell})}(R\tilde{f}_!g'^*K, \mathbb{F}_{\ell}) . \end{array}$$

If we apply this to  $K = \mathbb{L}[n]$ , then by invariance of étale cohomology under change of algebraically closed base field, the upper left arrow is the isomorphism  $R\Gamma(Y'_{\acute{e}t}, \mathbb{F}_{\ell}) \cong R\Gamma(\tilde{Y}'_{\acute{e}t}, \mathbb{F}_{\ell})$ , which is nonzero. Then the diagram implies that the map  $g'^*\mathbb{L}[n] \rightarrow \tilde{\mathbb{L}}[\tilde{n}]$  in  $D_{\acute{e}t}(\tilde{Y}', \mathbb{F}_{\ell})$  cannot be the zero map. This implies that  $\tilde{n} \geq n$ , and that it is an isomorphism if  $\tilde{n} = n$ .

Thus, if for any  $\tilde{C}/C$  as above, one has  $\tilde{n} = n$ , then it follows that base change holds. In general,  $\tilde{n} \leq 3 \dim. \mathrm{trg} f$  is bounded in terms of  $f$ , so at most finitely many values appear, and we can choose some  $\tilde{C}/C$  achieving the maximal value. It follows that  $\tilde{f} : \tilde{Y}' \rightarrow \tilde{X} = \mathrm{Spa}(\tilde{C}, \mathcal{O}_{\tilde{C}})$  has the property that  $D_{\tilde{f}}$  commutes with any base change to a strictly totally disconnected perfectoid space over  $\tilde{X}$ .

Now choose a  $v$ -hypercover of  $X = \mathrm{Spa}(C, \mathcal{O}_C)$  by strictly totally disconnected perfectoid spaces  $\tilde{X}_{\bullet} \rightarrow X$ , with  $\tilde{X}_0 = \tilde{X}$ , and let  $\tilde{f}_{\bullet} : \tilde{Y}'_{\bullet} \rightarrow \tilde{X}_{\bullet}$  be the pullback. Then we have the derived category  $D(\tilde{X}_{\bullet}, \Lambda)$  of sheaves of  $\Lambda$ -modules on the simplicial space  $\tilde{X}_{\bullet}$ , and the full subcategory  $D_{\mathrm{cart}, \acute{e}t}(\tilde{X}_{\bullet}, \Lambda) \subset D(\tilde{X}_{\bullet}, \Lambda)$  where all pullback maps are isomorphisms, and all terms lie in  $D_{\acute{e}t}(\tilde{X}_i, \Lambda)$ . By Proposition 17.3, one has an equivalence of categories

$$D_{\acute{e}t}(X, \Lambda) \cong D_{\mathrm{cart}, \acute{e}t}(\tilde{X}_{\bullet}, \Lambda) .$$

Similarly,

$$D_{\text{ét}}(Y', \Lambda) \cong D_{\text{cart,ét}}(\tilde{Y}'_{\bullet}, \Lambda) .$$

Moreover, the functor  $Rf_! : D_{\text{ét}}(Y', \Lambda) \rightarrow D_{\text{ét}}(X, \Lambda)$  gets identified with the functor

$$R\tilde{f}_{\bullet} : D_{\text{cart,ét}}(\tilde{Y}'_{\bullet}, \Lambda) \rightarrow D_{\text{cart,ét}}(\tilde{X}_{\bullet}, \Lambda)$$

which is termwise given by  $R\tilde{f}_{i!}$ . It preserves the “cartesian” condition by proper base change.

Now note that all terms  $R\tilde{f}_{i!}$  admit right adjoints  $R\tilde{f}_i^!$ , and these right adjoints preserve the “cartesian” condition by the base change already established. This implies that the right adjoint to

$$R\tilde{f}_{\bullet} : D_{\text{cart,ét}}(\tilde{Y}'_{\bullet}, \Lambda) \rightarrow D_{\text{cart,ét}}(\tilde{X}_{\bullet}, \Lambda)$$

is given by the functor

$$R\tilde{f}_{\bullet}^! : D_{\text{cart,ét}}(\tilde{X}_{\bullet}, \Lambda) \rightarrow D_{\text{cart,ét}}(\tilde{Y}'_{\bullet}, \Lambda)$$

that is termwise  $R\tilde{f}_i^!$ . This gives a commutative diagram

$$\begin{array}{ccc} D_{\text{ét}}(X, \Lambda) & \xrightarrow{Rf^!} & D_{\text{ét}}(Y', \Lambda) \\ \downarrow \cong & & \downarrow \cong \\ D_{\text{cart,ét}}(\tilde{X}_{\bullet}, \Lambda) & \xrightarrow{R\tilde{f}_{\bullet}^!} & D_{\text{cart,ét}}(\tilde{Y}'_{\bullet}, \Lambda) \\ \downarrow & & \downarrow \\ D_{\text{ét}}(\tilde{X}, \Lambda) & \xrightarrow{R\tilde{f}^!} & D_{\text{ét}}(\tilde{Y}', \Lambda) , \end{array}$$

where the vertical maps are given by pullback. This gives the desired base change result along  $\text{Spa}(\tilde{C}, \mathcal{O}_{\tilde{C}}) \rightarrow \text{Spa}(C, \mathcal{O}_C)$  in general, which finishes the proof of the base change result for  $Rf^!$  along maps of strictly totally disconnected perfectoid spaces.

In the general case, we can now find a v-hypercover  $\tilde{X}_{\bullet} \rightarrow Y$  of  $Y$  such that all  $\tilde{X}_i$  are disjoint unions of strictly totally disconnected perfectoid spaces. Applying the previous discussion again, we get the desired base change for  $Rf^!$  along the map  $\tilde{X}_0 \rightarrow Y$ ; this implies also that  $Rf^! \Lambda \otimes_{\Lambda} f^* \rightarrow Rf^!$  is an equivalence, as this can now be checked after base change to  $\tilde{X}_0$ , where we know it.

Now, for the general base change identity, we can cover any map  $\tilde{Y} \rightarrow Y$  by a map between disjoint unions of strictly totally disconnected perfectoid spaces  $\tilde{X} \rightarrow X$ ; we know base change for  $\tilde{X} \rightarrow \tilde{Y}$ ,  $X \rightarrow Y$  and  $\tilde{X} \rightarrow X$ , which implies it for  $\tilde{Y} \rightarrow Y$ .  $\square$

Using Proposition 23.12, one can show that composites of cohomologically smooth morphisms are cohomologically smooth.

**Proposition 23.13.** *Let  $g : Y'' \rightarrow Y'$  and  $f : Y' \rightarrow Y$  be separated morphisms of small v-stacks and let  $\ell \neq p$  be a prime. If  $f$  and  $g$  are representable in locally spatial diamonds and  $\ell$ -cohomologically smooth, then  $f \circ g$  is representable in locally spatial diamonds and  $\ell$ -cohomologically smooth. Conversely, if  $g$  and  $f \circ g$  are representable in locally spatial diamonds and  $\ell$ -cohomologically smooth, and  $g$  is surjective, then  $f$  is representable in locally spatial diamonds and  $\ell$ -cohomologically smooth.*

**Remark 23.14.** The first part of the proof shows more generally that if  $f : Y' \rightarrow Y$  is a separated 0-truncated map of small v-stacks and  $g : Y'' \rightarrow Y'$  is a universally open (e.g.,  $\ell$ -cohomologically smooth), separated and surjective map of small v-stacks such that  $f \circ g$  is representable in locally spatial diamonds, then  $f$  is representable in locally spatial diamonds.

*Proof.* The first part follows from Proposition 23.12. For the second part, we check first that  $f$  is representable in locally spatial diamonds. By Proposition 13.4, we can check this after pullback to a strictly totally disconnected  $Y$ . In that case,  $Y''$  is separated and locally spatial over  $Y$ . For any quasicompact open subspace  $V \subset Y''$ , the image  $U \subset Y'$  is open by Proposition 23.11. Moreover, it is quasicompact, as  $V$  surjects onto  $U$ . Also,  $|U|$  is the quotient of the spectral space  $|V|$  by an open qcqs equivalence relation (given by the image of  $|V \times_U V|$ ), so  $|U|$  is spectral and  $|V| \rightarrow |U|$  is a spectral map by Lemma 2.10. Thus,  $U$  is spatial, and  $Y'$  is locally spatial.

One easily checks  $\dim.\mathrm{trg} f \leq \dim.\mathrm{trg}(f \circ g) < \infty$  (where one uses the modified  $\widetilde{\mathrm{tr}.c}$  in the definition of  $\dim.\mathrm{trg}$ ), and  $f$  is compactifiable by Proposition 22.3. We may assume  $Y = X$  is strictly totally disconnected, and we have to check that

$$Rf^! \mathbb{F}_\ell \otimes_{\mathbb{F}_\ell} f^* \rightarrow Rf^!$$

is an equivalence, and that  $Rf^! \mathbb{F}_\ell$  is invertible. But as  $f \circ g$  is  $\ell$ -cohomologically smooth,

$$R(f \circ g)^! \mathbb{F}_\ell = Rg^!(Rf^! \mathbb{F}_\ell) = Rg^! \mathbb{F}_\ell \otimes_{\mathbb{F}_\ell} g^* Rf^! \mathbb{F}_\ell$$

is invertible, and  $Rg^! \mathbb{F}_\ell$  is invertible as  $g$  is  $\ell$ -cohomologically smooth; this implies that  $g^* Rf^! \mathbb{F}_\ell$  is invertible, which shows that  $Rf^! \mathbb{F}_\ell$  is invertible. Similarly, to check whether  $Rf^! \mathbb{F}_\ell \otimes_{\mathbb{F}_\ell} f^* \rightarrow Rf^!$  is an equivalence, we can check after applying  $Rg^! = Rg^! \mathbb{F}_\ell \otimes_{\mathbb{F}_\ell} g^*$ . But

$$Rg^! Rf^! = R(f \circ g)^! = R(f \circ g)^! \mathbb{F}_\ell \otimes_{\mathbb{F}_\ell} (f \circ g)^* = Rg^! \mathbb{F}_\ell \otimes_{\mathbb{F}_\ell} g^* Rf^! \mathbb{F}_\ell \otimes_{\mathbb{F}_\ell} g^* f^* = Rg^!(Rf^! \mathbb{F}_\ell \otimes_{\mathbb{F}_\ell} f^*),$$

as desired.  $\square$

Also, we can check cohomological smoothness v-locally on the target.

**Proposition 23.15.** *Let  $f : Y' \rightarrow Y$  be a separated map of small v-stacks that is representable in locally spatial diamonds, and let  $\ell \neq p$  be a prime. Let  $g : \tilde{Y} \rightarrow Y$  be map of small v-stacks with pullback  $\tilde{f} : \tilde{Y}' = Y' \times_Y \tilde{Y} \rightarrow \tilde{Y}$ .*

*If  $f$  is  $\ell$ -cohomologically smooth, then  $\tilde{f}$  is  $\ell$ -cohomologically smooth. Conversely, if  $\tilde{f}$  is  $\ell$ -cohomologically smooth,  $g$  is a surjective map of v-sheaves, and  $\dim.\mathrm{trg} f < \infty$ , then  $f$  is  $\ell$ -cohomologically smooth.*

It is not clear to us whether the condition  $\dim.\mathrm{trg} f < \infty$  can be checked v-locally on the target.

*Proof.* The first statement is clear from the definition. For the converse, we can assume that  $Y = X$  and  $\tilde{Y} = \tilde{X}$  are strictly totally disconnected. Then it follows from the argument involving a simplicial v-hypercover  $\tilde{X}_\bullet \rightarrow X$  with  $\tilde{X}_0 = \tilde{X}$  in the proof of Proposition 23.12.  $\square$

Finally, let us note that we get the expected smooth base change results.

**Proposition 23.16.** *Let*

$$\begin{array}{ccc} Y' & \xrightarrow{\tilde{g}} & Y \\ \downarrow f' & & \downarrow f \\ X' & \xrightarrow{g} & X \end{array}$$

be a cartesian diagram of small  $v$ -stacks, and assume that  $n\Lambda = 0$  for some  $n$  prime to  $p$ .

(i) Assume that  $g$  is compactifiable and representable in locally spatial diamonds with  $\dim. \operatorname{trg} f < \infty$ , and  $A \in D_{\text{ét}}(Y, \Lambda)$ . Then

$$Rg^! Rf_* A \cong Rf'_* \widetilde{Rg}^! A .$$

(ii) Assume that  $g$  is separated, representable in locally spatial diamonds and  $\ell$ -cohomologically smooth, and  $\Lambda$  is  $\ell$ -power torsion. Then for all  $A \in D_{\text{ét}}(Y, \Lambda)$ , the base change morphism

$$g^* Rf_* A \rightarrow Rf'_* \widetilde{g}^* A$$

is an isomorphism.

*Proof.* Part (i) follows from Proposition 22.19 by passing to right adjoints. Now part (ii) follows from part (i) and Proposition 23.12.  $\square$

**Proposition 23.17.** *Let  $f : Y \rightarrow X$  be a separated map of small  $v$ -stacks that is representable in locally spatial diamonds and  $\ell$ -cohomologically smooth. There is a functorial isomorphism*

$$f^* R\mathcal{H}om(A, B) \cong R\mathcal{H}om(f^* A, f^* B)$$

in  $A, B \in D_{\text{ét}}(X, \Lambda)$ .

*Proof.* This follows from Proposition 23.3 (ii) together with the identification  $Rf^! = f^* \otimes_{\Lambda} Rf^! \Lambda$ , where  $Rf^! \Lambda$  is invertible.  $\square$

## 24. EXAMPLES OF SMOOTH MORPHISMS

In the previous section, we have defined cohomologically smooth morphisms, and checked that in this case  $!$ -pullback takes a very simple form. Moreover, the class of cohomologically smooth morphisms has good stability properties. However, we have not seen any example of cohomologically smooth morphisms. In this section, we will give criteria guaranteeing cohomological smoothness.

We start with the following important theorem, which uses the main results of Huber's book.

**Theorem 24.1.** *Let  $f : \mathbb{B} \rightarrow *$  be the projection from the ball to the point. Then  $f$  is  $\ell$ -cohomologically smooth for any prime  $\ell \neq p$ , and for any ring  $\Lambda$  with  $n\Lambda = 0$  for some  $n$  prime to  $p$ , one has a canonical isomorphism*

$$D_f = Rf^! \Lambda \cong \Lambda(1)[2] .$$

*Proof.* Fix a prime  $\ell \neq p$ . We check the conditions of Proposition 23.10. Condition (i) is easy (in fact,  $\dim. \operatorname{trg} f = 1$ ). For condition (ii), let  $X$  be a strictly totally disconnected perfectoid space, and let  $\mathcal{F}$  be a constructible  $\ell$ -torsion sheaf on  $\mathbb{B} \times X$ . We can find a map  $X \rightarrow \operatorname{Spa}(K, \mathcal{O}_K)$ , where  $K$  is a perfectoid field (given by  $\mathbb{F}_p((\varpi^{1/p^\infty}))$  for a pseudo-uniformizer  $\varpi$ ). Then we can write  $X$  as a cofiltered inverse limit of spaces  $X_i \rightarrow \operatorname{Spa}(K, \mathcal{O}_K)$  which are topologically of finite type, cf. [Sch12, Lemma 6.13], and  $\mathcal{F}$  is the preimage of some constructible sheaf  $\mathcal{F}_i$  on  $\mathbb{B} \times X_i$  for  $i$  large enough by Proposition 20.7. Consider the cartesian diagram

$$\begin{array}{ccc} \mathbb{B} \times X & \xrightarrow{h} & \mathbb{B} \times X_i \\ \downarrow f_X & & \downarrow f_{X_i} \\ X & \xrightarrow{g} & X_i . \end{array}$$

Then for all  $j \geq 0$ , one has

$$R^j f_{X!} \mathcal{F} = R^j f_{X!} h^* \mathcal{F}_i = g^* R^j f_{X_i!} \mathcal{F}_i .$$

By [Hub96, Theorem 6.2.2], the sheaf  $R^j f_{X_i!} \mathcal{F}_i$  is constructible on  $X_i$ , and thus its pullback along  $g^*$  is a constructible sheaf on  $X$ , as desired.

Condition (iii) is a condition about  $f_X : \mathbb{B} \times X \rightarrow X$  in case  $X = \mathrm{Spa}(C, C^+)$  is strongly Noetherian. Thus, Huber's results and in particular [Hub96, Theorem 7.5.3] applies.

For condition (iv), take again a strictly totally disconnected perfectoid space  $X$ . We can in fact find a map  $X \rightarrow \mathrm{Spa}(C, \mathcal{O}_C)$ , where  $C$  is algebraically closed: As above, we have a map  $X \rightarrow \mathrm{Spa}(K, \mathcal{O}_K)$  where  $K = \mathbb{F}_p((\varpi^{1/p^\infty}))$ , but in fact this factors over  $\mathrm{Spa}(C, \mathcal{O}_C)$ , where  $C$  is the completed algebraic closure of  $K$ , as  $X$  has no nonsplit finite étale covers. Consider the cartesian diagram

$$\begin{array}{ccc} \mathbb{B} \times X & \xrightarrow{h} & \mathbb{B} \times \mathrm{Spa}(C, \mathcal{O}_C) \\ \downarrow f_X & & \downarrow f_C \\ X & \xrightarrow{g} & \mathrm{Spa}(C, \mathcal{O}_C) . \end{array}$$

There is a trace map

$$Rf_{C!} \mathbb{F}_\ell(1) \rightarrow \mathbb{F}_\ell[-2]$$

by [Hub96, Theorem 7.2.2] (noting that  $R^i f_{C!} \mathbb{F}_\ell(1) = 0$  for  $i > 2$  by [Hub96, Proposition 5.5.8]). By base change, we get a trace map

$$Rf_{X!} \mathbb{F}_\ell(1) = g^* Rf_{C!} \mathbb{F}_\ell(1) \rightarrow \mathbb{F}_\ell[-2] ,$$

which by adjunction gives a map

$$\alpha : \mathbb{F}_\ell(1)[2] \rightarrow Rf_X^! \mathbb{F}_\ell .$$

We claim that  $\alpha$  is an isomorphism. Note that we have already proved that condition (iii) of Proposition 23.4 is satisfied, so also condition (iv) is satisfied. In particular,  $Rf_X^! \mathbb{F}_\ell$  commutes with quasi-pro-étale base change, and so we can check whether  $\alpha$  is an equivalence after pullback to connected components of  $X$ . Thus, we can assume  $X = \mathrm{Spa}(C, C^+)$ . In this case, the result follows from [Hub96, Theorem 7.5.3].  $\square$

We also need the following result.

**Proposition 24.2.** *Let  $f : Y' \rightarrow Y$  be a separated map of small  $v$ -stacks that is representable in locally spatial diamonds and  $\ell$ -cohomologically smooth for some prime  $\ell \neq p$ . Assume that there is a profinite group  $K$  of pro-order prime to  $\ell$  with a free action  $\underline{K} \times Y' \rightarrow Y'$  over  $Y$ , i.e. such that  $\underline{K} \times Y' \rightarrow Y' \times_Y Y'$  is an injection.*

*Then the quotient map  $f/\underline{K} : Y'/\underline{K} \rightarrow Y$  is separated, representable in locally spatial diamonds, and  $\ell$ -cohomologically smooth. Moreover, if we fix the  $\Lambda$ -valued Haar measure on  $K$  with total volume 1, there is a natural equivalence of functors*

$$Rf^! = q^* R(f/\underline{K})^! : D_{\text{ét}}(Y, \Lambda) \rightarrow D_{\text{ét}}(Y', \Lambda) ,$$

where  $q : Y' \rightarrow Y'/\underline{K}$  denotes the quotient map.



One also has the identity

$$Rf^! = Rq^!R(f/\underline{K})^! .$$

However, it is not true that  $Rq^! = q^*$ . For example, after a base change,  $q$  becomes  $\underline{S} \times \mathrm{Spa}(C, \mathcal{O}_C) \rightarrow \mathrm{Spa}(C, \mathcal{O}_C)$  for some profinite set  $S$  and algebraically closed perfectoid field  $C$ . In this case,  $Rq^!\Lambda$  can be identified with the sheaf sending any open and closed subset  $T \subset S$  to the distributions  $\mathrm{Hom}(C^0(T, \Lambda), \Lambda)$ , as follows easily from the adjunction defining  $Rq^!$ . Under the choice of a Haar measure, we get however a natural map  $q^* \rightarrow Rq^!$ , as in the proof below.

*Proof.* We can assume that  $Y = X$  is a strictly totally disconnected perfectoid space. Then  $Y'/\underline{K}$  is locally spatial (and quasiseparated) by Lemma 2.10 and Lemma 10.13. It is also separated by the valuative criterion. Its canonical compactification  $\overline{Y'}/\underline{K}^{/Y}$  is given by  $\overline{Y'}^{/Y}/\underline{K}$ . To check whether the inclusion is an open immersion, one uses that  $\overline{Y'}^{/Y} \rightarrow \overline{Y'}^{/Y}/\underline{K}$  is a quotient map.

It remains to see that  $f/\underline{K}$  is  $\ell$ -cohomologically smooth. It is clear that  $\dim.\mathrm{trg} f/\underline{K} = \dim.\mathrm{trg} f < \infty$ . Now, fixing the  $\Lambda$ -valued Haar measure on  $K$  with total volume 1, we first construct a natural transformation

$$q^* \rightarrow Rq^! : D_{\acute{\mathrm{e}}\mathrm{t}}(Y'/\underline{K}, \Lambda) \rightarrow D_{\acute{\mathrm{e}}\mathrm{t}}(Y', \Lambda) .$$

This is adjoint to a map  $Rq_!q^* = Rq_*q^* = q_*q^* \rightarrow \mathrm{id}$ . For any open subgroup  $H \subset K$ , consider the projection  $q_{H,K} : Y'/\underline{H} \rightarrow Y'/\underline{K}$ . Then  $q_*q^*$  is the filtered colimit of  $q_{H,K}q_{H,K}^*$ , and there are natural trace maps  $q_{H,K}q_{H,K}^* \rightarrow \mathrm{id}$ , which one can divide by  $[K : H]$  to make them compatible. (Note that here, we are implicitly using the choice of the Haar measure.) This gives the desired natural transformation  $q_*q^* \rightarrow \mathrm{id}$ , and thus  $q^* \rightarrow Rq^!$ .

In particular, we get a natural transformation

$$q^*R(f/\underline{K})^! \rightarrow Rq^!R(f/\underline{K})^! = Rf^! .$$

We claim that this is an equivalence. It suffices to check this for  $\Lambda = \mathbb{F}_\ell$ . Note that once this is known, it follows that  $f/\underline{K}$  satisfies condition (iii) of Proposition 23.4, and  $R(f/\underline{K})^!\mathbb{F}_\ell$  is invertible, which shows that  $f/\underline{K}$  is  $\ell$ -cohomologically smooth.

Thus, it remains to check that

$$q^*R(f/\underline{K})^! \rightarrow Rq^!R(f/\underline{K})^! = Rf^! : D_{\acute{\mathrm{e}}\mathrm{t}}(X, \mathbb{F}_\ell) \rightarrow D_{\acute{\mathrm{e}}\mathrm{t}}(Y', \mathbb{F}_\ell)$$

is an equivalence. This can be checked locally on  $Y'$ , so we can assume that  $Y'$  is spatial. By Proposition 20.10, it suffices to check that for any constructible sheaf  $\mathcal{F}$  on  $Y'$  and any  $A \in D_{\acute{\mathrm{e}}\mathrm{t}}(X, \mathbb{F}_\ell)$ , one has

$$\mathrm{Hom}_{D_{\acute{\mathrm{e}}\mathrm{t}}(Y', \mathbb{F}_\ell)}(\mathcal{F}, q^*R(f/\underline{K})^!A) = \mathrm{Hom}_{D_{\acute{\mathrm{e}}\mathrm{t}}(Y', \mathbb{F}_\ell)}(\mathcal{F}, Rf^!A) .$$

Let  $q_H : Y' \rightarrow Y'/\underline{H}$  be the projection. By Proposition 20.7, we can assume that  $\mathcal{F} = q_H^* \mathcal{F}_H$  for some constructible sheaf  $\mathcal{F}_H$  on  $Y'/\underline{H}$ . Then

$$\begin{aligned}
\mathrm{Hom}_{D_{\acute{e}t}(Y', \mathbb{F}_\ell)}(\mathcal{F}, q^* R(f/\underline{K})^! A) &= \mathrm{Hom}_{D_{\acute{e}t}(Y', \mathbb{F}_\ell)}(q_H^* \mathcal{F}, q^* R(f/\underline{K})^! A) \\
&= \mathrm{Hom}_{D_{\acute{e}t}(Y'/\underline{H}, \mathbb{F}_\ell)}(\mathcal{F}_H, q_{H*} q^* R(f/\underline{K})^! A) \\
&= \mathrm{Hom}_{D_{\acute{e}t}(Y'/\underline{H}, \mathbb{F}_\ell)}(\mathcal{F}_H, \varinjlim_{H' \subset H} q_{H', H*} q_{H', K}^* R(f/\underline{K})^! A) \\
&= \varinjlim_{H' \subset H} \mathrm{Hom}_{D_{\acute{e}t}(Y'/\underline{H}, \mathbb{F}_\ell)}(\mathcal{F}_H, q_{H', H*} q_{H', K}^* R(f/\underline{K})^! A) \\
&= \varinjlim_{H' \subset H} \mathrm{Hom}_{D_{\acute{e}t}(Y'/\underline{H}', \mathbb{F}_\ell)}(q_{H', H}^* \mathcal{F}_H, q_{H', K}^* R(f/\underline{K})^! A) \\
&= \varinjlim_{H' \subset H} \mathrm{Hom}_{D_{\acute{e}t}(Y'/\underline{H}', \mathbb{F}_\ell)}(q_{H', H}^* \mathcal{F}_H, Rq_{H', K}^! R(f/\underline{K})^! A) \\
&= \varinjlim_{H' \subset H} \mathrm{Hom}_{D_{\acute{e}t}(Y'/\underline{H}', \mathbb{F}_\ell)}(q_{H', H}^* \mathcal{F}_H, R(f/\underline{H}')^! A) \\
&= \varinjlim_{H' \subset H} \mathrm{Hom}_{D_{\acute{e}t}(X, \mathbb{F}_\ell)}(R(f/\underline{H}')^! q_{H', H}^* \mathcal{F}_H, A) .
\end{aligned}$$

Let  $\mathcal{F}_{H'} = q_{H', H}^* \mathcal{F}_H$ . We claim that the filtered direct system

$$R(f/\underline{H}')^! \mathcal{F}_{H'} \in D_{\acute{e}t}(X, \mathbb{F}_\ell)$$

is eventually constant, i.e. for all sufficiently small  $H'$ , all transition maps are isomorphisms. For this, note first that all transition maps are split injective (using the trace maps as above). On the other hand,  $Rf_! \mathcal{F} \in D_{\acute{e}t}(X, \mathbb{F}_\ell)$  is bounded with constructible cohomology sheaves (i.e., compact), and

$$\begin{aligned}
Rf_! \mathcal{F} &= Rf_! q_H^* \mathcal{F} = R(f/\underline{H})^! Rq_H^! q_H^* \mathcal{F} \\
&= R(f/\underline{H})^! q_{H*} q_H^* \mathcal{F} \\
&= R(f/\underline{H})^! \varinjlim_{H' \subset H} q_{H', H*} q_{H', H}^* \mathcal{F} \\
&= \varinjlim_{H' \subset H} R(f/\underline{H})^! q_{H', H*} \mathcal{F}_{H'} \\
&= \varinjlim_{H' \subset H} R(f/\underline{H}')^! \mathcal{F}_{H'} .
\end{aligned}$$

The map to the direct limit factors over some term in the direct limit by compactness of  $Rf_! \mathcal{F}$ . As all transition maps are split injective, this shows that indeed the system  $R(f/\underline{H}')^! \mathcal{F}_{H'}$  is eventually constant, and eventually equal to  $Rf_! \mathcal{F}$ . Continuing the equations above, we see that

$$\begin{aligned}
\mathrm{Hom}_{D_{\acute{e}t}(Y', \mathbb{F}_\ell)}(\mathcal{F}, q^* R(f/\underline{K})^! A) &= \varinjlim_{H' \subset H} \mathrm{Hom}_{D_{\acute{e}t}(X, \mathbb{F}_\ell)}(R(f/\underline{H}')^! q_{H', H}^* \mathcal{F}_H, A) \\
&= \mathrm{Hom}_{D_{\acute{e}t}(X, \mathbb{F}_\ell)}(Rf_! \mathcal{F}, A) \\
&= \mathrm{Hom}_{D_{\acute{e}t}(Y', \mathbb{F}_\ell)}(\mathcal{F}, Rf^! A) ,
\end{aligned}$$

as desired.  $\square$

There is a variant of the proposition when the action is not free, but we know already that all fibres are smooth.

**Proposition 24.3.** *Let  $f : Y' \rightarrow Y$  be a separated map of small  $v$ -stacks that is representable in locally spatial diamonds and  $\ell$ -cohomologically smooth for some prime  $\ell \neq p$ . Assume that there is a profinite group  $K$  of pro-order prime to  $\ell$  with an action  $\underline{K} \times Y' \rightarrow Y'$  over  $Y$  for which the map  $\underline{K} \times Y' \rightarrow Y' \times_Y Y'$  is 0-truncated and qcqs.*

*Let  $Y'/\underline{K}$  be the quotient of  $Y'$  by the equivalence relation which is given as the image of  $\underline{K} \times Y' \rightarrow Y' \times_Y Y'$ . Then the quotient map  $f/\underline{K} : Y'/\underline{K} \rightarrow Y$  is separated and representable in locally spatial diamonds. Moreover, if for all complete algebraically closed fields  $C$  with an open and bounded valuation subring  $C^+ \subset C$ , and all maps  $\mathrm{Spa}(C, C^+) \rightarrow Y$ , the pullback  $Y'/\underline{K} \times_Y \mathrm{Spa}(C, C^+) \rightarrow \mathrm{Spa}(C, C^+)$  is  $\ell$ -cohomologically smooth, then  $f/\underline{K} : Y'/\underline{K} \rightarrow Y$  is  $\ell$ -cohomologically smooth, and if we fix the  $\Lambda$ -valued Haar measure on  $K$  with total volume 1, there is a natural equivalence of functors*

$$Rf^! = q^* R(f/\underline{K})^! : D_{\text{ét}}(Y, \Lambda) \rightarrow D_{\text{ét}}(Y', \Lambda) ,$$

where  $q : Y' \rightarrow Y'/\underline{K}$  denotes the quotient map.

*Proof.* First, we check that  $Y'/\underline{K} \rightarrow Y$  is compactifiable, representable in locally spatial diamonds, with  $\dim. \mathrm{trg} < \infty$ . For this, we can assume that  $Y = X$  is a strictly totally disconnected perfectoid space. Then  $Y'/\underline{K}$  is locally spatial (and quasiseparated) by Lemma 2.10 and Lemma 10.13. It is also separated by the valuative criterion. Its canonical compactification  $\overline{Y'/\underline{K}}^{/Y}$  is given by  $\overline{Y'}^{/Y}/\underline{K}$ . To check whether the inclusion is an open immersion, one uses that  $\overline{Y'}^{/Y} \rightarrow \overline{Y'}^{/Y}/\underline{K}$  is a quotient map. It is clear that  $\dim. \mathrm{trg} f/\underline{K} = \dim. \mathrm{trg} f < \infty$ .

It remains to see that  $f/\underline{K}$  is  $\ell$ -cohomologically smooth, and that  $Rf^! = q^* R(f/\underline{K})^!$ . We start by constructing a natural transformation

$$q^* \rightarrow Rq^! : D_{\text{ét}}(Y'/\underline{K}, \Lambda) \rightarrow D_{\text{ét}}(Y', \Lambda) .$$

Recall that there is always a natural transformation

$$Rq^! \Lambda \otimes_{\Lambda}^{\mathbb{L}} q^* \rightarrow Rq^! ;$$

thus, it is enough to construct a natural map  $\Lambda \rightarrow Rq^! \Lambda$ . This is adjoint to a trace map  $Rq_! \Lambda \rightarrow \Lambda$ . But note that  $q$  is proper and quasi-pro-étale, so  $Rq_! = Rq_*$ , and  $Rq_* = q_*$  is of cohomological dimension 0. Thus, it remains to construct a map of sheaves  $q_* \Lambda \rightarrow \Lambda$ . This map can be constructed  $v$ -locally, and is given by integrating over the  $K$ -action, fixing the  $\Lambda$ -valued Haar measure on  $K$  with total volume 1.

Now we prove that  $f/\underline{K}$  is  $\ell$ -cohomologically smooth. For this, we can assume that  $X$  is strictly totally disconnected, and  $Y'$  is quasicompact. We check the criteria of Proposition 23.10. Part (i) is clear. For part (ii), note that for any constructible sheaf  $\mathcal{F}$  on  $Y'/\underline{K}$ , one has natural maps

$$\mathcal{F} \rightarrow q_* q^* \mathcal{F} = Rq_! q^* \mathcal{F} \rightarrow Rq_! Rq^! \mathcal{F} \rightarrow \mathcal{F}$$

whose composite is an isomorphism. This implies that  $R(f/\underline{K})_! \mathcal{F}$  is a direct summand of  $Rf_! q^* \mathcal{F} = R(f/\underline{K})_! Rq_! q^* \mathcal{F}$ , which is thus constructible. Now part (iii) is a question about geometric fibres that holds by hypothesis. In particular, condition (iii) of Proposition 23.4 holds true, so  $R(f/\underline{K})^!$  commutes with quasi-pro-étale base change. Thus, checking whether

$$q^* R(f/\underline{K})^! \rightarrow Rf^!$$

is an equivalence can be done on geometric fibres; but here it holds by assumption. In particular, this implies that condition (iv) of Proposition 23.10 is satisfied. Thus,  $f/\underline{K}$  is  $\ell$ -cohomologically smooth; moreover, we have already checked that  $q^* R(f/\underline{K})^! \rightarrow Rf^!$  is an equivalence, as desired.  $\square$

In particular, we note that smooth morphisms of analytic adic spaces give examples of cohomologically smooth morphisms.

**Proposition 24.4.** *Let  $f : Y' \rightarrow Y$  be a separated smooth morphism of analytic adic spaces over  $\mathrm{Spa} \mathbb{Z}_p$ , i.e.  $f$  is locally on  $Y'$  the composition of an étale map  $Y' \rightarrow Y \times \mathbb{B}^n$  and the projection  $Y \times \mathbb{B}^n \rightarrow Y$ , cf. [Hub96, Corollary 1.6.10]. Then  $f^\diamond : (Y')^\diamond \rightarrow Y^\diamond$  is  $\ell$ -cohomologically smooth.*

*Proof.* By Lemma 15.6,  $f$  is representable in locally spatial diamonds. Using Proposition 22.3, one sees that  $f$  is compactifiable.

By Proposition 23.13, it suffices to check that  $Y \times \mathbb{B}^n \rightarrow Y$  is  $\ell$ -cohomologically smooth. By induction, we can assume that  $n = 1$ . As we can cover  $\mathbb{B}$  by two copies of  $\mathbb{T} \subset \mathbb{B}$  given by  $\mathbb{T}(R, R^+) = (R^+)^\times$ , it suffices to show that  $Y \times \mathbb{T} \rightarrow Y$  is  $\ell$ -cohomologically smooth. If we work over  $\mathrm{Spa} \mathbb{Q}_p$  instead, we can even base change to  $Y = \mathrm{Spa} \mathbb{C}_p$ .

In that case, there is a  $\mathbb{Z}_p$ -torsor

$$\widetilde{\mathbb{T}}_{\mathbb{C}_p} = \mathrm{Spa} \mathbb{C}_p \langle T^{\pm 1/p^\infty} \rangle \rightarrow \mathbb{T}_{\mathbb{C}_p} = \mathrm{Spa} \mathbb{C}_p \langle T^{\pm 1} \rangle ,$$

so that

$$(Y \times \mathbb{T})^\diamond = \mathbb{T}_{\mathbb{C}_p}^\diamond = \widetilde{\mathbb{T}}_{\mathbb{C}_p}^\diamond / \underline{\mathbb{Z}}_p \rightarrow Y^\diamond = (\mathrm{Spa} \mathbb{C}_p)^\diamond$$

is  $\ell$ -cohomologically smooth by Proposition 24.2 and Theorem 24.1.

On the other hand, if  $Y$  is of characteristic  $p$ , the result follows directly from Theorem 24.1. In mixed characteristic, we argue as follows. We need to see that  $Y \times \mathbb{T} \rightarrow Y$  is  $\ell$ -cohomologically smooth. We can assume, by a  $v$ -cover, that  $Y$  lives over  $\mathrm{Spa} \mathbb{Z}_p^{\mathrm{cycl}}$ . In that case,  $Y \times \mathbb{T}$  is a quotient of  $Y \times \widetilde{\mathbb{T}}$  by a nonfree  $\underline{\mathbb{Z}}_p$ -action. The result follows from Proposition 24.3, as we have already verified that all geometric fibres are  $\ell$ -cohomologically smooth.  $\square$

Another example is the following.

**Proposition 24.5.** *The map  $(\mathrm{Spa} \mathbb{Q}_p)^\diamond \rightarrow *$  is  $\ell$ -cohomologically smooth.*

*Proof.* Let  $K_\infty$  be the cyclotomic  $\mathbb{Z}_p$ -extension of  $\mathbb{Q}_p$ . Then  $K_\infty^\flat \cong \mathbb{F}_p((t^{1/p^\infty}))$ , and

$$(\mathrm{Spa} \mathbb{Q}_p)^\diamond = \mathrm{Spa} \mathbb{F}_p((t^{1/p^\infty})) / \underline{\mathbb{Z}}_p .$$

In particular, after pullback along the  $v$ -cover  $\mathrm{Spa}(C, \mathcal{O}_C) \rightarrow *$ , where  $C$  is some algebraically closed nonarchimedean field of characteristic  $p$ , one has

$$(\mathrm{Spa} \mathbb{Q}_p)^\diamond \times \mathrm{Spa}(C, \mathcal{O}_C) = \mathbb{D}_C^\times / \underline{\mathbb{Z}}_p ,$$

where  $\mathbb{D}_C^\times = \mathrm{Spa} \mathbb{F}_p((t^{1/p^\infty})) \times \mathrm{Spa}(C, \mathcal{O}_C)$  denotes the punctured open unit disc over  $\mathrm{Spa}(C, \mathcal{O}_C)$ , which is an open subset of  $\mathbb{B} \times \mathrm{Spa}(C, \mathcal{O}_C)$ . Using Proposition 24.2, this shows that  $(\mathrm{Spa} \mathbb{Q}_p)^\diamond \times \mathrm{Spa}(C, \mathcal{O}_C) \rightarrow \mathrm{Spa}(C, \mathcal{O}_C)$  is  $\ell$ -cohomologically smooth, and thus also  $(\mathrm{Spa} \mathbb{Q}_p)^\diamond \rightarrow *$ .  $\square$

Let us end with a question, which gives one way of making precise the intuition that “regular rings are smooth over an absolute base”.

**Question 24.6.** *Let  $R$  be a regular  $\mathbb{Z}_p$ -algebra that is  $I$ -adically complete for some ideal  $I$  containing  $p$ , and topologically of finite type over  $\mathbb{Z}_p$ . Is the map  $(\mathrm{Spa} R)^\diamond \rightarrow *$  cohomologically smooth?*

Here,  $(\mathrm{Spa} R)^\diamond$  is the small  $v$ -sheaf parametrizing untilts over  $R$ . Even the case  $R = \mathbb{Z}_p$  is not known to us. We note that in general, this question is related to Grothendieck’s purity conjecture (the theorem of Thomason–Gabber).

## REFERENCES

- [Bou98] Nicolas Bourbaki, *Commutative algebra. Chapters 1–7*, Elements of Mathematics (Berlin), Springer-Verlag, Berlin, 1998, Translated from the French, Reprint of the 1989 English translation. MR 1727221
- [BS15] Bhargav Bhatt and Peter Scholze, *The pro-étale topology for schemes*, Astérisque (2015), no. 369, 99–201. MR 3379634
- [Col02] Pierre Colmez, *Espaces de Banach de dimension finie*, J. Inst. Math. Jussieu **1** (2002), no. 3, 331–439. MR 1956055 (2004b:11160)
- [dJvdP96] Johan de Jong and Marius van der Put, *Étale cohomology of rigid analytic spaces*, Doc. Math. **1** (1996), No. 01, 1–56 (electronic). MR 1386046
- [Elk73] Renée Elkik, *Solutions d'équations à coefficients dans un anneau hensélien*, Ann. Sci. École Norm. Sup. (4) **6** (1973), 553–603 (1974). MR 0345966 (49 #10692)
- [Fal02] Gerd Faltings, *Almost étale extensions*, Astérisque (2002), no. 279, 185–270, Cohomologies  $p$ -adiques et applications arithmétiques, II. MR 1922831 (2003m:14031)
- [Far16] Laurent Fargues, *Geometrization of the local Langlands correspondence: an overview*, arXiv:1602.00999.
- [FF11] Laurent Fargues and Jean-Marc Fontaine, *Courbes et fibrés vectoriels en théorie de Hodge  $p$ -adique*, preprint, 2011.
- [Fon13] Jean-Marc Fontaine, *Perfectoïdes, presque pureté et monodromie-poids (d'après Peter Scholze)*, Astérisque (2013), no. 352, Exp. No. 1057, x, 509–534, Séminaire Bourbaki. Vol. 2011/2012. Exposés 1043–1058. MR 3087355
- [FS] Laurent Fargues and Peter Scholze, *Geometrization of the local Langlands correspondence*, in preparation.
- [GR03] Ofer Gabber and Lorenzo Ramero, *Almost ring theory*, Lecture Notes in Mathematics, vol. 1800, Springer-Verlag, Berlin, 2003. MR 2004652 (2004k:13027)
- [Gro68] Alexander Grothendieck, *Le groupe de Brauer. III. Exemples et compléments*, Dix exposés sur la cohomologie des schémas, Adv. Stud. Pure Math., vol. 3, North-Holland, Amsterdam, 1968, pp. 88–188. MR 244271
- [Han16] David Hansen, *Period morphisms and variations of  $p$ -adic hodge structure*, <http://www.math.columbia.edu/~hansen/periodmapmod.pdf>.
- [Hub93a] R. Huber, *Continuous valuations*, Math. Z. **212** (1993), no. 3, 455–477. MR 1207303 (94e:13041)
- [Hub93b] Roland Huber, *Étale cohomology of Henselian rings and cohomology of abstract Riemann surfaces of fields*, Math. Ann. **295** (1993), no. 4, 703–708. MR 1214956
- [Hub96] ———, *Étale cohomology of rigid analytic varieties and adic spaces*, Aspects of Mathematics, E30, Friedr. Vieweg & Sohn, Braunschweig, 1996. MR 1734903
- [Ked16] Kiran S. Kedlaya, *On commutative nonarchimedean Banach fields*, arXiv:1602.09004.
- [KL15] Kiran S. Kedlaya and Ruochuan Liu, *Relative  $p$ -adic Hodge theory: foundations*, Astérisque (2015), no. 371, 239. MR 3379653
- [KL16] ———, *Relative  $p$ -adic Hodge theory, II: Imperfect period rings*, arXiv:1602.06899.
- [Lur09] Jacob Lurie, *Higher topos theory*, Annals of Mathematics Studies, vol. 170, Princeton University Press, Princeton, NJ, 2009. MR 2522659 (2010j:18001)
- [Lur16] ———, *Higher algebra*, 2016, <http://www.math.harvard.edu/~lurie/papers/HA.pdf>.
- [LZ14] Yifeng Liu and Weizhe Zheng, *Enhanced six operations and base change theorem for Artin stacks*, arXiv:1211.5948v2.
- [RV14] Michael Rapoport and Eva Viehmann, *Towards a theory of local Shimura varieties*, Münster J. Math. **7** (2014), no. 1, 273–326. MR 3271247
- [Sch92] Claus Scheiderer, *Quasi-augmented simplicial spaces, with an application to cohomological dimension*, J. Pure Appl. Algebra **81** (1992), no. 3, 293–311. MR 1179103
- [Sch12] Peter Scholze, *Perfectoid spaces*, Publ. Math. Inst. Hautes Études Sci. **116** (2012), 245–313. MR 3090258
- [Sch15] ———, *On torsion in the cohomology of locally symmetric varieties*, Ann. of Math. (2) **182** (2015), no. 3, 945–1066. MR 3418533
- [Sta] The Stacks Project Authors, *Stacks Project*, <http://stacks.math.columbia.edu>.
- [SW13] Peter Scholze and Jared Weinstein, *Moduli of  $p$ -divisible groups*, Cambridge Journal of Mathematics **1** (2013), no. 2, 145–237.