

Small eigenvalues of geometrically finite manifolds

Ursula Hamenstädt*

Mathematisches Institut der Universität
Berlingstraße 1, D-53115 Bonn, Germany

Abstract

Let M be a complete geometrically finite manifold of bounded negative curvature, infinite volume and dimension at least 3. We give both a lower bound for the bottom of the spectrum of M and an upper bound for the number of the small eigenvalues of M . These bounds only depend on the dimension, curvature bounds and the volume of the one-neighborhood of the convex core.¹

1 Introduction

Let M be a complete Riemannian manifold of dimension $n \geq 3$ with sectional curvature $-\kappa \leq K \leq -1$ for some $\kappa \geq 1$. Denote by X the universal covering of M ; then there is a discrete torsion free subgroup Γ of the isometry group $\text{Iso}(X)$ of X such that $M = X/\Gamma$. The group Γ acts by homeomorphisms on the *ideal boundary* ∂X of X . The *limit set* Λ of Γ is the smallest nonempty closed Γ -invariant subset of ∂X . The quotient under the action of Γ of the closed convex hull of Λ in X is a closed convex subset $C(M)$ of M , the so-called *convex core* of M .

*Research partially supported by SFB 611.

¹AMS Subject classification: 58J50.

Keywords: Geometrically finite manifolds, bottom of the L^2 -spectrum, bottom of the essential spectrum, small eigenvalues.

The manifold M is called *geometrically finite* if the volume of the one-neighborhood $C_1(M)$ of $C(M)$ in M is finite. This is in particular the case if the manifold M is of finite volume itself [B].

The Laplace-Beltrami operator Δ of M acts on the space of smooth functions with compact support and admits a unique extension to an unbounded self-adjoint operator on $L^2(M)$. The spectrum of $-\Delta$ is a closed subset $\sigma(M)$ of the half-line $[0, \infty)$ which is the disjoint union of the *essential spectrum* $\sigma_{\text{ess}}(M)$ and the *discrete spectrum* $\sigma_{\text{disc}}(M)$. The essential spectrum is a closed subset of $[0, \infty)$. The discrete spectrum consists of eigenvalues of finite multiplicity; they are isolated points in $\sigma(M)$.

In this note we are interested in the *small part* of the spectrum of M , i.e. the intersection of $\sigma(M)$ with the interval $[0, (n-1)^2/4)$. Notice that $(n-1)^2/4$ is just the bottom of the L^2 -spectrum of hyperbolic n -space \mathbf{H}^n . Our first result is a positive lower bound for the bottom $\lambda_0(M)$ of the spectrum of M under the assumption that the volume of M is infinite. This bound only depends on the dimension n , the curvature bounds and the volume of $C_1(M)$ and generalizes an earlier result of Burger and Canary [BC] for geometrically finite hyperbolic manifolds, with a simpler proof.

We also show that the bottom of the essential spectrum of M is not smaller than $(n-1)^2/4$. Thus the small part of the spectrum consists of a collection of eigenvalues. We give an upper bound for the number of eigenvalues contained in a closed interval of the form $[0, (n-1)^2/4 - \chi]$ for some $\chi > 0$. This bound only depends on the dimension n , curvature bounds, the volume of $C_1(M)$ and the choice of χ and generalizes in a weaker form an estimate of Buser, Colbois and Dodziuk [BCD] for manifolds of finite volume and bounded negative curvature. Our results can be summarized as follows.

Theorem: *Let $n \geq 3, \kappa \geq 1$ and let M be a geometrically finite manifold of dimension n , infinite volume and sectional curvature contained in $[-\kappa, -1]$. Denote by $\text{vol}(C_1(M))$ the volume of the neighborhood of radius 1 of the convex core of M .*

1. *There is a constant $c_0 = c_0(n, \kappa) > 0$ such that*

$$\lambda_0(M) \geq \frac{c_0}{\text{vol}(C_1(M))^2}.$$

2. *The essential spectrum of M is contained in $[(n-1)^2/4, \infty)$.*

3. For every $\chi > 0$ there is a constant $c_1 = c_1(n, \kappa, \chi) > 0$ such that the number of eigenvalues of M contained in the interval $(0, (n-1)^2/4 - \chi]$ is bounded from above by $c_1^{\text{vol}(C_1(M))}$.

In the case of geometrically finite subgroups Γ of the simple rank-one Lie groups $O_{\mathbb{F}}(n, 1)$ where \mathbb{F} is one of the Archimedean fields $\mathbb{R}, \mathbb{C}, \mathbb{H}, \mathbb{O}$ the bottom of the spectrum of the corresponding rank-one locally symmetric space M corresponds precisely to the *critical exponent* of Γ . In particular, the bottom of the spectrum of M is 0 if and only if this critical exponent equals $\delta_{\mathbb{F}}(n) = \dim(M) + \dim_{\mathbb{R}}\mathbb{F} - 2$. The proof of our theorem also yields the following result.

Corollary: *Let Γ be a geometrically finite subgroup of $O_{\mathbb{F}}(n, 1)$ and denote by M the locally symmetric space defined by Γ .*

1. *The critical exponent of Γ equals $\delta_{\mathbb{F}}(n)$ if and only if Γ is a lattice.*
2. *The essential spectrum of M is contained in $[\delta_{\mathbb{F}}(n), \infty)$, and for every $\chi > 0$ the number of eigenvalues of M contained in $[0, \delta_{\mathbb{F}}(n) - \chi]$ is bounded from above by $c(\chi)^{\text{vol}(C_1(M))}$.*

The first part of our corollary is well known for real hyperbolic manifolds (see [BC] and the references there) and, in a much stronger form, for general discrete subgroups of $O_{\mathbb{H}}(n, 1), O_{\mathbb{O}}(2, 1)$ (see [C]).

Our method is very general and can be applied to give a lower bound for the spectrum and the essential spectrum for the Laplacian acting on forms; however we do not address this question here. For a discussion of further related results we refer to [BC] and [BCD].

2 Tubes, cusps and ends

In this section we collect some preliminary results on the spectrum of tubes, cusps and ends in a geometrically finite Riemannian manifold M of dimension $n \geq 3$ and of bounded negative curvature $-\kappa \leq K \leq -1$. Recall that such a manifold M can be written in the form $M = X/\Gamma$ where Γ is a discrete torsion free subgroup of the isometry group $\text{Iso}(X)$ of the universal covering X of M . The group Γ is called *elementary* if its action on the ideal boundary ∂X

of X has a finite orbit. An elementary torsion free discrete isometry group Γ either is an infinite cyclic group of loxodromic isometries or Γ is a parabolic group which fixes a point $\xi \in \partial X$ (see [B]).

We begin with estimating the bottom of the spectrum for the quotient of X under a torsion free elementary group Γ .

Lemma 2.1: *Let Γ be an infinite cyclic group of loxodromic isometries of X ; then $\lambda_0(X/\Gamma) \geq (n-1)^2/4$.*

Proof: Since the bottom of the L^2 -spectrum of $M = X/\Gamma$ for the operator $-\Delta$ equals the top of the positive spectrum of $-\Delta$ it is enough to find a positive $\Delta + (n-1)^2/4$ -superharmonic function on M [S]. For this let $\tilde{\gamma}$ be the axis of the action of Γ on X ; then $\tilde{\gamma}$ is a Γ -invariant geodesic in X , and Γ acts as a group of translations on $\tilde{\gamma}$. Denote by $\tau > 0$ the translation length on $\tilde{\gamma}$ of a generator Ψ of Γ ; then for every $x \in X$ and every $k \in \mathbb{Z}$ we have $\text{dist}(x, \Psi^k x) \geq k\tau$.

Fix a point $x \in X$ and denote by $\rho_x(y)$ the distance of a point $y \in X$ to x . Write $\delta = (n-1)/2$; using Rauch's comparison theorem we have

$$\Delta(e^{-\delta\rho_x}) \leq e^{-\delta\rho_x}(-(n-1)\delta + \delta^2)$$

at least on $X - \{x\}$. Therefore the function $e^{-\delta\rho_x}$ is $\Delta + \delta^2$ -superharmonic, moreover it is positive. On the other hand, by our above observation, for each $y \in X$ the series

$$\sum_k e^{-\delta\rho_{\Psi^k x}(y)}$$

converges and hence the assignment $y \rightarrow \sum_k e^{-\delta\rho_{\Psi^k x}(y)}$ defines a Γ -invariant positive $\Delta + \delta^2$ -superharmonic function on X which projects to a positive $\Delta + \delta^2$ -superharmonic function on $M = X/\Gamma$. This shows the lemma. \square

Next we look at the bottom of the spectrum of an elementary parabolic group. For this recall that for every complete Riemannian manifold N the bottom $\lambda_0(N)$ of the spectrum of N is the infimum of the *Rayleigh quotients* $\mathcal{R}(f) = \frac{\int |df|^2}{\int f^2}$ over all nontrivial smooth functions on N with compact support.

Lemma 2.2: *Let $\Gamma \subset \text{Iso}(X)$ be an elementary parabolic group; then $\lambda_0(X/\Gamma) \geq (n-1)^2/4$.*

Proof: Let $\Gamma \subset \text{Iso}(X)$ be an elementary parabolic group. Then Γ stabilizes a horosphere H in X , and $M = X/\Gamma$ is diffeomorphic to $H/\Gamma \times \mathbb{R}$ with

a diffeomorphism which maps each line $\{z\} \times \mathbb{R}$ to a geodesic in M . If we denote by dx the volume element on H/Γ of the restriction of the Riemannian metric on M then the volume element ω on M can be represented in the form $\omega = dt \times hdx$ where $h : H/\Gamma \times \mathbb{R} \rightarrow (0, \infty)$ satisfies

$$\frac{\partial}{\partial t} h(x, t) \geq (n-1)h(x, t)$$

(see [HIH]).

Now let $f : M \rightarrow \mathbb{R}$ be any smooth function with compact support. We use an idea of Kean (see [BCD]). Namely, for $x \in H/\Gamma$ we have

$$\begin{aligned} & \left(\int_{-\infty}^{\infty} \left| \frac{\partial f}{\partial t} \right|^2 h(x, t) dt \right)^{1/2} \left(\int_{-\infty}^{\infty} f^2 h(x, t) dt \right)^{1/2} \geq \int_{-\infty}^{\infty} \left| \frac{\partial f}{\partial t} \right| f h(x, t) dt \\ &= \frac{1}{2} \int_{-\infty}^{\infty} \left| \frac{\partial}{\partial t} (f^2) \right| h(x, t) dt \geq \frac{1}{2} \int_{-\infty}^{\infty} f^2 \frac{\partial}{\partial t} h(x, t) dt \geq \frac{n-1}{2} \int_{-\infty}^{\infty} f^2 h(x, t) dt \end{aligned}$$

and therefore

$$\int_{H/\Gamma} \int_{-\infty}^{\infty} \left| \frac{\partial f}{\partial t} \right|^2 h(x, t) dt dx \geq \left(\frac{n-1}{2} \right)^2 \int_{H/\Gamma} \int_{-\infty}^{\infty} f^2 h(x, t) dt dx.$$

Thus the Rayleigh quotient of f is at least $(n-1)^2/4$. \square

Now consider an arbitrary geometrically finite manifold M of curvature contained in $[-\kappa, -1]$. As in the introduction, let $C(M)$ be the convex core of M . Then $M - C(M)$ is an open subset of M which consists of finitely many connected components (see [B]). Recall that for each open subset Ω of a Riemannian manifold M the smallest Rayleigh quotient $\mu_1(\Omega)$ of Ω is defined to be the infimum of all Rayleigh quotients for all smooth functions f with compact support in Ω ; in particular we have $\mu_1(\Omega) \geq \mu_1(M)$.

The next lemma is similar to Lemma 2.2. For its formulation, denote for a subset A of M and $r > 0$ by $B(A, r)$ the open r -neighborhood of A in M .

Lemma 2.3: *Let M be geometrically finite with curvature $-\kappa \leq K \leq -1$; then for every $r > 0$ we have $\mu_1(M - \overline{B(C(M), r)}) \geq (\tanh r)^2 (n-1)^2/4$.*

Proof: Since $B(C(M), r)$ is the r -neighborhood of a convex subset of M , its boundary is a smooth hypersurface $\partial B(C(M), r)$ in M . The second fundamental form of this boundary is positive definite. If we denote by N the outer normal of $\partial B(C(M), r)$ then the normal exponential map \exp is a

diffeomorphism of $\{tN \mid t \in (0, \infty)\}$ onto $M - \overline{B(C(M), r)}$. The pullback under \exp of the volume element ω on $M - \overline{B(C(M), r)}$ can be written in the form $\omega = dt \times h(x, t)dx$ where dx is the volume element on $\partial B(C(M), r)$ and where $h : \partial B(C(M), r) \times [0, \infty) \rightarrow [1, \infty)$ is a smooth function. By comparison with a manifold of constant curvature -1 , this function satisfies

$$\frac{\partial}{\partial t} h(x, t) \geq (n-1)h(x, t) \tanh(t+r)$$

for all $t \geq 0$. The lemma now follows from the arguments in the proof of Lemma 2.2. \square

Remark: Using the notations from the introduction, if Γ is a geometrically finite subgroup of the group $O_{\mathbb{F}}(n, 1)$ for $\mathbb{F} = \mathbb{R}, \mathbb{C}, \mathbb{H}, \mathbb{O}$ then using the standard computations for Jacobi fields in rank-one symmetric spaces we observe that the proofs of Lemma 2.1-2.3 remain valid if we replace the lower bound $(n-1)^2/4$ for the spectrum of our tubes, cusps and ends by the (sharp) bound $\delta_{\mathbb{F}}(n)$.

3 Proof of the theorem

We continue to look at a geometrically finite Riemannian manifold (M, g) of dimension $n \geq 3$ and sectional curvature $-\kappa \leq K \leq -1$. As before, we denote by X the universal covering of M and by Γ the fundamental group of M . For a number $\delta > 0$ let $M_{\text{thin}(\delta)}$ be the set of all points in M at which the injectivity radius is smaller than δ . Following Margulis, there is a constant $\epsilon_0 > 0$ only depending on the dimension and the curvature bounds such that for every $\epsilon \leq \epsilon_0$ each connected component of $M_{\text{thin}(\epsilon)}$ is isometric to an open connected subset of the quotient of X under an elementary torsion-free subgroup G of Γ . In the sequel we always denote by ϵ_0 this constant. For $\epsilon \in (0, \epsilon_0]$ define $M_{\text{thick}(\epsilon)}$ to be the ϵ -thick part of M , i.e. the set of all points in M with injectivity radius at least ϵ . Since M is geometrically finite, the intersection $M_{\text{thick}(\epsilon)} \cap C(M)$ is compact [B].

For an open subset Ω of M the Laplacian acting on smooth functions with compact support in Ω admits a unique self-adjoint extension Δ_{Ω} to an unbounded operator on $L^2(\Omega)$. The domain of $\Delta_{\Omega}^{1/2}$ is the usual Sobolev space $H^1(\Omega)$ which is the completion of the space of smooth functions with compact support in Ω with respect to the Sobolev norm $\|f\|^2 = \int f^2 + \int |df|^2$. Let $\sigma(\Omega)$ be the spectrum of $-\Delta_{\Omega}$.

We use the spectral theorem for $-\Delta_\Omega$ in the following form (see [D]). There is a finite measure μ on $\sigma(\Omega) \times \mathbb{N}$ and a unitary operator $U : L^2(\Omega) \rightarrow L^2(\sigma(\Omega) \times \mathbb{N}, d\mu)$ as follows. Define $h(s, n) = s$; then $f \in L^2(\Omega)$ is contained in the domain of $-\Delta_\Omega$ if and only if $hU(f) \in L^2(\sigma(\Omega) \times \mathbb{N}, d\mu)$, and if this is the case we have $-U\Delta_\Omega U^{-1}(Uf) = hU(f)$. The spectral measure of such a function f is supported in an interval $[\lambda - \kappa, \lambda + \kappa]$ if and only if the function Uf is supported in $[\lambda - \kappa, \lambda + \kappa] \times \mathbb{N}$. Since $(u, q) \rightarrow \int g(du, dq)$ is the quadratic form of $-\Delta_\Omega^{1/2}$ this implies that for every $q \in H^1(\Omega)$ we have

$$\left| \int g(df, dq) - \lambda \int fq \right| \leq \kappa \int |fq|.$$

Using this inequality for $u = f$ we obtain in particular that the Raleigh quotient of f is contained in the interval $[\lambda - \kappa, \lambda + \kappa]$. Moreover, if f and q are contained in the domain of $-\Delta_\Omega$ and if their spectral measures are supported on disjoint subsets of $\sigma(\Omega)$ then $\int fq = \int g(df, dq) = 0$.

Call a square integrable function f on M *normalized* if $\int f^2 = 1$. We have.

Lemma 3.1: *For every $\sigma > 0$ there is a number $R_0 = R_0(\sigma, \kappa, n) > 0$, a number $\delta = \delta(\sigma, \kappa, n) > 0$ and a number $\epsilon = \epsilon(\sigma, \kappa, n) < \epsilon_0$ with the following properties. Let $f \in H^1(M)$ be a normalized function with spectral measure supported in a subinterval of $[0, (n-1)^2/4 - \sigma]$ of length at most δ ; then there is a normalized function $Lf \in H^1(B(M_{\text{thick}(\epsilon)} \cap C(M), R_0))$ whose spectral measure is supported in $[0, (n-1)^2/4 - \sigma/2]$ and such that*

$$\int (f - Lf)^2 < 1/4.$$

Proof: By the result of Margulis, for $\epsilon \leq \epsilon_0$ each connected component of $M_{\text{thin}(\epsilon)}$ is isometric to an open subset of the quotient of X under an elementary subgroup of the isometry group of X . Thus by domain monotonicity and Lemma 2.1, Lemma 2.2 the smallest Raleigh quotient $\mu_1(M_{\text{thin}(\epsilon_0)})$ of $M_{\text{thin}(\epsilon_0)}$ is not smaller than $(n-1)^2/4$. Lemma 2.3 also shows that $\mu_1(M - \overline{B(C(M), r)}) \geq (\tanh r)^2(n-1)^2/4$.

Let $\sigma > 0$ and define $\chi = \min\{1/16(12+3(n-1)^2), \sigma/2\}$ and $\nu = 4\chi^3/(n+1)^2$. By Lemma 2.3 of [FH], applied to the closed set $\overline{B(C(M), r)}$ for a number $r > 0$ which is sufficiently big that $(\tanh r)^2(n-1)^2/4 \geq (n-1)^2/4 - \sigma/2$ and to the closed set $M_{\text{thick}(\epsilon_0)}$, there is a number $\delta = \delta(\sigma) < \nu$ and a

constant $R = R(\sigma) > 0$ with the following properties. Let $f \in H^1(M)$ be a normalized function whose spectral measure is supported in a closed subinterval of $[0, (n-1)^2/4 - \sigma]$ of length at most δ ; then

$$\int_{M-B(C(M),R)} f^2 + |df|^2 < \nu^2/4, \quad \int_{M-B(M_{\text{thick}(\epsilon_0)},R)} f^2 + |df|^2 < \nu^2/4.$$

Since each connected component of the ϵ_0 -thin part of M is isometric to a connected subset of the quotient of X under an elementary group, there are numbers $\epsilon_1 \in (0, \epsilon_0]$, $\epsilon \in (0, \epsilon_1]$ only depending on the dimension, the curvature bounds and the radius R (which only depends on σ) such that $B(M_{\text{thick}(\epsilon_0)}, R) \subset M_{\text{thick}(\epsilon_1)}$ and $B(M_{\text{thick}(\epsilon_1)}, R) \subset M_{\text{thick}(\epsilon)}$. But this just means that $B(C(M), R) \cap B(M_{\text{thick}(\epsilon_0)}, R) \subset B(C(M), R) \cap M_{\text{thick}(\epsilon_1)} \subset B(C(M) \cap M_{\text{thick}(\epsilon)}, R)$; thus our function f as above satisfies

$$\int_{M-B(C(M) \cap M_{\text{thick}(\epsilon)}, R)} f^2 + |df|^2 < \nu^2/2.$$

Choose a smooth function $\tilde{u} : \mathbb{R} \rightarrow [0, 1]$ which equals 1 on $(-\infty, 0]$, vanishes on $[2, \infty)$ and whose gradient is pointwise bounded in norm by 1. Define $u(x) = \tilde{u}(\text{dist}(x, C(M) \cap M_{\text{thick}(\epsilon)}) - R)$; then u is supported in the $R+2$ -neighborhood $\Omega = B(C(M) \cap M_{\text{thick}(\epsilon)}, R+2)$ of $C(M) \cap M_{\text{thick}(\epsilon)}$ and its gradient is pointwise bounded in norm by 1. The function uf is contained in $H^1(\Omega)$ and it satisfies

$$\begin{aligned} & \left| \int f^2 - (uf)^2 \right| + \int (f - uf)^2 \leq 2 \int_{M-B(C(M) \cap M_{\text{thick}(\epsilon)}, R)} f^2 < \nu^2 \quad \text{and} \\ & \int |df - d(uf)|^2 + \left| \int |df|^2 - |d(uf)|^2 \right| \\ & \leq 2 \int_{M-B(C(M) \cap M_{\text{thick}(\epsilon)}, R)} |df|^2 + f^2 + |2ufg(df, du)| < 3\nu^2. \end{aligned}$$

Let λ be the midpoint of an interval of length at most $\delta \leq \nu$ which contains the support of the spectral measure for f . The function uf lies in the domain of the operator Δ_Ω and therefore uf admits an L^2 -orthogonal decomposition of the form $uf = \alpha + \varphi + \beta$ where the spectral measure of α is supported in $[0, \lambda - \chi^2]$, the spectral measure of φ is supported in $[\lambda - \chi^2, \lambda + \chi]$ and the spectral measure of β is supported in $[\lambda + \chi, \infty)$. Denote by $\|\psi\|$ the L^2 -norm of a square integrable function ψ on Ω or M

and denote similarly by $\|\xi\|$ the L^2 -norm of a square integrable one-form ξ on Ω or M . Define $Lf = \alpha + \varphi/\|\alpha + \varphi\|$. For the proof of the lemma it is now enough to show that $\|\beta\| < 1/8$. Namely, if this is the case then we have $|\|\alpha + \varphi\| - 1| \leq \|\alpha + \varphi - f\| < 1/4$ and hence $\|Lf - \alpha - \varphi\| < 1/4$ and $\|f - Lf\| \leq \|f - \alpha - \varphi\| + \|\alpha + \varphi - Lf\| < 1/2$.

For this estimate of $\|\beta\|$ we proceed as in [FH]. Observe first that we have

$$\|\alpha\|^2 = \int \alpha(uf) = \int \alpha f + \int \alpha(uf - f) \leq \int \alpha f + \nu\|\alpha\|.$$

Now f is normalized and the Raleigh quotient of α is not bigger than $\lambda - \chi^2 < (n-1)^2/4$ and consequently we obtain that

$$\begin{aligned} (\lambda - \chi^2)\|\alpha\|^2 &\geq \|d\alpha\|^2 = \int g(d\alpha, d(uf)) \geq \int g(d\alpha, df) - \sqrt{3\nu}\|d\alpha\| \\ &\geq \lambda \int \alpha f - \nu(\|\alpha\| + \sqrt{3\lambda}\|\alpha\|) \geq \lambda\|\alpha\|^2 - \nu\|\alpha\|(\lambda + 1 + \sqrt{3\lambda}). \end{aligned}$$

Since $\lambda + 1 + \sqrt{3\lambda} \leq (n+1)^2/4$ we conclude that $\|\alpha\| \leq \nu(n+1)^2/4\chi^2 = \chi$ by our choice of ν .

On the other hand, the square norm of β can be estimated as follows. By construction we have

$$\begin{aligned} (1 + 3\nu^2)(\lambda + \nu) &\geq (1 + 3\nu^2)\|df\|^2 \geq \|d(uf)\|^2 \\ &= \|d\alpha\|^2 + \|d\varphi\|^2 + \|d\beta\|^2 \geq (\lambda - \chi^2)\|\varphi\|^2 + (\lambda + \chi)\|\beta\|^2. \end{aligned}$$

Since $\|\varphi\|^2 + \|\beta\|^2 = \|uf\|^2 - \|\alpha\|^2 \geq 1 - \nu^2 - \|\alpha\|^2 \geq 1 - 2\chi^2$ we obtain from this that

$$(1 + 3\nu^2)(\lambda + \nu) \geq (1 - 2\chi^2)(\lambda - \chi^2) + \chi\|\beta\|^2$$

and hence $\chi\|\beta\|^2 \leq \nu + 3\nu^2(\lambda + \nu) + \chi^2 + 2\chi^2(\lambda - \chi^2)$ and $\|\beta\|^2 \leq \chi(3 + 3\lambda)$. By our choice of χ this means that $\|\beta\| < 1/8$ from which the lemma follows. \square

Recall that if $\Omega \subset M$ is an open set with compact closure then the spectrum of Ω consists of a countable number of eigenvalues going to ∞ . In the sequel we always count our eigenvalues with multiplicities. We have.

Corollary 3.2:

1. *The essential spectrum of M is contained in $[(n-1)^2/4, \infty)$.*

2. *There is a constant $\chi > 0$ with the following property. For $\sigma > 0$ let $R_0 = R_0(\sigma, \kappa, n) > 0$ be as in Lemma 3.1 and let $q \geq 0$ be the number of eigenvalues of $\Omega = B(M_{\text{thick}(\epsilon)} \cap C(M), R_0)$ which are contained in the interval $(0, (n-1)^2/4 - \sigma/2]$; then the number of eigenvalues of M contained in the interval $(0, (n-1)^2/4 - \sigma]$ is not bigger than χ^q .*

Proof: Let $\sigma > 0$ and let $R_0 = R_0(n, \kappa, \sigma) > 0$ be as in Lemma 3.1. Since the closure of $\Omega = B(M_{\text{thick}(\epsilon)} \cap C(M), R_0)$ is compact, its spectrum consists of a countable number of eigenvalues which increase to ∞ . Let E be the sum of the eigenspaces for $-\Delta_\Omega$ with respect to those eigenvalues which are not bigger than $(n-1)^2/4 - \sigma/2$. Then E is a finite-dimensional linear subspace of $H^1(\Omega)$, and its orthogonal complement E^\perp in $H^1(\Omega)$ is the closed vector subspace which admits a Hilbert basis consisting of the complete set of eigenfunctions with respect to all eigenvalues contained in $((n-1)^2/4 - \sigma/2, \infty)$.

Let $\delta = \delta(\sigma) > 0$ be as in Lemma 3.1 and let $f_1, \dots, f_\ell \subset H^1(M)$ be a collection of mutually L^2 -orthogonal normalized functions whose spectral measures are supported in a subinterval of $[0, (n-1)^2/4 - \sigma]$ of length at most δ . By Lemma 3.1 there is a collection $g_1, \dots, g_\ell \subset E$ of normalized functions with $\int (f_i - g_i)^2 < 1/4$. Then the functions g_i are contained in the unit sphere S in the finite-dimensional vector space E equipped with the L^2 -inner product, and their pairwise euclidean distance is at least $1/4$. In other words, the balls in S of radius $1/8$ which are centered at the points g_1, \dots, g_ℓ are pairwise disjoint. But this implies that there is a universal constant $\chi > 0$ such that the cardinality of the set $\{g_1, \dots, g_\ell\}$ and hence the cardinality of the set $\{f_1, \dots, f_\ell\}$ is bounded from above by $\chi^{\dim(E)}$.

Thus the dimension of the vector-subspace of $H^1(M)$ which is spanned by all functions whose spectral measures are supported in a subinterval of $[0, (n-1)^2/4 - \sigma]$ of length at most δ is bounded from above by $\chi^{\dim(E)}$. This means that the essential spectrum of M does not intersect the interval $[0, (n-1)^2/4 - \sigma]$, and the dimension of the sum of the eigenspaces of $-\Delta$ for eigenvalues contained in $[0, (n-1)^2/4 - \sigma]$ is bounded from above by $\chi^{\dim(E)}$. Since $\sigma > 0$ was arbitrary the corollary follows. \square

Corollary 3.2 shows the second part of the theorem from the introduction and easily implies the third.

Corollary 3.3: *For every $\sigma > 0$ there is a constant $c_1 = c_1(\sigma, n, \kappa) > 0$ such that the number of eigenvalues of M which are contained in the interval*

$[0, (n-1)^2/4 - \sigma]$ is not bigger than $c_1^{\text{vol}(C_1(M))}$.

Proof: Let $\sigma > 0$ and let $\epsilon = \epsilon(\sigma) > 0$ and $R_0 = R_0(\sigma) > 0$ be as in Lemma 3.1. Notice that ϵ and R_0 only depend on σ and the curvature bounds. Denote by Ω the open R_0 -neighborhood of $M_{\text{thick}(\epsilon)} \cap C(M)$; then the closure of Ω is compact. We claim that there is a constant $\zeta > 0$ only depending on R_0 , the dimension and the curvature bounds such that the volume of Ω is bounded from above by $\zeta \text{vol}(C_1(M))$.

To see this recall that $B(C(M), 1/2)$ is the $1/2$ -neighborhood of a convex subset of M and therefore the second fundamental form of the boundary $\partial B(C(M), 1/2)$ of $B(C(M), 1/2)$ at a point $x \in \partial B(C(M), 1/2)$ is not bigger than the second fundamental form of a ball of radius $1/2$ in M which meets $\partial B(C(M), 1/2)$ tangentially at x . Standard comparison arguments then imply that the second fundamental form of $\partial B(C(M), 1/2)$ is globally uniformly bounded. Comparison with a manifold of constant curvature κ then shows that the volume of $B(C(M), R_0)$ is bounded from above by a fixed multiple of $\text{vol}(C_1(M))$.

As a consequence of this and Corollary 3.2 we only have to show that the number of eigenvalues of $-\Delta_\Omega$ contained in the interval $[0, (n-1)^2/4]$ is bounded from above by a constant multiple of $\text{vol}(\Omega)$.

For this recall from Lemma 3.1 and its proof that there is a number $\rho > 0$ only depending on R_0, ϵ and the curvature bounds such that for every $z \in \bar{\Omega}$ the injectivity radius of M at z is at least ρ . Choose a collection of points $\{p_1, \dots, p_\ell\} \subset \Omega$ of maximal cardinality whose pairwise distances are at least ρ . Then the closed balls of radius ρ centered at the points p_i cover Ω , and the balls of radius $\rho/2$ centered at the points p_i are pairwise disjoint. Since the injectivity radius at each of the points p_i is at least ρ , the volume of the balls of radius $\rho/2$ about the points p_i is bounded from below by a universal constant only depending on the curvature bounds. Thus the cardinality of our collection is bounded from above by a constant multiple of $\text{vol}(\Omega)$.

For $i \geq 1$ define $U_i = \{z \in \Omega \mid \text{dist}(z, p_i) \leq \text{dist}(z, p_j) \text{ for } j \neq i\}$. The sets U_i form a partition of Ω , and each of the sets U_i is contained in the ball of radius ρ about p_i . Now the smallest nonzero eigenvalue with Neumann boundary conditions of a ball of radius ρ in a simply connected manifold of curvature contained in $[-\kappa, -1]$ is not smaller than $(n-1)^2/4$ (compare [BCD]) and therefore by domain monotonicity, the second eigenvalue for each of the sets U_i with mixed boundary conditions (i.e. Neumann boundary con-

ditions on $\partial U_i \cap \Omega$ and Dirichlet boundary conditions on $\partial \Omega$) is not smaller than $(n-1)^2/4$ as well. On the other hand, the number of small eigenvalues on Ω is bounded from above by the total number of small eigenvalues with Neumann boundary conditions for any partition of Ω and therefore this number is bounded from above by a constant multiple of the volume of $C_1(M)$. Together with Corollary 3.2, this yields the corollary. \square

We are left with showing the first part of the theorem from the introduction. For this denote by $\lambda_0 = \lambda_0(M)$ the bottom of the spectrum for $-\Delta$ on M . By Corollary 3.2, if $\lambda_0 < (n-1)^2/4$ then λ_0 is an eigenvalue of $-\Delta$, and there is up to a constant a unique solution f of the equation $\Delta + \lambda_0(M) = 0$. This function f is square integrable and positive. For the purpose of our theorem we then may assume that such an eigenfunction f exists. For $\sigma = (n-1)^2/8$ denote by $R_0 = R_0(n, \kappa, \sigma)$ the constant as in Lemma 3.1. Let $\epsilon_1 > 0$ be sufficiently small that $B(M_{\text{thick}(\epsilon_0)}, R_0) \subset M_{\text{thick}(\epsilon_1)}$; notice once more that ϵ_1 can be chosen to depend only on the dimension and the curvature bounds. We have.

Lemma 3.4: *There is a number $\beta > 0$ only depending on the dimension and the curvature bounds of M with the following property. Let f be a positive normalized eigenfunction on M with respect to the eigenvalue $\lambda_0 < (n-1)^2/8$; then $f(x) \leq \beta\sqrt{\lambda_0}$ for all $x \in M_{\text{thick}(\epsilon_1)} \cap \partial B(C(M), R_0)$.*

Proof: By the infinitesimal Harnack inequality of Cheng and Yau [CY], applied to our positive normalized eigenfunction f with eigenvalue $\lambda_0 \in (0, (n-1)^2/8]$, there is a number $c_1 > 0$ only depending on the curvature bounds such that the function f satisfies $|d \log f| \leq c_1$ on $M_{\text{thick}(\epsilon_1)}$. In particular, if for a point $x \in M_{\text{thick}(\epsilon_1)} \cap \partial B(C(M), R_0)$ and a number $\alpha > 0$ we have $f(x) \geq \alpha\sqrt{\lambda_0}$ then there is a ball $B \subset \partial B(C(M), R_0)$ about x with the following property. Let N be the exterior normal of $B(C(M), R_0)$ and let \exp be the normal exponential map; then $\int_{\exp(B \times (0, \infty))} f^2 \geq c_2 \alpha^2 \lambda_0$ where $c_2 > 0$ is a universal constant.

We now use the arguments in the proof of Lemma 2.2 and Lemma 2.3. Namely, let $h(y, t)$ be the function on $\partial B(C(M), R_0) \times [0, \infty)$ which determines the pullback under \exp of the volume element of M with respect to the product measure on $\partial B(C(M), R_0) \times [0, \infty)$ as in Lemma 2.2. Then $h(y, 0) = 1$ for all $y \in \partial B(C(M), R_0)$ and moreover $\frac{\partial}{\partial t} h(y, t) \geq (n-1)h(y, t)/\sqrt{2}$ for every $y \in B$ and $t \geq 0$. For $y \in \partial B(C(M), R_0)$ we conclude as in the proof

of Lemma 2.2 that

$$\begin{aligned}
& \left(\int_0^\infty \left| \frac{\partial f}{\partial t} \right|^2 h(y, t) dt \right)^{1/2} \left(\int_0^\infty f^2 h(y, t) dt \right)^{1/2} \geq \int_0^\infty \left| \frac{\partial f}{\partial t} \right| f |h(y, t)| dt \\
& = \frac{1}{2} \int_0^\infty \left| \frac{\partial}{\partial t} (f^2) \right| h(y, t) dt \geq -\frac{1}{2} \int_0^\infty \frac{\partial}{\partial t} (f^2) h(y, t) dt \\
& = \frac{1}{2} f^2(y) + \frac{1}{2} \int_0^\infty f^2 \frac{\partial}{\partial t} h(y, t) dt \geq \frac{1}{2} f^2(y) + \frac{n-1}{2\sqrt{2}} \int_0^\infty f^2 h(y, t) dt.
\end{aligned}$$

Integration over B then shows that $\int_{\exp(B \times (0, \infty))} \left| \frac{\partial f}{\partial t} \right|^2 \geq \frac{(n-1)^2}{8} c_2 \alpha^2 \lambda_0$ which implies that $\alpha^2 \leq \frac{8}{c_2(n-1)^2}$. This shows the lemma. \square

Now we are ready for the proof of the first part of our theorem. Recall that it is enough to show the existence of a constant $\nu > 0$ only depending on the curvature bounds such that with $R_0 = R_0(n, \kappa, (n-1)^2/8) > 0$ as in Lemma 3.1 we have $\lambda_0 \geq \nu / (\text{vol} B(C(M), R_0))^2$. For this we follow [DR] and argue by contradiction. Assume that $\lambda_0 = \lambda_0(M) < \alpha / \text{vol}(B(C(M), R_0))^2$ for a small constant $\alpha > 0$. Denote by K_1, \dots, K_m the connected components of $M_{\text{thick}(\epsilon_1)} \cap B(C(M), R_0)$. Then for every $j \in \{1, \dots, m\}$ any two points in K_j can be connected by a chain B_1, \dots, B_k of overlapping balls of radius $\epsilon_1/2$ where $k \leq c_1 \text{vol}(K_j)$ for a universal constant $c_1 > 0$ and for which each B_i intersects at most a fixed number ℓ of other B_j 's. Now if f is an eigenfunction on M with respect to the eigenvalue λ_0 , then by the arguments of [DR] the oscillation of f on each of the sets K_j is not bigger than $c_2(\alpha / \text{vol}(B(C(M), R_0)))^{1/2}$ where $c_2 > 0$ is a universal constant. On the other hand, each of the components K_j intersects $\partial B(C(M), R_0)$ (compare [BC]) and by Lemma 3.4 the value of f at such a boundary point does not exceed $\beta \sqrt{\lambda_0}$ where $\beta > 0$ is a universal constant. As a consequence, we have $|f| \leq c_3(\alpha / \text{vol}(B(C(M), R_0)))^{1/2}$ on $\cup_{i=1}^m K_i$ where $c_3 > 0$ is another universal constant and hence $\int_{\cup K_j} f^2 \leq c_3^2 \alpha$. For sufficiently small α this value is strictly smaller than $1/2$. However Lemma 3.1 implies that $\int_{\cup K_j} f^2 \geq 3/4$ which is a contradiction.

Remark: The constant $(n-1)^2/4$ for our spectral bounds appears in our above argument only in the form of a lower bound for the spectrum of the tubes, cusps and ends of our geometrically finite manifold. Thus using the remark at the end of Section 2, in the case of a geometrically finite locally symmetric space we can use the sharp bound $\delta_{\mathbb{F}}(n)$ in our proof and obtain

the same statements, with $(n-1)^2/4$ replaced by $\delta_{\mathbb{F}}(n)$. This then yields the corollary from the introduction.

4 References

- [B] B. Bowditch, *Geometrical finiteness with variable negative curvature*, Duke Math. J. 77 (1995), 229-274.
- [BC] M. Burger, R. Canary, *A lower bound on λ_0 for geometrically finite hyperbolic n -manifolds*, J. reine angew. Math. 454 (1994), 37-57.
- [BCD] P. Buser, B. Colbois, J. Dodziuk, *Tubes and eigenvalues for negatively curved manifolds*, J. Geom. Anal. 3 (1993), 1-26.
- [CY] S.Y. Cheng, S.T. Yau, *Differential equations on Riemannian manifolds and their geometric applications*, Comm. Pure Appl. Math. 28 (1975), 333-354.
- [C] K. Corlette, *Hausdorff dimension of limit sets I*, Inv. Math. 102 (1990), 521-542.
- [D] E.B. Davies, *Spectral theory and differential operators*, Cambridge University Press, Cambridge 1995.
- [DR] J. Dodziuk, B. Randol, *Lower bounds for λ_1 on a finite-volume hyperbolic manifold*, J. Diff. Geo. 24 (1986), 133-139.
- [FH] K. Fissmer, U. Hamenstädt, *Spectral convergence of manifold pairs*, to appear in Comm. Math. Helv.
- [HIH] E. Heintze, H.J. Im Hof, *Geometry of horospheres*, J. Diff. Geo. 12 (1977), 481-491.
- [S] D. Sullivan, *Related aspects of positivity in Riemannian geometry*, J. Diff. Geo. 25 (1987), 327-351.