

Prof. Dr. Günter Harder

# Cohomology of Arithmetic Groups

Zu erscheinen 2019  
Erstellt mit L<sup>A</sup>T<sub>E</sub>X 2<sub>ε</sub>

# Preface

Finally this is now the book on "Cohomology of Arithmetic Groups" which was announced in my "Lectures on Algebraic Geometry I and II," (LAG I,II) Originally the purpose of these two volumes was to assemble basic material for this volume III. This applies especially to the chapters I-IV in LAG I, where we provide the necessary background in homological algebra.

During the years 1980-2000 I gave various advanced courses on number theory, algebraic geometry and also on "Cohomology of arithmetic Groups" at the university of Bonn. I prepared some notes for the lectures, because there was essentially no literature covering this subject.

At some point I had the idea to use these notes as a basis for a book on this subject, a book that introduced into the subject but that also covered the applications to number theory.

It was clear that a self-contained exposition needs some preparation, we need homological algebra and later if we treat Shimura varieties, we need also a lot of algebraic geometry, especially the concept of moduli spaces. Since the cohomology groups of arithmetic groups are sheaf cohomology groups, and since the theory of sheaves and sheaf cohomology is ubiquitous in algebraic geometry I had the idea to write a volume "Lectures on Algebraic Geometry" where I discuss the impact of sheaf theory and the cohomology of sheaves to algebraic geometry. This volume eventually became the two volumes mentioned above and the writing of these volume is at last partly responsible for the delay.

The applications to number theory concern the relationship between special values of  $L$ -functions and the integral structure of the cohomology as module under the Hecke algebra. On the one hand we can prove rationality statements for special values (Manin and Shimura), on the other hand these special values tell us something about the denominators of the Eisenstein classes. These connections was already discussed in the original notes in 1985 for the special case of  $Sl_2(\mathbb{Z})$ . and the precise results in this special case are stated at the end of Chapter 5 of this book.

This (conjectural) relationship between special values of  $L$ -functions and the denominators of Eisenstein classes is one of the central themes of this book. The conjectures concerning the denominators imply congruences between eigenvalues of Hecke operators on different groups. It was extremely important for me that these conjectures on congruences got some support by experimental calculations by G. van der Geer and C. Faber and others.

These conjectures about the connections between denominators of Eisenstein classes and divisibility of special values can be verified in principle in any given case. In Chapter 3 we discuss a strategy to compute the cohomology and the Hecke endomorphisms explicitly for any specific example. Hence we can check the conjecture in such a situation. For the group  $Sl_2(\mathbb{Z})$  such explicit calculations have been done by my former student X.-D. Wang in his Bonn dissertation and are now resumed in Chapter 3.

On the other hand in this special case these conjectures are proved in chapter 5. There should be some more cases in which these conjectures can not be proved at the present moment, but where we can verify them experimentally, without using too much computing time.

The denominator question is not only an interesting problem in itself, we will also indicate how these denominators allow us to produce non trivial elements in certain Selmer groups. This means that we can construct elements in various Selmer groups which owe their existence certain divisibility of special  $L$ -values. Such a connection between  $L$ -values and the structure of the Galois group is a central theme in number theory and starts with Kummer.

I hope that this book can serve as an introduction into the field "Cohomology of arithmetic Groups", on the other hand many questions are left open and may initiate interesting research.

# Contents

0.1	Introduction . . . . .	x
<b>1</b>	<b>Basic Notions and Definitions</b>	<b>1</b>
1.1	Affine algebraic groups over $\mathbb{Q}$ . . . . .	1
1.1.1	Affine group schemes . . . . .	3
1.1.2	The classical groups and their realisation as split semi-simple group schemes over $\text{Spec}(\mathbb{Z})$ . . . . .	9
1.1.3	$k$ -forms of algebraic groups . . . . .	14
1.1.4	The Lie-algebra . . . . .	19
1.1.5	Structure of semisimple groups over $\mathbb{R}$ and the symmetric spaces . . . . .	22
1.1.6	Special low dimensional cases . . . . .	27
1.2	Arithmetic groups . . . . .	30
1.2.1	Affine group schemes over $\mathbb{Z}$ . . . . .	32
1.2.2	$\Gamma$ -modules . . . . .	33
1.2.3	The locally symmetric spaces . . . . .	34
1.2.4	Low dimensional examples . . . . .	36
1.2.5	Fixed point sets and stabilizers for $\text{Gl}_2(\mathbb{Z}[i]) = \tilde{\Gamma}$ . . . . .	41
1.2.6	Compactification of $\Gamma \backslash X$ . . . . .	43
1.2.7	The Borel-Serre compactification of $\text{Sl}_2(\mathbb{Z}) \backslash \mathbb{H}$ . . . . .	45
1.2.8	The Borel-Serre compactification, reduction theory of arithmetic groups . . . . .	46
<b>2</b>	<b>The Cohomology groups</b>	<b>63</b>
2.1	Cohomology of arithmetic groups as cohomology of sheaves on $\Gamma \backslash X$ . . . . .	63
2.1.1	The relation between $H^\bullet(\Gamma, \mathcal{M})$ and $H^\bullet(\Gamma \backslash X, \mathcal{M})$ . . . . .	64
2.1.2	How to compute the cohomology groups $H^i(\Gamma \backslash X, \tilde{\mathcal{M}})$ ? . . . .	67
2.1.3	Special examples in low dimensions. . . . .	73
2.1.4	The homology as singular homology . . . . .	77
2.1.5	The fundamental exact sequence . . . . .	80
2.1.6	How to compute the cohomology groups $H_c^q(\Gamma \backslash X, \tilde{\mathcal{M}})$ . . . . .	84
2.1.7	Modified cohomology groups . . . . .	84
2.1.8	The case $\Gamma = \text{Sl}_2(\mathbb{Z})$ . . . . .	85
<b>3</b>	<b>Hecke Operators</b>	<b>89</b>
3.1	The construction of Hecke operators . . . . .	89
3.1.1	Commuting relations . . . . .	92
3.1.2	More relations between Hecke operators . . . . .	95

3.2	Some results on semi-simple $\mathfrak{A}$ modules . . . . .	99
3.2.1	Explicit formulas for the Hecke operators, a general strategy.	101
3.3	Hecke operators for $\mathrm{Gl}_2$ : . . . . .	103
3.3.1	The boundary cohomology . . . . .	103
3.3.2	The explicit description of the cohomology . . . . .	107
3.3.3	The map to the boundary cohomology . . . . .	108
3.3.4	The first interesting example . . . . .	113
3.3.5	Computing mod $p$ . . . . .	117
3.3.6	The denominator and the congruences . . . . .	118
3.3.7	The $p$ -adic $\zeta$ -function . . . . .	126
3.3.8	The Wieferich dilemma . . . . .	128
<b>4</b>	<b>Representation Theory, Eichler-Shimura Isomorphism</b>	<b>129</b>
4.1	Harish-Chandra modules with cohomology . . . . .	129
4.1.1	The finite rank highest weight modules . . . . .	129
4.1.2	The principal series representations . . . . .	132
4.1.3	Intertwining operators . . . . .	137
4.1.4	Reducibility and representations with non trivial cohomology . . . . .	140
4.1.5	The Eichler-Shimura Isomorphism . . . . .	148
4.1.6	Local Whittaker models . . . . .	150
4.1.7	The $L$ -functions . . . . .	157
4.1.8	The Periods . . . . .	159
4.1.9	The Eisenstein cohomology class . . . . .	169
<b>5</b>	<b>Application to Number Theory</b>	<b>171</b>
5.1	Modular symbols, $L$ - values and denominators of Eisenstein classes. . . . .	171
5.1.1	Modular symbols attached to a torus in $\mathrm{Gl}_2$ . . . . .	171
5.1.2	Evaluation of cuspidal classes on modular symbols . . . . .	172
5.1.3	Evaluation of Eisenstein classes on capped modular symbols . . . . .	176
5.1.4	The capped modular symbol . . . . .	178
<b>6</b>	<b>Cohomology in the adelic language</b>	<b>191</b>
6.1	The spaces . . . . .	191
6.1.1	The (generalized) symmetric spaces . . . . .	191
6.1.2	The locally symmetric spaces . . . . .	194
6.1.3	The group of connected components, the structure of $\pi_0(S_{K_f}^G)$ . . . . .	197
6.1.4	The Borel-Serre compactification . . . . .	198
6.1.5	The easiest but very important example . . . . .	200
6.2	The sheaves and their cohomology . . . . .	201
6.2.1	Basic data and simple properties . . . . .	201
6.3	The action of the Hecke-algebra . . . . .	206
6.3.1	The action on rational cohomology . . . . .	206
6.3.2	The integral cohomology as a module under the Hecke algebra . . . . .	208
6.3.3	Excursion: Finite dimensional $\mathcal{H}$ -modules and representations. . . . .	211
6.3.4	The Satake isomorphism . . . . .	217

6.3.5	The case $\mathrm{Gl}_2$ .	220
6.3.6	Back to cohomology	222
6.3.7	The Manin-Drinfeld principle	223
6.3.8	The Gauss-Bonnet formula	225
6.3.9	Some (philosophical) remarks	238
6.3.10	Some questions and and some general facts	243
6.3.11	Poincaré duality	243
<b>7</b>	<b>The fundamental question</b>	<b>249</b>
7.0.1	The Langlands philosophy	249
<b>8</b>	<b>Analytic methods</b>	<b>267</b>
8.1	The representation theoretic de-Rham complex	267
8.1.1	Rational representations	267
8.1.2	Harish-Chandra modules and $(\mathfrak{g}, K_\infty)$ -cohomology.	268
8.1.3	The representation theoretic de-Rham isomorphism	269
8.1.4	Input from representation theory of real reductive groups.	271
8.1.5	Representation theoretic Hodge-theory.	272
8.1.6	Input from the theory of automorphic forms	274
8.1.7	Consequences.	283
8.1.8	Growth of cohomology classes	287
8.1.9	Franke's Theorem	294
8.2	Modular symbols	295
8.2.1	The general pattern	295
8.2.2	Model spaces	300
8.2.3	Rationality and integrality results	303
8.2.4	The special case $\mathrm{Gl}_2/F_0$	305
8.2.5	Poincare duality and modular symbols.	331
8.2.6	The $L$ -functions	348
8.2.7	The special values of $L$ -functions	349
<b>9</b>	<b>Eisenstein cohomology</b>	<b>351</b>
9.1	The Borel-Serre compactification	351
9.1.1	The two spectral sequences	353
9.1.2	Induction	354
9.1.3	A review of Kostants theorem	354
9.1.4	The inverse problem	358
9.2	The goal of Eisenstein cohomology	359
9.2.1	Induction and the local intertwining operator at finite places	361
9.3	The Eisenstein intertwining operator	361
9.4	The special case $\mathrm{Gl}_n$	366
9.4.1	Rationality of $L$ -values and some questions	371
9.5	Residual classes	371
9.6	Detour: $(\mathfrak{g}, K_\infty)$ - modules with cohomology for $G = \mathrm{Gl}_n$	374
9.6.1	The tempered representation at infinity	375
9.6.2	The lowest $K_\infty$ type in $\mathbb{D}_\lambda$	382
9.6.3	The unitary modules with cohomology, cohomological induction.	384
9.6.4	The $A_{\mathfrak{q}_{u,v}}(\lambda)$ as Langlands quotients	390
9.6.5	Congruences	394

9.7	The example $G = \mathrm{Sp}_2/\mathbb{Z}$ . . . . .	398
9.7.1	Some notations and structural data . . . . .	398
9.7.2	The cuspidal cohomology of the Siegel-stratum . . . . .	399

## 0.1 Introduction

This book is meant to be an introduction into the cohomology of arithmetic groups.

An arithmetic group  $\Gamma$  is a discrete subgroups of a Lie group  $G(\mathbb{R}) \subset \mathrm{GL}_n(\mathbb{R})$  whose matrix entries satisfy certain rationality and integrality condition. The most basic example of such a group is the group  $\mathrm{SL}_n(\mathbb{Z}) \subset \mathrm{SL}_n(\mathbb{R})$ . More generally we can start from an algebraic subgroup  $G/\mathbb{Q} \subset \mathrm{GL}_n/\mathbb{Q}$ , for instance the orthogonal group of a quadratic form. Then we get arithmetic groups  $\Gamma \subset \mathbb{G}(\mathbb{Q}) \subset G(\mathbb{R})$  if we impose certain integrality conditions on the matrix coefficients of the elements of  $\Gamma$ .

Now we consider rational representations  $\rho : G/\mathbb{Q} \rightarrow \mathcal{M}_{\mathbb{Q}}$ , where  $\mathcal{M}_{\mathbb{Q}}$  is a finite dimensional  $\mathbb{Q}$ -vector space. We can find finitely generated  $\mathbb{Z}$  modules  $\mathcal{M}$  such that  $\mathcal{M}_{\mathbb{Q}} = \mathcal{M} \otimes_{\mathbb{Z}} \mathbb{Q}$  which are  $\Gamma$ -invariant and hence  $\Gamma$ -modules.

The main object of interest are the cohomology groups

$$H^{\bullet}(\Gamma, \mathcal{M}) = \bigoplus_q H^q(\Gamma, \mathcal{M}).$$

These cohomology groups are defined in terms of homological algebra, they are the derived functors of the functor  $\mathcal{M} \rightarrow \mathcal{M}^{\Gamma}$  (= invariants under  $\Gamma$ .)

These cohomology groups are of course defined for any group  $\Gamma$  and any  $\Gamma$ -module, for us it is important that  $\Gamma \subset G(\mathbb{R})$  where  $G$  is a reductive or even semi simple group, for instance  $\mathrm{SL}_n(\mathbb{R})$ .

Let  $K_{\infty} \subset G(\mathbb{R})$  be a maximal compact subgroup, for example  $\mathrm{SO}(n) \subset \mathrm{SL}_n(\mathbb{R})$ . The quotient  $X = \mathbb{G}(\mathbb{R})/K_{\infty}$  is a symmetric space, it carries a Riemannian metric which is  $G(\mathbb{R})$ -invariant under the left action, it may have finitely many connected components, each connected component is diffeomorphic to  $\mathbb{R}^d$ , hence contractible.

Our arithmetic group  $\Gamma$  acts properly discontinuously on  $X$  we can form the quotient  $\Gamma \backslash X$ , this quotient is an orbifold. We can pass to a suitable subgroup of finite index  $\Gamma' \subset \Gamma$  such that  $\Gamma'$  has no non trivial elements of finite order (i.e. is torsion free). Then  $\Gamma' \backslash X$  is a Riemannian manifold, it is a so called locally symmetric space. The map  $\Gamma' \backslash X \rightarrow \Gamma \backslash X$  is a finite covering with some ramifications. If  $\Gamma$  has elements of finite order then  $\Gamma \backslash X$  is only a Riemannian orbifold. These spaces  $\Gamma \backslash X$  provide a very interesting class of spaces, which are of interest for differential geometers, mathematicians interested in analysis on manifolds and topologists. But they are in a sense of arithmetic origin and therefore they are of interest for number theorists.

Our  $\Gamma$  module  $\mathcal{M}$  endues the space  $\Gamma \backslash X$  with a sheaf  $\tilde{\mathcal{M}}$  (section 6.2) with values in finitely generated abelian groups. If  $\Gamma$  is torsion free then  $\tilde{\mathcal{M}}$  is a locally constant sheaf, or in other words a local system.

We introduce the sheaf- cohomology groups

$$H^\bullet(\Gamma \backslash X, \tilde{\mathcal{M}}) = \bigoplus_q H^q(\Gamma \backslash X, \tilde{\mathcal{M}})$$

these cohomology groups are "essentially" the same as the above group cohomology groups, these two versions of cohomology become equal, if  $X$  is connected and  $\Gamma$  is torsion free. We will see that these cohomology groups are finitely generated  $\mathbb{Z}$  modules.

We have some additional structure on these cohomology groups. In general the quotient space  $\Gamma \backslash X$  is not compact. Therefore we may also consider the cohomology with compact supports  $H_c^\bullet(\Gamma \backslash X, \tilde{\mathcal{M}})$ . Moreover we will construct a neighborhood  $\dot{\mathcal{N}}(\Gamma \backslash X)$  of "infinity" (see (1.2.8)) and from this we get a long exact sequence

$$\cdots \rightarrow H_c^q(\Gamma \backslash X, \tilde{\mathcal{M}}) \xrightarrow{i_c} H^q(\Gamma \backslash X, \tilde{\mathcal{M}}) \xrightarrow{r} H^q(\dot{\mathcal{N}}(\Gamma \backslash X), \tilde{\mathcal{M}}) \xrightarrow{\delta} H_c^{q+1}(\Gamma \backslash X, \tilde{\mathcal{M}}) \rightarrow \cdots \quad (1)$$

This sequence will be called the fundamental ( long ) exact sequence. We also introduce the "inner cohomology"

$$H_!^q(\Gamma \backslash X, \tilde{\mathcal{M}}) := \ker(r) = \text{Im}(i_c)$$

A second structural ingredient is the Hecke algebra. Under certain conditions we have an action of a big algebra of operators acting on all these cohomology groups and the action commutes with arrows in the fundamental exact sequence.

This is the so called Hecke algebra  $\mathcal{H}$  ( or  $\mathcal{H}_\Gamma$  ), it originates from the structure of the arithmetic group  $\Gamma$ . The group  $\Gamma$  has many subgroups  $\Gamma'$  of finite index, for which we can construct two arrows

$$\Gamma' \backslash X \begin{array}{c} \xrightarrow{p_1} \\ \xrightarrow{p_2} \end{array} \Gamma \backslash X. \quad (2)$$

Such a pair of arrows is also called a correspondence between on  $\Gamma \backslash X$ . Such a correspondence, together with a suitable map  $u : p_1^*(\tilde{\mathcal{M}}) \rightarrow p_2^*(\tilde{\mathcal{M}})$ , induces an endomorphism in the cohomology. These endomorphisms act on all the modules in the exact sequence above and are compatible with the arrows.

*The basic objects of interest in this book are the various cohomology groups, which appear in the fundamental exact sequence, together with the the action of the Hecke algebra  $\mathcal{H}$  on them.*

It is my intention is to keep the exposition as elementary as possible, the text should be readable by graduate students. We will need some background material from algebraic topology and from homological algebra ( cohomology and homology of groups, spectral sequences, sheaf cohomology). This material

is expounded in the first four chapters in [28], of course it can be found in many other textbooks.

In the later chapters (starting from chapter 6) we also need results and concepts from the theory of algebraic groups, the theory of symmetric spaces, arithmetic groups, and reduction theory for arithmetic groups. Furthermore we need results from the theory of representations of real semi-simple groups.

This material is somewhat more advanced, but in the In the first five chapters we all these concepts and results are explained in terms in terms of special examples. Especially the sections on the general reduction theory and the Borel-Serre compactification (section (1.2.8)) could be skipped in a first reading.

For the the Lie groups  $\mathrm{Sl}_2(\mathbb{R})$  and  $\mathrm{Sl}_2(\mathbb{C})$  and their arithmetic subgroups  $\mathrm{Sl}_2(\mathbb{Z})$  and  $\mathrm{Sl}_2(\mathbb{Z}[\sqrt{-1}])$  these prerequisite concepts are easy to explain and we will do so in this book. For instance if  $\Gamma = \mathrm{Sl}_2(\mathbb{Z})$  or more generally a congruence subgroup of finite index the symmetric space  $\mathrm{Sl}_2(\mathbb{R})/K_\infty$  is the upper half plane  $\mathbb{H}_2 = \{z \in \mathbb{C} \mid \Im(z) = y > 0\} = \mathrm{Sl}_2(\mathbb{R})/\mathrm{SO}(2)$ . The quotient space  $\Gamma \backslash \mathbb{H}_2$  is punctured Riemann surface. In this special case we have the  $\Gamma$  module  $\mathcal{M}_n = \{\sum a_\nu X^\nu Y^{n-\nu} \mid a_\nu \in \mathbb{Z}\}$ . We will study the the cohomology groups  $H^1(\Gamma \backslash \mathbb{H}_2, \mathcal{M}_n)$  and their module structure under the Hecke algebra in detail. We will prove some very specific results for these cohomology groups.

In Chapter four we discuss results from the theory of representations of the Lie- groups  $\mathrm{Sl}_2(\mathbb{R})$  and  $\mathrm{Sl}_2(\mathbb{C})$ , and we explain the impact of these results on the cohomology. With these results at hand we formulate the famous Eichler-Shimura isomorphism, and we can sketch its proof. This Eichler-Shimura isomorphism also establishes the connection between  $H^1(\Gamma \backslash \mathbb{H}_2, \tilde{\mathcal{M}}_n) \otimes \mathbb{C}$  and the space of modular forms of weight  $n+2$ . In the second half of this book in Chapter 8 we discuss what is called "Representation theoretic Hodge theory" and the Eichler-Shimura theorem becomes a special case of a much more general theorem.

On the other hand we will show that the results for the special groups  $\mathrm{Sl}_2(\mathbb{Z})$ ,  $\mathrm{Sl}_2(\mathbb{Z}[\sqrt{d}])$ , or suitable subgroups of finite index of them, have deep and interesting consequences. We will discuss the Eisenstein cohomology for these special groups and explain the interaction between special values of  $L$ -functions and the structure of the cohomology. A prototype of such a result is the formula for the denominator of the Eisenstein class (Theorem 5.1.1). It is clear that this result should be a special case of a much more general theorem At this moment it is not clear how far these generalisations reach (See section 9.6.5).

In Chapter 5 we discuss some applications of these results to number theory, and we have to accept some even more advanced topics. We concentrate on the case that  $\Gamma \subset \mathrm{Sl}_2(\mathbb{Z})$  and we will use the fact that- with a grain of salt - the quotient  $\Gamma \backslash \mathbb{H}_2$  is the set of  $\mathbb{C}$ -valued points the moduli space of elliptic curves (with some additional structure). This is also explained in [28],[29].

Then for any prime  $\ell$  the cohomology groups  $H^1(\Gamma \backslash \mathbb{H}_2, \tilde{\mathcal{M}}_n) \otimes \mathbb{Z}_\ell$  are actually  $\ell$ -adic etale cohomology groups, especially we get an action of the Galois-group  $\mathrm{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$  on these  $\ell$ -adic cohomology groups. This action commutes with the action of the Hecke algebra. The insights into the structure of the cohomology groups as Hecke modules provides insights into the structure of the

Galois group  $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ , for instance we discuss the theorem of Herbrand-Ribet ([12], [65])

In Chapter 6 we study the cohomology groups of arithmetic groups in a more general framework. We start from arbitrary reductive groups  $G/\mathbb{Q}$ , we assume some familiarity with the theory of semi-simple real groups and the theory of symmetric spaces. There will be some overlap with the earlier chapters.

We use the adelic language, our locally symmetric spaces will be double coset spaces  $\mathcal{S}_{K_f}^G = G(\mathbb{Q}) \backslash G(\mathbb{A}) / K_\infty K_f$ . Here  $K_f$  is an open compact subgroup of  $G(\mathbb{A}_f)$ , it is the so called level subgroup. These locally symmetric spaces turn out to be disjoint unions of the previous ones.

Again define sheaves  $\tilde{\mathcal{M}}$  on these spaces, this will be sheaves with values in the category of finitely generated  $\mathbb{Z}$ -modules, and we are interested in the various cohomology groups  $H_i^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})$ . in our fundamental exact sequence.(1) We know that all these cohomology groups are finitely generated  $\mathbb{Z}$ -modules. (Raghunathan)

Here we have to work a little bit to define the integral cohomology and to define the action of the Hecke operators on these integral cohomology groups.

In this context the Hecke -algebra becomes a restricted product of local Hecke -algebras, this means  $\mathcal{H}_{K_f} = \bigotimes'_p \mathcal{H}_p$ . The local algebras  $\mathcal{H}_p$  have an identity. The level subgroup  $K_f$  determines a finite set  $\Sigma = \Sigma_{K_f}$  of ramified primes. The sub algebra  $\mathcal{H}^{(\Sigma)} = \bigotimes_{p \notin \Sigma} \mathcal{H}_p$  is a central sub-algebra of  $\mathcal{H}_{K_f}$ . For an unramified prime  $p \notin \Sigma$  the structure of  $\mathcal{H}_p$  is given by the Satake isomorphism. (Theorem 6.3.1).

We may pass to the rational cohomology groups  $H_i^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}} \otimes \mathbb{Q})$ , these are finite dimensional  $\mathbb{Q}$  vector space together with the action of  $\mathcal{H}$ . We will show in section 8.1.7 that the action of  $\mathcal{H}$  on the inner cohomology  $H_i^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}} \otimes \mathbb{Q})$  is semi simple, i.e. each  $\mathcal{H}$  invariant submodule has a  $\mathcal{H}$ -invariant complement. This implies that we can find a finite (normal) extension  $F/\mathbb{Q}$  such that  $H_i^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}} \otimes F)$  is a direct sum of absolutely irreducible  $\mathcal{H}$  module. Therefore we get an isotypical decomposition

$$H_i^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}} \otimes F) = \bigoplus_{\pi_f \in \text{Coh}_i(G, K_f, \mathcal{M})} H_i^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}} \otimes F)(\pi_f)$$

where the  $\pi_f$  denote isomorphism classes of absolutely irreducible  $\mathcal{H}$ -modules. Such an absolutely irreducible Hecke module is the restricted tensor product:  $\pi_f = \bigotimes'_p \pi_p$ . The restriction of  $\pi_f$  to  $\mathcal{H}^{(\Sigma)}$  gives us a homomorphism  $\pi^{(\Sigma)} = \bigotimes_{p \notin \Sigma} \pi_p : \mathcal{H}^{(\Sigma)} \rightarrow \mathcal{O}_F$ .

After that we discuss some general facts concerning these cohomology groups (Poincare duality, homology, adjunction formulas for Hecke operators) and we have a section on the Gauss-Bonnet theorem.

Chapter 7 is somewhat philosophical. We have seen in the previous Chapter 4 and we will also see in Chapter 8 how the cohomology groups after tensoring by  $\mathbb{C}$  are related to the space of automorphic forms. In 1967 R. Langlands formulated a visionary program concerning automorphic forms, this is the Langlands

program. In this Chapter 7 we discuss some of the aspects of this program in the context of cohomology of arithmetic groups. The main player is the Langlands dual group  ${}^{\vee}G/\mathbb{Q}$ .

The Langlands dual group  ${}^{\vee}G/\mathbb{Q}$  has the following purpose: For any absolutely irreducible  $\pi_f$  which occurs non trivially in the cohomology  $H^?(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}} \otimes F)$  and any  $p \notin \Sigma$  the theorem of Satake provides a canonical semi-simple conjugacy class  $\omega_p(\pi_p) \in {}^{\vee}G(F)$ . For any representation  $r : {}^{\vee}G/\mathbb{Q} \rightarrow \mathrm{Gl}(V)$  of the algebraic group  ${}^{\vee}G/\mathbb{Q}$  we can attach an  $L$ -function which is defined as an infinite product

$$L^{(\Sigma)}(\pi_f, r, s) := \prod_{p \notin \Sigma} \frac{1}{\det(\mathrm{Id} - r(\omega_p(\pi_p))p^{-s}|V)} = \prod_{p \notin \Sigma} L(\pi_p, r, s)$$

With some extra effort we can also attach local Euler factors  $L(\pi_p, r, s)$  to the ramified primes  $p \in \Sigma$  and then the  $L$  function is defined as  $L(\pi_f, r, s) = \prod_p L(\pi_p, r, s)$ .

A very bold prediction of the Langlands philosophy says that to any absolutely irreducible  $\pi_f$  which occurs somewhere in the cohomology  $H^?(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}} \otimes F)$  and any representation  $r$  we can find a motive  $\{\mathbb{M}(\pi_f, r)\}$  such that we an equality of  $L$ -functions

$$L(\pi_f, r, s) = L(\mathbb{M}(\pi_f, r), s).$$

It is one of the central themes in this book to investigate the relationship between the  $L$ -functions  $L(\pi_f, r, s)$  (analytic properties, special values) and the structure of the integral cohomology as modules under the Hecke-algebra, a first instance is theorem 5.1.1.

In Chapter 8 we develop the analytic tools for the computation of the cohomology. Here we do not use the language of adèles. We assume that the  $\Gamma$ -module  $\mathcal{M}$  is a  $\mathbb{C}$ -vector space and it is obtained from a rational representation of the underlying algebraic group. In this case one may interpret the sheaf  $\tilde{\mathcal{M}}$  as the sheaf of locally constant sections in a flat bundle, and this implies that the cohomology is computable from the de-Rham-complex associated to this flat bundle. We could even go one step further and introduce a Laplace operator so that we get some kind of Hodge-theory and we can express the cohomology in terms of harmonic forms. Here we encounter serious difficulties since the quotient space  $\Gamma \backslash X$  is not compact. But we will proceed in a slightly different way. Instead of doing analysis on  $\Gamma \backslash X$  we work on  $\mathcal{C}_{\infty}(\Gamma \backslash G(\mathbb{R}))$ . This space is a module under the group  $G(\mathbb{R})$ , which acts by right translations, but we rather consider it as a module under the Lie algebra  $\mathfrak{g}$  of  $G(\mathbb{R})$  on which also the group  $K_{\infty}$  acts, it is a  $(\mathfrak{g}, K)$ -module.

Since our module  $\mathcal{M}$  comes from a rational representation of the underlying group  $G$ , we may replace the de-Rham-complex by another complex

$$H^{\bullet}(\mathfrak{g}, K_{\infty}, \mathcal{C}_{\infty}(\Gamma \backslash G(\mathbb{R})) \otimes \tilde{\mathcal{M}}),$$

this complex computes the so called  $(\mathfrak{g}, K)$ -cohomology. The general principle will be to "decompose" the  $(\mathfrak{g}, K)$ -module  $\mathcal{C}_{\infty}(\Gamma \backslash G(\mathbb{R}))$  into irreducible submodules and therefore to compute the cohomology as the sum of the contributions

of the individual submodules. This is a group theoretic version of the classical approach by Hodge-theory. Again we have to overcome two difficulties. The first one is that the quotient  $\Gamma \backslash G(\mathbb{R})$  is not compact and hence the above decomposition does not make sense.

The second problem is that we have to understand the irreducible  $(\mathfrak{g}, K)$ -modules and their cohomology.

The first problem is of analytical nature, we will give some indication how this can be solved by passing to certain subspaces of the cohomology the so called cuspidal or better the inner cohomology. The central result is the Theorem??.

This result is a partial generalisation of the theorem of Eichler-Shimura, it describes the cuspidal part of the cohomology in terms of irreducible representations occurring in the space of cusp forms and contains some information on the discrete cohomology, which is slightly weaker. (See proposition 8.1.4) We shall also give some indications how it can be proved.

We shall also state some general results concerning the second problem, we briefly recall the construction of the irreducible modules with non trivial  $(\mathfrak{g}, K_\infty)$  cohomology.

We discuss some consequences of Theorem 8.1.1. It implies some vanishing theorems in cohomology, it implies that the inner cohomology is a semi simple module for the Hecke-algebra, and it helps to understand the  $K$ -theory of algebraic number theory.

In the next section we use reduction theory-or better the description of  $\dot{\mathcal{N}}(\Gamma \backslash X, \tilde{\mathcal{M}})$ - to discuss some growth conditions for cohomology classes, basically we show that cohomology classes which given by a weight can be represented by differential forms which have essentially the same weight.

In the second half of this chapter we will resume the discussion of modular symbols.

In the last chapter 9 we discuss the Eisenstein-cohomology. The theorem of Eichler-Shimura describes only a certain part of the cohomology, the Eisenstein-cohomology is meant to fill the gap, it is complementary to the cuspidal cohomology. These Eisenstein classes are obtained by an infinite summation process, which sometimes does not converge and is made convergent by analytic continuation.

In the beginning of this chapter 9 we recall the Borel-Serre compactification, we discuss the spectral sequences induced by the stratification of the Borel-Serre boundary. We continue by recalling the process of constructing Eisenstein cohomology classes by infinite summations and analytic (or meromorphic) continuation. We already discussed Eisenstein cohomology in this book for the case of the special group  $Sl_2(\mathbb{R})$  in chapter 4. For the group  $Gl_2/K$  over a number field we refer to [24]. We have the general theorem of Franke [17], but I think that Franke's theorem is still far away from a final answer, there are many questions open and we have to exploit the various possibilities for applications in number theory. In the rest of this chapter we give an outline of these possible application, we formulate some results and we also formulate some speculative ideas.

Under certain conditions (if the Manin Drinfeld principle is valid) these Eisenstein cohomology classes are actually rational classes ( or classes over some

specific number field). Then we may for instance evaluate on certain cycles and it happens that the result is a special value of an  $L$ -function divided by a period (See for instance chapter 4. ) Hence we can prove rationality results for these modified  $L$ - values. This allows us to prove rationality results for special  $L$ -values. (See [26], [35]).

The central theme of this book is the understanding of the integral cohomology  $H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})$  as a module under the Hecke algebra, for instance we want to understand the denominators of the Eisenstein classes.

In Chapter 9 we formulate the general principle that under suitable conditions this denominator should be related ( divisible?, equal ?) to a certain special value of an  $L$ -function, which occurs in the constant term of the Eisenstein series. The prototype of such a relationship occurs in [31], (actually the "abelian" case is discussed in chapter 5).

This principle ( or conjecture ) can be verified (or falsified) experimentally, on the other hand there is a strategy to prove assuming certain finiteness for mixed Grothendieck motives.

# Chapter 1

## Basic Notions and Definitions

Affgr

### 1.1 Affine algebraic groups over $\mathbb{Q}$ .

A linear algebraic group  $G/\mathbb{Q}$  is a subgroup  $G \subset GL_n$ , which is defined as the set of common zeroes of a set of polynomials in the matrix coefficients where in addition these polynomials have coefficients in  $\mathbb{Q}$ . Of course we cannot take just any set of polynomials the set has to be somewhat special before its common zeroes form a group. The following examples will clarify what I mean:

1.) The group  $GL_n$  is an algebraic group itself, the set of equations is empty. It has the subgroup  $Sl_n \subset GL_n$ , which is defined by the polynomial equation

$$Sl_n = \{x \in GL_n \mid \det(x) = 1\}$$

2.) Another example is given by the orthogonal group of a quadratic form

$$f(x_1, \dots, x_n) = \sum_{i=1}^n a_i x_i^2$$

where  $a_i \in \mathbb{Q}$  and all  $a_i \neq 0$  (this is actually not necessary for the following). We look at all matrices

$$\alpha = \begin{pmatrix} a_{11} & \dots & a_{1n} \\ \vdots & & \vdots \\ a_{n1} & \dots & a_{nn} \end{pmatrix}$$

which leave this form invariant, i.e.

$$f(\alpha \underline{x}) = f(\underline{x})$$

for all vectors  $\underline{x} = (x_1, \dots, x_n)$ . This defines a set of polynomial equations for the coefficient  $a_{ij}$  of  $\alpha$ .

These  $\alpha$  form a group, this is the linear algebraic group  $SO(f)$ .

3.) Instead of taking a quadratic form — which is the same as taking a symmetric bilinear form — we could take an alternating bilinear form

$$\begin{aligned} \langle \underline{x}, \underline{y} \rangle &= \langle x_1, \dots, x_{2n}, y_1, \dots, y_{2n} \rangle = \\ &= \sum_{i=1}^n (x_i y_{i+n} - x_{i+n} y_i) = f(\underline{x}, \underline{y}). \end{aligned}$$

This form defines the symplectic group:

$$Sp_n = \{ \alpha \in GL_{2n} \mid \langle \alpha \underline{x}, \alpha \underline{y} \rangle = \langle \underline{x}, \underline{y} \rangle \}.$$

An Important remark: The reader may have observed that we did not specify a field (or a ring) from which we take the entries of the matrices. This is done intentionally, because we may take the entries from any commutative ring  $R$  which contains the rational numbers  $\mathbb{Q}$  and for which  $1 \in \mathbb{Q}$  is the identity element (this means that  $R$  is a  $\mathbb{Q}$ -algebra). In other words: for any algebraic group  $G/\mathbb{Q} \subset GL_n$  and any  $\mathbb{Q}$ -algebra  $R$  we may define

$$G(R) \subset GL_n(R)$$

as the group of those matrices whose coefficients satisfy the required polynomial equations.

Adopting this point of view we can say that

*A linear algebraic group  $G/\mathbb{Q}$  defines a functor from the category of  $\mathbb{Q}$ -algebras (i.e. commutative rings containing  $\mathbb{Q}$ ) into the category of groups.*

4.) Another example is obtained by the so-called restriction of scalars. Let us assume we have a finite extension  $K/\mathbb{Q}$ , we consider the vector space  $V = K^n$ . This vector space may also be considered as a  $\mathbb{Q}$ -vector space  $V_0$  of dimension  $n[K:\mathbb{Q}] = N$ . We consider the group

$$GL_N/\mathbb{Q}.$$

Our original structure as a  $K$ -vector space may be considered as a map

$$\Theta : K \longrightarrow \text{End}_{\mathbb{Q}}(V_0),$$

and the group  $GL_n(K)$  is then the subgroup of elements in  $GL_N(\mathbb{Q})$  which commute with all the elements of  $\Theta(x), x \in K$ . Hence we define the subgroup

$$G/\mathbb{Q} = R_{K/\mathbb{Q}}(GL_n) = \{ \alpha \in GL_N \mid \alpha \text{ commutes with } \Theta(K) \}. \quad (1.1)$$

Then  $G(\mathbb{Q}) = GL_n(K)$ . For any  $\mathbb{Q}$ -algebra  $R$  we get

$$G(R) = GL_n(K \otimes_{\mathbb{Q}} R).$$

This can also be applied to an algebraic subgroup  $H/K \hookrightarrow GL_n/K$ , i.e. a subgroup that is defined by polynomial equations with coefficients in  $K$ .

Our definition of a linear algebraic group is a little bit provisorial. If we consider for instance the two linear algebraic groups

$$\begin{aligned} G_1/\mathbb{Q} &= \left\{ \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \right\} \subset \mathrm{Gl}_2 \\ G_2/\mathbb{Q} &= \left\{ \begin{pmatrix} 1 & 0 & x \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right\} \subset \mathrm{GL}_3 \end{aligned}$$

then we would like to say, that these two groups are isomorphic. They are two different “realizations” of the additive group  $G_a/\mathbb{Q}$ . We see that the same linear algebraic group may be realized in different ways as a subgroup of different  $\mathrm{GL}_N$ 's.

Of course there is a concept of linear algebraic group which does not rely on embeddings. The understanding of this concept requires a little bit of affine algebraic geometry. The drawback of our definition here is that we cannot define morphism between linear algebraic group. Especially we do not know when they are isomorphic.

We assert the reader that the general theory implies that a morphism between two algebraic groups is the same thing as a morphism between the two functors from  $\mathbb{Q}$ -algebras to groups. In some sense it is enough to give this functor. For instance, we have the multiplicative group  $\mathbb{G}_m/\mathbb{Q}$  given by the functor

$$R \longrightarrow R^\times$$

and the additive group  $G_a/\mathbb{Q}$  given by  $R \rightarrow R^+$ .

We can realise (represent is the right term) the the group  $\mathbb{G}_m/\mathbb{Q}$  as

$$\mathbb{G}_m/\mathbb{Q} = \left\{ \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix} \right\} \subset \mathrm{Gl}_2$$

AGS

### 1.1.1 Affine group schemes

We say just a few words concerning the systematic development of the theory of linear algebraic groups. This is not directly used in the next few chapters but it will be useful in Chapter 8.

For any field  $k$  an affine  $k$ -algebra is a finitely generated algebra  $A/k$ , i.e. it is a commutative ring with identity, containing  $k$ , the identity of  $k$  is equal to the identity of  $A$ , which is finitely generated over  $k$  as an algebra. In other words

$$A = k[x_1, x_2, \dots, x_n] = k[X_1, X_2, \dots, X_n]/I,$$

where they  $X_i$  are independent variables and where  $I$  is a finitely generated ideal of polynomials in  $k[X_1, \dots, X_n]$ .

Such an affine  $k$ -algebra defines a functor from the category of  $k$ -algebras to the category of sets, namely  $B \mapsto \mathrm{Hom}_k(A, B)$ . A structure of a group scheme on  $A/k$  consists of the following data:

- a) A  $k$  homomorphism  $m : A \rightarrow A \otimes_k A$  (the comultiplication)
- b) A  $k$ -valued point  $e : A \rightarrow k$  (the identity element)

c) An inverse  $inv : A \rightarrow A$ ,

which satisfy the following requirement: For any  $k$ -algebra  $B$  our homomorphism  $m$  induces a map

$${}^t m : \text{Hom}_k(A \otimes_k A, B) = \text{Hom}_k(A, B) \times \text{Hom}_k(A, B) \rightarrow \text{Hom}_k(A, B)$$

and we require that this induces a group structure on  $\text{Hom}_k(A, B)$ . We also require that the  $k$  valued point  $e$  is the identity and that  $inv$  yields the inverse.

We leave it to the reader to figure out what this means for  $m, e, inv$ , especially what does associativity mean (Hint: Choose  $B = A$ ).

An affine  $k$ -algebra  $A$  together with such a collection  $m, e, inv$  is called an affine group scheme  $G/k = (A, m, e, inv)$ . The  $k$ -algebra  $A$  is the coordinate ring, or the ring of *regular functions* of the group scheme. We will denote it by  $A(G)$ . The group of  $B/k$  valued points will be denoted by  $G(B) = \text{Hom}_k(A(G), B)$ . For  $g \in G(B)$  and  $f \in A(G) \otimes B$  we write  $g(f) = f(g)$ , we evaluate the regular function at the point  $g \in G(B)$ .

The group  $\mathbb{G}_m$  has the coordinate ring  $A(\mathbb{G}_m) = k[t, t^{-1}]$ ,  $m(t) = t \otimes t$ ,  $e(t) = 1$ ,  $inv(t) = t^{-1}$  and the coordinate ring of the additive group  $\mathbb{G}_a$  is  $A(\mathbb{G}_a) = k[x]$  and  $m(x) = x \otimes 1 + 1 \otimes x$ ,  $e(x) = 0$ ,  $inv(x) = -x$ .

The group scheme  $\text{Gl}_n/k$  has the coordinate ring

$$A = k[\dots, x_{i,j}, \dots, y]/(\det(x_{i,j})y - 1); 1 \leq i, j, \leq n$$

and the comultiplication is given by

$$m(x_{i,j}) = \sum_{\nu=1}^n x_{i,\nu} \otimes x_{\nu,j} \tag{1.2}$$

It is clear what a homomorphism between affine group schemes is. A homomorphism  $\phi : G \rightarrow H$  is surjective (resp. injective) if the homomorphism  ${}^t \phi : A(H) \rightarrow A(G)$  is injective (resp.) surjective.

A *rational representation* of  $G/k$  is a homomorphism of group schemes  $\rho : G/k \rightarrow \text{Gl}_n/k$ .

If for instance  $V/k$  is a vector space of dimension  $n$  then we can define the group scheme  $\text{Gl}(V)$ , if we choose a  $k$ -basis on  $V$ , then we can identify  $\text{Gl}(V)/k = \text{Gl}_n/k$ . If  $G/k$  is any affine group scheme, we say that  $V/k$  is a  $G$ -module if we have a homomorphism  $\rho : G/k \rightarrow \text{Gl}(V)$ . Hence we know that for any  $k$ -algebra  $B/k$  we have a homomorphism  $\rho(B) : G(B) \rightarrow \text{Gl}(V \otimes_k B)$ . Of course this is functorial in  $B/k$ , i.e. a homomorphism  $\psi : B/k \rightarrow B'/k$  induces a homomorphism  $G(B) \rightarrow G(B')$ .

We may also consider actions of  $G/k$  on vector spaces  $W/k$  which are not of finite dimension, here we require a certain finiteness condition. As before we have an action

$$\rho_B : G(B) \times (W \otimes B) \rightarrow W \otimes B \tag{1.3}$$

which is functorial in  $B/k$ . But now we assume in addition that for any  $w \in W$  there is a finite set of elements  $w_1, w_2, \dots, w_d$  such that for any  $B/k$  and any  $g \in G(B)$

$$\rho_B(g)w = \sum_{i=1}^d w_i \otimes b_i(g) \text{ with } b_i \in A(G).$$

It suffices to check this for the "universal" element  $\text{Id} \in \text{Hom}_k(A(G), A(G)) = G(A(G))$ , this means we have to find  $w_1, w_2, \dots, w_d \in W$  such that

$$\rho_{A(G)}(\text{Id})w = \sum_{i=1}^d h_i \otimes w_i \text{ with } h_i \in A(G).$$

This implies of course that the  $k$ -subspace  $W' = \sum kw_i$  which is generated by these  $w_i$  is invariant under the action  $\rho$  and it contains  $w$ . Hence we see that our  $k$ -vector space  $W$  is a union of finite dimensional subspaces which are invariant under the action of  $G/k$ .

Therefore we say that a vector space  $W/k$  with an action of  $G/k$  is a  $G$ -module if it satisfies the above finiteness condition.

The ring of regular functions  $A(G)$  is a  $G \times_k G$  module: For  $(g_1, g_2) \in G \times_k G(B) = G(B) \times G(B)$  the action and  $f \in A(G), x \in G(B)$  the action is defined by

$$\rho(g_1, g_2)f(x) = f(g_1^{-1}xg_2).$$

We have to verify the finiteness condition. To do this we write a formula for  $\rho(g_1, g_2)f \in A(G) \otimes B$ . We have the comultiplication  $m : A(G) \rightarrow A(G) \otimes_k A(G)$ , we apply it to the first factor on the right hand side and get  $m_{1,2} \circ m : A(G) \rightarrow A(G) \otimes_k A(G) \otimes_k A(G)$ . Then

$$m_{1,2} \circ m(f) = \sum_{\mu} h'_{\mu} \otimes h_{\mu} \otimes h''_{\mu}$$

Then by definition

$$\rho(g_1, g_2)f = \sum_{\mu} h_{\mu} \otimes \text{inv}(h'_{\mu})(g_1)h''_{\mu}(g_2)$$

and this says that  $\rho(g_1, g_2)f$  lies in the submodule generated by the  $h_{\mu}$ .

Of course we may restrict the action to each the two factors, we simply choose  $g_1 = e$ , -we get the action by right translations- or we choose  $g_2 = e$ , this gives the action by left translations.

It is not difficult to show that for an affine group scheme we can find a collection of elements  $e_0, e_1, \dots, e_r \in A(G)$  such that  $e_i^2 = e_i \forall i, e_i e_j = 0 \forall i \neq j$  such that  $1_A = \sum_i e_i$  and such that the subalgebras  $A(G)e_i$  are integral. Then there is exactly one element (say  $e_0$ ) such that  $e(e_0) = 1$ . Then  $A(G)e_0$  is a subgroup scheme, it is called the *connected component of the identity*. (See for instance [29], Chap. 7 , 7.2)

A group scheme  $G/k$  is *connected*, if its affine algebra  $A(G) = A(G)e_0$  is integral.

**Base change**

If we have a field  $L \supset k$  and a linear group  $G/k$  then the group  $G/L = G \times_k L$  is the group over  $L$  where we forget that the coefficients of the equations are contained in  $k$ . The group  $G \times_k L$  is the *base extension* from  $G/k$  to  $L$ .

**Tori, their character module,...**

A special class of algebraic groups is given by the *tori*. An algebraic group  $T/k$  over a field  $k$  is called a *split torus* if it is isomorphic to a product of  $\mathbb{G}_m/k$ -s,

$$T/k \xrightarrow{\sim} \mathbb{G}_m^d.$$

The algebraic group  $T/k$  is called a torus if it becomes a split torus after a suitable finite extension of the ground field, i.e we have  $T \times_k L \xrightarrow{\sim} \mathbb{G}_m^r/L$ .

If we take an arbitrary finite field extension  $L/\mathbb{Q}$  we may consider the functor

$$R \rightarrow (L \otimes_{\mathbb{Q}} R)^{\times}.$$

It is not hard to see that this functor can be represented by an algebraic group over  $\mathbb{Q}$ , which is denoted by  $R_{L/\mathbb{Q}}(\mathbb{G}_m/L)$  and called the Weil restriction of  $\mathbb{G}_m/L$ . We propose the notation

$$R_{L/\mathbb{Q}}(\mathbb{G}_m/L) = \mathbb{G}_m^{L/\mathbb{Q}} \tag{1.4}$$

The reader should try to prove that for a finite extension  $\tilde{L}/L$  which is normal over  $\mathbb{Q}$  we have

$$\mathbb{G}_m^{L/\mathbb{Q}} \times_{\mathbb{Q}} \tilde{L} \xrightarrow{\sim} (\mathbb{G}_m/\tilde{L})^{[L:\mathbb{Q}]}$$

and this shows that  $\mathbb{G}_m^{L/\mathbb{Q}}$  is a torus .

A torus  $T/k$  is called *anisotropic* if it does not contain a non trivial split torus. Any torus  $C/k$  contains a maximal split torus  $S/k$  and a maximal anisotropic torus  $C_1/k$ . The multiplication induces a map

$$m : S \times C_1 \rightarrow C$$

this is a surjective (in the sense of algebraic groups) homomorphism whose kernel is a finite algebraic group. We call such map an *isogeny* and we write that  $C = S \cdot C_1$ , we say that  $C$  is the product of  $S$  and  $C_1$  up to isogeny.

We give an example. Our torus  $R_{L/\mathbb{Q}}(\mathbb{G}_m/L)$  contains  $\mathbb{G}_m/\mathbb{Q}$  as a subtorus: For any ring  $R$  containing  $\mathbb{Q}$  we have  $R^{\times} = \mathbb{G}_m(R) \subset (R \otimes L)^{\times}$ . On the other hand we have the norm map  $N_{L/\mathbb{Q}} : (R \otimes L)^{\times} \rightarrow R^{\times}$  and the kernel defines a subgroup

$$R_{L/\mathbb{Q}}^{(1)}(\mathbb{G}_m/L) \subset R_{L/\mathbb{Q}}(\mathbb{G}_m/L)$$

and it is clear that

$$m : \mathbb{G}_m \times R_{L/\mathbb{Q}}^{(1)}(\mathbb{G}_m/L) \rightarrow R_{L/\mathbb{Q}}(\mathbb{G}_m/L)$$

has a finite kernel which is the finite algebraic group of  $[L : \mathbb{Q}]$ -th roots of unity.

For any torus  $T = \mathbb{G}_m^r$  we define the character module as the group of homomorphisms

$$X^*(T) = \text{Hom}(T, \mathbb{G}_m).. \quad (1.5)$$

If the torus is split, i.e.  $T = \mathbb{G}_m^r$  then  $X^*(T) = \mathbb{Z}^r$  and the identification is given by  $(n_1, n_2, \dots, n_r) \mapsto \{(x_1, x_2, \dots, x_r) \mapsto x_1^{n_1} x_2^{n_2} \dots x_r^{n_r}\}$ . We write the group structure on  $X^*(T)$  additively, this means that  $(\gamma_1 + \gamma_2)(x) = \gamma_1(x)\gamma_2(x)$ .

It is a theorem that for any torus  $T/K$  we can find a finite, separable, normal extension  $L/K$  such that  $T \times_K L$  splits. Then it is easy to see that we have an action of the Galois group  $\text{Gal}(L/K)$  on  $X^*(T \times_K L) = \mathbb{Z}^r$ . If we have two tori  $T_1/K, T_2/K$  which split over  $L$

$$\text{Hom}_K(T_1, T_2) \xrightarrow{\sim} \text{Hom}_{\text{Gal}(L/K)}(X^*(T_2 \times_K L), X^*(T_1 \times_K L)) \quad (1.6)$$

To any  $\text{Gal}(L/K)$ -action on  $\mathbb{Z}^n$  we can find a torus  $T/K$  which splits over  $L$  and which realises this action.

A homomorphism  $\phi : T_1/K \rightarrow T_2$  is called an isogeny if  $\dim(T_1) = \dim(T_2)$  and if  ${}^t\phi : X^*(T_2) \rightarrow X^*(T_1)$  is injective.

### Semi-simple groups, reductive groups,.

An important class of linear algebraic groups is formed by the *semisimple* and the *reductive* groups. (For a general reference [80].) We do not want to give the precise definition here. Roughly, a linear group is reductive if it is connected and if it does not contain a non trivial normal subgroup which is isomorphic to a product of groups of type  $G_a$ . A group is called semisimple, if it is reductive and does not contain a non trivial torus in its centre.

A semi-simple group  $G/k$  is simple, if it does not contain any normal subgroup of dimension  $> 0$ . Any semi-simple group is up to isogeny a product of simple groups. Any semi simple group  $G/\mathbb{Q}$  contains a maximal torus  $T/\mathbb{Q} \subset G/\mathbb{Q}$  such a maximal torus is equal to its own centraliser. A semi simple group is split if it contains a split maximal torus, i.e. a maximal torus which is split. There is a finite extension  $L/\mathbb{Q}$  such that  $T \times_{\mathbb{Q}} L$  is split, then  $G \times_{\mathbb{Q}} L$  is also split.

For example the groups  $\text{Sl}_n, \text{Sp}_n$  are (split) semi simple, the groups  $\text{SO}(f)$  are semi-simple provided  $n \geq 3$ . (See next subsection 1.1.2 ) groups  $\text{Gl}_n$  and especially the multiplicative group  $\text{Gl}_1/\mathbb{Q} = \mathbb{G}_m/\mathbb{Q}$  are reductive. Any reductive group  $G/\mathbb{Q}$  (or over any field of characteristic zero) has a central torus  $C/\mathbb{Q}$  and this central torus contains a maximal split torus  $S$ . The derived group  $G^{(1)}/\mathbb{Q}$  is semi simple and we get an isogeny

$$G^{(1)} \times C_1 \times S \rightarrow G$$

or briefly  $G = G^{(1)} \cdot C_1 \cdot S$ .

If for instance  $G = R_{L/\mathbb{Q}}(\text{Gl}_n/L)$  then  $G^{(1)} = R_{L/\mathbb{Q}}(\text{Sl}_n/L)$  and  $C = R_{L/\mathbb{Q}}(\mathbb{G}_m/L)$  and this yields the product decomposition up to isogeny

$$G = G^{(1)} \cdot R_{L/\mathbb{Q}}^{(1)}(\mathbb{G}_m/L) \cdot \mathbb{G}_m. \quad (1.7)$$

For  $\mathrm{Gl}_n/\mathbb{Q}$  the central torus is the group  $\mathbb{G}_m/\mathbb{Q}$  the center of  $\mathrm{Sl}_n/\mathbb{Q}$  is the finite group scheme  $\mu_n$  of  $n$ -th roots of unity. The coordinate ring of  $\mu_n$  is the finite algebra  $A(\mu_n) = \mathbb{Q}[t]/(t^n - 1)$ . Of course we may replace  $\mathbb{Q}$  by any ring commutative ring  $R$ .

We can form the quotient group scheme

$$\mathrm{PGL}_n/\mathbb{Q} = (\mathrm{Gl}_n/\mathbb{G}_m)/\mathbb{Q} \xrightarrow{\sim} (\mathrm{Sl}_n/\mathbb{Q})/\mu_n \quad (1.8)$$

this is also the adjoint group of  $\mathrm{Gl}_n/\mathbb{Q}$  and  $\mathrm{Sl}_n/\mathbb{Q}$ , i.e.

$$\mathrm{Ad}(\mathrm{Gl}_n) = \mathrm{PGL}_n = \mathrm{Gl}_n/\mathbb{G}_m = \mathrm{Sl}_n/\mu_n. \quad (1.9)$$

We could certainly drop the assumption that a reductive group should be connected, we could simply say that  $G/\mathbb{Q}$  is reductive (semi-simple...) if its connected component of the identity is reductive (semi-simple...).

Another important class of semi simple groups is given by the *quasisplit* groups. A group  $G/\mathbb{Q}$  is called quasisplit if it contains a Borel subgroup  $B/\mathbb{Q} \subset G/\mathbb{Q}$ . A Borel subgroup  $B/\mathbb{Q}$  is a maximal solvable subgroup, it contains a maximal torus  $T/\mathbb{Q} \subset B/\mathbb{Q}$ , this torus is also a maximal torus in  $G/\mathbb{Q}$ . Then  $B = U \rtimes T$  is the semidirect product of this torus and the *unipotent radical*  $U/\mathbb{Q}$ . We discuss a special example which is of great relevance for our subject.

Let  $L/\mathbb{Q}$  be a quadratic extension, let us denote the non trivial automorphism by  $a \mapsto \bar{a}$ . Let  $V/L$  be a finite dimensional vector space together with a hermitian form  $h : V \times_L V \rightarrow L$ , i.e. we have

$$h(v, w) = \overline{h(w, v)}; h(\lambda u + \mu v, w) = \lambda h(u, w) + \mu h(v, w) \quad \forall u, v, w \in V, \lambda, \mu \in L.$$

Furthermore we assume that  $h$  is non degenerate, i.e. for any  $v \in V, v \neq 0$  we find a  $w \in V$  such that  $h(v, w) \neq 0$ . Then we can define the group  $\mathrm{SU}(h)/\mathbb{Q}$ : For any commutative  $\mathbb{Q}$ -algebra  $R$  we have

$$\mathrm{SU}(h)(R) = \{g \in \mathrm{Sl}(V \otimes_{\mathbb{Q}} R) \mid h(gv, gw) = h(v, w) \text{ and } \det(g) = 1\} \quad (1.10)$$

Then  $\mathrm{SU}(h)/\mathbb{Q}$  is a semi simple group over  $\mathbb{Q}$ . We can also define the unitary group  $\mathrm{U}(h)/\mathbb{Q}$  where we drop the condition that the determinant is one and the group of hermitian similitudes  $\mathrm{GU}(h)$  where

$$\mathrm{GU}(h)(R) = \{g \in \mathrm{Gl}(V \otimes_{\mathbb{Q}} R) \mid h(gv, gw) = d(g)h(v, w) \quad \forall v, w \in V \otimes_{\mathbb{Q}} R\}, \quad (1.11)$$

here  $d : \mathrm{GU}(h) \rightarrow R_{L/\mathbb{Q}}(\mathbb{G}_m)$  is a homomorphism, the kernel of  $d$  is the group  $\mathrm{U}(h)$ .

We consider the special case where

$$V = Le_1 \oplus \cdots \oplus Le_n \oplus (Le_0) \oplus Lf_n \oplus \cdots \oplus Lf_1$$

the summand  $Le_0$  is left out if  $\dim_L V$  is even. The hermitian scalar product is given by

$$h_1(e_i, f_i) = h_1(f_i, e_i) = 1 \quad \forall i = 1, \dots, n \quad (h_1(e_0, e_0) = 1)$$

and all other scalar products equal to zero. Then  $SU(h_1)$  is a quasi split semi simple group over  $\mathbb{Q}$ : The elements  $t \in \text{Gl}(V)$  for which

$$t = \{t : e_i \mapsto t_i e_i; t : f_i \mapsto \bar{t}_i^{-1}; (t : e_0 \mapsto t_0 e_0 \text{ with } t_0 \bar{t}_0 = 1)\}$$

are the  $\mathbb{Q}$ -valued points of a maximal torus  $T_1/\mathbb{Q} \subset SU(h_1)$ . The vector space  $V/L$  comes with a natural flag

$$\begin{aligned} \mathcal{F} := \{0\} \subset Le_1 \subset \dots \subset \oplus Le_1 \oplus \dots \oplus Le_n \subset (Le_1 \oplus \dots \oplus Le_n + Le_0) \subset \\ (Le_1 \oplus \dots \oplus Le_n \oplus Le_0 \oplus Lf_n) \subset \dots \subset (Le_1 \oplus \dots \oplus Le_n \oplus Le_0 \oplus Lf_n \oplus \dots \oplus Lf_2) \subset V \end{aligned} \quad (1.12)$$

Now the subgroup  $B_1/\mathbb{Q} \subset SU(h_1)/\mathbb{Q}$  which fixes  $\mathcal{F}$  is a maximal solvable subgroup in  $SU(h_1)$ .

**To do: classification of quasi-split groups**

### 1.1.2 The classical groups and their realisation as split semi-simple group schemes over $\text{Spec}(\mathbb{Z})$

We will not discuss the general notion of a semi-simple group scheme over a base  $S$ , instead we will discuss the examples of classical groups and explain the main structure theorems in examples.

#### The group scheme $\text{Sl}_n/\text{Spec}(\mathbb{Z})$

We consider a free module  $M$  of rang  $n$  over  $\text{Spec}(\mathbb{Z})$ . We define the group scheme  $\text{Sl}(M)/\text{Spec}(\mathbb{Z})$ : for any  $\mathbb{Z}$  algebra  $R$  we have  $\text{Sl}(M)(R) = \text{Sl}(M \otimes_{\mathbb{Z}} R)$ .

This is clearly a semi simple group scheme over  $\text{Spec}(\mathbb{Z})$  because :

- a) The group scheme is smooth over  $\text{Spec}(\mathbb{Z})$
- b) For any field  $k$  -which is of course a  $\mathbb{Z}$ -algebra we have

$$\text{Sl}(M) \times_{\text{Spec}(\mathbb{Z})} \text{Spec}(k) = \text{Sl}(M \otimes_{\mathbb{Z}} k)/\text{Spec}(k)$$

and for any  $k$  this group scheme does not contain a normal subgroup scheme, which is isomorphic to  $G_a^r/\text{Spec}(k)$  (hence it is reductive) and its center is a finite group scheme.

If we fix a basis  $e_1, e_2, \dots, e_n$  then we get a split maximal torus  $T/\text{Spec}(\mathbb{Z})$  this is the sub group scheme which fixes the lines  $\mathbb{Z}e_i$ , with respect to this basis we have

$$T(R) = \left\{ \begin{pmatrix} t_1 & 0 & \dots & 0 \\ 0 & t_2 & \dots & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & t_n \end{pmatrix} \mid t_i \in R^\times, \prod_i t_i = 1 \right\}$$

With respect to this torus  $T/\text{Spec}(\mathbb{Z})$  we define root subgroups. This are smooth subgroup schemes  $U \subset G$  which are isomorphic to the additive group

scheme  $G_a/\text{Spec}(\mathbb{Z})$  and which are normalized by  $T$ . It is clear that these root subgroups are given by

$$\tau_{ij} : G_a \rightarrow \text{Sl}(M)$$

$$\tau_{ij} : x \rightarrow \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & 1 & \dots & 0 & 0 \\ 0 & 0 & \ddots & x & 0 \\ 0 & 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

where the entry  $x$  is placed in the  $i$ -th row and  $j$ -th column. Let us denote the image by  $U_{\alpha_{ij}}$ .

Then we get the relation

$$t\tau_{ij}(x)t^{-1} = \tau_{ij}((t_i/t_j)x)$$

(If I write such a relation then I always mean that  $t, x..$  are elements in  $T(R), G_a(R)...$  for some unspecified  $\mathbb{Z}$ - algebra  $R$ .)

### The root system

The characters

$$\alpha_{ij} : T \rightarrow G_m$$

$$\alpha_{ij} : \begin{pmatrix} t_1 & 0 & \dots & 0 \\ 0 & t_2 & \dots & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & t_n \end{pmatrix} \rightarrow t_i/t_j$$

are from the set  $\Delta$  of simple roots in the character module of the torus. We may select a subset of positive roots

$$\Delta^+ = \{\alpha_{ij} \mid i < j\}.$$

Then the torus  $T$  and the  $U_{\alpha_{ij}}$  with  $\alpha_{ij} \in \Delta^+$  stabilize the flag

$$\mathcal{F} = (0) \subset \mathbb{Z}e_1 \subset \mathbb{Z}e_1 \oplus \mathbb{Z}e_2 \subset \dots \subset M.$$

The stabilizer of the flag is a smooth sub group scheme  $B/\text{Spec}(\mathbb{Z})$ . It is so-but not entirely obvious- that  $B$  is a maximal solvable sub group scheme. These maximal subgroup schemes are called Borel subgroups.

It is clear that the morphism

$$T \times \prod_{\alpha_{ij}, i < j} U_{\alpha_{ij}} \rightarrow B,$$

which is induced by the multiplication is an isomorphism of schemes.

The set  $\Delta^+$  of positive roots contains the subset  $\pi \subset \Delta$  of simple roots  $t_i/t_{i+1}$ . Every positive root can be written as a sum of simple roots with positive coefficients.

**The flag variety**

It is not so difficult to see that the flags form a projective scheme  $\text{Gr}/\text{Spec}(\mathbb{Z})$ . From this it follows:

For any Dedekind ring  $A$  and its quotient field  $K$  we have

$$\text{Gr}(K) = \text{Gr}(A).$$

If  $A$  is even a discrete valuation ring then we can show easily  
The group  $\text{Sl}_n(A)$  acts transitively on  $\text{Gr}(A)$ .

The whole point is, that results of this type are true for arbitrary split semi simple groups  $\mathcal{G}/\text{Spec}(\mathbb{Z})$ . This is not so easy to explain and also much more difficult to prove. But we have the series of so called classical groups and for those these results are again easy to see. ( The main problem in the general approach is that we have to start from an abstract definition of a semi simple group and not from a group which is given to us in a rather explicit way like  $\text{Sl}_n$  or the classical groups)

**The group scheme  $\text{Sp}_g/\text{Spec}(\mathbb{Z})$** 

Now we choose again a free  $\mathbb{Z}$  module  $M$  but we assume that we have a non degenerate alternating pairing

$$\langle \cdot, \cdot \rangle: M \times M \rightarrow \mathbb{Z}$$

where non degenerate means: If  $x \in M$  and  $\langle x, M \rangle \subset a\mathbb{Z}$  with some integer  $a > 1$ , then  $x = ay$  with  $y \in M$ . It is well known and also very easy to prove that  $M$  is of even rank  $2g$  and that we can find a basis

$$\{e_1, \dots, e_g, f_g, \dots, f_1\}$$

such that  $\langle e_i, f_i \rangle = -\langle f_i, e_i \rangle = 1$  and all other values of the pairing on basis elements are zero.

The automorphism group scheme of  $\mathbb{G} = \text{Aut}((M, \langle \cdot, \cdot \rangle))$  is the symplectic group  $\text{Sp}_g/\text{Spec}(\mathbb{Z})$ . Again it is easy to find out how a maximal torus must look like. With respect to our basis we can take

$$T = \left\{ \begin{pmatrix} t_1 & 0 & & \dots & 0 \\ 0 & \ddots & & \dots & 0 \\ 0 & 0 & t_g & & 0 \\ 0 & 0 & 0 & t_g^{-1} & \dots \\ 0 & & & & \ddots & 0 \\ & 0 & & & & t_1^{-1} \end{pmatrix} \right\}$$

We can say that the torus is the stabilizer of the ordered collection of rank 2 submodules  $\mathbb{Z}e_i, \mathbb{Z}f_i$ . We can define a Borel subgroup  $B/\mathbb{Z}$  which is the stabilizer of the flag

$$\mathcal{F} = (0) \subset \mathbb{Z}e_1 \subset \dots \subset \mathbb{Z}e_1 \cdots \oplus \dots \mathbb{Z}e_g \subset \mathbb{Z}e_1 \cdots \oplus \dots \mathbb{Z}e_g \oplus \mathbb{Z}f_g \subset \dots \subset M$$

(A flag starts with isotropic subspaces until we reach half the rank of the module. But then this lower part of the flag determines the upper half, because it is given by the orthogonal complements of the members in the lower half).

We can define the root subgroups (with respect to  $T$ )

$$\tau_\alpha : G_\alpha \xrightarrow{\sim} U_\alpha \subset \mathbb{G}$$

which are normalized by  $T$ . As before we have the relation

$$t\tau(x)t^{-1} = \tau(\alpha(t)x),$$

where  $\alpha \in \Delta \subset X^*(T)$ .

Now it is not quite so easy to write down what these root subgroups are, we write down the simple positive roots in the the case  $g = 2$ : We have the maximal torus

$$T = \left\{ \begin{pmatrix} t_1 & 0 & 0 & 0 \\ 0 & t_2 & 0 & 0 \\ 0 & 0 & t_2^{-1} & 0 \\ 0 & 0 & 0 & t_1^{-1} \end{pmatrix} \right\}$$

and we want to find one-parameter subgroups  $U_\alpha \subset \mathbb{G}$  which stabilize the flag.

The one parameter subgroups corresponding to the simple roots are

$$\tau_{\alpha_1} : x \mapsto \{e_1 \mapsto e_1, e_2 \mapsto e_2 + xe_1, f_2 \mapsto f_2, f_1 \mapsto f_1 - xf_2\}$$

$$\tau_{\alpha_2} : y \mapsto \{e_1 \mapsto e_1, e_2 \mapsto e_2 + ye_2, f_2 \mapsto f_2 + ye_2, f_1 \mapsto f_1\}$$

where  $\alpha_1(t) = t_1/t_2, \alpha_2(t) = t_2^2$ . The unipotent radical is then

$$\left\{ \begin{pmatrix} 1 & x & v & u \\ 0 & 1 & y & -v \\ 0 & 0 & 1 & -x \\ 0 & 0 & 0 & 1 \end{pmatrix} \right\}$$

As before it is not so difficult to show that the flags form a smooth projective scheme  $\mathcal{X}/\text{Spec}(\mathbb{Z})$  (see also [book], V.2.4.3). Show that for any discrete valuation ring  $A$  the group  $\mathbb{G}(A)$  acts transitively on  $\mathcal{X}(A) = \mathcal{X}(K)$ . It is also easy to verify the statements in 1.1.

### The group scheme $\text{SO}(n, n)/\text{Spec}(\mathbb{Z})$

We can play the same game with symmetric forms. Let  $M$  together with its basis as above, we replace  $g$  by  $n$ . But now we take the quadratic form  $F$

$$F : M \rightarrow \mathbb{Z}$$

which is defined by

$$F(x_1e_1 \cdots + x_n e_n + y_n f_n + \cdots + y_1 f_1) = \sum x_i y_i$$

and all other values of the pairing on basis elements are zero. We define the group scheme of isomorphisms but in addition we require the the determinant is one. Hence

$$\text{SO}(n, n)/\text{Spec}(\mathbb{Z}) = \text{Aut}(M, F, \det = 1).$$

The maximal torus and the flags look pretty much the same as in the previous case. But the set of roots looks different. For  $n = 2$  the torus and the unipotent radical are given by

$$T = \left\{ \begin{pmatrix} t_1 & 0 & 0 & 0 \\ 0 & t_2 & 0 & 0 \\ 0 & 0 & t_2^{-1} & 0 \\ 0 & 0 & 0 & t_1^{-1} \end{pmatrix} \right\}, \quad U = \left\{ \begin{pmatrix} 1 & x & y & -xy \\ 0 & 1 & 0 & -y \\ 0 & 0 & 1 & -x \\ 0 & 0 & 0 & 1 \end{pmatrix} \right\}.$$

The system of positive roots consists of two roots  $\alpha_1(t) = t_1/t_2, \alpha_2(t) = t_1 t_2$ . This is the Dynkin diagram  $A_1 \times A_1$  hence there exists a homomorphism (isogeny) between group schemes over  $\text{Spec}(\mathbb{Z})$ :

$$\text{Sl}_2 \times \text{Sl}_2 \rightarrow \text{SO}(2, 2).$$

It is an amusing exercise to write down this isogeny.

Another even more interesting exercise is the computation of the roots for the torus (here  $n = 3$ )

$$T = \left\{ \begin{pmatrix} t_1 & 0 & 0 & 0 & 0 & 0 \\ 0 & t_2 & 0 & 0 & 0 & 0 \\ 0 & 0 & t_3 & 0 & 0 & 0 \\ 0 & 0 & 0 & t_3^{-1} & 0 & 0 \\ 0 & 0 & 0 & 0 & t_2^{-1} & 0 \\ 0 & 0 & 0 & 0 & 0 & t_1^{-1} \end{pmatrix} \right\}.$$

In this case we have the root subgroups

$$\tau_{\alpha_1} : x \mapsto \begin{pmatrix} 1 & x & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & -x \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad \tau_{\alpha_2} : x \mapsto \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & x & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & -x & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

and

$$\tau_{\alpha_3} : x \mapsto \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & x & 0 & 0 \\ 0 & 0 & 1 & 0 & -x & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

where

$$\alpha_1(t) = t_1/t_2, \quad \alpha_2(t) = t_2/t_3, \quad \alpha_3(t) = t_2 t_3$$

Use the result of this computation to show that we have an isogeny

$$\text{Sl}_4 \rightarrow \text{SO}(3, 3).$$

How can we give a linear algebra interpretation of this isogenies.

**The group scheme  $\mathrm{SO}(n+1, n)/\mathrm{Spec}(\mathbb{Z})$** 

Of course we can also consider quadratic forms in an odd number of variables. We take a free  $\mathbb{Z}$ -module of rank  $2n+1$  with a basis

$$\{e_1, \dots, e_n, h, f_n, \dots, f_1\}.$$

On this module we consider the quadratic form

$$F : M \rightarrow \mathbb{Z}$$

$$F\left(\sum x_i e_i + zh + \sum y_i f_i\right) = \sum x_i y_i + z^2.$$

From this quadratic form we get the bilinear form

$$B(u, v) = F(u+v) - F(u) - F(v).$$

We have the relation

$$F(u) = 2B(u, u),$$

hence we can reconstruct the quadratic form from the bilinear form if we extend  $\mathbb{Z}$  to a larger ring where 2 is invertible.

We consider the automorphism scheme

$$\mathcal{G}/\mathrm{Spec}(\mathbb{Z}) = \mathrm{SO}(n+1, n)/\mathrm{Spec}(\mathbb{Z}) = \mathrm{Aut}(M, F, \det = 1)/\mathrm{Spec}(\mathbb{Z})$$

and I claim that this is indeed a semi simple group scheme over  $\mathrm{Spec}(\mathbb{Z})$ . To see this I strongly recommend to discuss the case  $n=1$ .

We have of course the maximal torus

$$T = \left\{ \begin{pmatrix} t & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & t^{-1} \end{pmatrix} \right\}.$$

It is also the stabilizer of the collection of three subspaces  $\mathbb{Z}e, \mathbb{Z}h, \mathbb{Z}f$ , here we use the determinant condition.

Now one has to discuss the root subgroups with respect to this torus.

From this we can derive that we have an isogeny

$$\mathrm{Sl}_2 \rightarrow \mathrm{SO}(2, 1)$$

It is also interesting to look at the case  $n=2$ . In this case we can compare the root systems of  $\mathrm{Sp}_2$  and  $\mathrm{SO}(3, 2)$  they are isomorphic. Now it is a general theorem in the theory of split semi simple group schemes that the root system determines the group scheme up to isogeny. Hence we should be able to construct an isogeny between  $\mathrm{Sp}_2$  and  $\mathrm{SO}(3, 2)$ . Who can do it?

**1.1.3  $k$ -forms of algebraic groups**

**Exercise:** 1) Consider the following two quadratic forms over  $\mathbb{Q}$ :

$$f(x, y, z) = x^2 + y^2 - z^2, \quad f_1(x, y, z) = x^2 + y^2 - 3z^2.$$

Prove that the first form is isotropic. This means there exists a vector  $(a, b, c) \in \mathbb{Q}^3 \setminus \{0\}$  with

$$f(a, b, c) = 0.$$

Show that the second form is anisotropic, i.e. it has no such vector.

2) Prove that the two linear algebraic group  $G/\mathbb{Q} = SO(f)/\mathbb{Q}$  and  $G_1/\mathbb{Q} = SO(f_1)/\mathbb{Q}$  cannot be isomorphic. (Hint: This is not so easy since we did not define when two groups are isomorphic.)

Here is some advice: In general we call an element  $e \neq u \in G(\mathbb{Q})$  unipotent if it is unipotent in  $GL_n(\mathbb{Q})$  where we consider  $G/\mathbb{Q} \hookrightarrow GL_n/\mathbb{Q}$ . It turns out that this notion of unipotence does not depend on the embedding.

Now it is possible to show that our first group  $G(\mathbb{Q}) = SO(f)(\mathbb{Q})$  has unipotent elements, and  $G_1(\mathbb{Q})$  does not. Hence these two groups cannot be isomorphic.

3) Prove that the two algebraic groups  $G \times_{\mathbb{Q}} \mathbb{R}$  and  $G_1 \times_{\mathbb{Q}} \mathbb{R}$  are isomorphic, and therefore the two groups  $G(\mathbb{R})$  and  $G_1(\mathbb{R})$  are isomorphic.

In this example we see, that we may have two groups  $G/k, G_1/k$  which are not isomorphic but which become isomorphic over some extension  $L/k$ . Then we say that the groups are  $k$ -forms of each other. To determine the different forms of a given group  $G/k$  is sometimes difficult one has to use the concepts of Galois cohomology. For a separable normal extension  $L/k$  we have the almost tautological description

$$G(k) = \{g \in G(L) \mid \sigma(g) = g \text{ for all elements in the Galois group } \text{Gal}(L/k)\}.$$

Now let we can consider the functor  $\text{Aut}(G)$ : It attaches to any field extension  $L/k$  the group of automorphisms  $\text{Aut}(G)(L)$  of the algebraic group  $G \times_k L$ . We denote this action by  $g \mapsto \sigma(g) = g^\sigma$ . Note that this notation gives us the rule  $g^{(\sigma\tau)} = (g^\tau)^\sigma$ . A 1-cocycle of  $\text{Gal}(L/k)$  with values in  $\text{Aut}(G)$  is a map  $c : \sigma \mapsto c_\sigma \in \text{Aut}(G)(L)$  which satisfies the cocycle rule

$$c_{\sigma\tau} = c_\sigma c_\tau^\sigma \quad (1.13)$$

Now we define a new action of  $\text{Gal}(L/k)$  on  $G(L)$ : An element  $\sigma$  acts by

$$g \mapsto c_\sigma g^\sigma c_\sigma^{-1}$$

We define a new algebraic group  $G_1/k$ : For any extension  $E/k$  we have an action of  $\text{Gal}(L/k)$  on  $E \otimes_k L$  and we put

$$G_1(E) = \{g \in G(E \otimes_k L) \mid g = c_\sigma g^\sigma c_\sigma^{-1}\} \quad (1.14)$$

For the trivial cocycle  $\sigma \mapsto 1$  this gives us back the original group.

It is plausible and in fact not very difficult to show that  $E \rightarrow G_1(E)$  is in fact represented by an algebraic group  $G_1/k$ . This group is clearly a  $k$ -form of  $G/k$ .

We can define an equivalence relation on the set of cocycles, we say that

$$\{\sigma \mapsto c_\sigma\} \sim \{\sigma \mapsto c'_\sigma\}$$

if and only if we can find a  $a \in G(L)$  such that

$$c'_\sigma = a^{-1} c_\sigma a^\sigma \text{ for all } \sigma \in \text{Gal}(L/k)$$

We define  $H^1(L/k, \text{Aut}(G))$  as the set of 1-cocycles modulo this equivalence relation. If we have a larger normal separable extension  $L' \supset L \supset k$  then we get an inclusion  $H^1(L/k, \text{Aut}(G)) \hookrightarrow H^1(L'/k, \text{Aut}(G))$ . If  $\bar{k}_s$  is a separable closure of  $k$  we can form the limit over all finite extensions  $k \subset L \subset \bar{k}_s$  and put

$$H^1(\bar{k}_s/k, \text{Aut}(G)) = \varinjlim H^1(L/k, \text{Aut}(G))$$

This set is isomorphic to the set of isomorphism classes of  $k$ -forms of  $G/k$ .

If  $L/k$  is a cyclic extension and if  $\sigma \in \text{Gal}(L/k)$  is a generator, then a cocycle  $c : \text{Gal}(L/k) \rightarrow \text{Aut}(G)(L)$  is determined by its value  $g = c(\sigma) \in \text{Aut}(G)(L)$ . But we have to satisfy the cocycle relation. We have a useful little

**Lemma 1.1.1.** *The assignment  $\sigma \mapsto c(\sigma) = g$  provides a 1-cocycle if and only iff*

$$\text{Norm}(g) = gg^\sigma \dots g^{\sigma^{n-1}} = \text{Id}$$

and

$$H^1(\text{Gal}(L/k), \text{Aut}(G)(L)) = \{g \in \text{Aut}(G)(L) \mid \text{Norm}(g) = \text{Id}\} / \text{hgh}^{-\sigma} \sim g\}.$$

*Proof.* Straightforward calculation □

We may apply the same concepts in a slightly different situation. A  $k$ -algebra  $\mathcal{D}$  over the field  $k$  is called a *central simple algebra*, if it has a unit element  $\neq 0$ , if it is finite dimensional over  $k$ , if its centre is  $k$  (embedded via the unit element) and if it has no non trivial two sided ideals. It is a classical theorem, that such an algebra over a separably closed field  $\bar{k}_s$  is isomorphic to a full matrix algebra  $M_n(\bar{k}_s)$ . Hence we can say that over an arbitrary field  $k$  any central simple algebra of dimension  $n^2$  is a  $k$ -form of  $M_n(k)$ .

For any algebraic group  $G/k$  we may consider the adjoint group  $\text{Ad}(G)$ , this is the quotient of  $G/k$  by its center. It can be shown, that this is again an algebraic group over  $k$ . It is clear that we have an embedding

$$\text{Ad}(G) \rightarrow \text{Aut}(G)$$

which for any  $g \in \text{Ad}(G)(L)$  is given by

$$g \mapsto \{x \mapsto g^{-1}xg\}.$$

A  $k$ -form  $G_1/k$  of a group  $G/k$  is called an *inner  $k$ -form*, if it is in the image of

$$H^1(\bar{k}_s/k, \text{Ad}(G)) \rightarrow H^1(\bar{k}_s/k, \text{Aut}(G)).$$

We call a semi simple group  $G/k$  *anisotropic*, if it does not contain a non trivial split torus (See exercise in (1.1.3)) In our example below the group of elements of norm 1 is always semi simple and anisotropic if and only if  $D(a, b)$  is a field.

I want to give an example, we consider the algebraic group  $\text{Gl}_2/\mathbb{Q}$  we consider two integers  $a, b \neq 0$ , for simplicity we assume that  $b$  is not a square. Then

we have the quadratic extension  $L = \mathbb{Q}(\sqrt{b})$ , let  $\sigma$  be its non trivial automorphism. The element  $\begin{pmatrix} 0 & a \\ 1 & 0 \end{pmatrix}$  defines the inner automorphism

$$\text{Ad}\left(\begin{pmatrix} 0 & a \\ 1 & 0 \end{pmatrix}\right) : g \mapsto \begin{pmatrix} 0 & a \\ 1 & 0 \end{pmatrix} g \begin{pmatrix} 0 & a \\ 1 & 0 \end{pmatrix}^{-1}$$

of the group  $\text{GL}_2$ , Then  $\sigma \mapsto \text{Ad}\left(\begin{pmatrix} 0 & a \\ 1 & 0 \end{pmatrix}\right)$  and  $\text{Id}_{\text{Gal}(L/k)} \mapsto \text{Id}_{\text{Aut}(\text{GL}_2)(L)}$  is a 1-cocycle and we get a  $\mathbb{Q}$  form of our group.

Hence we get a  $\mathbb{Q}$  form  $G_1 = G(a, b)/\mathbb{Q}$  of our group  $\text{GL}_2$ . It is an inner form.

Now we can see easily that group of rational points of our above group  $G(a, b)(\mathbb{Q})$  is the multiplicative group of a central simple algebra  $D(a, b)/\mathbb{Q}$ . To get this algebra we consider the algebra  $M_2(L)$  of (2,2)-matrices over  $L$ . We define

$$D(a, b) = \{x \in M_2(L) \mid x = \text{Ad}\left(\begin{pmatrix} 0 & a \\ 1 & 0 \end{pmatrix}\right) x^\sigma \text{Ad}\left(\begin{pmatrix} 0 & a \\ 1 & 0 \end{pmatrix}\right)^{-1}\}. \quad (1.15)$$

We have an embedding of the field  $L$  into this algebra, which is given by

$$u \mapsto \begin{pmatrix} u & 0 \\ 0 & u^\sigma \end{pmatrix}$$

Let  $u_b$  the image of  $\sqrt{b}$  under this map. We also have the element  $u_a = \begin{pmatrix} 0 & a \\ 1 & 0 \end{pmatrix}$  in this algebra.

Now I leave it as an exercise to the reader that as a  $\mathbb{Q}$  vector space

$$D(a, b) = \mathbb{Q} \oplus \mathbb{Q}u_b \oplus \mathbb{Q}u_a \oplus \mathbb{Q}u_a u_b$$

We have the relation  $u_a^2 = a, u_b^2 = b, u_a u_b = -u_b u_a$ .

Of course we should ask ourselves: When is  $D(a, b)$  split, this means isomorphic to  $M_2(\mathbb{Q})$ ? To answer this question we consider the norm homomorphism, which is defined by

$$x + yu_b + zu_a + wa_a u_b \mapsto (x + yu_b + zu_a + wa_a u_b)(x - yu_b - zu_a - wa_a u_b) = x^2 - y^2 b - z^2 a + w^2 ab.$$

It is easy to see that  $D(a, b)$  splits if and only if we can find a non zero element whose norm is zero.

If we do this over  $\mathbb{R}$  as base field and if we take  $a = -1, b = -1$  then we get the Hamiltonian quaternions, which is non split.

We may also look at the  $p$ -adic completions  $\mathbb{Q}_p$  of our field. Then it is not difficult to see that  $D(a, b)$  splits over  $\mathbb{Q}_p$  if  $p \neq 2$  and  $p \nmid ab$ . Hence it is clear that there is only a finite number of primes  $p$  for which  $D(a, b)$  does not split.

If we consider  $\mathbb{R}$  as completion at the infinite place, and the  $\mathbb{Q}_p$  as the completions at the finite places, then we have

The algebra  $D(a, b)$  splits if and only if it splits at all places. The number of places where it does not split is always even.

The first assertion is the so called Hasse-Minkowski principle, the second assertion is essentially equivalent to the quadratic reciprocity law.

### Construction of division algebras and anisotropic groups

We give some indication how to construct anisotropic groups over  $\mathbb{Q}$  (or even over any number field). We choose a cyclic extension  $L/\mathbb{Q}$  of degree  $n$  and we pick a number  $a \in \mathbb{Q}^\times$ , let  $A(a) \in \mathrm{GL}_n(\mathbb{Q})$  be the following matrix

$$A(a) = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ & & \vdots & \ddots & \\ 0 & 0 & \dots & 0 & 1 \\ a & 0 & 0 & 0 & 0 \end{pmatrix} \quad (1.16)$$

Let  $\sigma \in \mathrm{Gal}(L/\mathbb{Q})$  be a generator then  $\sigma^\nu \mapsto A(a)^\nu \pmod{\mathbb{G}_m}$  is a homomorphism from  $\mathrm{Gal}(L/\mathbb{Q})$  to  $\mathrm{PGL}_n(\mathbb{Q})$  and since  $A(a) \in \mathrm{GL}_n(\mathbb{Q})$  this is also a 1-cocycle  $c: \mathrm{Gal}(L/\mathbb{Q}) \rightarrow \mathrm{PGL}_n(\mathbb{Q}) := \{\sigma^\nu \mapsto A(a)^\nu\}$ . It defines a cohomology class  $[A(a)] \in H^1(L/\mathbb{Q}, \mathrm{Ad}(\mathrm{GL}_n))$  and hence an inner  $\mathbb{Q}$ -form  $G/\mathbb{Q}$  of  $\mathrm{GL}_n/\mathbb{Q}$ . In Galois cohomology we have the boundary map

$$\delta: H^1(L/\mathbb{Q}, \mathrm{Ad}(\mathrm{GL}_n)) \rightarrow H^2(L/\mathbb{Q}, \mathbb{G}_m) = \mathbb{Q}^\times / N_{L/\mathbb{Q}}(L^\times)$$

and it is clear that

$$\delta([A(a)]) = a \in \mathbb{Q}^\times / N_{L/\mathbb{Q}}(L^\times)$$

Now it is well known that the  $\mathbb{Q}$ -form  $G/\mathbb{Q}$  of  $\mathrm{GL}_n/\mathbb{Q}$  is anisotropic if and only if the class  $a \in \mathbb{Q}^\times / N_{L/\mathbb{Q}}(L^\times)$  is an element of order  $n$ . We know from algebraic number theory that there are infinitely many primes  $p$  which are inert, i.e.  $p$  is unramified in  $L$  and the prime ideal  $(p)$  stays prime in the ring of integers  $\mathcal{O}_L$ . Then it is easy to see that the order of  $p \in \mathbb{Q}^\times / N_{L/\mathbb{Q}}(L^\times)$  is  $n$ . Hence we see that the set of isomorphism classes of anisotropic  $\mathbb{Q}$ -forms of  $\mathrm{GL}_n/\mathbb{Q}$  is abundant.

Obviously the group  $M_n(\mathbb{Q})^\times = \mathrm{GL}_n(\mathbb{Q})$  and we also know that any automorphism of  $M_n(\mathbb{Q})^\times$  is inner, hence  $\mathrm{Aut}(M_n(\mathbb{Q})) = \mathrm{PGL}_n(\mathbb{Q})$ . Therefore the isomorphism classes of  $\mathbb{Q}$ -forms of  $M_n(\mathbb{Q})$  are equal to the set  $H^1(\mathbb{Q}, \mathrm{PGL}_n)$ . Such a  $\mathbb{Q}$ -form  $\mathcal{D}/\mathbb{Q}$  is a central simple algebra over  $\mathbb{Q}$ . The central simple algebra  $\mathcal{D}$  defined by the class  $[A(a)]$  can be described explicitly:

It contains the field  $L/\mathbb{Q}$  as a maximal commutative subalgebra and it is generated by  $L$  and another element  $a_\sigma \in \mathcal{D}$  which satisfies the following relations

$$\forall x \in L \text{ we have } a_\sigma x a_\sigma^{-1} = \sigma(x); \quad a_\sigma^n = a$$

If we modify  $a_\sigma$  and put  $a'_\sigma = a_\sigma y$  with  $y \in L^\times$  then the first relation still holds and the second relation becomes  $(a'_\sigma)^n = a N_{L/\mathbb{Q}}(y)$ . Hence the isomorphism class of  $\mathcal{D}$  is determined by the class  $a \in \mathbb{Q}^\times / N_{L/\mathbb{Q}}(L^\times)$ . It is easy to see that for  $a = 1$  the central simple algebra is equal to the endomorphism ring of the  $\mathbb{Q}$ -vector space  $L/\mathbb{Q}$ . (This is the linear independence of the elements  $\sigma^\nu$  in  $\mathrm{End}(L/\mathbb{Q})$ .)

### Quasisplit $\mathbb{Q}$ -forms

We recall that a semi-simple group  $G/\mathbb{Q}$  is quasisplit, if it contains a Borel subgroup  $B/\mathbb{Q}$ . This Borel subgroup contains its unipotent radical  $U/\mathbb{Q}$  and a maximal torus  $T/\mathbb{Q}$ . Two such maximal tori  $T/\mathbb{Q}, T_1/\mathbb{Q}$  are conjugate by an element  $u \in U(\mathbb{Q})$ . Let  $G_0/\mathbb{Q}$  be a split group which is a  $\mathbb{Q}$ -form of  $G/\mathbb{Q}$ . We pick a maximal split torus  $T_0/\mathbb{Q}$  and a Borel  $B_0/\mathbb{Q} \supset T_0/\mathbb{Q}$ . Then we see that the triple  $(G, B, T)/\mathbb{Q}$  is a  $\mathbb{Q}$ -form of  $(G_0, B_0, T_0)/\mathbb{Q}$ . Hence it can be constructed from a 1-cocycle representing a cohomology class  $\xi \in H^1(\mathbb{Q}, \text{Aut}(((G_0, B_0, T_0))))$ , where of course  $\text{Aut}(((G_0, B_0, T_0)))$  is the subgroup of  $\text{Aut}(G_0)$  which fixes  $T_0, B_0$ . Obviously we have an exact sequence

$$1 \rightarrow T_0^{(\text{ad})} \rightarrow \text{Aut}(((G_0, B_0, T_0))) \rightarrow \text{Autext}((G_0, B_0, T_0)) \rightarrow 1, \quad (1.17)$$

here  $\text{Autext}((G_0, B_0, T_0))$  is the very "small" group of automorphisms of the Dynkin diagram  $\Phi$ . It is well known- and easy to see in the examples of classical groups- that this sequence has a section  $s_0 : \text{Autext}((G_0, B_0, T_0)) \rightarrow \text{Aut}(((G_0, B_0, T_0)))$  and this gives us a map in Galois cohomology

$$s_0^\bullet : H^1(\mathbb{Q}, \text{Autext}((G_0, B_0, T_0))) = \text{Hom}(\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}), \text{Autext}((G_0, B_0, T_0))) \rightarrow H^1(\mathbb{Q}, \text{Autext}((G_0))) \quad (1.18)$$

Hence we see that the isomorphism classes of quasisplit  $\mathbb{Q}$ -forms of  $G/\mathbb{Q}$  are given by homomorphisms  $\psi : \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{Gal}(L/\mathbb{Q}) \hookrightarrow \text{Autext}((G_0))$ .

In the special case  $G_0/\mathbb{Q} = \text{SL}_n/\mathbb{Q}$  with  $T_0/\mathbb{Q}, B_0/\mathbb{Q}$  being the standard diagonal torus and the standard Borel subgroup of upper triangular matrices this looks as follows: We have the element

$$w_0 = \begin{pmatrix} 0 & 0 & 0 & \dots & 1 \\ 0 & 0 & \dots & 1 & 0 \\ & & \vdots & \ddots & \\ 0 & 1 & \dots & 0 & 0 \\ \pm 1 & 0 & 0 & 0 & 0 \end{pmatrix} \in \text{SL}_n(\mathbb{Q}) \quad (1.19)$$

this element  $w_0$  conjugates  $B_0$  into its opposite  $B_0^-$  the group of lower triangular matrices. The standard Cartan involution  $\Theta : g \rightarrow {}^t g^{-1}$  does the same and therefore the composition  $\text{Ad}(w_0) \circ \Theta$  is an automorphism of  $G_0/\mathbb{Q}$  which fixes  $B_0, T_0$ . It is an outer automorphism if  $n \geq 3$  and gives us the non trivial element of  $\text{Autext}(G_0)$ . Hence we get a 1-cocycle if we choose a quadratic extension  $L/\mathbb{Q}$  and send the non trivial element in  $\text{Gal}(L/\mathbb{Q})$  to  $\text{Ad}(w_0) \circ \Theta$ .

We leave it an exercise to the reader to show that the  $\mathbb{Q}$  form obtained from this cocycle (cohomology class) is isomorphic to the above group  $\text{SU}(h_1)/\mathbb{Q}$ .

#### 1.1.4 The Lie-algebra

We need some basic facts about the Lie-algebras of algebraic groups.

For any algebraic group  $G/k$  we can consider its group of points with values in  $k[\epsilon] = k[X]/(X^2)$ . We have the homomorphism  $k[\epsilon] \rightarrow k$  sending  $\epsilon$  to zero and hence we get an exact sequence

$$0 \rightarrow \mathfrak{g} \rightarrow G(k[\epsilon]) \rightarrow G(k) \rightarrow 1.$$

The kernel  $\mathfrak{g}$  is a  $k$ -vector space, if the characteristic of  $k$  is zero, then its dimension is equal to the dimension of  $G/k$ . It is denoted by  $\mathfrak{g} = \text{Lie}(G)$ .

Let us consider the example of the group  $G = SO(f)$ , where  $f : V \times V \rightarrow k$  is a non degenerate symmetric bilinear form. In this case an element in  $G(k[\epsilon])$  is of the form  $\text{Id} + \epsilon A, A \in \text{End}(V)$  for which

$$f((\text{Id} + \epsilon A)v, (\text{Id} + \epsilon A)w) = f(v, w)$$

for all  $v, w \in V$ . Taking into account that  $\epsilon^2 = 0$  we get

$$\epsilon(f(Av, w) + f(v, Aw)) = 0,$$

i.e.  $A$  is skew with respect to the form, and  $\mathfrak{g}$  is the  $k$ -vector space of skew endomorphisms. If we give  $V$  a basis and if  $f = \sum x_i^2$  with respect to this basis then this means the the matrix of  $A$  is skew symmetric.

If we consider  $G = \text{Gl}_n/k$  then  $\mathfrak{g} = M_n(k)$ , the Lie-bracket is given by

$$(A, B) \mapsto AB - BA \quad (1.20)$$

We have some kind of a standard basis for our Lie algebra

$$\mathfrak{g} = \bigoplus_{i=1}^n kH_i \oplus \bigoplus_{i,j,i \neq j} kE_{i,j} \quad (1.21)$$

where  $H_i$  (resp.  $E_{i,j}$ ) are the matrices

$$H_i = \begin{pmatrix} 0 & 0 & \dots & 0 & 0 & 0 \\ 0 & 0 & \dots & 0 & 0 & 0 \\ 0 & 0 & \ddots & & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & \dots & \ddots & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \text{ resp. } E_{i,j} = \begin{pmatrix} 0 & 0 & \dots & 0 & 0 & 0 \\ 0 & 0 & \dots & 0 & 0 & 0 \\ 0 & 0 & \ddots & & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \dots & \ddots & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

and the only non zero entries ( $=1$ ) is at  $(i, i)$  on the diagonal (resp. and  $(i, j)$  off the diagonal.)

For the group  $\text{Sl}_n/k$  the Lie-algebra is  $\mathfrak{g}^{(0)} = \{A \in M_n(k) \mid \text{tr}(A) = 0\}$  and again we have a standard basis

$$\mathfrak{g}^{(0)} = \bigoplus_{i=1}^{n-1} k(H_i - H_{i+1}) \oplus \bigoplus_{i,j,i \neq j} kE_{i,j} \quad (1.22)$$

If  $\rho : G \rightarrow \text{Gl}(V)$  is a rational representation of our group  $G/k$  then it is clear from our considerations above that we have a "derivative" of this representation

drho

$$d\rho : \mathfrak{g} = \text{Lie}(G/k) \rightarrow \text{Lie}(\text{Gl}(V)) = \text{End}(V) \quad (1.23)$$

this is  $k$ -linear.

Every group scheme  $G/k$  has a very special representation, this is the the *Adjoint representation*. We observe that the group acts on itself by conjugation, this is the morphism

$$\text{Inn} : G \times_k G \rightarrow G$$

which on  $T$  valued points is given by

$$\text{Inn}(g_1, g_2) \mapsto g_1 g_2 (g_1)^{-1}.$$

This action clearly induces a representation

$$\text{Ad} : G/k \rightarrow \text{Gl}(\mathfrak{g})$$

and this is the adjoint representation. This adjoint representation has a derivative and this is a homomorphism of  $k$  vector spaces

$$D_{\text{Ad}} = \text{ad} : \mathfrak{g} \rightarrow \text{End}(\mathfrak{g}).$$

We introduce the notation: For  $T_1, T_2 \in \mathfrak{g}$  we put

$$[T_1, T_2] := \text{ad}(T_1)(T_2).$$

Now we can state the famous and fundamental result

**Theorem 1.1.1.** *The map  $(T_1, T_2) \mapsto [T_1, T_2]$  is bilinear and antisymmetric. It induces the structure of a Lie-algebra on  $\mathfrak{g}$ , i.e. we have the Jacobi identity*

$$[T_1, [T_2, T_3]] + [T_2, [T_3, T_1]] + [T_3, [T_1, T_2]] = 0.$$

We do not prove this here. In the case  $G/k = \text{Gl}(V)$  and  $T_1, T_2 \in \text{Lie}(\text{Gl}(V) = \text{End}(V))$  we have  $[T_1, T_2] = T_1 T_2 - T_2 T_1$  and in this case the Jacobi Identity is a well known identity.

On any Lie algebra we have a symmetric bilinear form (the Killing form)

$$B : \mathfrak{g} \times \mathfrak{g} \rightarrow k \tag{1.24}$$

which is defined by the rule

$$B(T_1, T_2) = \text{trace}(\text{ad}(T_1) \circ \text{ad}(T_2))$$

A simple computation shows that for the examples  $\mathfrak{g} = \text{Lie}(\text{Gl}_n)$  and  $\mathfrak{g}^{(0)} = \text{Lie}(\text{Sl}_n)$  we have

$$B(T_1, T_2) = 2n \text{tr}(T_1 T_2) - 2 \text{tr}(T_1) \text{tr}(T_2) \tag{1.25}$$

we observe that in case that one of the  $T_i$  is central, i.e.  $= u\text{Id}$  we have  $B(T_1, T_2) = 0$ . In the case of  $\mathfrak{g}^{(0)}$  the second term is zero.

*It is well known that a linear algebraic group is semi-simple if and only if the Killing form  $B$  on its Lie algebra is non degenerate.*

### 1.1.5 Structure of semisimple groups over $\mathbb{R}$ and the symmetric spaces

We need some information concerning the structure of the group  $G_\infty = G(\mathbb{R})$  for semisimple groups over  $G/\mathbb{R}$ . We will provide this information simply by discussing a series of examples.

Of course the group  $G(\mathbb{R})$  is a topological group, actually it is even a Lie group. This means it has a natural structure of a  $\mathcal{C}_\infty$ -manifold with respect to this structure. Instead of  $G(\mathbb{R})$  we will very often write  $G_\infty$ . Let  $G_\infty^0$  be the connected component of the identity in  $G_\infty$ . It is an open subgroup of finite index. We will discuss the

**Theorem of E. Cartan:** *The group  $G_\infty^0$  always contains a maximal compact subgroup  $K_\infty \subset G_\infty^0$  and all maximal compact subgroups are conjugate under  $G_\infty^0$ . The quotient space  $X = G_\infty^0/K_\infty$  is again a  $\mathcal{C}_\infty$ -manifold. It is diffeomorphic to an  $\mathbb{R}^N$  and carries a Riemannian metric which is invariant under the operation of  $G_\infty^0$  from the left. It has sectional curvature  $\leq 0$  and therefore any two points can be joined by a unique geodesic. The maximal compact subgroup  $K \subset G_\infty^0$  is connected and equal to its own normalizer. Therefore the space  $X$  can be viewed as the space maximal compact subgroups in  $G_\infty^0$ .*

For any maximal compact subgroup  $K_x \subset G_\infty$  exists an unique automorphism  $\Theta_x$  with  $\Theta_x^2 = e$  such that  $K_x = \{g \in G_\infty^0 \mid \Theta(g) = g\}$ , this is the Cartan involution corresponding to  $K_x$ . The Cartan involutions are in one-to-one correspondence with the maximal compact subgroups.

A Cartan involution  $\Theta_x$  induces an involution also called  $\Theta_x$  on the Lie algebra  $\mathfrak{g}_\mathbb{R}$  of  $G_\infty$  and we get a decomposition into  $\pm$  eigenspaces

$$\mathfrak{g} = \mathfrak{k}_x \oplus \mathfrak{p}_x; \quad \mathfrak{k}_x = \{U \in \mathfrak{g} \mid \Theta_x(U) = U\}; \quad \mathfrak{p}_x = \{V \in \mathfrak{g} \mid \Theta_x(V) = -V\}$$

where of course  $\mathfrak{k}_x$  is the Lie algebra of  $K_x$ . The differential of the action of  $G_\infty$  on  $G(\mathbb{R})/K_x$  provides an isomorphism  $D_x : \mathfrak{p}_x \xrightarrow{\sim} T_x^X$  (then tangent space at  $x$ ). For  $V_1, V_2 \in \mathfrak{p}_x$  we have  $[V_1, V_2] \in \mathfrak{k}_x$  the map  $R : \mathfrak{p}_x \times \mathfrak{p}_x \rightarrow \mathfrak{k}_x$  is the curvature tensor. The  $\mathbb{R}$ -vector space  $\mathfrak{g}_c := \mathfrak{k}_x + \sqrt{-1}\mathfrak{p}_x \subset \mathfrak{g} \otimes_\mathbb{R} \mathbb{C}$  is a Lie algebra, for  $U_1 + \sqrt{-1}V_1, U_2 + \sqrt{-1}V_2 \in \mathfrak{g}_c$  we get for the Lie-bracket

$$[U_1 + \sqrt{-1}V_1, U_2 + \sqrt{-1}V_2] = [U_1, U_2] - [V_1, V_2] + \sqrt{-1}([U_1, V_2] + [U_2, V_1]) \in \mathfrak{g}_c$$

To this Lie algebra  $\mathfrak{g}_c$  corresponds an algebraic group  $G_c/\mathbb{R}$  which is a  $\mathbb{R}$ -form of  $G/\mathbb{R}$ , the group  $G_c(\mathbb{R})$  is compact. The group  $G_c/\mathbb{R}$  is called the compact dual of  $G/\mathbb{R}$ . On  $G_c/\mathbb{R}$  we have only one Cartan involution  $\Theta = Id$ .

This theorem is fundamental. To illustrate this theorem we consider a series of examples:

#### The groups $Sl_d(\mathbb{R})$ and $Gl_n(\mathbb{R})$ :

The group  $Sl_d(\mathbb{R})$  is connected. If  $K \subset Sl_d(\mathbb{R})$  is a closed compact subgroup, then we can find a positive definite quadratic form

$$f : \mathbb{R}^n \rightarrow \mathbb{R},$$

such that  $K \subset SO(f, \mathbb{R})$ . since the group  $SO(f, \mathbb{R})$  itself is compact, we have equality. Two such forms  $f_1, f_2$  define the same maximal compact subgroup if

there is a  $\lambda > 0$  in  $\mathbb{R}$  such that  $\lambda f_1 = f_2$ . We say that  $f_1$  and  $f_2$  are conformally equivalent.

This is rather clear, if we believe the first assertion about the existence of  $f$ . The existence of  $f$  is also easy to see if one believes in the theory of integration on  $K$ . This theory provides a positive invariant integral

$$\begin{aligned} \mathcal{C}_c(K) &\longrightarrow \mathbb{R} \\ \varphi &\longrightarrow \int_K \varphi(k) dk \end{aligned}$$

with  $\int \varphi > 0$  if  $\varphi \geq 0$  and not identically zero (positivity),  $\int \varphi(kk_0) dk = \int \varphi(k_0k) dk = \int \varphi(k) dk$  (invariance).

To get our form  $f$  we start from any positive definite form  $f_0$  on  $\mathbb{R}^n$  and put

$$f(\underline{x}) = \int_K f_0(k\underline{x}) dk.$$

A positive definite quadratic form on  $\mathbb{R}^n$  is the same as a symmetric positive definite bilinear form. Hence the space of positive definite forms is the same as the space of positive definite symmetric matrices

$$\tilde{X} = \{A = (a_{ij}) \mid A = {}^t A, A > 0\}.$$

Hence we can say that the space of maximal compact subgroups in  $\mathrm{Sl}_d(\mathbb{R})$  is given by

$$X = \tilde{X}/\mathbb{R}_{>0}^*.$$

It is easy to see that a maximal compact subgroup  $K \subset \mathrm{Sl}_d(\mathbb{R})$  is equal to its own normalizer (why?). If we view  $X$  as the space of positive definite symmetric matrices with determinant equal to one, then the action of  $\mathrm{Sl}_d(\mathbb{R})$  on  $X = \mathrm{Sl}_d(\mathbb{R})/K$  is given by

$$(g, A) \longrightarrow g A {}^t g,$$

and if we view it as the space of maximal compact subgroups, then the action is conjugation.

There is still another interpretation of the points  $x \in X$ . In our above interpretation a point was a symmetric, positive definite bilinear form  $\langle \cdot, \cdot \rangle_x$  on  $\mathbb{R}^n$  up to a homothety. From this we get a transposition  $g \mapsto {}^t_x g$ , which is defined by the rule  $\langle gv, u \rangle_x = \langle v, {}^t_x gu \rangle_x$  and from this we get the involution

$$\Theta_x : g \mapsto ({}^t_x g)^{-1} \tag{1.26}$$

Then the corresponding maximal compact subgroup is

$$K_x = \{g \in \mathrm{Sl}_n(\mathbb{R}) \mid \Theta_x(g) = g\} \tag{1.27}$$

This involution  $\Theta_x$  is a Cartan involution, it also induces an involution also called  $\Theta_x$  on the Lie-algebra and it has the property that (See 6.4)

$$(u, v) \mapsto B(u, \Theta_x(v)) = B_{\Theta_x}(u, v) \tag{1.28}$$

is negative definite. This bilinear form is  $K_x$  invariant. All these Cartan involutions are conjugate.

If we work with  $\mathrm{Gl}_n(\mathbb{R})$  instead then we have some freedom to define the symmetric space. In this case we have the non trivial center  $\mathbb{R}^\times$  and it is sometimes useful to define

$$X = \mathrm{Gl}_n(\mathbb{R})/\mathrm{SO}(\mathbb{R}) \cdot \mathbb{R}_{>0}^\times, \quad (1.29)$$

then our symmetric space has two components, a point is pair  $(\Theta_x, \epsilon)$  where  $\epsilon$  is an orientation. If we do not divide by  $\mathbb{R}_{>0}^\times$  then we multiply the Riemannian manifold  $X$  by a flat space and we get the above space  $\tilde{X}$ .

A Cartan involution on  $\mathrm{Gl}_n(\mathbb{R})$  is an involution which induces a Cartan involution on  $\mathrm{Sl}_n(\mathbb{R})$  and which is trivial on the center.

**Proposition 1.1.1.** *The Cartan involutions on  $\mathrm{Gl}_n(\mathbb{R})$  are in one to one correspondence to the euclidian metrics on  $\mathbb{R}^n$  up to conformal equivalence.*

Finally we recall the Iwasawa decomposition. Inside  $\mathrm{Gl}_n(\mathbb{R})$  we have the standard Borel- subgroup  $B(\mathbb{R})$  of upper triangular matrices and it is well known that

$$\mathrm{Gl}_n(\mathbb{R}) = B(\mathbb{R}) \cdot \mathrm{SO}(\mathbb{R}) \cdot \mathbb{R}_{>0}^\times \quad (1.30)$$

and hence we see that  $B(\mathbb{R})$  acts transitively on  $X$ .

### The compact dual of $\mathrm{Sl}_n(\mathbb{R})$

If  $G/\mathbb{R}$  is a semi simple group, then  $G_c/\mathbb{R}$  is a  $\mathbb{R}$ -form of  $G/\mathbb{R}$ . Therefore we find a cohomology class  $\xi_c \in H^1\mathbb{C}/(\mathbb{R}, \mathrm{Aut}(G))$  corresponding to  $G_c$ . It is clear from the Theorem of Cartan how we get a cocycle representing this class: We choose a Cartan involution  $\Theta \in \mathrm{Aut}(G)$ , the Galois group  $\mathrm{Gal}(\mathbb{C}/\mathbb{R})$  is cyclic of order 2 let  $\mathbf{c}$  be the generator (the complex conjugation). Then  $\mathbf{c} \mapsto \mathbf{c} \circ \Theta$  yields a 1-cocycle in  $C^1(\mathrm{Gal}(\mathbb{C}/\mathbb{R}), \mathrm{Aut}(G)(\mathbb{C}))$ . (Lemma 1.1.1 ) and this 1-cocycle represents the class  $\xi_c$ .

This means for the group  $\mathrm{Sl}_n/\mathbb{R}$  that

$$G_c(\mathbb{R}) = \{g \in \mathrm{Sl}_n(\mathbb{C}) | \mathbf{c}(g^{-1}) = g\}$$

and if we go back to the usual notion and write  $\mathbf{c}(g) = \bar{g}$  then we get

$$G_c(\mathbb{R}) = \{g \in \mathrm{Sl}_n(\mathbb{C}) | g \bar{g} = \mathrm{Id}\} = \mathrm{SU}(n)$$

Here of course  $\mathrm{SU}(n) = \mathrm{SU}(h_c)$  where  $h_c(z_1, z_2, \dots, z_n) = \sum_{i=1}^n z_i \bar{z}_i$  is the standard positive definite hermitian form on  $\mathbb{C}^n$ .

We know that for  $G/\mathbb{R} = \mathrm{Sl}_n/\mathbb{R}$  and  $n > 2$  the Cartan involution  $\Theta$  is the generator of  $\mathrm{Aut}(G)/\mathrm{Ad}(G)$  and hence it is clear that  $\xi_c$  is not in the image of  $H^1\mathbb{C}/(\mathbb{R}, \mathrm{Ad}(G)) \rightarrow H^1\mathbb{C}/(\mathbb{R}, \mathrm{Aut}(G))$ . This means that in this case  $G_c/\mathbb{R} = \mathrm{SU}(n)/\mathbb{R}$  is not an inner  $\mathbb{R}$ -form of  $\mathrm{Sl}_n/\mathbb{R}$ , in turn this also means that  $\mathrm{Sl}_n/\mathbb{R}$  is not an inner  $\mathbb{R}$ -form of  $\mathrm{SU}(n)/\mathbb{R}$ .

In this context the following general proposition is of importance

**Proposition 1.1.2.** *A semi simple group scheme  $G/\mathbb{R}$  is an inner  $\mathbb{R}$  form of its compact dual  $G_c/\mathbb{R}$  if and only if*

- a) *The Cartan involution  $\Theta$  of  $G/\mathbb{R}$  is an inner automorphism of  $G/\mathbb{R}$ .*
- b) *The group  $G/\mathbb{R}$  has a compact maximal torus  $T_c/\mathbb{R} \subset G/\mathbb{R}$ .*

**Give a name to this class of groups ? Examples?**

**The Arakelow- Chevalley scheme**  $(\mathrm{Gl}_n/\mathbb{Z}, \Theta_0)$ 

We start from the free lattice  $L = \mathbb{Z}e_1 \oplus \mathbb{Z}e_2 \oplus \cdots \oplus \mathbb{Z}e_n$  and we think of  $\mathrm{Gl}_n/\mathbb{Z}$  as the scheme of automorphism of this lattice. If we choose an euclidian metric  $\langle \cdot, \cdot \rangle$  on  $L \otimes \mathbb{R}$ , then we call the pair  $(L, \langle \cdot, \cdot \rangle)$  an *Arakelow vector bundle*. From the (conformal class of) metric we get a Cartan involution  $\Theta$  on  $\mathrm{Gl}_n(\mathbb{R})$ , and the pair  $(\mathrm{Gl}_n/\mathbb{Z}, \Theta)$  is an Arakelow group scheme.

We may choose the standard euclidian metric with respect to the given basis, i.e.  $\langle e_i, e_j \rangle = \delta_{i,j}$ . The the resulting Cartan involution is the standard one:  $\Theta_0 : g \mapsto ({}^t g)^{-1}$ . This pair  $(\mathrm{Gl}_n/\mathbb{Z}, \Theta_0)$  is called an Arakelow- Chevalley scheme. (In a certain sense the integral structure of  $\mathrm{Gl}_n/\mathbb{Z}$  and the choice of the Cartan involution are "optimally adapted")

In this case we find for our basis elements in (1.21)

$$B_{\Theta_0}(H_i, H_j) = -2n\delta_{i,j} + 2; B_{\Theta_0}(E_{i,j}, E_{k,l}) = -2n\delta_{i,k}\delta_{j,l} \quad (1.31)$$

hence the  $E_{i,j}$  are part of an orthonormal basis.

We propose to call a pair  $(L, \langle \cdot, \cdot \rangle_x)$  an Arakelow vector bundle over  $\mathrm{Spec}(\mathbb{Z}) \cup \{\infty\}$  and  $(\mathrm{Gl}_n, \Theta_x)$  an Arakelow group scheme. The Arakelow vector bundles modulo conformal equivalence are in one-to one correspondence with the Arakelow group schemes of type  $\mathrm{Gl}_n$ .

**The group  $\mathrm{Sl}_d(\mathbb{C})$** 

We now consider the group  $G/\mathbb{R}$  whose group of real points is  $G(\mathbb{R}) = \mathrm{Sl}_d(\mathbb{C})$  (see 1.1 example 4)).

A completely analogous argument as before shows that the maximal compact subgroups are in one to one correspondence to the positive definite hermitian forms on  $\mathbb{C}^n$  (up to multiplication by a scalar). Hence we can identify the space of maximal compact subgroup  $K \subset G(\mathbb{R})$  to the space of positive definite hermitian matrices

$$X = \{A \mid A = {}^t \bar{A}, A > 0, \det A = 1\}.$$

The action of  $\mathrm{Sl}_d(\mathbb{C})$  by conjugation on the maximal compact subgroups becomes

$$A \longrightarrow g A {}^t \bar{g}$$

on the space of matrices.

**The orthogonal group:**

The next example we want to discuss is the orthogonal group of a non degenerate quadratic form

$$f(x_1, \dots, x_n) = x_1^2 + \dots + x_m^2 - x_{m+1}^2 - \dots - x_n^2,$$

since at this moment we consider only groups over the real numbers, we may assume that our form is of this type. In this case one has the usual notation

$$SO(f, \mathbb{R}) = SO(m, n - m).$$

Of course we can use the same argument as before and see that for any maximal compact subgroup  $K \subset SO(f, \mathbb{R})$  we may find a positive definite form  $\psi$

$$\psi : \mathbb{R}^n \longrightarrow \mathbb{R}$$

such that  $K = SO(f, \mathbb{R}) \cap SO(\psi, \mathbb{R})$ . But now we cannot take all forms  $\psi$ , i.e. only special forms  $\psi$  provide maximal compact subgroup.

We leave it to the reader to verify that any compact subgroup  $K$  fixes an orthogonal decomposition  $\mathbb{R}^n = V_+ \oplus V_-$  where our original form  $f$  is positive definite on  $V_+$  and negative definite on  $V_-$ . Then we can take a  $\psi$  which is equal to  $f$  on  $V_+$  and equal to  $-f$  on  $V_-$ .

Exercise 3 a) Let  $V/\mathbb{R}$  be a finite dimensional vector space and let  $f$  be a symmetric non degenerate form on  $V$ . Let  $K \subset SO(f)$  be a compact subgroup. If  $f$  is not definite then the action of  $K$  on  $V$  is not irreducible.

b) We can find a  $K$  invariant decomposition  $V = V_- \oplus V_+$  such that  $f$  is negative definite on  $V_-$  and positive definite on  $V_+$ .

In this case the structure of the quotient space  $G(\mathbb{R})/K$  is not so easy to understand. We consider the special case of the form

$$x_1^2 + \dots + x_n^2 - x_{n+1}^2 = f(x_1, \dots, x_{n+1}).$$

We consider in  $\mathbb{R}^{n+1}$  the open subset

$$X_- = \{v = (x_1 \dots x_{n+1}) \mid f(v) < 0\}.$$

It is clear that this set has two connected components, one of them is

$$X_-^+ = \{v \in X_- \mid x_{n+1} > 0\}$$

Since it is known that  $SO(n, 1)$  acts transitively on the vectors of a given length, we find that  $SO(n, 1)$  cannot be connected. Let  $G_\infty^0 \subset SO(n, 1)$  be the subgroup leaving  $X_-^+$  invariant.

Now it is not difficult to show that for any maximal compact subgroup  $K \subset G_\infty^0$  we can find a ray  $\mathbb{R}_{>0}^* \cdot v \subset X_-^{(+)}$  which is fixed by  $K$ .

(Start from  $v_0 \in X_-^{(+)}$  and show that  $\mathbb{R}_{>0}^* K v_0$  is a closed convex cone in  $X_-^{(+)}$ . It is  $K$  invariant and has a ray which has a ‘‘centre of gravity’’ and this is fixed under  $K$ .)

For a vector  $v = (x_1, \dots, x_{n+1}) \in X_-^{(+)}$  we may normalize the coordinate  $x_{n+1}$  to be equal to one; then the rays  $\mathbb{R}_{>0}^+ v$  are in one to one correspondence with the points of the ball

$$\overset{\circ}{D}_n = \{(x_1, \dots, x_n) \mid x_1^2 + \dots + x_n^2 < 1\} \subset X_-^{(+)}$$

This tells us that we can identify the set of maximal compact subgroups  $K \subset G_\infty^0$  with the points of this ball. The first conclusion is that  $G_\infty^0/K \simeq D^n$  is topologically a cell (diffeomorphic to  $\mathbb{R}^n$ ). Secondly we see that for a  $v \in X_-^+$  we have an orthogonal decomposition with respect to  $f$

$$\mathbb{R}^{n+1} = \langle v \rangle + \langle v \rangle^\perp,$$

and the corresponding maximal compact subgroup is the orthogonal group on  $\langle v \rangle^\perp$ .

**Give Cartan Involutions?**

### 1.1.6 Special low dimensional cases

1) We consider the ( semi-simple ) group  $\mathrm{Sl}_2(\mathbb{R})$ . It acts on the upper half plane

$$\mathbb{H} = \{z \mid z \in \mathbb{C}, \Im(z) > 0\}$$

by

$$(g, z) \longrightarrow \frac{az + b}{cz + d}, \quad g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{Sl}_2(\mathbb{R}).$$

It is clear that the stabilizer of the point  $i \in H$  is the standard maximal compact subgroup

$$K_\infty = \mathrm{SO}(2) = \left\{ \begin{pmatrix} \cos \varphi & \sin \varphi \\ -\sin \varphi & \cos \varphi \end{pmatrix} \right\}.$$

Hence we have  $\mathbb{H} = \mathrm{Sl}_2(\mathbb{R})/K_\infty$ . But this quotient has been realized as the space of symmetric positive definite  $2 \times 2$ -matrices with determinant equal to one

$$x = \left\{ \begin{pmatrix} y_1 & x_1 \\ x_1 & y_2 \end{pmatrix} \mid y_1 y_2 - x_1^2 = 1, y_1 > 0 \right\}.$$

It is clear how to find an isomorphism between these two explicit realizations. The map

$$\begin{pmatrix} y_1 & x_1 \\ x_1 & y_2 \end{pmatrix} \longrightarrow \frac{i + x_1}{y_2},$$

is compatible with the action of  $\mathrm{Sl}_2(\mathbb{R})$  on both sides and sends the identity

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \text{ to the point } i.$$

If we start from a point  $z \in H$  the corresponding metric is as follows: We identify the lattices  $\langle 1, z \rangle = \{a + bz \mid a, b \in \mathbb{Z}\} = \Omega$  to the lattice  $\mathbb{Z}^2 \subset \mathbb{R}^2$  by sending  $1 \rightarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix}$  and  $z \rightarrow \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ . The standard euclidian metric on  $\mathbb{C} = \mathbb{R}^2$  induces a metric on  $\Omega \subset \mathbb{C}$ , and this metric is transported to  $\mathbb{R}^2$  by the identification  $\Omega \otimes \mathbb{R} \rightarrow \mathbb{R}^2$ .

We may also start from the (reductive) group  $\mathrm{Gl}_2(\mathbb{R})$ , it has the centre  $C(\mathbb{R}) = \left\{ \begin{pmatrix} t & 0 \\ 0 & t \end{pmatrix} \right\}$ . Let  $C(\mathbb{R})^{(0)}$  be the connected component of the identity of  $C(\mathbb{R})$ .

In this case we define  $K_\infty = \mathrm{SO}(2) \times C(\mathbb{R})^{(0)}$ . Then the quotient

$$\mathrm{Gl}_2(\mathbb{R})/K_\infty = \mathbb{H} \cup \mathbb{H}_-$$

where  $\mathbb{H}_-$  is the lower half plane.

2) The two groups  $\mathrm{Sl}_2(\mathbb{R})$  and  $\mathrm{PSl}_2(\mathbb{R})^{(0)} = \mathrm{Sl}_2(\mathbb{R})/\{\pm \mathrm{Id}\}$  give rise to the same symmetric space. The group  $\mathrm{PSl}_2(\mathbb{R})$  acts on the space  $M_2(\mathbb{R})$  of  $2 \times 2$ -matrices by conjugation (the group  $\mathrm{Gl}_2(\mathbb{R})$  acts by conjugation and the centre acts trivially) and leaves invariant the space

$$\{A \in M_2(\mathbb{R}) \mid \mathrm{trace}(A) = 0\} = M_2^0(\mathbb{R}).$$

On this three-dimensional space we have a symmetric quadratic form

$$\begin{aligned} B & : M_2^0(\mathbb{R}) \longrightarrow \mathbb{R} \\ B & : A \mapsto \frac{1}{2} \mathrm{trace}(A^2) \end{aligned}$$

and with respect to the basis

$$H = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, V = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, Y = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad (1.32)$$

this form is  $x_1^2 + x_2^2 - x_3^2$ .

Hence we see that  $SO(M_2^0(\mathbb{R}), B) = SO(2, 1)$ , and hence we have an isomorphism between  $PSL_2(\mathbb{R})$  and the connected component of the identity  $G_\infty^0 \subset SO(2, 1)$ . Hence we see that our symmetric space  $H = \mathrm{Sl}_2(\mathbb{R})/K_\infty = PSL_2(\mathbb{R})/\overline{K}_\infty$  can also be realized (see ..... ) as disc

$$D = \{(x_1, x_2) \mid x_1^2 + x_2^2 < 1\}$$

where we normalized  $x_3 = 1$  on  $X_-^{(+)}$  as in ..... .

### The group $\mathrm{Sl}_2(\mathbb{C})$

Recall that in this case the symmetric space is given by the positive definite hermitian matrices

$$A = \left\{ \begin{pmatrix} y_1 & z \\ \bar{z} & y_2 \end{pmatrix} \mid \det(A) = 1, y_1 > 0 \right\}.$$

In this case we have also a realization of the symmetric space as an upper half space. We send

$$\begin{pmatrix} y_1 & w \\ \bar{w} & y_2 \end{pmatrix} \mapsto \left( \frac{w}{y_2}, \frac{1}{y_2} \right) = (z, \zeta) \in \mathbb{C} \times \mathbb{R}_{>0}$$

The inverse of this isomorphism is given by

$$(z, \zeta) \mapsto \begin{pmatrix} \zeta + z\bar{z}/\zeta & z/\zeta \\ \bar{z}/\zeta & 1/\zeta \end{pmatrix}$$

As explained earlier, the action of  $\mathrm{Gl}_2(\mathbb{C})$  on the maximal compact subgroup given by conjugation yields the action

$$G(\mathbb{R}) \times X \longrightarrow X,$$

$$(g, A) \longrightarrow gA^t\bar{g},$$

on the hermitian matrices. Translating this into the realization as an upper half space yield the slightly scaring formula

$$G \times (\mathbb{C} \times \mathbb{R}_{>0}) \longrightarrow \mathbb{C} \times \mathbb{R}_{>0},$$

$$(g, (z, \zeta)) \longrightarrow \left( \frac{(az + b) \overline{(cz + d)} + a\bar{c} \zeta^2}{(cz + d) \overline{(cz + d)} + c\bar{c} \zeta^2}, \frac{\zeta}{(cz + d) \overline{(cz + d)} + c\bar{c} \zeta^2} \right)$$

**1.3.4. The Riemannian metric:** It was already mentioned in the statement of the theorem of Cartan that we always have a  $G_\infty^0$  invariant Riemannian

metric on  $X$ . It is not too difficult to construct such a metric which in many cases is rather canonical.

In the general case we observe that the maximal compact subgroup is the stabilizer of the point  $x_0 = e \cdot K \in G_\infty^0/K_\infty = X$ . Hence it acts on the tangent space of  $x_0$ , and we can construct a  $K$ -invariant positive definite quadratic form on this tangent space. Then we use the action of  $G_\infty^0$  on  $X$  to transport this metric to an arbitrary point in  $X$ : If  $x \in X$  we find a  $g$  so that  $x = gx_0$ , it defines an isomorphism between the tangent space at  $x_0$  and the tangent space at  $x$ . Hence we get a form on the tangent space at  $x$ , which will not depend on the choice of  $g \in G_\infty^0$ .

In our examples this metric is always unique up to scalars.

a) In the case of the group  $\mathrm{Sl}_d(\mathbb{R})$  we may take as a base point  $x_0 \in X$  the identity  $\mathrm{Id} \in \mathrm{Sl}_d(\mathbb{R})$ . The corresponding maximal compact subgroup is the orthogonal group  $\mathrm{SO}(n)$ . The tangent space at  $\mathrm{Id}$  is given by the space

$$\mathrm{Sym}_n^0(\mathbb{R}) = T_{\mathrm{Id}}^X$$

of symmetric matrices with trace zero. On this space we have the form

$$Z \longrightarrow \mathrm{trace}(Z^2),$$

which is positive definite (a symmetric matrix has real eigenvalues). It is easy to see that the orthogonal group acts on this tangent space by conjugation, hence the form is invariant.

b) A similar argument applies to the group  $G_\infty = \mathrm{Sl}_d(\mathbb{C})$ . Again the identity  $\mathrm{Id}$  is a nice positive definite hermitian matrix. The tangent space consists of the hermitian matrices

$$T_{\mathrm{Id}}^X = \{A \mid A = {}^t \bar{A} \text{ and } \mathrm{tr}(A) = 0\},$$

and the invariant form is given by

$$A \longrightarrow \mathrm{tr}(A\bar{A}).$$

c) In the case of the group  $G_\infty^0 \subset \mathrm{SO}(f)$  where  $f$  is the quadratic form

$$f(x_1, \dots, x_{n+1}) = x_1^2 + \dots + x_n^2 - x_{n+1}^2.$$

We realized the symmetric space as the open ball

$$\overset{\circ}{D}_n = \{(x_1, \dots, x_n) \mid x_1^2 + \dots + x_n^2 < 1\}.$$

The orthogonal group  $\mathrm{SO}(n, 1)$  is the stabilizer of  $0 \in \overset{\circ}{D}_n$ , and hence it is clear that the Riemannian metric has to be of the form

$$h(x_1^2 + \dots + x_n^2)(dx_1^2 + \dots + dx_n^2)$$

(in the usual notation). A closer look shows that the metric has to be

$$\frac{dx_1^2 + \dots + dx_n^2}{\sqrt{1 - x_1^2 - \dots - x_n^2}}.$$

In our two low dimensional spacial examples the metric is easy to determine. For the action of the group  $\mathrm{Sl}_2(\mathbb{R})$  on the upper half plane  $H$  we observe that for any point  $z_0 = x + iy \in H$  the tangent vectors  $\frac{\partial}{\partial x}|_{z_0}$ ,  $\frac{\partial}{\partial y}|_{z_0}$  form a basis of the tangent spaces at  $z_0$ .

If we take  $z_0 = i$  then the stabilizer is the group  $SO(2)$  and for

$$e(\varphi) = \begin{pmatrix} \cos \varphi & \sin \varphi \\ -\sin \varphi & \cos \varphi \end{pmatrix}.$$

We have

$$\begin{aligned} e(\varphi) \cdot \left( \frac{\partial}{\partial x} \Big|_i \right) &= \cos 2\varphi \cdot \frac{\partial}{\partial x} \Big|_i + \sin 2\varphi \frac{\partial}{\partial y} \Big|_i \\ e(\varphi) \cdot \left( \frac{\partial}{\partial y} \Big|_i \right) &= \sin 2\varphi \cdot \frac{\partial}{\partial x} \Big|_i + \cos 2\varphi \frac{\partial}{\partial y} \Big|_i. \end{aligned}$$

Hence we find that  $\frac{\partial}{\partial x} \Big|_i$  and  $\frac{\partial}{\partial y} \Big|_i$  have to be orthogonal and of the same length.

Now the matrix

$$\begin{pmatrix} y & x \\ 0 & 1 \end{pmatrix}$$

sends  $i$  into the point  $z = x + iy$ . It sends  $\frac{\partial}{\partial x} \Big|_i$  and  $\frac{\partial}{\partial y} \Big|_i$  into  $y \cdot \frac{\partial}{\partial x} \Big|_z$  and  $y \cdot \frac{\partial}{\partial y} \Big|_z$ , and hence we must have for our invariant metric

$$\left\langle \frac{\partial}{\partial x} \Big|_z, \frac{\partial}{\partial y} \Big|_z \right\rangle = 0; \quad \left\langle \frac{\partial}{\partial x} \Big|_z, \frac{\partial}{\partial x} \Big|_z \right\rangle = \frac{1}{y^2}; \quad \left\langle \frac{\partial}{\partial y} \Big|_z, \frac{\partial}{\partial y} \Big|_z \right\rangle = \frac{1}{y^2},$$

and this is in the usual notation the metric

$$ds^2 = \frac{1}{y^2}(dx^2 + dy^2).$$

A completely analogous argument yields the metric

$$ds^2 = \frac{1}{\zeta^2}(d\zeta^2 + dx^2 + dy^2).$$

for the space  $\mathbb{H}_3$ .

## 1.2 Arithmetic groups

If we have a linear algebraic group  $G/\mathbb{Q} \hookrightarrow GL_n$  we may consider the group  $\Gamma = G(\mathbb{Q}) \cap GL_n(\mathbb{Z})$ . This is the first example of an *arithmetic* group. It has the following fundamental property:

**Proposition:** *The group  $\Gamma$  is a discrete subgroup of the topological group  $G(\mathbb{R})$ .*

This is rather easily reduced to the fact that  $\mathbb{Z}$  is discrete in  $\mathbb{R}$ . Actually our construction provides a big family of arithmetic groups. For any integer  $m > 0$  we have the homomorphism of reduction mod  $m$ , namely

$$GL_n(\mathbb{Z}) \longrightarrow GL_n(\mathbb{Z}/m\mathbb{Z}).$$

The kernel  $GL_n(\mathbb{Z})(m)$  of this homomorphism has finite index in  $GL_n(\mathbb{Z})$  and hence the intersection  $\Gamma' = GL_n(\mathbb{Z})(m) \cap \Gamma$  has finite index in  $\Gamma$ .

**Definition 2.1.:** A subgroup  $\Gamma''$  of  $\Gamma$  is called a congruence subgroup, if we can find an integer  $m$  such that

$$GL_n(\mathbb{Z})(m) \cap \Gamma \subset \Gamma'' \subset \Gamma.$$

At this point a remark is in order. We explained already that a linear algebraic group  $G/\mathbb{Q}$  may be embedded in different ways into different groups  $GL_n$ , i.e.

$$\begin{array}{c} \hookrightarrow GL_{n_1} \\ G \\ \hookrightarrow GL_{n_2} \end{array}$$

In this case we may get two different congruence subgroups

$$\Gamma_1 = G(\mathbb{Q}) \cap GL_{n_1}(\mathbb{Z}), \Gamma_2 = G(\mathbb{Q}) \cap GL_{n_2}(\mathbb{Z}).$$

It is not hard to show that in such a case we can find an  $m > 0$  such that

$$\begin{array}{l} \Gamma_1 \supset \Gamma_2 \cap GL_{n_2}(\mathbb{Z})(m) \\ \Gamma_2 \supset \Gamma_1 \cap GL_{n_1}(\mathbb{Z})(m) \end{array} .$$

From this we conclude that the notion of congruence subgroup does not depend on the way we realized the group  $G/\mathbb{Q}$  as a subgroup in the general linear group.

Now we may also define the notion of an *arithmetic* subgroup. A subgroup  $\Gamma' \subset G(\mathbb{Q})$  is called arithmetic if for any congruence subgroup  $\Gamma \subset G(\mathbb{Q})$  the group  $\Gamma' \cap \Gamma$  is of finite index in  $\Gamma'$  and  $\Gamma$ . (We say that  $\Gamma'$  and  $\Gamma$  are commensurable.) By definition all congruence subgroups are arithmetic subgroups.

The most prominent example of an arithmetic group is the group

$$\Gamma = \mathrm{Sl}_2(\mathbb{Z}).$$

Another example is obtained as follows. We defined for any number field  $K/\mathbb{Q}$  the group

$$G/\mathbb{Q} = R_{K/\mathbb{Q}}(\mathrm{Sl}_d)$$

for which  $G(\mathbb{Q}) = \mathrm{Sl}_d(K)$ . If  $\mathcal{O}_K$  is the ring of integers in  $K$ , then  $\Gamma = \mathrm{Sl}_d(\mathcal{O}_K)$  (and also  $\tilde{\Gamma} = GL_n(\mathcal{O}_K)$ ) is a congruence (and hence arithmetic) subgroup of  $G(\mathbb{Q})$ .

It is very interesting that the groups  $\Gamma = \mathrm{Sl}_2(\mathbb{Z})$  and  $\mathrm{Sl}_2(\mathcal{O}_K)$  for imaginary quadratic  $K/\mathbb{Q}$  always contain arithmetic subgroups  $\Gamma' \subset \Gamma$  which are not congruence subgroups. This means that in general the class of arithmetic subgroups is larger than the class of congruence subgroups. We will prove this assertion in **Non Congruence subgroups**).

If only the group  $G(\mathbb{R})$  is given (as the group of real points of a group  $G/\mathbb{R}$  or perhaps only as a Lie group, then the notion of arithmetic group  $\Gamma \subset G(\mathbb{R})$

is not defined. The notion of an arithmetic subgroup  $\Gamma \subset G(\mathbb{R})$  requires the choice of a group scheme  $G/\mathbb{Q}$  such that the group  $G(\mathbb{R})$  is the group of real points of this group over  $\mathbb{Q}$ . The exercise in 1.1.2. shows that different  $\mathbb{Q}$ - forms provide different arithmetic groups.

*Exercise 2 If  $\gamma \in \text{Gl}_n(\mathbb{Z})$  is a nontrivial torsion element and if  $\gamma \equiv \text{Id} \pmod{m}$  then  $m = 1$  or  $m = 2$ . In the latter case the element  $\gamma$  is of order 2.*

*This implies that for  $m \geq 3$  the congruence subgroup  $\text{Gl}_n(\mathbb{Z})(m)$  of  $\text{Gl}_n(\mathbb{Z})$  is torsion free.*

This implies of course that any arithmetic group has a subgroup of finite index, which is torsion free.

Affgroup

### 1.2.1 Affine group schemes over $\mathbb{Z}$

There is a slightly more sophisticated view of arithmetic groups. In our book [29] section 7.5.6 and on p. 50,51 we discuss briefly the general notion of a group scheme over an arbitrary base scheme  $S$ . An affine group scheme over  $G/\mathbb{Z}$  is a finitely generated  $\mathbb{Z}$ -algebra  $A(G)$  together with a comultiplication  $m : A(G) \rightarrow A(G) \otimes A(G)$ . For any  $\mathbb{Z}$ -algebra  $B$  (commutative and with identity) the comultiplication  $m$  induces a multiplication on the  $B$ -valued points

$${}^t m : \text{Hom}_{\text{alg}}(A, B) \times \text{Hom}_{\text{alg}}(A, B) \rightarrow \text{Hom}_{\text{alg}}(A, B)$$

and the requirement is that this multiplication defines a group structure on  $G(B) = \text{Hom}_{\text{alg}}(A, B)$ . In educated language :  $G/\mathbb{Z}$  is a functor from the category of affine schemes into the category of groups.

For instance we can define the group scheme  $\text{Gl}_n/\mathbb{Z}$ . The affine algebra is

$$A(\text{Gl}_n) = \mathbb{Z}[X_{11}, X_{12}, \dots, X_{1n}, X_{21}, \dots, X_{nn}, Y]/(Y \det(X_{ij}) - 1)$$

Then the group  $\text{Gl}_n(\mathbb{Z})$  of  $\mathbb{Z}$ -valued points of  $\text{Gl}_n/\mathbb{Z}$  is our group  $\text{Gl}_n(\mathbb{Z})$ .

If  $G/\mathbb{Q} \subset \text{Gl}_n/\mathbb{Q}$  is a subgroup, then the affine algebra  $A(G) = A(\text{Gl}_n) \otimes \mathbb{Q}/I$ , where  $I$  is an ideal in  $A(\text{Gl}_n) \otimes \mathbb{Q}$ . Since  $G/\mathbb{Q}$  is a subgroup this ideal must satisfy

$$m_{\text{Gl}_n}(I) \subset A(\text{Gl}_n) \otimes \mathbb{Q} \otimes I + I \otimes A(\text{Gl}_n) \otimes \mathbb{Q}.$$

Let  $J = A(\text{Gl}_n) \cap I$ , then it is easy to check that the comultiplication of  $A(\text{Gl}_n)$  satisfies

$$m_{\text{Gl}_n}(J) \subset A(\text{Gl}_n) \otimes J + J \otimes A(\text{Gl}_n)$$

and this tells us that  $m_{\text{Gl}_n}$  induces a comultiplication

$$m : A(\text{Gl}_n)/J \rightarrow A(\text{Gl}_n)/J \otimes A(\text{Gl}_n)/J$$

which provides a group scheme structure. This means that we have extended the group scheme  $G/\mathbb{Q}$  to a group scheme over  $\mathcal{G}/\mathbb{Z}$ . The affine algebra  $A(\mathcal{G}) = A(\text{Gl}_n)/J$ . This extension depends on the choice of the embedding into  $\text{Gl}_n/\mathbb{Q}$  and it is called the *flat extension*. Then the base extension  $\mathcal{G} \times_{\mathbb{Z}} \mathbb{Q} = G/\mathbb{Q}$ , this base extension is called the *generic fiber* of  $\mathcal{G}/\mathbb{Z}$ .

We now may understand our arithmetic group  $\Gamma = G(\mathbb{Q}) \cap \mathrm{Gl}_n(\mathbb{Z})$  as the group  $\mathcal{G}(\mathbb{Z})$  of  $\mathbb{Z}$  valued points of a group scheme over  $\mathbb{Z}$ . Since we know what  $\mathcal{G}(\mathbb{Z}/m\mathbb{Z})$  is we can define congruence subgroups  $\Gamma_H$  as inverse images of subgroups  $H \subset \mathcal{G}(\mathbb{Z}/m\mathbb{Z})$  under the projection  $\mathcal{G}(\mathbb{Z}) \rightarrow \mathcal{G}(\mathbb{Z}/m\mathbb{Z})$ .

There is the special class of semi-simple or reductive group schemes. Roughly speaking an affine group scheme  $G/\mathbb{Z}$  is semi-simple (resp. reductive), if its generic fiber  $G \times_{\mathbb{Z}} \mathbb{Q}$  is semi-simple (resp. reductive) and if for all primes  $p$  the group scheme  $G \times_{\mathbb{Z}} \mathbb{F}_p$  (the reduction mod  $p$ ) is a semi-simple ((resp. reductive)) group scheme over  $\mathbb{F}_p$ .

Of course the simplest example of a semi-simple (resp. reductive) group (scheme) over  $\mathbb{Z}$  is the group  $\mathrm{Sl}_n/\mathbb{Z}$  (resp.  $\mathbb{G}_n/\mathbb{Z}$ ).

We can also construct semi-simple group-schemes by taking flat extensions of orthogonal (resp. symplectic) groups over  $\mathbb{Q}$ , (see section 1.2.1, example 2) and 3). Here the symmetric (resp. alternating) form has to satisfy certain arithmetic conditions (See chap4.pdf).

lattices

### 1.2.2 $\Gamma$ -modules

We consider modules  $\mathcal{M}$  (i.e. abelian groups) with an action of  $\Gamma$ , (see [28], Chap. 2). We want to discuss briefly some special classes of such  $\Gamma$ -modules.

The most important classes of  $\Gamma$ -modules are the modules of *arithmetic origin*. To construct such modules we realise our arithmetic group as  $\Gamma = G(\mathbb{Q}) \cap \mathrm{Gl}_n(\mathbb{Z})$ . Then we take any rational representation  $\rho : G/\mathbb{Q} \rightarrow \mathrm{Gl}(V)$ , where  $V$  is a finite dimensional  $\mathbb{Q}$ -vector space. Now we look for finitely generated submodules  $\mathcal{M} \subset V$  such that  $\mathcal{M} \otimes \mathbb{Q} = V$  which are invariant under the action of  $\Gamma$ . Such a module is a  $\Gamma$ -module of arithmetic origin.

It is not difficult to show that given any finitely generated module  $\mathcal{M}'$  which is a full sublattice, i.e.  $\mathcal{M}' \otimes \mathbb{Q} = V$ , we can find a congruence subgroup  $\Gamma_1 \subset \Gamma$  such that  $\Gamma_1 \mathcal{M}' = \mathcal{M}'$ . Then

$$\mathcal{M} = \bigcap_{\gamma \in \Gamma/\Gamma_1} \gamma \mathcal{M}'$$

is a  $\Gamma$  module (of arithmetic origin).

A second class of  $\Gamma$  modules are those of *congruence origin*. To get such a module we simply pick a congruence subgroup  $\Gamma(N) \subset \Gamma$  and then we simply look at finitely generated abelian groups  $V$  with an action of  $\Gamma/\Gamma(N)$  on  $V$ .

We get some important examples of  $\Gamma$  modules of congruence origin if we start from a  $\Gamma$ -module  $\mathcal{M}$  of arithmetic origin. Then we choose an integer  $N$  and consider the  $\Gamma$  module  $\mathcal{M} \otimes \mathbb{Z}/N\mathbb{Z}$ . On this module  $\Gamma(N)$  acts trivially, hence this module is a  $\Gamma/\Gamma(N)$  module of congruence origin.

We go back to the more sophisticated point of view above, our arithmetic group is the group  $\Gamma = \mathcal{G}(\mathbb{Z})$  of  $\mathbb{Z}$  valued points of the flat extension  $\mathcal{G}/\mathbb{Z}$ .

Now we pick a torsion free finitely generated module  $\mathcal{M}$ , we know what it means that  $\mathcal{M}$  is a  $\mathcal{G}/\mathbb{Z}$  module: It simply means that for any commutative ring  $B$  with identity we have a  $B$ -linear action of  $\mathcal{G}(B)$  on the  $B$ -module  $\mathcal{M} \otimes B$ , or

in other words we have a homomorphism  $\mathcal{G}(B) \rightarrow \mathrm{Gl}_B(\mathcal{M} \otimes_{\mathbb{Z}} B)$ . Of course we require that this action is functorial in  $B$ .

For this book -especially for the first half- the group scheme  $\mathrm{Gl}_2/\mathbb{Z}$  plays a dominant role. In this case the irreducible representations of  $\mathrm{Gl}_n \times_{\mathbb{Z}} \mathbb{Q}$  are well known. We consider the  $\mathbb{Q}$  vector space of homogenous polynomials in two variables and of degree  $n$

$$\mathcal{M}_{n,\mathbb{Q}} := \{P(X, Y) = \sum_{\nu=0}^n a_{\nu} X^{\nu} Y^{n-\nu} \mid a_{\nu} \in \mathbb{Q}\}.$$

We choose an integer  $m$  define an action of  $\mathrm{Gl}_2(\mathbb{Q})$ :

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} P(X, Y) = P(aX + cY, bX + dY) \det \begin{pmatrix} a & b \\ c & d \end{pmatrix}^m, \quad (1.33)$$

this gives us the  $\mathrm{Gl}_2/\mathbb{Q}$ -module  $\mathcal{M}_{n,\mathbb{Q}}[m]$ .

But now it is easy to get  $\mathrm{Gl}_2/\mathbb{Z}$ -modules, we simply define

$$\mathcal{M}_n := \{P(X, Y) = \sum_{\nu=0}^n a_{\nu} X^{\nu} Y^{n-\nu} \mid a_{\nu} \in \mathbb{Z}\} \quad (1.34)$$

and then we define the  $\mathrm{Gl}_2/\mathbb{Z}$  modules  $\mathcal{M}_n[m]$  by the same formula as above.

At this point a small remark is in order. If look at  $\mathcal{M}_n[m]$  only as  $\mathrm{Gl}_2(\mathbb{Z})$ -module then the module "knows" what  $n$  is, clearly  $n = \mathrm{rank}(\mathcal{M}_n) - 1$ . But this  $\mathrm{Gl}_2(\mathbb{Z})$ - module does not "know" what  $m$  is. The only information we get is

$$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} P = (-1)^m P$$

and from this we only get the value of  $m \pmod{2}$ .

### 1.2.3 The locally symmetric spaces

We start from a semisimple group  $G/\mathbb{Q}$ . To this group we attached the the group of real points  $G(\mathbb{R}) = G_{\infty}$ . In  $G_{\infty}$  we have the connected component  $G_{\infty}^0$  of the identity and in this group we choose a maximal compact subgroup  $K$ . The quotient space  $X = G_{\infty}/K$  is a symmetric space which now may have several connected components. On this space we have the action of an arithmetic group  $\Gamma$ .

We have a fundamental fact:

*The action of  $\Gamma$  on  $X$  is properly discontinuous, i.e. for any point  $x \in X$  there exists an open neighborhood  $U_x$  such that for all  $\gamma \in \Gamma$  we have*

$$\gamma U_x \cap U_x = \emptyset \quad \text{or} \quad \gamma x = x.$$

Moreover for all  $x \in X$  the stabilizer

$$\Gamma_x = \{\gamma \mid \gamma x = x\}$$

is finite.

This is easy to see: If we consider the projection  $p : G(\mathbb{R}) \rightarrow G(\mathbb{R})/K = X$ , then the inverse image  $p^{-1}(U_x)$  of a relatively compact neighborhood  $U_x$  of  $x = g_0K$  is of the form  $V_{g_0} \cdot K$ , where  $V_{g_0}$  is a relatively compact neighborhood of  $g_0$ . Hence we look for the solutions of the equation

$$\gamma vk = v'k', \gamma \in \Gamma, v, v' \in V_{g_0}, k, k' \in K.$$

Since  $\Gamma$  is discrete in  $G(\mathbb{R})$  there are only finitely many possibilities for  $\gamma$  and they can be ruled out by shrinking  $U_x$  with the exception of those  $\gamma$  for which  $\gamma x = x$ . If  $\gamma x = x$  this means that  $\gamma g_0K = g_0K$  and hence  $\gamma \in \Gamma \cap g_0K g_0^{-1}$  this intersection is a compact discrete set, hence finite.

If  $\Gamma$  has no torsion then the projection

$$\pi : X \longrightarrow \Gamma \backslash X$$

is locally a  $\mathcal{C}_\infty$ -diffeomorphism. To any point  $x \in \Gamma \backslash X$  and any point  $\tilde{x} \in \pi^{-1}(x)$  we find a neighborhood  $U_{\tilde{x}}$  such that

$$\pi : U_{\tilde{x}} \xrightarrow{\sim} U_x.$$

Hence the space  $\Gamma \backslash X$  inherits the Riemannian metric and the quotient space is a locally symmetric space.

If our group  $\Gamma$  has torsion, then a point  $\tilde{x} \in X$  may have a nontrivial stabilizer  $\Gamma_{\tilde{x}}$ . Then it is not difficult to prove that  $\tilde{x}$  has a neighborhood  $U_{\tilde{x}}$  which is invariant under  $\Gamma_{\tilde{x}}$  and that for all  $\tilde{y} \in U_{\tilde{x}}$  the stabilizer  $\Gamma_{\tilde{y}} \subset \Gamma_{\tilde{x}}$ . This gives us a diagram

$$\begin{array}{ccc} U_{\tilde{x}} & \longrightarrow & \Gamma_{\tilde{x}} \backslash U_{\tilde{x}} = U_x \\ \downarrow & & \downarrow \\ X & \xrightarrow{\pi} & \Gamma \backslash X \end{array}$$

i.e. the point  $x \in \Gamma \backslash X$  has a neighborhood which is the quotient of a neighborhood  $U_{\tilde{x}}$  by a finite group.

In this case the quotient space  $\Gamma \backslash X$  may have singularities. Such spaces are called orbifolds. They have a natural stratification. Any point  $x$  defines a  $\Gamma$  conjugacy class  $[\Gamma_{\tilde{x}}]$  of finite subgroups  $\Gamma_{\tilde{x}} \subset \Gamma$ . On the other hand a conjugacy class  $[c]$  of finite subgroups  $H \subset \Gamma$  defines the (non empty) subset (stratum)  $\Gamma \backslash X([c])$  of those points  $x \in \Gamma \backslash X$  for which  $\Gamma_{\tilde{x}} \in [c]$ .

These strata are easy to describe. We observe that for any finite  $H \subset \Gamma$  the fixed point set  $X^H$  intersected with a connected component of  $X$  is contractible. Let  $x_0 \in X^H$  be a point with  $\Gamma_{x_0} = H$ . Then any other point  $x \in X^H$  is of the form  $x = gx_0$  with  $g \in G(\mathbb{R})$ . This implies that  $g \in N(H)(\mathbb{R})$ , where  $N(H)$  is the normaliser of  $H$ , it is an algebraic subgroup. Then  $N(H)(\mathbb{R}) \cap K = K^H$  is compact subgroup, put  $\Gamma^H = \Gamma \cap N(H)(\mathbb{R})$ , and we get an embedding

$$\Gamma^H \backslash X^H \hookrightarrow \Gamma \backslash X.$$

This space contains the open subset  $(\Gamma^H \backslash X^H)^{(0)}$  of those  $x$  where  $H \in [\Gamma_x]$  and this is in fact the stratum attached to the conjugacy class of  $H$ .

We have an ordering on the set of conjugacy classes, we have  $[c_1] \leq [c_2]$  if for any  $H_1 \in [c_1]$  there exists a subgroup  $H_2 \in [c_2]$  such that  $H_1 \subset H_2$ . These strata are not closed, the closure  $\overline{\Gamma \backslash X([c])}$  is the union of lower dimensional strata.

If we start investigating the stratification above we immediately hit upon number theoretic problems. Let us pick a prime  $p$  and we consider the group  $\Gamma = \mathrm{Sl}_{p-1}[\mathbb{Z}]$  and the ring of  $p$ -th roots of unity  $\mathbb{Z}[\zeta_p]$  as a  $\mathbb{Z}$ -module is free of rank  $p-1$  and hence we get an element

$$\zeta_p \in \mathrm{Sl}(\mathbb{Z}[\zeta_p]) = \mathrm{Sl}_{p-1}(\mathbb{Z})$$

and hence a cyclic subgroup of order  $p$ . But clearly we have many conjugacy classes of elements of order  $p$  in  $\Gamma$  because any ideal  $\mathfrak{a}$  is a free  $\mathbb{Z}$ -module. If we want to understand the conjugacy classes of elements of order  $p$  or the conjugacy classes of cyclic subgroups of order  $p$  in  $\mathrm{Sl}_{p-1}(\mathbb{Z})$  we need to understand the ideal class group. In the next section we will discuss some simple examples.

These quotient spaces  $\Gamma \backslash X$  attract the attention of various different kinds of mathematicians. They provide interesting examples of Riemannian manifolds and they are intensively studied from that point of view. On the other hand number theoretic data enter into their construction. Hence any insight into the structure of these spaces contains number theoretic information.

It is not difficult to see that any arithmetic group  $\Gamma$  contains a normal congruence subgroup  $\Gamma'$  which does not have torsion. This can be deduced easily from the exercise ... at the end of this section. Hence we see that  $\Gamma' \backslash X$  is a Riemannian manifold which is a finite cover of  $\Gamma \backslash X$  with covering group  $\Gamma/\Gamma'$ . The following general theorem is due to Borel and Harish-Chandra:

*The quotient  $\Gamma \backslash X$  always has finite volume with respect to the Riemannian metric. The quotient space  $\Gamma \backslash X$  is compact if and only if the group  $G/\mathbb{Q}$  is anisotropic.*

We will give some further explanation below.

#### 1.2.4 Low dimensional examples

We consider the action of the group  $\Gamma = \mathrm{Sl}_2(\mathbb{Z}) \subset \mathrm{Sl}_2(\mathbb{R})$  on the upper half plane

$$X = \mathbb{H} = \{z \mid \Im(z) = y > 0\} = \mathrm{Sl}_2(\mathbb{R})/SO(2).$$

As we explained in ... we may consider the point  $z = x+iy$  as a positive definite euclidian metric on  $\mathbb{R}^2$  up to a positive scalar. We saw already that this metric can be interpreted as the metric on  $\mathbb{C}$  induced on the lattice  $\Omega = \langle 1, z \rangle$ . The action of  $\mathrm{Sl}_2(\mathbb{Z})$  on the upper half plane corresponds to changing the basis  $1, z$  of  $\Omega$  into another basis and then normalizing the first vector of the new basis to length equal one.

This means that under the action of  $\mathrm{Sl}_2(\mathbb{Z})$  we may achieve that the first vector  $1$  in the lattice is of shortest length. In other words  $\Omega = \langle 1, z \rangle$  where now  $|z| \geq 1$ .


Since we can change the basis by  $1 \rightarrow 1$  and  $z \rightarrow z + n$ . We still have  $|z + n| \geq 1$ . Hence see that this condition implies that we can move  $z$  by these translation into the strip  $-1/2 \leq \Re(z) \leq 1/2$  and since 1 is still the shortest vector we end up in the classical fundamental domain:

$$\mathcal{F} = \{z \mid -1/2 \leq \Re(z) \leq 1/2, |z| \geq 1\} \quad (1.35)$$

Two points  $z_1, z_2 \in \mathcal{F}$  are inequivalent under the action of  $\mathrm{Sl}_2(\mathbb{Z})$  unless they differ by a translation. i.e.

$$z_1 = -\frac{1}{2} + it, \quad z_2 = z_1 + 1 = \frac{1}{2} + it,$$

or we have  $|z_1| = 1$  and  $z_2 = -\frac{1}{z_1}$ . Hence the quotient  $\mathrm{Sl}_2(\mathbb{Z}) \backslash \mathbb{H}$  is given by the following picture



It turns out that this quotient is actually a Riemann surface, i.e. the finite stabilizers at  $i$  and  $\rho$  do not produce singularities. As a Riemann surface the quotient is the complex plane or better the projective line  $\mathbb{P}^1(\mathbb{C})$  minus the point at infinity.

It is clear that the points  $i$  and  $\rho = +\frac{1}{2} + \frac{1}{2}\sqrt{-3}$  in the upper half plane are -up to conjugation by an element  $\gamma \in \text{Sl}_2(\mathbb{Z})$ - the only points with non-trivial stabiliser . Actually the stabilisers are given by

$$\Gamma_i = \left\{ \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right\} , \quad \Gamma_\rho = \left\{ \begin{pmatrix} -1 & 1 \\ -1 & 0 \end{pmatrix} \right\} .$$

We denote the matrices

$$S = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}; R = \begin{pmatrix} -1 & 1 \\ -1 & 0 \end{pmatrix}.$$

The second example is given by the group  $\Gamma = \mathrm{Sl}_2(\mathbb{Z}[i]) \subset \mathrm{Sl}_2(\mathbb{C}) = G_\infty = R_{\mathbb{C}/\mathbb{R}}(\mathrm{Gl}_2/\mathbb{C})(\mathbb{R})$  (See(1.1)). Here we should remember that the choice of  $G_\infty$  allows a whole series of arithmetic groups. For any imaginary quadratic extension  $K = \mathbb{Q}(\sqrt{-d})$  with  $\mathcal{O}_K$  as its ring of integers we may embed  $K$  into  $\mathbb{C}$  and get

$$\mathrm{Sl}_2(\mathcal{O}_K) = \Gamma \subset G_\infty.$$

If the number  $d$  becomes larger then the structure of the group  $\Gamma$  becomes more and more complicated. We discuss only the simplest case.

We will construct a fundamental domain for the action of  $\Gamma$  on the three-dimensional hyperbolic space  $\mathbb{H}_3 = \mathbb{C} \times \mathbb{R}_{>0}$ .

We identify  $\mathbb{H}_3$  with the space of positive definite hermitian matrices

$$X = \{A \in M_2(\mathbb{C}) \mid A = {}^t \bar{A}, A > 0, \det(A) = 1\}.$$

We consider the lattice

$$\Omega = \mathbb{Z}[i] \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \mathbb{Z}[i] \cdot \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

in  $\mathbb{C}^2$  and view  $A$  as a hermitian metric on  $\mathbb{C}^2$  where  $\mathbb{C}/\Omega$  has volume 1. Let  $e'_1 = \begin{pmatrix} \alpha \\ \beta \end{pmatrix}$  be a vector of shortest length. We can find a second vector  $e'_2 = \begin{pmatrix} \gamma \\ \delta \end{pmatrix}$  so that  $\det \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = 1$ . This argument is only valid because  $\mathbb{Z}[i]$  is a principal ideal domain. We consider the vectors  $e'_2 + \nu e'_1$  where  $\nu \in \mathbb{Z}[i]$ . We have

$$\langle e'_2 + \nu e'_1, e'_2 + \nu e'_1 \rangle_A = \langle e'_2 + \nu e'_1 : 1' \rangle_A + \nu \langle e'_1, e'_2 \rangle_A + \bar{\nu} \langle e'_2, e'_1 \rangle_A + \nu \bar{\nu} \langle e'_1, e'_1 \rangle_A.$$

Since we have the euclidean algorithm in  $\mathbb{Z}[i]$  we can choose  $\nu$  such that

$$-\frac{1}{2} \langle e'_1, e'_1 \rangle \leq \mathrm{Re} \langle e'_1, e'_2 \rangle_A, \Im \langle e'_1, e'_2 \rangle_A \leq \frac{1}{2} \langle e'_1, e'_1 \rangle_A.$$

If we translate this to the action of  $\mathrm{Sl}_2(\mathbb{Z}[i])$  on  $\mathbb{H}_3$  then we find that every point  $x = (z; \zeta) \in \mathbb{H}_3$  is equivalent to a point in the domain

$$\tilde{F} = \{(z, \zeta) \mid -\frac{1}{2} \leq \mathrm{Re}(z), \Im(z) \leq \frac{1}{2}; z\bar{z} + \zeta^2 \geq 1\}.$$

Since we have still the action of the matrix  $\begin{pmatrix} i & 0 \\ 0 & -1 \end{pmatrix}$  we even find a smaller fundamental domain

$$F = \{(z, \zeta) \mid -\frac{1}{2} \leq \mathrm{Re}(z), \Im(z) \leq \frac{1}{2}; z\bar{z} + \zeta^2 \geq 1 \text{ and } \mathrm{Re}(z) + \Im(z) \geq 0\}.$$

I want to discuss also the extension of our considerations to the case of the reductive group  $\mathrm{Gl}_2(\mathbb{C})$ . In such a case we have to enlarge the maximal compact

subgroup. In this case the group  $\tilde{K} = \mathrm{Sl}_1(2) \cdot \mathbb{C}^* = K \cdot \mathbb{C}^*$  is a good choice where  $\mathbb{C}^*$  is the centre of  $\mathrm{Gl}_2(\mathbb{C})$ . Then we get

$$\mathbb{H}_3 = \mathrm{Sl}_2(\mathbb{C})/K = \mathrm{Gl}_2(\mathbb{C})/\tilde{K}$$

i.e. we have still the same symmetric space. But the group  $\tilde{\Gamma} = \mathrm{Gl}_2(\mathbb{Z}[i])$  is still larger. We have an exact sequence

$$1 \rightarrow \Gamma \rightarrow \tilde{\Gamma} \rightarrow \{i^\nu\} \rightarrow 1.$$

The centre  $Z_{\tilde{\Gamma}}$  of  $\tilde{\Gamma}$  is given by the matrices  $\left\{ \begin{pmatrix} i^\nu & 0 \\ 0 & i^\nu \end{pmatrix} \right\}$ . The centre  $Z_\Gamma$  has index 2 in  $Z_{\tilde{\Gamma}}$ . Since the centre acts trivially on the symmetric space, hence the above fundamental domain will be “cut into two halves” by the action of  $\tilde{\Gamma}$ . The matrices  $\begin{pmatrix} i^\nu & 0 \\ 0 & 1 \end{pmatrix}$  induce rotation of  $\nu \cdot 90^\circ$  around the axis  $z = 0$  and therefore it becomes clear that the region

$$F_0 = \{(z, \zeta) \mid 0 \leq \Im(z), \mathrm{Re}(z) \leq \frac{1}{2}, z\bar{z} + \zeta^2 \geq 1\}$$

is a fundamental domain for  $\tilde{\Gamma}$ .

The translations  $z \rightarrow z + 1$  and  $z \rightarrow z + i$  identify the opposite faces of  $F$ . This induces an identification on  $F_0$ , namely

$$\left(\frac{1}{2} + iy, \zeta\right) \longrightarrow \left(-\frac{1}{2} + iy, \zeta\right) \longrightarrow \left(y + \frac{i}{2}, \zeta\right).$$

On the bottom of the domain  $F_0$ , namely

$$F_0(1) = \{(z, \zeta) \in F_0 \mid z\bar{z} + \zeta^2 = 1\}$$

we have the further identification

$$(z, \zeta) \longrightarrow (i\bar{z}, \zeta).$$

Hence we see that the quotient space  $\tilde{\Gamma} \backslash \mathbb{H}_3$  is given by the following figure.

I want to discuss the fixed points and the stabilizers of the fixed points of  $\tilde{\Gamma}$ . Before I can do that, I need some simple facts concerning the structure of  $\mathrm{Gl}_2$ .

The group  $\mathrm{Gl}_2(K)$  acts upon the projective line  $\mathbb{P}^1(K) = (K^2 \setminus \{0\})/K^*$ . We write

$$\mathbb{P}^1(K) = (K) \cup \{\infty\}; \quad K(xe_1 + e_2) = x, Ke_1 = \infty.$$

It is quite clear that the action of  $g = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \mathrm{Gl}_2(K)$  is given by

$$gx = \frac{\alpha x + \beta}{\gamma x + \delta}.$$

The action of  $\mathrm{Gl}_2(K)$  on  $\mathbb{P}^1(K)$  is transitive. For a point  $x \in \mathbb{P}^1(K)$  the stabilizer  $B_x$  is clearly a linear subgroup of  $\mathrm{Gl}_2/K$ . If  $x = \infty$ , then this stabilizer is the subgroup

$$B_\infty = \left\{ \begin{pmatrix} a & u \\ 0 & b \end{pmatrix} \right\},$$

and for  $x = 0$  we get

$$B_0 = \left\{ \begin{pmatrix} a & 0 \\ u & b \end{pmatrix} \right\}.$$

It is clear that these subgroups  $B_x$  are conjugate under the action of  $\mathrm{Gl}_2(K)$ . They are in fact maximal solvable subgroups of  $\mathrm{Gl}_2$ .

If we have two different points  $x_1, x_2 \in \mathbb{P}^1(K)$ , then this corresponds to a choice of a basis where the basis vectors are only determined up to scalars. Then the intersection of the two groups  $B_{x_1} \cap B_{x_2}$  is a so-called maximal torus. If we choose  $x_1 = Ke_1, x_2 = Ke_2$ , then

$$B_{x_1} \cap B_{x_2} = \left\{ \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} \right\}.$$

Any other maximal torus of the form  $B_{x_1}, B_{x_2}$  is conjugate to  $T_0$  under  $\mathrm{Gl}_2(K)$ .

Now we assume  $K = \mathbb{C}$ . We compactify the three dimensional hyperbolic space by adding  $\mathbb{P}^1(\mathbb{C})$  at infinity, i.e.

$$\mathbb{H}_3 \hookrightarrow \overline{\mathbb{H}}_3 = \mathbb{H}_3 \cup \mathbb{P}^1(\mathbb{C}) = \mathbb{C} \times \mathbb{R}_{\geq 0} \cup \{\infty\}.$$

(The reader should verify that there is a natural topology on  $\overline{\mathbb{H}}_3$  for which the space is compact and for which  $\mathrm{Gl}_2(\mathbb{C})$  acts continuously.)

Now let us assume that  $a \in \mathrm{Gl}_2(\mathbb{C})$  is an element which has a fixed point on  $\mathbb{H}_3$  and which is not central. Since it lies in a maximal compact subgroup times  $\mathbb{C}^x$  we see that this element  $a$  can be diagonalized

$$a \longrightarrow g_0 a g_0^{-1} = \begin{pmatrix} \alpha & 0 \\ 0 & \beta \end{pmatrix} = a'$$

with  $\alpha \neq \beta$  and  $|\alpha/\beta| = 1$ .

Then it is clear that the fixed point set for  $a'$  is the line

$$\mathrm{Fix}(a') = \{(0, \zeta) \mid \zeta \in \mathbb{R}_{>0}\},$$

i.e. we do not get an isolated fixed point but a full fixed line.

The element  $a'$  has the two fixed points  $\infty, 0$  in  $\mathbb{P}^1(\mathbb{C})$ , and hence it defines the torus  $T_0(\mathbb{C})$ . Then it is clear that

$$\mathrm{Fix}(a') = \{(0, \zeta) \mid \zeta > 0\} = T_0(\mathbb{C}) \cdot (0, 1)$$

i.e. the fixed point set is an orbit under the action of  $T_0(\mathbb{C})$ .

### 1.2.5 Fixed point sets and stabilizers for $\mathrm{Gl}_2(\mathbb{Z}[i]) = \tilde{\Gamma}$

If we want to describe the stabilizers up to conjugation, we can focus our attention on  $F_0$ .

If we have an element  $\gamma \in \tilde{\Gamma}$ ,  $\gamma$  not central and if we assume that  $\gamma$  has fixed points on  $\mathbb{H}_3$ , then we know that  $\gamma$  defines a torus  $T_\gamma = \mathrm{centralizer}_{\mathrm{Gl}_2}(\gamma) = \mathrm{stabilizer}$  of  $x_\gamma, x_{\gamma'} \in \mathbb{P}^1(\mathbb{C})$ . This torus is defined over  $\mathbb{Q}(i)$ , but it is not necessarily diagonalizable over  $\mathbb{Q}(i)$ , it may be that the coordinates of  $x_\gamma, x_{\gamma'}$  lie in a quadratic extension of  $F/\mathbb{Q}(i)$ . This is the quadratic extension defined by the eigenvalues of  $\gamma$ .

We look at the edges of the fundamental domain  $F_0$ . We saw that they consist of connected pieces of the straight lines

$$G_1 = \{(z, \zeta) \mid z = 0\}, G_2 = \{(z, \zeta) \mid z = \frac{1}{2}\}, G_3 = \{(z, \zeta) \mid z = \frac{1+i}{2}\},$$

and the circles (these circles are euclidean circles and geodesics for the hyperbolic metric)

$$D_1 = \{(z, \zeta) \mid z\bar{z} + \zeta^2 = 1, \Im(z) = \operatorname{Re}(z)\}, D_2 = \{(z, \zeta) \mid z\bar{z} + \zeta^2 = 1, \Im(z) = 0\},$$

$$D_3 = \{(z, \zeta) \mid z\bar{z} + \zeta^2 = 1, \operatorname{Re}(z) = \frac{1}{2}\}.$$

The pair of points  $(\infty, (z_0, 0)) \in \mathbb{P}^1(\mathbb{C}) \times \mathbb{P}^1(\mathbb{C})$  has as its stabilizer

$$T_{z_0}(\mathbb{C}) = \begin{pmatrix} 1 & z_0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \alpha & 0 \\ 0 & \beta \end{pmatrix} \begin{pmatrix} 1 & -z_0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} \alpha & z_0(\beta - \alpha) \\ 0 & \beta \end{pmatrix},$$

the straight line  $\{(z_0, \zeta) \mid \zeta > 0\}$  is an orbit under  $T_{z_0}(\mathbb{C})$  and it consists of fixed points for

$$T_{z_0}(\mathbb{C})(1) = \left\{ \begin{pmatrix} \alpha & z_0(\beta - \alpha) \\ 0 & \beta \end{pmatrix} \mid \alpha/\beta \in S^1 \right\}.$$

We can easily compute the pointwise stabilizer of  $G_1, G_2, G_3$  in  $\tilde{\Gamma}$ . They are

$$\begin{aligned} \tilde{\Gamma}_{G_1} &= \left\{ \begin{pmatrix} i^\nu & 0 \\ 0 & i^\mu \end{pmatrix} \right\} = \left\{ \begin{pmatrix} i^\nu & 0 \\ 0 & i \end{pmatrix} \right\} \cdot z_{\tilde{\Gamma}} \\ \Gamma_{\tilde{G}_2} &= \left\{ \begin{pmatrix} i^\nu & \frac{1-i^\nu}{2} \\ 0 & 1 \end{pmatrix} \mid \frac{1-i^\nu}{2} \in \mathbb{Z}[i] \right\} \cdot Z_{\tilde{\Gamma}} = \left\{ \begin{pmatrix} \pm 1 & \frac{1\pm 1}{2} \\ 0 & 1 \end{pmatrix} \right\} \cdot Z_{\tilde{\Gamma}} \\ \Gamma_{\tilde{G}_3} &= \left\{ \begin{pmatrix} i^\nu & \frac{(1-i^\nu)(1+i)}{2} \\ 0 & 1 \end{pmatrix} \right\} \cdot Z_{\tilde{\Gamma}}, \end{aligned}$$

where in the last case we have to take into account that  $\frac{(1-i^\nu)(1+i)}{2} \in \mathbb{Z}[i]$  for all  $\nu$ .

Hence modulo the centre  $Z_{\tilde{\Gamma}}$  these stabilizers are cyclic groups of order 4, 2, 4.

The arcs  $D_i$  are also pointwise fixed under the action of certain cyclic groups, namely

$$\begin{aligned} D_1 &= \operatorname{Fix} \left( \begin{pmatrix} 0 & i \\ 1 & 0 \end{pmatrix} \right) \\ D_2 &= \operatorname{Fix} \left( \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right) \\ D_3 &= \operatorname{Fix} \left( \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix} \right), \end{aligned}$$

and we check easily that these arcs are geodesics joining the following points in the boundary

$$\begin{aligned} D_1 &\text{ runs from } \sqrt{i} \text{ to } -\sqrt{i} \\ D_2 &\text{ runs from } i \text{ to } -i \\ D_3 &\text{ runs from } e = e^{\frac{1\pi i}{6}} = e^{\frac{\pi i}{3}} \text{ to } \bar{\rho}. \end{aligned}$$

The corresponding tori are

$$\begin{aligned} T_1 = \text{Stab}(-1, 1) &= \left\{ \begin{pmatrix} \alpha & i\beta \\ \beta & \alpha \end{pmatrix} \right\} \\ T_2 = \text{Stab}(-\sqrt{i}, \sqrt{i}) &= \left\{ \begin{pmatrix} \alpha & \beta \\ \beta & \alpha \end{pmatrix} \right\} \\ T_3 = \text{Stab}(\rho, \bar{\rho}) &= \left\{ \begin{pmatrix} \delta - \beta & \beta \\ -\beta & \delta \end{pmatrix} \right\}. \end{aligned}$$

The torus  $T_2$  splits over  $\mathbb{Q}(i)$ , the other two tori split over a quadratic extension of  $\mathbb{Q}(i)$ .

Now it is not difficult anymore to describe the finite stabilizers and the corresponding fixed point sets. If  $x \in \mathbb{H}_3$  for which the stabilizer is bigger than  $Z_{\tilde{\Gamma}}$ , then we can conjugate  $x$  into  $F_0$ . It is very easy to see that  $x$  cannot lie in the interior of  $F_0$  because then we would get an identification of two points nearby  $x$  and hence still in  $F_0$  under  $\tilde{\Gamma}$ .

If  $x$  is on one of the lines  $D_1, D_2, D_3$  or on one of the arcs  $G_1, G_2, G_3$  but not on the intersection of two of them, then the stabilizer  $\Gamma_x$  is equal to  $Z_{\tilde{\Gamma}}$  times the cyclic group we attached to the line or the arc earlier. Finally we are left with the three special points

$$\begin{aligned} x_{12} &= D_1 \cap D_2 \cap G_1 = \{(0, 1)\} \\ x_{13} &= D_1 \cap D_3 \cap G_3 = \left\{ \left( \frac{1+i}{2}, \frac{\sqrt{2}}{2} \right) \right\} \\ x_{23} &= D_2 \cap D_3 \cap G_2 = \left\{ \left( \frac{1}{2}, \frac{\sqrt{3}}{2} \right) \right\}. \end{aligned}$$

In this case it is clear that the stabilizers are given by

$$\begin{aligned} \tilde{\Gamma}_{x_{12}} &= \left\langle \begin{pmatrix} 0 & i \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} i & 0 \\ 0 & 1 \end{pmatrix} \right\rangle = D_4 \\ \tilde{\Gamma}_{x_{13}} &= \left\langle \begin{pmatrix} 0 & i \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} i & 1 \\ 0 & 1 \end{pmatrix} \right\rangle = S_4 \\ \tilde{\Gamma}_{x_{23}} &= \left\langle \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix} \right\rangle = S_3. \end{aligned}$$

### 1.2.6 Compactification of $\Gamma \backslash X$

Our two special low dimensional examples show clearly that the quotient spaces  $\Gamma \backslash X$  are not compact in general. There exist various constructions to compactify them.

If, for instance,  $\Gamma \subset \text{Sl}_2(\mathbb{Z})$  is a subgroup of finite index, then the quotient  $\Gamma \backslash \mathbb{H}$  is a Riemann surface. It can be embedded into a compact Riemann surface by adding a finite number of points. This is a special case of a more general theorem of Satake and Baily-Borel: If the symmetric space  $X$  is actually hermitian symmetric (this means it has a complex structure) then we have the

structure of a quasi-projective variety on  $\Gamma \backslash X$ . This is the so-called Baily-Borel compactification. It exists only under special circumstances.

I will discuss the process of compactification in some more detail for our special low dimensional examples.


### Compactification of $\mathrm{Sl}_2(\mathbb{Z}) \backslash \mathbb{H}$ by adding points

Let  $\Gamma \subset \mathrm{Sl}_2(\mathbb{Z})$  be any subgroup of finite index. The group  $\Gamma$  acts on the rational projective line  $\mathbb{P}^1(\mathbb{Q})$ . We add it to the upper half plane and form

$$\bar{\mathbb{H}} = \mathbb{H} \cup \mathbb{P}^1(\mathbb{Q}),$$

and we extend the action of  $\Gamma$  to this space. Since the full group  $\mathrm{Sl}_2(\mathbb{Z})$  acts transitively on  $\mathbb{P}^1(\mathbb{Q})$  we find that  $\Gamma$  has only finitely many orbits on  $\mathbb{P}^1(\mathbb{Q})$ .

Now we introduce a topology on  $\bar{\mathbb{H}}$ . We defined a system of neighborhoods of points  $\frac{p}{q} = r \in \mathbb{P}^1(\mathbb{Q})$ . We define the Farey circles  $S\left(c, \frac{p}{q}\right)$  which touch the real axis in the point  $r = p/q$  ( $p, q = 1$ ) and have the radius  $\frac{c}{2q^2}$ . For  $c = 1$  we get the picture



Let us denote by  $D\left(c, \frac{p}{q}\right) = \cup_{c': 0 < c' \leq c} S\left(c', \frac{p}{q}\right)$  the Farey disks. For  $c \rightarrow 0$  these Farey disks  $D\left(c, \frac{p}{q}\right)$  define a system of neighborhoods of the point  $r = p/q$ . The Farey disks at  $\infty \in \mathbb{P}^1(\mathbb{Q})$  are given by the regions

$$D(T, \infty) = \{z \mid \Im(z) \geq T\}.$$

It is easy to check that an element  $\gamma \in \mathrm{Sl}_2(\mathbb{Z})$  which sends  $\infty \in \mathbb{P}^1(\mathbb{Q})$  into the point  $r = \frac{p}{q}$  sends  $D(T, \infty)$  to  $D\left(\frac{1}{T}, \frac{p}{q}\right)$ . These Farey disks  $D(c, r)$  do not meet provided we take  $c < 1$ . The considerations in 1.6.1 imply that the complement of the union of Farey disks is relatively compact modulo  $\Gamma$ , and since  $\Gamma$  has finitely many orbits on  $\mathbb{P}^1(\mathbb{Q})$ , we see easily that

$$Y_\Gamma = \Gamma \backslash \bar{\mathbb{H}}$$

is compact (which means of course also Hausdorff).

It is essential that the set of Farey circles  $D(c, r)$  and  $D\left(\frac{1}{c}, \infty\right)$  is invariant under the action of  $\Gamma$  on the one hand and decomposes into several connected components (which are labeled by the point  $r \in \mathbb{P}^1(\mathbb{Q})$ ) on the other hand. Hence

$$\Gamma \backslash \bigcup_r D(c, r) = \bigcup_{\Gamma r_i} \Gamma \backslash D(c, r_i)$$


where  $r_i$  is a set of representatives for the action of  $\Gamma$  on  $\mathbb{P}^1(\mathbb{Q})$  and where  $\Gamma_{r_i}$  is the stabilizer of  $r_i$  in  $\Gamma$ .

It is now clear that  $\Gamma_{r_i} \backslash D(c, r_i)$  is holomorphically equivalent to a punctured disc and hence the above compactification is obtained by filling the point into this punctured disc and this makes it clear that  $Y_\Gamma$  is a Riemann surface.

BSC

### 1.2.7 The Borel-Serre compactification of $\mathrm{Sl}_2(\mathbb{Z}) \backslash \mathbb{H}$

There is another construction of a compactification. We look at the disks  $D(c, r)$  and divide them by the action of  $\Gamma_r$ . For any point  $y \in S(c', r) - \{r\}$  there exists a unique geodesic joining  $r$  and  $y$ , passing orthogonally through  $S(c', r)$  and hitting the projective line in another point  $y_\infty$  ( $= -1/4$  in the picture below)



If  $r = \infty$ , then this system of geodesics is given by the vertical lines  $\{y \cdot I + x \mid x \in \mathbb{R}\}$ . This allows us to write the set

$$D(c, r) - \{r\} = X_{\infty, r} \times [c, 0]$$

where  $X_{\infty, r} = \mathbb{P}^1(\mathbb{R}) - \{r\}$ . The stabilizer  $\Gamma_r$  acts  $D(c, r)$  and on the right hand side of the identification it acts on the first factor, the quotient  $\Gamma_r \backslash X_{\infty, r}$  is a circle. Hence we can compactify the quotient

$$\Gamma_r \backslash D(c, r) - \{r\} \hookrightarrow \Gamma_r \backslash X_{\infty, r} \times [c, 0].$$

This gives us a second way to compactify  $\Gamma \backslash \mathbb{H}$ , we apply this process to a finite set of representatives of  $\mathbb{P}^1(\mathbb{Q}) \bmod \Gamma$ .

There is a slightly different way of looking at this. We may form the union

$$\mathbb{H} \cup \bigcup_r X_{\infty, r} = \tilde{\mathbb{H}}$$

and topologize it in such a way that

$$D(c, r) = X_{\infty, r} \times [c, 0] \subset X_{\infty, r} \times [c, 0]$$

is a local homeomorphism. Then we see that the compactification above is just the quotient

$$\Gamma \backslash \tilde{\mathbb{H}}$$

and the boundary is simply

$$\partial(\Gamma \backslash \mathbb{H}) = \Gamma \backslash \bigcup_{r \in \mathbb{P}^1(\mathbb{Q})} X_{\infty, r}.$$

This compactification is called the Borel-Serre compactification. Its relation to the Baily-Borel is such that the latter is obtained by the former by collapsing the circles at infinity to a point.

It is quite clear that a similar construction applies to the action of a group  $\Gamma \subset \mathrm{Sl}_2(\mathbb{Z}[i])$  on the three-dimensional hyperbolic space. The Farey circles will be substituted by spheres  $S(c, \alpha)$  which touch the complex plane  $\{(z, 0) \mid z \in \mathbb{C}\} \subset \mathbb{H}_3$  in the point  $(\alpha, 0)$ ,  $\alpha \in \mathbb{P}^1(\mathbb{Q}(i))$  and for  $\alpha = \infty$  the Farey sphere is the horizontal plane  $S(\infty, \zeta_0) = \{(z, \zeta_0) \mid z \in \mathbb{C}\}$ . An element  $\gamma \in \Gamma$  which maps  $(0, \infty)$  to  $\alpha$  maps  $S(\infty, \zeta_0)$  to  $S(c, \alpha)$ , where  $c = 1/\zeta_0$ . For a given  $\alpha$  we may identify the different spheres if we vary  $c$  and for any point  $\alpha \in \mathbb{P}^1(\mathbb{Q}(i))$  we define  $X_{\infty, \alpha} = \mathbb{P}^1(\mathbb{C}) \setminus \{\alpha\}$ . Again we can identify

$$D(c, \alpha) \setminus \{\alpha\} = X_{\infty, \alpha} \times (0, c] \subset \overline{D(c, \alpha) \setminus \{\alpha\}} = \partial(\Gamma \backslash \mathbb{H}) = X_{\infty, \alpha} \times [0, c]$$

The stabilizer  $\Gamma_\alpha$  acts on  $D(c, \alpha) \setminus \{\alpha\}$  and again this yields an action on the first factor. If we choose  $\alpha = \infty$  then

$$\Gamma_\infty = \left\{ \begin{pmatrix} \zeta & a \\ 0 & \zeta^{-1} \end{pmatrix} \mid \zeta \text{ root of unity, } a \in M_\infty \right\}$$

where  $M_\infty$  is a free rank 2 module in  $\mathbb{Z}[i]$ . If  $\zeta$  does not assume the value  $i$  then  $\Gamma_\infty \backslash X_{\infty, \infty}$  is a two-dimensional torus, a product of two circles. If  $\zeta$  assumes the value  $i$  then  $\Gamma_\infty \backslash X_{\infty, \infty}$  is a two dimensional sphere. If course we get the same result for an arbitrary  $\alpha$ .

Then we get an action of the group  $\Gamma$  on  $\tilde{\mathbb{H}}_3 = \mathbb{H}_3 \cup \bigcup_{\alpha \in \mathbb{P}^1(K)} \overline{D(c, \alpha) \setminus \{\alpha\}}$

and the quotient is compact.

The the set of orbits of  $\Gamma$  on  $\mathbb{P}^1(\mathbb{Q}(i))$  is finite, these orbits are called the cusps.

BSC0

### 1.2.8 The Borel-Serre compactification, reduction theory of arithmetic groups

This section could be skipped in a first reading. For the particular groups  $\mathrm{Sl}_2/\mathbb{Q}$  or  $\mathrm{Sl}_2(\mathbb{Z}[\sqrt{-d}])$  this compactification has been discussed in detail in the previous sections. A reader who is interested in the specific applications to number theory which will be discussed in the following chapters 2-5 only needs the results from section 1.2.7.

The Borel-Serre compactification works in complete generality for any semi-simple or reductive group  $G/\mathbb{Q}$ . To explain it, we need the notion of a parabolic subgroup of  $G/\mathbb{Q}$ .

A subgroup  $P/\mathbb{Q} \hookrightarrow G/\mathbb{Q}$  is parabolic if the quotient variety in the sense of algebraic geometry is a projective variety. We mentioned already earlier that

for the group  $\mathrm{Gl}_2/\mathbb{Q}$  we have an action of  $\mathrm{Gl}_2$  on the projective line  $\mathbb{P}^1$  and the stabilizers  $B_x$  of the points  $x \in \mathbb{P}^1(\mathbb{Q})$  are the so-called Borel subgroups of  $\mathrm{Gl}_2/\mathbb{Q}$ . They are maximal solvable subgroups and

$$\mathrm{Gl}_2/B_x = \mathbb{P}^1,$$

hence they are also parabolic.

More generally we get parabolic subgroups of  $\mathrm{Gl}_n/\mathbb{Q}$ , if we choose a flag on the vector space  $V = \mathbb{Q}^n = \mathbb{Q}e_1 \oplus \cdots \oplus \mathbb{Q}e_n$ . This is an increasing sequence of subspaces

$$\mathcal{F} : (0) = \{(0)\} = V_0 \subset V_1 \subset V_2 \subset \cdots \subset V_k = V.$$

The stabilizer  $P$  of such a flag is always a parabolic subgroup; the quotient space

$$G/P = \text{Variety of all flags of the given type,}$$

where the type of the flag is the sequence of the dimensions  $n_i = \dim V_i$ .

These flag varieties (the Grassmannians) are smooth projective schemes over  $\mathrm{Spec}(\mathbb{Z})$  and this implies that any flag  $\mathcal{F}$  is induced by a flag

$$\mathcal{F}_{\mathbb{Z}} : (0) = \{(0)\} = L_0 \subset L_1 \subset L_2 \subset \cdots \subset L_k = L = \mathbb{Z}^n \quad (1.36)$$

where  $L_i = V_i \cap L$ , and of course  $L_i \otimes \mathbb{Q} = V_i$ . This is the elementary fact which will be used later.

If our group  $G/\mathbb{Q}$  is the orthogonal group of a quadratic form

$$f(x_1, \dots, x_n) = \sum_{i=1}^n a_i x_i^2$$

with  $a_i \in K^*$ . Then we have to replace the flags by sequences of subspaces

$$\mathcal{F} : 0 \subset W_1 \subset W_2 \cdots \subset W_2^\perp \subset W_1^\perp \subset V,$$

where the  $W_i$  are isotropic spaces for the form  $f$ , i.e.  $f|_{W_i} \equiv 0$ , and where the  $W_i^\perp$  are the orthogonal complements of the subspaces. Again the stabilizers of these flags are the parabolic subgroups defined over  $\mathbb{Q}$ .

Especially, if the form  $f$  is anisotropic over  $\mathbb{Q}$ , i.e. there is no non-zero vector  $\underline{x} \in K^n$  with  $f(\underline{x}) = 0$ , then the group  $G/\mathbb{Q}$  does not have any parabolic subgroup over  $\mathbb{Q}$ . This equivalent to the fact that  $G(\mathbb{Q})$  does not have unipotent elements.

These parabolic subgroups always have a unipotent radical  $U_P$  which is always the subgroup which acts trivially on the successive quotients of the flag. The unipotent radical is a normal subgroup, the quotient  $P/U_P = M$  is a reductive group again, it is called the Levi-quotient of  $P$ .

We go back to the group  $\mathrm{Gl}_n/\mathbb{Q}$ . It contains the standard maximal torus whose  $R$  valued points are

$$T_0(R) = \{t = \begin{pmatrix} t_1 & 0 & \cdots & 0 \\ 0 & t_2 & \cdots & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & t_n \end{pmatrix} \mid t_i \in R^\times\} \quad (1.37)$$

It is a subgroup of the Borel subgroup (maximal solvable subgroup or minimal parabolic subgroup) whose  $R$ -valued points are

$$B_0(R) = \{ \underline{b} = \begin{pmatrix} t_1 & u_{1,2} & \cdots & u_{1,n} \\ 0 & t_2 & \cdots & u_{2,n} \\ 0 & 0 & \ddots & u_{n-1,n} \\ 0 & 0 & 0 & t_n \end{pmatrix} \mid t_i \in R^\times \} \quad (1.38)$$

and its unipotent radical  $U_0$  consists of those  $b \in B_0$  where all the  $t_i = 1$ . This unipotent radical contains the one dimensional root subgroups

$$U_{i,j} = \left\{ \begin{pmatrix} 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ 0 & 0 & \ddots & x & 0 \\ 0 & 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} \right\}, x \in R \quad (1.39)$$

where  $i < j$ , these one dimensional subgroups are isomorphic to the one dimensional additive group  $\mathbb{G}_a$ . They are normalized by the torus, for an element  $t \in T(R)$  and  $x_{i,j} \in U_{i,j}(R) = R$  we have

$$\underline{t} x_{i,j} \underline{t}^{-1} = t_i/t_j x_{i,j}. \quad (1.40)$$

For  $i = 1, \dots, n, j = 1, \dots, n, i \neq j$  (resp.  $i < j$ ) the characters  $\alpha_{i,j}(\underline{t}) = t_i/t_j$  are called the roots (resp. positive roots) of  $T_0$  in  $\mathrm{Gl}_n$ . We denote these systems of roots by  $\Delta^{\mathrm{Gl}_n}$  (resp)  $\Delta_+^{\mathrm{Gl}_n}$ . The one dimensional subgroups  $U_{i,j}, i \neq j$  are called the root subgroups.

Inside the set of positive roots we have the set of simple roots

$$\pi = \pi^{\mathrm{Gl}_n} = \{ \alpha_{1,2}, \dots, \alpha_{i,i+1}, \dots, \alpha_{n-1,n} \} \quad (1.41)$$

If we pass to the semi-simple subgroup  $\mathrm{Sl}_n/\mathbb{Q}$  then the torus and the Borel-subgroup has to be replaced by  $T_0^{(1)}, B_0^{(1)}$ , where we have  $\prod_i t_i = 1$ . The system of roots does not change, we have  $\pi = \pi^{\mathrm{Gl}_n} = \pi^{\mathrm{Sl}_n}$ .

We change the notation slightly, for  $i = 1, \dots, n-1$  we define  $\alpha_i := \alpha_{i,i+1}$  then for  $i < j$  we get  $\alpha_{i,j} = \alpha_i + \dots + \alpha_{j-1}$ , and  $\pi = \{ \alpha_1, \alpha_2, \dots, \alpha_{n-1} \}$

The Borel subgroup  $B_0$  is the stabilizer of the "complete" flag

$$\{0\} \subset \mathbb{Q}e_1 \subset \mathbb{Q}e_1 \oplus \mathbb{Q}e_2 \subset \cdots \subset \mathbb{Q}e_1 \oplus \mathbb{Q}e_2 \oplus \cdots \oplus \mathbb{Q}e_n, \quad (1.42)$$

the parabolic subgroups  $P_0 \supset B_0$  are the stabilizers of "partial" flags

$$\{0\} \subset \mathbb{Q}e_1 \oplus \cdots \oplus \mathbb{Q}e_{n_1} \subset \mathbb{Q}e_1 \oplus \cdots \oplus \mathbb{Q}e_{n_1} \oplus \mathbb{Q}e_{n_1+1} \oplus \cdots \oplus \mathbb{Q}e_{n_1+n_2} \subset \cdots \subset \mathbb{Q}^n. \quad (1.43)$$

The parabolic subgroup  $P_0$  also acts on the direct sum of the successive quotients

$$(\mathbb{Q}e_1 \oplus \cdots \oplus \mathbb{Q}e_{n_1}) \bigoplus (\mathbb{Q}e_{n_1+1} \oplus \cdots \oplus \mathbb{Q}e_{n_1+n_2}) \bigoplus \cdots \quad (1.44)$$

and this yields a homomorphism

$$r_{P_0} : P_0 \rightarrow M_0 = \mathrm{Gl}_{n_1} \times \mathrm{Gl}_{n_2} \times \cdots \quad (1.45)$$

hence  $M_0$  is the Levi quotient of  $P_0$ . By definition the unipotent radical  $U_{P_0}$  of  $P_0$  is the kernel of  $r_0$ . The semi-simple component will be  $M_0^{(1)} = \mathrm{Sl}_{n_1} \times \mathrm{Sl}_{n_2} \times \dots$

A parabolic subgroups  $P_0 \supset B_0$  defines a subset

$$\Delta^{P_0} = \{\alpha_{i,j} \in \Delta^{\mathrm{Gl}_n} \mid U_{i,j} \subset P_0\}$$

and the set decomposes into two sets

$$\Delta^{M_0} = \{\alpha_{i,j} \mid U_{i,j} \text{ and } U_{j,i} \subset \Delta^{P_0}\}; \quad \Delta^{U_{P_0}} = \Delta^{P_0} \setminus \Delta^{M_0}. \quad (1.46)$$

Intersecting this decomposition with the set  $\pi^{\mathrm{Gl}_n}$  yields a disjoint decomposition

$$\pi^{\mathrm{Gl}_n} = \pi^{M_0} \cup \pi^U \quad (1.47)$$

where  $\pi^U = \{\alpha_{n_1}, \alpha_{n_1+n_2}, \dots\}$ . In turn any such decomposition of  $\pi^{\mathrm{Gl}_n}$  yields a well defined parabolic  $P_0 \supset B_0$ .

If we choose another maximal split torus  $T_1$  and a Borel subgroup  $B_1 \supset T_1$  then this amounts to the choice of a second ordered basis  $v_1, v_2, \dots, v_n$  the  $v_i$  are given up to a non zero scalar factor. We can find a  $g \in \mathrm{Gl}_n(\mathbb{Q})$  which maps  $e_1, e_2, \dots, e_n$  to  $v_1, v_2, \dots, v_n$ , and hence we can conjugate the pair  $(B_0, T_0)$  to  $(B_1, T_1)$  and hence the parabolic subgroups containing  $B_0$  into the parabolic subgroups containing  $B_1$ . The conjugating element  $g$  also identifies

$$i_{T_0, B_0, T_1, B_1} : X^*(T_0) \xrightarrow{\sim} X^*(T_1)$$

and this identification does not depend on the choice of the conjugating element  $g$ . This allows us to identify the two set of positive simple roots  $\pi^{\mathrm{Gl}_n} \subset X^*(T_0)$  and  $\pi \subset X^*(T_1)$ . Eventually we can speak of the set  $\pi$  of simple roots of  $\mathrm{Gl}_n$ . Hence we have the fundamental fact

*The  $\mathrm{Gl}_n(\mathbb{Q})$  conjugacy classes of parabolic subgroups  $P/\mathbb{Q}$  are in one to one correspondence with the subsets  $\pi' = \pi^M$ . Then number of elements in  $\pi \setminus \pi' = \pi^U$  is called the rank of  $P$ , the set  $\pi'$  is called the type of  $P$ .*

We will denote the unipotent radical of  $P$  by  $U_P$  and the reductive quotient of  $P$  by  $U_P$  will be denoted by  $M_P = P/U_P$ . Then  $\pi' = \pi^{M_P}$ . We will also use a slightly different notation: If  $P$  is given then we also use  $U (= U_P)$  for the unipotent radical and  $M = P/U$  for the reductive quotient.

We formulated this result for  $\mathrm{Gl}_n/\mathbb{Q}$  but we can replace  $\mathbb{Q}$  by any field  $k$  and  $\mathrm{Gl}_n$  by any reductive group  $G/k$ . We have to define the system of relative simple positive roots  $\pi^G$  for any  $G/k$  (See [B-T]).

The group  $G/k$  itself is also a parabolic subgroup it corresponds to  $\pi' = \pi$ . We decide that we do not like it and hence we consider only proper parabolic subgroups  $P \neq G$ , i.e.  $\pi' \neq \emptyset$ . We can define the Grassmann variety  $\mathrm{Gr}^{[\pi']}$  of parabolic subgroups of type  $\pi'$ . This is a smooth projective variety and  $\mathrm{Gr}^{[\pi']}(\mathbb{Q})$  is the set of parabolic subgroups of type  $\pi'$ .

There is always a unique minimal conjugacy class it corresponds to  $\pi' = \emptyset$ . (In our examples this minimal class is given by the maximal flags, i.e. those flags where the dimension of the subspaces increases by one at each step (until we reach a maximal isotropic space in the case of an orthogonal group)). The

(proper) maximal parabolic subgroups are those for which  $\pi' = \pi \setminus \{\alpha_i\}$ , i.e.  $\pi^{U_{P_i}} = \{\alpha_i\}$

For any parabolic subgroup  $P/\mathbb{Q} \subset G/\mathbb{Q}$  we consider the character module  $X^*(P) := \text{Hom}(P/\mathbb{Q}, \mathbb{G}_m)$ . Since we do not have any non trivial homomorphisms from the unipotent  $U_P$  to  $\mathbb{G}_m$  we have  $\text{Hom}(P/\mathbb{Q}, \mathbb{G}_m) = \text{Hom}(M_P, \mathbb{G}_m)$ .

The reductive quotient  $M_P = M_P^{(1)} \cdot C_P$  where  $C_P$  is the central torus and  $M_P^{(1)}$  the semi-simple part ( the derived group). The quotient  $M_P/M_P^{(1)} = C'_P$  is a torus and  $C_P \rightarrow C'_P$  is an isogeny. Hence we have

$$\text{Hom}(P/\mathbb{Q}, \mathbb{G}_m) \otimes \mathbb{Q} = \text{Hom}(M_P, \mathbb{G}_m) \otimes \mathbb{Q} = \text{Hom}(C_P, \mathbb{G}_m) \otimes \mathbb{Q} = \text{Hom}(C'_P, \mathbb{G}_m) \otimes \mathbb{Q} \quad (1.48)$$

For a maximal parabolic subgroup  $P$  of type  $\pi' = \{\alpha_i\}$  we consider the module  $\text{Hom}(P, \mathbb{G}_m) \otimes \mathbb{Q} \subset X^*(T) \otimes \mathbb{Q}$ . Of course it always contains the determinant and

$$\text{Hom}(P, \mathbb{G}_m) \otimes \mathbb{Q} = \mathbb{Q}\gamma_i \oplus \mathbb{Q} \det$$

where  $\gamma_i$  is

$$\gamma_i(t) = \left( \prod_{\nu=1}^{\nu=i} t_\nu \right) \det(t)^{-i/n}. \quad (1.49)$$

These  $\gamma_i$  are called the dominant fundamental weights.

If our maximal parabolic subgroup is  $P/\mathbb{Q}$  is defined as the stabilizer of a flag  $0 \subset W \subset V = \mathbb{Q}^n$ , then the unipotent radical is  $U = \text{Hom}(V/W, W)$ . An element  $y \in P(\mathbb{Q})$  induces linear maps  $y_W, y_{V/W}$  and hence  $\text{Ad}(y)$  on  $U = \text{Hom}(V/W, W)$ . We get a character  $\gamma_P(y) = \det(\text{Ad}(y)) \in \text{Hom}(P, \mathbb{G}_m)$  which is called the sum of the positive roots. An easy computation shows that

$$n\gamma_i = \gamma_P \quad (1.50)$$

We add points at infinity to our symmetric space: We consider the disjoint union  $\cup_{\pi \neq \pi_G} \text{Gr}^{[\pi']}(\mathbb{Q})$  and form the space

$$\overline{X} = X \cup \bigcup_{\pi' \neq \emptyset} \text{Gr}^{[\pi']}(\mathbb{Q}).$$

This is the analogue of  $\mathbb{H} \cup \mathbb{P}^1(\mathbb{Q})$  in our first example, it is now more complicated because we have several Grassmannians, and we also have maps

$$r_{\pi_1, \pi_2} \text{Gr}^{[\pi_1]}(\mathbb{Q}) \rightarrow \text{Gr}^{[\pi_2]}(\mathbb{Q}) \text{ if } \pi_2 \subset \pi_1.$$

Our first aim is to put a topology on this space such that  $\Gamma \backslash \overline{X}$  becomes a compact Hausdorff space.

In our first example we interpreted the Farey circle  $D\left(c, \frac{p}{q}\right)$  with  $0 < c < 1$  as an open subset of points in  $\mathbb{H}$ , which are close to the point  $\frac{p}{q}$ , but far away from any other point in  $\mathbb{P}^1(\mathbb{Q})$ .

The point of reduction theory is that for any parabolic  $P \in \text{Gr}^{[\pi']}(\mathbb{Q})$  (here we also allow  $P = G$ ) we will define open sets

$$X^P(\underline{c}_{\pi'}, r(\underline{c}_{\pi'})) \subset X \quad (1.51)$$

which depend on certain parameters  $\underline{c}_{\pi'}, r(\underline{c}_{\pi'})$ . The points in  $X^P(\underline{c}_{\pi'}, r(\underline{c}_{\pi'}))$  should be viewed as the points, which are "very close" to the parabolic subgroup  $P$  (controlled by  $\underline{c}_{\pi'}$ ) but "keep a certain distance" (controlled by  $r(\underline{c}_{\pi'})$ ) to the parabolic subgroups  $Q \not\supset P$ . They are the analogues of the Farey circles. We will see:

a) This system of open sets is invariant under the action  $\mathrm{Gl}_n(\mathbb{Z})$

b) For  $P = G$  the set  $X^G(\emptyset, r_0)$  is relatively compact modulo the action of  $\mathrm{Gl}_n(\mathbb{Z})$ .

c) Any subgroup  $\Gamma \subset \mathrm{Gl}_n(\mathbb{Z})$  has only finitely many orbits on any  $\mathrm{Gr}^{[\pi]}(\mathbb{Q})$

d) For a suitable choice of the parameters  $\underline{c}_{\pi'}$ , and  $r(\underline{c}_{\pi'})$  we have :

$$X = \bigcup_P X^P(\underline{c}_{\pi'}, r(\underline{c}_{\pi'})) = X^G(\emptyset, r_0) \cup \bigcup_{P: P \text{ proper}} X^P(\underline{c}_{\pi'}, r(\underline{c}_{\pi'}))$$

and if  $P$  and  $P_1$  are conjugate and  $P \neq P_1$  then  $X^P(\underline{c}_{\pi'}, r(\underline{c}_{\pi'})) \cap X^{P_1}(\underline{c}_{\pi'}, r(\underline{c}_{\pi'})) = \emptyset$ .

Let us assume that we have constructed such a system of open sets, then c) and d) imply that for a given type  $\pi'$  we have

$$\Gamma \backslash \bigcup_{P: \text{type}(\pi')=\pi} X^P(\underline{c}_{\pi'}, r(\underline{c}_{\pi'})) = \bigcup \Gamma_{P_i} \backslash X^{P_i}(\underline{c}_{\pi'}, r(\underline{c}_{\pi'}))$$

where  $\{\dots, P_i, \dots\} = \Sigma(\pi, \Gamma)$  is a set of representatives of  $\mathrm{Gr}^{[\pi]}(\mathbb{Q})$  modulo the action of  $\Gamma$  and  $\Gamma_{P_i} = \Gamma \cap P_i(\mathbb{Q})$ .

This tells us that we have a covering

$$\Gamma \backslash X = \Gamma \backslash X^G(\emptyset, r_0) \cup \bigcup_{\pi' \neq \emptyset} \bigcup_{P \in \Sigma(\pi', \Gamma)} \Gamma_P \backslash X^P(\underline{c}_{\pi'}, r(\underline{c}_{\pi'})) \quad (1.52)$$

*The philosophy of reduction theory is that  $\Gamma \backslash X^G(\emptyset, r_0)$  is relatively compact and that we have an explicit description of the sets  $\Gamma_P \backslash X^P(\underline{c}_{\pi'}, r(\underline{c}_{\pi'}))$  as fiber bundles with nil manifolds as fiber over the locally symmetric spaces  $\Gamma_M \backslash X^M$ .*

We give the definition of the sets  $X^P(\underline{c}_{\pi'}, r(\underline{c}_{\pi'}))$ . We stick to the case that  $G = \mathrm{Gl}_n/\mathbb{Q}$  and  $\Gamma \subset \Gamma_0 = \mathrm{Gl}_n(\mathbb{Z})$  is a (congruence) subgroup of finite index. We defined the positive definite bilinear form (See 1.28)

$$\tilde{B}_{\Theta_x} = -\frac{1}{2n} B_{\Theta_x} : \mathfrak{g}_{\mathbb{R}} \times \mathfrak{g}_{\mathbb{R}} \rightarrow \mathbb{R}$$

and we have the identification  $\mathfrak{g}_{\mathbb{R}} \xrightarrow{\sim} T_e^{\mathrm{G}(\mathbb{R})}$ , and hence we get a euclidian metric on the tangent space  $T_e^{\mathrm{G}(\mathbb{R})}$  at the identity  $e$ . This extends to a left invariant Riemannian metric on  $G(\mathbb{R})$ , we denote it by  $d_{\Theta_x} s^2$ . Hence we get a volume form  $d_{\mathrm{vol}_H}^{\Theta_x}$  on any closed subgroup  $H(\mathbb{R}) \subset G(\mathbb{R})$ .

For any point  $x \in X$  and any parabolic subgroup  $P/\mathbb{Q}$  with unipotent radical  $U/\mathbb{Q}$  we define

$$p_P(P, x) = \text{vol}_U^{\Theta_x}(\Gamma_0 \cap U(\mathbb{R}) \backslash U(\mathbb{R})) \quad (1.53)$$

For the Arakelov-Chevalley scheme  $(\text{Gl}_n/\mathbb{Z}, \Theta_0)$  See(1.1.5) we have that  $\tilde{B}_{\Theta_0}(E_{i,j}) = 1$ . We have by construction

$$U_{i,j}(\mathbb{Z}) \backslash U_{i,j}(\mathbb{R}) = \mathbb{R}/\mathbb{Z} \quad (1.54)$$

and under this identification  $E_{i,j}$  maps to  $\frac{\partial}{\partial x}$ . Hence we get

$$d_{\text{vol}_{U_{i,j}}}^{\Theta_0}(U_{i,j}(\mathbb{Z}) \backslash U_{i,j}(\mathbb{R})) = 1$$

and from this we get immediately

**Proposition 1.2.1.** *For any parabolic subgroup  $P_0$  containing the torus  $T_0$  we have*

$$p_P(P_0, \Theta_0) = 1.$$

Let  $(L, <, >_x)$  be an Arakelov vector bundle and  $(\text{Gl}_n, \Theta_x)$  the corresponding Arakelov group scheme (of type  $\text{Gl}_n$ ) let

$$\mathcal{F}_{\mathbb{Z}} : (0) = \{(0)\} = L_0 \subset L_1 \subset L_2 \subset \dots \subset L_k = L = \mathbb{Z}^n$$

be a flag and  $P/\mathbb{Z}$  the corresponding parabolic subgroup. Then we have the homomorphism

$$r_P : P/\text{Spec}(\mathbb{Z}) \rightarrow M/\mathbb{Z} = \prod_{i=1}^{i=k} \text{Gl}(L_i/L_{i-1}) \quad (1.55)$$

with kernel  $U_P/\mathbb{Z}$ . The metric  $<, >_x$  on  $L \otimes \mathbb{R}$  yields an orthogonal decomposition

$$L \otimes \mathbb{R} = \bigoplus_{i=1}^{i=k} L_i/L_{i-1} \otimes \mathbb{R}$$

and hence an Arakelov bundle structure  $(L_i/L_{i-1}, (\Theta_x)_i)$  for all  $i$ , and therefore an Arakelov group scheme structure on  $M/\mathbb{Z}$ .

Hence we get

**Proposition 1.2.2.** *If  $(\text{Gl}_n, \Theta)$  is an Arakelov group scheme then  $\Theta$  induces an Arakelov group scheme structure  $\Theta^M$  on any reductive quotient  $M = P/U$ .*

**Definition :** A pair  $(\text{Gl}_n/\mathbb{Z}, \Theta)$  is called stable (resp. semi stable) if for any proper parabolic subgroup  $P/\mathbb{Q} \subset \text{Gl}_n/\mathbb{Q}$  we have

$$p_P(P, \Theta) > 1 \quad (1.56)$$

In our example in (1.2.6) the stable points are those outside the union of the closed Farey circles.

To get a better understanding of these numbers we have to do some computations with roots and weights. Let us start from an Arakelov vector bundle  $(L = \mathbb{Z}^d, \langle, \rangle)$  and let us assume that  $L$  is equipped with a complete flag

$$\mathcal{F}_0 = \{0\} = L_0 \subset L_1 \subset \cdots \subset L_{d-1} \subset L_d \quad (1.57)$$

which defines a Borel subgroup  $B/\mathbb{Z}$ . The quotients  $(L_i/L_{i-1}, \langle, \rangle_i)$  are Arakelov line bundles over  $\mathbb{Z}$  or in a less sophisticated language they are free modules of rank one and the generating vector  $\bar{e}_i$  has a length  $\sqrt{\langle \bar{e}_i, \bar{e}_i \rangle_i}$ . This length is of course also the volume of  $(L_i/L_{i-1} \otimes \mathbb{R})/(L_i/L_{i-1})$ .

The unipotent radical  $U/\mathbb{Z} \subset B/\mathbb{Z}$  has a filtration  $\{(0)\} \subset V_1 \subset \cdots \subset V_{n(n-1)/2-1} \subset V_{n(n-1)/2} = U$  by normal subgroups, the successive quotients are isomorphic to  $\mathbb{G}_a$  and the torus  $T = B/U$  acts by a positive root  $\alpha_{i,j}$  and this is a one to one correspondence between the subquotients and the positive roots. Then it is clear: If  $\nu$  corresponds to  $(i, j)$  then

$$(V_\nu/V_{\nu+1}, \Theta_\nu) = (L_i/L_{i-1}, \langle, \rangle_i) \otimes (L_j/L_{j-1}, \langle, \rangle_j)^{-1}. \quad (1.58)$$

Moreover the quotients  $(V_\nu/V_{\nu+1}, \Theta_\nu)$  depend only on the conformal class of  $\langle, \rangle$  and hence only on the resulting Cartan involution  $\Theta$ .

The unipotent subgroup  $U/\mathbb{Z}$  contains the one parameter subgroup  $U_{i,j}/\mathbb{Z}$  and this one parameter subgroup maps isomorphically to  $(V_\nu/V_{\nu+1})$ . Hence our construction defines the Arakelov line bundle  $(U_{i,j}, \Theta_{i,j})$ .

If we now define  $n_{\alpha_{i,j}}(B, x) = \text{vol}_{\Theta_{i,j}}(U_{i,j}(\mathbb{R})/U_{i,j}(\mathbb{Z}))$  then it is clear that

$$p_B(B, x) = \prod_{i < j} n_{\alpha_{i,j}}(B, x) \quad (1.59)$$

If  $P \supset B$  then its unipotent radical  $U_P \subset U$  and we defined the set  $\Delta^{U_P}$  as the set of positive roots for which  $U_{i,j} \subset U_P$ . Then we have

$$p_P(B, x) = \prod_{(i,j) \in \Delta^{U_P}} n_{\alpha_{i,j}}(B, x) \quad (1.60)$$

Here it is important to notice the right hand side does not depend on the choice of  $B \subset P$ .

We follow a convention and put  $2\rho_P = \sum_{(i,j) \in \Delta^{U_P}} \alpha_{i,j}$  so that  $\rho_P$  is the half sum of positive roots in the unipotent radical. Formula (1.50) tells us that for any maximal parabolic subgroup  $P_{i_0}$

$$2\rho_{P_{i_0}} = \sum_{i \leq i_0, j \geq i_0+1} \alpha_{i,j} = n\gamma_{i_0}. \quad (1.61)$$

For any  $\gamma = \sum z_i \alpha_{i,i+1} \in X^*(T) \otimes \mathbb{C}$  we define the homomorphism

$$|\gamma| : T(\mathbb{R}) \rightarrow \mathbb{C}^\times : |\gamma| : t \rightarrow \prod_i |\alpha_{i,i+1}(t)|^{z_i} \quad (1.62)$$

Since the numbers  $n_{\alpha_{i,j}}(B, x)$  are positive real numbers we define for any

$$n_\gamma(B, x) = \prod_{i=1}^{n-1} n_{\alpha_{i,j}}(B, x)^{x_i}. \quad (1.63)$$

Here we see that the second argument is a Borel-subgroup  $B$ . But if the above character  $\gamma : B(\mathbb{R}) \rightarrow \mathbb{R}_{>0}^\times$  extends to a character  $\gamma : P(\mathbb{R}) \rightarrow \mathbb{R}_{>0}^\times$  then we can define

$$n_\gamma(P, x) := n_\gamma(B, x)$$

and this number only depends on  $P$  and not on the Borel subgroup  $B \subset P$ . The characters in  $\gamma \in X^*(T)$  for which  $|\gamma|$  extend to  $P(\mathbb{R})$  are exactly the linear combinations (See (1.65) below)  $\gamma = \sum_{\alpha_i \in \pi^U} x_i \gamma_i$ . The characters  $\gamma_P = \sum_{\alpha_i \in \pi^U} r_i \gamma_i$  where the  $r_i > 0$  are rational numbers. Let  $P_i$  be the maximal parabolic subgroup of type  $\pi \setminus \{\alpha_i\}$  containing  $P$  then the above formula implies that

$$p_P(P, x) = \prod_{\alpha_i \in \pi^U} n_{\gamma_i}(P_i, x)^{r_i} = \prod_{\alpha_i \in \pi^U} p_{P_i}(P_i, x)^{\frac{r_i}{n}} \quad (1.64)$$

This tells us

*The Arakelov scheme  $(\mathrm{Gl}_n/\mathbb{Z}, \Theta)$  is stable if for all maximal parabolic subgroups  $p_{P_i}(P_i, \Theta) = n_{\gamma_i}(P_i, \Theta)^n > 1$ .*

We need a few more formulas relating roots and weights. For any parabolic subgroup we have the division of the set of simple roots into two parts

$$\pi = \pi^M \cup \pi^{U_P}.$$

This induces a splitting of the character module split

$$X^*(T) \otimes \mathbb{Q} = \bigoplus_{\alpha_i \in \pi^M} \mathbb{Q}\alpha_i \oplus \bigoplus_{\alpha_i \in \pi^{U_P}} \mathbb{Q}\gamma_i \quad (1.65)$$

where  $\gamma_i$  is the dominant fundamental weight attached to  $\alpha_i$  (See (1.49)).

If now  $\alpha_i \in \pi^{U_P}$  then we can project  $\alpha_i$  to the second component, this projection

$$\alpha_i^P = \alpha_i + \sum_{\alpha_\nu \in \pi^M} c_{i,\nu} \alpha_\nu \quad (1.66)$$

Here an elementary - but not completely trivial - computation shows that

$$c_{i,\nu} \geq 0 \quad (1.67)$$

Since  $\alpha_i^P \in \bigoplus_{\alpha_i \in \pi^{U_P}} \mathbb{Q}\gamma_i$  these characters extend to  $P(\mathbb{R})$  and hence  $n_{\alpha_i^P}(P, x)$  is defined.

We state the two fundamental theorems of reduction theory

**Theorem 1.2.1.** *For any Arakelov group scheme  $(\mathrm{Gl}_n, \Theta_x)$  we can find a Borel subgroup  $B \subset \mathrm{Gl}_n$  for which*

$$n_{\alpha_i}(B, \Theta_x) = n_{\alpha_i}(B, x) < \frac{2}{\sqrt{3}} \text{ for all } i = 1, \dots, n-1$$

**Theorem 1.2.2.** *For any Arakelov group scheme  $(\mathrm{Gl}_n, \Theta)$  we can find a unique parabolic subgroup  $P$  such that for all  $\alpha_i \in \pi^{U_P}$  we have*

$$n_{\alpha_i^P}(P, \Theta) < 1 \text{ and he reductive quotient } (M, \Theta^M) \text{ is semi stable.}$$

The first theorem is due to Minkowski, the second theorem is proved in [Stu], [Gray].

This parabolic subgroup is called the canonical destabilizing group. We denote it by  $P(x)$ , if  $(G, x)$  is semi stable then  $P(x) = G$ . This gives us a dissection of  $X$  into the subsets

$$X = \bigcup_{P: \text{parabolic subgroups of } G/\mathbb{Q}} X^{[P]} = \{x \in X \mid P(x) = P\} \quad (1.68)$$

Clearly  $\gamma X^{[P]} = X^{[\gamma P \gamma^{-1}]}$ , if we divide by the group  $\Gamma$  the we get

$$\Gamma \backslash X = \bigcup_{P \in \mathrm{Par}(\Gamma)} \Gamma_P \backslash X^{[P]} \quad (1.69)$$

where  $\mathrm{Par}(\Gamma)$  is a set of representatives of  $\Gamma$  conjugacy classes of parabolic subgroups of  $\mathrm{Gl}_n/\mathbb{Q}$ . This is a decomposition of  $\Gamma \backslash X$  into a disjoint union of subsets. The subset  $\Gamma \backslash X^{[\mathrm{Gl}_n]}$  is compact, it is the set of semi stable pairs  $(x, \mathrm{Gl}_n)$ , the subsets  $\Gamma_P \backslash X^{[P]}$  for  $P \neq G$  are in a certain sense "open in some directions" and "closed in some other direction". Therefore this decomposition is not so useful for the study of cohomology groups.

To remedy this we introduce larger subsets. For a real number  $r, 0 < r < 1$  we define  $\boxed{\mathrm{Gstable}}$

$$X^{\mathrm{Gl}_n}(r) = \{x \in X \mid n_{\gamma_\alpha}(P(x), x) > r, \text{ for all } \alpha \in \pi^{U_{P(x)}}\}. \quad (1.70)$$

It contains the set of semi-stable  $(\mathrm{Gl}_n, x)$  If we choose  $r < 1$  but close to one then some of the elements in  $X^{\mathrm{Gl}_n}(r)$  may be unstable but only a "little bit".

Together with the first theorem this has a consequence

**Proposition 1.2.3.** *The quotient  $X^{\mathrm{Gl}_n}(r) = \Gamma \backslash X^{\mathrm{Gl}_n}(r)$  is relatively compact open subset of  $\Gamma \backslash X$ , It contains the set of semi-stable  $(\mathrm{Gl}_n, x)$ .*

We start from a parabolic subgroup  $P$  and let  $M = P/U_P$  be its Levi-quotient. Our considerations above also apply to  $M/\mathbb{Q}$ . The group  $P(\mathbb{R})$  acts transitively on  $X$  and we put (See (1.55))

$$X^M = U_P(\mathbb{R}) \backslash X \text{ and let } q_M : X \rightarrow X^M \text{ be the projection ,}$$

here  $X^M = M(\mathbb{R})/K_\infty^M$  where  $K_\infty^M$  is the image of  $P(\mathbb{R}) \cap K_\infty$  in  $M(\mathbb{R})$ . Let  $S \subset M$  be the maximal split torus in the center of  $M$  then we define

$$X^{M^{(1)}} := M(\mathbb{R})/K_\infty^M \cdot S^{(0)}(\mathbb{R}) \quad (1.71)$$

where of course  $S^{(0)}(\mathbb{R})$  is the connected component of the identity of  $S(\mathbb{R})$ , For a simple roots  $\alpha \in \pi^M$ , a Borel subgroup  $\bar{B} \subset M/\mathbb{Q}$  and a point  $x^M = q_M(x)$

we can define the numbers  $n_\alpha(\bar{B}, x^M)$  essentially in the same way as before and clearly

$$n_\alpha(\bar{B}, x^M) = n_\alpha(B, x)$$

if  $B$  is the inverse image of  $\bar{B}$ .

We have to be a little bit careful with the numbers  $p_{\bar{Q}}(\bar{Q}, x^M)$  because the for the inverse image  $Q$  the unipotent radical  $U_Q$  is larger than  $U_{\bar{Q}}$ . Therefore we have to look at the dominant fundamental weights  $\gamma_\alpha^M \in \bigoplus_{\alpha_i \in \pi^M} \mathbb{Q}\alpha_i$ , and formulate the stability condition for  $x^M$  in terms of these  $\gamma_\alpha^M$ :

*The point  $x^M$  is stable, if for all  $\alpha_i \in \pi^M$  the inequality  $n_{\gamma_{\alpha_i}^M}(\bar{P}_{\alpha_i}, x^M) > 1$  holds. Again we denote the destabilizing group by  $P(x^M)$ .*

Hence we see that for a number  $r_M < 1$  we can define regions

$$X^M(r_M) = \{x^M | n_{\gamma_{\alpha_i}^M}(\bar{P}_{\alpha_i}, x^M) > r_M \text{ whenever } \bar{P}_{\alpha_i} \supset \bar{P}(x^M)\} \quad (1.72)$$

We choose numbers  $0 < c_P < 1$ , furthermore we choose a number  $r(c_P) < 1$  and define

$${}^*X^P(c_P, r(c_P)) = \{x | n_{\alpha^P}(P, x) < c_P \text{ for all } \alpha \in \pi^{U_P}; x^M \in X^M(r(c_P))\} \quad (1.73)$$

**Proposition 1.2.4.** *For a given  $r(c_P) < 1$  we can find numbers  $c_P$  such that that for any  $x \in {}^*X^P(c_P, r(c_P))$  the destabilising parabolic subgroup  $P(x) \subset P$ . The same is true in the other direction: For a given  $0 < c_P < 1$  we can find  $r < 1$  such that for  $x \in {}^*X^P(c_P, r)$  the destabilising parabolic subgroup  $P(x) \subset P$ .*

To see this we have to look at the destabilising subgroup  $\bar{Q} \subset (M, x_M)$ . Its inverse image  $Q \subset P$  is a parabolic subgroup of  $\text{Gl}_n$ . The reductive quotient  $(\bar{M}, x_{\bar{M}})$  of  $Q$  is semi-stable. We want to show that  $Q$  is the destabilising parabolic of  $(\text{Gl}_n, x)$ . We have to show that

$$n_{\alpha^Q}(Q, x) < 1 \quad \forall \alpha \in \pi^{U_Q} = \pi^{U_P} \cup \pi^{U_{\bar{Q}}}.$$

For  $\alpha \in \pi^{U_{\bar{Q}}}$  this is true by definition. For  $\alpha \in \pi^{U_P}$  we have

$$\alpha^P = \alpha + \sum_{\beta \in \pi^M} a_{\alpha, \beta} \beta \quad \text{and} \quad \alpha^Q = \alpha + \sum_{\beta' \in \pi^M} a'_{\alpha, \beta'} \beta,$$

where  $a_{\alpha, \beta} \geq 0$ . The roots  $\beta \in \pi^{U_{\bar{Q}}}$  can be expressed in terms of the  $\beta^{\bar{Q}} = \beta^Q$ :

$$\beta^Q = \beta + \sum_{\beta' \in \pi^{\bar{M}}} a_{\beta, \beta'}^* \beta' \quad (1.74)$$

and hence

$$\alpha^Q = \alpha^P - \sum_{\beta \in \pi^{U_{\bar{Q}}}} a_{\alpha, \beta} \beta^Q + \sum_{\beta' \in \pi^{\bar{M}}} c_{\alpha \beta'} \beta'. \quad (1.75)$$

The last sum is zero because  $\alpha^Q, \alpha^P, \beta^Q$  are orthogonal to the module  $\bigoplus_{\beta'} \mathbb{Z}\beta'$ .

We get the relation

$$n_{\alpha Q}(Q, x) = n_{\alpha P}(P, x) \cdot \prod_{\beta \in \pi^{U_Q}} n_{\beta Q}(Q, x)^{-a_{\alpha, \beta}}. \quad (1.76)$$

Now it comes down to show that  $\boxed{\text{wc}}$

$$\begin{aligned} n_{\alpha P}(P, x) < c_\alpha, \forall \alpha \in \pi^{U_P} \text{ and } n_{\beta Q}(Q, x) > r, \forall \beta \in \pi^{U_Q} \\ \implies n_{\alpha Q}(P, x) < 1; \forall \alpha \in \pi^{U_P} \end{aligned} \quad (1.77)$$

This is certainly true if either the  $c_\alpha$  are small enough or if  $r$  is sufficiently close to one. In this case we say that  $(c_P, r)$  is well chosen.

Therefore we define

$$X^P(c_P, r(c_P)) = \{x \in {}^*X^P(c_P, r(c_P)) \mid P(x) \subset P\}$$

we have  $X^P(c_P, r(c_P)) = {}^*X^P(c_P, r(c_P))$ , if  $(c_P, r(c_P))$  is well chosen.

We claim that we can find a family of parameters

$$(\dots, (c_P, r(c_P)), \dots)_{P: \text{parabolic over } \mathbb{Q}}$$

where  $(c_P, r(c_P))$  only depend on the type of  $P$ , such that we get a covering

$\boxed{\text{COV}}$

$$X = \bigcup_P X^P(c_P, r(c_P)) \quad (1.78)$$

and hence

$$\Gamma \backslash X = \Gamma \bigcup_P \backslash X^P(c_P, r(c_P)) = \bigcup_{P \in \text{Par}(\Gamma)} \Gamma_P \backslash X^P(c_P, r(c_P))$$

We change the notation slightly, since these numbers only depend on the type  $\pi' = \pi^M = t(P)$  we replace  $c_P$  by  $c_{\pi'}$  and  $r(c_P)$  by  $r(c_{\pi'})$ .

To prove the claim we choose a number  $0 < c_\emptyset < 1$ . In this case  $r_0 = r(c_\emptyset)$  can be any number. Then we choose a number  $0 < r_1 < c_\emptyset$ . For any  $\pi_i = \{\alpha_i\}$  we choose a  $c_{\pi_i} < 1$  such that  $(c_{\pi_i}, r_1)$  is well chosen. We continue and chose  $0 < r_2 < c_{\pi_i}$  for all  $i$  and for any two element subset  $J \subset \pi$  we choose numbers  $0 < c_J < 1$  such that  $(c_J, r_2)$  is well chosen. This goes until we reach top parabolic.

Now we get a covering of  $X$  by the open sets  $X^P(c_\pi, r(\pi))$ . To see this we pick a point  $x \in X$ , we have to show that it lies in at least one of the sets  $X^P(c_P, r(c_P))$ . If it is not in  $X^{\text{Gl}_n}(r_{n-1})$  then we find a maximal parabolic  $P_i$  such that  $n_{\alpha_i}(P_i, x) < c_{\pi \setminus \{\alpha_i\}}$ . We project  $x$  to the point  $x^{M_i} \in X^{M_i}$ . If this point is in  $X^{M_i}(r_{n-2})$  then  $x \in X^{P_i}(c_{\pi \setminus \{\alpha_i\}}, r_{n-2})$  and we are done. If not we apply our argument above to  $x^{M_i}$  and  $\pi' = \pi \setminus \{\alpha_i\}$ . We continue the same reasoning and at latest it stops for  $\pi' = \emptyset$ .

We have a very explicit description of these sets  $\Gamma_P \backslash X^P(c_{\pi'}, r(c_{\pi'}))$ . We consider the evaluation map

$$\begin{aligned} n^{\pi_{U_P}} : \Gamma_P \backslash X^P(c_{\pi'}, r(c_{\pi'})) &\rightarrow \prod_{\alpha \in \pi_{U_P}} (0, c_\alpha) \\ x &\mapsto (\dots, n_{\alpha^P}(P, x), \dots)_{\alpha \in \pi_{U_P}} \end{aligned} \quad (1.79)$$

Of course we also have the homomorphism

$$|\alpha^{\pi_{U_P}}| : P(\mathbb{R}) \rightarrow \{\dots, |\alpha^P|, \dots\}_{\alpha \in \pi_{U_P}} \quad (1.80)$$

and the multiplication by an element  $y \in P(\mathbb{R})$  induces an isomorphism of the fibers

$$(n^{\pi_{U_P}})^{-1}(c_1) \xrightarrow{\sim} (n^{\pi_{U_P}})^{-1}(c_2) \text{ if } |\alpha^{\pi_{U_P}}|(y) \cdot c_1 = c_2$$

where the multiplication is taken componentwise. This identification depends on the choice of  $y$ .

To get a canonical identification we use the geodesic action which is introduced in the paper by Borel and Serre. We define an action of  $A = (\prod_{\alpha \in \pi \setminus \pi'} \mathbb{R}_{>0}^\times)$  on  $X$ . This action depends on  $P$  and we denote it by

$$(a, x) \mapsto a \bullet x \quad (1.81)$$

A point  $x \in X$  defines a Cartan involution  $\Theta_x$  and then the parabolic subgroup  $P^{\Theta_x}$  of  $G \times \mathbb{R}$  is opposite to  $P \times \mathbb{R}$  and  $P \times \mathbb{R} \cap P^{\Theta_x} = M_x$  is a Levi factor, the projection  $P \rightarrow M$  induces an isomorphism

$$\phi_x : M \times \mathbb{R} \xrightarrow{\sim} M_x. \quad (1.82)$$

The character  $\alpha^{\pi'}$  induces an isomorphism

$$s_x : A \xrightarrow{\sim} S_x(\mathbb{R})^{(0)}$$

where  $S_x$  is the maximal Hence we  $S_x(\mathbb{R})^{(0)}$  is the connected component of the identity of the center  $M_x(\mathbb{R}) \cap \text{Sl}_n(\mathbb{R})$  and we put

$$a \bullet x = s_x(a)x$$

We have to verify that this is indeed an action. This is clear because for the Cartan-involution  $\Theta_{a \bullet x}$  we obviously have

$$P^{\Theta_x} = P^{\Theta_{a \bullet x}}.$$

It is also clear that this action commutes with the action of  $P(\mathbb{R})$  on  $X$  because

$$y s_x(a)x = s_{yx}(a)yx \text{ for all } y \in P(\mathbb{R}), x \in X.$$

It follows from the construction that the semigroup  $A_- = \{\dots, a_\nu, \dots\}$  - where  $0 < a_\nu \leq 1$  - acts via the geodesic action on  $X^P(c_\pi, r(c_{\pi'}))$  and that for  $a \in A_-$  we get an isomorphism

$$(n^{\pi_{U_P}})^{-1}(c_1) \xrightarrow{\sim} (n^{\pi_{U_P}})^{-1}(ac_1).$$

This yields a decomposition

$$X^P(c_{\pi'}, r(\underline{c}_{\pi'})) = (n^{\pi^{U_P}})^{-1}(c_0) \times \prod_{\alpha \in \pi'} (0, c_\alpha]$$

where  $c_0$  is an arbitrary point in the product.

Since we know that  $|\alpha^{\pi'}|$  is trivial on  $\Gamma_P$  and since the action of  $P$  commutes with the geodesic action we conclude

$$\Gamma_P \backslash X^P(c_{\pi'}, r(\underline{c}_{\pi'})) = \Gamma_P \backslash (n^{\pi'})^{-1}(c_0) \times \prod_{\alpha \in \pi'} (0, c_\alpha] \quad (1.83)$$

Let  $P^{(1)}(\mathbb{R}) = \ker(|\alpha^{\pi^{U_P}}|)$  then the fiber  $(n^{\pi'})^{-1}(c_0)$  is a homogenous space under  $P^{(1)}(\mathbb{R})$ . We have the symmetric space  $X^M$  attached to  $M$ , to be precise this is

$$X^M = M(\mathbb{R})/K_\infty$$

We have the projection map  $p_{P,M} : X \rightarrow X^M$  where  $X^M$  is the space of Cartan involutions on the reductive quotient  $M$ . Hence we get a map

$$p_{P,M}^* = p_{P,M} \times n^{\pi^{U_P}} : X \rightarrow X^M \times \prod_{\alpha \in \pi'} (0, c_\alpha] \quad (1.84)$$

The geodesic action only acts on the second factor of the product  $X^M \times \prod_{\alpha \in \pi'} (0, c_\alpha]$ , the map  $p_{P,M}^*$  commutes with the geodesic action.

The group  $U_P(\mathbb{R})$  acts simply transitively on the fibers of this projection, and hence

$$q_{P,M} : \Gamma_P \backslash X^P(c_{\pi'}, r(\underline{c}_{\pi'})) \rightarrow \Gamma_M \backslash X^M(r(\underline{c}_P)) \times \prod_{\alpha \in \pi'} (0, c_\alpha] \quad (1.85)$$

is a fiber bundle with fiber isomorphic  $\Gamma_U \backslash U(\mathbb{R})$ . If we pick a point  $\Gamma_M \backslash X^M(r(\underline{c}_P)) \times \prod_{\alpha \in \pi'} (0, c_\alpha]$  then the identification of  $q_{P,M}^{-1}(\cdot)$  with  $\Gamma_U \backslash U(\mathbb{R})$  depends on the choice of a point  $\tilde{x} \in X^P(c_{\pi'}, r(\underline{c}_{\pi'}))$  which maps to  $x$ .

This can now be compactified, we define the closure

$$\overline{\Gamma_P \backslash X^P(c_{\pi'}, r(\underline{c}_P))} := \Gamma_P \backslash (n^{\pi^{U_P}})^{-1}(c_0) \times \prod_{\alpha \in \pi_G \backslash \pi} [0, c_{\pi'}],$$

and

$$\partial \overline{\Gamma_P \backslash X^P(c_{\pi'}, \Omega_\pi)} = \overline{\Gamma_P \backslash X^P(c_{\pi'}, \Omega_\pi)} \setminus \Gamma_P \backslash X^P(c_\pi, \Omega_\pi)$$

this is equal to

$$\partial \overline{\Gamma_P \backslash X^P(c_{\pi'}, r(\underline{c}_P))} = \Gamma_P \backslash (n^{\pi^{U_P}})^{-1}(c_0) \times \partial \left( \prod_{\nu \in \pi_G \backslash \pi} [0, c_\pi] \right)$$

where of course  $\partial \left( \prod_{\nu \in \pi_G \backslash \pi} [0, c_\pi] \right) \subset \prod_{\nu \in \pi_G \backslash \pi} [0, c_\pi]$  is the subset where at least one of the coordinates is equal to zero.

We form the disjoint union of of these boundaries over the  $\pi$  and set of representatives of  $\Gamma$  conjugacy classes, this is a compact space. Now there is

still a minor technical point. If we have two parabolic subgroups  $Q \subset P$  then the intersection  $X^P(\underline{c}_P, r(\underline{c}_P)) \cap X^Q(\underline{c}_Q, r(\underline{c}_Q)) \neq \emptyset$ . If we now have points

$$x \in \overline{\partial\Gamma_P \backslash X^P(c_\pi, r(\underline{c}_P))}, y \in \overline{\partial\Gamma_Q \backslash X^Q(c_{\pi'}, r(\underline{c}_{P'}))}$$

then we identify these two points if we have a sequence of points  $\{x_n\}_{n \in \mathbb{N}}$  which lies in the intersection  $X^P(c_\pi, r(\underline{c}_P)) \cap X^Q(c_{\pi'}, r(\underline{c}_{P'}))$  and which converges to  $x$  in  $\overline{\Gamma_P \backslash X^P(c_\pi, r(\underline{c}_P))}$  and to  $y$  in  $\overline{\Gamma_Q \backslash X^Q(c_{\pi'}, r(\underline{c}_{P'}))}$ . A careful inspection shows that this provides an equivalence relation  $\sim$ , and we define

$$\partial(\Gamma \backslash X) = \bigcup_{\pi', P \in \text{Par}(\Gamma)} \overline{\partial\Gamma_P \backslash X^P(c_\pi, r(\underline{c}_P))} / \sim$$

and the Borel-Serre compactification will be the manifold with corners

$$\overline{\Gamma \backslash X} = \Gamma \backslash (X \cup \bigcup_{P: P \text{ proper}} \overline{X^P(c_{\pi'}, r(\underline{c}_P))}). \quad (1.86)$$

We define a "tubular" neighborhood of the boundary we put

$$\mathcal{N}(\Gamma \backslash X) = \Gamma \backslash \bigcup_{P: P \text{ proper}} \overline{X^P(c_{\pi'}, r(\underline{c}_P))} \quad (1.87)$$

and then we define the "punctured tubular" neighborhood as

$$\dot{\mathcal{N}}(\Gamma \backslash X) = \Gamma \backslash \bigcup_{P: P \text{ proper}} X^P(c_{\pi'}, r(\underline{c}_P)) = \Gamma \backslash X \cap \mathcal{N}(\Gamma \backslash X) \quad (1.88)$$

Eventually we want to use the above covering as a tool to understand cohomology (See section 8.1.8) For this it is also necessary to understand the intersections

$$X^{P_1}(c_{\pi_1}, r(\underline{c}_{\pi_1})) \cap \cdots \cap X^{P_k}(c_{\pi_k}, r(\underline{c}_{\pi_k})) \quad (1.89)$$

Our proposition 1.2.4 implies that for any point  $x$  in the intersection the destabilizing parabolic subgroup  $P(x) \subset P_1 \cap \cdots \cap P_k$ . Hence we see that the above intersection can only be non empty if  $Q = P_1 \cap \cdots \cap P_k$  is a parabolic subgroup. Then  $\pi^{U_Q} = \cup_{\nu=1}^k \pi^{U_{P_\nu}}$ . Let  $M$  be the reductive quotient of  $Q$ .

Now we look at the product  $\prod_{\alpha \in \pi^{U_Q}} \mathbb{R}_{>0}^\times$ , here it seems to be helpful to identify it - using the logarithm - with  $\mathbb{R}^{d_Q}$ :

$$\log : \prod_{\alpha \in \pi^{U_Q}} \mathbb{R}_{>0}^\times \xrightarrow{\sim} \mathbb{R}^{d_Q} \quad (1.90)$$

We consider the map

$$\begin{aligned} N^Q : X^{P_1}(c_{\pi_1}, r(\underline{c}_{\pi_1})) \cap \cdots \cap X^{P_k}(c_{\pi_k}, r(\underline{c}_{\pi_k})) &\rightarrow \mathbb{R}^{d_Q} \\ N^Q : x &\mapsto (\dots, -\log(n_{\alpha Q}(Q, x)), \dots)_{\alpha Q \in \pi^{U_Q}} \end{aligned} \quad (1.91)$$

Consider a point  $x \in X^{P_\nu}(c_{\pi_\nu}, r(\underline{c}_{\pi_\nu}))$ , for  $\alpha \in \pi^{U_{P_\nu}}$  we have

$$-\log(n_{\alpha P_\nu}(P_\nu, x)) \geq -\log(c_{\pi_\nu})$$

We can express  $-\log(n_{\alpha^{P_\nu}}(P_\nu, x))$  as a linear combination of the  $-\log(n_{\alpha^Q}(Q, x))$ , with  $\alpha \in \pi^{U_Q}$ . This means that the root  $\alpha \in \pi^{U_{P_\nu}}$  defines a half space  $H_\nu^+(\alpha)$  in  $\mathbb{R}^{d_Q}$  and  $N^Q(x) \subset H_\nu^+(\alpha)$  in  $\mathbb{R}^{d_Q}$ .

Now we assume that  $x$  is in the intersection (1.89). For the roots  $\alpha \in \pi \setminus \pi^{U_{P_\nu}}$  we have the condition (1.72). For the roots  $\alpha \in \pi^{U_Q} \setminus \pi^{U_{P_\nu}}$  this yields

$$-\log(n_{\gamma_\alpha^{M_\nu}}(P_\nu, x)) \leq -\log(r(\pi_\nu)).$$

Therefore we see that the image of  $N^Q$  is contained in the intersection of a finite number of half spaces, which are obtained from a finite family of hyperplanes. These hyperplanes depend on the parameters  $c_{\pi_\nu}, r(\pi_\nu)$ , let us call this intersection  $C(\underline{c}, \underline{r})$ , it is a convex -possibly empty- subset of  $\mathbb{R}^{d_Q}$ .

We investigate the restriction

$$N^Q : X^{P_1}(c_{\pi_1}, r(\underline{c}_{\pi_1})) \cap \cdots \cap X^{P_\nu}(c_{\pi_k}, r(\underline{c}_{\pi_k})) \rightarrow C(\underline{c}, \underline{r})$$

We observe that the unipotent radical  $U_Q(\mathbb{R})$  acts by left translations on the intersection, we get a diagram

$$\begin{array}{ccc} X^{P_1}(c_{\pi_1}, r(\underline{c}_{\pi_1})) \cap \cdots \cap X^{P_\nu}(c_{\pi_k}, r(\underline{c}_{\pi_k})) & \rightarrow & C(\underline{c}, \underline{r}) \\ \downarrow p_M & & \\ X^M \times \mathbb{R}^{d_Q} & \rightarrow & \mathbb{R}^{d_Q} \end{array} \quad (1.92)$$

Now it is clear from the definitions that the image of  $p_M$  is a set

$$\text{Im}(p_M) = \Omega^M(\underline{c}, \underline{r}) \times C(\underline{c}, \underline{r})$$

where  $\Omega^M(\underline{c}, \underline{r}) \subset X^M$  is a subset containing the set  $X^{M, st}$  of semi stable points and is described by certain inequalities as in (1.70). This subset is  $\Gamma_M$  invariant and  $\Gamma_M \backslash \Omega^M(\underline{c}, \underline{r})$  is relatively compact.

Hence we see that we have essentially the same situation as in (1.85). The map

$$q_M : X^{P_1}(c_{\pi_1}, r(\underline{c}_{\pi_1})) \cap \cdots \cap X^{P_\nu}(c_{\pi_k}, r(\underline{c}_{\pi_k})) \rightarrow \Gamma_M \backslash \Omega^M(\underline{c}, \underline{r}) \times C(\underline{c}, \underline{r}) \quad (1.93)$$

is a fiber bundle with fiber isomorphic to  $\Gamma_{U_Q} \backslash U_Q(\mathbb{R})$ .

In the following we refer to the book of S. Helgason [39].

We mention an important property of the sets  $X^P(c'_\pi, r(c_P))$ . We assume that our symmetric space  $X$  is connected, then it is well known that it is convex, any two points  $p, q \in X$  can be joined by a unique geodesic  $[p, q]$ . We say that a subset  $U \subset X$  is convex if for any two points  $p, q \in U$  also the geodesic  $[p, q] \subset U$ .

**Proposition 1.2.5.** *Let  $\Omega \subset \Omega^M(\underline{c}, \underline{r})$  be a convex subset. Then the inverse image  $p_M^{-1}(\Omega \times C(\underline{c}, \underline{r}))$  is a convex subset of  $X^{P_1}(c_{\pi_1}, r(\underline{c}_{\pi_1})) \cap \cdots \cap X^{P_\nu}(c_{\pi_k}, r(\underline{c}_{\pi_k}))$*

*Proof.* The assertion is easily reduced to the following:

Let  $P$  be a maximal parabolic subgroup, let  $M$  be its reductive quotient, let  $\alpha$  be the simple root not in  $\pi^M$  and  $\Omega \subset X^{M^{(1)}}$ . Then the set for any choice

of We choose a  $c_\alpha > 0$  and claim that  $X^P(c_\alpha, \Omega) = \{x \in X \mid n_{\alpha^P}(P, x) \leq c_\alpha ; q_M(x) \in \Omega\}$  is convex .

To see this we pick a point  $x \in X^P(c_\alpha, \Omega)$ , let  $T_x^X$  be the tangent space at  $x$ . The action of  $G(\mathbb{R})$  on  $X$  gives us a surjective map  $D_x : \mathfrak{g}_{\mathbb{R}} \rightarrow T_{x_0}^X$  and this induces an isomorphism  $D_x : \mathfrak{g}_{\mathbb{R}}/\mathfrak{k}_x \xrightarrow{\sim} T_x^X$ , here of course  $\mathfrak{k}_x$  is the Lie-algebra of  $K_x$ . We get the well known Cartan decomposition of the Lie-algebra

$$\mathfrak{g}_{\mathbb{R}} = \mathfrak{k}_x \oplus \mathfrak{p}_x \text{ where } \mathfrak{p}_x = \{V \in \mathfrak{g}_{\mathbb{R}} \mid \Theta_x(V) = -V\} \quad (1.94)$$

and we get the isomorphism  $D_x : \mathfrak{p}_x \xrightarrow{\sim} T_x^X$ . Starting from our parabolic subgroup  $P$  we get a finer decomposition of  $\mathfrak{p}_x$ .

Let  $\mathfrak{P}_{\mathbb{R}}$  be the the Lie algebra of  $P \times \mathbb{R}$ . The intersection  $P \times \mathbb{R} \cap \Theta_x(P \times \mathbb{R}) = M_x$  and we get for the Lie algebras  $\mathfrak{m}_x = \mathfrak{m}^{(0)} \oplus \mathfrak{a}$  and this gives the finer decoposition  $\mathfrak{m}_x = \mathfrak{k}_x^M \oplus \mathfrak{p}^{(M_x)} \oplus \mathfrak{a}$  and then

$$\mathfrak{p}_x = \mathfrak{p}^{(M_x)} \oplus \mathfrak{a} \oplus \{V - \Theta_x(V)\}_{V \in \mathfrak{u}} \quad (1.95)$$

where  $V \in \mathfrak{u}_{\mathbb{R}}$  and  $\mathfrak{a} = \mathbb{R}Y_A$ . We normalise  $Y_A$  such that  $d\alpha^P(Y_A) = 1$ . Then we can write a tangent vector  $T_x^X$  as image of

$$Y = Y_M + aY_A + (V - \theta(V));$$

We know that there is a unique geodesic  $c : \mathbb{R} \rightarrow X$  starting at  $x$  with  $c'(t) = Y$ . The theorem 3.3 in Chapter IV in [39] says that this geodesic is  $c(t) = \exp(tY) \cdot x$ . A tedious computation using the Iwasawa decomposition and the Campell-Hausdorff formula shows that

$$-\log(n_{\alpha^P}(\exp(tY) \cdot x)) = -\log(n_{\alpha^P}(x)) + at - a^2q(Y_A, V)t^2 \quad (1.96)$$

where  $q(Y_A, V)$  is a positive definite form in  $V$ .

If now  $x_1 \in X^P(c_\alpha, \Omega)$  is a second point, We find a tangent vector  $Y = Y_M + aY_A + (V - \theta(V))$  such that  $t \mapsto \exp(tY) \cdot x$  is the geodesic joining  $x$  and  $x_1 = \exp(Y) \cdot x$ . If we project these two points to  $X^{M^{(1)}}$  then the images  $\bar{x}, \bar{x}_1 \in \Omega$  and  $\exp(t(Y_M)\bar{x})$  is the geodesic in  $X^{M^{(1)}}$ . and hence for  $t \in [0, 1]$  we have  $\exp(t(Y_M)\bar{x})$ . But now

$$-\log(n_{\alpha^P}(x)) \geq -\log(c_\alpha); \quad -\log(n_{\alpha^P}(\exp(Y) \cdot x)) = -\log((n_{\alpha^P}(x_1)) \geq -\log(c_\alpha).$$

Since the second derivative is always  $> 0$  (see(1.96) it follows that  $-\log(n_{\alpha^P}(\exp(tY) \cdot x)) \geq -\log(c_\alpha) \forall t \in [0, 1]$ . □

We formulated the main theorems of reduction theory only for  $\text{Gl}_n/\mathbb{Q}$  because we did not want to much from the theory of reductive groups ( for instance [?] ). But actually these results extend to general reductive groups, basically in the same formulation. Especially we get

**Theorem 1.2.3.** (*Borel-Harish-Chandra*): *If  $G/\mathbb{Q}$  is an anisotropic reductive group and  $\Gamma \subset G(\mathbb{Q})$  is an arithmetic subgroup then*

$$\Gamma \backslash X = \Gamma \backslash G(\mathbb{R})/K_\infty$$

*is compact.*

## Chapter 2

# The Cohomology groups

### 2.1 Cohomology of arithmetic groups as cohomology of sheaves on $\Gamma \backslash X$ .

We are now in the position to unify — for the special case of arithmetic groups — the two cohomology theories from our chapter II and chapter IV in [28].

We start from a semi simple group  $G/\mathbb{Q}$  and we choose an arithmetic congruence subgroup  $\Gamma \subset G(\mathbb{Q})$ . Let  $X = G(\mathbb{R})/K$  as before. A second datum will be a  $\Gamma$ -module  $\mathcal{M}$ , in principle this can be any  $\Gamma$ -module.

Let  $\mathcal{M}$  is a  $\Gamma$ -module then we can attach a sheaf  $\tilde{\mathcal{M}}$  on  $\Gamma \backslash X$  to it, this sheaf has values in the category of abelian groups. To do this we have to define for any open subset  $U \subset X$  the group of sections  $\tilde{\mathcal{M}}(U)$ . We start from the projection

$$\pi : X \longrightarrow \Gamma \backslash X$$

and define sheaf

$$\tilde{\mathcal{M}}(U) = \{f : \pi^{-1}(U) \rightarrow \mathcal{M} \mid f \text{ is locally constant } f(\gamma u) = \gamma f(u)\}. \quad (2.1)$$

This is clearly a sheaf. For any point  $x \in \Gamma \backslash X$  we can find a neighborhood  $V_x$  with the following property: We choose a point  $\tilde{x} \in \pi^{-1}(x)$ , then  $\tilde{x}$  has a convex  $\Gamma_{\tilde{x}}$ -invariant neighbourhood  $U_{\tilde{x}}$ , for which  $\gamma U_{\tilde{x}} \cap U_{\tilde{x}} \neq \emptyset \iff \gamma \notin \Gamma_{\tilde{x}}$  and then we put  $V_x = \Gamma_{\tilde{x}} \backslash U_{\tilde{x}}$ . We call such a neighbourhood  $V_x$  an *orbiconvex* neighbourhood. It is clear that

$$\tilde{\mathcal{M}}(V_x) = \mathcal{M}^{\Gamma_{\tilde{x}}}.$$

Since  $x$  has a cofinal system of neighbourhoods of this kind, we see that we get an isomorphism

$$j_{\tilde{x}} : \tilde{\mathcal{M}}(V_x) = \tilde{\mathcal{M}}_x \xrightarrow{\sim} \mathcal{M}^{\Gamma_{\tilde{x}}}.$$

The last isomorphism depends on the choice of  $\tilde{x}$ . If we are in the special case that  $\Gamma$  has no fixed points then we can cover  $\Gamma \backslash X$  by open sets  $U$  so that  $\tilde{\mathcal{M}}/U$  is isomorphic to a constant sheaf  $\underline{\mathcal{M}}_U$ . These sheaves are called *local systems*. If we have fixed points we call them *orbilocal systems*.

We will denote the functor, which sends  $\mathcal{M}$  to  $\tilde{\mathcal{M}}$  by

$$\mathrm{sh}_\Gamma : \mathbf{Mod}_\Gamma \rightarrow \mathcal{S}_{\Gamma \backslash X},$$

occasionally we will write  $\mathrm{sh}_\Gamma(\mathcal{M})$  instead of  $\tilde{\mathcal{M}}$ , especially in situations where we work with several discrete subgroups.

For the following we refer to [28] Chapter 2

The motivations for these constructions are

1) The spaces  $\Gamma \backslash X$  are interesting examples of so-called locally symmetric spaces (provided  $\Gamma$  has no torsion). Hence they are of interest for differential geometers and analysts.

2) If we have some understanding of the geometry of the quotient space  $\Gamma \backslash X$  we gain some insight into the structure of  $\Gamma$ . This will become clear when we discuss the examples in ...x.y.z.

3) The cohomology groups  $H^\bullet(\Gamma, \mathcal{M})$  are closely related and in many cases even isomorphic to the sheaf cohomology groups  $H^\bullet(\Gamma \backslash X, \tilde{\mathcal{M}})$ . Again the geometry provides tools to compute these cohomology groups in some cases (see x.y.z.).

4) If the  $\Gamma$ -module  $\mathcal{M}$  is a  $\mathbb{C}$ -vector space and obtained from a rational representation of  $G/\mathbb{Q}$ , then we can apply analytic tools to get insight (de Rham cohomology, Hodge theory).

### 2.1.1 The relation between $H^\bullet(\Gamma, \mathcal{M})$ and $H^\bullet(\Gamma \backslash X, \mathcal{M})$

In this section we assume that  $X$  is connected.

The functor

$$\mathcal{M} \rightarrow H^0(\Gamma \backslash X, \tilde{\mathcal{M}}) = \mathcal{M}^\Gamma.$$

is a functor from the category of  $\Gamma$ -modules to the category  $\mathbf{Ab}$  of abelian groups. We can write our functor  $\mathcal{M} \rightarrow \mathcal{M}^\Gamma$  as a composition of

$$\mathrm{sh}_\Gamma : \mathcal{M} \longrightarrow \tilde{\mathcal{M}} \text{ and } H^0 : \tilde{\mathcal{M}} \rightarrow H^0(\Gamma \backslash X, \tilde{\mathcal{M}}).$$

We want to apply the composition rule from [28] 4.6.4.

*In a first step we have to convince ourselves that  $\mathrm{sh}_\Gamma$  sends injective  $\Gamma$ -modules to acyclic sheaves.*

In [28], 2.2.4. we constructed the induced  $\Gamma$ -module  $\mathrm{Ind}_{\{1\}}^\Gamma \mathcal{M}$ , for any  $\Gamma$ -module  $\mathcal{M}$ . This is the module of all functions  $f : \Gamma \rightarrow \mathcal{M}$  and  $\gamma_1 \in \Gamma$  acts on this module by  $(\gamma_1 f)(\gamma) = f(\gamma \gamma_1)$ . The map

$$m \mapsto f_m = \{\gamma \mapsto \gamma m\} \tag{2.2}$$

is an injective  $\Gamma$ -module homomorphism.

In a first step we prove that for any such induced module the sheaf  $\mathrm{sh}_\Gamma(\mathrm{Ind}_{\{1\}}^\Gamma \mathcal{M})$  is acyclic.

We have a little

2.1. COHOMOLOGY OF ARITHMETIC GROUPS AS COHOMOLOGY OF SHEAVES ON  $\Gamma \backslash X$ .65

**Lemma 2.1.1.** *Let us consider the projection  $\pi : X \rightarrow \Gamma \backslash X$  and the constant sheaf  $\underline{\mathcal{M}}_X$  on  $X$ . Then we have a canonical isomorphism of sheaves*

$$\pi_*(\underline{\mathcal{M}}_X) \xrightarrow{\sim} \widetilde{\text{Ind}}_{\{1\}}^{\Gamma} \mathcal{M}.$$

*Proof.* This is rather obvious. Let us consider a small neighborhood  $U_x$  of a point  $x$ , such that  $\pi^{-1}(U_x)$  is the disjoint union of small contractible neighborhoods  $U_{\tilde{x}}$  for  $\tilde{x} \in \pi^{-1}(x)$ . Then for all points  $\tilde{x}$  we have  $\underline{\mathcal{M}}_X(U_{\tilde{x}}) = \mathcal{M}$  and

$$\pi_*(\underline{\mathcal{M}}_X)(U_x) = \prod_{\tilde{x} \in \pi^{-1}(x)} \mathcal{M}.$$

On the other hand

$$\widetilde{\text{Ind}}_{\{1\}}^{\Gamma} \mathcal{M}(U_x) = \left\{ h : \pi^{-1}(U_x) \rightarrow \text{Ind}_{\{1\}}^{\Gamma} \mathcal{M} \mid h \text{ is locally constant } h(\gamma u) = \gamma h(u) \right\}$$

For  $u \in \pi^{-1}(U_x)$  the element  $h(u)$  itself is a map

$$h(u) : \Gamma \longrightarrow \mathcal{M},$$

and  $(\gamma h(u))(\gamma_1) = h(u)(\gamma_1 \gamma)$  (here  $\gamma_1 \in \Gamma$  is the variable.)

Hence we know the function  $u \rightarrow h(u)$  from  $\pi^{-1}(U_x)$  to  $\text{Ind}_{\{1\}}^{\Gamma} \mathcal{M}$  if we know its value  $h(u)(1)$  and this value can be prescribed on the connected components of  $\pi^{-1}(U_x)$ . On these connected components it is constant, we may take its value at  $\tilde{x}$  and hence

$$h \longrightarrow (\dots, h(\tilde{x})(1), \dots)_{\tilde{x} \in \pi^{-1}(x)}$$

yields the desired isomorphism.

Now acyclicity is clear.. We apply example d) in [28], 4.6.3 to this situation. The fibre of  $\pi$  is a discrete space and hence

$$\pi_*(\underline{\mathcal{M}}_X) = \widetilde{\text{Ind}}_{\{1\}}^{\Gamma} \mathcal{M}$$

and  $R^q(\pi_*)(\underline{\mathcal{M}}_X) = 0$  for  $q > 0$ . Therefore the spectral sequence yields

$$H^q(X, \underline{\mathcal{M}}_X) = H^q(\Gamma \backslash X, \pi_*(\underline{\mathcal{M}}_X)) = H^q\left(\Gamma \backslash X, \widetilde{\text{Ind}}_{\{1\}}^{\Gamma} \mathcal{M}\right),$$

and since  $X$  is a cell, we see that this is zero for  $q \geq 1$ . □

We apply this to the case that  $\mathcal{M} = \mathcal{I}$  is an injective  $\Gamma$ -module. Clearly we can always embed  $\mathcal{I} \rightarrow \text{Ind}_{\{1\}}^{\Gamma} \mathcal{I}$ . But this is now a direct summand; hence it follows from the acyclicity of  $\widetilde{\text{Ind}}_{\{1\}}^{\Gamma} \mathcal{I}$  that also  $\tilde{\mathcal{I}}$  must be acyclic.

Hence we can apply the composition rule and get spectral sequence with  $E_2$  term

$$H^p(\Gamma \backslash X, R^q(\text{sh}_{\Gamma})(\mathcal{M})) \Rightarrow H^n(\Gamma, \mathcal{M}).$$

The edge homomorphism yields a homomorphism

$$H^n(\Gamma \backslash X, \text{sh}_{\Gamma}(\mathcal{M})) \rightarrow H^n(\Gamma, \mathcal{M}) \tag{2.3}$$

which in general is neither injective nor surjective.

We have seen in section (1.2.2) that -under our assumption that  $G/\mathbb{Q}$  is semisimple- the stabilisers  $\Gamma_x$  are finite. This implies that the stalks  $R^q(\mathrm{sh}_\Gamma)(\mathcal{M})_x = H^q(\Gamma_{\bar{x}}, \mathcal{M})$  for  $q > 0$  are torsion groups actually they are annihilated by  $\#\Gamma_x$ . This implies that the edge homomorphism has finite kernel and cokernel.

In this book we are mainly interested in the cohomology groups  $H^n(\Gamma \backslash X, \mathrm{sh}_\Gamma(\mathcal{M}))$  and not so much in the group cohomology  $H^\bullet(\Gamma, \mathcal{M})$ .

### Functorial properties of cohomology

We investigate the functorial properties of the cohomology with respect to the change of  $\Gamma$ . If  $\Gamma' \subset \Gamma$  is a subgroup of finite index, then we have, of course, the functor

$$\mathbf{Mod}_\Gamma \longrightarrow \mathbf{Mod}_{\Gamma'},$$

which is obtained by restricting the  $\Gamma$ -module structure to  $\Gamma'$ . Since for any  $\Gamma$ -module  $\mathcal{M}$  we have  $\mathcal{M}^\Gamma \longrightarrow \mathcal{M}^{\Gamma'}$ , we obtain a homomorphism

$$\mathrm{res} : H^i(\Gamma, \mathcal{M}) \longrightarrow H^i(\Gamma', \mathcal{M}).$$

We give an interpretation of this homomorphism in terms of sheaf cohomology. We have the diagram

$$\begin{array}{ccc} & X & \\ \pi_{\Gamma'} \swarrow & & \searrow \pi_\Gamma \\ \pi_1 = \pi_{\Gamma, \Gamma'} : \Gamma' \backslash X & \longrightarrow & \Gamma \backslash X \end{array}$$

and a  $\Gamma$ -module  $\mathcal{M}$  produces sheaves  $\mathrm{sh}_\Gamma(\mathcal{M}) = \tilde{\mathcal{M}}$  and  $\mathrm{sh}_{\Gamma'}(\mathcal{M}) \cong \tilde{\mathcal{M}}'$  on  $\Gamma' \backslash X$  and  $\Gamma \backslash X$  respectively. It is clear that we have a homomorphism

$$\pi_1^*(\tilde{\mathcal{M}}) \longrightarrow \tilde{\mathcal{M}}'.$$

To get this homomorphism we observe that for  $y_1 \in \Gamma' \backslash X$  we have  $\pi_1^*(\tilde{\mathcal{M}})_{y_1} = \tilde{\mathcal{M}}_{\pi_1(y_1)}$ , and this is

$$\{f : \pi^{-1}(\pi_1(y)) \rightarrow \mathcal{M} \mid f(\gamma\tilde{y}) = \gamma f(\tilde{y}) \text{ for all } \gamma \in \Gamma, \tilde{y} \in \pi^{-1}(\pi(y))\}$$

and

$$\tilde{\mathcal{M}}'_{y_1} = \{fg : (\pi')^{-1}(y_1) \rightarrow \mathcal{M} \mid f(\gamma'\tilde{y}) = \gamma' f(\tilde{y}) \text{ for all } \gamma' \in \Gamma', \tilde{y} \in (\pi')^{-1}(y_1)\},$$

and if we pick a point  $\tilde{y} \in (\pi')^{-1}(y_1) \subset \pi^{-1}(\pi_1(y_1))$  then

$$\pi_1^*(\mathcal{M})_{y_1} \simeq \mathcal{M}^{\Gamma_{\tilde{y}_1}} \subset \tilde{\mathcal{M}}'_{y_1} = \mathcal{M}^{\Gamma'_{\tilde{y}_1}}.$$

Hence we get (or define) our restriction homomorphism as (see I, ....)

$$H^i(\Gamma \backslash X, \mathrm{sh}_\Gamma(\mathcal{M})) \longrightarrow H^i(\Gamma' \backslash X, \pi_1^*(\mathrm{sh}_\Gamma(\mathcal{M}))) \longrightarrow H^i(\Gamma' \backslash X, \mathrm{sh}_{\Gamma'}(\mathcal{M})).$$

There is also a map in the opposite direction.

## 2.1. COHOMOLOGY OF ARITHMETIC GROUPS AS COHOMOLOGY OF SHEAVES ON $\Gamma \backslash X$ .67

Since the fibres of  $\pi_1$  are discrete we have

$$H^i(\Gamma' \backslash X, \tilde{\mathcal{M}}) \xrightarrow{\sim} H^i(\Gamma \backslash X, \pi_{1,*}(\tilde{\mathcal{M}})).$$

But the same reasoning as in the previous section yields an isomorphism

$$\pi_{1,*}(\tilde{\mathcal{M}}) \xrightarrow{\sim} \widetilde{\text{Ind}}_{\Gamma'}^{\Gamma} \mathcal{M}.$$

Hence we get an isomorphism

$$H^i(\Gamma' \backslash X, \tilde{\mathcal{M}}) \xrightarrow{\sim} H^i(\Gamma \backslash X, \widetilde{\text{Ind}}_{\Gamma'}^{\Gamma} \mathcal{M})$$

which is well known as Shapiro's lemma. But we have a  $\Gamma$ -module homomorphism

$$e : \text{Ind}_{\Gamma'}^{\Gamma} \mathcal{M} \longrightarrow \mathcal{M}$$

which sends an  $f : \Gamma \rightarrow \mathcal{M}$ , in  $f \in \text{Ind}_{\Gamma'}^{\Gamma} \mathcal{M}$  to the sum

$$\text{tr}(f) = \sum \gamma_i^{-1} f(\gamma_i)$$

where the  $\gamma_i$  are representatives for the classes of  $\Gamma' \backslash \Gamma$ . This homomorphism induces a map in the cohomology. We get a composition

$$\pi_{1,\bullet} : H^i(\Gamma' \backslash X, \tilde{\mathcal{M}}) \longrightarrow H^i(\Gamma \backslash X, \tilde{\mathcal{M}}).$$

It is not difficult to check that

$$\pi_{1,\bullet} \circ \pi_1^{\bullet} = [\Gamma : \Gamma'].$$

### 2.1.2 How to compute the cohomology groups $H^i(\Gamma \backslash X, \tilde{\mathcal{M}})$ ?

#### The Čech complex of an orbiconvex Covering

We consider a point  $\tilde{x} \in X$  and an open neighbourhood  $\tilde{U}_{\tilde{x}} \subset X$ . We say that  $\tilde{U}_{\tilde{x}}$  is an *orbiconvex* neighborhood of  $\tilde{x}$  if

a) The set  $\tilde{U}_{\tilde{x}}$  is convex, i.e. for any two points in  $\tilde{x}_1, \tilde{x}_2 \in \tilde{U}_{\tilde{x}}$  the geodesic joining  $\tilde{x}_1$  and  $\tilde{x}_2$  lies in  $\tilde{U}_{\tilde{x}}$ .

**irgendwo früher was zu Geodäten sagen, )**

b) We have  $\gamma \tilde{U}_{\tilde{x}} \cap \tilde{U}_{\tilde{x}} = \emptyset$  unless  $\gamma \tilde{x} = \tilde{x}$  and in this case we even have  $\gamma \tilde{U}_{\tilde{x}} = \tilde{U}_{\tilde{x}}$ .

A family of orbiconvex neighborhoods  $\{\tilde{U}_{\tilde{x}_i}\}_{i=1,\dots,r}$  of points  $\tilde{x}_1, \dots, \tilde{x}_r$  will be called an *orbiconvex covering*, if

$$\bigcup_{i=1}^r \bigcup_{\gamma \in \Gamma} \gamma \tilde{U}_{\tilde{x}_i} = X. \quad (2.4)$$

We will show later that we can always find a finite orbiconvex covering of  $X$ .

If now  $\{\tilde{U}_{\tilde{x}_i}\}_{i=1,\dots,r}$  is an orbiconvex covering we put  $U_{x_i} = \pi(\tilde{U}_{\tilde{x}_i})$ , and then we get finite covering by open sets

$$\bigcup_{x_i} U_{x_i} = \Gamma \backslash X$$

We call  $\mathfrak{U} = \{U_{x_i}\}$  an orbiconvex covering of  $\Gamma \backslash X$ .

We will see further down that the intersections  $U_{\underline{i}} = U_{x_{i_1}} \cap U_{x_{i_2}} \cap \cdots \cap U_{x_{i_q}}$  are acyclic, i.e.  $H^k(U_{\underline{i}}, \tilde{\mathcal{M}}) = 0$  for  $k > 0$ .

This implies that the Čzech complex (See [28], Chap. 4)

$$C^\bullet(\mathfrak{U}, \tilde{\mathcal{M}}) := 0 \rightarrow \bigoplus_{i \in I} \tilde{\mathcal{M}}(U_{x_i}) \xrightarrow{d_0} \bigoplus_{i < j} \tilde{\mathcal{M}}(U_{x_i} \cap U_{x_j}) \rightarrow \quad (2.5)$$

computes the cohomology.

For the implementation on a computer we need to resolve the definition of the spaces of sections and the definition of the boundary maps. (By this I mean that we have to write explicitly

$$\tilde{\mathcal{M}}(U_{\underline{i}}) = \bigoplus_{\eta} \mathcal{M}_{\eta}$$

where  $\eta$  runs through an index set and  $\mathcal{M}_{\eta}$  are explicit subspaces of  $\mathcal{M}$  and then we have to write down certain explicit linear maps  $\mathcal{M}_{\eta} \rightarrow \mathcal{M}_{\eta'}$ .)

To be more precise: We have to write  $U_{\underline{i}} = \cup U_{\eta}$  as the union of its connected components, we have to choose a connected component  $\tilde{U}_{\eta}$  in  $\pi^{-1}(U_{\eta})$  for each value of  $\eta$ , and then the evaluation of a section  $m \in \tilde{\mathcal{M}}(U_{\underline{i}})$  on these  $\tilde{U}_{\eta}$  yields an isomorphism

$$\oplus ev_{\tilde{U}_{\eta}} : \tilde{\mathcal{M}}(U_{\underline{i}}) \xrightarrow{\sim} \bigoplus_{\eta} \mathcal{M}^{\Gamma_{\eta}}.$$

If we replace  $\tilde{U}_{\eta}$  by  $\gamma \tilde{U}_{\eta}$  then we get for  $m \in \tilde{\mathcal{M}}(\pi(\tilde{U}_{\eta}))$  the equality

$$\gamma ev_{\tilde{U}_{\eta}}(m) = ev_{\gamma \tilde{U}_{\eta}} \quad (2.6)$$

Especially the choice of the  $\tilde{x}_i$  yields an identification

$$ev_{U_{x_i}} : \tilde{\mathcal{M}}(U_{x_i}) \xrightarrow{\sim} \mathcal{M}^{\Gamma_{\tilde{x}_i}} \quad (2.7)$$

this gives us the first term in the complex.

The computation the second term is a little bit more delicate. The point is that the intersections  $U_{x_i} \cap U_{x_j}$  may not be connected. To get these connected components we have to find the elements  $\gamma \in \Gamma$  for which

$$\tilde{U}_{\tilde{x}_i} \cap \gamma(\tilde{U}_{\tilde{x}_j}) \neq \emptyset \quad (2.8)$$

It is clear that this gives us a finite set  $G_{i,j}$  of elements  $\gamma \in \Gamma/\Gamma_{x_j}$ . We have a little lemma

**Lemma 2.1.2.** *The images  $\pi(\tilde{U}_{\tilde{x}_i} \cap \gamma(\tilde{U}_{\tilde{x}_j}))$  are the connected components of  $U_{x_i} \cap U_{x_j}$ , two elements  $\gamma, \gamma_1$  give the same connected component if and only if  $\gamma_1 \in \Gamma_{x_i} \gamma \Gamma_{x_j}$ .*

Let  $F_{i,j} \subset G_{i,j}$  be a set of representatives for the action of  $\Gamma_{x_i}$  on  $G_{i,j}$  this set can be identified to the set of connected components. Of course the set  $\tilde{U}_{\tilde{x}_i} \cap \gamma(\tilde{U}_{\tilde{x}_j})$  may have a non trivial stabilizer  $\Gamma_{i,j,\gamma}$  and then we get an identification

$$\oplus_{\gamma \in F_{i,j}} ev_{\tilde{U}_{x_i} \cap \gamma \tilde{U}_{x_j}} : \tilde{\mathcal{M}}(U_{x_i} \cap U_{x_j}) \xrightarrow{\sim} \bigoplus_{\gamma \in F_{i,j}} \mathcal{M}^{\Gamma_{i,j,\gamma}} \quad (2.9)$$

This is now an explicit (i.e. digestible for a computer) description of the second term in our complex above. We still need to give the explicit formula for  $d_0$  in the complex

$$0 \rightarrow \bigoplus_{i \in I} \mathcal{M}^{\Gamma_{\tilde{x}_i}} \xrightarrow{d_0} \bigoplus_{i < j} \bigoplus_{\gamma \in F_{i,j}} \mathcal{M}^{\Gamma_{i,j,\gamma}} \quad (2.10)$$

Looking at the definition it is clear that this map is given by

$$(\dots, m_i, \dots, m_j, \dots) \mapsto (\dots, m_i - \gamma m_j, \dots) \quad (2.11)$$

Here we have to observe that  $\gamma \in \Gamma/\Gamma_{x_j}$  but this does not matter since  $m_j \in \mathcal{M}^{\Gamma_{\tilde{x}_j}}$ . So we have an explicit description of the beginning of the Čech complex.

A little reasoning shows of course that a different choice  $F'_{i,j}$  of the representatives provides an isomorphic complex.

Now it is clear, how to proceed. At first we have to understand the combinatorics of the covering  $\mathfrak{U} = \{U_{x_i}\}_{i \in I}$ .

We consider sets

$$G_{\underline{i}} = \{\underline{\gamma} = (e, \gamma_1, \dots, \gamma_q) \mid \gamma_i \in \Gamma/\Gamma_{x_i}; \tilde{U}_{\tilde{x}_0} \cap \dots \cap \gamma_i \tilde{U}_{\tilde{x}_i} \cap \gamma_q \tilde{U}_{\tilde{x}_q} \neq \emptyset\}$$

on these sets we have an action of  $\Gamma_{x_0}$  by multiplication from the left. Again let  $F_{\underline{i}}$  be a system of representatives modulo the action of  $\Gamma_{x_0}$ .

We abbreviate

$$\tilde{U}_{\underline{i}, \underline{\gamma}} = \tilde{U}_{\tilde{x}_0} \cap \dots \cap \gamma_i \tilde{U}_{\tilde{x}_i} \cap \gamma_q \tilde{U}_{\tilde{x}_q},$$

let  $\Gamma_{\underline{i}, \underline{\gamma}}$  be the stabilizer of  $\tilde{U}_{\underline{i}, \underline{\gamma}}$ .

The images  $\pi(\tilde{U}_{\underline{i}, \underline{\gamma}})$  under the projection map  $\pi$  are the connected components  $\pi(\tilde{U}_{\underline{i}, \underline{\gamma}}) = U_{\underline{i}, \underline{\gamma}} \subset U_{\underline{i}} = U_{x_{i_0}} \cap \dots \cap U_{x_{i_\nu}} \cap \dots \cap U_{x_{i_q}}$ . On the other hand each set  $\tilde{U}_{\underline{i}, \underline{\gamma}}$  is a connected component in  $\pi^{-1}(U_{\underline{i}, \underline{\gamma}})$ . We get an isomorphism

$$\bigoplus_{\underline{\gamma} \in F_{\underline{i}}} ev_{\tilde{U}_{\underline{i}, \underline{\gamma}}} : \tilde{\mathcal{M}}(U_{\underline{i}}) = \tilde{\mathcal{M}}(U_{x_{i_0}} \cap \dots \cap U_{x_{i_\nu}} \cap \dots \cap U_{x_{i_q}}) \xrightarrow{\sim} \bigoplus_{\underline{\gamma} \in F_{\underline{i}}} \mathcal{M}^{\Gamma_{\underline{i}, \underline{\gamma}}}. \quad (2.12)$$

We need to give explicit formulas for the boundary maps

$$\bigoplus_{\underline{i} \in I^q} \tilde{\mathcal{M}}(U_{\underline{i}}) \xrightarrow{d_q} \bigoplus_{\underline{i} \in I^{q+1}} \tilde{\mathcal{M}}(U_{\underline{i}}).$$

Abstractly this boundary operator is defined as follows: We look at pairs  $\underline{i} \in I^{q+1}, \underline{i}^{(\nu)} \in I^q$  where  $\underline{i}^{(\nu)}$  is obtained from  $\underline{i}$  by deleting the  $\nu$ -th entry. Then we have  $U_{\underline{i}} \subset U_{\underline{i}^{(\nu)}}$  and from this we get the resulting restriction homomorphism  $R_{\underline{i}^{(\nu)}, \underline{i}} : \tilde{\mathcal{M}}(U_{\underline{i}^{(\nu)}}) \rightarrow \tilde{\mathcal{M}}(U_{\underline{i}})$ . Then

$$d_q = \sum_{\underline{i}} \sum_{\nu=0}^q (-1)^\nu R_{\underline{i}^{(\nu)}, \underline{i}}$$

and hence we have to give an explicit description of  $R_{\underline{i}^{(\nu)}, \underline{i}}$  with respect to the isomorphism in the diagram (2.12).

We pick two connected components  $\pi(\tilde{U}_{\underline{i},\underline{\gamma}}) \subset U_{\underline{i}}$  and  $\pi(\tilde{U}_{\underline{i}(\nu),\underline{\gamma}'}) \subset U_{\underline{i}(\nu)}$ , then we know that

$$\tilde{U}_{\underline{i},\underline{\gamma}} \subset \tilde{U}_{\underline{i}(\nu),\underline{\gamma}'} \iff \exists \eta_{\gamma,\gamma'} \in \Gamma \text{ such that } \eta_{\gamma,\gamma'} \gamma'_\mu = \gamma_\mu \text{ for all } \mu \neq \nu$$

and then the restriction of  $R_{\underline{i}(\nu),\underline{i}}$  to these two components is given by

$$\begin{array}{ccc} \tilde{\mathcal{M}}(\pi(\tilde{U}_{\underline{i}(\nu),\underline{\gamma}'})) & \xrightarrow{\text{ev}_{\tilde{U}_{\underline{i}(\nu),\underline{\gamma}'}}} & \mathcal{M}^{\Gamma_{\underline{i}(\nu),\underline{\gamma}'}} \\ \downarrow R_{\underline{i}(\nu),\underline{i}} & & \downarrow \eta_{\gamma,\gamma'} \\ \tilde{\mathcal{M}}(\pi(\tilde{U}_{\underline{i},\underline{\gamma}})) & \xrightarrow{\text{ev}_{\tilde{U}_{\underline{i},\underline{\gamma}}}} & \mathcal{M}^{\Gamma_{\underline{i},\underline{\gamma}}} \end{array} \quad (2.13)$$

Here the two horizontal maps are isomorphisms, we observe that  $\eta_{\gamma,\gamma'}$  is unique up to an element in  $\Gamma_{\underline{i}(\nu),\underline{\gamma}'}$  and hence the vertical arrow  $\eta_{\gamma,\gamma'}$  is well defined.

Hence we conclude:

*Once we have found a finite orbiconvex covering of  $\Gamma \backslash X$ , we can write down an explicit complex, which computes the cohomology groups  $H^\bullet(\Gamma \backslash X, \tilde{\mathcal{M}})$ .*

We may also look at this situation from a different point of view: If  $x \in X$  is any point and  $\Gamma_x \subset \Gamma$  its stabilizer, then we define the induced  $\Gamma$  module

$$\text{Ind}_{\Gamma_x}^\Gamma \mathbb{Z} := \{f : \Gamma \rightarrow \mathbb{Z} \mid f \text{ has finite support and } f(a\gamma) = f(\gamma), \forall a \in \Gamma_x, \gamma \in \Gamma\} \quad (2.14)$$

If  $V_x$  is an open neighbourhood of  $x$  which satisfies b) an c) then we have  $\pi^{-1}(\pi(V_x)) = \bigcup_{\gamma \in \Gamma/\Gamma_x} \gamma V_x$  and

$$\pi^*(\tilde{\mathcal{M}})\left(\bigcup_{\gamma \in \Gamma/\Gamma_x} \gamma V_x\right) = \text{Hom}(\text{Ind}_{\Gamma_x}^\Gamma \mathbb{Z}, \mathcal{M}).$$

We have the covering

$$\tilde{\mathcal{U}} = \bigcup_{i,\gamma \in \Gamma/\Gamma_{\tilde{x}_i}} \gamma \tilde{U}_{\tilde{x}_i} = X$$

of the symmetric space. The Čzech-complex  $C^\bullet(\tilde{\mathcal{U}}, \pi^*(\tilde{\mathcal{M}}))$  computes the cohomology groups  $H^q(X, \pi^*(\tilde{\mathcal{M}}))$  which are trivial for  $q > 0$ . Our considerations above yield

$$C^\bullet(\tilde{\mathcal{U}}, \pi^*(\tilde{\mathcal{M}})) = 0 \rightarrow \bigoplus_{i=1}^r \text{Hom}(\text{Ind}_{\Gamma_{x_i}}^\Gamma \mathbb{Z}, \mathcal{M}) \xrightarrow{d^1} \bigoplus_{i < j, \tilde{x}_{i,j}} \text{Hom}(\text{Ind}_{\Gamma_{\tilde{x}_{i,j}}}^\Gamma \mathbb{Z}, \mathcal{M}) \xrightarrow{d^2} \dots$$

Now it is easy to see that the boundary maps are induced by maps between the induced modules

$$\xrightarrow{\delta^2} \bigoplus_{i < j, \tilde{x}_{i,j}} \text{Ind}_{\Gamma_{\tilde{x}_{i,j}}}^\Gamma \mathbb{Z} \xrightarrow{\delta^1} \bigoplus_{i=1}^r \text{Ind}_{\Gamma_{\tilde{x}_i}}^\Gamma \mathbb{Z} \rightarrow 0,$$

2.1. COHOMOLOGY OF ARITHMETIC GROUPS AS COHOMOLOGY OF SHEAVES ON  $\Gamma \backslash X$ .71

where for  $f \in \bigoplus \text{Ind}_{\tilde{x}_j}^{\Gamma} \mathbb{Z}$ , in degree  $\nu$  and  $\omega \in C^{\nu-1}(\tilde{\mathfrak{U}}, \pi^*(\tilde{\mathcal{M}}))$  the relation  $\omega(\delta^\nu(f)) = d^{\nu-1}(\omega)(f)$  defines  $\delta^\nu$ . We get an augmented complex

$$P^\bullet := \rightarrow \bigoplus_{\tilde{x}_j} \text{Ind}_{\tilde{x}_j}^{\Gamma} \mathbb{Z} \rightarrow \cdots \rightarrow \bigoplus_{\tilde{x}_{i,j}} \text{Ind}_{\tilde{x}_{i,j}}^{\Gamma} \mathbb{Z} \rightarrow \bigoplus_{i=1}^r \text{Ind}_{\tilde{x}_i}^{\Gamma} \mathbb{Z} \rightarrow \mathbb{Z} \rightarrow 0 \quad (2.15)$$

and since  $C^\bullet(\tilde{\mathfrak{U}}, \pi^*(\tilde{\mathcal{M}}))$  is acyclic in degree  $> 0$ , we get that  $P^\bullet$  is an acyclic resolution of the trivial module  $\mathbb{Z}$ .

Let  $N = \prod_i \#\Gamma_{\tilde{x}_i}$  and  $R := \mathbb{Z}[\frac{1}{N}]$  then the  $R[\Gamma]$  module  $\text{Ind}_{\tilde{x}_i}^{\Gamma} \otimes R$  is a direct summand in  $R[\Gamma]$  and hence a projective  $R[\Gamma]$  module. This implies of course that

$$P^\bullet \otimes R \Rightarrow \bigoplus_{\tilde{x}_j} \text{Ind}_{\tilde{x}_j}^{\Gamma} R \rightarrow \cdots \rightarrow \bigoplus_{\tilde{x}_{i,j}} \text{Ind}_{\tilde{x}_{i,j}}^{\Gamma} R \rightarrow \bigoplus_{i=1}^r \text{Ind}_{\tilde{x}_i}^{\Gamma} R \rightarrow R \rightarrow 0 \quad (2.16)$$

is indeed a projective resolution of the trivial  $\Gamma$ -module  $R$ . Therefore we know that

$$H^\bullet(\Gamma, \mathcal{M}_R) = H^\bullet(0 \rightarrow \text{Hom}_{\Gamma}(\bigoplus_{i=1}^r \text{Ind}_{\tilde{x}_i}^{\Gamma} R, \mathcal{M}_R) \rightarrow \bigoplus_{i < j, \tilde{x}_{i,j}} \text{Hom}_{\Gamma}(\text{Ind}_{\tilde{x}_{i,j}}^{\Gamma} R, \mathcal{M}_R) \rightarrow \cdots) \quad (2.17)$$

where now on the left hand side we have the group cohomology.

If we do not tensor by  $R$  then the Čech-complex

$$0 \rightarrow \bigoplus_{i=1}^r \text{Hom}_{\Gamma}(\text{Ind}_{\tilde{x}_i}^{\Gamma} \mathbb{Z}, \mathcal{M}) \rightarrow \bigoplus_{i < j, \tilde{x}_{i,j}} \text{Hom}_{\Gamma}(\text{Ind}_{\tilde{x}_{i,j}}^{\Gamma} \mathbb{Z}, \mathcal{M}) \rightarrow \cdots \quad (2.18)$$

is isomorphic to the Čech complex (2.5) and it computes the sheaf cohomology  $H^\bullet(\Gamma \backslash X, \mathcal{M})$ .

It follows from reduction theory that

**Theorem 2.1.1.** *We can construct a finite covering  $\Gamma \backslash X = \bigcup_{i \in E} U_{x_i} = \mathfrak{U}$  by orbiconvex sets.*

*Proof.* This is rather clear. We start from the covering by the sets  $X^P(c'_\pi, r(c'_\pi))$ . The set of "almost stable" points  $X^G(r) \subset X$  is relatively compact modulo  $\Gamma$ . For any point  $\tilde{x} \in X$  we look at the minimum distance

$$d(\tilde{x}) := \min_{\gamma \in \Gamma \backslash \Gamma_{\tilde{x}}} d(\tilde{x}, \gamma \tilde{x}).$$

since the action of  $\Gamma$  is properly discontinuous this minimum distance  $d(\tilde{x}) > 0$ . Let  $D(\tilde{x}, d(\tilde{x})/2) := \{\tilde{y} \mid d(\tilde{y}, \tilde{x}) < d(\tilde{x})/2\}$ , (-the Dirichlet-ball around  $\tilde{x}$ -) then

$D(\tilde{x}, d(\tilde{x})/2)$  is an orbiconvex neighborhood of  $\tilde{x}$ . Then we can find finitely many points  $\tilde{x}_1, \dots, \tilde{x}_r$  such that

$$\bigcup_{i=1}^r \bigcup_{\gamma \in \Gamma} \gamma D(\tilde{x}_i, d(\tilde{x}_i)/2) \supset X^G(r).$$

We have to find a covering for the  $X^P(c_P, r(c_P))$ . We recall the fibration (See (1.84))

$$p_{P,M}^* : X^P(c_{\pi'}, r(c_{\pi'})) \rightarrow X^M(r(c_{\pi'})) \times \prod_{\alpha \in \pi'} (0, c_\alpha].$$

We apply our previous argument and find a finite covering

$$\bigcup_{i=1}^s \bigcup_{\gamma \in \Gamma_M} \gamma D(\tilde{y}_i, d(\tilde{y}_i)/2) \supset X^M(r(c_{\pi'})).$$

We pick a point  $c_0 \in \prod_{\alpha \in \pi'} (0, c_\alpha]$  then the inverse image  $(p_{P,M}^*)^{-1}(D(\tilde{y}_i, d(\tilde{y}_i)/2)) \times c_0$  is relatively compact and we can find an orbiconvex covering  $\{\mathfrak{B}\{V_{\tilde{x}_\mu}\}\}$  of this set. Then the products  $V_{\tilde{x}_\mu} \times \prod_{\alpha \in \pi'} (0, c_\alpha]$  provide an orbiconvex covering of  $X^P(c_{\pi'}, r(c_{\pi'}))$ . Of course these sets are not (relatively) compact anymore.  $\square$

This of course implies the following theorem of Raghunathan

**Theorem 2.1.2.** *If  $R$  is any commutative ring with identity and if  $\mathcal{M}$  is a finitely generated  $R - \Gamma -$  module then the total cohomology*

$$\bigoplus_{q \in \mathbb{N}} H^q(\Gamma \backslash X, sh_\Gamma(\mathcal{M}))$$

*is a finitely generated  $R$ -module*

We think that it is a very important problem to have computer programs which compute the cohomology effectively. One way to get such a software would be to write a procedure which effectively finds an orbiconvex covering for which the sets  $U_{\tilde{x}_\nu}$  are big, so that we need only few of them.

A first step would be to find effectively an optimal orbiconvex covering  $\{U_{\tilde{x}_\nu}\}$  of the set  $X^G(r)$  of almost stable points. The covering sets must not necessarily be Dirichlet balls. We could proceed and apply this also to the different  $X^M(r(c_{\pi'}))$  and find orbiconvex covers  $\{V_{\tilde{y}_\mu}^M\}$  for them. Then we may consider the inverse images  $(p_{P,M}^*)^{-1}(V_{\tilde{y}_\mu}^M \times \prod_{\alpha \in \pi'} (0, c_\alpha]) = \tilde{V}_{\tilde{y}_\mu}^M$ . This family of sets  $\{\{\gamma U_{\tilde{x}_\nu}\}, \dots, \gamma_1 \tilde{V}_{\tilde{y}_\mu}^M, \dots\}$  provide a covering of  $X$  by open sets, hence the images under the projection provide a covering

$$\mathfrak{W} = \{W_i\}_{i \in I} = \{\{U_{x_\nu}\}, \dots, \{\tilde{V}_{y_\mu}^M\}, \dots\}$$

of  $\Gamma \backslash X$ , here the index set  $I$  is the union of the  $x] \tilde{x}_\nu, \dots, {}^M y_{\tilde{y}_\mu}$ .

Of course we have a problem: The sets  $\tilde{V}_{\tilde{y}_\mu}^M$  are not acyclic anymore, so we can not use the Čech complex of this covering for the computation of the cohomology. But we know that

$$\tilde{V}_{\tilde{y}_\mu}^M \rightarrow V_{\tilde{y}_\mu}^M \times \prod_{\alpha \in \pi'} (0, c_\alpha]$$

is a fiber bundle with fiber  $U(\mathbb{Z}) \backslash U(\mathbb{R})$ , Since the base  $V_{\tilde{y}_\mu}^M \times \prod_{\alpha \in \pi'} (0, c_\alpha]$  is acyclic we know that

$$H^\bullet(\tilde{V}_{\tilde{y}_\mu}^M) \xrightarrow{\sim} \mathbb{H}^\bullet(U(\mathbb{Z}) \backslash U(\mathbb{R}), \tilde{\mathcal{M}}) \quad (2.19)$$

and we have a good understanding of the cohomology on the right. If for instance we tensor by the rationals the Theorem of Kostant (See section ??) gives us a complete description of the cohomology  $H^\bullet(U(\mathbb{Z}) \backslash U(\mathbb{R}), \tilde{\mathcal{M}} \otimes \mathbb{Q})$ .

For  $\underline{i} \in I^{p+1}$  we put  $\mathfrak{W}_{\underline{i}} = W_{i_0} \cap W_{i_1} \cap \dots \cap W_{i_p}$ . Now we follow [28], 4.6.6, for any  $q \geq 0$  write the Čech complex

$$C^\bullet(\mathfrak{W}, \mathcal{H}^q) := \prod_{\underline{i} \in I^{p+1}} H^q(W_{\underline{i}}) \rightarrow \prod_{\underline{i} \in I^{p+2}} H^q(W_{\underline{i}}) \quad (2.20)$$

and then we know that we get a spectral sequence

$$H^p(C^\bullet(\mathfrak{W}, \mathcal{H}^q)) = E_1^{p,q} \implies H^{p+q}(\Gamma \backslash X, \tilde{\mathcal{M}}) \quad (2.21)$$

### 2.1.3 Special examples in low dimensions.

We consider the group  $\Gamma = \mathrm{Sl}_2(\mathbb{Z}) / \{\pm \mathrm{Id}\}$  and its action on the upper half plane  $\mathbb{H}$ . We want to investigate the cohomology groups  $H^i(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})$  for any module  $\Gamma$ -module  $\mathcal{M}$ . Let  $p : \mathbb{H} \rightarrow \Gamma \backslash \mathbb{H}$  be the projection. We have the two special points  $i$  and  $\rho$  in  $\mathbb{H}$  they are up to conjugation by  $\Gamma$  the only points which have a non trivial stabilizer. We construct two nice orbiconvex neighborhoods of these two points. The stabilizers  $\Gamma_i$ , resp.  $\Gamma_\rho$  are cyclic and generated by the two elements

$$S = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, R = \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix}$$

respectively.

We begin with  $i$ . We consider the strip  $V_i = \{z \mid -1/2 < \Re(z) < 1/2\}$ , the element  $S$  maps the two vertical boundary lines  $\Re(z) = \pm \frac{1}{2}$  into geodesic circles starting from 0 and ending in  $\pm 2$ . Then the intersection  $\tilde{U}_i = V_i \cap S(V_i)$  is an orbiconvex neighbourhood of  $i$ .

Let us look at  $\rho$ . We consider the strip  $V_\rho = \{z \mid -0 < \Re(z) < 1\}$  and now we define  $\tilde{U}_\rho = V_\rho \cap R(V_\rho) \cap R^2(V_\rho)$ . This is a nice orbiconvex neighbourhood of  $\rho$ .

Now it is clear that these two sets provide an orbiconvex covering of  $\mathbb{H}$ , if  $U_i = p(\tilde{U}_i), U_\rho = p(\tilde{U}_\rho)$  then

$$\Gamma \backslash \mathbb{H} = U_i \cup U_\rho. \quad (2.22)$$

We have  $\tilde{\mathcal{M}}(U_i) = \text{sh}_\Gamma(\mathcal{M})(U_i) = \mathcal{M}^{\Gamma_i}$ ,  $\tilde{\mathcal{M}}(U_\rho) = \mathcal{M}^{\Gamma_\rho}$  and hence the cohomology groups are given by the cohomology of the complex

$$0 \rightarrow \mathcal{M}^{\Gamma_i} \oplus \mathcal{M}^{\Gamma_\rho} \rightarrow \mathcal{M} \rightarrow 0 \quad (2.23)$$

Then  $H^0(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) = \mathcal{M}^\Gamma = \mathcal{M}^{\Gamma_i} \cap \mathcal{M}^{\Gamma_\rho}$ . Since this is true for any  $\Gamma$  module we easily conclude that  $\Gamma$  is generated by  $\Gamma_i, \Gamma_\rho$ . And we get

$$H^1(\text{Sl}_2(\mathbb{Z}) \backslash \mathbb{H}, \tilde{\mathcal{M}}_{\mathbb{Z}}) = \mathcal{M} / (\mathcal{M}^{\langle S \rangle} \oplus \mathcal{M}^{\langle R \rangle}), \quad (2.24)$$

and the cohomology vanishes in higher degrees.

**Exercise 1:** Let  $\Gamma' \subset \Gamma = \text{Sl}_2(\mathbb{Z}) / \pm \text{Id}$  be a subgroup of finite index. Prove  
ii) We have (Shapiros lemma)

$$H^1(\Gamma' \backslash \mathbb{H}, \mathbb{Z}) = H^1(\Gamma \backslash \mathbb{H}, \widetilde{\text{Ind}}_{\Gamma'}^{\Gamma} \mathbb{Z}).$$

These cohomology groups are free of rank

$$[\Gamma : \Gamma'] - n_i - n_\rho + 1$$

where  $n_i$  (resp.  $n_\rho$ ) is the number of orbits of  $\Gamma_i$  (resp.  $\Gamma_\rho$ ) on  $\Gamma' \backslash \Gamma$ . If  $\Gamma'$  is torsion free then

$$\text{rank}(H^1 \Gamma' \backslash \mathbb{H}, \widetilde{\text{Ind}}_{\Gamma'}^{\Gamma} \mathbb{Z}) = \frac{1}{6} [\Gamma : \Gamma'] + 1$$

The Euler-characteristic of  $\Gamma' \backslash \mathbb{H}$  is  $\frac{1}{6} [\Gamma : \Gamma']$ .

**Exercise 2:** Let  $\mathcal{M}_n$  be the  $\text{Sl}_2(\mathbb{Z})$ -module of homogenous polynomials in the two variables  $X, Y$  and coefficients in  $\mathbb{Z}$ . (See 1.2.2). We have the usual action of  $\text{Sl}_2(\mathbb{Z})$  on this module by

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} P(X, Y) = P(aX + cY, bX + dY).$$

these modules define a sheaf  $\tilde{\mathcal{M}}_n$  on  $\Gamma \backslash \mathbb{H}$ , and we want to investigate their cohomology groups.

Prove:

i) If  $n$  is odd, then  $\tilde{\mathcal{M}}_n = 0$ .

Hence we assume  $n \geq 2$  and  $n$  even from now on.

ii) For  $n > 0$  we have  $H^0(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n) = 0$ .

iii) If we tensorize by  $\mathbb{Q}$ , then  $H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathbb{Q})$  is a vector space of rank  $n - 1 - 2 \left[ \frac{n}{4} \right] - 2 \left[ \frac{n}{6} \right]$ .

2.1. COHOMOLOGY OF ARITHMETIC GROUPS AS COHOMOLOGY OF SHEAVES ON  $\Gamma \backslash X$ .75

**Hint:** Diagonalise the action of  $\Gamma_i$  and  $\Gamma_\rho$  on  $\mathcal{M}_n \otimes \bar{\mathbb{Q}}$  separately and look at the eigenspaces. To say it differently: Over  $\bar{\mathbb{Q}}$  we can conjugate the matrices  $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ ,  $\begin{pmatrix} 1 & 1 \\ -1 & 0 \end{pmatrix}$  into the diagonal maximal torus  $\begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix}$ , and then look at the decomposition of  $\mathcal{M}_n$  into weight spaces.

iv) Investigate the torsion in  $H^1(\Gamma \backslash \mathbb{H}, \mathcal{M}_n)$ . (Start from the sequence  $0 \rightarrow \mathcal{M}_n \rightarrow \mathcal{M}_n \rightarrow \mathcal{M}_n/\ell\mathcal{M}_n \rightarrow 0$ .)

v) Now we consider  $\Gamma = \mathrm{Sl}_2(\mathbb{Z})$ . The two matrices  $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$  and  $R = \begin{pmatrix} -1 & 1 \\ -1 & 0 \end{pmatrix}$  are generators of the stabilisers of  $i$  and  $\rho$  respectively.

We take for our module  $\mathcal{M}$  the cyclic group  $\mathbb{Z}/12\mathbb{Z}$ , consider the spectral sequence

$$H^p(\Gamma \backslash \mathbb{H}, R^q(\mathrm{sh}_\Gamma)(\mathbb{Z}/12\mathbb{Z})).$$

Show that  $H^0(\Gamma \backslash \mathbb{H}, R^1(\mathrm{sh}_\Gamma)(\mathbb{Z}/12\mathbb{Z})) = \mathbb{Z}/12\mathbb{Z}$ . Show that the differential

$$H^0(\Gamma \backslash \mathbb{H}, R^1(\mathrm{sh}_\Gamma)(\mathbb{Z}/12\mathbb{Z})) \rightarrow H^2(\Gamma \backslash \mathbb{H}, \mathrm{sh}_\Gamma(\mathbb{Z}/12\mathbb{Z}))$$

vanishes and conclude

$$H^1(\Gamma, \mathbb{Z}/12\mathbb{Z}) = \mathbb{Z}/12\mathbb{Z}.$$

???

**The group  $\Gamma = \mathrm{Sl}_2(\mathbb{Z}[i])$**

A similar computation can be made up to compute the cohomology in the case of  $\tilde{\Gamma} = \mathrm{Gl}_2(\mathcal{O})$ . We have the three special points  $x_{12}, x_{13}$  and  $x_{23}$  (See(1.2.5)), and we choose closed sets  $A_{ij}$  containing these points which just leave out a small open strip containing the opposite face. If  $\tilde{A}_{ij}$  is a component of the inverse image of  $A_{ij}$  in  $\mathbb{H}_3$ , then

$$A_{ij} = \Gamma_{ij} \backslash \tilde{A}_{ij}.$$

The intersections  $A_{ij} \cap A_{i'j'} = A_\nu$  are closed sets. They are of the form

$$A_\nu = \Gamma_\nu \backslash \tilde{A}_\nu$$

where  $\Gamma_\nu$  is the stabilizer of the arc joining  $x_{ij}$  and  $x_{i'j'}$ . The restrictions of our sheaves  $\tilde{\mathcal{M}}$  to the  $A_{ij}$  and  $A_\nu$  and to  $A = A_{12} \cap A_{23} \cap A_{13}$  are acyclic and hence we get a complex

$$0 \longrightarrow \tilde{\mathcal{M}} \longrightarrow \bigoplus_{(i,j)} \tilde{\mathcal{M}}_{A_{ij}} \longrightarrow \bigoplus \tilde{\mathcal{M}}_{A_\nu} \longrightarrow \tilde{\mathcal{M}}_A \longrightarrow 0$$

where the  $\tilde{\mathcal{M}}_?$  are the restrictions of  $\tilde{\mathcal{M}}$  to ??? and then extended to the space again.

Hence we find that our cohomology groups are equal to the cohomology groups of the complex

$$0 \longrightarrow \bigoplus_{(i,j)} \mathcal{M}^{\Gamma_{ij}} \xrightarrow{d^1} \bigoplus_{\nu} \mathcal{M}^{\Gamma_{\nu}} \xrightarrow{d^2} \mathcal{M} \longrightarrow 0$$

with boundary maps

$$\begin{aligned} d^1 : (m_{12}, m_{13}, m_{23}) &\longmapsto (m_{12} - m_{13}, m_{23} - m_{12}, m_{13} - m_{23}) \\ d^2 : (m_1, m_2, m_3) &\longmapsto m_1 + m_2 + m_3. \end{aligned}$$

If we take for instance  $\tilde{\mathcal{M}} = \mathbb{Z}$  then we get  $H^0(\tilde{\Gamma} \backslash \mathbb{H}_3, \mathbb{Z}) = \mathbb{Z}$  and  $H^i(\tilde{\Gamma} \backslash \mathbb{H}_3, \mathbb{Z}) = 0$  for  $i > 0$  as it should be.

### Homology, Cohomology with compact support and Poincaré duality.

Here we have to use the theory of compactifications. For any locally symmetric space we can embed  $\Gamma \backslash X$  into its Borel-Serre compactification

$$i : \Gamma \backslash X \longrightarrow \Gamma \backslash \bar{X}_{BS},$$

and this process was explained in detail for our low dimensional examples. Especially we give an explicit description of a neighborhood of a point  $x \in \partial(\Gamma \backslash \bar{X}_{BS})$ . If we have a sheaf  $\tilde{\mathcal{M}}$  on  $\Gamma \backslash X$ , we can extend it to the compactification by using the functor  $i_*$ . We get a sheaf

$$i_*(\tilde{\mathcal{M}}) \quad \text{on} \quad \Gamma \backslash \bar{X}_{BS},$$

it is clear from the description of a neighborhood of a point in the boundary, that  $i_*$  is exact. ( This is not true for the Baily-Borel compactification.)

Our construction  $\mathcal{M} \rightarrow \tilde{\mathcal{M}}$  can be extended to the action of  $\Gamma$  on  $\bar{X}_{BS}$  and clearly

$$i_*(\tilde{\mathcal{M}}) = \text{result of the construction } \mathcal{M} \rightarrow \tilde{\mathcal{M}} \text{ on } \Gamma \backslash \bar{X}_{BS}.$$

Hence we get from our general results in Chapter I, ..... that

$$H^\bullet(\Gamma \backslash X, \tilde{\mathcal{M}}) = H^\bullet(\Gamma \backslash \bar{X}_{BS}, i_*(\tilde{\mathcal{M}})).$$

But we have another construction of extending the sheaf  $\tilde{\mathcal{M}}$  from  $\Gamma \backslash X$  to  $\Gamma \backslash \bar{X}_{BS}$ . This is the so called extension by zero. We define the sheaf  $i_!(\tilde{\mathcal{M}})$  on  $\Gamma \backslash \bar{X}_{BS}$  by giving the stalks. For  $x \in \Gamma \backslash \bar{X}_{BS}$  we put

$$i_!(\tilde{\mathcal{M}})_x = \begin{cases} \tilde{\mathcal{M}}_x & \text{if } x \in \Gamma \backslash X \\ 0 & \text{if } x \notin \Gamma \backslash X \end{cases}.$$

It is clear that  $i_!$  is an exact functor sending sheaves on  $\Gamma \backslash X$  to sheaves on  $\Gamma \backslash \bar{X}_{BS}$ , and we have for an arbitrary sheaf

$$H^0(\Gamma \backslash \bar{X}_{BS}, i_!(\mathcal{F})) = H_c^0(\Gamma \backslash X, \mathcal{F})$$

## 2.1. COHOMOLOGY OF ARITHMETIC GROUPS AS COHOMOLOGY OF SHEAVES ON $\Gamma \backslash X$ .77

where  $H_c^0(\Gamma \backslash X, \mathcal{F})$  is the abelian group of those sections  $s \in H^0(\Gamma \backslash X, \mathcal{F})$  for which the support

$$\text{supp}(s) = \{x \mid s_x \neq 0\}$$

is compact.

Hence we define the cohomology with compact supports as

$$H_c^q(\Gamma \backslash X, \mathcal{F}) = H^q(\Gamma \backslash \bar{X}_{BS}, i_! \mathcal{F}).$$

If  $\tilde{\mathcal{M}}$  is a sheaf on  $\Gamma \backslash X$  which is obtained from a  $\Gamma$ -module  $\mathcal{M}$ , then it is quite clear that

$$H_c^0(\Gamma \backslash X, \tilde{\mathcal{M}}) = 0,$$

provided our quotient  $\Gamma \backslash X$  is not compact.

The cohomology with compact supports is actually related to the homology of the group: I want to indicate that we have a natural isomorphism

$$H_i(\Gamma, \mathcal{M}) \simeq H_c^{d-i}(\Gamma \backslash X, \tilde{\mathcal{M}})$$

under the assumption that  $X$  is connected and the orders of the stabilizers are invertible in  $R$ .

This is the analogous statement to the theorem .... which we discussed when we introduced cohomology.

Our starting point is the fact that the projective  $\Gamma$ -modules have analogous vanishing properties as the induced modules.

**Lemma:** *Let us assume that  $\Gamma$  acts on the connected symmetric space  $X$ . If  $P$  is a projective module then*

$$H_c^i(\Gamma \backslash X, \tilde{P}) = \begin{cases} 0 & \text{if } i \neq \dim X \\ P_\Gamma & \text{if } i = \dim X. \end{cases}$$

Let us believe this lemma. Then it is quite clear that

$$H_i(\Gamma, \mathcal{M}) \simeq H_c^{d-i}(\Gamma \backslash X, \tilde{P}),$$

because both sides can be computed from a projective resolution.

### 2.1.4 The homology as singular homology

We have still another description of the homology. We form the singular chain complex

$$\rightarrow C_i(X) \rightarrow C_{i-1}(X) \rightarrow \dots \rightarrow C_0(X) \rightarrow 0.$$

This is a complex of  $\Gamma$ -modules, and we can form the tensor product with  $\mathcal{M}$ . We get a complex of  $\Gamma$ -modules

$$\xrightarrow{d_{i+1}} C_i(X) \otimes \mathcal{M} \xrightarrow{d_i} C_{i-1}(X) \otimes \mathcal{M} \rightarrow \dots$$

We define the chain complex

$$C_\bullet(\Gamma \backslash X, \underline{\mathcal{M}}),$$

simply as the resulting complex of  $\Gamma$ -coinvariants. The homology groups are defined as

$$H_i(\Gamma \backslash X, \underline{\mathcal{M}}) = \ker(d_i) / \text{Im}(d_{i+1}) \quad (2.25)$$

### The cosheaves

The symbol  $\underline{\mathcal{M}}$  should be interpreted as the cosheaf attached to our  $\Gamma$ -module, this is an object which is dual to the sheaf  $\tilde{\mathcal{M}}$ . For a point  $\bar{x} \in \Gamma \backslash X$  costalk  $\underline{\mathcal{M}}_{\bar{x}}$  is given as follows: As in (2.1) we consider the projection  $\pi_\Gamma : X \rightarrow \Gamma \backslash X$  and maps with finite support

$$\mathcal{C}(\bar{x}, \mathcal{M}) := \{f : \pi_\Gamma^{-1}(\bar{x}) \rightarrow \mathcal{M}\}. \quad (2.26)$$

On this module we have an action of  $\Gamma$  which is given by act

$$(\gamma f)(x) = \gamma(f(\gamma^{-1}x)). \quad (2.27)$$

Then our costalk is given by the coinvariants

$$\underline{\mathcal{M}}_{\bar{x}} = \mathcal{C}(\bar{x}, \mathcal{M})_\Gamma = \mathcal{C}(\bar{x}, \mathcal{M}) / \{f - \gamma f, \gamma \in \Gamma, f \in \mathcal{C}(\bar{x}, \mathcal{M})\} \quad (2.28)$$

We have the homomorphism  $\int : \mathcal{M}_{\bar{x}} \rightarrow \mathcal{M}$  which is given by summation  $f \mapsto \sum_{x \in \pi_\Gamma^{-1}(\bar{x})} f(x)$  and this induces an isomorphism invint

$$\int : \mathcal{C}(\bar{x}, \mathcal{M})_\Gamma \xrightarrow{\sim} \underline{\mathcal{M}}_{\bar{x}} \quad (2.29)$$

We pick a point  $x \in \pi_\Gamma^{-1}(\bar{x})$  and an open neighborhood  $U_x$  of  $x$  such that  $\gamma U_x \cap U_x \neq \emptyset$  implies  $\gamma \in \Gamma_x$ . We consider the space  $\mathcal{C}(\bar{x}, x, \mathcal{M})$  of those maps, which are supported in the point  $x$ . This space is of course equal to  $\mathcal{M}$  and the composition

$$\delta_x : \mathcal{C}(\bar{x}, x, \mathcal{M}) \rightarrow \mathcal{C}(\bar{x}, \mathcal{M}) \rightarrow \underline{\mathcal{M}}_{\bar{x}}$$

induces an isomorphism

$$\delta_x : \mathcal{M}_{\Gamma_x} \xrightarrow{\sim} \underline{\mathcal{M}}_{\bar{x}} \quad (2.30)$$

If we pick a second point  $\bar{y} \in \pi_\Gamma(U_x)$  and a  $y \in \pi_\Gamma^{-1}(\bar{y}) \cap U_x$  then clearly  $\Gamma_y \subset \Gamma_x$  and therefore we get a specialization map

$$r_{\bar{y}, \bar{x}} : \underline{\mathcal{M}}_{\bar{y}} \rightarrow \underline{\mathcal{M}}_{\bar{x}}. \quad (2.31)$$

Now it becomes clear why these objects are called cosheaves. For the sheaf  $\tilde{\mathcal{M}}$  we get in the corresponding situation a map in the opposite direction

$$\tilde{\mathcal{M}}_{\bar{x}} \rightarrow \tilde{\mathcal{M}}_{\bar{y}} \quad (2.32)$$

as a specialization map between the stalks of  $\tilde{\mathcal{M}}$ . An element  $f^* \in \underline{\mathcal{M}}_{\bar{x}}$  can be

represented as an array refcos

$$f^* = \{\dots, f(x), \dots\}_{x \in \pi^{-1}(\bar{x})} \quad (2.33)$$

where  $f(x) \in (\underline{\mathcal{M}}_{\bar{x}})_{\Gamma_x}$  and  $f(\gamma x) = \gamma f(x)$ .

Now we can give a different description of the group of  $i$ -chains  $C_i(\Gamma \backslash X, \underline{\mathcal{M}})$ : An  $i$ -chain with values in the cosheaf  $\underline{\mathcal{M}}$  is of the form  $\sigma \otimes f$  where  $\sigma : \Delta^i \rightarrow \Gamma \backslash X$  is a continuous (differentiable) map from the  $i$  dimensional simplex  $\Delta^i$  to  $\Gamma \backslash X$  and where  $f$  is a section in the cosheaf, i.e.  $f_x \in \underline{\mathcal{M}}_{\sigma(x)}$  and where  $f_x$  varies continuously. (This means: If  $\sigma(y)$  specializes to  $\sigma(x)$  then  $r_{\sigma(y), \sigma(x)}(f_y) = f_x$ .)

Then  $C_i(\Gamma \backslash X, \underline{\mathcal{M}})$  is the free abelian group generated by these  $i$  chains with values in  $\underline{\mathcal{M}}$ . Then the boundary maps  $d_i$  are defined in the usual way and we get a slightly different description of the homology groups  $H_i(\Gamma \backslash X, \underline{\mathcal{M}})$ .

But we may choose for our module  $\mathcal{M}$  simply the group ring. Then

$$(C_\bullet(X) \otimes \mathbb{Z}[\Gamma])_\Gamma \simeq C_\bullet(X),$$

and hence we have, since  $X$  is a cell, that

$$H_i(\Gamma \backslash X, \underline{\mathbb{Z}[\Gamma]}) = 0 \quad \text{for } i > 0.$$

On the other hand we have

$$H_0(\Gamma \backslash X, \underline{\mathcal{M}}) = \mathcal{M}_\Gamma.$$

This follows directly from looking at the complex

$$(C_1(X) \otimes \mathcal{M})_\Gamma \longrightarrow (C_0(X) \otimes \mathcal{M})_\Gamma.$$

First of all we observe that 0-cycles

$$x_1 \otimes m - x_0 \otimes m$$

are boundaries since  $X$  is pathwise connected. On the other hand we have that

$$x_0 \otimes m - \gamma x_0 \otimes \gamma m \in C_0(X) \otimes \mathcal{M}$$

becomes zero if we go to the coinvariants and this implies the assertion.

If we have in addition that the orders of the stabilizers are invertible in  $R$  than it is clear that a short exact sequence of  $R$ - $\Gamma$ -modules

$$0 \longrightarrow \mathcal{M}' \longrightarrow \mathcal{M} \longrightarrow \mathcal{M}'' \longrightarrow 0$$

leads to an exact sequence of complexes

$$0 \longrightarrow C_\bullet(\Gamma \backslash X, \underline{\mathcal{M}'}) \longrightarrow C_\bullet(\Gamma \backslash X, \underline{\mathcal{M}}) \longrightarrow C_\bullet(\Gamma \backslash X, \underline{\mathcal{M}''}) \longrightarrow 0,$$

and hence to a long exact cohomology sequence

$$H_i(\Gamma \backslash X, \underline{\mathcal{M}'}) \longrightarrow H_i(\Gamma \backslash X, \underline{\mathcal{M}}) \longrightarrow H_i(\Gamma \backslash X, \underline{\mathcal{M}''}) \longrightarrow H_{i-1}(\Gamma \backslash X, \underline{\mathcal{M}'}).$$

Now it is clear that

$$H_i(\Gamma, \mathcal{M}) \simeq H_i(\Gamma \backslash X, \underline{\mathcal{M}}) \simeq H_c^{d-i}(\Gamma \backslash X, \tilde{\mathcal{M}}).$$

fundex

### 2.1.5 The fundamental exact sequence

By construction we have the exact sequence

$$0 \rightarrow i_!(\tilde{\mathcal{M}}) \rightarrow i_*(\tilde{\mathcal{M}}) \rightarrow i_*(\tilde{\mathcal{M}})/i_!(\tilde{\mathcal{M}}) \rightarrow 0$$

of sheaves and clearly  $i_*(\tilde{\mathcal{M}})/i_!(\tilde{\mathcal{M}})$  is simply the restriction of  $i_*(\tilde{\mathcal{M}})$  to the boundary extended by zero to the entire space. This yields the *fundamental exact sequence*

$$\rightarrow H^{q-1}(\partial(\Gamma \backslash X), \tilde{\mathcal{M}}) \rightarrow H_c^q(\Gamma \backslash X, \tilde{\mathcal{M}}) \rightarrow H^q(\Gamma \backslash \bar{X}, \tilde{\mathcal{M}}) \xrightarrow{r} H^q(\partial(\Gamma \backslash X), \tilde{\mathcal{M}}) \rightarrow \dots$$

We define the “inner cohomology”  $\boxed{\text{inncoh}}$

$$H_!^q(\Gamma \backslash X, \tilde{\mathcal{M}}) := \text{Im}(H_c^q(\Gamma \backslash X, \tilde{\mathcal{M}}) \rightarrow H^q(\Gamma \backslash \bar{X}, \tilde{\mathcal{M}})) = \ker H^q(\Gamma \backslash \bar{X}, \tilde{\mathcal{M}}) \xrightarrow{r} H^q(\partial(\Gamma \backslash X), \tilde{\mathcal{M}}) \quad (2.34)$$

( This a little bit misleading because these groups are not honest cohomology groups, they are not the cohomology groups of a space with coefficients in a sheaf. An exact sequence of sheaves  $0 \rightarrow \mathcal{M}' \rightarrow \mathcal{M} \rightarrow \mathcal{M}'' \rightarrow 0$  does not provide an exact sequence for these  $H_!$  groups. )

In the special case that the underlying group  $G/\mathbb{Q}$  is anisotropic the fundamental exact sequence becomes trivial, in this case the quotient  $\Gamma \backslash X$  is compact and we have

$$H^\bullet(\Gamma \backslash X, \tilde{\mathcal{M}}) = H_c^\bullet(\Gamma \backslash X, \tilde{\mathcal{M}}) = H_!^\bullet(\Gamma \backslash X, \tilde{\mathcal{M}}).$$

Many authors prefer to consider the case of a compact quotient  $\Gamma \backslash X$ , but I think we lose some very interesting phenomena if we concentrate on this case. On the other hand we do not need to read the next subsection. Also readers who are more interested in the low dimensional cases and the more specific results in these cases may well skip reading the next subsection.

#### The cohomology of the boundary

We want to have a slightly different look at this sequence. We recall the covering (See 1.87, 8.31)

$$\Gamma \backslash X = \Gamma \backslash X(r) \cup \overset{\bullet}{\mathcal{N}}(\Gamma \backslash X) = \Gamma \backslash X(r) \cup \bigcup_{P: P \text{ proper}} \Gamma_P \backslash X^P(\underline{c}_{\pi'}, r(\underline{c}_{\pi'})) \quad (2.35)$$

where the union runs over  $\Gamma$  conjugacy classes of parabolic subgroups over  $\mathbb{Q}$  and  $\overset{\bullet}{\mathcal{N}}(\Gamma \backslash X)$  is a punctured tubular neighborhood of  $\infty$ , i.e. the boundary of the Borel-Serre compactification.

It is well known (See for instance [book] vol I , 4.5 ) that from a covering  $\Gamma \backslash X = \bigcup_i V_i$  we get a Čech complex and a spectral sequence with  $E_1^{p,q}$ - term

$$\prod_{i=\{i_0, i_1, \dots, i_p\}} H^q(V_{\underline{i}}, \tilde{\mathcal{M}}) \quad (2.36)$$

2.1. COHOMOLOGY OF ARITHMETIC GROUPS AS COHOMOLOGY OF SHEAVES ON  $\Gamma \backslash X$ .81

where  $V_{\underline{i}} = V_{i_0} \cap \dots \cap V_{i_p}$ . The boundary in the Čzech complex gives us the differential

$$d_1^{p,q} : \prod_{\underline{i}=\{i_0, i_1, \dots, i_p\}} H^q(V_{\underline{i}}, \tilde{\mathcal{M}}) \rightarrow \prod_{\underline{j}=\{j_0, j_1, \dots, j_{p+1}\}} H^q(V_{\underline{j}}, \tilde{\mathcal{M}}) \quad (2.37)$$

Here we work with the alternating Čzech complex, we also assume that we have an ordering on the set of simple positive roots. If such a  $V_{\underline{i}}$  is non empty then it of the form  $\Gamma_Q \backslash X^Q(C(\underline{\tilde{c}}))$ .

We return to the diagram (??), on the left hand side we can divide by  $\Gamma_Q$ . We have the map which maps a Cartan involution on  $X$  to a Cartan-involution on  $M$ . Then we get a diagram

$$\begin{array}{ccc} f^\dagger : X^Q(C(\underline{\tilde{c}})) & \rightarrow & X^M(r) \times C_{U_Q}(\underline{\tilde{c}}) \\ \downarrow p_Q & & \downarrow p_M \\ f : \Gamma_Q \backslash X^Q(C(\underline{\tilde{c}})) & \rightarrow & \Gamma_M \backslash X^M(r) \times C_{U_Q}(\underline{\tilde{c}}) \end{array} \quad (2.38)$$

where the bottom line is a fibration. To describe the fiber in a point  $\tilde{x}$  we pick a point  $x \in (p_m \circ f^\dagger)^{-1}$ . Then  $U_Q(\mathbb{R})$  acts simply transitively on the fiber  $(f^\dagger)^{-1}(f^\dagger(x))$  hence  $U_Q(\mathbb{R}) = (f^\dagger)^{-1}(f^\dagger(x))$ . Then  $p_Q : U_Q(\mathbb{R}) \rightarrow \Gamma_{U_Q} \backslash U_Q(\mathbb{R})$  yields the identification  $i_x : \Gamma_{U_Q} \backslash U_Q(\mathbb{R}) \xrightarrow{\sim} f^{-1}(\tilde{x})$ . If we replace  $x$  by  $\gamma x = x_1$  with  $\gamma \in \Gamma_{U_Q}$  then we get  $i_{x_1} = \text{Ad}(\gamma) \circ i_x$  where for  $u \in U_Q$   $\text{Ad}(\gamma)(u) = \gamma u \gamma^{-1}$  where for  $u \in U_Q(\mathbb{R})$ , under this action of  $\Gamma_Q$ .

We have the spectral sequence

$$H^p(\Gamma_M \backslash X^M(r), R^q f_*(\tilde{\mathcal{M}})) \Rightarrow H^{p+q}(\Gamma_Q \backslash X^Q(C(c_{\pi_1}, \dots, c_{\pi_\nu})), \tilde{\mathcal{M}})$$

and clearly  $R^q f_*(\tilde{\mathcal{M}})$  is a locally constant sheaf. This sheaf is easy to determine. Under the above identification we get an isomorphism

$$i_x^\bullet : H^\bullet(\Gamma_{U_Q} \backslash U_Q(\mathbb{R}), \tilde{\mathcal{M}}) \xrightarrow{\sim} R^\bullet(\tilde{\mathcal{M}})_{\tilde{x}}.$$

The adjoint action  $\text{Ad} : \Gamma_Q \rightarrow \text{Aut}(\Gamma_{U_Q} \backslash U_Q(\mathbb{R}))$  induces an action of  $\Gamma_Q$  on the cohomology  $H^\bullet((\Gamma_{U_Q} \backslash U_Q(\mathbb{R})), \tilde{\mathcal{M}})$ . Since the functor cohomology is the derived functor of taking  $\Gamma_{U_Q}$  invariants it follows that the restriction of  $\text{Ad}$  to  $\Gamma_{U_Q}$  acts trivially on  $H^\bullet(\Gamma_{U_Q} \backslash U_Q(\mathbb{R}), \tilde{\mathcal{M}})$ . Consequently  $H^\bullet((\Gamma_{U_Q} \backslash U_Q(\mathbb{R})), \tilde{\mathcal{M}})$  is a  $\Gamma_M$ -module. We get

$$R^\bullet f_*(\tilde{\mathcal{M}}) \xrightarrow{\sim} H^\bullet(\widetilde{\Gamma_{U_Q} \backslash U_Q(\mathbb{R})}, \tilde{\mathcal{M}})$$

and hence our spectral sequence becomes

vEst

$$H^p(\Gamma_M \backslash X^M(r), H^\bullet(\widetilde{\Gamma_{U_Q} \backslash U_Q(\mathbb{R})}, \tilde{\mathcal{M}})) \Rightarrow H^{p+q}(\Gamma_Q \backslash X^Q(C(\underline{\tilde{c}})), \tilde{\mathcal{M}}) \quad (2.39)$$

We can take the composition  $r_Q \circ f$ . Then it is obvious that for any point  $c_0 \in C_{U_Q}(\underline{\tilde{c}})$  the restriction map

$$H^\bullet(X^Q(C(\underline{\tilde{c}})), \tilde{\mathcal{M}}) \rightarrow H^\bullet(X^Q((r_Q \circ f)^{-1}(c_0)), \tilde{\mathcal{M}}) \quad (2.40)$$

is an isomorphism. On the other hand it is clear that we may vary our parameter  $\tilde{c}$  we may assume that the  $C_{U_Q}(\tilde{c})$  go to infinity. Then we may enlarge the parameter  $r$  without violating the assumptions in proposition 1.2.3. Hence we get that the inclusion  $\Gamma_Q \backslash X^Q(C(\tilde{c})) \subset \Gamma_Q \backslash X^Q$  induces an isomorphism in cohomology

$$H^\bullet(\Gamma_Q \backslash X^Q(C(\tilde{c}), \tilde{\mathcal{M}})) \xrightarrow{\sim} H^\bullet(\Gamma_Q \backslash X, \tilde{\mathcal{M}}) \quad (2.41)$$

We choose a total ordering on the set of  $\Gamma$  conjugacy classes of parabolic subgroups, i.e. we enumerate them by a finite interval of integers  $[1, N]$ . We also enumerate the set of simple roots  $\{\alpha_1, \dots, \alpha_d\}$  in our special case  $\alpha_i = \alpha_{i, i+1}$ . For any conjugacy class  $[P]$  we define the type of  $P$  to be  $t(P) = \pi^{U_P}$  the subset of unipotent simple roots and  $d(P) = \#\pi^{U_P}$  the cardinality of this set. If  $P_{i_1}, \dots, P_{i_r}$  are maximal,  $i_1 < i_2 < \dots < i_r$  and if  $P_{i_1} \cap \dots \cap P_{i_r} = Q$  is a parabolic subgroup then we require that  $t(P_{i_1}) < \dots < t(P_{i_r})$ .

The indexing set  $\text{Par}(\Gamma)$  of our covering is the  $\Gamma$  conjugacy classes of parabolic subgroups over  $\mathbb{Q}$ . If we have a finite set  $[P_{i_0}], [P_{i_1}], \dots, [P_{i_p}]$  of conjugacy classes then we say  $[Q] \in [P_{i_0}], [P_{i_1}], \dots, [P_{i_p}]$  if we can find representatives  $P'_{i_\nu} \in [P_{i_\nu}]$  and  $Q' \in [Q]$  such that  $Q' = P'_{i_0} \cap \dots \cap P'_{i_p}$ .

Hence we see that the  $E_1^{\bullet, q}$  complex in our spectral sequence (2.37) is given by

$$\prod_i H^q(\Gamma_{Q_i} \backslash X^{Q_i}(C(\tilde{c}), \tilde{\mathcal{M}})) \rightarrow \prod_{i < j} \prod_{[R] \in [Q_i] \cap [Q_j]} H^q(\Gamma_R \backslash X^R(C(\tilde{c}), \tilde{\mathcal{M}})) \rightarrow \quad (2.42)$$

this obtained from our covering (8.31). Now we replace our covering by a simplicial space, i.e. we consider the diagram of maps between spaces

$$\mathfrak{Par} := \prod_i \Gamma_{Q_i} \backslash X \begin{array}{c} \xleftarrow{p_1} \\ \xleftarrow{p_2} \end{array} \prod_{i < j} \prod_{[R] \in [Q_i] \cap [Q_j]} \Gamma_R \backslash X \begin{array}{c} \xleftarrow{\quad} \\ \xleftarrow{\quad} \\ \xleftarrow{\quad} \end{array} \quad (2.43)$$

this yields a spectral sequence with  $E_1^{\bullet, q}$  term

$$\prod_i H^q(\Gamma_{Q_i} \backslash X, \tilde{\mathcal{M}}) \xrightarrow{d^{(0)}} \prod_{i < j} \prod_{[R] \in [P_i] \cap [P_j]} H^q(\Gamma_R \backslash X^R, \tilde{\mathcal{M}}) \xrightarrow{d^{(1)}} \quad (2.44)$$

Our covering also yields a simplicial space which is a subspace of (2.43) we get a map from (2.37) to (2.44) and this map is an isomorphism of complexes.

We replace  $\mathfrak{Par}$  by another simplicial complex

$$\mathfrak{Parma} := \prod_{[P]: d(P)=1} \Gamma_P \backslash X \begin{array}{c} \xleftarrow{p_1} \\ \xleftarrow{p_2} \end{array} \prod_{[Q]: d(Q)=2} \Gamma_Q \backslash X \begin{array}{c} \xleftarrow{\quad} \\ \xleftarrow{\quad} \\ \xleftarrow{\quad} \end{array} \quad (2.45)$$

We have an obvious projection  $\Pi : \mathfrak{Par} \rightarrow \mathfrak{Parma}$  which induces a homomorphism

$$\begin{array}{ccc}
 \prod_i H^q(\Gamma_{Q_i} \backslash X, \tilde{\mathcal{M}}) & \xrightarrow{d^{(0)}} & \prod_{i < j} \prod_{[R] \in [P_i] \cap [P_j]} H^q(\Gamma_R \backslash X^R, \tilde{\mathcal{M}}) & \xrightarrow{d^{(1)}} \\
 \uparrow & & \uparrow & \\
 \prod_{[P]:d(P)=1} H^q(\Gamma_P \backslash X, \tilde{\mathcal{M}}) & \xrightarrow{d^{(0)}} & \prod_{[R]:d(R)=2} H^q(\Gamma_R \backslash X^R, \tilde{\mathcal{M}}) & \xrightarrow{d^{(1)}} \\
 & & & (2.46)
 \end{array}$$

and an easy argument in homological algebra shows that this induces an isomorphism in cohomology or in other words an isomorphism of the  $E_2^{p,q}$  terms of the two spectral sequences.

We had the covering

$$\dot{\mathcal{N}}(\Gamma \backslash X) = \bigcup_{P: P \text{ proper}} \Gamma_P \backslash X^P(\underline{c}_{\pi'}, r(\underline{c}_{\pi'})) \quad (2.47)$$

which gives us the spectral sequence converging to  $H^\bullet(\dot{\mathcal{N}}(\Gamma \backslash X), \tilde{\mathcal{M}})$  with

$$E_1^{p,q} = \bigoplus_{i_0 < i_1 < \dots < i_p} \bigoplus_{[Q] \in [P_{i_0}] \cap [P_{i_1}] \cap \dots \cap [P_{i_p}]} H^q(\Gamma_Q \backslash X^Q(\underline{c}_{\pi'}, r(\underline{c}_{\pi'}), \tilde{\mathcal{M}})) \quad (2.48)$$

Our covering of  $\dot{\mathcal{N}}(\Gamma \backslash X)$  gives us a simplicial space  $\mathfrak{Cov}(\dot{\mathcal{N}}(\Gamma \backslash X))$  and we have maps

$$\mathfrak{Cov}(\dot{\mathcal{N}}(\Gamma \backslash X)) \hookrightarrow \mathfrak{Par} \rightarrow \mathfrak{Par}_{\text{max}}. \quad (2.49)$$

We saw that the resulting maps induced an isomorphism in the  $E_2^{p,q}$  terms of the spectral sequences. Hence we see that  $\mathfrak{Par}_{\text{max}}$  yields a spectral sequence

$$E_1^{p,q} = \bigoplus_{[P]:d(P)=p+1} H^q(\Gamma_P \backslash X, \tilde{\mathcal{M}}) \Rightarrow H^{p+q}(\dot{\mathcal{N}}(\Gamma \backslash X), \tilde{\mathcal{M}}) \quad (2.50)$$

At this point we want to raise an interesting question

*Does this spectral sequence degenerate at  $E_2^{p,q}$  level?*

The author of this book is hoping that the answer to this question is no! And this is so for interesting reasons! We come back to this question when we discuss the Eisenstein cohomology.

The complement of  $\dot{\mathcal{N}}(\Gamma \backslash X)$  is a relatively compact open set  $V \subset \Gamma \backslash X$ , this set contains the stable points. We define  $\tilde{\mathcal{M}}_V^! = i_{V,!}(\tilde{\mathcal{M}})$  then we get an exact sequence

$$0 \rightarrow \tilde{\mathcal{M}}_V^! \rightarrow \tilde{\mathcal{M}} \rightarrow \tilde{\mathcal{M}}/\tilde{\mathcal{M}}_V^! \rightarrow 0 \quad (2.51)$$

and  $\tilde{\mathcal{M}}/\tilde{\mathcal{M}}_V^!$  is obviously the extension of the restriction of  $\tilde{\mathcal{M}}$  to  $\dot{\mathcal{N}}(\Gamma \backslash X)$  and the extended by zero to  $\Gamma \backslash X$ . We claim (easy proof later) that

$$H_c^\bullet(\Gamma \backslash X, \tilde{\mathcal{M}}) = H^\bullet(\Gamma \backslash X, \tilde{\mathcal{M}}_V^!) \quad (2.52)$$

and this gives us again the fundamental exact sequence fix

$$H^{q-1}(\dot{\mathcal{N}}(\Gamma \backslash X), \tilde{\mathcal{M}}) \rightarrow H^q(\Gamma \backslash X, \tilde{\mathcal{M}}_V^!) \rightarrow H^q(\Gamma \backslash X, \tilde{\mathcal{M}}) \rightarrow H^q(\dot{\mathcal{N}}(\Gamma \backslash X), \tilde{\mathcal{M}}) \rightarrow \quad (2.53)$$

### 2.1.6 How to compute the cohomology groups $H_c^q(\Gamma \backslash X, \tilde{\mathcal{M}})$

We apply the considerations in 4.8 from the [book]. Again we cover  $\Gamma \backslash X$  by orbiconvex open neighborhoods  $U_{x_i}$ , and now we define

$$\tilde{\mathcal{M}}_{\underline{x}}^! = (i_{\underline{x}})_! i_{\underline{x}}^*(\tilde{\mathcal{M}}).$$

These sheaves have properties, which are dual to those of the sheaves  $\tilde{\mathcal{M}}_{\underline{x}}$ . If  $\underline{x} = (x_1, \dots, x_s)$  and if we add another point  $\underline{x}' = (x_1, \dots, x_s, x_{s+1})$  then we have the restriction  $\tilde{\mathcal{M}}_{\underline{x}} \rightarrow \tilde{\mathcal{M}}_{\underline{x}'}$ , which were used to construct the Čech resolution.

Now let  $d = \dim(X)$ . For the  $!$  sheaves we get (See [book], loc. cit.) get a morphism  $\tilde{\mathcal{M}}_{\underline{x}'}^! \rightarrow \tilde{\mathcal{M}}_{\underline{x}}^!$ . For  $\underline{x} = (x_1, \dots, x_s)$  we define the degree  $d(\underline{x}) = d+1-s$ . Then we construct the Čech-coresolution (See [book], 4.8.3)

$$\rightarrow \prod_{\underline{x}:d(\underline{x})=q} \tilde{\mathcal{M}}_{\underline{x}}^! \rightarrow \cdots \rightarrow \prod_{(x_i, x_j)} \tilde{\mathcal{M}}_{x_i, x_j}^! \rightarrow \prod_{x_i} \tilde{\mathcal{M}}_{x_i}^! \rightarrow i_!(\tilde{\mathcal{M}}) \rightarrow 0.$$

Now we have a dual statement to the proposition with label **acyc**

Proposition: (**acyc!**) If  $d = \dim(X)$  then

$$H^q(U_{\underline{x}}, \tilde{\mathcal{M}}_{\underline{x}}^!) = \begin{cases} \mathcal{M}_{\Gamma_{\bar{y}}} & q = d \\ 0 & q \neq d \end{cases}$$

Hence the above complex of sheaves provides a complex of modules  $C_!^*(\mathfrak{U}, \tilde{\mathcal{M}})$ :

$$\rightarrow \prod_{\underline{x}:d(\underline{x})=q} H^d(U_{\underline{x}}, \tilde{\mathcal{M}}_{\underline{x}}^!) \rightarrow \cdots \rightarrow \prod_{(x_i, x_j)} H^d(U_{x_i, x_j}, \tilde{\mathcal{M}}_{x_i, x_j}^!) \rightarrow \prod_{x_i} \tilde{H}^d(U_{x_i}, \tilde{\mathcal{M}}_{x_i}^!) \rightarrow 0.$$

Now it is clear that

$$H^q(\Gamma \backslash X, i_!(\tilde{\mathcal{M}})) = H_c^q(\Gamma \backslash X, \tilde{\mathcal{M}}) = H^q(C_!^*(\mathfrak{U}, \tilde{\mathcal{M}})).$$

Now let us assume that  $\mathcal{M}$  is a finitely generated module over some commutative noetherian ring  $R$  with identity. Then clearly all our cohomology groups will be  $R$ -modules.

Our Theorem A above implies

**Theorem** (Raghunathan) *Under our general assumptions all the cohomology groups  $H_c^q(\Gamma \backslash X, \tilde{\mathcal{M}})$ ,  $H^q(\Gamma \backslash X, \tilde{\mathcal{M}})$ ,  $H_!^q(\Gamma \backslash X, \tilde{\mathcal{M}})$ ,  $H^q(\partial(\Gamma \backslash X), \tilde{\mathcal{M}})$  are finitely generated  $R$  modules.*

### 2.1.7 Modified cohomology groups

Most of the time our module  $\mathcal{M}$  will be a finitely generated  $\mathbb{Z}$  module and the theorem of Raghunathan says that the cohomology groups are also finitely generated  $\mathbb{Z}$  modules. Sometimes we replace  $\mathbb{Z}$  ring of integers  $\mathcal{O}_F$  of a finite extension  $F/\mathbb{Q}$  and then we will even invert some finite numbers of primes.

Hence we our coefficient modules will be finitely generated  $R$ -modules where  $\mathcal{O}_F \subset R \subset F$ . In any case these rings  $R$  will be Dedekind rings.

Starting from the fundamental exact sequence we have introduced the modified cohomology groups  $H_!^q(\ )$ . There is a second process of modification: If  $H^\bullet(\ )$  is any of these cohomology groups then Hint

$$H^\bullet(\ )_{\text{int}} := H^\bullet(\ ) / \text{Tors} = \text{Im}(H^\bullet(\ ) \rightarrow H^\bullet(\ ) \otimes \mathbb{Q}) \quad (2.54)$$

We have to discuss a minor problem: These two processes of modification do not quite commute. This is due to the fact that the resulting sequence

$$\rightarrow H^{q-1}(\partial(\Gamma \backslash X), \tilde{\mathcal{M}}_R)_{\text{int}} \rightarrow H_c^q(\Gamma \backslash X, \tilde{\mathcal{M}}_R)_{\text{int}} \xrightarrow{j} H^q(\Gamma \backslash \bar{X}, \tilde{\mathcal{M}}_R)_{\text{int}} \xrightarrow{r} H^q(\partial(\Gamma \backslash X), \tilde{\mathcal{M}}_R)_{\text{int}}$$

is not necessarily exact anymore. Clearly we have  $H_!^q(\Gamma \backslash X, \tilde{\mathcal{M}}_R)_{\text{int}} = \text{Im}(j)$  and if we now define  $H^q \Gamma \backslash X, \tilde{\mathcal{M}}_R)_{\text{int},!} := \ker(r)$  then we have

$$H_!^q(\Gamma \backslash X, \tilde{\mathcal{M}}_R)_{\text{int}} \subset H^q \Gamma \backslash X, \tilde{\mathcal{M}}_R)_{\text{int},!}$$

but this inclusion may be proper. The following proposition is an elementary exercise in homological algebra supqbd

**Proposition 2.1.1.** *The quotient  $H^q(\Gamma \backslash X, \tilde{\mathcal{M}}_R)_{\text{int},!} / H_!^q(\Gamma \backslash X, \tilde{\mathcal{M}}_R)_{\text{int}}$  is finite and isomorphic to a subquotient of  $H^q(\partial(\Gamma \backslash X), \tilde{\mathcal{M}}_R)$*

We will discuss an example in section 3.3.1

### 2.1.8 The case $\Gamma = \text{Sl}_2(\mathbb{Z})$

In this book we study intensively the special case  $\Gamma = \text{Sl}_2(\mathbb{Z})$ . In this case we can formulate and prove some very specific results, especially we understand the denominators of the Eisenstein classes (Theorem 3.67).

In the following  $\mathcal{M}$  can be any  $\Gamma$ -module. We investigate the fundamental exact sequence for this special group.

Of course we start again from our covering  $\Gamma \backslash \mathbb{H} = U_i \cup U_\rho$ . The cohomology with compact supports is the cohomology of the complex

$$0 \rightarrow H^2(U_i \cap U_\rho, \tilde{\mathcal{M}}_{i,\rho}^!) \rightarrow H^2(U_i, \tilde{\mathcal{M}}_i^!) \oplus H^2(U_\rho, \tilde{\mathcal{M}}_\rho^!) \rightarrow 0.$$

Now we have  $H^2(U_i \cap U_\rho, \tilde{\mathcal{M}}_{i,\rho}^!) = M, H^2(U_i, \tilde{\mathcal{M}}_i^!) = \mathcal{M}_{\Gamma_i} = \mathcal{M}/(\text{Id} - S)M, H^2(U_\rho, \tilde{\mathcal{M}}_\rho^!) = \mathcal{M}_{\Gamma_\rho} = \mathcal{M}/(\text{Id} - R)\mathcal{M}$  and hence we get the complex

$$0 \rightarrow \mathcal{M} \rightarrow \mathcal{M}_{\Gamma_i} \oplus \mathcal{M}_{\Gamma_\rho} \rightarrow 0$$

and from this we obtain

$$H^1(\Gamma \backslash \mathbb{H}, i_!(\mathcal{M})) = \ker(\mathcal{M} \rightarrow (M/(\text{Id} - S)M \oplus \mathcal{M}/(\text{Id} - R)M))$$

and

$$H^0(\Gamma \backslash \mathbb{H}, i_!(\mathcal{M})) = 0, H^2(\Gamma \backslash \mathbb{H}, i_!(\mathcal{M})) = \mathcal{M}_\Gamma$$

We discuss the fundamental exact sequence in this special case. To do this we have to understand the cohomology of the boundary  $H^\bullet(\partial(\Gamma \backslash \mathbb{H}), \tilde{M})$ . We discussed the Borel-Serre compactification and saw that in this case we get this compactification if we add a circle at infinity to our picture of the quotient. But we may as well cut the cylinder at any level  $c > 1$ , i.e. we consider the level line  $\mathbb{H}(c) = \{z = x + ic | z \in \mathbb{H}\}$  and divide this level line by the action of the translation group

$$\Gamma_U = \left\{ \begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix} \mid n \in \mathbb{Z} \right\} = \left\{ \begin{pmatrix} \epsilon & n \\ 0 & \epsilon \end{pmatrix} \mid n \in \mathbb{Z}, \epsilon = \pm 1 \right\} / \{\pm \text{Id}\}.$$

But this quotient is homotopy equivalent to the cylinder

$$\Gamma_U \backslash \mathbb{H} \simeq \Gamma_U \backslash \mathbb{H}(c).$$

We apply our general consideration on cohomology of arithmetic groups to this situation and find

$$H^\bullet(\partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}}) = H^\bullet(\Gamma_U \backslash \mathbb{H}, \text{sh}_{\Gamma_U}(\mathcal{M})) = H^\bullet(\Gamma_U \backslash \mathbb{H}(c), \text{sh}_{\Gamma_U}(\mathcal{M})).$$

This cohomology is easy to compute. The group  $\Gamma_U$  is generated by the element  $T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ . It is rather clear that

$$H^0(\Gamma_U \backslash \mathbb{H}, \text{sh}_{\Gamma_U}(\mathcal{M})) = \mathcal{M}^{\Gamma_U}, H^1(\Gamma_U \backslash \mathbb{H}, \text{sh}_{\Gamma_U}(\mathcal{M})) = \mathcal{M}_{\Gamma_U} = \mathcal{M}/(\text{Id} - T)\mathcal{M}.$$

Then our fundamental exact sequence becomes (See (2.24)) fundexsq

$$0 \rightarrow \mathcal{M}^\Gamma \rightarrow \mathcal{M}^{\Gamma_U} \rightarrow \ker(\mathcal{M} \rightarrow (\mathcal{M}/(\text{Id} - S)\mathcal{M} \oplus \mathcal{M}/(\text{Id} - R)\mathcal{M})) \xrightarrow{j} \mathcal{M}/(\mathcal{M}^{\Gamma_i} \oplus \mathcal{M}^{\Gamma_\rho}) \xrightarrow{r} \mathcal{M}/(\text{Id} - T)\mathcal{M} \rightarrow \mathcal{M}_\Gamma \rightarrow 0 \quad (2.55)$$

Now it may come as a little surprise to the readers, that we can formulate a little exercise which is not entirely trivial

**Exercise:** Write down explicitly all the arrows in the above fundamental sequence

We give the answer without proof. I change notation slightly and work with the matrices

$$S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, R = \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix}$$

and we have the relation

$$RS = T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

Then  $\Gamma_i = \langle S \rangle, \Gamma_\rho = \langle R \rangle$ . The map

$$\mathcal{M}/(\mathcal{M}^{\langle S \rangle} \oplus \mathcal{M}^{\langle R \rangle}) \rightarrow \mathcal{M}/(\text{Id} - T)\mathcal{M}$$

is given by

$$m \mapsto m - Sm$$

2.1. COHOMOLOGY OF ARITHMETIC GROUPS AS COHOMOLOGY OF SHEAVES ON  $\Gamma \backslash X$ .87

We have to show that this map is well defined: If  $m \in \mathcal{M}^{<S>}$  then  $m \mapsto 0$ . If  $m \in \mathcal{M}^{<R>}$  then

$$m - Sm = m - SR^{-1}m = m - Tm$$

and this is zero in  $\mathcal{M}/(\text{Id} - T)\mathcal{M}$ .

The map

$$\ker(\mathcal{M} \rightarrow (\mathcal{M}/(\text{Id} - S)\mathcal{M} \oplus \mathcal{M}/(\text{Id} - R)\mathcal{M})) \rightarrow \mathcal{M}/(\mathcal{M}^{<S>} \oplus \mathcal{M}^{<R>})$$

is a little bit delicate. We pick an element  $m$  in the kernel, hence we can write it as

$$m = m_1 - Sm_1 = m_2 - R^{-1}m_2$$

and send  $m \mapsto m_1 - m_2$  (Here we have to use the orientation). If we modify  $m_1, m_2$  to  $m'_1 = m_1 + n_1, m'_2 = m_2 + n_2$  then  $m'_1 - m'_2$  gives the same element in  $\mathcal{M}/(\mathcal{M}^{<S>} \oplus \mathcal{M}^{<R>})$ .

This answer can only be right if  $m_1 - m_2$  goes to zero under the map  $r$ , i.e. we have to show that

$$m_1 - m_2 - S(m_1 - m_2) \in (\text{Id} - T)\mathcal{M}$$

We compute

$$\begin{aligned} m_1 - m_2 - S(m_1 - m_2) &= m - m_2 + Sm_2 = m - m_2 + R^{-1}m_2 - R^{-1}m_2 + Sm_2 = \\ &= -R^{-1}m_2 + Sm_2 = -T^{-1}Sm_2 + Sm_2 \in (\text{Id} - T)\mathcal{M} \end{aligned}$$

Finally we claim that the map  $\mathcal{M}^{<T>} \rightarrow \ker(\mathcal{M} \rightarrow (\mathcal{M}/(\text{Id} - S)\mathcal{M} \oplus \mathcal{M}/(\text{Id} - R)\mathcal{M}))$  is given by  $m \mapsto m - Sm = m - R^{-1}T^{-1}m = m - R^{-1}m$ .

There is still another element of structure. The map  $c : z \mapsto -\bar{z}$  induces an (differentiable) involution of  $\mathbb{H}$ . We put  $S_1 = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$  then  $\gamma cz = cS_1\gamma S_1^{-1}z$  and therefore  $c$  induces an involution on  $\Gamma \backslash \mathbb{H}$ . We get an isomorphism of cohomology groups

$$c^{(1)} : H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) \xrightarrow{\sim} H^1(\Gamma \backslash \mathbb{H}, c_*(\tilde{\mathcal{M}})) \quad (2.56)$$

The direct image sheaf  $c_*(\tilde{\mathcal{M}})$  is by definition the sheaf attached to the  $\Gamma$  module  $\mathcal{M}^{(S_1)}$  : This module is equal to  $\mathcal{M}$  as an abstract module, but the action is twisted by a conjugation by the above matrix  $S_1$ , i.e.

$$\gamma * m = S_1\gamma S_1^{-1}m \quad (2.57)$$

Now we assume that  $\mathcal{M}$  is actually a  $\text{Gl}_2(\mathbb{Z})$  module. Then the map  $m \rightarrow S_1m$  provides an isomorphism  $\mathcal{M}^{(S_1)} \xrightarrow{\sim} \mathcal{M}$  and hence we get an involution on the cohomology groups

$$c^\bullet : H^\bullet(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) \rightarrow H^\bullet(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) \quad (2.58)$$

We have the explicit description of the cohomology groups  $H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})$  and we can compute this involution in terms of this description. We observe that the matrix  $SS_1$  fixes the two points  $i, \rho$  and hence the two open sets  $U_i, U_\rho$  of the covering. Hence it also fixes  $\mathcal{M}^{\Gamma_i}$  and  $\mathcal{M}^{\Gamma_\rho}$  and therefore the map  $m \mapsto SS_1m$

induces an involution on  $\mathcal{M}/\mathcal{M}^{\Gamma_i} \oplus \mathcal{M}^{\Gamma_\rho} = H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})$  and this is our map  $c^{(1)}$ .

The cohomology has a  $+$  and a  $-$  eigen submodule under this involution, and

$$H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) \supset H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})_+ \oplus H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})_-$$

the sum of the two eigen modules has finite index which is a power of 2.

### Poincare' duality

We assume that our  $\Gamma$  module  $\mathcal{M}$  is a finitely generated and locally free module over  $R$ , where  $R$  is a Dedekind ring or a field. We assume  $\frac{1}{2} \in R$ . In section 6.3.11 we discuss Poincare duality in greater generality, here we consider the pairing (see 6.109)

$$H_!^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})_{\text{int}} \times H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}^\vee)_{\text{int},!} \rightarrow H_!^2(\Gamma \backslash \mathbb{H}, R) = R \quad (2.59)$$

It is clear that the involution  $c$  induces multiplication by  $-1$  on  $H_!^2(\Gamma \backslash \mathbb{H}, R)$ . On the other hand we have the decompositions of the above cohomology groups into  $\pm$  eigen modules. The pairings of the  $+, +$  parts and the  $-, -$  give zero and then we get pairings

$$\begin{aligned} H_!^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})_{\text{int},+} \times H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}^\vee)_{\text{int},!,-} &\rightarrow R \\ H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})_{\text{int},!,+} \times H_!^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}^\vee)_{\text{int},-} &\rightarrow R \end{aligned} \quad (2.60)$$

both of them are partially non degenerate.

If we have  $\mathcal{M} = \mathcal{M}^\vee$  then we get eqranks

$$\text{rank}(H_!^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})_{\text{int},+}) = \text{rank}(H_!^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})_{\text{int},-}) \quad (2.61)$$

**Final remark:** The reader may get the impression that - at least in the case  $\Gamma = \text{Sl}_2(\mathbb{Z})$ -it is easy to compute the cohomology, but the contrary is true. In the case  $\Gamma = \text{Sl}_2(\mathbb{Z})/\pm \text{Id}$  we found formulae for the rank of the cohomology groups, this seems to be a satisfactory answer, but it is not. The point is that in the next section we will introduce the Hecke operators, these Hecke operators form an algebra of endomorphisms of the cohomology groups. It is a fundamental question (see further down) to understand the cohomology as a module under the action of this Hecke algebra. It is difficult to write down the effect of a Hecke operator on a module like  $\mathcal{M}/(\mathcal{M}^{\Gamma_i} + \mathcal{M}^{\Gamma_\rho})$ . We will discuss an explicit example in section 3.3.

The situation is even worse if we consider the case  $\Gamma = \text{Gl}_2(\mathbb{Z}[i])/\{(i^\nu \text{Id})\}$ . First of all we notice that it is not possible to read off the dimensions of the individual groups  $H^i(\Gamma \backslash \mathbb{H}_3, \tilde{\mathcal{M}})$  from the complex in 2.1.3). Of course we can compute them in any given case, but our method does not give any kind of theoretical insight.

We will see later that we can prove vanishing theorems  $H^i(\tilde{\Gamma} \backslash \mathbb{H}_3, \tilde{\mathcal{M}}_{\mathbb{C}})$  for certain coefficient systems  $\tilde{\mathcal{M}}_{\mathbb{C}}$  by transcendental means. These results can not be obtained by our elementary methods.

# Chapter 3

## Hecke Operators

### 3.1 The construction of Hecke operators

We mentioned already that the cohomology and homology groups of an arithmetic group has an additional structure. We have the action of the so-called Hecke algebra. The following description of the Hecke algebra is somewhat provisional, we get a richer Hecke algebra, if we work in the adelic context (See Chapter 6 ). But the description here is more intuitive.

We start from the arithmetic group  $\Gamma \subset G(\mathbb{Q})$  and an arbitrary  $\Gamma$ -module  $\mathcal{M}$ . The module  $\mathcal{M}$  is also a module over a ring  $R$  which in the beginning may be simply  $\mathbb{Z}$ . More generally  $R$  may be the ring of integers in an algebraic number field, where we also inverted a finite number of primes.

At this point it is better to have a notation for this action

$$\Gamma \times \mathcal{M} \rightarrow \mathcal{M}, (\gamma, m) \mapsto r(\gamma)(m)$$

where now  $r : \Gamma \rightarrow \text{Aut}_R(\mathcal{M})$ .

We assume that  $\mathcal{M}$  is a module over a ring  $R$  in which we can invert the orders of the stabilizers of fixed points of elements  $\gamma \in \Gamma$ .

If we have a subgroup  $\Gamma' \subset \Gamma$  of finite index, then we constructed maps

$$\begin{aligned} \pi_{\Gamma', \Gamma}^\bullet : H^\bullet(\Gamma \backslash X, \tilde{\mathcal{M}}) &\longrightarrow H^\bullet(\Gamma' \backslash X, \tilde{\mathcal{M}}) \\ \pi_{\Gamma', \Gamma, \bullet} : H^\bullet(\Gamma \backslash X, \tilde{\mathcal{M}}) &\longrightarrow H^\bullet(\Gamma' \backslash X, \tilde{\mathcal{M}}) \end{aligned}$$

(see 2.1.1).

We pick an element  $\alpha \in G(\mathbb{Q})$ . The group

$$\Gamma(\alpha^{-1}) = \alpha^{-1}\Gamma\alpha \cap \Gamma$$

is a subgroup of finite index in  $\Gamma$  and the conjugation by  $\alpha$  induces an isomorphism

$$\text{inn}(\alpha) : \Gamma(\alpha^{-1}) \longrightarrow \Gamma(\alpha).$$

We get an isomorphism

$$j(\alpha) : \Gamma(\alpha^{-1}) \backslash X \longrightarrow \Gamma(\alpha) \backslash X$$

which is induced by the map  $x \rightarrow \alpha x$  on the space  $X$ . This yields an isomorphism of cohomology groups

$$j(\alpha)^\bullet : H^\bullet(\Gamma(\alpha^{-1}) \backslash X, \tilde{\mathcal{M}}) \longrightarrow H^\bullet(\Gamma(\alpha) \backslash X, j(\alpha)_*(\tilde{\mathcal{M}})).$$

We compute the sheaf  $j(\alpha)_*(\tilde{\mathcal{M}})$ . For a point  $x \in \Gamma(\alpha) \backslash X$  we have  $j(\alpha)_*(\tilde{\mathcal{M}})_x = \tilde{\mathcal{M}}_{x'}$  where  $j(\alpha)(x') = X$ . We have the projection  $\pi_{\Gamma(\alpha^{-1})} : X \rightarrow \Gamma(\alpha^{-1}) \backslash X$ , and the definition yields

$$(\tilde{\mathcal{M}})'_x = \left\{ s : \pi_{\Gamma(\alpha^{-1})}^{-1}(x') \rightarrow \mathcal{M} \mid s(\gamma m) = \gamma s(m) \text{ for all } \gamma \in \Gamma(\alpha^{-1}) \right\}$$

The map  $z \rightarrow \alpha z$  provides an identification  $\pi_{\Gamma(\alpha^{-1})}^{-1}(x') \xrightarrow{\sim} \pi_{\Gamma(\alpha)}^{-1}(x)$  in terms of this fibre we can describe the stalk at  $x$  as

$$j(\alpha)_*(\tilde{\mathcal{M}})_x = \left\{ s : \pi_{\Gamma(\alpha)}^{-1}(x) \rightarrow \mathcal{M} \mid s(\gamma v) = \alpha^{-1} \gamma \alpha s(v) \text{ for all } \gamma \in \Gamma(\alpha) \right\}.$$

Hence we see: We may use  $\alpha$  to define a new  $\Gamma(\alpha)$ -module  $\mathcal{M}^{(\alpha)}$ : The underlying abelian group of  $\mathcal{M}^{(\alpha)}$  is  $\mathcal{M}$  but the operation of  $\Gamma(\alpha)$  is given by

$$(\gamma, m) \longrightarrow (\alpha^{-1} \gamma \alpha) m = \gamma *_\alpha m.$$

Then the sheaf  $j(\alpha)_*(\tilde{\mathcal{M}})$  is equal to  $\tilde{\mathcal{M}}^{(\alpha)}$ . Hence we see that every element

$$u_\alpha \in \text{Hom}_{\Gamma(\alpha)}(\mathcal{M}^{(\alpha)}, \mathcal{M})$$

defines a map  $\tilde{u}_\alpha : j(\alpha)_*(\tilde{\mathcal{M}}) \rightarrow \tilde{\mathcal{M}}$ . Now we get a commuting diagram

$$\begin{array}{ccccc} H^\bullet(\Gamma(\alpha^{-1}) \backslash X, \tilde{\mathcal{M}}) & \xrightarrow{j(\alpha)^\bullet} & H^\bullet(\Gamma(\alpha) \backslash X, j(\alpha)_*(\tilde{\mathcal{M}})) & \xrightarrow{\tilde{u}_\alpha^\bullet} & H^\bullet(\Gamma(\alpha) \backslash X, \mathcal{M}) \\ \uparrow \pi^\bullet & & & & \downarrow \pi^\bullet \\ H^\bullet(\Gamma \backslash X, \tilde{\mathcal{M}}) & \xrightarrow{T(\alpha, u_\alpha)} & & & H^\bullet(\Gamma \backslash X, \mathcal{M}) \end{array} \quad (3.1)$$

where the operator on the bottom line is the Hecke operator. It depends on two data:

- a) the element  $\alpha \in G(\mathbb{Q})$ ,
- b) the choice of  $u_\alpha \in \text{Hom}_{\Gamma(\alpha)}(\mathcal{M}^{(\alpha)}, \mathcal{M})$ .

It is not difficult to show that the operator  $T(\alpha, u_\alpha)$  only depends on the double coset  $\Gamma \alpha \Gamma$ , provided we adapt the choice of  $u_\alpha$ . To be more precise if

$$\alpha_1 = \gamma_1 \alpha \gamma_2 \quad \gamma_1, \gamma_2 \in \Gamma,$$

then we have an obvious bijection

$$\Phi_{\gamma_1, \gamma_2} : \text{Hom}_{\Gamma(\alpha)}(\mathcal{M}^{(\alpha)}, \mathcal{M}) \longrightarrow \text{Hom}_{\Gamma(\alpha_1)}(\mathcal{M}^{\alpha_1}, \mathcal{M})$$

which is given by

$$\Phi_{\gamma_1, \gamma_2}(u_\alpha) = u_{\alpha_1} = \gamma_1 u_\alpha \gamma_2.$$

The reader will verify without difficulties that

$$T(\alpha, u_\alpha) = T(\alpha_1, u_{\alpha_1}).$$

(Verify this for  $H^0$  and then use some kind of resolution (See next section) )

There is are cases where we have also a rather obvious choice of  $u_\alpha$ .

In the first case our  $\Gamma$ -module  $\mathcal{M}$  is of arithmetic origin Then we have a canonical choice of an

$$u_{\alpha, \mathbb{Q}} : \mathcal{M}_{\mathbb{Q}}^{(\alpha)} \longrightarrow \mathcal{M}_{\mathbb{Q}},$$

which is given by  $m \mapsto \alpha m$ . But this morphism will not necessarily map the lattice  $\mathcal{M}^{(\alpha)}$  into  $\mathcal{M}$ . We also do not want that  $u_{\alpha, \mathbb{Q}}\mathcal{M} \subset b\mathcal{M}$ , where  $b$  is an integer  $b > 1$ . Clearly we can find a rational number  $d(\alpha) > 0$  for which

$$d(\alpha) \cdot u_{\alpha, \mathbb{Q}} : \mathcal{M}^{(\alpha)} \longrightarrow \mathcal{M} \text{ and } d(\alpha) \cdot u_{\alpha, \mathbb{Q}}(\mathcal{M}^{(\alpha)}) \not\subset b\mathcal{M} \text{ for any integer } b > 1.$$

Then  $u_\alpha = d(\alpha) \cdot u_{\alpha, \mathbb{Q}}$  is called the *normalized* choice. The canonical choice defines endomorphisms on the rational cohomology, i.e. the cohomology with coefficients in  $\mathcal{M}_{\mathbb{Q}}$  whereas the normalized Hecke operators induce endomorphism of the integral cohomology. We will resume this theme in section 6.3.2.

In the second case we assume that  $\Gamma = G(\mathbb{Z})$ , let  $\Gamma(N) \subset \Gamma$  be the full congruence subgroup mod  $N$ . Now we assume that  $\mathcal{M}$  is a  $\Gamma/\Gamma(N)$  module. Then we can pick an element  $\alpha \in G(\mathbb{Z}[\frac{1}{M}])$  where  $M$  is any integer prime to  $N$ . Since we have the homomorphism  $\mathbb{Z}[\frac{1}{M}] \rightarrow \mathbb{Z}/N\mathbb{Z}$  and our module  $\mathcal{M}$  is also a  $G(\mathbb{Z}[\frac{1}{M}])$  module. Therefore we can simply choose  $u_\alpha := m \mapsto \alpha m$ . Perhaps it is reasonable to call such a module a *module of finite type*. Hence we see that we have an essentially canonical way to define Hecke operators on tensor products of modules obtained by a rational representation and modules of finite type.

We see that we can construct many endomorphisms  $T(\alpha, u_\alpha) : H^\bullet(\Gamma \backslash X, \tilde{\mathcal{M}}) \rightarrow H^\bullet(\Gamma \backslash X, \tilde{\mathcal{M}})$ . These endomorphisms will generate an algebra

$$\mathcal{H}_{\Gamma, \tilde{\mathcal{M}}} \subset \text{End}(H^\bullet(\Gamma \backslash X, \tilde{\mathcal{M}})). \quad (3.2)$$

This is the so-called Hecke algebra. We can also define endomorphisms  $T(\alpha, u_\alpha)$  on the cohomology with compact supports, on the inner cohomology and the cohomology of the boundary. Since the operators are compatible with all the arrows in the fundamental exact sequence we denote them by the same symbol.

We now assume that  $\mathcal{M}$  is a finitely generated  $R$  module where  $R$  is the ring of integers in an algebraic number field  $K/\mathbb{Q}$ . Then our cohomology groups  $H^q(\Gamma \backslash X, \mathcal{M})$  are finitely generated  $R$ -modules with an action of the algebra  $\mathcal{H}$  on it. The Hecke algebra also acts on the inner cohomology  $H_!^q(\Gamma \backslash X, \mathcal{M})$  If we tensorize our coefficient system with any number field  $L \supset K$ , then we write  $M_L = M \otimes L$ .

We state without proof the following fundamental theorem :

He-ss

**Theorem 3.1.1.** *Let  $\mathcal{M}$  be a module of arithmetic origin. For any extension  $L/K/\mathbb{Q}$  the  $\mathcal{H}_\Gamma \otimes L$  module  $H_!^q(\Gamma \backslash X, M_L)$  is semi simple, i.e. a direct sum of irreducible  $\mathcal{H}_\Gamma$  modules.*

The proof of this theorem will be discussed in Chapter 8 ( section 8.1.7 it requires some input from analysis. We give a brief sketch. We tensorize our coefficient system by  $\mathbb{C}$ , i.e. we consider  $\mathcal{M}_L \otimes_L \mathbb{C} = \mathcal{M}_{\mathbb{C}}$ . Let us assume that  $\Gamma$  is torsion free. First of all start from the well known fact, that the cohomology  $H^\bullet(\Gamma \backslash X, \tilde{\mathcal{M}}_{\mathbb{C}})$  can be computed from the de-Rham-complex

$$H^\bullet(\Gamma \backslash X, \tilde{\mathcal{M}}_{\mathbb{C}}) = H^\bullet(\Omega^\bullet \otimes \tilde{\mathcal{M}}_{\mathbb{C}}(\Gamma \backslash X)).$$

We introduces some specific positive definite hermitian form on  $\mathcal{M}_{\mathbb{C}}$  and this allows us to define a hermitian scalar product between two  $\tilde{\mathcal{M}}_{\mathbb{C}}$  -valued  $p$ -forms

$$\langle \omega_1, \omega_2 \rangle = \int_{\Gamma \backslash X} \omega_1 \wedge * \omega_2,$$

provided one of the forms is compactly supported.

This will allow us a positive definite scalar product on  $H_1^p(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{n, \mathbb{C}})$ , We apply theorem 8.1.1 , this theorem tells us that we can find representatives  $\omega_1^h, \omega_2^h$  which are harmonic (they satisfy certain differential equations) and then

$$\langle [\omega_1], [\omega_2] \rangle := \int_{\Gamma \backslash X} \omega_1^h \wedge * \omega_2^h, \quad (3.3)$$

defines a positive definite hermitian scalar product on  $H_1^q(\Gamma \backslash X, \tilde{\mathcal{M}}_{\mathbb{C}})$ . Finally we show that  $\mathcal{H}_\Gamma$  is self adjoint with respect to this scalar product, and then semi-simplicity follows from the standard argument.

In the classical case of  $\mathrm{Gl}_2$  this is the Peterson scalar product.

**Verweis auf später?**

### 3.1.1 Commuting relations

We want to say some words concerning the structure of the Hecke algebra.

To begin we discuss the action of the Hecke-algebra on  $H^0(\Gamma \backslash X, \tilde{\mathcal{M}})$ . We have to do this since we defined the cohomology in terms of injective (or acyclic) resolutions and therefore the general results concerning the structure of the Hecke algebra can be reduced to this special case.

If we have a  $\Gamma$ -module  $\mathcal{M}$  and if we look at the diagram defining the Hecke operators, then we see that we get in degree 0

$$\begin{array}{ccc} \mathcal{M}^{\Gamma(\alpha^{-1})} & \xrightarrow{(\mathcal{M}^{(\alpha)})^{\Gamma(\alpha)} \xrightarrow{u_\alpha}} & \mathcal{M}^{\Gamma(\alpha)} \\ \uparrow & & \downarrow \\ \mathcal{M}^\Gamma & \xrightarrow{T(\alpha, u_\alpha)} & \mathcal{M}^\Gamma \end{array}$$

where the first arrow on the top line is induced by the identity map  $\mathcal{M} \rightarrow \mathcal{M}^{(\alpha)} = \mathcal{M}$  and the second by a map  $u_\alpha \in \mathrm{Hom}_{\mathbf{Ab}}(\mathcal{M}, \mathcal{M})$  which satisfies  $u_\alpha((\alpha \gamma \alpha^{-1})m) = \gamma u_\alpha(m)$ . Recalling the definition of the vertical arrow on the right, we find

$$T(\alpha, u_\alpha)(v) = \sum_{\gamma \in \Gamma/\Gamma(\alpha)} \gamma \cdot u_\alpha(v).$$

We are interested to get formulae for the product of Hecke operators, so, for instance, we would like to show that under certain assumptions on  $\alpha, \beta$  and certain adjustment of  $u_\alpha, u_\beta$  and  $u_{\alpha\beta}$  we can show

$$T(\alpha, u_\alpha) \cdot T(\beta, u_\beta) = T(\beta, u_\beta) \cdot T(\alpha, u_\alpha) = T(\alpha\beta, u_{\alpha\beta}).$$

It is easy to see what the conditions are if we want such a formula to be true. We look at what happens in  $H^0$ . For  $v \in \mathcal{M}^\Gamma$  we get

$$T(\alpha, u_\alpha) \cdot T(\beta, u_\beta)(v) = \sum_{\gamma \in \Gamma/\Gamma(\alpha)} \gamma u_\alpha \left( \sum_{\eta \in \Gamma/\Gamma(\beta)} \eta u_\beta(v) \right)$$

We assume that the following three conditions hold

(i) for each  $\eta$  we can find an  $\eta' \in \Gamma$  such that

$$\eta' \circ u_\alpha = u_\alpha \circ \eta,$$

(ii) The elements  $\gamma\eta'$  form a system of representatives for  $\Gamma/\Gamma(\alpha\beta)$

(iii)  $u_\alpha u_\beta(v) = u_\beta u_\alpha(v) = u_{\alpha\beta}(v)$ .

Then we get

$$\begin{aligned} T(\alpha, u_\alpha) \cdot T(\beta, u_\beta)(v) &= \sum_{\gamma \in \Gamma/\Gamma(\alpha)} \sum_{\eta' \in \Gamma/\Gamma(\beta)} \gamma \eta' u_\alpha u_\beta(v) = \sum_{\xi \in \Gamma/\Gamma(\alpha\beta)} \xi u_{\alpha\beta}(v) = \\ &T(\alpha\beta, u_{\alpha\beta})(v) \end{aligned}$$

We want to explain in a special case that we may have relations like the one above.

Let  $S$  be a finite set of primes, let  $|S|$  be the product of these primes. Then we define  $\Gamma_S = G(\mathbb{Z}[\frac{1}{|S|}])$ . We say that  $\alpha \in G(\mathbb{Q})$  has support in  $S$  if  $\alpha \in G(\mathbb{Z}[\frac{1}{|S|}])$ .

We take the group  $\Gamma = \mathrm{Sl}_d(\mathbb{Z})$ , and we take two disjoint sets of primes  $S_1, S_2$ . For the group  $\Gamma$  one can prove the so-called strong approximation theorem which asserts that for any natural number  $m$  the map

$$\mathrm{Sl}_d(\mathbb{Z}) \longrightarrow \mathrm{Sl}_d(\mathbb{Z}/m\mathbb{Z})$$

is surjective. (This special case is actually not so difficult. The theorem holds for many other arithmetic groups, for instance for simply connected Chevalley schemes over  $\mathrm{Spec}(\mathbb{Z})$ .)

We consider the case

$$\alpha = \begin{pmatrix} a_1 & & & \\ & a_2 & & \\ & & \ddots & \\ & & & a_d \end{pmatrix} \in \Gamma_{S_1}, \beta = \begin{pmatrix} b_1 & & & \\ & b_2 & & \\ & & \ddots & \\ & & & b_d \end{pmatrix} \in \Gamma_{S_2},$$

where  $a_d | a_{d-1} \dots | a_1$  and  $b_d | b_{d-1} \dots | b_1$ . It is clear that we can find integers  $n_1$  and  $n_2$  which are only divisible by the primes in  $S_1$  and  $S_2$  respectively, so that

$$\Gamma(n_i) \subset \Gamma(\alpha^{-1}), \Gamma(n_2) \subset \Gamma(\beta^{-1}),$$

where the  $\Gamma(n_i)$  are the full congruence subgroups  $\pmod{n_1}$  and  $n_2$  respectively. Since we have

$$\mathrm{Sl}_d(\mathbb{Z}/n\mathbb{Z}) = \mathrm{Sl}_d(\mathbb{Z}/n_1\mathbb{Z}) \times \mathrm{Sl}_d(\mathbb{Z}/n_2\mathbb{Z})$$

we get

$$\Gamma/\Gamma(\alpha^{-1}\beta^{-1}) \xrightarrow{\sim} \Gamma/\Gamma(\alpha^{-1}) \times \Gamma/\Gamma(\beta^{-1}).$$

On the right hand side we can choose representatives  $\gamma$  for  $\Gamma/\Gamma(\alpha^{-1})$  which satisfy  $\gamma \equiv \mathrm{Id} \pmod{n_2}$  and  $\eta$  for  $\Gamma/\Gamma(\beta^{-1})$  which satisfy  $\eta \equiv \mathrm{Id} \pmod{n_1}$ . Then the products  $\gamma\eta$  will form a system of representatives for  $\Gamma/\Gamma(\alpha^{-1}\beta^{-1})$ . But then we clearly have  $u_\alpha\eta = \eta u_\alpha$  and we see that (i) and (ii) above are true. Then we can put  $u_{\alpha\beta} = u_\alpha u_\beta$ .

We consider the case that our module  $\mathcal{M}$  is a  $R$ -lattice in  $\mathcal{M}_{\mathbb{Q}}$ , where  $\mathcal{M}_{\mathbb{Q}}$  is a rational  $G(\mathbb{Q})$ -module. Then we saw that we can write

$$u_\alpha = d(\alpha) \cdot \alpha$$

where  $d(\alpha)$  will be a product of powers of the primes  $p$  dividing  $n_1$  and an analogous statement can be obtained for  $\beta$  and  $n_2$ .

Since we have  $\alpha\beta = \beta\alpha$  and since clearly  $d(\alpha)d(\beta) = d(\alpha\beta)$  we also get the commutation relation.

So far we only proved this relation only for the action on  $H^0(\Gamma \backslash X, \tilde{\mathcal{M}})$ . If we want to prove it for cohomology in higher degrees, we have to choose an acyclic resolution

$$0 \longrightarrow \mathcal{M} \longrightarrow A^0 \longrightarrow A^1 \longrightarrow \dots = 0 \longrightarrow \mathcal{M} \longrightarrow A^\bullet$$

and compute the cohomology from this resolution. We have to extend the maps  $u_\alpha, u_\beta$  to this complex

$$\begin{array}{ccc} 0 \longrightarrow & \mathcal{M}^{(\alpha)} & \longrightarrow & (A^\bullet)^{(\alpha)} \\ & \downarrow u_\alpha & & \downarrow u_\alpha^{(\bullet)} \\ 0 \longrightarrow & \mathcal{M} & \longrightarrow & A^\bullet, \end{array}$$

and we have to prove that the relation

$$u_\alpha \eta u_\beta = \eta' u_\alpha u_\beta = \eta' u_{\alpha\beta}$$

also holds on the complex. Once we can prove this, it becomes clear that the commutation rule also holds in higher degrees.

We choose the special resolution

$$\begin{aligned} 0 \rightarrow \mathcal{M} \rightarrow \mathrm{Ind}^\bullet(\mathcal{M}) = \\ 0 \longrightarrow \mathcal{M} \longrightarrow \mathrm{Ind}_{\{1\}}^\Gamma \mathcal{M} \longrightarrow \mathrm{Ind}_{\{1\}}^\Gamma (\mathrm{Ind}_{\{1\}}^\Gamma \mathcal{M} / \mathcal{M}) \longrightarrow \end{aligned} \tag{3.4}$$

It is clear that it suffices to show: If we selected the  $u_\alpha, u_\beta$  in such a way that we have the condition (i), (ii) and (iii) above satisfied, then we can choose extensions  $u_\alpha, u_\beta, u_{\alpha\beta}$  to  $\mathrm{Ind}_{\{1\}}^\Gamma \mathcal{M}$  so that (i), (ii) and (iii) are also satisfied. Once we have done this we can proceed by induction.

In other words we have the diagram of  $\Gamma(\alpha)$ -modules

$$\begin{array}{ccccc} 0 & \longrightarrow & \mathcal{M}^{(\alpha)} & \longrightarrow & (\text{Ind}_{\{1\}}^{\Gamma} \mathcal{M})^{(\alpha)} \\ & & \downarrow u_{\alpha} & & \downarrow ? \\ 0 & \longrightarrow & \mathcal{M} & \longrightarrow & \text{Ind}_{\{1\}}^{\Gamma} \mathcal{M}, \end{array}$$

and we are searching for a suitable vertical arrow  $?$ . The horizontal arrows are given by (as before see (2.2)) by  $i : m \longrightarrow f_m : \{\gamma \longrightarrow \gamma m\}$ .

We make another assumption concerning our  $\alpha, \beta$ . We assume that there exists an automorphism  $\Theta$  of  $G/\mathbb{Q}$  such that  $\Theta(\alpha) = \alpha^{-1}, \Theta(\beta) = \beta^{-1}$  and  $\Theta\Gamma = \Gamma$ . This assumption is certainly fulfilled in the case above, we simply take  $\Theta(g) = {}^t g^{-1}$ , i.e. transpose inverse.

We choose representatives  $\xi_1, \dots, \xi_r$  for  $\Gamma/\Gamma(\alpha^{-1})$ , then  $\Theta\xi_1, \dots, \Theta\xi_r$  is a system of representatives for  $\Gamma/\Gamma(\alpha)$ . To define the vertical arrow  $? = u_{\alpha}^{(0)}$  we require

$$u_{\alpha}^{(0)}(f)(\Theta\xi_{\nu}) = u_{\alpha}(f(\xi_{\nu})) \quad \forall \nu = 1, \dots, r$$

and this yields a unique  $\Gamma(\alpha)$ - module isomorphism, for all  $\gamma \in \Gamma(\alpha)$  we must have

$$u_{\alpha}^{(0)}(f)(\Theta\xi_{\nu}\gamma) = u_{\alpha}(f(\xi_{\nu}\alpha^{-1}\gamma\alpha)) \quad \forall \nu = 1, \dots, r.$$

Iterating this construction gives us the  $u_{\alpha}^{(\bullet)}$ , by construction these morphisms satisfy (i), (ii), (iii). Since the complex  $H^0(\Gamma \backslash X, \text{Ind}(\mathcal{M}))_c$  computes the cohomology groups  $H^{\bullet}(\Gamma \backslash X, \mathcal{M})$  the commutation rules hold in all degrees.

HHO

### 3.1.2 More relations between Hecke operators

We look at the algebra of Hecke operators in the special case that  $G/\mathbb{Z} = \text{Gl}_2/\mathbb{Z}$ , we consider the action on  $H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})$  where  $\Gamma = \text{Sl}_2(\mathbb{Z})$ , we assume  $n$  even and  $\mathcal{M} = \mathcal{M}[-\frac{n}{2}]$ . This has the effect that the centre of  $G/\mathbb{Z}$  acts trivially on  $\mathcal{M}$  and this makes life simpler.

We attach a Hecke operator to any coset  $\Gamma\alpha\Gamma$  where  $\alpha \in \text{Gl}_2^+(\mathbb{Q})$  (i.e.  $\det(\alpha) > 0$ , we want  $\alpha$  to act on the upper half plane). Then  $\alpha$  and  $\lambda\alpha$  with  $\lambda \in \mathbb{Q}^*$  define the same operator. Hence we may assume that the matrix entries of  $\alpha$  are integers. The theorem of elementary divisors asserts that the double cosets

$$\Gamma \cdot M_n(\mathbb{Z})_{\det \neq 0} \cdot \Gamma \subset \text{Gl}_2^+(\mathbb{Q})$$

are represented by matrices of the form

$$\begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$$

where  $b \mid a$ . But here we can divide by  $b$ , and we are left with the matrix

$$\alpha = \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix}, \quad a \in \mathbb{N}.$$

We can attach a Hecke operator to this matrix provided we choose  $u_\alpha$ . We see that  $\alpha$  induces on the basis vectors

$$X^\nu Y^{n-\nu} \longrightarrow a^{\nu-n/2} \cdot X^\nu Y^{n-\nu}.$$

Hence we see that we have the following natural choice for  $u_\alpha$

$$u_\alpha : P(X, Y) \longrightarrow a^{n/2} \alpha \cdot P(X, Y).$$

(See the general discussion of the Hecke operators)

Hence we get a family of endomorphisms

$$T \left( \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix}, u \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \right) = T(a) \quad (3.5)$$

of the cohomology  $H^i(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})$ .

We have seen already that we have  $T_a T_b = T_{ab}$  if  $a, b$  are coprime.

Hence we have to investigate the local algebra  $\mathcal{H}_p$  which is generated by the

$$T_{p^r} = T \left( \begin{pmatrix} p^r & 0 \\ 0 & 1 \end{pmatrix}, u \begin{pmatrix} p^r & 0 \\ 0 & 1 \end{pmatrix} \right)$$

for the special case of the group  $\Gamma = \text{Sl}_2(\mathbb{Z})$  and the coefficient system  $\mathcal{M} = \mathcal{M}_n[-\frac{n}{2}]$ . To do this we compute the product

$$T_{p^r} \cdot T_p = T \left( \begin{pmatrix} p^r & 0 \\ 0 & 1 \end{pmatrix}, u_{\alpha_p^r} \right) \cdot T \left( \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}, u_{\alpha_p} \right)$$

where the  $u_{\alpha_p^r}$  are the canonical choices.

Again we investigate first what happens in degree zero, i.e. on  $H^0(\Gamma \backslash \mathbb{H}, \tilde{I})$  here  $I$  is any  $\Gamma$ -module.

Let  $\alpha = \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}, \xi \in H^0(\Gamma \backslash X, \tilde{I})$  then

$$T(\alpha^r, u_{\alpha^r}) T(\alpha, u_\alpha) \xi = \left( \sum_{\gamma \in \Gamma/\Gamma(\alpha^r)} \gamma u_{\alpha^r} \right) \left( \sum_{\eta \in \Gamma/\Gamma(\alpha)} \eta u_\alpha \right) (\xi)$$

We have the classical system of representatives

$$\Gamma/\Gamma(\alpha^r) = \bigcup_{j \pmod{p^r}} \begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix} \Gamma(\alpha^r) \cup \bigcup_{j' \pmod{p^{r-1}}} \begin{pmatrix} 1 & 0 \\ j'p & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \Gamma(\alpha^r),$$

and our product of Hecke operators becomes

$$\left( \sum_{j \pmod{p^r}} \begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix} \right) \left( \sum_{j' \pmod{p^{r-1}}} \begin{pmatrix} 1 & 0 \\ j'p & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} u_{\alpha^r} \right) \left( \sum_{j_1 \pmod{p}} \begin{pmatrix} 1 & j_1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} u_\alpha \right) (\xi) =$$

$$\begin{aligned}
& \left[ \sum_{j \bmod p^r, j_1 \bmod p} \begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix} u_{\alpha^r} \begin{pmatrix} 1 & j_1 \\ 0 & 1 \end{pmatrix} u_{\alpha}(\xi) \right. \\
& + \left. \sum_{j' \bmod p^{r-1}, j_1 \bmod p} \begin{pmatrix} 1 & 0 \\ j'p & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} u_{\alpha^r} \begin{pmatrix} 1 & j_1 \\ 0 & 1 \end{pmatrix} u_{\alpha}(\xi) \right] + \\
& + \left[ \sum_{j \bmod p^r} \begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix} u_{\alpha^r} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right] u_{\alpha}(\xi) + \\
& \left( \sum_{j' \bmod p^{r-1}} \begin{pmatrix} 1 & 0 \\ j'p & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} u_{\alpha^r} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} u_{\alpha}(\xi) \right)
\end{aligned}$$

Now we have to assume that  $u_{\alpha^r}$  satisfy commutation rules

$$u_{\alpha^r} u_{\alpha} = u_{\alpha^{r+1}}$$

$$u_{\alpha^r} \begin{pmatrix} 1 & j_1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & j_1 p^r \\ 0 & 1 \end{pmatrix} u_{\alpha^r} \quad (3.6)$$

$$u_{\alpha^r} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} u_{\alpha} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = c_I(p) u_{\alpha^{r-1}}$$

where  $c_I(p)$  is a non zero integer. If we exploit the first two commutation relation then we get as the sum in the first [...]

$$\begin{aligned}
& \left[ \sum_{j \bmod p^r, j_1 \bmod p} \begin{pmatrix} 1 & j + p^r j_1 \\ 0 & 1 \end{pmatrix} \right. \\
& \left. \sum_{j' \bmod p^{r-1}, j_1 \bmod p} \begin{pmatrix} 1 & 0 \\ (j' + p^{r-1} j_1)p & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right] u_{\alpha^{r+1}}(\xi) \quad (3.7) \\
& = T(p^{r+1}, u_{\alpha^{r+1}})(\xi).
\end{aligned}$$

To compute the contribution of second [...] we observe that  $w = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \in \Gamma$  and hence we have  $w\xi = \xi$ . Then the second commutation relation yields for the sum of the terms in the the second [...]

$$c_I(p) \left( \sum_{j \bmod p^r} \begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix} + \sum_{j' \bmod p^{r-1}} \begin{pmatrix} 1 & 0 \\ j'p & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right) u_{\alpha^{r-1}}(\xi). \quad (3.8)$$

We observe that for  $j \equiv 0 \pmod{p^{r-1}}$  we get

$$\begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix} u_{\alpha^{r-1}}(\xi) = u_{\alpha^{r-1}} \begin{pmatrix} 1 & \frac{j}{p^{r-1}} \\ 0 & 1 \end{pmatrix}(\xi) = u_{\alpha^{r-1}}(\xi)$$

and in case  $r > 1$  for  $j' \equiv 0 \pmod{p^{r-2}}$

$$\begin{pmatrix} 1 & 0 \\ j'p & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} u_{\alpha^{r-1}}(\xi) = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} u_{\alpha^{r-1}} \begin{pmatrix} 1 & \frac{pj'}{p^{r-1}} \\ 0 & 1 \end{pmatrix}(\xi) = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} u_{\alpha^{r-1}}(\xi) \quad (3.9)$$

here we used again (3.6) and  $\xi \in H^0(\Gamma \backslash X, \tilde{I})$ . In other words in the summation (??) the first term only depends on  $j \pmod{p^{r-1}}$  and the second only on  $j' \pmod{p^{r-2}}$ . For  $r > 1$  this yields for the second term (3.8)

$$pc_I(p) \left( \sum_{j \pmod{p^{r-1}}} \begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix} + \sum_{j' \pmod{p^{r-2}}} \begin{pmatrix} 1 & 0 \\ j'p & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right) u_{\alpha^{r-1}}(\xi) = pc_I(p) T(p^{r-1}) \xi$$

If  $r = 1$  the value for (3.8) is  $c_I(p)(p+1)u_{\alpha^0}$  and hence we get the general formula

$$T_{p^r} \cdot T_p = T_{p^{r+1}} + (p + \epsilon(p))c_I(p)T_{p^{r-1}} \quad (3.10)$$

where  $\epsilon(r) = 0$  if  $r > 1$  and  $\epsilon(r) = 1$  for  $r = 1$ .

This formula is valid for all values of  $r \geq 0$  if we put  $T_{p^{-1}} = 0$ .

We want to know what this means for the action on  $H^1(\Gamma \backslash \mathbb{H}, \mathcal{M})$ , we start again from our special resolution. (3.4). A simple calculation gives that the  $u_{\alpha^r}$  satisfy the relations (3.6) with  $c_{\mathcal{M}}(p) = p^n$ . Hence we get for the action on  $H^1(\Gamma \backslash \mathbb{H}, \mathcal{M})$

$$T_{p^r} \cdot T_p = T_{p^{r+1}} + p^{n+1}T_{p^{r-1}} + \epsilon(r)p^n T_{p^{r-1}} \quad (3.11)$$

where  $\epsilon(r) = 0$  if  $r > 1$  and  $\epsilon(r) = 1$  for  $r = 1$ .

### Interlude

We assume that a majority of the readers has seen Hecke operators in the context of modular forms and has seen formulas for these Hecke operators acting on spaces of modular forms, which look very similar to the formulas above. (See [Serre], [Hecke], [Ogg]). This is of course not accidental, in the following chapter we will discuss the Eichler-Shimura isomorphism, which provides an injection of the space of modular forms of weight  $k$  into the cohomology  $H^1(\Gamma \backslash \mathbb{H}, \mathcal{M}_{k-2} \otimes \mathbb{C})$ . (See Thm. 4.1.2). This is a Hecke-module isomorphism and this explains the relation between the classical Hecke operators and the "cohomological" Hecke operators.

There is a slight difference between the formulas here and in (HSO), the reason is that our  $T_{p^r}$  differ slightly from the classical Hecke operators. But we always have  $T_p$  defined as above is equal to  $T_p$  in (HSO).

We want to stress that in this text so far -except in the introduction- there is no mentioning of modular forms, this is intentional.

### End Interlude

This can be generalised. We choose an integer  $N > 1$  and we take as our arithmetic group the full congruence group  $\Gamma = \Gamma(N)$ . For any prime  $p \nmid N$  the  $T(\alpha, u_\alpha)$  with  $\alpha \in \text{Gl}_2^+(\mathbb{Z}[1/p])$  form a commutative subalgebra  $\mathcal{H}_p$  which is generated by  $T_p$ . This is the so called *unramified* Hecke algebra.

For  $p|N$  we can also consider the  $T(\alpha, u_\alpha)$  with  $\alpha \in \text{Gl}_2^+(\mathbb{Z}[1/p])$ . They will also generate a local algebra  $\mathcal{H}_p$  of endomorphisms in any of our cohomology groups, but this algebra will not necessarily be commutative. But if we have two different primes  $p, p_1$  then we saw that the  $\mathcal{H}_p, \mathcal{H}_{p_1}$  commute with each other. All these algebras  $\mathcal{H}_p$  have an identity element  $e_p$ , we form the algebra

$$\mathcal{H}_\Gamma = \bigotimes_p' \mathcal{H}_p$$

where the superscript indicates that a tensor  $h_f = \bigotimes_p h_p \in \mathcal{H}_\Gamma$  has a factor  $e_p$  for almost all  $p$ . This algebra acts on all our cohomology groups. We recall that the action of  $\mathcal{H}_\Gamma$  on the inner cohomology groups is semi-simple (See Thm. 3.1.1). This has important consequences, which we discuss after a brief recapitulation of the theory of semi simple modules.

## 3.2 Some results on semi-simple $\mathfrak{A}$ modules

We need a few results from the theory of algebras  $\mathfrak{A}$  acting on finite dimensional vector spaces over a field  $L$ . Let  $\bar{L}$  be an algebraic closure of  $L$ .

Let  $V$  be a finite dimensional vector space over some field  $L$  and an  $L$ -algebra  $\mathfrak{A}$  with identity acting on  $V$  by endomorphisms. We say that the action of  $\mathfrak{A}$  on  $V$  is semi-simple, if the action of  $\mathfrak{A} \otimes \bar{L}$  on  $V \otimes \bar{L}$  is semi simple and this means that any  $\mathfrak{A}$  submodule  $W \subset V \otimes \bar{L}$  has a complement, i.e. we can find an  $\mathfrak{A}$ -submodule  $W^\dagger \subset V \otimes \bar{L}$  such that  $V \otimes \bar{L} = W + W^\dagger$ .

Then it is clear that we get a decomposition indexed by a finite set  $E$

$$V \otimes \bar{L} = \bigoplus_{i \in E} W_i$$

where the  $W_i$  are irreducible submodules, i.e. they do not contain any non trivial  $\mathfrak{A}$  submodule.

In general this decomposition will not be unique. For any two  $W_i, W_j$  of these submodules we have ( Schur's lemma)

$$\text{Hom}_{\mathfrak{A}}(W_i, W_j) = \begin{cases} \bar{L} & \text{if they are isomorphic as } \mathfrak{A} \text{-modules} \\ 0 & \text{else} \end{cases}$$

We decompose the indexing set  $E = E_1 \cup E_2 \cup \dots \cup E_k$  according to isomorphism types. For any  $E_\nu$  we choose an  $\mathfrak{A}$  module  $W_{[\nu]}$  of this given isomorphism type. Then by definition

$$\text{Hom}_{\mathfrak{A}}(W_{[\nu]}, W_j) = \begin{cases} \bar{L} & \text{if } j \in E_\nu \\ 0 & \text{else} \end{cases}.$$

Now we define  $H_{[\nu]} = \text{Hom}_{\mathfrak{A}}(W_{[\nu]}, V \otimes \bar{L})$  we get an inclusion  $H_{[\nu]} \otimes W_{[\nu]}$  whose image  $X_\nu$  will be an  $\mathfrak{A}$  submodule, which is a direct sum of copies of  $W_{[\nu]}$ .

We get a direct sum decomposition

$$V \otimes \bar{L} = \bigoplus_{\nu} \bigoplus_{i \in E_\nu} W_i = \bigoplus_{\nu} X_\nu$$

then this last decomposition is easily seen to be unique, it is called the *isotypical* decomposition.

If  $V$  is a semi simple  $\mathfrak{A}$  module then any submodule  $W \subset V$  also has a complement ( this is not entirely obvious because by definition only  $W_{\bar{L}}$  has a complement in  $V_{\bar{L}}$ . But a small moment of meditation gives us that finding such a complement is the same as solving an inhomogeneous system of linear

equations over  $L$ . If this system has a solution over  $\bar{L}$  it also has a solution over  $L$ .) and hence we also can decompose the  $\mathfrak{A}$  module  $V$  into irreducibles. Again we can group the irreducibles according to isomorphism types and we get an isotypical decomposition

$$V = \bigoplus_{i \in E} U_i = \bigoplus_{\nu} \bigoplus_{i \in E_{\nu}} U_i = \bigoplus_{\nu} Y_{\nu}.$$

But an irreducible  $\mathfrak{A}$  module  $W$  may become reducible if we extend the scalars to  $\bar{L}$ . So it may happen that some of our  $U_i$  decompose further. Since it is clear that for any two  $\mathfrak{A}$ -modules  $V_1, V_2$  we have

$$\mathrm{Hom}_{\mathfrak{A}}(V_1, V_2) \otimes \bar{L} = \mathrm{Hom}_{\mathfrak{A} \otimes \bar{L}}(V_1 \otimes \bar{L}, V_2 \otimes \bar{L})$$

we know that we get the isotypical decomposition of  $V \otimes \bar{L}$  by taking the isotypical decomposition of the  $Y_{\nu} \otimes \bar{L}$  and then taking the direct sum over  $\nu$ .

Example: Let  $L_1/L$  be a finite extension of degree  $> 1$ , then we put  $\mathfrak{A} = L_1$  and  $V = L_1$ , the action is given by multiplication. Clearly  $V$  is irreducible, but  $V \otimes \bar{L}$  is not. If  $L_1/L$  is separable then the module is semisimple, otherwise it is not.

We say that the  $\mathfrak{A}$ -module  $V$  is absolutely irreducible, if the  $\mathfrak{A} \otimes \bar{L}$ -module  $V \otimes \bar{L}$  is irreducible. In this case it we have a classical result:

**Proposition.** *Let  $V$  be a semi simple  $\mathfrak{A}$  module. Then the following assertions are equivalent*

- i) *The  $\mathfrak{A}$  module  $V$  is absolutely irreducible*
- ii) *The image of  $\mathfrak{A}$  in the ring of endomorphisms is  $\mathrm{End}(V)$*
- iii) *The vector space of  $\mathfrak{A}$  endomorphisms  $\mathrm{End}_{\mathfrak{A}}(V) = L$ .*

This can be an exercise for an algebra class. Where do we need the assumption that  $V$  is semi simple?

**Proposition:** *For any semi-simple  $\mathfrak{A}$  module  $V$  we can find a finite extension  $L_1/L$  such that the irreducible sub modules in the decomposition into irreducibles are absolutely irreducible.*

Let us now assume that we have two algebras  $\mathfrak{A}, \mathfrak{B}$  acting on  $V$ , let us assume that these two operations commute i.e. for  $A \in \mathfrak{A}, B \in \mathfrak{B}, v \in V$  we have  $A(Bv) = B(Av)$ . This structure is the same as having a  $\mathfrak{A} \otimes_L \mathfrak{B}$  structure on  $V$ . Let us assume that  $\mathfrak{A}$  acts semi simply on  $V$  and let us assume that the irreducible  $\mathfrak{A}$  submodules of  $V$  are absolutely irreducible. Then it is clear that the isotypical summands  $Y_{\nu} = \bigoplus W_i$  are invariant under the action  $\mathfrak{B}$ . Now we pick an index  $i_0$  then the evaluation maps gives us a homomorphism

$$W_{i_0} \otimes \mathrm{Hom}_{\mathfrak{A}}(W_{i_0}, Y_{\nu}) \rightarrow Y_{\nu}.$$

Under our assumptions this is an isomorphism. Then we see that we get

$$V = \bigoplus_{\nu} W_{i_{\nu}} \otimes \mathrm{Hom}_{\mathfrak{A}}(W_{i_0}, Y_{\nu})$$

where  $i_\nu$  is any element in  $E_\nu$  and where  $\mathfrak{A}$  acts upon the first factor and  $\mathfrak{B}$  acts upon the second factor via the action of  $\mathfrak{B}$  on  $Y_\nu$ .

Especially we see:

**Proposition** *If  $V$  is an absolutely irreducible  $\mathfrak{A} \otimes_L \mathfrak{B}$  module then  $V \xrightarrow{\sim} X \otimes Y$ , where  $X$  (resp,  $Y$ ) is an absolutely  $\mathfrak{A}$  (resp.  $\mathfrak{B}$ ) module*

We apply these considerations to get

Heckess

**Theorem 3.2.1.** *For any  $L$  we can decompose :*

$$H_!^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{n,L}) = \bigoplus H_!^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{n,L})(\Pi_f)$$

where this is the isotypical decomposition and the  $\Pi_f$  are isomorphism classes of irreducible modules. There is a finite extension  $L/\mathbb{Q}$  such that all the isomorphism classes of isotypical modules which occur are actually absolutely irreducible.

If  $\Pi_f$  is an absolutely irreducible  $\mathcal{H}_\Gamma$  module then it is the tensor product  $\Pi_f = \otimes \pi_p$  where the  $\pi_p$  are absolutely irreducible  $\mathcal{H}_p$  modules. For  $p \nmid N$  the modules  $\pi_p$  are of dimension one (see above theorem) and they are determined by a number  $\lambda(\pi_p) \in \mathcal{O}_L$  which is the eigenvalue of  $T_p$  on  $\pi_p$ .

This follows easily from our previous considerations. The eigenvalues  $\lambda(\pi_p)$  are algebraic integers because  $T_p$  induces an endomorphism of  $H_!^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{n,\mathcal{O}_L})$  which after tensorization with  $L$  becomes the  $T_p$  on the rational vector space. The above field extension is called the splitting field of  $\mathcal{H}_\Gamma$ .

These two theorems 3.1.1 and 3.2.1 are special cases of more general results. We can start from an arbitrary reductive groups over  $\mathbb{Q}$ , arbitrary congruence subgroups  $\Gamma \subset G(\mathbb{Q})$  and arbitrary coefficient systems  $\mathcal{M}$  obtained from a rational representation of  $G/\mathbb{Q}$ , they are finitely generated modules over  $\mathbb{Z}$ . Then we can consider certain symmetric spaces  $X = G(\mathbb{R})/K_\infty$  and we have the cohomology groups  $H^\bullet(\Gamma \backslash X, \mathcal{M})$ , they are finitely generated  $\mathbb{Z}$  modules. Again we can define an action of the Hecke algebra  $\mathcal{H}_\Gamma$  and this Hecke algebra acts semi simply on the inner cohomology  $H_!^\bullet(\Gamma \backslash X, \tilde{\mathcal{M}}_\mathbb{Q})$ . (Theorem 3.1.1) Again this Hecke algebra is the tensor product of local Hecke algebras where for almost all primes these local Hecke algebras  $\mathcal{H}_p$  are polynomial rings in a certain number of variables. Then the theorem ??) is also valid in this situation. We resume this theme in Chapter 6

HEOP

### 3.2.1 Explicit formulas for the Hecke operators, a general strategy.

In the following section we discuss the Hecke operators and for numerical experiments it is useful to have an explicit procedure to compute them in a given

case. The main obstruction to get such an explicit procedure is to find an explicit way to compute the arrow  $j^\bullet(\alpha)$  in the top line of the diagram (3.1). (we change notation  $j(\alpha)$  to  $m(\alpha)$ ).

Let us assume that we have computed the cohomology groups on both sides by means of orbiconvex coverings  $\mathfrak{V} : \cup_{i \in I} V_{y_i} = \Gamma(\alpha^{-1}) \backslash X$  and  $\mathfrak{U} : \cup_{j \in J} U_{y_j} = \Gamma(\alpha) \backslash X$ .

The map  $m(\alpha)$  is an isomorphism between spaces and hence  $m(\alpha)(\mathfrak{V})$  is an acyclic covering of  $\Gamma(\alpha) \backslash X$ . This induces an identification

$$C^\bullet(\mathfrak{V}, \tilde{\mathcal{M}}) = C^\bullet(m(\alpha)(\mathfrak{V}), \tilde{\mathcal{M}}^{(\alpha)})$$

and the complex on the right hand side computes  $H^\bullet(\Gamma(\alpha) \backslash X, \tilde{\mathcal{M}}^{(\alpha)})$ . But this cohomology is also computable from the complex  $C^\bullet(\mathfrak{U}, \tilde{\mathcal{M}}^{(\alpha)})$ . We take the disjoint union of the two indexing sets  $I \cup J$  and look at the covering  $m_\alpha(\mathfrak{V}) \cup \mathfrak{U}$ . (To be precise: We consider the disjoint union  $\tilde{I} = I \cup J$  and define a covering  $\mathfrak{W}_i$  indexed by  $\tilde{I}$ . If  $i \in \tilde{I}$  then  $W_i = m(\alpha)(V_{y_i})$  and if  $i \in J$  then we put  $W_i = U_{x_i}$ . We get a diagram of Čzech complexes

$$\begin{array}{ccccc} \rightarrow & \bigoplus_{\underline{i} \in I^q} \tilde{\mathcal{M}}^{(\alpha)}(W_{\underline{i}}) & \rightarrow & \bigoplus_{\underline{i} \in I^{q+1}} \tilde{\mathcal{M}}^{(\alpha)}(W_{\underline{i}}) & \rightarrow \\ & \uparrow & & \uparrow & \\ \rightarrow & \bigoplus_{\underline{i} \in \tilde{I}^q} \tilde{\mathcal{M}}^{(\alpha)}(W_{\underline{i}}) & \rightarrow & \bigoplus_{\underline{i} \in \tilde{I}^{q+1}} \tilde{\mathcal{M}}^{(\alpha)}(W_{\underline{i}}) & \rightarrow \\ & \downarrow & & \downarrow & \\ \rightarrow & \bigoplus_{\underline{i} \in J^q} \tilde{\mathcal{M}}^{(\alpha)}(W_{\underline{i}}) & \rightarrow & \bigoplus_{\underline{i} \in J^{q+1}} \tilde{\mathcal{M}}^{(\alpha)}(W_{\underline{i}}) & \rightarrow \end{array} \quad (3.12)$$

The sets  $I^\bullet, J^\bullet$  are subsets of  $\tilde{I}^\bullet$  and the up- and down-arrows are the resulting projection maps. We know that these up- and down-arrows induce isomorphisms in cohomology.

Hence we can start from a cohomology class  $\xi \in H^q(\Gamma(\alpha) \backslash X, \tilde{\mathcal{M}}^{(\alpha)})$ , we represent it by a cocycle

$$c_\xi \in \bigoplus_{\underline{i} \in I^q} \tilde{\mathcal{M}}^{(\alpha)}(W_{\underline{i}}).$$

Then we can find a cocycle  $\tilde{c}_\xi \in \bigoplus_{\underline{i} \in \tilde{I}^q} \tilde{\mathcal{M}}^{(\alpha)}(W_{\underline{i}})$  which maps to  $c_\xi$  under the uparrow. To get this cocycle we have to do the following: our cocycle  $c_\xi$  is an array with components  $c_\xi(\underline{i})$  for  $\underline{i} \in I^q$ . We have  $d_q(c_\xi) = 0$ . To get  $\tilde{c}_\xi$  we have to give the values  $\tilde{c}_\xi(\underline{i})$  for all  $\underline{i} \in \tilde{I}^q \setminus I^q$ . We must have

$$d_q \tilde{c}_\xi = 0.$$

this yields a system of linear equations for the remaining entries. We know that this system of equations has a solution -this is then our  $\tilde{c}_\xi$  - and this solution is unique up to a boundary  $d_{q-1}(\xi')$ . Then we apply the downarrow to  $\tilde{c}_\xi$  and get a cocycle  $c_\xi^\dagger$ , which represents the same class  $\xi$  but this class is now represented by a cocycle with respect to the covering  $\mathfrak{U}$ . We apply the map  $\tilde{u}^\alpha : \tilde{\mathcal{M}}^{(\alpha)} \rightarrow \tilde{\mathcal{M}}$  to this cocycle and then we get a cocycle which represents the image of our class  $\xi$  under  $T_\alpha$ .

In the following section we discuss the explicit computation of a Hecke operator in a very specific situation. We start from our computation in section (2.1.3) and write down some  $H^\bullet(\Gamma \mathcal{X}, \tilde{\mathcal{M}})$  explicitly. On these modules we give explicit procedures to compute a Hecke operator. We get some supply of data and we look for some interesting laws or we try to verify some conjectures (see (3.67)).

### 3.3 Hecke operators for $GL_2$ :

For the rest of this chapter we discuss a very specific case. The algebraic group scheme will be  $GL_2/\mathbb{Z}$ . The symmetric space will be

$$X = GL_2(\mathbb{R})/K_\infty \text{ where } K_\infty = SO(2) \times \left\{ \begin{pmatrix} t & 0 \\ 0 & t \end{pmatrix} \mid t \in \mathbb{R}^\times, t > 0 \right\}.$$

Then the space  $X$  is the union of an upper and a lower half plane. We choose  $\tilde{\Gamma} = GL_2(\mathbb{Z})$ , then

$$\tilde{\Gamma} \backslash G_\infty / K_\infty = \Gamma \backslash \mathbb{H},$$

where  $\Gamma = SL_2(\mathbb{Z})$  and  $\mathbb{H}$  is the upper half plane. Earlier we defined the  $\Gamma$ -modules  $\mathcal{M}_n[m]$  (See 1.2.2), in the following we put  $\mathcal{M} = \mathcal{M}_n[0]$ .

We refer to Chapter 2 2.1.3. We have the two open sets  $\tilde{U}_i$ , resp.  $\tilde{U}_\rho \subset \mathbb{H}$ , they are fixed under

$$S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \text{ and } R = \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix},$$

respectively. We also will use the elements

$$T_+ = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, S_1^+ = T_- S T_-^{-1} = \begin{pmatrix} -1 & 1 \\ -2 & 1 \end{pmatrix} \in \Gamma_0^+(2)$$

$$T_- = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, S_1^- = T_+ S T_+^{-1} = \begin{pmatrix} -1 & 2 \\ -1 & 1 \end{pmatrix} \in \Gamma_0^-(2)$$

The elements  $S_1^+$  and  $S_1^-$  are elements of order four, i.e.  $(S_1^+)^2 = (S_1^-)^2 = -\text{Id}$ , the corresponding fixed points are  $\frac{i+1}{2}$  and  $i+1$  respectively. Hence  $S_1^-$  fixes the sets  $\alpha \tilde{U}_{\frac{i+1}{2}}$  and  $\tilde{U}_{i+1}$ , this is the only occurrence of a non trivial stabilizer.

#### 3.3.1 The boundary cohomology

It is easier to compute the action of the Hecke operator  $T_p$  on the cohomology of the boundary, i. e. to compute the endomorphism

$$T_p : H^1(\partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}}) \rightarrow H^1(\partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}}).$$

We know (see 2.55) that  $H^1(\partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}}) = \mathcal{M}/(1-T_+)\mathcal{M}$ , we collect some easy facts concerning this module. For  $n \geq k \geq 0$  we define the submodules

$$\mathcal{M}^{(k)} = \mathbb{Z} X^k Y^{n-k} \oplus \mathbb{Z} X^{k+1} Y^{n-k-1} \oplus \dots \oplus \mathbb{Z} X^n$$

for  $k = 0$  (resp.  $k = n$ ) we have  $\mathcal{M}^{(0)} = \mathcal{M}$  (resp.  $\mathcal{M}^{(n)} = \mathbb{Z} X^n$ ). These modules are invariant under the action of  $T_+$  we have  $(1-T_+)\mathcal{M}^{(k)} \subset \mathcal{M}^{(k+1)}$ , and  $\mathcal{M}^{(k)}/\mathcal{M}^{(k+1)} \xrightarrow{\sim} \mathbb{Z}$ . The map  $(1-T_+)$  induces a map

$$\partial_k : \mathcal{M}^{(k)}/\mathcal{M}^{(k+1)} \rightarrow \mathcal{M}^{(k+1)}/\mathcal{M}^{(k+2)}$$

which is given by multiplication with  $n-k$ . Hence it is clear that

$$\mathcal{M}/(1-T_+)\mathcal{M} = \mathbb{Z}[Y^n] \oplus \mathcal{M}^{(1)}/(1-T_+)\mathcal{M}$$

and the second summand is a finite module. The filtration of  $\mathcal{M}$  by the  $\mathcal{M}^{(k)}$  induces a filtration  $H^1(\partial(\Gamma\backslash\mathbb{H}), \tilde{\mathcal{M}})$ , we put

$$H^1(\partial(\Gamma\backslash\mathbb{H}), \tilde{\mathcal{M}})^{(k)} := \text{Im}(H^1(\partial(\Gamma\backslash\mathbb{H}), \tilde{\mathcal{M}}^{(k)}) \rightarrow H^1(\partial(\Gamma\backslash\mathbb{H}), \tilde{\mathcal{M}})) \quad (3.13)$$

Then pn1

**Proposition 3.3.1.** *For  $k > 0$  the quotient*

$$H^1(\partial(\Gamma\backslash\mathbb{H}), \tilde{\mathcal{M}})^{(k)} / H^1(\partial(\Gamma\backslash\mathbb{H}), \tilde{\mathcal{M}})^{(k+1)} \xrightarrow{\sim} \mathbb{Z}/(n-k+1)\mathbb{Z}$$

The Hecke operator  $T_p$  acts on  $H^1(\partial(\Gamma\backslash\mathbb{H}), \tilde{\mathcal{M}})^{(k)} / H^1(\partial(\Gamma\backslash\mathbb{H}), \tilde{\mathcal{M}})^{(k+1)}$  by multiplication with  $p^k + p^{n-k+1}$ . Especially we have

$$T_p[Y^n] = (p^{n+1} + 1)[Y^n]$$

*Proof.* We introduce the polynomials

$$\epsilon_k(X, Y) := X^n \frac{Y}{X} \left(\frac{Y}{X} - 1\right) \dots \left(\frac{Y}{X} - k + 1\right) = X^n \prod_{\nu=0}^{k-1} \left(\frac{Y}{X} - \nu\right) = k! X^n \binom{\frac{Y}{X}}{k} =$$

$$X^{n-k} (Y - X) \dots (Y - (k-1)X) = X^{n-k} Y^k + \dots + (-1)^k k! X^n$$

Obviously these  $\epsilon_k(X, Y)$  form a basis of  $\mathcal{M}$ . Pascal's rule for binomial coefficient says  $\binom{\frac{Y}{X}+1}{k} = \binom{\frac{Y}{X}}{k} + \binom{\frac{Y}{X}}{k-1}$  and this yields

$$T_+ \epsilon_k(X, Y) = \epsilon_k(X, X+Y) = \epsilon_k(X, Y) + k \epsilon_{k-1}(X, Y)$$

and from this we get

$$\mathcal{M}/(1 - T_+) \mathcal{M} = \mathbb{Z} \epsilon_n(X, Y) \oplus \bigoplus_{k=n-1}^0 (\mathbb{Z}/(k+1)\mathbb{Z}) \epsilon_k(X, Y) \quad (3.14)$$

this is the first assertion.

We pick a prime  $p$  and investigate the action of  $T_p$  on  $H^1(\partial(\Gamma\backslash\mathbb{H}), \tilde{\mathcal{M}})$ . We recall the definition of the Hecke operator, we start from the matrix  $\alpha = \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}$  and we consider the diagram (3.1) adapted to our situation

$$\begin{array}{ccc} H^1(\partial(\Gamma(\alpha^{-1})\backslash\mathbb{H}), \tilde{\mathcal{M}}) & \xrightarrow{j(\alpha)^{(1)}} & H^1(\partial(\Gamma(\alpha)\backslash\mathbb{H}), j(\alpha)_*(\tilde{\mathcal{M}})) \xrightarrow{\tilde{u}_\alpha^{(1)}} & H^1(\partial(\Gamma(\alpha)\backslash\mathbb{H}), \mathcal{M}) \\ \uparrow \pi^{(1)} & & & \downarrow \pi^{(1)} \\ H^1(\Gamma\backslash X, \tilde{\mathcal{M}}) & \xrightarrow{T(\alpha, u_\alpha)} & & H^1(\partial(\Gamma\backslash\mathbb{H}), \tilde{\mathcal{M}}) \end{array} \quad (3.15)$$

The group  $\Gamma(\alpha^{-1}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid c \equiv 0 \pmod{p} \right\}$ , it acts on  $\mathbb{P}^1(\mathbb{Q})$  and has two orbits which can be represented by  $\infty$  and  $0$ . The stabilisers of these two cusps are  $\Gamma_\infty = \{\pm \text{Id } T_+^\nu\}$  and  $\Gamma_0 = \{\pm \text{Id } T_-^{p\nu}\}$  respectively. Hence we get

$$H^1(\partial(\Gamma(\alpha^{-1})\backslash\mathbb{H}), \tilde{\mathcal{M}}) = \mathcal{M}/(\text{Id} - T_+) \mathcal{M} \oplus \mathcal{M}/(\text{Id} - T_-^p) \mathcal{M} \quad (3.16)$$

We identify  $H^1(\partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}}) = \mathcal{M}/(\text{Id} - T_+) \mathcal{M} \xrightarrow{w_0} \mathcal{M}/(\text{Id} - T_-) \mathcal{M}$  where the last arrow is induced by the map  $m \mapsto w_0 m$  with  $w_0 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ . Then

$$\pi^{(1)}(m) = (m, \sum_{j=0}^{p-1} \begin{pmatrix} 1 & 0 \\ j & 1 \end{pmatrix} w_0 m) \quad (3.17)$$

For the composition

$$u_\alpha^{(1)} \circ j(\alpha)^{(1)} : \mathcal{M}/(\text{Id} - T_+) \mathcal{M} \oplus \mathcal{M}/(\text{Id} - T_-^p) \mathcal{M} \rightarrow \mathcal{M}/(\text{Id} - T_+^p) \mathcal{M} \oplus \mathcal{M}/(\text{Id} - T_-) \mathcal{M}$$

is given by  $u_\alpha^{(1)} \circ j(\alpha)^{(1)}(m_\infty, m_0) \mapsto (\alpha m_\infty, \alpha m_0)$ . and  $\pi_{(1)}((n_\infty, n_0)) = n_\infty + w_0 n_0$ . This yields

$$T_p(m) = \alpha m + w_0 \alpha w_0^{-1} \sum_{j=0}^{p-1} \begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix} m$$

On  $\mathcal{M}^{(k)}/\mathcal{M}^{(k+1)}$  the element  $\begin{pmatrix} 1 & j \\ 0 & 1 \end{pmatrix}$  acts as identity,  $\alpha$  is multiplication by  $p^k$  and  $w_0 \alpha w_0^{-1}$  is multiplication  $p^{n-k}$ .  $\square$

Here we encounter a situation where the quotient  $H^1(\Gamma \backslash \mathbb{H}, \mathcal{M})_{\text{int},!} / H_!^1(\Gamma \backslash \mathbb{H}, \mathcal{M})_{\text{int}}$  may become non trivial and interesting (see(2.54)). We have to consider the exact sequence

$$0 \rightarrow H_!^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) \rightarrow H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) \xrightarrow{r} H^1(\partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}}) \quad (3.18)$$

Our cohomology groups may have some torsion  $\mathcal{T}_1 \subset H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})$ ,  $\mathcal{T}_2 \subset H^1(\partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}})$  and the map  $r$  maps the torsion  $\mathcal{T}_1$  to a submodule  $r(\mathcal{T}_1) \subset \mathcal{T}_2$ . But it will happen that  $r(r^{-1}(\mathcal{T}_2))$  is strictly larger than  $r(\mathcal{T}_1)$  this means that some non torsion elements are mapped to torsion elements under  $r$ . By definition  $H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})_{\text{int},!} = r^{-1}(\mathcal{T}_2)$  and therefore

$$H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})_{\text{int},!} / H_!^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})_{\text{int}} = r(r^{-1}(\mathcal{T}_2)) / \mathcal{T}_1 \quad (3.19)$$

This has been investigated extensively by Taiwang Deng in [14].

Let  $\pi_1 : \mathbb{H} \rightarrow \Gamma \backslash \mathbb{H}$  be the projection. We get a covering  $\Gamma \backslash \mathbb{H} = \pi_1(\tilde{U}_i) \cup \pi_1(\tilde{U}_\rho) = U_i \cap U_\rho$ . From this covering we get the Czech complex

$$\begin{aligned} 0 &\rightarrow \tilde{\mathcal{M}}(U_i) \oplus \tilde{\mathcal{M}}(U_\rho) \rightarrow \tilde{\mathcal{M}}(U_i \cap U_\rho) \rightarrow 0 \\ &\quad \downarrow ev_{\tilde{U}_i} \oplus ev_{\tilde{U}_\rho} \quad \quad \downarrow ev_{\tilde{U}_i \cap \tilde{U}_\rho} \quad (3.20) \\ \mathcal{M}^{<S>} \oplus \mathcal{M}^{<R>} &\rightarrow \mathcal{M} \rightarrow 0 \end{aligned}$$

and this gives us our formula for the first cohomology

$$H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) = \mathcal{M}/(\mathcal{M}^{<S>} \oplus \mathcal{M}^{<R>}) \quad (3.21)$$

We want to discuss the Hecke operator  $T_2$ . To do this we pass to the subgroups

$$\begin{aligned}\Gamma_0^+(2) &= \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid c \equiv 0 \pmod{2} \right\} \\ \Gamma_0^-(2) &= \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid b \equiv 0 \pmod{2} \right\}\end{aligned}\tag{3.22}$$

we form the two quotients and introduce the projection maps  $\pi_2^\pm : \mathbb{H} \rightarrow \Gamma_0^\pm(2)\backslash\mathbb{H}$ . We have an isomorphism between the spaces

$$\Gamma_0^+(2)\backslash\mathbb{H} \xrightarrow{\alpha_2} \Gamma_0^-(2)\backslash\mathbb{H}$$

which is induced from the map  $m_2 : z \mapsto \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix} z = 2z$ . This map induces an isomorphism

$$\alpha_2^\bullet : H^1(\Gamma_0^+(2)\backslash\mathbb{H}, \tilde{\mathcal{M}}) \xrightarrow{\sim} H^1(\Gamma_0^-(2)\backslash\mathbb{H}, \tilde{\mathcal{M}}^{(\alpha)}).\tag{3.23}$$

We also have the map between sheaves  $u_2 : m \mapsto \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix} m$  and the composition with this map induces a homomorphism in cohomology

$$H^1(\Gamma_0^+(2)\backslash\mathbb{H}, \tilde{\mathcal{M}}) \xrightarrow{u_2 \circ \alpha_2^\bullet} H^1(\Gamma_0^-(2)\backslash\mathbb{H}, \tilde{\mathcal{M}}).\tag{3.24}$$

This is the homomorphism we need for the computation of the Hecke operator; it is easy to define but it may be difficult in practice to compute it.

Each of the spaces  $\Gamma_0^+(2)\backslash\mathbb{H}, \Gamma_0^-(2)\backslash\mathbb{H}$  has two cusps which can be represented by the points  $\infty, 0 \in \mathbb{P}^1(\mathbb{Q})$ . The stabilizers of these two cusps in  $\Gamma_0^+(2)$  resp.  $\Gamma_0^-(2)$  are

$$\langle T_+ \rangle \times \{\pm \text{Id}\} \text{ and } \langle T_-^2 \rangle \times \{\pm \text{Id}\} \subset \Gamma_0^+(2)$$

resp.

$$\langle T_+^2 \rangle \times \{\pm \text{Id}\} \text{ and } \langle T_- \rangle \times \{\pm \text{Id}\} \subset \Gamma_0^-(2)$$

the factor  $\{\pm \text{Id}\}$  can be ignored. Then we get

We know that

$$H^1(\partial(\Gamma_0^+(2)\backslash\mathbb{H}), \tilde{\mathcal{M}}) \xrightarrow{\sim} \mathcal{M}/(\text{Id} - T_+)\mathcal{M} \oplus \mathcal{M}/(\text{Id} - T_-^2)\mathcal{M}$$

$$H^1(\partial(\Gamma_0^-(2)\backslash\mathbb{H}), \tilde{\mathcal{M}}) \xrightarrow{\sim} \mathcal{M}/(\text{Id} - T_+^2)\mathcal{M} \oplus \mathcal{M}/(\text{Id} - T_-)\mathcal{M}.$$

But now it is obvious that  $\alpha$  maps the cusp  $\infty$  to  $\infty$  and 0 to 0 and then it is also clear that for the boundary cohomology the map

$$\alpha_2^\bullet : \mathcal{M}/(\text{Id} - T_+)\mathcal{M} \oplus \mathcal{M}/(\text{Id} - T_-^2)\mathcal{M} \rightarrow \mathcal{M}/(\text{Id} - T_+^2)\mathcal{M} \oplus \mathcal{M}/(\text{Id} - T_-)\mathcal{M}$$

is simply the map which is induced by  $u_2 : \mathcal{M} \rightarrow \mathcal{M}$ . If we ignore torsion then the individual summands are infinite cyclic.

Our module  $\mathcal{M}$  is the module of homogenous polynomials of degree  $n$  in 2 variables  $X, Y$  with integer coefficients. Then the classes  $[Y^n], [X^n]$  of the polynomials  $Y^n$  (resp.)  $X^n$  are generators of  $(\mathcal{M}/(\text{Id} - T_+^\nu)\mathcal{M})/\text{tors}$  resp.  $(\mathcal{M}/(\text{Id} - T_+^\nu)\mathcal{M})/\text{tors}$  where  $\nu = 1$  resp. 2. Then we get for the homomorphism  $\alpha_2^\bullet$

$$\alpha_2^\bullet : [Y^n] \mapsto [Y^n], \alpha_2^\bullet : [X^n] \mapsto 2^n[X^n].\tag{3.25}$$

### 3.3.2 The explicit description of the cohomology

We give the explicit description of the cohomology  $H^1(\Gamma_0^+(2)\backslash\mathbb{H}, \tilde{\mathcal{M}})$ . We introduce the projections

$$\mathbb{H} \xrightarrow{\pi_2^+} \Gamma_0^+(2)\backslash\mathbb{H}; \quad \mathbb{H} \xrightarrow{\pi_2^-} \Gamma_0^-(2)\backslash\mathbb{H}$$

and get the covering  $\mathfrak{U}_2$

$$\Gamma_0^+(2)\backslash\mathbb{H} = \pi_2^+(\tilde{U}_i) \cup \pi_2^+(T_- \tilde{U}_i) \cup \pi_2^+(\tilde{U}_\rho) = \pi_2^+(\tilde{U}_i) \cup \pi_2^+(\tilde{U}_{\frac{i+1}{2}}) \cup \pi_2^+(\tilde{U}_\rho)$$

where we put  $T_- \tilde{U}_i = \tilde{U}_{\frac{i+1}{2}}$ . Our set  $\{x_\nu\}$  of indexing points is  $\mathbf{i}, \frac{i+1}{2}, \rho$ , we put  $U_{x_i}^+ = \pi_2^+(\tilde{U}_{x_i})$ . Note  $T_- \notin \Gamma_0^+(2), T_+ \in \Gamma_0^+(2)$ .

Again the cohomology is computed by the complex

$$0 \rightarrow \tilde{\mathcal{M}}(U_i^+) \oplus \tilde{\mathcal{M}}(T_- \tilde{U}_i^+) \oplus \tilde{\mathcal{M}}(U_\rho^+) \rightarrow \tilde{\mathcal{M}}(U_i^+ \cap U_\rho^+) \oplus \tilde{\mathcal{M}}(T_- \tilde{U}_i^+ \cap U_\rho^+) \rightarrow 0$$

we have to identify the terms as submodules of some  $\bigoplus \mathcal{M}$  and write down the boundary map explicitly. We have

$$\begin{array}{ccc} \tilde{\mathcal{M}}(U_i^+) \oplus \tilde{\mathcal{M}}(U_{\frac{i+1}{2}}^+) \oplus \tilde{\mathcal{M}}(U_\rho^+) & \xrightarrow{d_0} & \tilde{\mathcal{M}}(U_i^+ \cap U_\rho^+) \oplus \tilde{\mathcal{M}}(U_{\frac{i+1}{2}}^+ \cap U_\rho^+) \\ \downarrow \text{ev}_{\tilde{U}_i} \oplus \text{ev}_{T_- \tilde{U}_i} \oplus \text{ev}_{\tilde{U}_\rho} & & \downarrow \text{ev}_{\tilde{U}_i \cap \tilde{U}_\rho} \oplus \text{ev}_{\tilde{U}_i \cap T_+^{-1} \tilde{U}_\rho} \oplus \text{ev}_{T_- \tilde{U}_i \cap \tilde{U}_\rho} \\ \mathcal{M} \oplus \mathcal{M}^{\langle S_1^+ \rangle} \oplus \mathcal{M} & \xrightarrow{\bar{d}_0} & \mathcal{M} \oplus \mathcal{M} \oplus \mathcal{M} \end{array} \quad (3.26)$$

where the vertical arrows are isomorphisms. The boundary map  $\bar{d}_0$  in the bottom row is given by

$$(m_1, m_2, m_3) \mapsto (m_1 - m_3, m_1 - T_+^{-1} m_3, m_1 - m_2) = (x, y, z)$$

We may look at the (isomorphic) sub complex where  $x = z = 0$  and  $m_1 = m_2 = m_3$  then we obtain the complex

$$0 \rightarrow \mathcal{M}^{\langle S_1^+ \rangle} \rightarrow \mathcal{M} \rightarrow 0; \quad m_2 \mapsto m_2 - T_+^{-1} m_2$$

which provides an isomorphism

$$H^1(\Gamma_0^+(2)\backslash\mathbb{H}, \tilde{\mathcal{M}}) \xrightarrow{\sim} \mathcal{M}/(\text{Id} - T_+^{-1})\mathcal{M}^{\langle S_1^+ \rangle}. \quad (3.27)$$

A simple computation shows that the cohomology class represented by the class  $(x, y, z)$  is equal to the class represented by  $(0, y - x + T_+^{-1} z - z, 0)$  we write

$$[(x, y, z)] = [(0, y - x + T_+^{-1} z - z, 0)] \quad (3.28)$$

### 3.3.3 The map to the boundary cohomology

We have the restriction map for the cohomology of the boundary

$$\begin{array}{ccc}
H^1(\Gamma_0^+(2)\backslash\mathbb{H}, \tilde{\mathcal{M}}) & \xrightarrow{\sim} & \mathcal{M}/(\text{Id} - T_+^{-1})\mathcal{M}^{\langle S_1^+ \rangle} \\
\downarrow & & r^+ \oplus r^- \downarrow \\
H^1(\partial(\Gamma_0^+(2)\backslash\mathbb{H}), \tilde{\mathcal{M}}) & \xrightarrow{\sim} & \mathcal{M}/(\text{Id} - T_+)\mathcal{M} \oplus \mathcal{M}/(\text{Id} - T_-^2)\mathcal{M}
\end{array} \tag{3.29}$$

we give a formula for the second vertical arrow. We represent a class  $[m]$  by an element  $m \in \mathcal{M}$  and send  $m$  to its class in in each the two summands, respectively. This is well defined, for  $r^+$  it is obvious, while for  $r^-$  we observe that if  $m = x - T_+^{-1}x$  and  $S_1^+x = x$  then  $m = x - T_+^{-1}S_1^+x = x - T_-^2x$ .

#### Restriction and Corestriction

Now we have to give explicit formulas for the two maps  $\pi^*, \pi_*$  in the big diagram on p. 50 in Chap2.pdf. Here we should change notation: The map  $\pi$  in Chap.2 will now be denoted by :

$$\varpi_2^+ : \Gamma_0^+(2)\backslash\mathbb{H} \rightarrow \Gamma\backslash\mathbb{H} \tag{3.30}$$

We have the two complexes which compute the cohomology  $H^1(\Gamma_0^+(2)\backslash\mathbb{H}, \tilde{\mathcal{M}})$  and  $H^1(\Gamma\backslash\mathbb{H}, \tilde{\mathcal{M}})$ , and we have defined arrows between them. We realized these two complexes explicitly in (3.26) resp. (3.20) and we have

$$\begin{array}{ccc}
\tilde{\mathcal{M}}(U_i^+) \oplus \tilde{\mathcal{M}}(U_{\frac{i+1}{2}}^+) \oplus \tilde{\mathcal{M}}(U_\rho^+) & \xrightarrow{d_0} & \tilde{\mathcal{M}}(U_i^+ \cap U_\rho^+) \oplus \tilde{\mathcal{M}}(U_{\frac{i+1}{2}}^+ \cap U_\rho^+) \\
(\varpi_2^+)^{(0)} \uparrow \downarrow (\varpi_2^+)^{(0)} & & (\varpi_2^+)^{(1)} \uparrow \downarrow (\varpi_2^+)^{(1)} \\
\tilde{\mathcal{M}}(U_i) \oplus \tilde{\mathcal{M}}(U_\rho) & \xrightarrow{d_0} & \tilde{\mathcal{M}}(U_i \cap U_\rho)
\end{array} \tag{3.31}$$

and in terms of our explicit realization in diagram (3.26 ) this gives

$$\begin{array}{ccc}
\mathcal{M} \oplus \mathcal{M}^{\langle S_1 \rangle} \oplus \mathcal{M} & \xrightarrow{d_0} & \mathcal{M} \oplus \mathcal{M} \oplus \mathcal{M} \\
(\varpi_2^+)^{(0)} \uparrow \downarrow (\varpi_2^+)^{(0)} & & (\varpi_2^+)^{(1)} \uparrow \downarrow (\varpi_2^+)^{(1)} \\
\mathcal{M}^{\langle S \rangle} \oplus \mathcal{M}^{\langle R \rangle} & \xrightarrow{d_0} & \mathcal{M}
\end{array} \tag{3.32}$$

Looking at the definitions we find

$$\begin{array}{l}
(\varpi_2^+)^{(0)} : (m_1, m_2) \mapsto (m_1, T_-m_1, m_2) \\
(\varpi_2^+)^{(0)} : (m_1, m_2, m_3) \mapsto (m_1 + Sm_1 + T_-^{-1}m_2, (1 + R + R^2)m_3)
\end{array} \tag{3.33}$$

and we check easily that the composition  $(\varpi_2^+)^{(0)} \circ (\varpi_2^+)^{(0)}$  is the multiplication by 3 as it should be, since this is the index of  $\Gamma_0(2)^+$  in  $\Gamma$ .

For the two arrows in degree one we find

$$\begin{aligned} (\varpi_2^+)^{(1)} : m &\mapsto (m, Sm, T_-m) \\ (\varpi_2^+)^{(1)}_{(1)} : (m_1, m_2, m_3) &\mapsto (m_1 + Sm_2 + T_-^{-1}m_3) \end{aligned} \quad (3.34)$$

We apply equation (3.28) and we see that  $(\varpi_2^+)^{(1)}(m)$  is represented by

$$[(\varpi_2^+)^{(1)}(m)] = [0, Sm + T_+^{-1}T_-m - m - T_-m, 0] \quad (3.35)$$

We do the same calculation for  $\Gamma_0^-(2)$ . As before we start from a covering

$$\Gamma_0^-(2) \backslash \mathbb{H} = \pi_2^-(\tilde{U}_1) \cup \pi_2^-(T_+\tilde{U}_1) \cup \pi_2^-(\tilde{U}_\rho) = \pi_2^-(\tilde{U}_1) \cup \pi_2^-(\tilde{U}_{i+1}) \cup \pi_2^-(\tilde{U}_\rho)$$

and as before we put  $U_{y_\nu}^- = \pi_2^-(\tilde{U}_{y_\nu})$ . In this case  $\tilde{U}_{i+1} = T_+\tilde{U}_1$  is fixed by  $S_1^- = \begin{pmatrix} -1 & 2 \\ -1 & 1 \end{pmatrix} \in \Gamma_0^-(2)$  and we get a diagram for the Czech complex

$$\begin{array}{ccc} \tilde{\mathcal{M}}(U_{\mathbf{i}}^-) \oplus \tilde{\mathcal{M}}(U_{\mathbf{i}+1}^-) \oplus \tilde{\mathcal{M}}(U_\rho^-) & \xrightarrow{d_0} & \tilde{\mathcal{M}}(U_{\mathbf{i}}^- \cap U_\rho^-) \oplus \tilde{\mathcal{M}}(U_{\mathbf{i}+1}^- \cap U_\rho^-) \\ ev_{\tilde{U}_1} \oplus ev_{\tilde{U}_{i+1}} \downarrow \oplus ev_{\tilde{U}_\rho} & & ev_{\tilde{U}_1 \cap \tilde{U}_\rho} \oplus ev_{\tilde{U}_1 \cap T_-^{-1}\tilde{U}_\rho} \downarrow \oplus ev_{\tilde{U}_{i+1} \cap \tilde{U}_\rho} \\ \mathcal{M} \oplus \mathcal{M}^{\langle S_1^- \rangle} \oplus \mathcal{M} & \xrightarrow{\bar{d}_0} & \mathcal{M} \oplus \mathcal{M} \oplus \mathcal{M} \end{array} \quad (3.36)$$

Again we can modify this complex and get

$$H^1(\Gamma_0^-(2) \backslash \mathbb{H}, \tilde{\mathcal{M}}) \xrightarrow{\sim} \mathcal{M}/(\text{Id} - T_-^{-1})\mathcal{M}^{\langle S_1^- \rangle}. \quad (3.37)$$

We compute the arrows  $(\varpi_2^-)^*$ ,  $(\varpi_2^-)_*$  in degree one

$$\begin{aligned} (\varpi_2^-)^{(1)} : m &\mapsto (m, Sm, T_+m), \\ (\varpi_2^-)^{(1)}_{(1)} : (m_1, m_2, m_3) &\mapsto (m_1 + Sm_2 + T_+^{-1}m_3). \end{aligned} \quad (3.38)$$

### The computation of $\alpha_2^\bullet$ .

We recall our isomorphism  $\alpha$  between the spaces and the resulting isomorphism (3.23). The identity map of the module  $\mathcal{M}$  and the isomorphism  $\alpha$  on the space identifies the two complexes

$$\begin{array}{ccc} \tilde{\mathcal{M}}(U_{\mathbf{i}}^+) \oplus \tilde{\mathcal{M}}(U_{\frac{\mathbf{i}+1}{2}}^+) \oplus \tilde{\mathcal{M}}(U_\rho^+) & \xrightarrow{d_0} & \tilde{\mathcal{M}}(U_{\mathbf{i}}^+ \cap U_\rho^+) \oplus \tilde{\mathcal{M}}(U_{\frac{\mathbf{i}+1}{2}}^+ \cap U_\rho^+) \\ \tilde{\mathcal{M}}^{(\alpha)}(\alpha(U_{\mathbf{i}}^+)) \oplus \tilde{\mathcal{M}}^{(\alpha)}(\alpha(U_{\frac{\mathbf{i}+1}{2}}^+)) \oplus \tilde{\mathcal{M}}^{(\alpha)}(\alpha(U_\rho^+)) & \xrightarrow{d_0} & \tilde{\mathcal{M}}^{(\alpha)}(\alpha(U_{\mathbf{i}}^+ \cap U_\rho^+)) \oplus \tilde{\mathcal{M}}^{(\alpha)}(\alpha(U_{\frac{\mathbf{i}+1}{2}}^+ \cap U_\rho^+)) \end{array} \quad (3.39)$$

and if we consider their explicit realization then this identification is given by the equality of  $\mathbb{Z}$  modules  $\mathcal{M} = \mathcal{M}^{(\alpha)}$ . This equality of complexes expresses

the identification (3.23). We can compute the cohomology  $H^1(\Gamma_0^-(2)\backslash\mathbb{H}, \tilde{\mathcal{M}}^{(\alpha)})$  from any of the two coverings

$$\begin{aligned} \Gamma_0^-(2)\backslash\mathbb{H} &= \alpha(U_{\mathbf{i}}^+) \cup \alpha(U_{\frac{\mathbf{i}+1}{2}}^+) \cup \alpha(U_{\rho}^+) = U_{x_1} \cup U_{x_2} \cup U_{x_3} \\ \text{and} \\ \Gamma_0^-(2)\backslash\mathbb{H} &= U_{\mathbf{i}}^- \cup U_{\mathbf{i}+1}^- \cup U_{\rho}^- = U_{x_4} \cup U_{x_5} \cup U_{x_6}. \end{aligned} \quad (3.40)$$

We have to pick a class  $\xi \in H^1(\Gamma_0^-(2)\backslash\mathbb{H}, \tilde{\mathcal{M}}^{(\alpha)})$  and represent it by a cocycle

$$c_{\xi} \in \bigoplus_{1 \leq i < j \leq 3} \tilde{\mathcal{M}}^{(\alpha)}(U_{x_i} \cap U_{x_j})$$

(The cocycle condition is empty since  $U_{x_1} \cap U_{x_2} \cap U_{x_3} = \emptyset$ .)

Then we have to produce a cocycle

$$c_{\xi}^{\alpha} \in \bigoplus_{4 \leq i < j \leq 6} \tilde{\mathcal{M}}^{(\alpha)}(U_{x_i} \cap U_{x_j})$$

which represents the same class.

To get this cocycle we write down the three complexes

$$\begin{array}{ccc} \bigoplus_{1 \leq i < j \leq 3} \tilde{\mathcal{M}}^{(\alpha)}(U_{x_i} \cap U_{x_j}) & \rightarrow & 0 \\ \uparrow & & \\ \bigoplus_{1 \leq i < j \leq 6} \tilde{\mathcal{M}}^{(\alpha)}(U_{x_i} \cap U_{x_j}) & \rightarrow & \bigoplus_{1 \leq i < j < k \leq 6} \tilde{\mathcal{M}}^{(\alpha)}(U_{x_i} \cap U_{x_j} \cap U_{x_k}) \\ \downarrow & & \\ \bigoplus_{4 \leq i < j \leq 6} \tilde{\mathcal{M}}^{(\alpha)}(U_{x_i} \cap U_{x_j}) & \rightarrow & 0 \end{array} \quad (3.41)$$

for our cocycle  $c_{\xi}$  we find a cocycle  $c_{\xi}^{\dagger}$  in the complex in the middle which maps to  $c_{\xi}$  under the upwards arrow and this cocycle is unique up to a coboundary. Then we project it down by the downwards arrow, i.e. we only take its  $4 \leq i < j \leq 6$  components, and this is our cocycle  $c_{\xi}^{(\alpha)}$ .

We write down these complexes explicitly. For any pair  $\underline{i} = (i, j), i < j$  of indices we have to compute the set  $\mathcal{F}_{\underline{i}}$ . We drew some pictures and from these pictures we get (modulo errors) the following list (of lists):

$$\begin{array}{cccc} \mathcal{F}_{1,2} = \emptyset & \mathcal{F}_{1,3} = \{\text{Id}, T_+^{-2}\} & \mathcal{F}_{1,4} = \{\text{Id}\} & \mathcal{F}_{1,5} = \{\text{Id}, T_+^{-2}\} \\ \mathcal{F}_{1,6} := \{\text{Id}, T_-^{-1}\} & \mathcal{F}_{2,3} = \{\text{Id}\} & \mathcal{F}_{2,4} = \{\text{Id}, T_-\} & \mathcal{F}_{2,5} = \{\text{Id}\} \\ \mathcal{F}_{2,6} = \{\text{Id}\} & \mathcal{F}_{3,4} = \{\text{Id}, T_+^2\} & \mathcal{F}_{3,5} = \{\text{Id}\} & \mathcal{F}_{3,6} = \{\text{Id}, S_1^-\} \\ \mathcal{F}_{4,5} = \emptyset & \mathcal{F}_{4,6} = \{\text{Id}, T_-^{-1}\} & \mathcal{F}_{5,6} = \{\text{Id}\} & \end{array} \quad (3.42)$$

Now we have to follow the rules in the first section and we can write down an explicit version of the diagram (3.41). Here we have to be very careful,

because the sets  $\tilde{U}_{\tilde{x}_2}, \tilde{U}_{\tilde{x}_5}$  have the non-trivial stabilizer  $\langle S_1^- \rangle$  and we have to keep track of the action of  $\Gamma_{\tilde{x}_{2,5}}$ : the set  $\mathcal{F}_{i,j} \subset \Gamma_{\tilde{x}_i} \backslash \Gamma / \Gamma_{\tilde{x}_j}$ . Therefore we have to replace the group elements  $\gamma \in \mathcal{F}_{i,j}$  by sets  $\Gamma_{\tilde{x}_i} \gamma \Gamma_{\tilde{x}_j}$ . In the list above we have taken representatives.

$$\begin{array}{ccc}
\bigoplus_{1 \leq i < j \leq 3} \bigoplus_{\gamma \in \mathcal{F}_{i,j}} (\mathcal{M}^{(\alpha)})^{\Gamma_{i,j,\gamma}} & \rightarrow & 0 \\
\uparrow & & \\
\bigoplus_{1 \leq i < j \leq 6} \bigoplus_{\gamma \in \mathcal{F}_{i,j}} (\mathcal{M}^{(\alpha)})^{\Gamma_{i,j,\gamma}} & \rightarrow & \bigoplus_{1 \leq i < j < k \leq 6} \bigoplus_{\gamma \in \mathcal{F}_{i,j,k}} (\mathcal{M}^{(\alpha)})^{\Gamma_{i,j,k,\gamma}} \\
\downarrow & & \\
\bigoplus_{4 \leq i < j \leq 6} \bigoplus_{\gamma \in \mathcal{F}_{i,j}} (\mathcal{M}^{(\alpha)})^{\Gamma_{i,j,\gamma}} & \rightarrow & 0
\end{array} \tag{3.43}$$

Here we have to interpret this diagram. The module  $\mathcal{M}^{(\alpha)}$  is equal to  $\mathcal{M}$  as an abstract module, but an element  $\gamma \in \Gamma_0^-(2)$  acts by the twisted action (See ChapII, 2.2)

$$m \mapsto \gamma *_\alpha m = \alpha^{-1} \gamma \alpha * m$$

here the  $*$  denotes the original action. Hence we have to take the invariants  $(\mathcal{M}^{(\alpha)})^{\Gamma_{i,j,\gamma}}$  with respect to this twisted action. In our special situation this has very little effect since almost all the  $\Gamma_{i,j,\gamma}$  are trivial, except for the intersection  $\alpha(\tilde{U}_{\frac{i+1}{2}}) \cap \tilde{U}_i$  in which case  $\Gamma_{i,j,\gamma} = \langle S_1^- \rangle$ . Hence

$$(\mathcal{M}^{(\alpha)})^{\langle S_1^- \rangle} = \mathcal{M}^{\langle S_1^+ \rangle}.$$

Each of the complexes in (3.43) compute the cohomology group  $H^1(\Gamma_0^-(2) \backslash \mathbb{H}, \tilde{\mathcal{M}})$  and the diagram gives us a formula for the isomorphism in (3.23). To get  $u_\alpha^\bullet$  in (3.23) we apply the multiplication  $m_2: m \mapsto \alpha m$  to the complex in the middle and the bottom. Then the cocycle  $c_\xi^\alpha$  is now an element in  $\bigoplus \tilde{\mathcal{M}}^{(\alpha)}$  and  $\alpha c_\xi^\alpha$  represents the cohomology class  $u_\alpha^\bullet(\xi) \in H^1(\Gamma_0^-(2) \backslash \mathbb{H}, \tilde{\mathcal{M}})$ .

Now it is clear how we can compute the Hecke operator

$$T_2 = T \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix} : \mathcal{M} / (\mathcal{M}^{\langle S \rangle} \oplus \mathcal{M}^{\langle R \rangle}) \rightarrow \mathcal{M} / (\mathcal{M}^{\langle S \rangle} \oplus \mathcal{M}^{\langle R \rangle})$$

We pick a representative  $m \in \mathcal{M}$  of the cohomology class. We apply  $(\varpi_2^+)^{(1)}$  in the diagram (3.32) to it and this gives the element  $(Sm, m, T_- m) = c_\xi$ . We apply the above process to compute  $c_\xi^{(\alpha)}$ . Then  $\alpha c_\xi^{(\alpha)} = (m_1, m_2, m_3)$  is an element in  $\tilde{\mathcal{M}}(U_1^- \cap U_\rho^-) \oplus \tilde{\mathcal{M}}(U_{i+1}^- \cap U_\rho^-)$  and this module is identified with  $\mathcal{M} \oplus \mathcal{M} \oplus \mathcal{M}$  by the vertical arrow in (3.36). To this element we apply the trace

$$(\varpi_2^-)_{(1)}(m_1, m_2, m_3) = m_1 + m_2 + T_+^{-1} m_3$$

and the latter element in  $\mathcal{M}$  represents the class  $T_2([m])$ .

We have written a computer program which for a given  $\mathcal{M} = \mathcal{M}_n$ , i.e. for a given even positive integer  $n$ , computes the module  $H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})$  and the endomorphism  $T_2$  on it.

Looking our data we discovered the following (surprising?) fact: We consider the isomorphism in equation (3.23). We have the explicit description of the cohomology in (3.27)

$$H^1(\Gamma_0^+(2) \backslash \mathbb{H}, \tilde{\mathcal{M}}) \xrightarrow{\sim} \mathcal{M}/(\text{Id} - T_+^{-1})\mathcal{M}^{\langle S_1^+ \rangle}$$

and

$$H^1(\Gamma_0^-(2) \backslash \mathbb{H}, \tilde{\mathcal{M}}^{(\alpha)}) \xrightarrow{\sim} \mathcal{M}/(\text{Id} - T_-^{-1})(\mathcal{M}^{(\alpha)})^{\langle S_1^- \rangle}$$

We know that we may represent any cohomology class by a cocycle

$$c_\xi = (0, c_\xi, 0) \in \mathcal{M}^{(\alpha)}(\pi_2^-(\alpha(U_i) \cap \alpha(U_\rho))) \oplus \mathcal{M}^{(\alpha)}(\pi_2^-(\alpha(U_i) \cap \alpha(T_+^{-1}U_\rho))) \oplus \mathcal{M}^{(\alpha)}(\pi_2^-(\alpha(U_{\frac{i+1}{2}}) \cap \alpha(T_+^{-1}U_\rho)))$$

so it is non zero only in the middle component and then it is simply an element in  $\mathcal{M}$ . If we now look at our data, then it seems to be so that  $c_\xi^{(\alpha)}$  is also non zero only in the middle, hence

$$c_\xi^{(\alpha)} \in (0, c'_\xi, 0) \in 0 \oplus \mathcal{M}^{(\alpha)}(\pi_2^-(U_i \cap T_-^{-1}U_\rho)) \oplus 0$$

hence it is also in  $\mathcal{M}^{(\alpha)}$  and then our data seem to suggest that

$$c'_\xi = c_\xi$$

Hence we see that the homomorphism in equation (3.24) is simply given by

$$X^\nu Y^{n-\nu} \mapsto 2^\nu X^\nu Y^{n-\nu}.$$

Is there a kind of homotopy argument (- 2 moves continuously to 1)-, which explains this?

We get an explicit formula for the Hecke operator  $T_2$ : We pick an element  $m \in \mathcal{M}$  representing the class  $[m]$ . We send it by  $(\varpi_2^+)^{(1)}$  to  $H^1(\Gamma_0^+(2) \backslash \mathbb{H}, \tilde{\mathcal{M}})$ , i.e.

$$(\varpi_2^+)^{(1)} : m \mapsto (m, Sm, T_-m) \quad (3.44)$$

We modify it so that the first and the third entry become zero see( 3.28)

$$[(m, Sm, T_-m)] = [(0, Sm - m + T_+^{-1}T_-m - T_-m, 0)] \quad (3.45)$$

To the entry in the middle we apply  $M_2 = \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}$  and then apply  $(\varpi_2^-)_{(1)}$  and

get

$$T_2([m]) = [S \cdot M_2(Sm - m + T_+^{-1}T_-m - T_-m)] \quad (3.46)$$

Eisn

### 3.3.4 The first interesting example

We give an explicit formula for the cohomology in the case of  $\mathcal{M} = \mathcal{M}_{10}$ . We define the sub-module

$$\mathcal{M}^{\text{tr}} = \bigoplus_{\nu=0}^5 \mathbb{Z}Y^{10-\nu}X^\nu$$

and we have the truncation operator

$$\text{trunc} : Y^{10-\nu}X^\nu \mapsto \begin{cases} Y^{10-\nu}X^\nu & \text{if } \nu \leq 5, \\ (-1)^{\nu+1}Y^\nu X^{10-\nu} & \text{else,} \end{cases}$$

which identifies the quotient module  $\mathcal{M}/\mathcal{M}^{\langle S \rangle}$  to  $\mathcal{M}^{\text{tr}}$ . To get the cohomology we have to divide by the relations coming from  $\mathcal{M}^{\langle R \rangle}$ , i.e. we have to divide by the submodule  $\text{trunc}(\mathcal{M}^{\langle R \rangle})$ . The module of these relations is generated by

$$\begin{aligned} R_1 &= 10Y^9X + 20Y^7X^3 + Y^5X^5 \\ R_2 &= 9Y^8X^2 - 36Y^7X^3 + 14Y^6X^4 - 45Y^5X^5 \\ R_3 &= 8Y^7X^3 + 10Y^5X^5 \end{aligned}$$

and then

$$H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) = \bigoplus_{\nu=0}^5 \mathbb{Z}Y^{10-\nu}X^\nu / \{R_1, R_2, R_3\} \quad (3.47)$$

We simplify the notation and put  $e_\nu = Y^\nu X^{n-\nu}$ . Using  $R_1$  we can eliminate  $e_5 = -10e_9 - 20e_7$  and then

$$H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) = \bigoplus_{\nu=10}^{\nu=6} \mathbb{Z}e_\nu / \{-50e_9 + 9e_8 - 96e_7 + 14e_6, -100e_9 - 192e_7\} \quad (3.48)$$

introduce a new basis  $\{f_{10}, f_9, f_8, f_7, f_6, f_5\}$  of the  $\mathbb{Z}$  module  $\mathcal{M}^{\text{tr}}$ :

$$f_{10} = e_{10}; f_8 = -2e_8 - 3e_6; f_6 = 9e_8 + 14e_6 \quad (3.49)$$

$$f_9 = -12e_9 - 23e_7; f_7 = 25e_9 + 48e_7; f_5 = 10e_9 + 20e_7 + e_5$$

and hence in the quotient we get  $\bar{f}_5 = 0$  and  $2\bar{f}_7 = \bar{f}_6$  and therefore

$$H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) = \mathbb{Z}\bar{f}_{10} \oplus \mathbb{Z}\bar{f}_9 \oplus \mathbb{Z}\bar{f}_8 \oplus \mathbb{Z}/(4)\bar{f}_7 \quad (3.50)$$

(If we invert the primes  $< 12$  then we we can work with  $e_{10}, e_9, e_8$  and in cohomology  $e_6 = -\frac{9}{14}e_8, e_5 = \frac{5}{12}e_9, e_7 = -\frac{25}{48}e_9$ .)

If we can apply the above procedure to compute the action of  $T_2$  on cohomology we get the following matrix for  $T_2$ :

$$T_2 = \begin{pmatrix} 2049 & -68040 & 0 & 0 \\ 0 & -24 & 0 & 0 \\ 0 & 0 & -24 & 0 \\ 0 & 0 & 0 & 2 \end{pmatrix} \quad (3.51)$$

Hence we see that it is non trivial on the torsion subgroup. If we divide by the torsion then the matrix reduces to a (3,3)-matrix and this matrix gives us the endomorphism on the "integral" cohomology which is defined in generality by

$$H_{\text{int}}^{\bullet}(\Gamma \backslash X, \tilde{\mathcal{M}}) = H^{\bullet}(\Gamma \backslash X, \tilde{\mathcal{M}})/\text{tors} \subset H^{\bullet}(\Gamma \backslash X, \tilde{\mathcal{M}}_{\mathbb{Q}}) \quad (3.52)$$

here we should be careful: the functor  $H^{\bullet} \rightarrow H_{\text{int}}^{\bullet}$  is not exact. In our case we get (perhaps up to a little piece of 2-torsion) exact sequences of Hecke modules

$$\begin{array}{ccccccc} 0 & \rightarrow & \mathbb{Z}f_9 \oplus \mathbb{Z}f_8 & \rightarrow & \mathbb{Z}f_{10} \oplus \mathbb{Z}f_9 \oplus \mathbb{Z}f_8 & \xrightarrow{r} & \mathbb{Z}\bar{f}_{10} \rightarrow 0 \\ & & \parallel & & \parallel & & \parallel \\ 0 & \rightarrow & H_{\text{int},!}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) & \rightarrow & H_{\text{int}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) & \xrightarrow{r} & H_{\text{int}}^1(\partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}}) \rightarrow 0 \end{array} \quad (3.53)$$

where  $T_2(\bar{f}_{10}) = (2^{11} + 1)\bar{f}_{10}$ . If we tensor by  $\mathbb{Q}$  then we can find an unique element (the Eisenstein class)  $f_{10}^{\dagger} \in H_{\text{int}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) \otimes \mathbb{Q}$  which maps to  $\bar{f}_{10}$  and which satisfies  $T_2(f_{10}^{\dagger}) = (2^{11} + 1)f_{10}^{\dagger}$ . This element is not necessarily integral, in our case an easy computation shows that  $f^{\dagger} \notin H_{\text{int}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})$ . but  $691f^{\dagger} \in H_{\text{int}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})$ . This means that 691 is the denominator of  $f_{10}^{\dagger}$ , i.e. 691 is the denominator of the Eisenstein class  $f_{10}^{\dagger}$ .

Hence we see that

$$H_{\text{int}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) \supset H_{\text{int},!}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) \oplus \mathbb{Z}691f^{\dagger}$$

the quotient of these modules is isomorphic to  $\mathbb{Z}/691\mathbb{Z}$

The exact sequence  $\mathcal{X}_{10}$  in (3.53) is an exact sequence of modules for the Hecke algebra  $C \supset \mathbb{Z}[T_2]$  and hence it yields an element

$$[\mathcal{X}_{10}] \in \text{Ext}_{\mathcal{H}}^1(\mathbb{Z}f_{10}, H_{\text{int},!}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})), \quad (3.54)$$

and an easy calculation shows that this  $\text{Ext}^1$  group is cyclic of order 691 and that it is generated by  $\mathcal{X}_{10}$ .

Of course we should also look at the action of the full Hecke algebra  $\mathcal{H}$  on these cohomology groups. It turns out that for any prime  $p$  the Hecke operator  $T_p$  acts by the eigenvalue  $p^{11} + 1$  (See proposition 3.3.1). We will also see that a simple argument using Poincare duality and an adjointness of the Hecke operators shows that  $T_p$  acts by multiplication by a scalar  $\tau(p)$ . Then we can conclude (See below)

For all primes  $p$  we have

$$\tau(p) \equiv p^{11} + 1 \pmod{691}$$

### Interlude: Ramanujan's $\Delta(z)$

We want to stress that the previous considerations are purely algebraic and combinatorial, no analysis is involved. In the next chapter we will use analytic methods, especially we will use the results from the theory modular forms.

Ramanujan detected the function

$$\Delta(z) = e^{2\pi iz} \prod_{n=1}^{\infty} (1 - e^{2\pi inz})^{24}$$

this is the unique (up to a non zero scalar ) cusp form of weight 12 for  $Sl_2(\mathbb{Z})$ , (See [?]). We can expand

$$\Delta(z) = e^{2\pi iz} - 24e^{4\pi iz} + 252e^{6\pi iz} + \dots + a_n e^{2n\pi iz} + \dots$$

and the numbers  $\tau(n)$  are the coefficients we get by expanding. They satisfy (conjectured by Ramanujan)

$$a_{n_1 n_2} = a_{n_1} a_{n_2} \text{ if } n_1, n_2 \text{ are coprime; } a_{p^r} = \tau(p) \text{ if } p \text{ is a prime and } r \geq 2$$

These recursion formulas for the coefficients of the expansion were proved by Mordell (essentially by using Hecke operators) and later by Hecke in a more general framework.

In the next section we discuss the Eichler-Shimura isomorphism and in this special case it implies

$$\text{For any prime } p \text{ we have } a_p = \tau(p)$$

But for Ramanujan the numbers  $\tau(n) = a_n$  where simply the expansion coefficients and he has proved the famous congruence  $\tau(p) \equiv p^{11} + 1 \pmod{691}$ .

**Other congruences**

It is easy to check that  $H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})$  and  $H_c^2(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})$  do not have 5 or 7 torsion. Therefore we have we have (Prop. 3.3.1, see )

$$\mathbb{Z}/10\mathbb{Z}\epsilon_9(X, Y) \oplus \mathbb{Z}/5\mathbb{Z}\epsilon_4(X, Y) \oplus \mathbb{Z}/7\mathbb{Z}\epsilon_6(X, Y) \subset r(r^{-1}(\mathcal{T}_2))/\mathcal{T}_1 \quad (3.55)$$

and this implies the well known congruences

$$\tau(p) \equiv p^{10} + p \equiv p^6 + p^5 \pmod{5}; \tau(p) \equiv p^7 + p^5 \pmod{7} \quad (3.56)$$

[82] [14] These congruences are called congruences of *local origin* whereas the congruence mod 691 is a congruence of *global origin*.

**End of interlude**

We can go one step further and reduce mod 691. Since there is at most 2 torsion we get an exact sequence of Hecke-modules

$$0 \rightarrow H_{\text{int},1}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}} \otimes \mathbb{F}_{691}) \rightarrow H_{\text{int}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}} \otimes \mathbb{F}_{691}) \xrightarrow{r} H_{\text{int}}^1(\partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}} \otimes \mathbb{F}_{691}) \rightarrow 0. \quad (3.57)$$

The matrix giving the Hecke operator mod 691 becomes

$$T_2 = \begin{pmatrix} 667 & 369 & 0 \\ 0 & 667 & 0 \\ 0 & 0 & 667 \end{pmatrix} \quad (3.58)$$

This implies that the extension class  $[\mathcal{X}_{10} \otimes \mathbb{F}_{691}]$  is a element of order 691. This implies that 691 divides the order of  $[\mathcal{X}_{10}]$  and hence divides the order of the denominator of the Eisenstein class.

### The general case

Now we describe the general case  $\mathcal{M} = \mathcal{M}_n$  where  $n$  is an even integer. We define  $\mathcal{M}^{\text{tr}}$  as above, if  $n/2$  is even, then we leave out the summand  $X^{n/2}Y^{n/2}$ , we get

$$\mathcal{M}^{\text{tr}} = \mathcal{M}/\mathcal{M}^{\langle S \rangle}.$$

This gives us for the cohomology and the restriction to the boundary cohomology

$$\begin{array}{ccc} H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) & \xrightarrow{\sim} & \mathcal{M}^{\text{tr}}/\text{Rel} \\ \downarrow & & \downarrow \\ H^1(\partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}}) & \xrightarrow{\sim} & \mathcal{M}/(\text{Id} - T)\mathcal{M}. \end{array} \quad (3.59)$$

We have the basis

$$e_n = \text{trunc}(Y^n), e_{n-1} = \text{trunc}(Y^{n-1}X), \dots, \begin{cases} Y^{n/2}X^{n/2} & n/2 \text{ odd} \\ 0 & \text{else} \end{cases}$$

for  $\mathcal{M}^{\text{tr}}$ . Let us put  $n_2 = n/2$  or  $n/2 - 1$ . Then the algorithm *Smithnormalform* provides a second basis  $f_n = e_n, f_{n-1}, \dots, f_{n_2}$  such that the module of relations becomes

$$d_n f_n = 0, d_{n-1} f_{n-1} = 0, \dots, d_t f_t = 0, \dots, d_{n_2} f_{n_2} = 0$$

where  $d_{n_2} | d_{n_2+1} | \dots | d_n$ . We have  $d_n = d_{n-1} = \dots = d_{n-2s} = 0$  where  $2s + 1 = \dim H^1(\Gamma \backslash \mathbb{H}, \mathcal{M}) \otimes \mathbb{Q}$  and  $d_{n-2s-1} \neq 0$ .

Now we have a computer program which for a given  $n$  gives us an explicit matrix for  $T_2$ , it is of the form

$$T_2(f_i) = \sum_{j=n}^{j=n_2} t_{i,j}^{(2)} f_j \quad (3.60)$$

where we have (the numeration of the rows and columns is downwards from  $n$  to  $n_2$ )

$$\begin{aligned} t_{\nu,n}^{(2)} &= 0 \text{ for } \nu < n \text{ and } t_{i,j}^{(2)} \in \text{Hom}(\mathbb{Z}/(d_i), \mathbb{Z}/(d_j)) \\ \text{and } t_{i,j}^{(2)} &= 0 \text{ for } i \geq n - 2s, j < n - 2s \end{aligned} \quad (3.61)$$

If we divide by the torsion we get for the restriction map to the boundary cohomology

$$H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})_{\text{int}} = \bigoplus_{\nu=n}^{n-2s} \mathbb{Z} f_\nu \xrightarrow{r} H^1(\partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}})_{\text{int}} = \mathbb{Z} Y^n \quad (3.62)$$

where  $f_n \mapsto Y^n$  and  $T_2(Y^n) = (2^{n+1} + 1)Y^n$ . Now we will find that the endomorphism  $T_2 - (2^{n+1} + 1)\text{Id}$  of  $H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})_{\text{int}}$  is injective (Manin-Drinfeld principle see section 4.1.5) and this implies that we can find a vector

$$\text{Eis}_n = f_n + \sum_{\nu=n-1}^{\nu=n-2s} x_\nu f_\nu, \quad x_\nu \in \mathbb{Q} \quad (3.63)$$

which is an eigenvector for  $T_2$  i.e.

$$T_2(\text{Eis}_n) = (2^{n+1} + 1)\text{Eis}_n. \quad (3.64)$$

The least common multiple  $\Delta(n)$  of the denominators of the  $x_\nu$  is the denominator of the Eisenstein class, it is the smallest positive integer for which

$$\Delta(n)\text{Eis}_n \in H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})_{\text{int}}. \quad (3.65)$$

This denominator is of great interest and our computer program allows us to compute it for any given not too large  $n$ . We simply have to compute the  $x_\nu$ . We know that  $T_2(f_n) = (2^{n+1} + 1)f_n + \sum_{\mu=n-1}^{\mu=n-2s} t_{n,\mu}^{(2)} f_\mu$  and then the  $x_\nu$  are the unique solution of

$$\sum_{\nu=n-1}^{\nu=n-2s} ((2^{n+1} + 1)\delta_{\nu,\mu} - t_{\nu,\mu}^{(2)})x_\nu = t_{n,\mu}^{(2)}; \{\mu = n-1, \dots, n-2s\} \quad (3.66)$$

With the help of H. Gangl we carried the computation of the  $x_\nu$  and hence the  $\Delta(n)$  and we found for some not too large values of  $n$  (roughly  $n \leq 150$ ) that

$$\Delta(n) = \text{numerator}(\zeta(-1-n)). \quad (3.67)$$

Comment: Actually we will show that this is a theorem (Theorem 5.1.1). The reader might argue, why do you make such efforts to find out some experimental evidence for something you know to be true?

There are several reasons for doing this but the main motivation is the following. The Theorem 5.1.1 is hopefully a special case of a much more general assertion. The problem to determine (estimate) denominators of Eisenstein classes is ubiquitous in the cohomology of arithmetic groups. And we have many cases where we have conjectures relating these denominators to special values of  $L$ -functions. (See [31]) But in many of these cases the methods to prove theorems like Theorem 5.1.1 seem to fail. Therefore it seems to be of interest to develop algorithms which compute the cohomology and the action of Hecke operators explicitly in given cases. A toy model of such an algorithm has been written in the above case. We are aware that these algorithms may become very slow for more general reductive groups.

### 3.3.5 Computing mod $p$

Of course the coefficients  $t_{\nu,\mu}^{(2)}$  become very large if  $n$  becomes larger, hence we can verify (3.67) only in a very small range of degrees  $n$ .

But if we are a little bit more modest we may be able to check experimentally whether a given - perhaps large- prime  $p$ , which divides a numerator  $\zeta(-1-n)$  for a very large  $n$  actually divides  $\Delta(n)$ . Here we need a little bit of luck.

Assume that we have such a pair  $(p, n)$ . We want to show that the prime  $p$  divides the lcm of the denominators of the  $x_\nu$  in (3.66) and this means that

the equation (3.66) has no solution in  $\mathbb{Z}_{(p)}$ , the local ring at  $p$ . This is of course clear if the  $\pmod p$  reduced equation

$$\sum_{\nu=n-1}^{\nu \equiv n-2s} ((2^{n+1} + 1)\delta_{\nu,\mu} - t_{\nu,\mu}^{(2)})x_\nu \equiv t_{n,\mu}^{(2)} \pmod p \quad (3.68)$$

has no solution. (Of course the converse is not true, therefore we need just a little bit of luck!). In this computation the numbers become much smaller. In fact this has now been checked for all  $n \leq 100$  we can easily go much further.

higher

### Higher powers of $p$

This reasoning can also be applied if we look at higher powers of  $p$  dividing a numerator  $\zeta(-1-n)$ . Let us assume that  $p^{\delta_p(n)} | \text{numerator} \zeta(-1-n)$ . We have to show that  $p^{\delta_p(n)}$  divides the lcm of the denominators of the  $x_\nu$  in equation (3.66). This follows if we show that the equation

$$\sum_{\nu=n-1}^{\nu \equiv n-2s} ((2^{n+1} + 1)\delta_{\nu,\mu} - t_{\nu,\mu}^{(2)})x_\nu \equiv p^{\delta_p(n)-1} t_{n,\mu}^{(2)} \pmod{p^{\delta_p(n)}} \quad (3.69)$$

has no solution. This in turn means that the class

$$[\mathcal{X}_n \otimes \mathbb{Z}/p^{\delta_p(n)}\mathbb{Z}] \in \text{Ext}_{\mathcal{H}}^1((\mathbb{Z}/p^{\delta_p(n)}\mathbb{Z})(-1-n), H_{\text{int},!}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}} \otimes (\mathbb{Z}/p^{\delta_p(n)}\mathbb{Z})))$$

has exact order  $p^{\delta_p(n)}$ .

Interesting cases to check are  $p = 37, 59, 67, 101, \dots$  then we have

$$\zeta(-31) \equiv 0 \pmod{37}; \zeta(-283) \equiv 0 \pmod{37^2}; \zeta(-37579) \equiv 0 \pmod{37^3}; \zeta(-1072543) \equiv 0 \pmod{37^4}$$

$$\zeta(-43) \equiv 0 \pmod{59}; \zeta(-913) \equiv 0 \pmod{59^2}$$

Here our computations have a surprising outcome. For  $\zeta(-283)$  resp.  $\zeta(-913)$  it has been checked that the order of the extension class is 37 resp. 59 so it is smaller than expected. This is not in conflict with the assertion that the denominator is of order  $37^2, 59^2$ . In fact it turns out that the determinant of the matrix on the left hand side in (3.69) is  $(37^3)^2 = 37^6$  where the denominator only predicts  $37^4$ . Is this always so and is this also true for other Hecke operators?

### 3.3.6 The denominator and the congruences

For the following we assume that (3.67) is correct. We discuss the denominator of the Eisenstein class in this special case. In [Talk-Lille] this is discussed in a more abstract way, so here we treat basically the simplest example of 4.3 in [Talk-Lille]. Remember that in this section  $\mathcal{M} = \tilde{\mathcal{M}}_n$ , i.e. we have fixed an even positive integer  $n$ .

We have the fundamental exact sequence

fuex

$$0 \rightarrow H_{\text{int},!}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) \rightarrow H_{\text{int}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) \xrightarrow{r} H_{\text{int}}^1(\partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}}) = \mathbb{Z}e_n \rightarrow 0 \quad (3.70)$$

and we know that  $T_2(e_n) = (2^{n+1} + 1)e_n$ . We get a submodule

$$H_{\text{int},!}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) \oplus \mathbb{Z}\tilde{e}_n \subset H_{\text{int}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) \quad (3.71)$$

where  $\tilde{e}_n$  is primitive and  $T_2\tilde{e}_n = (2^{n+1} + 1)\tilde{e}_n$ . We have  $r(\tilde{e}_n) = \Delta(n)e_n$  and

$$H_{\text{int}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})/(H_{\text{int},!}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) \oplus \mathbb{Z}\tilde{e}_n) = \mathbb{Z}/\Delta(n)\mathbb{Z} \quad (3.72)$$

Any  $m \in \mathbb{Z}/\Delta(n)\mathbb{Z}$  can be written as

$$m = r\left(\frac{y' + m\tilde{e}_n}{\Delta(n)}\right) \quad (3.73)$$

and this yields an inclusion  $\mathbb{Z}/\Delta(n)\mathbb{Z} \hookrightarrow H_{\text{int},!}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) \otimes \mathbb{Z}/\Delta(n)\mathbb{Z}$ .

Hence

**Theorem 3.3.1.** *The Hecke module  $H_{\text{int}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) \otimes \mathbb{Z}/\Delta(n)\mathbb{Z}$  contains a cyclic submodule  $\mathbb{Z}/\Delta(n)\mathbb{Z}(-1-n)$  on which the Hecke operator  $T_p$  acts by the eigenvalue  $p^{n+1} + 1 \pmod{\Delta(n)}$  for all primes  $p$ .*

This theorem has interesting consequences which will be discussed in the following.

In section (4.1.6) we will review the famous multiplicity one theorem which follows from the theory of automorphic forms. This theorem implies that we can find a finite normal field extension  $F/\mathbb{Q}$  such that decoF

$$H_{\text{int},!}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) \otimes F = \bigoplus_{\pi_f} H_{\text{int},!}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}} \otimes F)[\pi_f] \quad (3.74)$$

where  $\pi_f$  runs over a finite set of homomorphisms  $\pi_f : \mathcal{H} \rightarrow \mathcal{O}_F$ , and where  $H^1..[\pi_f]$  is the rank 2 eigenspace for  $\pi_f$ . We also have the action of the complex conjugation on the cohomology (See sect. how) and under this action each eigenspace decomposes into a one dimensional + and a one dimensional - eigenspace, i.e.  $H^1..[\pi_f] = H_+^1..[\pi_f] \oplus H_-^1..[\pi_f]$ . Let us denote the set of  $\pi_f : \mathcal{H} \rightarrow \mathcal{O}_f$  which occur with positive multiplicity (then 2) in the above decomposition by  $\text{Coh}_1^{(n)}$ .

Our considerations at the beginning of this section imply that we also have a decomposition of

$$H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) \otimes F = H_{\text{int}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}) \otimes F \oplus Fe_n$$

where  $T_p e_n = (p^{n+1} + 1)e_n$ . Let  $\pi_f^{\text{Eis}} : \mathcal{H} \rightarrow \mathbb{Z}$  be the homomorphism  $\pi_f^{\text{Eis}} : T_p \rightarrow p^{n+1} + 1$ .

This decomposition induces a Jordan-Hölder filtration on the integral cohomology JH

$$(0) \subset \mathcal{JH}^{(1)} H_{\text{int},!}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{\mathcal{O}_F}) \subset \mathcal{JH}^{(2)} H_{\text{int},!}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{\mathcal{O}_F}) \subset \dots \subset \mathcal{JH}^{(r)} H_{\text{int},!}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{\mathcal{O}_F}) \quad (3.75)$$

where the subquotients are locally free  $\mathcal{O}_F$  modules of rank 2 and after tensoring with  $F$  they become isomorphic to the different eigenspaces.

We choose a prime  $p$  which divides  $\Delta(n)$ , let  $p^{\delta_p(n)} \parallel \Delta(n)$ . Let  $\mathfrak{p}$  be a prime in  $\mathcal{O}_F$  which lies above  $p$ . If  $e_p$  is the ramification index then we have

$$\{0\} \subset \mathcal{O}_F/\mathfrak{p}^{e_p \delta_p(n)}(-1-n) \subset H_{\text{int},!}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{\mathcal{O}_F}) \otimes \mathcal{O}_F/\mathfrak{p}^{e_p \delta_p(n)} \quad (3.76)$$

The above Jordan-Hölder filtration induces a Jordan-Hölder filtration on the cohomology  $\text{mod } \mathfrak{p}^{e_p \delta_p(n)}$  we have  $\boxed{\text{JHmod}}$

$$\{0\} \subset \mathcal{JH}^{(1)} H_{\text{int},!}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{\mathcal{O}_F}) \otimes \mathcal{O}_F/\mathfrak{p}^{e_p \delta_p(n)} \hookrightarrow \mathcal{JH}^{(2)} \dots \quad (3.77)$$

where again the subquotients are free  $\mathcal{O}_F/\mathfrak{p}^{e_p \delta_p(n)}$  modules of rank 2. A simple argument shows

$$\boxed{\text{congl}}$$

**Theorem 3.3.2.** *We can find  $\pi_{f,1}, \pi_{f,2}, \dots, \pi_{f,r}$  in the above decomposition and numbers  $f_1 > 0, f_2 > 0, \dots, f_r > 0$  such that  $\sum f_i = e_p \delta_p(n)$  and we have the congruence*

$$\pi_{f,i}(T_\ell) \equiv \ell^{n+1} + 1 \pmod{\mathfrak{p}^{f_i}} \quad (3.78)$$

for all primes  $\ell$ .

In the following section we look at this theorem from a slightly different point of view.

### $p$ -adic coefficients

In the previous section we decomposed the inner cohomology into eigenspaces under the action of the Hecke algebra. In our special situation - the underlying group  $G = \text{Gl}_2$ - this is also valid for the full cohomology. But our main object of interest is the cohomology with integral coefficients and our example above shows that the cohomology with integral coefficients does not split.

To investigate the structure of the cohomology groups  $H^\bullet(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})$  we choose a prime  $p$ . This prime will be fixed throughout this section, let  $\mathbb{Z}_{(p)} \subset \mathbb{Q}$  be the local ring at  $p$ . We are interested in the structure of the cohomology groups  $H^\bullet(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}} \otimes \mathbb{Z}_{(p)})$  as modules under the Hecke algebra. But now it is convenient to go still one step further, we tensorize our coefficient systems by  $\mathbb{Z}_p$ , the ring of  $p$ -adic integers. We want to simplify the notation: In this section we denote by  $\mathcal{M}_n$  the  $\mathbb{Z}_p$ -module  $\mathcal{M}_\lambda \otimes \mathbb{Z}_p$  where  $\lambda = n\gamma + d \det$  where the value of  $d$  is irrelevant it just has to have the right parity. (Comment?  $\mathbb{Z}_{(p)} \rightarrow \mathbb{Z}_p$  is flat hence it does preserve  $\text{Ext}^1$  groups.)

Let  $M$  be any finitely generated  $\mathbb{Z}_p$ -module, let  $T_p : M \rightarrow M$  be an endomorphism. Of course  $X$  is a topological module, the open neighborhoods of 0 are the modules  $p^r M$ . Following Hida we define two submodules

$$M_{\text{ord}} = \bigcap_{r \rightarrow \infty} T_p^r M; \quad M_{\text{nilpt}} = \{x \in M \mid T_p^r x \rightarrow 0\} \quad (3.79)$$

A simple compactness argument shows that

$$M = M_{\text{ord}} \oplus M_{\text{nilpt}} \quad (3.80)$$

and it is also clear that  $M \rightarrow M_{\text{ord}}$  is an exact functor.

We apply this to our cohomology groups, and we assume that  $\Gamma = \text{Sl}_2(\mathbb{Z})$ . We start from the exact sequence of  $\Gamma$  modules

$$0 \rightarrow \mathcal{M}_n \xrightarrow{\times p} \mathcal{M}_n \rightarrow \mathcal{M}_n \otimes \mathbb{F}_p \rightarrow 0. \quad (3.81)$$

Here we want to assume that  $p > 3$  then we get the resulting exact sequence of sheaves and hence a long exact sequence of cohomology groups

$$\begin{aligned} 0 \rightarrow (\mathcal{M}_n^\Gamma)_{\text{ord}} \xrightarrow{\times p} (\mathcal{M}_n^\Gamma)_{\text{ord}} \rightarrow (\mathcal{M}_n \otimes \mathbb{F}_p)_{\text{ord}} \rightarrow \\ \rightarrow H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n) \xrightarrow{\times p} H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n) \rightarrow H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathbb{F}_p) \rightarrow 0 \end{aligned} \quad (3.82)$$

and we can break this sequence into pieces

$$0 \rightarrow (\mathcal{M}_n^\Gamma)_{\text{ord}} \xrightarrow{\times p} (\mathcal{M}_n^\Gamma)_{\text{ord}} \rightarrow (\mathcal{M}_n \otimes \mathbb{F}_p)_{\text{ord}}^\Gamma \rightarrow H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n)[p] \rightarrow 0 \quad (3.83)$$

and

$$0 \rightarrow H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n)[p] \rightarrow H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n) \xrightarrow{\times p} H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n) \rightarrow H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathbb{F}_p) \rightarrow 0 \quad (3.84)$$

where of course  $\dots [p]$  means kernel of the multiplication by  $p$  and the far most 0 on the right is the vanishing of  $H^2$ .

We analyze these two sequences and get

ordtorfree

**Theorem 3.3.3.** *The cohomology  $H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n)$  is torsion free unless we have  $n > 0$  and  $n \equiv 0 \pmod{p(p-1)}$ . The cohomology groups  $H^1_{c, \text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n)$  are always torsion free and  $H^2_{c, \text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n) = 0$*

*Proof.* We consider the polynomial ring in two variables  $\mathbb{F}_p[X, Y]$ . On this ring we have the action of  $\text{Sl}_2(\mathbb{Z})$ . It is an old theorem of L.E. Dickson that the ring of invariants is generated by the two polynomials

$$f_1 = X^p Y - X Y^p \text{ and } f_2 = \frac{X^{p^2-1} - Y^{p^2-1}}{X^{p-1} - Y^{p-1}} = X^{(p-1)p} + X^{(p-1)(p-1)} Y^{p-1} + \dots \quad (3.85)$$

Now every element in  $(\mathcal{M}_n \otimes \mathbb{F}_p)_{\text{ord}}^\Gamma$  is a sum of monomials  $f_1^a f_2^b$  where  $a(p+1) + bp(p-1) = n$ . We see that

$$u \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} = u_\alpha : \mathcal{M}_n^{(\alpha)} \rightarrow \mathcal{M}_n$$

multiplies  $f_1$  with a multiple of  $p$  and hence we see that all the monomials with  $a > 0$  are multiplied by a multiple of  $p$ . This means that  $(\mathcal{M}_n \otimes \mathbb{F}_p)_{\text{ord}}^\Gamma \neq 0$  if and only if  $n = bp(p-1)$ . If  $n = 0$  we the map  $\mathcal{M}_n^\Gamma = \mathbb{Z}_p \rightarrow (\mathcal{M}_n \otimes \mathbb{F}_p)^\Gamma$  is surjective if  $n > 0$  we have  $\mathcal{M}_n^\Gamma = 0$  and hence the theorem.

For the assertions concerning the compactly supported cohomology we have to recall that  $H^2_c(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n) = (\mathcal{M}_n)_\Gamma = \mathcal{M}_n / I_\Gamma \mathcal{M}_n$  [book vol I, section 2 and 4.8.5]. We check easily that  $X^n, Y^n \in I_\Gamma \mathcal{M}_n$  and the assertion is clear.  $\square$

We write  $n = n_0 + (p-1)\alpha$  where we assume  $0 < n_0 < p-1$ , we know that

$$H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n) \otimes \mathbb{Z}/p^r\mathbb{Z} \xrightarrow{\sim} H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathbb{Z}/p^r) \quad (3.86)$$

we have a second theorem

interpol

**Theorem 3.3.4.** *If  $n = n_0 + (p-1)\alpha, n' = n_0 + (p-1)\alpha'$  and  $\alpha \equiv \alpha' \pmod{p^{r-1}}$ , (i.e.  $n \equiv n' \pmod{(p-1)p^{r-1}}$ ) then we have a canonical Hecke invariant isomorphism*

$$\Phi(n, n')_r : H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathbb{Z}/p^r) \xrightarrow{\sim} H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{n'} \otimes \mathbb{Z}/p^r). \quad (3.87)$$

*This system of isomorphisms is consistent with change of the parameter  $\alpha, \alpha'$  and  $r$ .*

*Proof.* See paper on interpolation. □

We find a finite extension  $F/\mathbb{Q}_p$  such that we have a decomposition into eigenspaces

$$H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes F) = \bigoplus_{\pi_f} H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes F)[\pi_f] \oplus Fe_n \quad (3.88)$$

where the first summation goes over those  $\pi_f \in \text{Coh}_!^{(n)}$  for which  $\pi_f(T_p)$  is a unit in  $\mathcal{O}_{\mathfrak{p}}$ , the ring of integers in  $F$ . Let us denote this set by  $\text{Coh}_{!, \text{ord}}^{(n)}$ . Then the full summation goes over the set  $\text{Coh}_{\text{ord}}^{(n)} = \text{Coh}_{!, \text{ord}}^{(n)} \cup \{\pi_f^{\text{Eis}}\}$ . Intersecting this decomposition with  $H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathcal{O}_{\mathfrak{p}})$  gives us a submodule of finite index

$$H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathcal{O}_{\mathfrak{p}}) \supset \bigoplus_{\pi_f} H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathcal{O}_{\mathfrak{p}})[\pi_f] \oplus \mathcal{O}_{\mathfrak{p}}e_n \quad (3.89)$$

and this also gives us a Jordan-Hölder filtration as in (3.75).

We consider the reduction maps

redukdiag

$$\begin{array}{ccc} H^1_{!, \text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathcal{O}_{\mathfrak{p}}) & \rightarrow & H^1_{!, \text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathbb{F}(\mathfrak{p})) \\ \downarrow & & \downarrow \\ H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathcal{O}_{\mathfrak{p}}) & \rightarrow & H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathbb{F}(\mathfrak{p})) \end{array} \quad (3.90)$$

the right hand sides do not depend on  $\alpha$ . Any  $\pi_f \in \text{Coh}_{\text{ord}}^{(n)}$  we get a non zero homomorphism  $\bar{\pi}_f = \pi_f \times \mathbb{F}(\mathfrak{p}) : \mathcal{H} \rightarrow \mathbb{F}(\mathfrak{p})$ . The map  $\pi_f \rightarrow \bar{\pi}_f$  is not necessarily injective: we say that  $\pi_{1,f}$  and  $\pi_{2,f}$  are congruent  $\pmod{\mathfrak{p}}$  if  $\pi_{1,f}(T_\ell) \equiv \pi_{2,f}(T_\ell) \pmod{\mathfrak{p}}$  for all primes  $\ell$ , or in other words  $\bar{\pi}_{1,f} = \bar{\pi}_{2,f}$ . For a given  $\pi_f$  let  $\{\bar{\pi}_f\}$  be the set of all  $\pi_{i,f}$  which are congruent to the given  $\pi_f$ .

$$H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathbb{F}(\mathfrak{p}))\{\bar{\pi}_f\} = \{x \in H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathbb{F}(\mathfrak{p})) \mid (T_\ell - \bar{\pi}_f(T_\ell))^N x = 0\} \quad (3.91)$$

provided  $N \gg 0$ . Then it is easy to see that (See for instance [book,II], 7.2) that

$$H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathbb{F}(\mathfrak{p})) = \bigoplus_{\bar{\pi}_f} H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathbb{F}(\mathfrak{p}))\{\bar{\pi}_f\} \quad (3.92)$$

The kernel  $\mathfrak{m}_{\bar{\pi}_f}$  of  $\bar{\pi}_f$  is a maximal ideal, let  $\mathcal{H}_{\mathfrak{m}_{\bar{\pi}_f}}$  be the local ring at  $\mathfrak{m}_{\bar{\pi}_f}$ . Then the above decomposition can be written as

$$H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathbb{F}(\mathfrak{p})) = \bigoplus_{\bar{\pi}_f} H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathbb{F}(\mathfrak{p})) \otimes \mathcal{H}_{\mathfrak{m}_{\bar{\pi}_f}} / \mathfrak{m}_{\bar{\pi}_f}^N \quad (3.93)$$

Now we recall that we still have the action of complex conjugation (See sect.2.1.6) on the cohomology and it is clear (SEE(?)) that it commutes with the action of the Hecke algebra. Hence we see that the summands in the above decompose into a + and a - summand, i.e.

$$H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathbb{F}(\mathfrak{p})) \otimes \mathcal{H}_{\mathfrak{m}_{\bar{\pi}_f}} / \mathfrak{m}_{\bar{\pi}_f}^N = \bigoplus_{\pm} H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathbb{F}(\mathfrak{p})) \otimes \mathcal{H}_{\mathfrak{m}_{\bar{\pi}_f}} / \mathfrak{m}_{\bar{\pi}_f}^N[\pm] \quad (3.94)$$

Now we encounter some difficult questions. The first one asks whether we have some kind of multiplicity one theorem mod  $\mathfrak{p}$ . This question can be formulated as follows:

*Are the summands  $H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathbb{F}(\mathfrak{p})) \otimes \mathcal{H}_{\mathfrak{m}_{\bar{\pi}_f}} / \mathfrak{m}_{\bar{\pi}_f}^N[\pm]$  cyclic, i.e. are they - as  $\mathcal{H}_{\mathfrak{m}_{\bar{\pi}_f}} / \mathfrak{m}_{\bar{\pi}_f}^N$  modules - generated by one element ?*

To formulate the second question we regroup the decomposition (3.88)

$$H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes F) = \bigoplus_{\bar{\pi}_f} \left( \bigoplus_{\pi_f \in \{\bar{\pi}_f\}} H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes F)[\pi_f] \right) \quad (3.95)$$

pibar and define

$$\begin{aligned} & H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathcal{O}_{\mathfrak{p}})\{\bar{\pi}_f\} = \\ & \left( \bigoplus_{\pi_f \in \{\bar{\pi}_f\}} H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes F)[\pi_f] \right) \cap H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathcal{O}_{\mathfrak{p}}) \end{aligned} \quad (3.96)$$

and then we get a second variant of (3.94)

$$\bigoplus_{\bar{\pi}_f} H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathcal{O}_{\mathfrak{p}})\{\bar{\pi}_f\} = H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathcal{O}_{\mathfrak{p}}) \quad (3.97)$$

Now we are interested in the structure of the direct summands  $H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathcal{O}_{\mathfrak{p}})\{\bar{\pi}_f\}$ . It is clear that this is a free  $\mathcal{O}_{\mathfrak{p}}$  module of rank

$$r(\{\bar{\pi}_f\}) = \begin{cases} 2\#\{\bar{\pi}_f\} & \text{if } \{\bar{\pi}_f\} \neq \{\bar{\pi}_f^{\text{Eis}}\} \\ 2(\#\{\bar{\pi}_f\} - 1) + 1 & \text{if } \{\bar{\pi}_f\} = \{\bar{\pi}_f^{\text{Eis}}\} \end{cases} \quad (3.98)$$

Again we get a submodule

$$\bigoplus_{\pi_f \in \{\bar{\pi}_f\}} H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathcal{O}_{\mathfrak{p}})[\pi_f] \subset H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathcal{O}_{\mathfrak{p}})\{\bar{\pi}_f\} \quad (3.99)$$

Our second question is

*What can we say about the structure of the quotient*

$$H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathcal{O}_{\mathfrak{p}})\{\bar{\pi}_f\} / \bigoplus_{\pi_f \in \{\bar{\pi}_f\}} H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathcal{O}_{\mathfrak{p}})[\pi_f] ?$$

*For instance we may ask: Is this quotient non trivial if the cardinality of  $\{\bar{\pi}_f\}$  is greater than 1 ?*

For a subset  $\Sigma \subset \{\bar{\pi}_f\}$  we define in analogy with (3.96)

$$\begin{aligned} H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathcal{O}_{\mathfrak{p}})\{\Sigma\} = \\ (\bigoplus_{\pi_f \in \Sigma} H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes F)[\pi_f]) \cap H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathcal{O}_{\mathfrak{p}})\{\bar{\pi}_f\} \end{aligned} \quad (3.100)$$

and we call  $\Sigma$  a block if

$$H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathcal{O}_{\mathfrak{p}})\{\bar{\pi}_f\} = \quad (3.101)$$

$$H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathcal{O}_{\mathfrak{p}})\{\Sigma\} \oplus H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathcal{O}_{\mathfrak{p}})\{\{\bar{\pi}_f\} \setminus \Sigma\} \quad (3.102)$$

Then a slightly stronger version of our question above asks

*Can  $\{\bar{\pi}_f\}$  contain non trivial blocks?*

These two questions are closely related. We will come back later to these issues in this book. In the following we outline the general philosophy:

*The structure of the cohomology as module under the Hecke-algebra is influenced by divisibility of special values of certain  $L$  functions which are attached to the  $\pi_f$ .*

We have some partial results. ( For this see Herbrand -Ribet , Hida.. ).

If we consider the special case of  $\{\bar{\pi}_f^{\text{Eis}}\}$ . Our theorem 3.3.2 implies

$$p \mid \zeta(-1 - n) \Rightarrow \{\bar{\pi}_f^{\text{Eis}}\} > 1,$$

this has been proved by Ribet in [ ], he also proves the converse using a theorem of Herbrand [ ]. Our theorem 3.3.2 is stronger, because it implies higher congruences if  $\zeta(-1 - n)$  is divisible by a higher power of  $p$ . Moreover the existence of congruences do not imply anything about the denominator.

Of course the next question is: If we have  $p \mid \zeta(-1 - n)$ , what is the size of  $\{\bar{\pi}_f^{\text{Eis}}\}$  can it be  $> 2$ ? Let us pick a  $\pi_f \in \{\bar{\pi}_f^{\text{Eis}}\}$  which is not  $\pi_f^{\text{Eis}}$ . To this  $\pi_f$  we attach the so called symmetric square  $L$ -function  $L(\pi_f, \text{Sym}^2, s)$ . (See ...). This  $L$  function evaluated at a suitable "critical" point and divided by a carefully chosen period gives us a number

$$\mathcal{L}(\pi_f, \text{Sym}^2) \in \mathcal{O}_{F_0}$$

here  $F_0$  is a global field whose completion at  $\mathfrak{p}$  is our  $F$  above. Now a theorem Hida says (cum grano salis)

$$\#\pi_f^{\text{Eis}} > 2 \iff \mathfrak{p} \mid \mathcal{L}(\pi_f, \text{Sym}^2) \quad (3.103)$$

(See later) If we accept these two results then we get

Vand

**Theorem 3.3.5.** *If  $p^{\delta_p(n)} \mid \zeta(-1-n)$  and if  $\mathcal{L}(\pi_f, \text{Sym}^2) \not\subseteq \mathfrak{p}$ , then the number  $r$  in theorem 3.3.2 is equal to one, i.e.  $\{\bar{\pi}_f^{\text{Eis}}\} = \{\pi_f, \pi_f^{\text{Eis}}\}$  and we have the congruence*

$$\pi_f(T_\ell) \equiv \ell^{n+1} + 1 \pmod{\mathfrak{p}^{\delta_p(n)} \forall \text{ primes } \ell}$$

Finally we get  $\pi_f(T_\ell) \in \mathbb{Z}_p$  for all primes  $\ell$  and hence we may take  $\mathcal{O}_{\mathfrak{p}} = \mathbb{Z}_p$ .

We can find a basis  $f_0, f_1, f_3$  of  $H_{ord}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})$  where

a)  $f_1, f_2$  form a basis of  $H_{ord}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}})$

b) The complex conjugation  $c$  acts by  $c(f_i) = (-1)^{i+1} f_i$

and

c) the matrix  $T_\ell^{ord}$  with respect to this basis satisfies

$$T_\ell^{ord} \equiv \begin{pmatrix} \ell^{n+1} + 1 & 0 & 1 \\ 0 & \ell^{n+1} + 1 & 0 \\ 0 & 0 & \ell^{n+1} + 1 \end{pmatrix} \pmod{p^{\delta_p(n)}}$$

*Proof.* Clear □

If we drop the assumption  $\mathcal{L}(\pi_f, \text{Sym}^2) \not\subseteq \mathfrak{p}$  then the situation becomes definitely more complicated. In this case we have  $\{\bar{\pi}_f^{\text{Eis}}\} = \{\bar{\pi}_f^{\text{Eis}}, \pi_{1,f}, \dots, \pi_{r,f}\}$  where now  $r > 1$ . We apply theorem 3.3.2 to this situation where we replace the subscript  $!,_{\text{int}}$  by  $_{ord}$ . We have the filtration which is analogous to (3.75) but now the last quotient is of rank one and isomorphic to the cohomology of the boundary. We find a basis  $f_0, f_1, e_1, f_2, e_2, \dots, f_r, e_r$  adapted to this filtration and where  $c(f_i) = -f_i, c(e_i) = e_i$ . Then we get a matrix (we consider the case  $r = 2$ )

$$T_\ell^{ord} = \begin{pmatrix} \ell^{n+1} + 1 & 0 & 1 & 0 & 1 \\ 0 & \pi_{f,1}(T_\ell) & 0 & u & 0 \\ 0 & 0 & \pi_{f,1}(T_\ell) & 0 & v \\ 0 & 0 & 0 & \pi_{f,2}(T_\ell) & 0 \\ 0 & 0 & 0 & 0 & \pi_{f,2}(T_\ell) \end{pmatrix} \quad (3.104)$$

where  $u, v$  are units in  $\mathbb{Z}_p$  and where the diagonal entries satisfy some congruences  $\pi_{\nu,1}(T_\ell) \equiv \ell^{n+1} + 1 \pmod{\mathfrak{p}^{n_\nu}}$  where  $n_1 + n_2 = e_p \delta_p(n)$ . We come back to this later.

p-adic-zeta

### 3.3.7 The $p$ -adic $\zeta$ -function

We return to section 3.3.5. We are interested in the case that  $p$  is an irregular prime, i.e.  $p \mid \zeta(-1 - n_0)$ . We also assume that also  $\mathcal{L}(\pi_f, Sym^2) \notin \mathfrak{p}$ . We consider  $\zeta(-1 - n) = \zeta(-1 - n_0 - \alpha(p - 1))$  as function in the variable  $\alpha \in \mathbb{N}$  and we want to find values  $n = -1 - n_0 - \alpha(p - 1)$  such that  $\zeta(-1 - n)$  is divisible by higher powers of  $p$ . We know that there exist a  $p$ -adic  $\zeta$ -function and tells us - provided  $n_0 > 0$  - that

$$\boxed{\text{p-appr}}$$

$$\zeta(-1 - n) = \zeta(-1 - n_0 - \alpha(p - 1)) \equiv \zeta(-1 - n_0) + a(n_0, 1)\alpha p + a(n_0, 2)\alpha^2 p^2 \dots \quad (3.105)$$

where the coefficients  $a(n_0, \nu) \in \mathbb{Z}_p$ . Now several things can happen.

A) Our prime  $p$  does not divide the second coefficient  $a(n_0, 2)$ . Then we can apply Newton's method and we find a converging sequence  $\alpha_1, \alpha_2, \dots$  such that

$$\alpha_\nu \equiv \alpha_{\nu+1} \pmod{p^\nu} \text{ and } \zeta(-1 - n_0 - \alpha_\nu(p - 1)) \equiv 0 \pmod{p^{\nu+1}} \quad (3.106)$$

If now  $n_\nu = n_0 + \alpha_\nu(p - 1)$  then we can form the system of Hecke-modules (A Hida family)  $H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \mathcal{M}_{n_\nu})(\{\bar{\pi}_f^{\text{Eis}}\})$  and theorem 3.3.4 gives us Hecke-module morphisms

$$H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{n_\nu+1} \otimes \mathbb{Z}/p^{\nu+1}\mathbb{Z}) \xrightarrow{\Phi_\nu} H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{n_\nu} \otimes \mathbb{Z}/p^\nu\mathbb{Z}) \quad (3.107)$$

The sequence  $n_\nu$  converges to an  $p$ -adic integer  $n_\infty$ , we can form the projective limit and define

$$H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{n_\infty}) = \varprojlim H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{n_\nu} \otimes \mathbb{Z}/p^\nu\mathbb{Z}) \quad (3.108)$$

Under our assumptions this is a free  $\mathbb{Z}_p$ -module of rank 3. The Hecke operators  $T_\ell^{\text{ord}}$  acts on  $H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{n_\nu} \otimes \mathbb{Z}/p^\nu\mathbb{Z})$  by a matrix of the shape as in theorem 3.3.5, and the eigenvalues on the diagonal are

$$\ell^{n_\nu+1} + 1 = \ell^{n_0+(p-1)\alpha_\nu} + 1 \pmod{p^\nu}$$

For  $\ell \neq p$  we write  $\ell^{p-1} = 1 + \delta(\ell)p$ ,  $\delta(\ell) \in \mathbb{N}$  and then  $\ell^{n_0+(p-1)\alpha_\nu} = \ell^{n_0}(1 + \delta(\ell)p)^{\alpha_\nu}$  and hence it follows that  $\lim_{\nu \rightarrow \infty} \ell^{n_\nu} = \ell^{n_\infty}$  exists. Hence we see that  $T_\ell^{\text{ord}}$  acts on  $H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{n_\infty})$  by the matrix

$$T_\ell^{\text{ord}} \equiv \begin{pmatrix} \ell^{n_\infty+1} + 1 & 0 & 1 \\ 0 & \ell^{n_\infty+1} + 1 & 0 \\ 0 & 0 & \ell^{n_\infty+1} + 1 \end{pmatrix}$$

B) We have  $p \mid \zeta(-1 - n_0)$ ;  $p^2 \nmid \zeta(-1 - n_0)$  and  $p \mid a(n_0, 1)$ . In this case we can not increase the  $p$  power dividing  $\zeta(-1 - n)$ .

C) We have  $p^2 \mid \zeta(-1 - n_0)$ ;  $p \mid a(n_0, 1)$  and  $p \nmid a(n_0, 2)$

We rewrite (3.105)

$$\frac{\zeta(-1 - n)}{p^2} \equiv \frac{\zeta(-1 - n_0)}{p^2} + \frac{a(n_0, 1)}{p}\alpha + a(n_0, 2)\alpha^2 \pmod{p} \quad (3.109)$$

Now we get two numbers  $\alpha_\infty, \beta_\infty$  such that

$$\zeta(-1 - n_0 - \alpha_\infty(p-1)) = 0 ; \zeta(-1 - n_0 - \beta_\infty(p-1)) = 0$$

but these numbers are not necessarily in  $\mathbb{Z}_p$ , they lie in a quadratic extension  $\mathcal{O}_p$  of  $\mathbb{Z}_p$  hence they are not necessarily approximable by (positive ) integers. If we want to interpret these zeros in terms of cohomology modules with an action of the Hecke algebra we have to extend the range of coefficient systems. In [Ha-Int] we define "coefficient systems"  $\mathcal{M}_{n_0, \alpha}^\dagger$  where now  $\alpha$  is any element in  $\mathcal{O}_{\mathbb{C}_p}$ . (These coefficient systems are denoted  $\mathcal{P}_\chi$  in [Ha-Int]).

These coefficient systems are infinite dimensional  $\mathcal{O}_{\mathbb{C}_p}$ -modules, we can define the ordinary cohomology  $H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \mathcal{M}_{n_0, \alpha}^\dagger)$ . On these (ordinary) cohomology modules we have an action of the Hecke algebra and they satisfy the same interpolation properties as the previous ones, especially we have an extension of theorem 3.3.4 for these cohomology modules.

If  $\alpha = a$  is a positive integer then we have a natural homomorphism

$$\Psi_a : \mathcal{M}_{n_0+a(p-1)} \rightarrow \mathcal{M}_{n_0, \alpha}^\dagger$$

and this map induces an isomorphism on the ordinary part of the cohomology

isop

$$\Psi^{(1)} : H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \mathcal{M}_{n_0+a(p-1)}) \xrightarrow{\sim} H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \mathcal{M}_{n_0, \alpha}^\dagger) \quad (3.110)$$

We now allow any  $\alpha \in \mathcal{O}_p$ , our coefficient system will then be a system of  $\mathcal{O}_p$  modules and the cohomology modules will be  $\mathcal{O}_p$  modules. Of course we still have our fundamental exact sequence (3.111) of  $\mathcal{O}_p$  modules.

$$\rightarrow H_{\text{ord}, c}^1(\Gamma \backslash \mathbb{H}, \mathcal{M}_{n_0, \alpha}^\dagger) \rightarrow H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \mathcal{M}_{n_0, \alpha}^\dagger) \xrightarrow{r} H_{\text{ord}}^1(\partial(\Gamma \backslash \mathbb{H}), \mathcal{M}_{n_0, \alpha}^\dagger) = \mathcal{O}_p e_{n_0, \alpha} \rightarrow 0 \quad (3.111)$$

This is an exact sequence of Hecke-modules and we still have

$$T_\ell^{\text{ord}}(e_n) = (\ell^{n_0}(\ell^{p-1})^\alpha + 1)e_{n_0, \alpha} \quad (3.112)$$

Let  $\mathfrak{p} = (\varpi_p)$ , we define  $\delta_p(\alpha)$  by

$$\varpi_p^{\delta_p(\alpha)} || \zeta(-1 - n_0 - \alpha(p-1)).$$

In a forthcoming paper with Mahnkopf we will (hopefully) show that we can construct a section

Eissec

$$\text{Eis}_\alpha : \mathcal{O}_{\mathbb{C}_p} e_{n_0, \alpha} \otimes \mathbb{Q}_p \rightarrow H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \mathcal{M}_{n_0, \alpha}^\dagger) \otimes \mathbb{Q}_p \quad (3.113)$$

which is defined by analytic continuation and that  $\varpi_p^{\delta_p(\alpha)}$  is the exact denominator of  $\text{Eis}_\alpha$ .

If this turns out to be true then we can extend the results for ordinary cohomology modules  $H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \mathcal{M}_{n_0+(p-1)\alpha})$  to the extended class of cohomology modules  $H_{\text{ord}}^1(\Gamma \backslash \mathbb{H}, \mathcal{M}_{n_0, \alpha}^\dagger)$ . Especially if we look at our roots  $\alpha_\infty, \beta_\infty$  and assume that they are different then we get a theorem analogous to the theorem 3.3.5 for both of them. If these two roots are the same the situation is not clear to me.

### 3.3.8 The Wieferich dilemma

We are still assuming that our group  $\Gamma = \text{Sl}_2(\mathbb{Z})$ . We get a clean statement if we are in case A), i.e.

$$p \mid \zeta(-1 - n_0), p \nmid a(n_0, 1), p \nmid \mathcal{L}(\pi_f, \text{Sym}^2)$$

At the present moment we do not know of any prime  $p \mid \zeta(-1 - n_0)$  which does not satisfy A). This is not surprising: The primes  $p \mid \zeta(-1 - n_0)$  are called the irregular primes and they start with

$$37 \mid \zeta(-1 - 30), 59 \mid \zeta(-1 - 42) \dots$$

It is believable that for a prime  $p \mid \zeta(-1 - n_0)$  the numbers  $a(n_0, 1)$  and  $\mathcal{L}(\pi_f, \text{Sym}^2)$  are "unrelated" and in other words the residue classes  $a(n_0, 1) \pmod p$  and  $\mathcal{L}(\pi_f, \text{Sym}^2) \pmod p$  are randomly distributed. Hence we expect that the primes  $p \mid \zeta(-1 - n_0)$  which do not satisfy A) is a "sparsely distributed".

**Reference to Joe Buhler**

But this does not say that this never happens, actually depending on the probabilistic argument you prefer, it should happen eventually. But perhaps we will never find such a prime.

On the other hand

*The Wieferich dilemma: We do not know whether the set of primes which satisfy A) is infinite.*

We drop our assumption that  $\Gamma = \text{Sl}_2(\mathbb{Z})$  and replace it by a normal congruence subgroup of finite index. We choose a free  $\mathbb{Z}$ -module of finite rank  $\mathcal{V}$  with an action of  $\Gamma_0/\Gamma$ , i.e. we have a representation

$$\rho_{\mathcal{V}} : \Gamma_0/\Gamma \rightarrow \text{Gl}(\mathcal{V})$$

we assume that the matrix  $-\text{Id}$  acts by a scalar  $\omega_{\mathcal{V}}(-\text{Id}) = \pm 1$ . We look at the  $\Gamma$ -modules  $\widetilde{\mathcal{M}}_n \otimes \mathcal{V}$ , we assume that  $\mathcal{V}(-\text{Id}) \equiv n \pmod 2$ , These modules provide sheaves  $\widetilde{\mathcal{M}}_n \otimes \mathcal{V}$  and we can study the cohomology groups and especially we can study the fundamental exact sequence

$$\rightarrow H^1_{\text{ord},c}(\Gamma \backslash \mathbb{H}, \widetilde{\mathcal{M}}_n \otimes \mathcal{V}) \rightarrow H^1_{\text{ord}}(\Gamma \backslash \mathbb{H}, \widetilde{\mathcal{M}}_n \otimes \mathcal{V}) \xrightarrow{r} H^1_{\text{ord}}(\partial(\Gamma \backslash \mathbb{H}), \widetilde{\mathcal{M}}_n \otimes \mathcal{V}) \tag{3.114}$$

We have to compute  $H^1_{\text{ord}}(\partial(\Gamma \backslash \mathbb{H}), \widetilde{\mathcal{M}}_n \otimes \mathcal{V})$  as a module under the Hecke algebra and we can ask the denominator question again, provided this boundary cohomology is not trivial.

We may for instance choose a positive integer  $N$  and we consider the congruence subgroup  $\Gamma_0(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{Sl}_2(\mathbb{Z}) \mid c \equiv 0 \pmod N \right\}$ . Let  $\Gamma_1(N) \subset \Gamma_0(N)$  be the subgroup where  $a \equiv d \equiv 1 \pmod N$  then  $\Gamma_0(N)/\Gamma_1(N) = (\mathbb{Z}/N\mathbb{Z})^*$  We choose a character  $\chi : \Gamma_0(N)/\Gamma_1(N) \rightarrow \mathbb{C}^\times$  and consider the representation  $\mathcal{V} = \text{Ind}_{\Gamma_0(N)}^\Gamma \chi$ . In this case the denominator is essentially given by  $L$  values  $L(\chi, -1 - n)$  and these values will be divisible by smaller primes (compared to 37) and our chances to encounter cases of B) and or C) are much better.

## Chapter 4

# Representation Theory, Eichler-Shimura Isomorphism

HC

### 4.1 Harish-Chandra modules with cohomology

In Chapter 8 we will give a general discussion of the tools from representation theory and analysis which help us to understand the cohomology of arithmetic groups. Especially in Chapter 8 section 9.5 we will recall the results of Vogan-Zuckerman on the cohomology of Harish-Chandra modules.

Here we specialize these results to the specific cases  $G = \mathrm{Gl}_2(\mathbb{R})$  (case A) and  $G = \mathrm{Gl}_2(\mathbb{C})$  (case B)). For the general definition of Harish-Chandra modules and for the definition of  $(\mathfrak{g}, K_\infty)$  cohomology we refer to Chap.III, 4.

Mlambda

#### 4.1.1 The finite rank highest weight modules

We consider the case A), in this case our group  $G/\mathbb{R}$  is the base extension of the reductive group scheme  $\mathcal{G} = \mathrm{Gl}_2/\mathrm{Spec}(\mathbb{Z})$ . In principle this a pretentious language. At this point it simply means that we can speak of  $\mathcal{G}(R)$  for any commutative ring  $R$  with identity and that  $\mathcal{G}(R)$  depends functorially on  $R$ .

( Sometimes in the following we will replace  $\text{Spec}(\mathbb{Z})$  by  $\mathbb{Z}$ .) Then  $\mathcal{G}^{(1)}/\mathbb{Z}$  is the kernel of the determinant map  $\det : \mathcal{G}/\mathbb{Z} \rightarrow \mathbb{G}_m/\mathbb{Z}$ . We have the maximal torus  $\mathcal{T}/\mathbb{Z}$  and the Borel subgroup  $\mathcal{B}/\mathbb{Z}$ . We consider the character module  $X^*(\mathcal{T}) = X^*(T \times \mathbb{C})$ . This character module is  $\mathbb{Z}e_1 \oplus \mathbb{Z}e_2$  where

$$e_i : \begin{pmatrix} t_1 & 0 \\ 0 & t_2 \end{pmatrix} \mapsto t_i \tag{4.1}$$

Any character can be written as  $\lambda = n\gamma + d \det$  where  $\gamma = \frac{e_1 - e_2}{2} (\notin X^*(\mathcal{T}) !)$ ,  $\det = e_1 + e_2$  and where  $n \in \mathbb{Z}$ ,  $d \in \frac{1}{2}\mathbb{Z}$  and  $n \equiv 2d \pmod{2}$ . We assume that  $\lambda$  is dominant, i.e.  $n \geq 0$  and drop the assumption that  $n$  should be even.

To any such character  $\lambda$  we want to attach a highest weight module  $\mathcal{M}_\lambda$ . and consider the  $\mathbb{Z}$ - module of polynomials

$$\mathcal{M}_n = \{P(X, Y) \mid P(X, Y) = \sum_{\nu=0}^n a_\nu X^\nu Y^{n-\nu}, a_\nu \in \mathbb{Z}\}.$$

To a polynomial  $P \in \mathcal{M}_n$  we attach the regular function (See Chap. IV)

$$f_P\left(\begin{pmatrix} x & y \\ u & v \end{pmatrix}\right) = P(u, v) \det\left(\begin{pmatrix} x & y \\ u & v \end{pmatrix}\right)^{\frac{n}{2}+d_1} \tag{4.2}$$

and we obviously have

$$f_P\left(\begin{pmatrix} t_1 & w \\ 0 & t_2 \end{pmatrix} \begin{pmatrix} x & y \\ u & v \end{pmatrix}\right) = t_2^n (t_1 t_2)^d f_P\left(\begin{pmatrix} x & y \\ u & v \end{pmatrix}\right) = \lambda^-\left(\begin{pmatrix} t_1 & w \\ 0 & t_2 \end{pmatrix}\right) f_P\left(\begin{pmatrix} x & y \\ u & v \end{pmatrix}\right) \tag{4.3}$$

where  $\lambda^- = -n\gamma + (\frac{n}{2} + d_1) \det = -n\gamma + d \det$  considered as a character on  $\mathcal{B}$ . On this module the group scheme  $\mathcal{G}/\mathbb{Z}$  acts by right translations:

$$\rho_\lambda\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}\right)(f_P)\left(\begin{pmatrix} x & y \\ u & v \end{pmatrix}\right) = f_P\left(\begin{pmatrix} x & y \\ u & v \end{pmatrix}\right) \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

This is a module for the group scheme  $\mathcal{G}/\mathbb{Z}$  it is called the highest weight module for  $\lambda$  and is denoted by  $\mathcal{M}_\lambda$ .

Comment: When we say that  $\mathcal{M}_\lambda$  is a module for the group scheme  $\mathcal{G}/\mathbb{Z}$  we mean nothing more than that for any commutative ring  $R$  with identity we have an action of  $\mathcal{G}(R)$  on  $\mathcal{M}_n \otimes R$ , which is given by (4.2) and depends functorially on  $R$ . We can "evaluate" at  $R = \mathbb{Z}$  and get the  $\Gamma = \text{Gl}_2(\mathbb{Z})$  module  $\mathcal{M}_{\lambda, \mathbb{Z}}$ . (Actually we should not so much distinguish between the  $\text{Gl}_2(\mathbb{Z})$  module  $\mathcal{M}_{\lambda, \mathbb{Z}}$  and  $\mathcal{M}_\lambda$ )

Remark: There is a slightly more sophisticated interpretation of this module. We can form the flag manifold  $\mathcal{B} \backslash \mathcal{G} = \mathbb{P}^1/\mathbb{Z}$  and the character  $\lambda$  yields a line bundle  $\mathcal{L}_{\lambda^-}$ . The group scheme  $\mathcal{G}$  is acting on the pair  $(\mathcal{B} \backslash \mathcal{G}, \mathcal{L}_{\lambda^-})$  and hence on  $H^0(\mathcal{B} \backslash \mathcal{G}, \mathcal{L}_{\lambda^-})$  which is tautologically equal to  $\mathcal{M}_\lambda$  (Borel-Weil theorem).

We can do essentially the same in the case B) . In this case we start from an imaginary quadratic extension  $F/\mathbb{Q}$  and let  $\mathcal{O} = \mathcal{O}_F \subset F$  its ring of integers. We

form the group scheme  $\mathcal{G}/\mathbb{Z} = R_{\mathcal{O}/\mathbb{Z}}(\mathrm{Gl}_2/\mathcal{O})$ . Again  $\mathcal{G}^{(1)}/\mathbb{Z}$  will be the kernel of  $\det : \mathcal{G}/\mathbb{Z} \rightarrow \mathcal{Z}/\mathbb{Z} = R_{\mathcal{O}/\mathbb{Z}}(\mathbb{G}_m)$ . Then  $\mathcal{G}(\mathcal{O}) = \mathrm{Gl}_2(\mathcal{O} \otimes \mathcal{O}) \subset \mathrm{Gl}_2(\mathcal{O}) \times \mathrm{Gl}_2(\mathcal{O})$ . The base change of the maximal torus  $T/\mathbb{Q} \subset \mathcal{G} \times_{\mathbb{Z}} \mathbb{Q}$  is the product  $T_1 \times T_2/F$  where the two factors are the standard maximal tori in the two factors  $\mathrm{Gl}_2/F$ .

We get for the character module CHMsplit

$$X^*(T \times F) = X^*(T_1) \oplus X^*(T_2) = \{n_1\gamma_1 + d_1 \det\} \oplus \{n_2\gamma_2 + d_2 \bar{\det}\} \quad (4.4)$$

where we have to observe the parity conditions  $n_1 \equiv 2d_1 \pmod{2}, n_2 \equiv 2d_2 \pmod{2}$ .

Then the same procedure as in case a) provides a free  $\mathcal{O}$ - module  $\mathcal{M}_\lambda$  with an action of  $\mathcal{G}(\mathbb{Z})$  on it. To see this action we embed the group  $\mathcal{G}(\mathbb{Z}) = \mathrm{Gl}_2(\mathcal{O})$  into  $\mathrm{Gl}_2(\mathcal{O}) \times \mathrm{Gl}_2(\mathcal{O})$  by the map  $g \mapsto (g, \bar{g})$  where  $\bar{g}$  is of course the conjugate. If now our  $\lambda = n_1\gamma_1 + d_1 \det_1 + n_2\gamma_2 + d_2 \det_2 = \lambda_1 + \lambda_2$  then we have our two  $\mathrm{Gl}_2(\mathcal{O})$  modules  $\mathcal{M}_{\lambda_1, \mathcal{O}}, \mathcal{M}_{\lambda_2, \mathcal{O}}$  and this provides the  $\mathrm{Gl}_2(\mathcal{O}) \times \mathrm{Gl}_2(\mathcal{O})$ -module  $\mathcal{M}_{\lambda_1, \mathcal{O}} \otimes \mathcal{M}_{\lambda_2, \mathcal{O}}$ , is now our  $\mathcal{M}_{\lambda, \mathcal{O}}$  is simply the restriction of this tensor product module to  $\mathcal{G}(\mathbb{Z})$ . Sometimes we will also write our character as the sum of the semi simple component and the central component, i.e.  $\lambda = \lambda^{(1)} + \delta = (n_1\gamma_1 + n_2\gamma_2) + (d_1 \det_1 + d_2 \det_2)$ . The relevant term is the semi simple component, the central component not important at all, it only serves to fulfill the parity condition. If we restrict the representation  $\mathcal{M}_\lambda$  to  $\mathcal{G}^{(1)}/\mathbb{Z}$  then the dependence on  $d$  disappears. In other words representations with the same semi simple highest weight component only differ by a twist, the role played by  $\delta$  is marginal.

We return to  $\mathrm{Gl}_2/\mathbb{Z}$ . Given  $\lambda = \lambda^{(1)} + \delta$  we define the dual character as  $\lambda^\vee = \lambda^{(1)} - \delta$ . For our finite dimensional modules we have

$$\mathcal{M}_\lambda^\vee \otimes \mathbb{Q} \xrightarrow{\sim} \mathcal{M}_{\lambda^\vee} \otimes \mathbb{Q} \quad (4.5)$$

If we consider the modules over the integers the above relation is not true. We define the submodule duallambda

$$\mathcal{M}_n^b = \{P(X, Y) \mid P(X, Y) = \sum_{\nu=0}^n \binom{n}{\nu} a_\nu X^\nu Y^{n-\nu}, a_\nu \in \mathbb{Z}\}. \quad (4.6)$$

This is a submodule of  $\mathcal{M}_n$  and the quotient  $\mathcal{M}_n/\mathcal{M}_n^b$  is finite. It is also clear that this submodule is invariant under  $\mathrm{Sl}_2/\mathbb{Z}$ . We introduce some notation

$$e_\nu := X^\nu Y^{n-\nu} \text{ and } e_\nu^b := \binom{n}{\nu} X^{n-\nu} Y^\nu, \quad (4.7)$$

the the  $e_\nu$  (resp.  $e_\nu^b$ ) for a basis of  $\mathcal{M}_n$  (resp.  $\mathcal{M}^b$ ).

An easy calculation shows that the pairing pairMn

$$\langle , \rangle_{\mathcal{M}}: (e_\nu, e_\mu^b) \mapsto \delta_{\nu, \mu} \quad (4.8)$$

is non degenerate over  $\mathbb{Z}$  and invariant under  $\mathrm{Sl}_2/\mathbb{Z}$ . We can also define the the twisted actions of  $\mathcal{G}/\mathbb{Z}$ . Of course we can define the twisted modules  $\mathcal{M}_\lambda^\vee$  and then we get a  $\mathcal{G}/\mathbb{Z}$  invariant non degenerate pairing over  $\mathbb{Z}$  :

$$\langle , \rangle_{\mathcal{M}}: \mathcal{M}_{\lambda^\vee}^b \times \mathcal{M}_\lambda \rightarrow \mathbb{Z}$$

In other words

$$(\mathcal{M}_\lambda)^\flat = \mathcal{M}_{\lambda^\vee}^\flat$$

We always consider  $\mathcal{M}_\lambda^\flat$  as the above submodule of  $\mathcal{M}_\lambda$ .

prinseries

### 4.1.2 The principal series representations

We consider the two real algebraic groups  $G = \mathrm{Gl}_2/\mathbb{R}$  ( case A ) and  $G = R_{\mathbb{C}/\mathbb{R}}(\mathrm{Gl}_2/\mathbb{C})$  ( case B). Let  $T/\mathbb{R}$ , resp.  $B/\mathbb{R}$  be the standard diagonal torus (resp. Borel subgroup of upper triangular matrices). Let us put  $Z/\mathbb{R} = \mathbb{G}_m$  (resp.  $R_{\mathbb{C}/\mathbb{R}}\mathbb{G}_m$ ). We have the determinant  $\det : G/\mathbb{R} \rightarrow Z/\mathbb{R}$  and moreover  $Z/\mathbb{R} = \text{center}(G/\mathbb{R})$ . If we restrict the determinant to the center then this becomes the map  $z \mapsto z^2$ . The kernel of the determinant is denoted by  $G^{(1)}/\mathbb{R}$ , of course  $G^{(1)} = \mathrm{Sl}_2$ , resp.  $R_{\mathbb{C}/\mathbb{R}}(\mathrm{Sl}_2/\mathbb{C})$ . Let us denote by  $\mathfrak{g}, \mathfrak{g}^{(1)}, \mathfrak{t}, \mathfrak{b}, \mathfrak{z}$  the corresponding Lie-algebras.

#### The Cartan decompositions

In both cases we fix a maximal compact subgroup  $K_\infty \subset G^{(1)}(\mathbb{R})$  :

$$K_\infty = e(\phi) = \left\{ \begin{pmatrix} \cos(\phi) & \sin(\phi) \\ -\sin(\phi) & \cos(\phi) \end{pmatrix} \mid \phi \in \mathbb{R} \right\} \text{ and } K_\infty = \left\{ \begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix} \mid \alpha\bar{\alpha} + \beta\bar{\beta} = 1 \right\} \quad (4.9)$$

We define extensions  $\tilde{K}_\infty = Z(\mathbb{R})^{(0)}K_\infty$ , where of course  $Z(\mathbb{R})^{(0)}$  is the connected component of the identity. In both cases the group  $K_\infty$  is the group of fixed points under the Cartan involution  $\Theta_0$  which is given by

$$\Theta_0 : g \mapsto {}^t g^{-1} \text{ resp. } g \mapsto {}^t \bar{g}^{-1} \text{ i.e. } \Theta_0 \left( \begin{pmatrix} a & b \\ c & d \end{pmatrix} \right) = \begin{pmatrix} \bar{d} & -\bar{c} \\ -\bar{b} & \bar{a} \end{pmatrix}. \quad (4.10)$$

This involution induces an involution on  $\mathfrak{g}^{(1)}$  we can extend it to an involution acting on  $\mathfrak{g} = \mathfrak{z} \oplus \mathfrak{g}^{(1)}$ , we let it act trivially on  $\mathfrak{z}$ . Then the fixed point Lie algebra  $\tilde{\mathfrak{k}} = \mathfrak{z} \oplus \mathfrak{k} \subset \mathfrak{z} \oplus \mathfrak{g}^{(1)}$  is the Lie-algebra of  $\tilde{K}_\infty$ .

Here are some arithmetic considerations, they may not be so relevant, but further down we make some choices of a basis in some of these algebras, and these choices can be justified by these considerations.

We can write our group scheme  $G/\mathbb{R}$  as a base extension of a group scheme  $\mathcal{G}/\mathbb{Z}$ , i.e.  $G/\mathbb{R} = \mathcal{G} \times_{\mathbb{Z}} \mathbb{R}$ . For this we simply take  $\mathcal{G}/\mathbb{Z} = \mathrm{Gl}_2/\mathbb{Z}$  in case A). In case B) we take  $\mathcal{G}/\mathbb{Z} = R_{\mathbb{Z}[i]/\mathbb{Z}}(\mathrm{Gl}_2/\mathbb{Z}[i])$ . In the case A) this gives a reductive group scheme over  $\mathbb{Z}$ , in case B) it is only a flat group scheme, but the base extension  $\mathcal{G} \times_{\mathbb{Z}} \mathbb{Z}[1/2]$  is reductive. ( This group scheme over  $\mathbb{Z}$  is not semi-simple since  $\mathbb{Z}[i]$  is ramified at the prime 2.)

Now it is clear that  $\Theta_0$  is actually an automorphism of  $\mathcal{G}/\mathbb{Z}$  and then it follows that the scheme of fixed points is again a group scheme  $\mathcal{K}/\mathbb{Z}$ . If we define  $R = \mathbb{Z}[1/2]$  then  $\mathcal{K} \times_{\mathbb{Z}} R$  is actually semi-simple. (If we replace  $\mathbb{Z}[i]$  by the ring of integers of another imaginary quadratic extension, we have to modify  $R$  accordingly.)

Consequently we see that the all the above Lie-algebras are defined over  $R$ , hence they actually are free  $R$  modules, we denote them by  $\mathfrak{g}_R$  and so on.

The Cartan  $\Theta_0$  involution induces an involution on the Lie algebras  $\mathfrak{g}_R, \mathfrak{g}_R^{(1)}$ , the module decomposes into a  $+$  and a  $-$  eigenspace CaDec

$$\mathfrak{g}_R = \tilde{\mathfrak{k}}_R \oplus \mathfrak{p}_R \text{ and } \mathfrak{g}_R^{(1)} = \mathfrak{k}_R \oplus \mathfrak{p}_R, \quad (4.11)$$

The  $+$  eigenspaces  $\tilde{\mathfrak{k}}_R, \mathfrak{k}_R$  are the Lie-algebras of  $\tilde{\mathcal{K}}, \mathcal{K}$ , both summands in the decompositions are  $\tilde{\mathcal{K}}$ -modules.

The Lie-algebra  $\mathfrak{b}_R$  is not stable under  $\Theta_0$ , it is clear that the intersection

$$\mathfrak{b}_R \cap \Theta_0(\mathfrak{b}_R) = \mathfrak{t}_R,$$

where  $\mathfrak{t}_R$  is the Lie-algebra of the standard maximal torus  $\mathcal{T}/R \subset \mathcal{G}/R$ . This torus is a product (up to isogeny)  $\mathcal{T}/R = \mathcal{Z} \cdot \mathcal{T}^{(1)}/R$ .

In case A) the torus  $\mathcal{T}^{(1)}/R \xrightarrow{\sim} \mathbb{G}_m/R$  and the Cartan involution  $\Theta_0$  acts by  $t \mapsto t^{-1}$ . Therefore it acts by  $-1$  on  $\mathfrak{t}_R^{(1)}$ . We write

$$\mathfrak{t}_R^{(1)} = R \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} = RH \quad (4.12)$$

the generator  $H$  is unique up to an element in  $R^\times$ , i.e. up to a sign and a power of 2.

In case B) the torus  $\mathcal{T}^{(1)}/R$  is (up to isogeny) a product  $\mathcal{T}_s^{(1)} \cdot \mathcal{T}_c^{(1)}/R$  the Cartan involution  $\Theta_0$  acts by  $t \rightarrow t^{-1}$  on the split component  $\mathcal{T}_s^{(1)}$  and by the identity on  $\mathcal{T}_c^{(1)}$ . The Lie-algebra decomposes accordingly into two summands of rank one:

$$\mathfrak{t}_R^{(1)} = R \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \oplus R \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} = RH \oplus RH_i.$$

In both cases the group scheme  $\mathcal{K}$  acts on  $\mathfrak{p}_R$  by the adjoint action, can describe this action explicitly.

In case A) the group scheme  $\mathcal{K}$  is the following group of matrices

$$\mathcal{K} = \left\{ \alpha = \begin{pmatrix} a & b \\ -b & a \end{pmatrix} \mid a^2 + b^2 = 1 \right\}$$

this is a torus over  $R$  which splits over  $R[i]$ . We have

$$\mathfrak{p}_R = RH \oplus R \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = RH \oplus RV$$

and  $\text{Ad}(\alpha)(H) = (a^2 - b^2)H - 2abV$ ,  $\text{Ad}(\alpha)(V) = 2abH + (a^2 - b^2)V$ . Since the torus splits over  $\mathbb{Z}[i]$  we can decompose  $\mathfrak{p} \otimes R[i]$  into weight spaces, we introduce the basis elements

$$P_+ := H - V \otimes i, \quad P_- := H + V \otimes i \in \mathfrak{p} \otimes R[i]$$

then Ppm

$$\text{Ad}(\alpha)P_+ = (a + bi)^2 P_+, \quad \text{Ad}(\alpha)P_- = (a - bi)^2 P_- \quad (4.13)$$

Hence we get - in case A) -the decomposition

$$\mathfrak{g}_R^{(1)} = \mathfrak{k}_R \oplus \mathfrak{p}_R = R \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \oplus RP_+ \oplus RP_- = RY \oplus RP_+ \oplus RP_- \quad (4.14)$$

where the generators are unique up to an element in  $R[i]^\times$ .

In case B) the group scheme  $\mathcal{K}/R$  is semi simple, it contains  $\mathcal{T}_c^{(1)}/R$  as maximal torus. The two  $\mathcal{K}/R$  modules  $\mathfrak{k}_R$  and  $\mathfrak{p}_R$  are highest weight modules of rank 3, since 2 is invertible in  $R$  they are even isomorphic. Again we can decompose them into rank one weight spaces and give almost canonical generators for these weight spaces. basisfkfp The Lie algebra

$$\mathfrak{k}_R = RH_i \oplus R \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \oplus R \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix} = RH_i \oplus RY \oplus F_i.$$

We introduce the elements  $P_{c+} = Y - F_i \otimes i$ ,  $P_{c-} = Y + F_i \otimes i$  and then

$$\mathfrak{k}_R \otimes R[i] = R[i]H_i \oplus R[i]P_{c+} \oplus R[i]P_{c-} \quad (4.15)$$

This is the decomposition into weight spaces under the action of  $\mathcal{T}_c^{(1)}/R$ , the element  $\alpha = \begin{pmatrix} x & 0 \\ 0 & \bar{x} \end{pmatrix}$  acts via the adjoint action

$$\text{Ad}(\alpha)P_{c+} = x^2P_{c+}, \text{Ad}(\alpha)H_i = H_i, \text{Ad}(\alpha)P_{c-} = x^{-2}P_{c-}$$

Essentially the same can be done for  $\mathfrak{p}_R \otimes R[i]$ . We define

$$P_{p,+} = V - \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix} \otimes i, P_{p,-} = V + \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix} \otimes i$$

then we get the weight decomposition basisfp

$$\mathfrak{p}_R \otimes R[i] = R[i]P_{p,+} \oplus R[i]H \oplus R[i]P_{p,-} \quad (4.16)$$

Our aim is to to construct certain irreducible (differentiable) representations of  $G(\mathbb{R})$  together with their "algebraic skeleton" the associated Harish-Chandra modules. Of course any homomorphism  $\eta : Z \rightarrow \mathbb{C}^\times$  yields via composition with the determinant a one dimensional  $G(\mathbb{R})$  module  $\mathbb{C}\eta$ . We want to construct infinite dimensional  $G(\mathbb{R})$  modules.

We start from a continuous homomorphism (a character)  $\chi : T(\mathbb{R}) \rightarrow \mathbb{C}^\times$ , of course this can also be seen as a character  $\chi : B(\mathbb{R}) \rightarrow \mathbb{C}^\times$ . This allows us to define the induced module

$$I_B^G \chi := \{f : G(\mathbb{R}) \rightarrow \mathbb{C} \mid f \in \mathcal{C}_\infty(G(\mathbb{R})), f(bg) = \chi(b)f(g), \forall b \in B(\mathbb{R}), g \in G(\mathbb{R})\} \quad (4.17)$$

where we require that  $f$  should be  $\mathcal{C}_\infty$ . Then this space of functions is a  $G(\mathbb{R})$ -module, the group  $G(\mathbb{R})$  acts by right translations: For  $f \in I_B^G \chi, g \in G(\mathbb{R})$  we put

$$R_g(f)(x) = f(xg)$$

We know that  $G(\mathbb{R}) = B(\mathbb{R}) \cdot \tilde{K}_\infty$ . This implies that a function  $f \in I_B^G \chi$  is determined by its restriction to  $K_\infty$ . In other words we have an identification of vector spaces

$$I_B^G \chi = \{f : \tilde{K}_\infty \rightarrow \mathbb{C} \mid f(t_c k) = \chi(t_c) f(k), t_c \in \tilde{K}_\infty \cap B(\mathbb{R}), k \in \tilde{K}_\infty\}. \quad (4.18)$$

We put  $T_c = B(\mathbb{R}) \cap \tilde{K}_\infty$  and define  $\chi_c$  to be the restriction of  $\chi$  to  $T_c$ . Then the module on the right in the above equation can be written as  $I_{T_c}^{\tilde{K}_\infty} \chi_c$ . By its very definition  $I_{T_c}^{\tilde{K}_\infty} \chi_c$  is only a  $K_\infty$  module.

Inside  $I_{T_c}^{\tilde{K}_\infty} \chi_c$  we have the submodule of vectors of finite type

$${}^\circ I_{T_c}^{\tilde{K}_\infty} \chi_c := \{f \in I_{T_c}^{\tilde{K}_\infty} \chi_c \mid \text{the translates } R_k(f) \text{ lie in a finite dimensional subspace}\} \quad (4.19)$$

The famous Peter-Weyl theorem tells us that all irreducible representations (satisfying some continuity condition) are finite dimensional and occur with finite multiplicity in  $I_{T_c}^{\tilde{K}_\infty} \chi_c$  and therefore we get

$${}^\circ I_{T_c}^{\tilde{K}_\infty} \chi_c = \bigoplus_{\vartheta \in \hat{K}_\infty} V_\vartheta^{m(\vartheta)} = \bigoplus_{\vartheta \in \hat{K}_\infty} {}^\circ I_{T_c}^{\tilde{K}_\infty} \chi_c[\vartheta] \quad (4.20)$$

where  $\hat{K}_\infty$  is the set of isomorphism classes of irreducible representations of  $K_\infty$ , where  $V_\vartheta$  is an irreducible module of type  $\vartheta$  and where  $m(\vartheta)$  is the multiplicity of  $\vartheta$  in  ${}^\circ I_{T_c}^{\tilde{K}_\infty} \chi_c$ . Of course  ${}^\circ I_{T_c}^{\tilde{K}_\infty} \chi_c$  is a submodule  $I_B^G \chi$ , but this submodule is not invariant under the operation of  $G(\mathbb{R})$ , in other words if  $0 \neq f \in {}^\circ I_{T_c}^{\tilde{K}_\infty} \chi_c$  and  $g \in G(\mathbb{R})$  a sufficiently general element then  $R_g(f) \notin {}^\circ I_{T_c}^{\tilde{K}_\infty} \chi_c$ .

We can differentiate the action of  $G(\mathbb{R})$  on  $I_B^G \chi$ . We have the well known exponential map  $\exp : \mathfrak{g} = \text{Lie}(G/\mathbb{R}) \rightarrow G(\mathbb{R})$  and for  $f \in I_B^G \chi, X \in \mathfrak{g}$  we define

$$Xf(g) = \lim_{t \rightarrow 0} \frac{f(g \exp(tX)) - f(g)}{t} \quad (4.21)$$

and it is well known and also easy to see, that this gives an action of the Lie-algebra on  $I_B^G \chi$ , we have  $X_1(X_2 f) - X_2(X_1 f) = [X_1, X_2]f$ . The Lie-algebra is a  $K_\infty$  module under the adjoint action and is obvious that for  $f \in {}^\circ I_{T_c}^{\tilde{K}_\infty} \chi_c[\vartheta]$  the element  $Xf$  lies in  $\bigoplus_{\vartheta' \in \hat{K}_\infty} {}^\circ I_{T_c}^{\tilde{K}_\infty} \chi_c[\vartheta']$  where  $\vartheta'$  runs over the finitely many isomorphism types occurring in  $V_\vartheta \otimes \mathfrak{g}$ .

**Proposition 4.1.1.** *The submodule  ${}^\circ I_{T_c}^{\tilde{K}_\infty} \chi_c \subset I_B^G \chi$  is invariant under the action of  $\mathfrak{g}$ .*

The submodule  ${}^\circ I_{T_c}^{\tilde{K}_\infty} \chi_c$  together with this action of  $\mathfrak{g}$  will now be denoted by  $\mathfrak{J}_B^G \chi$ . We should think of this module as an algebraic skeleton of  $I_B^G \chi$ .

Such a module will be called a  $(\mathfrak{g}, K_\infty)$ -module or a Harish-Chandra module this means that we have an action of the Lie-algebra  $\mathfrak{g}$ , an action of  $K_\infty$  and these two actions satisfy some obvious compatibility conditions.

We also observe that  ${}^\circ I_{T_c}^{\tilde{K}_\infty} \chi_c$  is also invariant under right translation  $R_z$  for  $z \in Z(\mathbb{R})$ . Hence we can extend the action of  $K_\infty$  to the larger group

$\tilde{K}_\infty = K_\infty \cdot Z(\mathbb{R})$ . Then  $\mathfrak{J}_B^G \chi$  becomes a  $(\mathfrak{g}, \tilde{K}_\infty)$  module. Finally observe that in the case A) the element complexcon

$$\mathbf{c} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \notin \tilde{K}_\infty, \tag{4.22}$$

but obviously for  $f \in \mathfrak{J}_B^G \chi$  the element  $R_{\mathbf{c}}(f) \in \mathfrak{J}_B^G \chi$ , hence  $R_{\mathbf{c}}$  induces an involution on  $\mathfrak{J}_B^G$ . We could also say that we can enlarge  $K_\infty$  (resp.  $\tilde{K}_\infty$ ) to subgroups  $K_\infty^*$  (resp.  $\tilde{K}_\infty^*$ ) which contain  $\mathbf{c}$  and contain  $K_\infty$  resp.  $\tilde{K}_\infty$  as subgroups of index two. Then  $\mathfrak{J}_B^G \chi$  also becomes a  $(\mathfrak{g}, \tilde{K}_\infty^*)$  module.

These  $(\mathfrak{g}, \tilde{K}_\infty)$  modules  $\mathfrak{J}_B^G \chi$  are called the principal series modules.

We denote the restriction of  $\chi$  to the central torus  $Z = \left\{ \begin{pmatrix} t & 0 \\ 0 & t \end{pmatrix} \right\}$  by  $\omega_\chi$ .

Then  $Z(\mathbb{R})$  acts on  $\mathfrak{J}_B^G \chi$  by the *central character* character  $\omega_\chi$ , i.e.  $R_z(f) = \omega_\chi(z)f$ . Once we fix the central character, then there is no difference between  $(\mathfrak{g}, \tilde{K}_\infty)$  and  $(\mathfrak{g}, K_\infty)$  modules.

### The decomposition into $K_\infty$ -types

Kutypes

We look briefly at the  $K_\infty$ -module  $\circ I_{T_c}^{\tilde{K}_\infty} \chi_c$ . In case A) the group

$$K_\infty = SO(2) = \left\{ \begin{pmatrix} \cos(\phi) & \sin(\phi) \\ -\sin(\phi) & \cos(\phi) \end{pmatrix} = e(\phi) \right\} \tag{4.23}$$

and  $T_c = T(\mathbb{R}) \cap K_\infty$  is cyclic of order two with generator  $e(\pi)$ . Then  $\chi_c$  is given by an integer  $m \pmod 2$ , i.e.  $\chi_c(e(\pi)) = (-1)^m$ . For any  $n \equiv m \pmod 2$  we define  $\psi_n \in \mathfrak{J}_B^G \chi$  by

$$\psi_n(e(\phi)) = e^{in\phi} \tag{4.24}$$

and then decoKuA

$$\mathfrak{J}_B^G \chi = \bigoplus_{k \equiv m \pmod 2} \mathbb{C} \psi_k \tag{4.25}$$

In the case B) the maximal compact subgroup is

$$U(2) \subset G(\mathbb{R}) = R_{\mathbb{C}/\mathbb{R}}(\mathrm{Gl}_2/\mathbb{C})(\mathbb{R}) \subset \mathrm{Gl}_2(\mathbb{C}) \times \mathbb{G}_2(\mathbb{C})$$

this is the group of real points of the reductive group  $U(2)/\mathbb{R}$ . The intersection

$$T_c = T(\mathbb{R}) \cap K_\infty = \left\{ \begin{pmatrix} e^{i\phi_1} & 0 \\ 0 & e^{i\phi_2} \end{pmatrix} = e(\underline{\phi}) \right\}.$$

The base change  $U(2) \times \mathbb{C} = \mathrm{Gl}_2/\mathbb{C}$  and  $T_c \times \mathbb{C}$  becomes the standard maximal compact torus. The irreducible finite dimensional  $U(2)$ -modules are labelled by dominant highest weights  $\lambda_c = n\gamma_c + d \det \in X^*(T_c \times \mathbb{C})$  (See section ( 4.1.1), here again  $n \geq 0, n \in \mathbb{Z}, n \equiv 2d \pmod 2$  and  $\gamma_c(e(\phi)) = e^{i(\phi_1 - \phi_2)/2}$ .)

We denote these modules by  $\mathcal{M}_{\lambda_c}$  after base change to  $\mathbb{C}$  they become the modules  $\mathcal{M}_{\lambda, \mathbb{C}}$ .

As a subgroup of  $G(\mathbb{R}) \subset \mathrm{Gl}_2(\mathbb{C}) \times \mathbb{G}_2(\mathbb{C})$  our torus is

$$T_c = \left\{ \begin{pmatrix} e^{i\phi_1} & 0 \\ 0 & e^{i\phi_2} \end{pmatrix} \times \begin{pmatrix} e^{-i\phi_1} & 0 \\ 0 & e^{-i\phi_2} \end{pmatrix} \right\} \xrightarrow{\sim} \left\{ \begin{pmatrix} e^{i\phi_1} & 0 \\ 0 & e^{i\phi_2} \end{pmatrix} \right\} \quad (4.26)$$

and the restriction of  $\chi$  to  $T_c$  is of the form

$$\chi_c(e(\underline{\phi})) = e^{ia\phi_1 + ib\phi_2} = e^{\frac{a-b}{2}(\phi_1 - \phi_2)} e^{\frac{a+b}{2}(\phi_1 + \phi_2)} \quad (4.27)$$

and this character is  $(a - b)\gamma_c + \frac{a+b}{2} \det$ . Then we know

decoKuB

$${}^\circ I_{T_c}^{\tilde{K}_\infty} \chi_c = \mathfrak{J}_B^G \chi = \bigoplus_{\substack{\mu_c = k\gamma_c + \frac{a+b}{2} \det; k \equiv (a-b) \\ \text{mod } 2; k \geq |a-b|}} \mathcal{M}_{\mu_c} \quad (4.28)$$

IndInt

### 4.1.3 Intertwining operators

Let  $N(T)$  the normalizer of  $T/\mathbb{R}$ , the quotient  $W = N(T)/T$  is a finite group scheme. The in our case the group  $W(\mathbb{R})$  is cyclic of order 2 and generated by

$$w_0 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

In case a) we have  $W(\mathbb{R}) = W(\mathbb{C})$  in case b) we have

$$G \times_{\mathbb{R}} \mathbb{C} = (\mathrm{Gl}_2 \times \mathrm{Gl}_2)/\mathbb{C} ; T \times_{\mathbb{R}} \mathbb{C} = T_1 \times T_2 ; \text{ and } W(\mathbb{C}) = \mathbb{Z}/2 \times \mathbb{Z}/2$$

where the two factors are generated by  $s_1 = (w_0, 1), s_2 = (1, w_0)$ . The group  $W(\mathbb{R})$  is the group of real points of the Weyl group, the group  $W = W(\mathbb{C})$  is the Weyl group or the absolute Weyl group.

We introduces the special character

$$|\rho|_{\mathbb{R}} : \begin{pmatrix} t_1 & u \\ 0 & t_2 \end{pmatrix} \rightarrow \left| \frac{t_1}{t_2} \right|^{\frac{1}{2}}$$

The group  $W(\mathbb{R})$  acts on  $T(\mathbb{R})$  by conjugation and hence it also acts on the group of characters, we denote this action by  $\chi \mapsto \chi^w$ . We define the twisted action

$$w \cdot \chi = (\chi|\rho|)^w |\rho|^{-1}$$

We recall some well known facts

- i) We have a non degenerate  $(\mathfrak{g}, K_\infty)$  invariant pairing

$$\mathfrak{J}_B^G \chi \times \mathfrak{J}_B^G \chi^{w_0} |\rho|_{\mathbb{R}}^2 \rightarrow \mathbb{C} \omega_\chi^2 \text{ given by } (f_1, f_2) \mapsto \int_{K_\infty} f_1(k) f_2(k) dk \quad (4.29)$$

We define the dual  $\mathfrak{J}_B^{G, \vee} \chi$  of a Harish-Chandra as a submodule of  $\mathrm{Hom}_{\mathbb{C}}(\mathfrak{J}_B^G \chi, \mathbb{C})$ , it consists of those linear maps which vanish on almost all  $K_\infty$  types. It is clear that this is again a  $(\mathfrak{g}, K_\infty)$ -module. The above assertion can be reformulated

ii) We have an isomorphism of  $(\mathfrak{g}, K_\infty)$  modules

$$\mathfrak{I}_B^G \chi \delta_\chi \rightarrow \mathfrak{I}_B^{G,\vee} \chi^{w_0} |\rho|_{\mathbb{R}}^2 \quad (4.30)$$

The group  $T(\mathbb{R}) = T_c \times (\mathbb{R}_{>0}^\times)^2$  and hence we can write any character  $\chi$  in the form

$$\chi(t) = \chi_c(t) |t_1|^{z_1} |t_2|^{z_2} \quad (4.31)$$

where  $z_1, z_2 \in \mathbb{C}$ .

For  $f \in \mathfrak{I}_B^G \chi, g \in G(\mathbb{R})$  we consider the integral

$$T_\infty^{\text{loc}}(f)(g) = \int_{U(\mathbb{R})} f(w_0 u g) du \quad (4.32)$$

It is well known and easy to check that these integrals converge absolutely and locally uniformly for  $\Re(z_1 - z_2) \gg 0$  and it is also not hard to see that they extend to meromorphic functions in the entire  $\mathbb{C}^2$ . We can "evaluate" them at all  $(z_1, z_2)$  by suitably regularizing at poles (for instance taking residues). This needs some explanation. To define the regularized intertwining operator we consider the "deformed" intertwining operator

$$T_\infty^{\text{loc}}(\chi^{w_0} |\gamma|^z) : \mathfrak{I}_B^G \chi^{w_0} |\gamma|^z \rightarrow \mathfrak{I}_B^G \chi |\rho|_{\mathbb{R}}^2 |\gamma|^{-z} \quad (4.33)$$

(See 4.32,  $\chi = \lambda_{\mathbb{R}}^{w_0} |\gamma|^z$ ) and this integral converges if  $\Re(z) \gg 0$ . We have decomposed

$$\mathfrak{I}_B^G \lambda_{\mathbb{R}}^{w_0} |\gamma|^z = \bigoplus_{\vartheta \in \tilde{K}_\infty} \circ I_{T_c}^{\tilde{K}_\infty} \chi_c[\vartheta] = \bigoplus_{\vartheta \in \tilde{K}_\infty} \mathfrak{I}_B^G \lambda_{\mathbb{R}}^{w_0} |\gamma|^z[\vartheta]$$

and our intertwining operator is a direct sum of linear maps between finite dimensional vector spaces

$$c(\lambda_{\mathbb{R}}^{w_0} |\gamma|^z, \vartheta) : \mathfrak{I}_B^G \chi^{w_0} |\gamma|^z[\vartheta] \rightarrow \mathfrak{I}_B^G \chi |\rho|_{\mathbb{R}}^2 |\gamma|^{-z}[\vartheta]$$

The finite dimensional vector spaces do not depend on  $z$  and the  $c(\lambda_{\mathbb{R}}^{w_0} |\gamma|^z, \vartheta)$  can be expressed in terms of values of the  $\Gamma$ -function. Especially they are meromorphic functions in the variable  $z$  (See sl2neu.pdf, ). Hence we can find an integer  $m \geq 0$  such that

$$z^m \mathfrak{I}_B^G \lambda_{\mathbb{R}}^{w_0} |\gamma|^z|_{z=0} : \mathfrak{I}_B^G \chi^{w_0} \rightarrow \mathfrak{I}_B^G \chi |\rho|_{\mathbb{R}}^2$$

is a non zero intertwining operator and this is now our regularized operator  $T_\infty^{\text{loc,reg}}(\chi^{w_0})$ .

iii) The regularized values define non zero intertwining operators

$$T_\infty^{\text{loc,reg}}(\chi) : \mathfrak{I}_B^G \chi \rightarrow \mathfrak{I}_B^G \chi^{w_0} |\rho|_{\mathbb{R}}^2 \quad (4.34)$$

These operators span the one dimensional space of intertwining operators

$$\text{Hom}_{(\mathfrak{g}, K_\infty)}(\mathfrak{I}_B^G \chi, \mathfrak{I}_B^G w_0 \cdot \chi).$$

Finally we discuss the question which of these representations are unitary. This means that we have to find a pairing

$$\psi : \mathfrak{I}_B^G \chi \times \mathfrak{I}_B^G \chi \rightarrow \mathbb{C} \quad (4.35)$$

which satisfies

- a) it is linear in the first and conjugate linear in the second variable
- b) It is positive definite, i.e.  $\psi(f, f) > 0 \forall f \in \mathfrak{I}_B^G \chi$
- c) It is invariant under the action of  $K_\infty$  and Lie-algebra invariant under the action of  $\mathfrak{g}$ , i.e. we have

$$\text{For } f_1, f_2 \in \mathfrak{I}_B^G \chi \text{ and } X \in \mathfrak{g} \text{ we have } \psi(Xf_1, f_2) + \psi(f_1, Xf_2) = 0.$$

We are also interested in quasi-unitary modules. This notion is perhaps best explained if and instead of c) we require

- d) There exists a continuous homomorphism (a character)  $\eta : G(\mathbb{R}) \rightarrow \mathbb{R}^\times$  such that  $\psi(gf_1, gf_2) = \eta(g)\psi(f_1, f_2)$ ,  $\forall g \in G(\mathbb{R})$ ,  $f_1, f_2 \in \mathfrak{I}_B^G \chi$

It is clear that a non zero pairing  $\psi$  which satisfies a) and c) is the same thing as a non zero  $(\mathfrak{g}, K_\infty)$ -module linear map

$$i_\psi : \mathfrak{I}_B^G \chi \rightarrow \overline{(\mathfrak{I}_B^G \chi)^\vee} \quad (4.36)$$

by definition  $i_\psi$  is a conjugate linear map from  $\mathfrak{I}_B^G \chi$  to  $(\mathfrak{I}_B^G \chi)^\vee$ . The map  $i_\psi$  and the pairing  $\psi$  are related by the formula  $\psi(v_1, v_2) = i_\psi(v_2)(v_1)$ .

Of course we know that (See (4.30))

$$\overline{(\mathfrak{I}_B^G \chi)^\vee} \xrightarrow{\sim} \mathfrak{I}_B^G \chi^{w_0} |\rho|_{\mathbb{R}}^2 \delta_\chi^{-1} \quad (4.37)$$

and we find such an  $i_\psi$  if

$$\chi = \overline{\chi^{w_0} |\rho|_{\mathbb{R}}^2 \delta_\chi^{-1}} \text{ or } \chi^{w_0} |\rho|_{\mathbb{R}}^2 = \overline{\chi^{w_0} |\rho|_{\mathbb{R}}^2 \delta_\chi^{-1}} \quad (4.38)$$

We write our  $\chi$  in the form (8.121). A necessary condition for the existence of a hermitian form is of course that all  $|\omega_\chi(x)| = 1$  for  $x \in Z(\mathbb{R})$  and this means that  $\Re(z_1 + z_2) = 0$ , hence we write

$$z_1 = \sigma + i\tau_1, z_2 = -\sigma + i\tau_2 \quad (4.39)$$

Then the two conditions in (4.38) simply say

$$(\text{un}_1) : \sigma = \frac{1}{2} \text{ or } (\text{un}_2) : \tau_1 = \tau_2 \text{ and } \chi_c = \chi^{w_0} \quad (4.40)$$

In both cases we can write down a pairing which satisfies a) and c). We still have to check b). In the first case, i.e.  $\sigma = \frac{1}{2}$  we can take the map  $i_\psi = \text{Id}$  and then we get for  $f_1, f_2 \in \mathfrak{I}_B^G \chi$  the formula

$$\psi(f_1, f_2) = \int_{K_\infty} f_1(k) \overline{f_2(k)} dk \quad (4.41)$$

and this is clearly positive definite. These are the representation of the unitary principal series.

In the second case we have to use the intertwining operator in (4.34) and write

$$\psi(f_1, f_2) = T_\infty^{\text{loc,reg}}(f_2)(f_1) \quad (4.42)$$

Now it is not clear whether this pairing satisfies b). This will depend on the parameter  $\sigma$ . We can twist by a character  $\eta : Z(\mathbb{R}) \rightarrow \mathbb{C}^\times$  and achieve that  $\chi_c = 1, \tau_1 = \tau_2 = 0$ . We know that for  $\sigma = \frac{1}{2}$  the intertwining operator  $T_\infty^{\text{loc}}$  is regular at  $\chi$  and since in addition under these conditions  $\mathfrak{J}_B^G \chi$  is irreducible we see that

$$T_\infty^{\text{loc}}(\chi) = \alpha \text{Id with } \alpha \in \mathbb{R}_{>0}^\times \quad (4.43)$$

Since we now are in case A) and B) at the same time we see that the two pairings defined by the rule in case (un<sub>1</sub>) and (un<sub>2</sub>) differ by a positive real number hence the pairing defined in (4.42) is positive definite if  $\sigma = \frac{1}{2}$ .

But now we can vary  $\sigma$ . It is well known that  $\mathfrak{J}_B^G \chi$  stays irreducible as long as  $0 < \sigma < 1$  (See next section) and since  $T_\infty^{\text{loc}}(\chi)(f)(f)$  varies continuously we see that (4.42) defines a positive definite hermitian product on  $\mathfrak{J}_B^G \chi$  as long as  $0 < \sigma < 1$ . This is the supplementary series. What happens if we leave this interval will be discussed in the next section.

nontriv

#### 4.1.4 Reducibility and representations with non trivial cohomology

As usual we denote by  $\rho \in X^*(T) \otimes \mathbb{Q}$  the half sum of positive roots we have  $\rho = \gamma$  ( resp.  $\rho = \gamma_1 + \gamma_2 \in X^*(T) \otimes \mathbb{Q}$  in case A) ( resp. B) ).

For any character  $\lambda \in X^*(T \times \mathbb{C})$  we define  $\lambda_{\mathbb{R}}$  to be the restriction (or evaluation)

$$\lambda_{\mathbb{R}} : T(\mathbb{R}) \rightarrow \mathbb{C}^\times.$$

This provides a homomorphism  $B(\mathbb{R}) \rightarrow T(\mathbb{R})$  and hence we get the Harish-Chandra modules  $\mathfrak{J}_B^G \lambda_{\mathbb{R}}$  which are of special interest for our subject namely the cohomology of arithmetic groups.

We just mention the fact that  $\mathfrak{J}_B^G \chi$  is always irreducible unless  $\chi = \lambda_{\mathbb{R}}$  (See sl2neu.pdf, Condition (red)).

We return to the situation discussed in section (4.1.1), especially we reintroduce the field  $F/\mathbb{Q}$ . Then we have  $X^*(T \times F) = X^*(T \times \mathbb{C})$  and hence  $\lambda \in X^*(T \times F)$ . We assume that  $\lambda$  is dominant, i.e.  $n \geq 0$  in case A) or  $n_1, n_2 \geq 0$  in case B). In this case we realized our modules  $\mathcal{M}_\lambda$  as submodules in the algebra of regular functions on  $\mathcal{G}/\mathbb{Z}$  and if we look at the definition (See (4.3)) we see immediately that  $\mathcal{M}_{\lambda, \mathbb{C}} \subset \mathfrak{J}_B^G \lambda_{\mathbb{R}}^{w_0}$  and hence we get an exact sequence of  $(\mathfrak{g}, K_\infty)$  modules seq

$$0 \rightarrow \mathcal{M}_{\lambda, \mathbb{C}} \rightarrow \mathfrak{J}_B^G \lambda_{\mathbb{R}}^{w_0} \rightarrow \mathcal{D}_\lambda \rightarrow 0 \quad (4.44)$$

Hence we see that  $\mathfrak{J}_B^G \lambda_{\mathbb{R}}^{w_0}$  is not irreducible. We can also look at the dual sequence. Here we recall that we wrote  $\lambda = n\gamma + d \det$ . Then we will see later that  $\mathcal{M}_{\lambda, \mathbb{C}}^{\vee} = \mathcal{M}_{\lambda - 2d \det, \mathbb{C}}$ . Hence after twisting the dual sequence becomes

$$0 \rightarrow \mathcal{D}_{\lambda}^{\vee} \otimes \det_{\mathbb{R}}^{2d} \rightarrow \mathfrak{J}_B^{G, \vee} \lambda_{\mathbb{R}}^{w_0} \rightarrow \mathcal{M}_{\lambda, \mathbb{C}} \rightarrow 0 \quad (4.45)$$

Equation (4.30) yields  $\mathfrak{J}_B^{G, \vee} \lambda_{\mathbb{R}}^{w_0} \xrightarrow{\sim} \mathfrak{J}_B^G \chi | \rho |_{\mathbb{R}}^2$  and our second sequence becomes

$$0 \rightarrow \mathcal{D}_{\lambda}^{\vee} \otimes \det_{\mathbb{R}}^{2d} \rightarrow \mathfrak{J}_B^G \lambda_{\mathbb{R}} | \rho |_{\mathbb{R}}^2 \rightarrow \mathcal{M}_{\lambda, \mathbb{C}} \rightarrow 0 \quad (4.46)$$

Now we consider the two middle terms in the two exact sequences (4.44, 4.46) above. The equation (4.34) claims that we have two non zero *regularized* intertwining operators

$$T_{\infty}^{\text{loc, reg}}(\lambda_{\mathbb{R}}^{w_0}) : \mathfrak{J}_B^G \lambda_{\mathbb{R}}^{w_0} \rightarrow \mathfrak{J}_B^G \lambda_{\mathbb{R}} | \rho |_{\mathbb{R}}^2 ; T_{\infty}^{\text{loc, reg}}(\lambda_{\mathbb{R}} | \rho |_{\mathbb{R}}^2) : \mathfrak{J}_B^G \lambda_{\mathbb{R}} | \rho |_{\mathbb{R}}^2 \rightarrow \mathfrak{J}_B^G \lambda_{\mathbb{R}}^{w_0} \quad (4.47)$$

If we now look more carefully at our two regularized intertwining operators above then a simple computation yields (see sl2neu.pdf)

**Proposition 4.1.2.** *The kernel of  $T_{\infty}^{\text{loc, reg}}(\lambda_{\mathbb{R}}^{w_0})$  is  $\mathcal{M}_{\lambda, \mathbb{C}}$  and this operator induces an isomorphism*

$$\bar{T}(\lambda_R) : \mathcal{D}_{\lambda} \xrightarrow{\sim} \mathcal{D}_{\lambda}^{\vee} \otimes \det_{\mathbb{R}}^{2d}$$

(Remember  $\lambda$  is dominant) The kernel of  $T_{\infty}^{\text{loc, reg}}(\lambda_{\mathbb{R}} | \rho |_{\mathbb{R}}^2)$  is  $\mathcal{D}_{\lambda}^{\vee} \otimes \det_{\mathbb{R}}^{2d}$  and it induces an isomorphism of  $\mathcal{M}_{\lambda, \mathbb{C}}$ .

The module  $\mathfrak{J}_B^G \chi$  is reducible if  $T_{\infty}^{\text{loc, reg}}(\chi)$  not an isomorphism and this happens if and only if  $\chi = \lambda_{\mathbb{R}}$  or  $\lambda_{\mathbb{R}}^{w_0} | \rho |_{\mathbb{R}}^2$  and  $\lambda$  dominant. (There is one exception to the converse of the above assertion, namely in the case A) and  $\sigma = \frac{1}{2}$  and  $\chi_c^{w_0} \neq \chi_c$ .)

### Unitarity

For us it is of relevance to know whether we have a positive definite hermitian form on the  $(\mathfrak{g}, K_{\infty})$ -modules  $\mathcal{D}_{\lambda}$ . To discuss this question we treat the cases A) and B) separately.

We look at the decomposition into  $K_{\infty}$ -types. (See (4.25)) In case A) (See (4.25)) it is clear that  $\mathcal{M}_{\lambda, \mathbb{C}}$  is the direct sum of the  $K_{\infty}$  types  $\mathbb{C}\psi_l$  with  $|l| \leq n$ . Hence KTA

$$\mathcal{D}_{\lambda} = \bigoplus_{k \leq -n-2, k \equiv m(2)} \mathbb{C}\psi_k \oplus \bigoplus_{k \geq n+2, k \equiv m(2)} \mathbb{C}\psi_k = \mathcal{D}_{\lambda}^{-} \oplus \mathcal{D}_{\lambda}^{+} \quad (4.48)$$

**Proposition 4.1.3.** *The representations  $\mathcal{D}_{\lambda}^{-}, \mathcal{D}_{\lambda}^{+}$  are irreducible, these are the discrete series representations. The element  $\mathbf{c}$  interchanges  $\mathcal{D}_{\lambda}^{-}, \mathcal{D}_{\lambda}^{+}$  hence  $\mathcal{D}_{\lambda}$  is an irreducible  $(\mathfrak{g}, \tilde{K}_{\infty}^*)$  module.*

The operator  $\bar{T}(\lambda_R)$  induces a quasi-unitary structure on the  $(\mathfrak{g}, \tilde{K}_{\infty})$ -module  $\mathcal{D}_{\lambda}$ . The two sets of  $K_{\infty}$  types occurring in  $\mathcal{M}_{\lambda, \mathbb{C}}$  and in  $\mathcal{D}_{\lambda}$  (resp.) are disjoint.

*Proof.* Remember that as a vector space  $\mathcal{D}_\lambda^\vee \otimes \det_{\mathbb{R}}^{2d} = \mathcal{D}_\lambda^\vee$ , only the way how  $\tilde{K}_\infty$  acts is twisted by  $\det_{\mathbb{R}}^{2d}$ . Then the form

$$h_\psi(f_1, f_2) = T_\infty^{\text{loc,reg}}(\lambda_{\mathbb{R}}^{w_0})(f_2)(f_1) \quad (4.49)$$

defines a quasi invariant hermitian form. It is positive definite (for more details see sl2neu.pdf).  $\square$

A similar argument works in case B). We restrict the  $\text{Gl}_2(\mathbb{C}) \times \text{Gl}_2(\mathbb{C})$  module  $\mathcal{M}_{\lambda, \mathbb{C}}$  to  $U(2) \times U(2)$  then it becomes the highest weight module  $\mathcal{M}_{\lambda_c} = \mathcal{M}_{\lambda_{1,c}} \otimes \mathcal{M}_{\lambda_{2,c}}$ . (See 4.1.1) Under the action of  $U(2) \subset U(2) \times U(2)$  it decomposes into  $U(2)$  types according to the Clebsch-Gordan formula CG

$$\mathcal{M}_{\lambda_c}|_{U(2)} = \bigoplus_{\substack{\mu_c = k\gamma_c + \frac{d_1+d_2}{2} \det; k \equiv (n_1-n_2) \pmod{2}; n_1+n_2 \geq k \geq |n_1-n_2|}} \mathcal{M}_{\mu_c} \quad (4.50)$$

Hence we get KTB

$$\mathcal{D}_{\lambda_c}|_{U(2)} = \bigoplus_{\substack{\mu_c = k\gamma_c + \frac{d_1+d_2}{2} \det; k \equiv (n_1-n_2) \pmod{2}; k \geq n_1+n_2+2}} \mathcal{M}_{\mu_c} \quad (4.51)$$

Again we have unitary

**Proposition 4.1.4.** *The operator  $T_\infty^{\text{loc,reg}}(\lambda_{\mathbb{R}}^{w_0})$  induces an isomorphism*

$$\bar{T}(\lambda_R) : \mathcal{D}_\lambda \xrightarrow{\sim} \mathcal{D}_\lambda^\vee \otimes \det_{\mathbb{R}}^{2d}$$

*The  $(\mathfrak{g}, K_\infty)$  modules are irreducible.*

*The operator  $T_\infty^{\text{loc,reg}}(\lambda_{\mathbb{R}}^{w_0})$  induces the structure of a quasi-unitary module on  $\mathcal{D}_\lambda$  if and only if  $n_1 = n_2$ . This is the only case when we have a quasi-unitary structure on  $\mathcal{D}_\lambda$ . The two sets of  $K_\infty$  types occurring in  $\mathcal{M}_{\lambda, \mathbb{C}}$  and in  $\mathcal{D}_\lambda$  (resp.) are disjoint.*

The Weyl  $W$  group acts on  $T$  by conjugation, hence on  $X^*(T \times \mathbb{C})$  and we define the twisted action by

$$s \cdot \lambda = s(\lambda + \rho) - \rho \quad (4.52)$$

Given a dominant  $\lambda$  we may consider the four characters  $w \cdot \lambda, w \in W(\mathbb{C}) = W$  and the resulting induced modules  $\mathcal{I}_B^G w \cdot \lambda_{\mathbb{R}}$ . We observe (notation from (4.1.1))

$$\begin{aligned} s_1 \cdot (n_1\gamma + d_1 \det + n_2\bar{\gamma} + d_2\overline{\det}) &= (-n_1 - 2)\gamma + d_1 \det + n_2\bar{\gamma} + d_2\overline{\det} \\ s_2 \cdot (n_1\gamma + d_1 \det + n_2\bar{\gamma} + d_2\overline{\det}) &= n_1\gamma + d_1 \det + (-n_2 - 2)\bar{\gamma} + d_2\overline{\det} \end{aligned} \quad (4.53)$$

Looking closely we see that that the  $K_\infty$  types occurring in  $\mathcal{I}_B^G s_1 \cdot \lambda$  or  $\mathcal{I}_B^G s_2 \cdot \lambda$  are exactly those which occur in  $\mathcal{D}_\lambda$ . This has a simple explanation, we have

exiso

**Proposition 4.1.5.** *For a dominant character  $\lambda$  we have isomorphisms between the  $(\mathfrak{g}, K_\infty)$  modules*

$$\mathcal{D}_\lambda \xrightarrow{\sim} \mathfrak{I}_B^G s_1 \cdot \lambda, \mathcal{D}_\lambda \xrightarrow{\sim} \mathfrak{I}_B^G s_2 \cdot \lambda. \tag{4.54}$$

The resulting isomorphism  $\mathfrak{I}_B^G s_1 \cdot \lambda \xrightarrow{\sim} \mathfrak{I}_B^G s_2 \cdot \lambda$  is of course given by  $T_\infty^{\text{loc}}(s_1 \cdot \lambda)$ .

**Interlude:** Here we see a fundamental difference between the two cases A) and B). In the second case the infinite dimensional subquotients of the induced representations are again induced representations. In the case A) this is not so, the representations  $\mathcal{D}_\lambda^\pm$  are not isomorphic to representations induced from the Borel subgroup.

These representation  $\mathcal{D}_\lambda^\pm$  are called *discrete series* representations and we want to explain briefly why. Let  $G$  be the group of real points of a reductive group over  $\mathbb{R}$  for example our  $G = G(\mathbb{R})$ , here we allow both cases. Let  $Z$  be the center of  $G$ , it can be written as  $Z_0(\mathbb{R}) \cdot Z_c$  where  $Z_c$  is maximal compact and  $Z_0 = (\mathbb{R}_{>0}^\times)^t$ . Let  $\omega^{(0)} : Z_0 \rightarrow \mathbb{R}_{>0}^\times$  be a character. Then we define the space

$$\mathcal{C}_\infty(G, \omega_R) := \{f \in \mathcal{C}(G) \mid f(zg) = \omega^{(0)}(z)f(g) ; \forall z \in Z_0, g \in G\} \tag{4.55}$$

and we define the subspace

$$L_\infty^2(G, \omega_R) := \{f \in \mathcal{C}_\infty(G, \omega_R) \mid \int_G f(g)\overline{f(g)}(\omega^{(0)}(g))^{-2}dg < \infty\} \tag{4.56}$$

where of course  $dg$  is a Haar measure. As usual  $L^2(G, \omega_R)$  will be the Hilbert space obtained by completion. This Hilbert space only depends in a very mild way on the choice of  $\omega^{(0)}$  we can find a character  $\delta : G \rightarrow \mathbb{R}_{>0}^\times$  such that  $\omega^{(0)}\delta|_{Z_0} = 1$ . Then  $f \mapsto f\delta$  provides an isomorphism  $L^2(G, \omega^{(0)}) \xrightarrow{\sim} L^2(G/Z_0)$ .

We have an action of  $G \times G$  on  $L^2(G, \omega^{(0)})$  by left and right translations. Then Harish-Chandra has investigated the question how this "decomposes" into irreducible submodules. Let  $\hat{G}_{\omega^{(0)}}$  be the set of isomorphism classes of irreducible unitary representations of  $G$ .

Then Harish-Chandra shows that there exist a positive measure  $\mu$  on  $\hat{G}_{\omega^{(0)}}$  and a measurable family  $H_\xi$  of irreducible unitary representations of  $G$  such that

$$L^2(G, \omega_R) = \int_{\hat{G}_{\omega_R}} H_\xi \otimes \overline{H_\xi} \mu(d\xi) \tag{4.57}$$

( If instead of a semi simple Lie group we take a finite group  $G$  then this is the fundamental theorem of Frobenius that the group ring  $\mathbb{C}[G] = \oplus_\theta V_\theta \otimes V_\theta^\vee$  where  $V_\theta$  are the irreducible representations.)

If we are in the case A), the sets consisting of just one point  $\{\mathcal{D}_\lambda^\pm\}$  have strictly positive measure, i.e.  $\mu(\{\mathcal{D}_\lambda^\pm\}) > 0$ . This means that the irreducible unitary  $G \times G$  modules  $\mathcal{D}_\lambda^\pm \otimes \mathcal{D}_{\lambda^\vee}^\pm$  occur as direct summand (i.e. discretely in  $L^2(G)$ ).

Such irreducible direct summands do not exist in the case B), in this case for any  $\xi \in \hat{G}$  we have  $\mu(\{\xi\}) = 0$ .

We return to the sequences (4.44),(4.46). We claim that both sequences do not split as sequences of  $(\mathfrak{g}, K_\infty)$ -modules. Of course it follows from the above proposition that these sequences split canonically as sequence of  $K_\infty$  modules. But then it follows easily that complementary summand is not invariant under the action of  $\mathfrak{g}$ . This means that the sequences provide non trivial classes in  $\text{Ext}_{(\mathfrak{g}, K_\infty)}^1(\mathcal{D}_\lambda, \mathcal{M}_{\lambda, \mathbb{C}})$  and hence these  $\text{Ext}^\bullet$  modules are interesting.

The general principles of homological algebra teach us that we can understand these extension groups in terms of relative Lie-algebra cohomology. Let  $\mathfrak{k}$  resp.  $\tilde{\mathfrak{k}}$  be the Lie-algebras of  $K_\infty$  resp.  $\tilde{K}_\infty$  the group  $\tilde{K}_\infty$  acts on  $\mathfrak{g}, \mathfrak{k}$  via the adjoint action (see 1.1.4) We start from a  $(\mathfrak{g}, \tilde{K}_\infty)$  module  $\mathfrak{I}_{B\chi}^G$  and a module  $\mathcal{M}_{\lambda, \mathbb{C}}$ .

Our goal is to compute the cohomology of the complex (See 8.1.2)

$$\text{Hom}_{\tilde{K}_\infty}(\Lambda^\bullet(\mathfrak{g}/\tilde{\mathfrak{k}}), \mathfrak{I}_{B\chi}^G \otimes \mathcal{M}_{\lambda, \mathbb{C}}). \quad (4.58)$$

There is an obvious condition for the complex to be non zero. The group  $Z(\mathbb{R}) \subset \tilde{K}_\infty$  acts trivially on  $\mathfrak{g}/\tilde{\mathfrak{k}}$  and hence we see that the complex is trivial unless we have

$$\omega_\chi^{-1} = \lambda_{\mathbb{R}}|_{Z(\mathbb{R})}$$

we assume that this relation holds.

We will derive a formula for these cohomology modules, which is a special case of a formula of Delorme. It will also be discussed in Chap. III. An element  $\omega \in \text{Hom}_{\tilde{K}_\infty}(\Lambda^n(\mathfrak{g}/\tilde{\mathfrak{k}}), \mathfrak{I}_{B\chi}^G \otimes \mathcal{M}_{\lambda, \mathbb{C}})$  attaches to any  $n$  tuple  $v_1, \dots, v_n$  of elements in  $\mathfrak{g}/\tilde{\mathfrak{k}}$  an element

$$\omega(v_1, \dots, v_n) \in \mathfrak{I}_{B\chi}^G \otimes \mathcal{M}_{\lambda, \mathbb{C}} \quad (4.59)$$

such that  $\omega(\text{Ad}(k)v_1, \dots, \text{Ad}(k)v_n) = k\omega(v_1, \dots, v_n)$  for all  $k \in \tilde{K}_\infty$ .

By construction

$$\omega(v_1, \dots, v_n) = \sum f_\nu \otimes m_\nu \text{ where } f_\nu \in \mathfrak{I}_{B\chi}^G, m_\nu \in \mathcal{M}_{\lambda, \mathbb{C}}$$

and  $f_\nu$  is a function in  $\mathcal{C}_\infty$  which is determined by its restriction to  $\tilde{K}_\infty$  ( and this restriction is  $\tilde{K}_\infty$  finite). We can evaluate this function at the identity  $e_G \in G(\mathbb{R})$  and then

$$\omega(v_1, \dots, v_n)(e_G) = \sum f_\nu(e) \otimes m_\nu \in \mathbb{C}\chi \otimes \mathcal{M}_{\lambda, \mathbb{C}}$$

The  $\tilde{K}_\infty$  invariance (4.59) implies that  $\omega$  is determined by this evaluation at  $e_G$ . Let  $\tilde{K}_\infty^T = T(\mathbb{R}) \cap \tilde{K}_\infty = Z(\mathbb{R}) \cdot T_c$ . Then it is clear that

$$\omega^* : \{v_1, \dots, v_n\} \mapsto \omega(v_1, \dots, v_n)(e_G) \quad (4.60)$$

is an element in

$$\omega^* \in \text{Hom}_{\tilde{K}_\infty^T}(\Lambda^n(\mathfrak{g}/\tilde{\mathfrak{k}}), \mathbb{C}\chi \otimes \mathcal{M}_{\lambda, \mathbb{C}}) \quad (4.61)$$

and we have: The map  $\omega \mapsto \omega^*$  is an isomorphism of complexes iso1

$$\text{Hom}_{\tilde{K}_\infty}(\Lambda^\bullet(\mathfrak{g}/\tilde{\mathfrak{k}}), \mathfrak{I}_{B\chi}^G \otimes \mathcal{M}_{\lambda, \mathbb{C}}) \xrightarrow{\sim} \text{Hom}_{\mathfrak{c}}(\Lambda^\bullet(\mathfrak{g}/\tilde{\mathfrak{k}}), \mathbb{C}\chi \otimes \mathcal{M}_{\lambda, \mathbb{C}}) \quad (4.62)$$

The Lie algebra  $\mathfrak{g}$  can be written as a sum of  $\mathfrak{c}$  invariant submodules

$$\mathfrak{g} = \mathfrak{b} + \tilde{\mathfrak{k}} = \mathfrak{t} + \mathfrak{u} + \tilde{\mathfrak{k}} \quad (4.63)$$

in case B) this sum is not direct, we have  $\mathfrak{b} \cap \tilde{\mathfrak{k}} = \mathfrak{t} \cap \tilde{\mathfrak{k}} = \mathfrak{c}$  and hence we get the direct sum decomposition into  $\tilde{K}_\infty^T$ -invariant subspaces

$$\mathfrak{g}/\tilde{\mathfrak{k}} = \mathfrak{t}/\mathfrak{c} \oplus \mathfrak{u}. \quad (4.64)$$

We get an isomorphism of complexes isodel

$$\mathrm{Hom}_{\tilde{K}_\infty}(\Lambda^\bullet(\mathfrak{g}/\tilde{\mathfrak{k}}), \mathfrak{J}_B^G \otimes \mathcal{M}_{\lambda, \mathbb{C}}) \xrightarrow{\sim} \mathrm{Hom}_{\tilde{K}_\infty^T}(\Lambda^\bullet(\mathfrak{t}/\tilde{\mathfrak{k}}), \mathbb{C}\chi \otimes \mathrm{Hom}(\Lambda^\bullet(\mathfrak{u}), \mathcal{M}_{\lambda, \mathbb{C}})) \quad (4.65)$$

the complex on the left is isomorphic to the total complex of the double complex on the right.

**Intermission: The theorem of Kostant** The next step is the computation of the cohomology of the complex  $\mathrm{Hom}(\Lambda^\bullet(\mathfrak{u}), \mathcal{M}_{\lambda, \mathbb{C}})$ .

Case A). Our group is  $G/\mathbb{Q} = \mathrm{Gl}_2/\mathbb{Q}$ . Then  $\mathfrak{u} = \mathbb{Q}E_+$  where  $E_+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$  and our module  $\mathcal{M}_{\lambda, \mathbb{Q}}$  has a decomposition into weight spaces

$$\mathcal{M}_{\lambda, \mathbb{Q}} = \bigoplus_{\nu=1}^{\nu=n-\nu} \mathbb{Q}X^{n-\nu}Y^\nu = \bigoplus_{\mu=-n, \mu \equiv n(2)}^{\mu=n} \mathbb{Q}e_\mu. \quad (4.66)$$

The torus  $T^{(1)} = \left\{ \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix} \right\}$  acts on  $e_\mu = X^{n-\nu}Y^\nu$  by

$$\rho_\lambda \left( \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix} \right) e_\mu = t^\mu e_\mu \quad (4.67)$$

We also have the action of the Lie algebra on  $\mathcal{M}_{\lambda, \mathbb{Q}}$  (See section ??) and by definition we get

$$d(\rho_\lambda)(E_+)e_\mu = E_+e_\mu = \frac{n-\mu}{2}e_{\mu+2} \quad (4.68)$$

Now we can write down our complex  $\mathrm{Hom}(\Lambda^\bullet(\mathfrak{u}), \mathcal{M}_{\lambda, \mathbb{C}})$  very explicitly. Let  $E_+^\vee \in \mathrm{Hom}(\mathfrak{u}, \mathbb{Q})$  be the element  $E_+^\vee(E_+) = 1$  then the complex becomes

$$0 \rightarrow \bigoplus_{\mu=-n, \mu \equiv n(2)}^{\mu=n} \mathbb{Q}e_\mu \xrightarrow{d} \bigoplus_{\mu=-n, \mu \equiv n(2)}^{\mu=n} \mathbb{Q}E_+^\vee \otimes e_\mu \rightarrow 0 \quad (4.69)$$

where  $d(e_\mu) = \frac{n-\mu}{2}E_+^\vee \otimes e_{\mu+2}$ . This gives us a decomposition of our complex into two sub complexes

$$\mathrm{Hom}(\Lambda^\bullet(\mathfrak{u}), \mathcal{M}_{\lambda, \mathbb{C}}) = \mathbb{H}^\bullet(\mathfrak{u}, \mathcal{M}_{\lambda, \mathbb{Q}}) \oplus AC^\bullet \quad (4.70)$$

where  $AC^\bullet$  as acyclic (it has no cohomology) and in

$$\mathbb{H}^\bullet(\mathfrak{u}, \mathcal{M}_{\lambda, \mathbb{Q}}) = \{0 \rightarrow \mathbb{Q}e_n \xrightarrow{d} \mathbb{Q}E_+^\vee \otimes e_{-n} \rightarrow 0\} \quad (4.71)$$

where the differential  $d$  is zero. Hence we get

$$H^\bullet(\mathfrak{u}, \mathcal{M}_{\lambda, \mathbb{Q}}) = H^\bullet(\text{Hom}(\Lambda^\bullet(\mathfrak{u}), \mathcal{M}_{\lambda, \mathbb{Q}})) = \mathbb{H}^\bullet(\mathfrak{u}, \mathcal{M}_{\lambda, \mathbb{Q}}) \quad (4.72)$$

We notice that the torus  $T$  acts on  $H^\bullet(\mathfrak{u}, \mathcal{M}_{\lambda, \mathbb{Q}})$  (The Borel subgroup  $B$  acts on the complex  $\text{Hom}(\Lambda^\bullet(\mathfrak{u}), \mathcal{M}_{\lambda, \mathbb{Q}})$  but since the Lie algebra cohomology is the derived functor of taking invariants under  $U$  (elements annihilated by  $\mathfrak{u}$ ) it follows that this action is trivial on  $U$ ).

Hence we see that  $T$  acts by the character  $\lambda$  on  $\mathbb{Q} e_n = H^0(\mathfrak{u}, \mathcal{M}_{\lambda, \mathbb{Q}})$  and by the character  $\lambda^- - \alpha = w_0 \cdot \lambda = \lambda^{w_0} - 2\rho$  on  $\mathbb{Q} E_+^\vee \otimes e_{-n} = H^1(\mathfrak{u}, \mathcal{M}_{\lambda, \mathbb{Q}})$ . Here we see the simplest example of the famous theorem of Kostant which will be discussed in Chap. III 6.1.3.

We discuss the case B). Again we want that our group  $G/\mathbb{R} = R_{\mathbb{C}/\mathbb{R}}(\text{Gl}_2/\mathbb{C})$  is a base change from a group  $G/\mathbb{Q}$  denoted by the same letter. We need an imaginary quadratic extension  $F/\mathbb{Q}$  and put  $G/\mathbb{Q} = R_{F/\mathbb{Q}}(\text{Gl}_2/F)$ . We choose a dominant weight  $\lambda = \lambda_1 + \lambda_2 = n_1\gamma_1 + d_1 \det_1 + n_2\gamma_2 + d_2 \det_2$  and then  $\mathcal{M}_{\lambda, F} = \mathcal{M}_{\lambda_1, F} \otimes \mathcal{M}_{\lambda_2, F}$  is an irreducible representation of  $G \times_{\mathbb{Q}} F = \text{Gl}_2 \times \text{Gl}_2/F$ . Now we have  $\mathfrak{u} \otimes F = FE_+^1 \oplus FE_+^2$ . Then basically the same computation yields:

The cohomology  $H^\bullet(\mathfrak{u}, \mathcal{M}_{\lambda, F})$  is equal the complex

$$\begin{aligned} \mathbb{H}^\bullet(\mathfrak{u}, \mathcal{M}_{\lambda, F}) = \{0 \rightarrow Fe_{n_1}^{(1)} \otimes Fe_{n_2}^{(2)} \xrightarrow{d} FE_+^{1, \vee} \otimes e_{-n_1}^{(1)} \otimes e_{n_2}^{(2)} \oplus FE_+^{1, \vee} \otimes e_{n_1}^{(1)} \otimes E_+^{2, \vee} \otimes e_{-n_2}^{(2)} \\ \xrightarrow{d} FE_+^{1, \vee} \otimes e_{-n_1}^{(1)} \otimes E_+^{2, \vee} \otimes e_{-n_2}^{(2)} \rightarrow 0\} \end{aligned} \quad (4.73)$$

where all the differentials are zero. The torus  $T$  acts by the weights

$$\lambda \text{ in degree 0, } s_1 \cdot \lambda, s_2 \cdot \lambda \text{ in degree 1, } w_0 \cdot \lambda \text{ in degree 2} \quad (4.74)$$

and we have a decomposition into one dimensional weight spaces

$$H^\bullet(\mathfrak{u}, \mathcal{M}_{\lambda, F}) = \bigoplus_{w \in W(\mathbb{C})} H^\bullet(\mathfrak{u}, \mathcal{M}_{\lambda, F})(w \cdot \lambda)$$

We go back to (9.116) and get a homomorphism of complexes

$$\text{Hom}_{\mathfrak{c}}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \mathbb{C}\chi \otimes \mathcal{M}_{\lambda, \mathbb{C}}) \rightarrow \text{Hom}_{\mathfrak{c}}(\Lambda^\bullet(\mathfrak{t}/\mathfrak{k}), \mathbb{C}\chi \otimes \mathbb{H}^\bullet(\mathfrak{u}, \mathcal{M}_{\lambda, \mathbb{C}})) \quad (4.75)$$

which induces an isomorphism in cohomology so that finally

$$H^\bullet(\mathfrak{g}, K_\infty, \mathfrak{I}_B^G \chi \otimes \mathcal{M}_{\lambda, \mathbb{C}}) \xrightarrow{\sim} H^\bullet(\text{Hom}_{\mathfrak{c}}(\Lambda^\bullet(\mathfrak{t}/\mathfrak{k}), \mathbb{C}\chi \otimes H^\bullet(\mathfrak{u}, \mathcal{M}_{\lambda, \mathbb{C}})) \quad (4.76)$$

and combining this with the results above we get cohlam

**Theorem 4.1.1.** *If we can find a  $w \in W(\mathbb{C})$  such that  $\chi^{-1} = w \cdot \lambda_{\mathbb{R}}$  then*

$$H^\bullet(\mathfrak{g}, K_\infty, \mathfrak{I}_B^G \chi \otimes \mathcal{M}_{\lambda, \mathbb{C}}) \xrightarrow{\sim} H^{l(w)}(\mathfrak{u}, \mathcal{M}_{\lambda, \mathbb{C}})(w \cdot \lambda) \otimes \Lambda^\bullet(\mathfrak{t}/\mathfrak{k})^\vee$$

*If there is no such  $w$  then the cohomology is zero.*

*Proof.* Our torus  $T(\mathbb{R}) = \mathfrak{c} \times \left\{ \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix}; t \in \mathbb{R}_{>0}^\times \right\} = \mathfrak{c} \times A$ . Hence we see that  $\dim \mathfrak{t}/\tilde{\mathfrak{k}} = 1$ , and the element  $H_0 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ . Of course we must have that  $\chi^{-1} \cdot \lambda_{\mathbb{R}}|_{\mathfrak{c}}$  is the trivial character. The second factor  $A$  does acts on  $\mathbb{C}\chi$  by the character  $\chi(t) = t^z$  and on  $H^{l(w)}(\mathfrak{u}, \mathcal{M}_{\lambda, \mathbb{C}})(w \cdot \lambda)$  by  $t \mapsto t^{m(w)}$ . Differentiating we get for the complex

$$0 \rightarrow H^{l(w)}(\mathfrak{u}, \mathcal{M}_{\lambda, \mathbb{C}})(w \cdot \lambda) \rightarrow \mathbb{C} \otimes H_0^\vee \otimes H^{l(w)}(\mathfrak{u}, \mathcal{M}_{\lambda, \mathbb{C}})(w \cdot \lambda) \rightarrow 0 \quad (4.77)$$

where the differential is multiplication by  $m(w) + z$ . Hence we see that the cohomology is trivial unless  $m(w) + z = 0$ , but this means  $\chi^{-1} = w \cdot \lambda_{\mathbb{R}}$ .  $\square$

### The cohomology of the modules $\mathcal{M}_{\lambda, \mathbb{C}}$ , $\mathcal{D}_\lambda$ and the cohomology of unitary modules

Again we start from a dominant character  $\lambda$ . We take the tensor product of the exact sequence (4.44) by  $\mathcal{M}_{\lambda^\vee, \mathbb{C}}$  and we get a long exact sequence of  $(\mathfrak{g}, K_\infty)$  cohomology modules

$$\begin{aligned} 0 \rightarrow H^0(\mathfrak{g}, K_\infty, \mathcal{M}_{\lambda, \mathbb{C}} \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}}) &\rightarrow H^0(\mathfrak{g}, K_\infty, \mathcal{J}_B^G \lambda_{\mathbb{R}}^{w_0} \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}}) \rightarrow H^0(\mathfrak{g}, K_\infty, \mathcal{D}_\lambda \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}}) \\ &\rightarrow H^1(\mathfrak{g}, K_\infty, \mathcal{M}_{\lambda, \mathbb{C}} \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}}) \rightarrow H^1(\mathfrak{g}, K_\infty, \mathcal{J}_B^G \lambda_{\mathbb{R}}^{w_0} \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}}) \rightarrow H^1(\mathfrak{g}, K_\infty, \mathcal{D}_\lambda \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}}) \\ &\rightarrow H^2(\mathfrak{g}, K_\infty, \mathcal{M}_{\lambda, \mathbb{C}} \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}}) \rightarrow H^2(\mathfrak{g}, K_\infty, \mathcal{J}_B^G \lambda_{\mathbb{R}}^{w_0} \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}}) \rightarrow H^2(\mathfrak{g}, K_\infty, \mathcal{D}_\lambda \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}}) \\ &\rightarrow H^3(\mathfrak{g}, K_\infty, \mathcal{M}_{\lambda, \mathbb{C}} \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}}) \rightarrow 0 \end{aligned} \quad (4.78)$$

We have seen that the modules  $\mathcal{D}_\lambda \xrightarrow{\sim} \mathcal{J}_B^G s_i \cdot \lambda_{\mathbb{R}}$  and hence know all the cohomology in this exact sequence except the the  $H^\bullet(\mathfrak{g}, K_\infty, \mathcal{M}_{\lambda, \mathbb{C}} \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}})$ . But then a careful analysis of  $K_\infty$ -types shows

#### Proposition 4.1.6.

$$\begin{aligned} H^0(\mathfrak{g}, K_\infty, \mathcal{M}_{\lambda, \mathbb{C}} \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}}) &= H^3(\mathfrak{g}, K_\infty, \mathcal{M}_{\lambda, \mathbb{C}} \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}}) = \mathbb{C}, \\ H^1(\mathfrak{g}, K_\infty, \mathcal{M}_{\lambda, \mathbb{C}} \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}}) &= H^2(\mathfrak{g}, K_\infty, \mathcal{M}_{\lambda, \mathbb{C}} \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}}) = 0 \\ H^3(\mathfrak{g}, K_\infty, \mathcal{D}_\lambda \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}}) &= H^2(\mathfrak{g}, K_\infty, \mathcal{J}_B^G \lambda_{\mathbb{R}}^{w_0} \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}}) = 0 \end{aligned}$$

If  $w \in W(\mathbb{C})$  is not  $= e, w_0$  (i.e. it is one of the elements of length one) then  $\mathcal{J}_B^G w \cdot \lambda_{\mathbb{R}} \xrightarrow{\sim} \mathcal{D}_\lambda$ . Looking at the  $K_\infty$  types occurring we see that the semi simple part of the lowest  $K_\infty$ -type is  $(n_1 + n_2 + 2)\gamma_c$ . The  $K_\infty$  type of  $\mathfrak{g}/\tilde{\mathfrak{k}}$  has highest weight  $2\gamma_c$  and  $\mathcal{M}_{\lambda, \mathbb{C}}$  has highest weight  $(n_1 + n_2)\gamma_c$ . This implies that our Lie -algebra complex becomes

$$0 \rightarrow \text{Hom}_{K_\infty}(\Lambda^1(\mathfrak{g}/\tilde{\mathfrak{k}}), \mathcal{J}_B^G w \cdot \lambda_{\mathbb{R}}) \xrightarrow{\partial} \text{Hom}_{K_\infty}(\Lambda^2(\mathfrak{g}/\tilde{\mathfrak{k}}), \mathcal{J}_B^G w \cdot \lambda_{\mathbb{R}}) \rightarrow 0 \quad (4.79)$$

in degree 1 and 2 the spaces are of dimension one and since the cohomology in these degrees is also one dimensional it follows that the boundary operator  $\partial = 0$ . EiShiso

### 4.1.5 The Eichler-Shimura Isomorphism

We want to apply these facts on representation theory to the study of cohomology groups  $H^\bullet(\Gamma \backslash X, \mathcal{M}_{\lambda, \mathbb{C}})$  where now  $\Gamma$  is a congruence subgroup of  $\mathrm{GL}_2(\mathbb{Z})$  or  $\mathrm{GL}_2(\mathcal{O})$ . (Discuss also quaternionic case- perhaps)

We start again from a dominant  $\lambda = n\gamma + d\det \in X^*(T \times \mathbb{C})$ . For every  $(\mathfrak{g}, K_\infty)$  invariant embedding  $\Psi_\lambda : \mathcal{I}_B^G w \cdot \lambda_{\mathbb{R}} \hookrightarrow \mathcal{C}_\infty(\Gamma \backslash G(\mathbb{R}))$  induces a homomorphism

$$\Psi_\lambda : \mathrm{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \mathcal{I}_B^G w \cdot \lambda_{\mathbb{R}} \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}}) \rightarrow \mathrm{Hom}_{K_\infty}((\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), (\mathcal{C}_\infty(\Gamma \backslash G(\mathbb{R})) \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}})) \quad (4.80)$$

We will show in section 8.1.3 Proposition 8.1.1 that the complex on the right is isomorphic to the de-Rham complex:

$$\mathrm{Hom}_{K_\infty}((\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), (\mathcal{C}_\infty(\Gamma \backslash G(\mathbb{R})) \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}}) \xrightarrow{\sim} \Omega^\bullet(\Gamma \backslash X, \mathcal{M}_{\lambda^\vee, \mathbb{C}}) \quad (4.81)$$

This de-Rham complex computes the cohomology and hence we get an homomorphism  $\boxed{\text{gkdeR}}$

$$\Psi_\lambda^\bullet : H^\bullet(\mathfrak{g}, K_\infty, \mathcal{I}_B^G w \cdot \lambda_{\mathbb{R}} \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}}) \rightarrow H^\bullet(\Gamma \backslash X, \mathcal{M}_{\lambda^\vee, \mathbb{C}} \otimes \mathbb{C}) \quad (4.82)$$

We denote by  $\omega^{(0)}$  the restriction of the central character of  $\mathcal{I}_B^G w \cdot \lambda_{\mathbb{R}}$  to the subgroup  $Z_0$ . (See above Interlude) and we introduce the spaces

$$\begin{aligned} \mathcal{E}^{\mathrm{mg}}(\lambda, w, \Gamma) &= \mathrm{Hom}_{(\mathfrak{g}, K_\infty)}(\mathcal{I}_B^G w \cdot \lambda_{\mathbb{R}}, \mathcal{C}_\infty^{\mathrm{mg}}(\Gamma \backslash G(\mathbb{R}), \omega^{(0)})) \\ &\cup \\ \mathcal{E}^{(2)}(\lambda, w, \Gamma) &= \mathrm{Hom}_{(\mathfrak{g}, K_\infty)}(\mathcal{I}_B^G w \cdot \lambda_{\mathbb{R}}, \mathcal{C}_\infty^{(2)}(\Gamma \backslash G(\mathbb{R}), \omega^{(0)})) \end{aligned} \quad (4.83)$$

where the superscripts mg resp. (2) mean moderate growth resp. square integrable.(Reference). From this we get two maps in cohomology

$$\Phi^\sharp : \mathcal{E}^\sharp(\lambda, w, \Gamma) \otimes H^\bullet(\mathfrak{g}, K_\infty, \mathcal{I}_B^G w \cdot \lambda_{\mathbb{R}} \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}}) \rightarrow H^\bullet(\Gamma \backslash X, \mathcal{M}_{\lambda^\vee, \mathbb{C}} \otimes \mathbb{C}) \quad (4.84)$$

Of course the module  $\mathcal{E}^{(2)}(\lambda, w, \lambda) = 0$  unless  $\mathcal{I}_B^G w \cdot \lambda_{\mathbb{R}}$  has a non trivial quotient module which admits a positive definite quasi unitary  $(\mathfrak{g}, K_\infty)$  invariant metric. This means that  $\mathcal{E}^{(2)}(\lambda, w, \lambda) \neq 0$  implies that in case B) the coefficients satisfy  $\boxed{\text{ul}}$

$$n_1 = n_2, \text{ i.e. } \lambda = n(\gamma_1 + \gamma_2) + d_1 \det + d_2 \det, \quad (4.85)$$

we will say that  $\lambda$  is unitary if this condition is fulfilled. Then the results in section (4.1.4) yield that these irreducible quasi unitary quotient modules are  $\mathcal{D}_\lambda^\pm$  in case A) and  $\mathcal{D}_\lambda$  in case B) .

If  $n = 0$  then  $\lambda$  extends to a character  $\tilde{\lambda} : G \rightarrow \mathbb{G}_m$  and  $\mathcal{M}_{\lambda, \mathbb{C}}$  is one dimensional we write  $\mathcal{M}_{\lambda, \mathbb{C}} = \mathbb{C}[\tilde{\lambda}]$ .

In the first two cases we know that

$$\mathcal{E}^{(2)}(\lambda, w, \Gamma) = \mathrm{Hom}_{(\mathfrak{g}, K_\infty)}(\mathcal{D}_\lambda, \mathcal{C}_\infty^{(2)}(\Gamma \backslash G(\mathbb{R}), \omega^{(0)}))$$

We have the fundamental  $\boxed{\text{ESI}}$

**Theorem 4.1.2.** (*Eichler-Shimura Isomorphism*) Assume  $\lambda$  unitary, then in degree 1 in case A, (resp. degree 1,2 in case B) the map

$$\Phi^{(2)} : \mathcal{E}^{(2)}(\lambda, w, \Gamma) \otimes H^\bullet(\mathfrak{g}, K_\infty, \mathcal{D}_\lambda \otimes \mathcal{M}_{\lambda^\vee, \mathbb{C}}) \rightarrow H_!^\bullet(\Gamma \backslash X, \mathcal{M}_{\lambda^\vee, \mathbb{C}} \otimes \mathbb{C}) \quad (4.86)$$

is an isomorphism.

If we are in the third case, i.e.  $n = 0$ , and if  $\lambda^2|_{\Gamma \cap Z} = 1$  then  $\text{Hom}_{(\mathfrak{g}, K_\infty)}(\mathbb{C}[\tilde{\lambda}], \mathcal{C}_\infty(G(\mathbb{R})))$  is one dimensional and generated by  $\Phi_\lambda : 1 \mapsto \tilde{\lambda}$ . The map

$$\mathbb{C}\Phi_\lambda \otimes H^\bullet(\mathfrak{g}, K_\infty, \mathbb{C}[\tilde{\lambda}] \otimes \mathbb{C}[\tilde{\lambda}^\vee]) \rightarrow H^\bullet(\Gamma \backslash X, \mathcal{M}_{\lambda^\vee, \mathbb{C}} \otimes \mathbb{C}) \quad (4.87)$$

is an isomorphism in degree zero and zero in all other degrees.

We want to relate this to the classical formulation in case A). The group  $\text{Sl}_2(\mathbb{R})$  acts transitively on the upper half plane  $\mathbb{H} = \text{Sl}_2(\mathbb{R})/\text{SO}(2)$ . For  $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  and  $z \in \mathbb{H}$  we put  $j(g, z) = cz + d$ . To any

$$\Phi \in \text{Hom}_{(\mathfrak{g}, K_\infty)}(\mathcal{D}_\lambda^+, \mathcal{C}_\infty^{(2)}(\Gamma \backslash G(\mathbb{R}), \omega^{(0)}))$$

we attach a function  $f_{n+2}^\Phi : \mathbb{H} \rightarrow \mathbb{C}$  : We write  $z = gi$  with  $g \in \text{Sl}_2(\mathbb{R})$  and put holWh

$$f_{n+2}^\Phi(z) = \Phi(\psi_{n+2})(g)j(g, i)^{n+2} \quad (4.88)$$

An easy calculation shows that  $f_{n+2}^\Phi$  is well defined and holomorphic (slzweineu.pdf)p.25-26) and for  $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{Sl}_2(\mathbb{Z})$  it satisfies

$$f_{n+2}^\Phi(\gamma z) = (cz + d)^{n+2} f_{n+2}^\Phi(z) \quad (4.89)$$

The condition that  $\Phi(\psi_{n+2})(g)$  is square integrable implies that  $f_{n+2}$  is a holomorphic cusp form of weight  $n + 2 = k$ . It is a special case of the theorem of Gelfand-Graev that this provides an isomorphism GelfGraev

$$\text{Hom}_{(\mathfrak{g}, K_\infty)}(\mathcal{D}_\lambda^+, \mathcal{C}_\infty^{(2)}(\Gamma \backslash G(\mathbb{R}))) \xrightarrow{\sim} S_k(\Gamma) \quad (4.90)$$

where of course  $S_k(\Gamma)$  is the space of holomorphic cusp forms for  $\Gamma$ .

We can do the same thing with  $\mathcal{D}_\lambda^-$  then we land in the spaces of anti holomorphic cusp forms, these two spaces are isomorphic under conjugation. Combining this with our results above gives the classical formulation of the Eichler-Shimura theorem:

We have a canonical isomorphism

$$S_k(\Gamma) \oplus \overline{S_k(\Gamma)} \xrightarrow{\sim} H_!^1(\Gamma \backslash X, \mathcal{M}_{\lambda^\vee, \mathbb{C}} \otimes \mathbb{C}) \quad (4.91)$$

### The-Manin-Drinfeld principle

Whittloc

### 4.1.6 Local Whittaker models

We recall some fundamental results from representation theory of groups  $\mathrm{Gl}_2(\mathbb{Q}_p)$ . Let  $F/\mathbb{Q}$  be a finite extension  $\mathbb{Q}$ . An admissible representation of  $\mathrm{Gl}_2(\mathbb{Q}_p)$  is an action of  $\mathrm{Gl}_2(\mathbb{Q}_p)$  on a  $F$ -vector space  $V$  which fulfills the following two additional requirements

a) For any open subgroup  $K_p \subset \mathrm{Gl}_2(\mathbb{Z}_p)$  the space of fixed vectors  $V^{K_p}$  is finite dimensional.

b) For any  $v \in V$  we find an open subgroup  $K_p \subset \mathrm{Gl}_2(\mathbb{Z}_p)$  such that  $v \in V^{K_p}$ .

We say that  $V$  is a  $\mathrm{Gl}_2(\mathbb{Q}_p)$  module, we denote the action of  $\mathrm{Gl}_2(\mathbb{Q}_p)$  on  $V$  by  $(g, v) \mapsto gv$ . In addition we want to assume that our module has a central character, this means that the center  $Z(\mathbb{Q}_p) = \mathbb{Q}_p^\times$  acts by a character  $\omega_V : Z(\mathbb{Q}_p) \rightarrow F^\times$ . Such a module is called irreducible if it does not contain a non trivial invariant submodule.

Again we dispose of a Hecke algebra, given  $K_p$  we consider the space of functions

$$\mathcal{H}_{K_p} = \{f : \mathrm{Gl}_2(\mathbb{Q}_p) \rightarrow F \mid f(zg) = \omega_V^{-1}(z)f(g) ; f \text{ has compact support mod } Z(\mathbb{Q}_p)\}$$

this gives as an algebra by convolution and this algebra acts on  $V^{K_p}$  by

$$f * v = \int_{\mathrm{Gl}_2(\mathbb{Q}_p)/Z(\mathbb{Q}_p)} f(x) x v dx$$

(See also section 6.3.3.) We normalize the measure  $dx$  such that it gives volume one to  $K_p$ .

We recall - and explain the meaning of - the fundamental fact that each isomorphism class of admissible irreducible modules has a unique Whittaker model. We assume that  $F \subset \mathbb{C}$ , then we define the (additive) character PSI

$$\psi_p : \mathbb{Q}_p \rightarrow \mathbb{C}^\times ; \psi_p : a/p^m \mapsto e^{\frac{2\pi i a}{p^m}} \tag{4.92}$$

it is clear that the kernel of  $\psi_p$  is  $\mathbb{Z}_p$ . Since we have  $U(\mathbb{Q}_p) = \mathbb{Q}_p$  we can view  $\psi_p$  as a character  $\psi_p : U(\mathbb{Q}_p) \rightarrow \mathbb{C}^\times$ . We introduce the space

$$\mathcal{C}_{\psi_p}(\mathrm{Gl}_2(\mathbb{Q}_p)) = \{f : \mathrm{Gl}_2(\mathbb{Q}_p) \rightarrow \mathbb{C} \mid f(ug) = \psi_p(u)f(g)\}$$

where in addition we require that our  $f$  is invariant under a suitable open subgroup  $K_f \subset \mathrm{Gl}_2(\mathbb{Z}_p)$ . The group  $\mathrm{Gl}_2(\mathbb{Q}_p)$  acts on this space by right translation (the action is not admissible but satisfies the above condition b) .

Now we can state the theorem about existence and uniqueness of the Whittaker model

Whittp

**Theorem 4.1.3.** *For any absolutely irreducible admissible  $\mathrm{Gl}_2(\mathbb{Q}_p)$  -module  $V$  we find a non trivial ( of course invariant under  $\mathrm{Gl}_2(\mathbb{Q}_p)$ ) homomorphism*

$$\Psi : V \rightarrow \mathcal{C}_{\psi_p}(\mathrm{Gl}_2(\mathbb{Q}_p)), \tag{4.93}$$

*it is unique up to multiplication by a non zero scalar.*

*Proof.* We refer to the literature. □

**Spherical representations, their Whittaker model and the Euler factor**

An absolutely irreducible  $\mathrm{Gl}_2(\mathbb{Q}_p)$  module is called spherical or unramified if for  $K_p = \mathrm{Gl}_2(\mathbb{Z}_p)$  we have  $V^{K_p} \neq \{0\}$ . In this case it is known that (*Reference*)

$$\dim_F(V^{\mathrm{Gl}_2(\mathbb{Z}_p)}) = 1; V^{\mathrm{Gl}_2(\mathbb{Z}_p)} = F\phi_0. \quad (4.94)$$

The Hecke algebra  $\mathcal{H}_{K_p}$  is commutative and generated by the two double cosets

$$T_p = \mathrm{Gl}_2(\mathbb{Z}_p) \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} \mathrm{Gl}_2(\mathbb{Z}_p) \text{ and } C_p = \mathrm{Gl}_2(\mathbb{Z}_p) \begin{pmatrix} p & 0 \\ 0 & p \end{pmatrix}. \quad (4.95)$$

The space  $V^{\mathrm{Gl}_2(\mathbb{Z}_p)}$  is an absolutely irreducible module for  $\mathcal{H}_{K_p}$  hence it is of rank one, let  $\psi_0$  be a generator. Our two operators act by scalars on  $V^{K_p}$ , we write

$$T_p(\psi_0) = \pi_V(T_p)\psi_0 \text{ and } C_p(\psi_0) = \pi_V(C_p)\psi_0 \quad (4.96)$$

The module  $V$  is completely determined by these two eigenvalues, of course  $\pi_V(C_p) = \omega_V(C_p)$ .

We can formulate this a little bit differently. Let  $\pi_p$  an isomorphism type of our  $\mathrm{Gl}_2(\mathbb{Q}_p)$  module  $V$ . Then our theorem above asserts that there is a unique  $\mathrm{Gl}_2(\mathbb{Q}_p)$ -module

$$\mathcal{W}(\pi_p) \subset \mathcal{C}_\psi(\mathrm{Gl}_2(\mathbb{R})) \quad (4.97)$$

with isomorphism-type equal to  $\pi_p \times_F \mathbb{C}$ . We call this module the Whittaker realization of  $\pi_p$ . If our isomorphism type is unramified then the resulting homomorphism of  $\mathcal{H}_p$  to  $F$  is also denoted by  $\pi_p$ .

We have the spherical vector  $h_{\pi_p}^{(0)} \in \mathcal{W}(\pi_p)^{\mathrm{Gl}_2(\mathbb{Q}_p)}$  which is unique up to a scalar. Since  $\mathrm{Gl}_2(\mathbb{Q}_p) = U(\mathbb{Q}_p)T(\mathbb{Q}_p)\mathrm{Gl}_2(\mathbb{Z}_p)$  this spherical vector is determined by its restriction to  $T(\mathbb{Q}_p)$ . We have a formula for this restriction. First of all we observe that

$$h_{\pi_p}^{(0)} \left( \begin{pmatrix} p^n & 0 \\ 0 & p^m \end{pmatrix} \right) = \pi_p(C_p^m) h_{\pi_p}^{(0)} \left( \begin{pmatrix} p^{n-m} & 0 \\ 0 & 1 \end{pmatrix} \right). \quad (4.98)$$

We claim that  $h_{\pi_p}^{(0)} \left( \begin{pmatrix} p^n & 0 \\ 0 & 1 \end{pmatrix} \right) = 0$  if  $n < 0$ . To see this we look at the equalities

$$h_{\pi_p}^{(0)} \left( \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix} \begin{pmatrix} p^n & 0 \\ 0 & 1 \end{pmatrix} \right) = \psi_p(u) h_{\pi_p}^{(0)} \left( \begin{pmatrix} p^n & 0 \\ 0 & 1 \end{pmatrix} \right) = h_{\pi_p}^{(0)} \left( \begin{pmatrix} p^n & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & p^{-n}u \\ 0 & 1 \end{pmatrix} \right)$$

and we can find an element  $u \in \mathbb{Q}_p$  such that  $p^{-n}u \in \mathbb{Z}_p$  and  $\psi_p(u) \neq 1$ , this implies the claim. Now we exploit the eigenvalue equation  $T_p(h_{\pi_p}^{(0)}) = \pi_p(T_p)h_{\pi_p}^{(0)}$ ,

we write the double coset  $K_p \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} K_p$  as union of right  $K_p$  cosets

$$K_p \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} K_p = \bigcup_{x \in \mathbb{Z}/p\mathbb{Z}} \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} K_p \cup \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} K_p.$$

Clearly

$$\begin{aligned} h_{\pi_p}^{(0)}\left(\begin{pmatrix} p^n & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}\right) &= h_{\pi_p}^{(0)}\left(\begin{pmatrix} p^{n+1} & 0 \\ 0 & 1 \end{pmatrix}\right) \\ h_{\pi_p}^{(0)}\left(\begin{pmatrix} p^n & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}\right) &= h_{\pi_p}^{(0)}\left(\begin{pmatrix} p^{n-1} & 0 \\ 0 & 1 \end{pmatrix}\right) \end{aligned}$$

and this implies the recursion formula recurs

$$\pi_p(T_p)h_{\pi_p}^{(0)}\left(\begin{pmatrix} p^n & 0 \\ 0 & 1 \end{pmatrix}\right) = \pi_p(C_p)h_{\pi_p}^{(0)}\left(\begin{pmatrix} p^{n-1} & 0 \\ 0 & 1 \end{pmatrix}\right) + \begin{cases} ph_{\pi_p}^{(0)}\left(\begin{pmatrix} p^{n+1} & 0 \\ 0 & 1 \end{pmatrix}\right) & \text{if } n \geq 0 \\ 0 & \text{if } n < 0 \end{cases} \quad (4.99)$$

We can normalize  $h_{\pi_p}^{(0)}\left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}\right) = 1$ , then the values for  $n > 0$  follow from the recursion.

There is a more elegant writing this recursion. For our unramified  $\pi_p$  we define the local Euler factor Euler

$$L(\pi_p, s) = \frac{1}{1 - \pi_p(T_p)p^{-s} + p\pi_p(C_p)p^{-2s}} \quad (4.100)$$

If we expand this into a power series in  $p^{-s}$  then Mellin

$$L(\pi_p, s) = \sum_{n=0}^{\infty} h_{\pi_p}^{(0)}\left(\begin{pmatrix} p^n & 0 \\ 0 & 1 \end{pmatrix}\right) p^n p^{-ns} \quad (4.101)$$

### Whittaker models for Harish-Chandra modules

We also have a theory of Whittaker models for the irreducible Harish-Chandra modules studied in section 4.1. The unipotent radical  $U(\mathbb{R}) = \mathbb{R}$  resp.  $U(\mathbb{R}) = \mathbb{C}$ . Again we fix characters  $\psi_{\infty} : U(\mathbb{R}) \rightarrow \mathbb{C}^{\times}$  we put

$$\psi_{\infty}(x) = \begin{cases} e^{-2\pi ix} & \text{in case A) } \\ e^{-2\pi i(x+\bar{x})} & \text{in case B) } \end{cases} \quad (4.102)$$

and as in the  $p$ -adic case we define

$$\mathcal{C}_{\psi_{\infty}}(G(\mathbb{R})) = \{f : G(\mathbb{R}) \rightarrow \mathbb{C} \mid f(ug) = \psi_{\infty}(u)f(g)\}$$

Then we have again Whittinf

**Theorem 4.1.4.** *For any infinite dimensional, absolutely irreducible admissible  $\mathrm{GL}_2(\mathbb{R})$ -module  $V$  we find a non trivial ( of course invariant under  $\mathrm{GL}_2(\mathbb{R})$ ) homomorphism*

$$\Psi : V \rightarrow \mathcal{C}_{\psi_{\infty}}(G(\mathbb{R})), \quad (4.103)$$

*This homomorphism is unique up to a scalar. The image of  $V$  under the homomorphism  $\Psi$  will be denoted by  $\tilde{V}$ .*

*Proof.* Again we refer to the literature. □

Hence we can say that for any isomorphism class  $\pi_\infty$  of irreducible infinite dimensional Harish-Chandra modules we have a unique Whittaker model  $\mathcal{W}(\pi_\infty) \subset \mathcal{C}_{\psi_\infty}(G(\mathbb{R}))$ . In the book of Godement we find explicit formulae for these Whittaker functions.

Actually it is easy to write down such maps  $\tilde{\Psi}_\pm$  resp.  $\tilde{\Psi}$  explicitly for our induced modules, we start from a dominant  $\lambda = n\gamma + \delta$  (resp.  $n(\gamma_1 + \gamma_2) + \delta$ ). We define

$$\tilde{\Psi} : \mathcal{I}_B^G \lambda_{\mathbb{R}} | \rho |_{\mathbb{R}}^2 \rightarrow \mathcal{C}_{\psi_\infty}(G(\mathbb{R}))$$

by the integral

$$\tilde{\Psi}(f)(g) = \int_{U(\mathbb{R})} f(wug) \psi_\infty(-u) du,$$

there is no problem with convergence. We apply this to the spherical function  $\psi_0 \in \mathcal{I}_B^G \lambda_{\mathbb{R}} | \rho |_{\mathbb{R}}^2$  (See ??) and a rather straightforward computation yields We choose a  $\lambda = n\gamma; n \geq 0$  and even and consider the induced module  $\mathcal{I}_B^G \lambda_{\mathbb{R}} \rho_{\mathbb{R}}^2 = \bigoplus_{\nu \equiv 0(2)} \mathbb{C} \phi_{\lambda, \nu}$ , we have the exact sequence

$$0 \rightarrow \mathcal{D}_\lambda^+ \oplus \mathcal{D}_\lambda^- \rightarrow \mathcal{I}_B^G \lambda_{\mathbb{R}} \rho_{\mathbb{R}}^2 \rightarrow \mathcal{M}_\lambda \rightarrow 0$$

We have the Whittaker map

$$\mathcal{F} : \mathcal{I}_B^G \lambda_{\mathbb{R}} \rho_{\mathbb{R}}^2 \rightarrow \mathcal{C}_\psi(G(\mathbb{R}))$$

which is defined by

$$\mathcal{F}(\phi_{\lambda, \nu}) \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) := \int_{-\infty}^{\infty} \phi_{\lambda, \nu} \left( w \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) e^{2\pi i x} dx$$

We write the Cartan decomposition

$$w \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & -1 \\ -t & -x \end{pmatrix} = \begin{pmatrix} \frac{t}{\sqrt{t^2+x^2}} & * \\ 0 & \sqrt{t^2+x^2} \end{pmatrix} \begin{pmatrix} \frac{-x}{\sqrt{t^2+x^2}} & \frac{t}{\sqrt{t^2+x^2}} \\ \frac{-t}{\sqrt{t^2+x^2}} & \frac{-x}{\sqrt{t^2+x^2}} \end{pmatrix}$$

and a straightforward computation gives us that we have to evaluate

$$t^{\frac{n}{2}+1} \int_{-\infty}^{\infty} \frac{e^{2\pi i x}}{(x+ti)^{n/2+\nu/2+1} (x-ti)^{n/2-\nu/2+1}} dx$$

This can be done by the Residue theorem, we integrate from  $-R$  to  $R$  and then from  $R$  back to  $-R$  along the circle in the upper half plane. Our function has only one pole in the upper half plane, namely for  $x = ti$  and therefore

$$\int_{-\infty}^{\infty} \phi_{\lambda, \nu} \left( w \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) e^{2\pi i x} dx = t^{\frac{n}{2}+1} \operatorname{Res}_{x=ti} \frac{e^{2\pi i x}}{(x+ti)^{n/2+\nu/2+1} (x-ti)^{n/2-\nu/2+1}}$$

If we put  $z := x - ti$  then our integral becomes

$$(2i)^{-n/2-\nu/2-1} t^{-\nu/2} e^{-2\pi t} \operatorname{Res}_{z=0} \frac{e^{2\pi i z}}{\left(1 + \frac{z}{2ti}\right)^{n/2+\nu/2+1} z^{n/2-\nu/2+1}} = P_{\lambda, \nu}(t) e^{-2\pi t},$$

where  $P_{\lambda,\nu}(t)$  is a Laurent polynomial in  $\mathbb{C}[t, t^{-1}]$ . This polynomial is zero if  $\nu \geq n + 2$  and this implies that  $\mathcal{F}$  maps  $\mathcal{D}_\lambda^+$  to zero.

Therefore our map  $\mathcal{F}$  induces an injection

$$\mathfrak{I}_B^G \lambda_{\mathbb{R}} \rho_{\mathbb{R}}^2 / \mathcal{D}_\lambda^+ \hookrightarrow \mathcal{C}_\psi(G(\mathbb{R}))$$

this is of course an intertwining operator. The module  $\mathcal{D}_\lambda^- \subset \mathfrak{I}_B^G \lambda_{\mathbb{R}} \rho_{\mathbb{R}}^2 / \mathcal{D}_\lambda^+$  it has  $\phi_{\lambda, -n-2}$  as a lowest weight vector. If we apply our above formula to  $\mathcal{F}(\phi_{\lambda, -n-2})$  then we see that the nasty factor  $(1 + \frac{z}{2ti})^{n/2+\nu/2+1}$  is equal to one in this case and hence we have up to a non zero constant

$$\mathcal{F}(\phi_{\lambda, -n-2})\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right) = c_\lambda t^{\frac{n}{2}+1} e^{-2\pi t}.$$

In the case A) the we the two discrete irreducible series representations  $\mathcal{D}_\lambda^+, \mathcal{D}_\lambda^-$  attached to a dominant weight  $\lambda$ . We have their Whittaker model

$$\tilde{\Psi}_\pm : \mathcal{D}_\lambda^\pm \hookrightarrow \mathcal{C}_{\psi_\infty}(\mathrm{Gl}_2(\mathbb{R})). \tag{4.104}$$

The group  $(\mathrm{Gl}_2(\mathbb{R}))$  has the two connected components  $\mathrm{Gl}_2(\mathbb{R})^+, \mathrm{Gl}_2(\mathbb{R})^-, (\det > 0, \det < 0)$  and we have

$$\tilde{\Psi}_+(\mathcal{D}_\lambda^+) = \tilde{\mathcal{D}}_\lambda^+ \text{ is supported on } \mathrm{Gl}_2(\mathbb{R})^+, \tilde{\mathcal{D}}_\lambda^- \text{ is supported on } \mathrm{Gl}_2(\mathbb{R})^- \tag{4.105}$$

Under the isomorphism  $\tilde{\Psi}_\pm$  the elements  $\psi_{\pm(n+2)}$  (See (4.24) ) are mapped to functions  $\tilde{\psi}_{\pm(n+2)}$ . We can normalize  $\tilde{\Psi}_\pm$  such that tpsin

$$\psi^\dagger_{n+2}\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right) = \begin{cases} t^{\frac{n}{2}+1} e^{-2\pi t} & \text{if } t > 0 \\ 0 & \text{else} \end{cases} \tag{4.106}$$

and  $\psi^\dagger_{-n-2}$  is given by the corresponding formula.

Whitt

**Global Whittaker models, Fourier expansions and multiplicity one**

We also have global Whittaker models. To define them we recall some results from Tate's thesis ([83]). We introduce the ring of adeles  $\mathbb{A} = \mathbb{A}_{\mathbb{Q}}$ , we write it as a product  $\mathbb{A} = \mathbb{Q}_\infty \times \mathbb{A}_f = \mathbb{R} \times \mathbb{A}_f$ . The ring of finite adeles contains the compact subring  $\hat{\mathbb{Z}} = \prod_p \mathbb{Z}_p$  of integral adeles.

We define a global character  $\psi : U(\mathbb{A})/U(\mathbb{Q}) = \mathbb{A}/\mathbb{Q} \rightarrow \mathbb{C}^\times$  as the product psiq

$$\psi(x_\infty, \dots, x_p, \dots) = \psi_\infty(x_\infty) \prod_p \psi_p(x_p) \tag{4.107}$$

where the local components  $\psi_p$  are as above, we have to check that  $\psi$  is trivial on  $U(\mathbb{Q})$ . (See [83], "note the minus sign") For any  $a \in \mathbb{Q}$  we define  $\psi^{[a]}(x) = \psi(ax)$ , so  $\psi = \psi^{[1]}$ . In ([83]) it is shown that the map

$$\mathbb{Q} \rightarrow \mathrm{Hom}(\mathbb{A}/\mathbb{Q}, \mathbb{C}^\times); a \mapsto \psi^{[a]} \tag{4.108}$$

is an isomorphism between  $\mathbb{Q}$  and the character group of  $\mathbb{A}/\mathbb{Q}$ . Hence we know that for any reasonable function  $h : \mathbb{A}/\mathbb{Q} \rightarrow \mathbb{C}$  we have a Fourier expansion

Fouex

$$h(\underline{u}) = \sum_{a \in \mathbb{Q}} \hat{h}(a) \psi(a\underline{u}) \tag{4.109}$$

where  $\hat{h}(a) = \int_{\mathbb{A}/\mathbb{Q}} h(\underline{u}) \psi(-a\underline{u}) d\underline{u}$ , and where  $\text{vol}_{d\underline{u}}(\mathbb{A}/\mathbb{Q}) = 1$ . Then we put

$$\mathcal{C}_\psi(\text{Gl}_2(\mathbb{R}) \times \text{Gl}_2(\mathbb{A}_f)/K_f) = \{f : \text{Gl}_2(\mathbb{R}) \times \text{Gl}_2(\mathbb{A}_f)/K_f \rightarrow \mathbb{C} \mid f(\underline{u}\underline{g}) = \psi(\underline{u})f(\underline{g})\}$$

this is a module for  $\text{Gl}_2(\mathbb{R}) \times \otimes' \mathcal{H}_p$

Let us start from the Harish-Chandra module  $\pi_\infty = \mathcal{D}_\lambda^+$  and a homomorphism  $\pi_f = \otimes' \pi_p : \otimes' \mathcal{H}_p \rightarrow F$  from the unramified Hecke algebra to  $F$ . We still assume for simplicity that  $K_f = \text{Gl}_2(\hat{\mathbb{Z}})$ .

The results on Whittaker-models imply that we have a unique Whittaker-model

$$\mathcal{W}(\pi) = \mathcal{W}(\pi_\infty) \otimes \mathbb{C}h_{\pi_f}^{(0)} \subset \mathcal{C}_\psi(\text{Gl}_2(\mathbb{R}) \times \text{Gl}_2(\mathbb{A}_f)/K_f) \tag{4.110}$$

for our isomorphism class  $\pi = \pi_\infty \times \pi_f$ . Here of course  $h_{\pi_f}^{(0)} = \otimes h_{\pi_p}^{(0)}$ .

We return to Theorem 4.1.2. On the space  $\mathcal{C}_\infty^{(2)}(\Gamma \backslash G(\mathbb{R}), \omega^{(0)})$  we have the action of the unramified Hecke algebra. To see this action we start from the observation that the map  $\text{Gl}_2(\mathbb{Q}) \rightarrow \text{Gl}_2(\mathbb{A}_f)/K_f$  (Chap. III , 1.5) is surjective and hence

$$\text{Gl}_2(\mathbb{Z}) \backslash \text{Gl}_2(\mathbb{R}) \xrightarrow{\sim} \text{Gl}_2(\mathbb{Q}) \backslash \text{Gl}_2(\mathbb{R}) \times \text{Gl}_2(\mathbb{A}_f)/K_f \tag{4.111}$$

and hence

$$\mathcal{C}_\infty^{(2)}(\text{Gl}_2(\mathbb{Z}) \backslash \text{Gl}_2(\mathbb{R})) = \mathcal{C}_\infty^{(2)}(\text{Gl}_2(\mathbb{Q}) \backslash \text{Gl}_2(\mathbb{R}) \times \text{Gl}_2(\mathbb{A}_f)/K_f) \tag{4.112}$$

and the space on the right is a  $\text{Gl}_2(\mathbb{R}) \times \otimes' \mathcal{H}_p$  module. Now we consider the  $\pi = \pi_\infty \times \pi_f$  isotypical submodule  $\mathcal{C}_\infty^{(2)}(\text{Gl}_2(\mathbb{Q}) \backslash \text{Gl}_2(\mathbb{R}) \times \text{Gl}_2(\mathbb{A}_f)/K_f)(\pi) \subset \mathcal{C}_\infty^{(2)}(\text{Gl}_2(\mathbb{Q}) \backslash \text{Gl}_2(\mathbb{R}) \times \text{Gl}_2(\mathbb{A}_f)/K_f)$ .

We have the famous Theorem which in the case  $\Gamma = \text{Sl}_2(\mathbb{Z})$  is due to Hecke

multone

**Theorem 4.1.5.** *If  $\mathcal{C}_\infty^{(2)}(\text{Gl}_2(\mathbb{Q}) \backslash \text{Gl}_2(\mathbb{R}) \times \text{Gl}_2(\mathbb{A}_f)/K_f)(\pi) \neq 0$  then have a canonical isomorphism*

$$\mathcal{F}_1 : \mathcal{W}(\pi) \xrightarrow{\sim} \mathcal{C}_\infty^{(2)}(\text{Gl}_2(\mathbb{Q}) \backslash \text{Gl}_2(\mathbb{R}) \times \text{Gl}_2(\mathbb{A}_f)/K_f)(\pi) \tag{4.113}$$

especially we know that  $\pi$  occurs with multiplicity one.

*Proof.* We give the inverse of  $\mathcal{F}_1$ . Given a function

$$h \in \mathcal{C}_\infty^{(2)}(\text{Gl}_2(\mathbb{Q}) \backslash \text{Gl}_2(\mathbb{R}) \times \text{Gl}_2(\mathbb{A}_f)/K_f)(\pi)$$

we define

$$h^\dagger((g_\infty, \underline{g}_f)) = \int_{U(\mathbb{Q}) \backslash U(\mathbb{A})} h(\underline{u}\underline{g}) \overline{\psi(\underline{u})} d\underline{u} \tag{4.114}$$

it is clear that  $h^\dagger(g_\infty, \underline{g}_f) \in \mathcal{W}(\pi)$ . It follows from the theory of automorphic forms that  $h$  is actually in the space of cusp forms, this means that the constant Fourier coefficient  $\int_{U(\mathbb{Q}) \backslash U(\mathbb{A})} h(\underline{u}g) d\underline{u} = 0$  and hence our Fourier expansion yields ((4.109), evaluated at  $u = 0$ )

$$h(\underline{g}) = \sum_{a \in \mathbb{Q}^\times} \int_{U(\mathbb{A})/U(\mathbb{Q})} h(\underline{u}g) \psi^{[a]}(\underline{u}) d\underline{u} \tag{4.115}$$

The measure  $d\underline{u}$  is invariant under multiplication by  $a \in \mathbb{Q}^\times$  and hence a individual term in the summation is

$$\int_{U(\mathbb{A})/U(\mathbb{Q})} h\left(\begin{pmatrix} 1 & \underline{u} \\ 0 & 1 \end{pmatrix} \underline{g}\right) \psi\left(\begin{pmatrix} 1 & a\underline{u} \\ 0 & 1 \end{pmatrix}\right) d\underline{u} = \int_{U(\mathbb{A})/U(\mathbb{Q})} h\left(\begin{pmatrix} 1 & a^{-1}\underline{u} \\ 0 & 1 \end{pmatrix} \underline{g}\right) \psi\left(\begin{pmatrix} 1 & \underline{u} \\ 0 & 1 \end{pmatrix}\right) d\underline{u} \tag{4.116}$$

Now

$$\begin{pmatrix} 1 & a^{-1}\underline{u} \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} a^{-1} & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & \underline{u} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix}$$

Since  $h$  is invariant under the action of  $G(\mathbb{Q})$  from the left we find

$$\int_{U(\mathbb{A})/U(\mathbb{Q})} h(\underline{u}g) \psi^{[a]}(\underline{u}) d\underline{u} = h^\dagger\left(\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} g_\infty\right) h_f^\dagger(\underline{a}_f) \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} (g_\infty, \underline{g}_f) \tag{4.117}$$

We evaluate at  $\underline{g} = (g_\infty, e)$  then

$$h^\dagger\left(\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} (g_\infty, e)\right) = h^\dagger\left(\begin{pmatrix} a_\infty & 0 \\ 0 & 1 \end{pmatrix} g_\infty, \begin{pmatrix} a_f & 0 \\ 0 & 1 \end{pmatrix}\right) \tag{4.118}$$

For a fixed  $g_\infty$  the function  $\underline{g}_f \mapsto h^\dagger(g_\infty, \underline{g}_f)$  is up to a factor equal to  $h_{\pi_f}^{(0)} = \bigotimes'_p h_{\pi_p}^{(0)}$  and hence we find

$$h^\dagger\left(\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} (g_\infty, e)\right) = h^\dagger\left(\begin{pmatrix} a_\infty & 0 \\ 0 & 1 \end{pmatrix} g_\infty, e\right) h_{\pi_f}^{(0)}\left(\begin{pmatrix} a_f & 0 \\ 0 & 1 \end{pmatrix}\right) \tag{4.119}$$

The recursion formulae ( 4.99),(4.134) imply that  $h_{\pi_f}^{(0)}\left(\begin{pmatrix} a_f & 0 \\ 0 & 1 \end{pmatrix}\right) = 0$  unless  $a \in \mathbb{Z}$ .

We restrict our functions to  $\text{Gl}_2^+(\mathbb{R})$ , i.e. we take  $g_\infty \in \text{Gl}_2(\mathbb{R})^+$  and we remember that our representation  $\pi_\infty$  is  $\mathcal{D}_\lambda^+$ . Then we know that for  $h_\infty \in \mathcal{D}_\lambda^+$  the value  $h^\dagger\left(\begin{pmatrix} a_\infty & 0 \\ 0 & 1 \end{pmatrix} g_\infty, e\right) = 0$  if  $a_\infty < 0$  and hence

$$h^\dagger\left(\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} (g_\infty, e)\right) = h^\dagger\left(\begin{pmatrix} a_\infty & 0 \\ 0 & 1 \end{pmatrix} g_\infty, \begin{pmatrix} a_f & 0 \\ 0 & 1 \end{pmatrix}\right) = 0 \text{ unless } a > 0, a \in \mathbb{Z},$$

and our Fourier expansion (4.109) becomes Fexpl

$$h(\underline{g}) = \sum_{a=1}^{\infty} h^\dagger\left(\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} (g_\infty, e)\right) h_{\pi_f}^{(0)}\left(\begin{pmatrix} a_f & 0 \\ 0 & 1 \end{pmatrix}\right) \tag{4.120}$$

□

We notice that there is never any problem with convergence. The Whittaker functions  $h_\infty^\dagger$  always decay very rapidly at infinity. We write  $g_\infty = \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix} \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} k$  with  $k \in K_\infty$ , then it is easy to see

$$|h_\infty^\dagger \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} g_\infty \right)| < P(t)e^{-2\pi t}$$

where  $P(t)$  is a polynomial in  $t$ . This implies that the series is really very rapidly converging (See remark below).

Now we choose for the component at infinity the function  $h_\infty^\dagger = \tilde{\psi}_{n+2}$  and we compute the corresponding holomorphic cusp form  $h^\Phi$  under the Eichler-Shimura isomorphism. We have the formula (4.88)

$$h^\Phi(z) = h^\Phi(x+iy) = h \left( \begin{pmatrix} y^{\frac{1}{2}} & \frac{x}{y^{\frac{1}{2}}} \\ 0 & y^{-\frac{1}{2}} \end{pmatrix} \right) j \left( \begin{pmatrix} y^{\frac{1}{2}} & \frac{x}{y^{\frac{1}{2}}} \\ 0 & y^{-\frac{1}{2}} \end{pmatrix}, i \right)^{n+2} = h \left( \begin{pmatrix} y^{\frac{1}{2}} & \frac{x}{y^{\frac{1}{2}}} \\ 0 & y^{-\frac{1}{2}} \end{pmatrix} \right) y^{-\frac{n}{2}-1}$$

and hence our Fourier expansion (4.120) becomes FouH

$$h^\Phi(z) = y^{-\frac{n}{2}-1} \sum_{a=1}^{\infty} \tilde{\phi}_{n+2} \left( \begin{pmatrix} ay & ax \\ 0 & 1 \end{pmatrix} \right) h_{\pi_f}^{(0)} \left( \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \right) \quad (4.121)$$

We have the formula (4.106) for  $\tilde{\phi}_{n+2}$  and then this becomes

$$h^\Phi(z) = \sum_{a=1}^{\infty} a^{\frac{n}{2}+1} h_{\pi_f}^{(0)} \left( \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \right) e^{2\pi i z a} \quad (4.122)$$

This is now the classical Fourier expansion of a holomorphic cusp eigenform of weight  $k = n + 2$ , ([38]). The numbers  $c(\pi_f, a) = a^{\frac{n}{2}+1} h_{\pi_f}^{(0)} \left( \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \right)$  are the Fourier coefficients and they also the eigenvalues of the operator  $T_a$ -defined in by Hecke in [38]- on  $h^\Phi$ . If we apply the the Eichler-Shimura isomorphism and interpret  $h^\Phi$  as a cohomology class then it is an eigenclass in  $H_1^1(\Gamma \backslash H, \mathcal{M}_n \otimes \mathbb{C})$  and for any prime  $p$  the number  $c(p\pi_f, p)$  is the eigenvalue of the operator  $T_p$  defined in 3.1.2.

We briefly come back to the question of convergence. Hecke proves in [38] that Estone

$$|c(\pi_f, a)| \leq Ca^{n+1+\epsilon} \quad (4.123)$$

and with this estimate the convergence becomes obvious.

Actually there is a much better estimate, which will be discussed in the "probably removed" section. Lfu

### 4.1.7 The $L$ -functions

We still assume that  $K_f = \text{Gl}_2(\hat{\mathbb{Z}})$  or what amounts to the same that  $\Gamma = \text{Sl}_2(\mathbb{Z})$ . We start from an eigenspace  $H_1^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_\lambda \otimes F)(\pi_f)$ , now  $\pi_f$  is simply a homomorphism  $\pi_f : \mathcal{H}_{K_f} \rightarrow \mathcal{O}_F$ . To this homomorphism we attach the cohomological  $L$ -function

$$L^{\text{coh}}(\pi_f, s) = \prod_p \frac{1}{1 - \pi_p(T_p)p^{-s} + p^{1+n-2s}} \quad (4.124)$$

here  $T_p$  is the Hecke operator defined in 3.1.2, it differs from the Hecke operator defined by convolution by a factor  $p^{\frac{n}{2}}$  in front. If we expand this product over all primes we get

$$L^{\text{coh}}(\pi_f, s) = \sum_{a=1}^{\infty} \frac{c(\pi_f, a)}{a^s} \quad (4.125)$$

and this is exactly the  $L$ -function Hecke attaches to the cusp form provided by  $\pi_f$ . But we want to stress that this cohomological  $L$ -function is defined in purely combinatorial terms (See section 3.2.1, and Chapter 7).

At this moment this  $L$  function is a formal expression, it is a formal Dirichlet series with coefficients in our field  $F$ , which is simply a finite extension of  $\mathbb{Q}$ . If we assume that  $F \subset \mathbb{C}$ . then we may interpret  $s$  as a complex variable and the above estimate of the size of the coefficients implies that this series converges absolutely and locally uniformly for  $\Re(s) > n + 2$  and hence gives a holomorphic function in this halfspace. But something much better is true. We define the completed  $L$  function

$$\Lambda^{\text{coh}}(\pi_f, s) = \frac{\Gamma(s)}{(2\pi)^s} L^{\text{coh}}(\pi_f, s), \quad (4.126)$$

for this completed  $L$ -function Hecke proved

**Theorem 4.1.6.** *The function  $\Lambda^{\text{coh}}(\pi_f, s)$  has holomorphic continuation into the entire complex plane and satisfies the functional equation*

$$\Lambda^{\text{coh}}(\pi_f, s) = (-1)^{\frac{n}{2}+1} \Lambda^{\text{coh}}(\pi_f, n + 2 - s)$$

*Proof.* We could refer to Hecke, but for some reason we give an outline of the argument. We have the integral representation (Mellin-transform)

$$\Lambda^{\text{coh}}(\pi_f, s) = \int_0^{\infty} \sum_{a=1}^{\infty} c(\pi_f, a) e^{-2\pi a y} y^s \frac{dy}{y} = \int_0^{\infty} h^{\Phi}(iy) y^s \frac{dy}{y}$$

of course here we have to be courageous ( or stupid ) enough to exchange integration and summation. But since  $e^{-2\pi a y}$  goes rapidly to zero if  $y \rightarrow \infty$  there is no problem with the upper integration limit  $\infty$ . If  $\Re(s) \gg 0$  the  $y^s$  also tends to zero fast enough, so that we do not have a problem with the lower integration limit. But now we can split the integration into two parts

$$\begin{aligned} & \int_0^{\infty} \sum_a^{\infty} c(\pi_f, a) e^{-2\pi a y} y^s \frac{dy}{y} = \\ & \int_0^1 \sum_a^{\infty} c(\pi_f, a) e^{-2\pi a y} y^s \frac{dy}{y} + \int_1^{\infty} \sum_a^{\infty} c(\pi_f, a) e^{-2\pi a y} y^s \frac{dy}{y} \end{aligned}$$

the second integration is converging for all values of  $s$ . To handle the first integral we observe that  $h^{\Phi}(-\frac{1}{z}) = z^{n+2} h^{\Phi}(z)$ , Hence we can substitute  $y \rightarrow \frac{1}{y}$  in the first integral and get

$$\begin{aligned} \Lambda^{\text{coh}}(\pi_f, s) = \\ \sum_a^{\infty} \left( \frac{1}{(2\pi)^s} \frac{c(\pi_f, a)}{a^s} \Gamma(s, 2\pi a) + \frac{(-1)^{\frac{n}{2}+1}}{(2\pi)^{n+2-s}} \frac{c(\pi_f, a)}{a^{n+2-s}} \Gamma(n + 2 - s, 2\pi a) \right). \end{aligned} \quad (4.127)$$

Here  $\Gamma(\cdot)$  is the incomplete  $\Gamma$  function, which defined by  $\Gamma(s, A) = \int_A^\infty e^{-y} y^s \frac{dy}{y}$ , it has the virtue that for any given value of  $s$  it decays rapidly if  $A$  goes to infinity.

Therefore it is clear that the summations in the above formula are converging very fast, hence it follows that  $\Lambda^{\text{coh}}(\pi_f, s)$  is holomorphic in the entire  $s$  plane and the functional equation also becomes obvious.  $\square$

We included the proof of the above theorem, because the above formula also gives us a very effective procedure to compute the numerical value of  $\Lambda^{\text{coh}}(\pi_f, s_0)$  with high accuracy. We will come back to this issue in section 5.6.

periods

### 4.1.8 The Periods

Together with the map  $\mathcal{F}_1$  comes the map

$$\begin{aligned} \tilde{\mathcal{F}}_1 &= \text{Id} \otimes \mathcal{F}_1 \otimes \text{Id} : \text{Hom}_{\tilde{K}_\infty}(\Lambda(\mathfrak{g}/\mathfrak{k}), \mathcal{W}(\pi) \otimes \tilde{\mathcal{M}}_\lambda) \rightarrow \\ &\text{Hom}_{\tilde{K}_\infty}(\Lambda(\mathfrak{g}/\mathfrak{k}), \text{Gl}_2(\mathbb{Q}) \backslash \mathcal{C}_\infty(\text{Gl}_2(\mathbb{R}) \times \text{Gl}_2(\mathbb{A}_f)/K_f) \otimes \tilde{\mathcal{M}}_\lambda) \end{aligned}$$

The purpose of the following computations is to fix a specific choice of basis elements  $\omega_\pm^\dagger \in \text{Hom}_{\tilde{K}_\infty}(\Lambda^1(\mathfrak{g}/\mathfrak{k}), \mathcal{D}_\lambda^\dagger \otimes \tilde{\mathcal{M}}_\lambda)$  (in case A)  $\omega_{1,2}^\dagger \in \text{Hom}_{\tilde{K}_\infty}(\Lambda^{1,2}(\mathfrak{g}/\mathfrak{k}), \mathcal{D}_\lambda^\dagger \otimes \tilde{\mathcal{M}}_\lambda)$  (in case B)) These "canonical" generators serve us to define the periods.

In case A) we have

$$\mathfrak{g}/\mathfrak{k} \xrightarrow{\sim} \mathbb{Q} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \oplus \mathbb{Q} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \mathbb{Q}H \oplus \mathbb{Q}V = \mathfrak{p} \quad (4.128)$$

If we put  $P = H + V \otimes i, \bar{P} = H - V \otimes i \in \mathfrak{g}/\mathfrak{k} \otimes \mathbb{Q}(i)$

**Notation abklaeren**  $V = E_+$  **auf S. 123 ?**) then

$$\mathfrak{g}/\mathfrak{k} \otimes \mathbb{Q}(i) = \mathbb{Q}(i)P \oplus \mathbb{Q}(i)\bar{P} \text{ and } e(\phi)Pe(-\phi) = e^{2i\phi}P; e(\phi)\bar{P}e(-\phi) = e^{-2i\phi}\bar{P} \quad (4.129)$$

Let  $P^\vee, \bar{P}^\vee \in \text{Hom}(\mathfrak{g}/\mathfrak{k}, \mathbb{Q}(i))$  be the dual basis. Then we check easily that Pvee

$$P^\vee(H) = \bar{P}^\vee(H) = \frac{1}{2} \text{ and } P^\vee(V) = -\frac{i}{2}, \bar{P}^\vee(V) = \frac{i}{2} \quad (4.130)$$

The module  $\tilde{\mathcal{M}}_\lambda \otimes \mathbb{Q}(i)$  decomposes under the action of  $\tilde{K}_\infty$  into eigenspaces under  $\tilde{K}_\infty$

$$\mathcal{M}_{\lambda^\vee} \otimes \mathbb{Q}(i) = \bigoplus_{\nu}^n \mathbb{Q}(i)(X + Y \otimes i)^{n-\nu}(X - Y \otimes i)^\nu \quad (4.131)$$

where

$$e(\phi)((X + Y \otimes i)^{n-\nu}(X - Y \otimes i)^\nu) = e^{\pi i(n-2\nu)\phi} \cdot (X + Y \otimes i)^{n-\nu}(X - Y \otimes i)^\nu.$$

Then we define the basis elements

$$\omega^\dagger = P^\vee \otimes \tilde{\psi}_{n+2} \otimes (X - Y \otimes i)^n; \bar{\omega}^\dagger = \bar{P}^\vee \otimes \tilde{\psi}_{-n-2} \otimes (X + Y \otimes i)^n \quad (4.132)$$

We still have our involution  $\mathbf{c} \in \tilde{K}_\infty^*$  (See (4.22)) and clearly we have  $\mathbf{c}\omega^\dagger = i^n \bar{\omega}^\dagger$  (Remember  $n \equiv 0 \pmod{2}$ .)

Now we put OPM

$$\omega_+^\dagger = \frac{1}{2}(\omega^\dagger + i^n \bar{\omega}^\dagger); \quad \omega_-^\dagger = \frac{1}{2}(\omega^\dagger - i^n \bar{\omega}^\dagger) \quad (4.133)$$

then these elements

$$\omega_\pm^\dagger = \frac{1}{2}(\omega^\dagger \pm i^n \bar{\omega}^\dagger) \in \text{Hom}_{K_\infty}(\Lambda^1(\mathfrak{g}/\mathfrak{k}), \tilde{\mathcal{D}}_\lambda \otimes \mathcal{M}_\lambda)_\pm$$

and they are generators of these one dimensional spaces. The choice of these generators seems to be somewhat arbitrary, in [33] we give some motivation for this choice.

There is an alternative way to select  $\omega_\pm^\dagger$ . If we evaluate  $\omega_\pm^\dagger$  on the element  $H \in \mathfrak{g}/\mathfrak{k} = \mathfrak{p}$  then

$$\omega_\pm^\dagger(H) = \frac{1}{4}(\psi_{n+2}^\dagger \otimes (X - Y \otimes i)^n \pm i^n (\psi_{-n-2}^\dagger \otimes (X + Y \otimes i)^n) \in \mathcal{D}_\lambda^\dagger \otimes \mathcal{M}_\lambda$$

These are functions on  $\text{Gl}_2(\mathbb{R})$  with values in  $\mathcal{M}_\lambda$ . We pair these functions with an  $\mathcal{M}_\lambda \otimes \mathbb{C}$  valued function, more precisely we consider the function  $g \mapsto \langle \omega_\pm^\dagger(\text{Ad}(g)H)(g), \rho_\lambda(g)X^\nu Y^\nu \rangle$ .

We restrict these scalar valued functions to the real points of the split torus

$$\langle \omega_\pm^\dagger(H)\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right), \rho_\lambda\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right)X^\nu Y^{n-\nu} \rangle =$$

$$\langle \frac{1}{4}(\psi_{n+2}^\dagger \otimes \left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right) \otimes (X - Y \otimes i)^n \pm i^n \psi_{-n-2}^\dagger \otimes \left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right) \otimes (X + Y \otimes i)^n), X^\nu Y^{n-\nu} \rangle t^{-\frac{n}{2} + \nu}$$

Now let  $\epsilon$  be a variable which can take the values  $+, -$ , then  $\epsilon 1 = +1, -1$ . Our formula (4.8) gives us  $\langle (X - \epsilon Y \otimes i)^n, X^\nu Y^{n-\nu} \rangle = (-\epsilon i)^{n-\nu}$  and combing this with the explicit formula (??) for the values of  $\psi_{\epsilon(n+2)}^\dagger\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right)$  we get

$$\langle \omega_\epsilon^\dagger(H)\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right), \rho_\lambda\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right)X^\nu Y^{n-\nu} \rangle = \begin{cases} (-i)^{n-\nu} t^{\frac{n}{2}+1} e^{-2\pi t} t^{-\frac{n}{2}+\nu} & \text{for } t > 0 \\ \epsilon i^{n-\nu} (-t)^{\frac{n}{2}+1} e^{2\pi t} (-t)^{-\frac{n}{2}+\nu} & \text{for } t < 0 \end{cases}$$

(Here we use that  $n$  is even, but with suitable minor modifications we can also treat the case  $n$  odd.) Then a straight forward computation yields Mellin

$$\int_{T^{\text{ad}}(\mathbb{R})} \langle \omega_\epsilon^\dagger(H)\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right), \rho_\lambda\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right)X^\nu Y^{n-\nu} \rangle \frac{dt}{t} = \frac{1}{2} \begin{cases} \frac{\Gamma(n+1-\nu)}{(2\pi)^{n+1-\nu}} & \text{if } (-1)^{\frac{n}{2}-\nu} = \text{sg}(\epsilon) \\ 0 & \text{else} \end{cases} \quad (4.134)$$

For each choice of the sign  $\epsilon = \pm 1$  one of these equation determines the generator  $\omega_\pm^\dagger$ . This formula will be of importance when we discuss the special values of  $L$ -functions.

In case B) we do basically the same, in some sense it is even simpler because  $K_\infty$  is maximal compact in this case, i.e.  $K_\infty = K_\infty^*$ . But on the other hand we need some very explicit information about the theory of irreducible representations of  $K_\infty$  and also about the decomposition of tensor products of these representations. We will also use some explicit formulas for Bessel functions.

The quotient  $\mathfrak{g}/\mathfrak{k}$  is a three-dimensional vector space over  $\mathbb{Q}$  the group  $K_\infty$  acts by the adjoint representation and this gives us the standard three dimensional representation of  $K_\infty = U(2)$ , which in addition is trivial on the center. (See 4.1.2). This module is given by the highest weight  $2\gamma_c$ . We must have  $\lambda = n(\gamma + \bar{\gamma}) + \dots$ , if we want  $\mathcal{E}^{(2)}(\lambda, w, \Gamma) \neq 0$ , and then the formulae 4.50 and 4.51 imply that for  $\bullet = 1, 2$

$$\dim_{\mathbb{C}} \operatorname{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \tilde{\mathcal{D}}_\lambda \otimes \mathcal{M}_{\lambda^\vee}) = 1 \quad (4.135)$$

Now we recall that we have defined a structure of a  $R = \mathbb{Z}[\frac{1}{2}]$  module on all the modules on the stage, hence we see that

$$\operatorname{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \tilde{\mathcal{D}}_\lambda \otimes \mathcal{M}_{\lambda^\vee}) = \operatorname{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k})_R, \tilde{\mathcal{D}}_{\lambda R} \otimes \mathcal{M}_{\lambda^\vee R}) \otimes \mathbb{C}, \quad (4.136)$$

here we are a little bit sloppy: The first subscript  $K_\infty$  is the compact group and the second subscript is a smooth groups scheme over  $R$ . For both choices of  $\bullet$  the second term in the above equation is a free  $R$  module of rank 1. We choose generators

$$\omega^{\dagger, \bullet} \in \operatorname{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k})_R, \tilde{\mathcal{D}}_{\lambda R} \otimes \mathcal{M}_{\lambda^\vee R}).$$

These generators  $\omega^{\dagger, 1}, \omega^{\dagger, 2}$  are well defined up to an element in  $R^\times$ .

Of course we may also fix these generators by prescribing values of certain Mellin transforms. To do this we need a little bit of representation theory. Of course we may replace  $K_\infty$  by  $SU(2)$  because the action of the center on the different modules cancels out. The modules  $\mathfrak{g}/\mathfrak{k} \otimes \mathbb{C}, \tilde{\mathcal{D}}_\lambda$  and  $\mathcal{M}_{\lambda^\vee} \otimes \mathbb{C}$  extend naturally to  $Sl_2(\mathbb{C})$  modules and hence we have to find an explicit generator in

$$\operatorname{Hom}_{Sl_2(\mathbb{C})}(\mathfrak{g}/\mathfrak{k} \otimes \mathbb{C}, \tilde{\mathcal{D}}_\lambda \otimes \mathcal{M}_{n\gamma} \otimes \mathcal{M}_{n\bar{\gamma}}).$$

We have an explicit basis for  $\mathfrak{g}/\mathfrak{k} \otimes \mathbb{C}$  (See (4.16), our module  $\mathcal{M}_{\lambda^\vee} = \mathcal{M}_{n\gamma}^b \otimes \mathcal{M}_{n\bar{\gamma}}^b \otimes_{\mathbb{C}} \mathbb{C}$  is given explicitly to us.

Our module  $\tilde{\mathcal{D}}_\lambda \subset \mathcal{I}_B^G \lambda_{\mathbb{R}} \rho_{\mathbb{R}}^2$ , and this last module decomposes into  $SU(2)$ -types (See( 4.28). These  $SU(2)$  modules canonically extend to  $Sl_2(\mathbb{C})$ -modules, we have

$$\mathcal{I}_B^G \lambda_{\mathbb{R}} \rho_{\mathbb{R}}^2 = \bigoplus_{\nu=0}^{\infty} \mathcal{M}_{2\nu\gamma} = \bigoplus_{\nu=0}^{\infty} \mathcal{I}_B^G \lambda_{\mathbb{R}} \rho_{\mathbb{R}}^2(2\nu)$$

and

$$\mathcal{I}_B^G \lambda_{\mathbb{R}} \rho_{\mathbb{R}}^2(2(n+1)) = \tilde{\mathcal{D}}_\lambda(2(n+1))$$

Now it is clear that our problem is to select a specific generator in

$$\operatorname{Hom}_{Sl_2(\mathbb{C})}(\mathfrak{g}/\mathfrak{k} \otimes \mathbb{C}, \tilde{\mathcal{D}}_\lambda(2(n+1)) \otimes \mathcal{M}_{n\gamma}^b \otimes \mathcal{M}_{n\bar{\gamma}}^b \otimes \mathbb{C}).$$

The modules  $\mathfrak{g}/\mathfrak{k} \otimes \mathbb{C}, \mathcal{M}_{n\gamma}^b, \mathcal{M}_{n\bar{\gamma}}^b$  come with an explicit basis (See ???), if we want to write down a specific  $\omega^{\dagger, \bullet}$  we have to write down a basis of  $\tilde{\mathcal{D}}_\lambda(2(n+1))$ .

To get such a basis we start from a basis element  $\Phi_\lambda \in \tilde{\mathcal{I}}_B^G \lambda_{\mathbb{R}} \rho_{\mathbb{R}}^2(0)$ . We recall the definition of  $\mathcal{I}_B^G \lambda_{\mathbb{R}} \rho_{\mathbb{R}}^2$  as an induced representation, the space of  $K_\infty$  invariant vectors is spanned by the spherical function

$$\Psi_\lambda(bk) = \Psi\left(\begin{pmatrix} t_1 & u \\ 0 & t_2 \end{pmatrix} \begin{pmatrix} a & b \\ -\bar{b} & \bar{a} \end{pmatrix}\right) = \lambda_{\mathbb{R}} \rho_{\mathbb{R}}^2(b).$$

We map the induced representation to its Whittaker model by

$$\mathcal{F} : \Psi \mapsto \left\{ g \mapsto \int \Psi\left(w \begin{pmatrix} 1 & x + iy \\ 0 & 1 \end{pmatrix} g\right) e^{2\pi i x} dx dy \right\} \quad (4.137)$$

our basis element will be  $\Phi_\lambda = \mathcal{F}(\Psi_\lambda)$ . A straightforward computation yields

$$\Phi_\lambda\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right) = \mathcal{F}(\Psi_\lambda)\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right) = \int_{-\infty}^{\infty} \frac{t^{n+2}}{(t^2 + x^2 + y^2)^{n+2}} e^{2\pi i x} dx dy$$

The educated reader knows that this function in the variable  $t$  is well known, we have

$$\Phi_\lambda\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right) = \frac{2\pi^{n+2}}{\Gamma(n+2)} t K_{n+1}(2\pi t)$$

where  $K_n(2\pi t)$  is the modified Bessel function. Of course  $\Phi_\lambda$  is a function on  $G(\mathbb{R}) = \mathrm{GL}_2(\mathbb{C})$ , it is right invariant under  $K_\infty$  and of course

$$\Phi_\lambda\left(\begin{pmatrix} 1 & x + iy \\ 0 & 1 \end{pmatrix} \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right) = e^{2\pi i x} \Phi_\lambda\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right)$$

hence it is defined by its restriction to  $T^{\mathrm{ad}}(\mathbb{R})_{>0}$ .

Starting from this function we construct the desired basis of  $\tilde{\mathcal{D}}_\lambda(2(n+1))$ . The Lie-algebra  $\mathfrak{g}$  acts on  $\mathcal{I}_B^G \lambda_{\mathbb{R}} \rho_{\mathbb{R}}^2$ , we restrict this action to  $\mathfrak{p}$  and it is clear that under this action

$$\mathfrak{p} \otimes \mathcal{I}_B^G \lambda_{\mathbb{R}} \rho_{\mathbb{R}}^2(2\nu) \rightarrow \mathcal{I}_B^G \lambda_{\mathbb{R}} \rho_{\mathbb{R}}^2(2\nu + 2) \oplus \mathcal{I}_B^G \lambda_{\mathbb{R}} \rho_{\mathbb{R}}^2(2\nu) \oplus \mathcal{I}_B^G \lambda_{\mathbb{R}} \rho_{\mathbb{R}}^2(2\nu - 2)$$

and if we extend this action to the tensor algebra we get a map

$$\mathfrak{p}^{\otimes n+1} \otimes \mathcal{I}_B^G \lambda_{\mathbb{R}} \rho_{\mathbb{R}}^2(0) \rightarrow \bigoplus_{\nu=0}^{n+1} \mathcal{I}_B^G \lambda_{\mathbb{R}} \rho_{\mathbb{R}}^2(2\nu).$$

The group  $K_\infty$  acts on  $\mathfrak{p}^{\otimes n+1}$  by the adjoint action and the above map is of course a  $K_\infty$  homomorphism. On the right hand side we can project to the highest  $K_\infty$  type  $\mathcal{I}_B^G \lambda_{\mathbb{R}} \rho_{\mathbb{R}}^2(2n+2) = \tilde{\mathcal{D}}_\lambda(2(n+1))$ , i.e. we get a surjective homomorphism

$$\Pi_{n+1} : \mathfrak{p}^{\otimes n+1} \otimes \mathcal{I}_B^G \lambda_{\mathbb{R}} \rho_{\mathbb{R}}^2(0) \rightarrow \tilde{\mathcal{D}}_\lambda(2(n+1)).$$

We have the standard surjective homomorphism  $\mathfrak{p}^{\otimes n+1} \rightarrow \mathrm{Sym}^{n+1}(\mathfrak{p})$ , let us denote its kernel by  $I_{n+1}$ . For any  $f \in \mathcal{I}_B^G \lambda_{\mathbb{R}} \rho_{\mathbb{R}}^2$  and  $X_1, X_2 \in \mathfrak{p}$  we have

$$(X_1 X_2 - X_2 X_1)f = [X_1, X_2]f.$$

Since the Lie bracket  $[X_1, X_2] \in \mathfrak{k}$  it follows easily that  $\Pi_{n+1}$  vanishes on the kernel  $I_{n+1}$ . Hence our homomorphism  $\Pi_{n+1}$  factors over the quotient, i.e.

$$\Pi_{n+1} : \text{Sym}^{n+1}(\mathfrak{p}) \rightarrow \tilde{\mathcal{D}}_\lambda(2(n+1)).$$

We change our notation for the basis of  $\mathfrak{p} \otimes \mathbb{C}$  (see 4.16) and put

$$\begin{aligned} X_1 &= \frac{1}{2} \left( \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} - i \otimes \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix} \right); X_0 = \frac{1}{\sqrt{2}} \left( \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \right) \\ X_{-1} &= \frac{1}{2} \left( \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + i \otimes \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix} \right) \end{aligned} \quad (4.138)$$

We have the following proposition

**Proposition 4.1.7.** *The  $2n + 3$  elements*

$$\{X_1^{n+1}, X_0 X_1^n, \dots, X_0^{n+1}, X_0^n X_{-1}, \dots, X_{-1}^{n+1}\}$$

form a basis of a  $K_\infty$  invariant subspace of  $\text{Sym}^{n+1}(\mathfrak{p}) \otimes \mathbb{C}$ . This subspace is irreducible, it is isomorphic to  $\mathcal{M}_{2n+2}$ . These basis elements are the weight eigenvectors for the action of  $T_c$ .

*Proof.* The representation of the algebraic group  $K_\infty$  on  $\mathfrak{p}$  extends to a representation of the algebraic group  $\text{Sl}_2/\mathbb{C}$  on  $\mathfrak{p} \otimes \mathbb{C}$ . As such it is isomorphic to the symmetric square  $\text{Sym}^2(\mathbb{C}^2)$  of the tautological representation, i.e. to the module  $\mathcal{M}_2$  of polynomials  $aU^2 + bUV + cV^2$ . We get an isomorphism  $\mathcal{M}_2 \xrightarrow{\sim} \mathfrak{p} \otimes \mathbb{C}$  by sending  $U^2 \mapsto X_1, UV \mapsto X_0, V^2 \mapsto X_{-1}$ . Now  $\text{Sym}^{2n+2}(\mathcal{M}_2) \subset \text{Sym}^{n+1}(\text{Sym}^2(\mathbb{C}^2)) = \text{Sym}^{n+1}(\mathfrak{p} \otimes \mathbb{C})$  is an invariant submodule. It has the basis  $U^{2n+2-\nu} V^\nu$  and clearly

$$U^{2n+2-\nu} V^\nu = X_1^{n+1-\nu} X_0^\nu \text{ if } \nu \leq n+1 \text{ and } X_0^{n+1\nu} X_1$$

and this implies the assertion.  $\square$

This implies that the elements

$$\{\Pi_{n+1}(X_1^{n+1}\Phi_\lambda), \Pi_{n+1}(X_0 X_1^n \Phi_\lambda), \dots, \Pi_{n+1}(X_0^n X_1 \Phi_\lambda), \Pi_{n+1}(X_0^{n+1} \Phi_\lambda), \Pi_{n+1}(X_0^n X_{-1} \Phi_\lambda), \dots, \Pi_{n+1}(X_{-1}^{n+1} \Phi_\lambda)\} \quad (4.139)$$

form a basis of  $\tilde{\mathcal{D}}_\lambda(2(n+1))$ .

We change our notation slightly. For  $m < 0$  we put  $X_1^m := X_{-1}^{-m}$  and for  $0 \leq \nu \leq 2n+2$  we put  $[\nu] = \nu$  if  $\nu \leq n+1$  and  $[\nu] = 2n+2-\nu$  if  $\nu \geq n+1$ . Then our above basis can be written as

$$\{\dots, \Pi_{n+1}(X_0^{[\nu]} X_1^{n+1-\nu} \Phi_\lambda), \dots\}_{\nu=0, \dots, \nu=2n+2}, \quad (4.140)$$

these are the weight vectors of weight  $2(n+1-\nu)\gamma$ . We introduce the notation

$$\Phi_{\lambda, \nu} := \Pi_{n+1}(X_0^{[\nu]} X_1^{n+1-\nu} \Phi_\lambda)$$

These functions  $\Phi_{\lambda, \nu}$  are Whittaker functions they satisfy

$$\Phi_{\lambda, \mu} \left( \begin{pmatrix} 1 & x + iy \\ 0 & 1 \end{pmatrix} g \right) = e^{2\pi i x} \Phi_{\lambda, \mu}(g).$$

But of course they are not  $K_\infty$  invariant, we have

$$\Phi_{\lambda,\nu}(g \begin{pmatrix} e^{i\phi} & 0 \\ 0 & e^{-i\phi} \end{pmatrix}) = e^{(2n+2-2\nu)i\phi} \Phi(g)$$

and more generally  $\Phi_{\lambda,\nu}(gk) = \sum_\mu a_{\nu,\mu}(k) \Phi_{\lambda,\mu}(g)$  where the  $a_{\nu,\mu}(k)$  are the matrix coefficients of  $\mathcal{M}_{2n+2}$ . (above proposition).

We express the restriction of these functions  $\Phi_{\lambda,\nu}$  to the torus  $T^{\text{ad}}(\mathbb{R})_{>0}$  in terms of Bessel functions. It is quite clear that for any Whittaker function  $\Phi$  of  $K_\infty$  type  $\leq k$  we have

$$\Pi_{k+1}(X_1 \Phi) \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) = \frac{\Phi \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \exp(\epsilon X_1) \right) - \Phi \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right)}{\epsilon}$$

We write  $X_1 = \frac{1}{2} \left( \begin{pmatrix} 0 & 2 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 2i \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} + \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \right)$  the last two matrices are in  $\mathfrak{k}$  so they preserve the  $K_\infty$  type and

$$\begin{aligned} & \Phi \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \exp(\epsilon \left( \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & i \\ 0 & 0 \end{pmatrix} \right)) \right) = \\ & \Phi \left( \exp(\epsilon \left( \begin{pmatrix} 0 & t \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & it \\ 0 & 0 \end{pmatrix} \right) \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) = \Phi \left( \left( \begin{pmatrix} 1 & \epsilon(t+it) \\ 0 & 1 \end{pmatrix} \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) \right) = \\ & e^{2\pi i \epsilon t} \Phi \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) = \Phi \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) (1 + it\epsilon) \end{aligned}$$

and hence

$$\Pi_{k+1}(X_1 \Phi) \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) = 2it \Phi \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right)$$

This gives us

$$\Phi_{\lambda,0} \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) = (X_1^{n+1} \Phi_\lambda) \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) = \frac{2^{2n+3} \pi^{2n+3}}{\Gamma(n+2)} t^{n+2} K_{n+1}(2\pi t) \quad (4.141)$$

Since this function is of weight  $2n+2$  we can forget the projection  $\Pi_{n+1}$ .

**Bessel formulas:**

$$\begin{aligned} \frac{d}{dt} K_n(t) &= -\frac{1}{2} (K_{n-1}(t) + K_{n+1}(t)) \\ K_{n+1}(t) &= K_{n-1}(t) + \frac{2n}{t} K_n(t) \end{aligned} \quad (4.142)$$

We also will need the Mellin transforms of these Bessel functions. Here we quote [1] .p.331,334

$$\begin{aligned} \int_0^\infty K_\nu(2\pi t) t^s \frac{dt}{t} &= 2^{s-2} (2\pi)^{-s} \Gamma\left(\frac{s-\nu}{2}\right) \Gamma\left(\frac{s+\nu}{2}\right) \\ \int_0^\infty K_\mu(2\pi t) K_\nu(2\pi t) t^s \frac{dt}{t} &= 2^{s-3} (2\pi)^{-s} \Gamma\left(\frac{s-\mu-\nu}{2}\right) \Gamma\left(\frac{s-\mu+\nu}{2}\right) \Gamma\left(\frac{s+\mu-\nu}{2}\right) \Gamma\left(\frac{s+\nu+\mu}{2}\right) \end{aligned} \quad (4.143)$$

This gives us

$$\begin{aligned}
 t \frac{d}{dt} t^\nu K_{n+1}(2\pi t) &= \nu t^\nu K_{n+1}(2\pi t) - \pi t^{\nu+1} (K_{n+2}(2\pi t) + K_n(2\pi t)) = \\
 \nu t^\nu K_{n+1}(2\pi t) - \pi t^{\nu+1} (K_n(2\pi t) + \frac{(n+1)}{\pi t} K_{n+1}(2\pi t)) &= \tag{4.144} \\
 (\nu - n - 1) t^\nu K_{n+1}(2\pi t) - \pi t^{\nu+1} K_n(2\pi t)
 \end{aligned}$$

We have  $\Phi_{\lambda, \nu} \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) = \Pi_{n+1} (X_0^{[\nu]} X_1^{n+1-\nu} \Phi_\lambda \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right))$ . The application of  $X_1^{n+1-\nu}$  gives us

$$\frac{(2\pi)^{n+2+|n+1-\nu|}}{\Gamma(n+2)} t^{1+|n+1-\nu|} K_{n+1}(2\pi t).$$

To this we apply  $X_0^{[\nu]}$ . The operator  $X_0$  is  $t \frac{d}{dt}$  then the formula above implies

$$\Pi_{n+1} (X_0^{[\nu]} X_1^{n+1-\nu} \Phi_\lambda \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right)) = \Pi_{n+1} (\dots + t^{[\nu]} \pi^{[\nu]} K_{n+1-\nu}(2\pi t)) \tag{4.145}$$

where the dots are a sum of terms  $at^m \pi^k K_m(2\pi t)$  where the  $m > |n+1-\nu|$ . For consistency reasons the  $\dots$  map to zero under  $\Pi_{n+1}$  this implies that

$$\Phi_{\lambda, \nu} \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) = \frac{2^{2n+3} \pi^{2n+3}}{\Gamma(n+2)} t^{n+2} K_{n+1-\nu} \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) \tag{4.146}$$

### Decompositions of tensor products

If  $\lambda_1 = n_1\gamma, \lambda_2 = n_2\gamma$  are two highest weights and if we consider the highest weight modules  $\mathcal{M}_{\lambda_1, \mathbb{Q}}, \mathcal{M}_{\lambda_2, \mathbb{Q}}$  then it is a classical theorem that

$$\mathcal{M}_{\lambda_1, \mathbb{Q}} \otimes \mathcal{M}_{\lambda_2, \mathbb{Q}} = \mathcal{M}_{(n_1+n_2)\gamma, \mathbb{Q}} \oplus \mathcal{M}_{(n_1+n_2-2)\gamma, \mathbb{Q}} \oplus \dots \oplus \mathcal{M}_{(n_1-n_2)\gamma, \mathbb{Q}} \dots$$

where we assume  $n_1 \geq n_2$ . Our next aim is to give an explicit homomorphism

$$j_{n_1, n_2} : \mathcal{M}_{(n_1+n_2)\gamma}^b \hookrightarrow \mathcal{M}_{n_1\gamma}^b \otimes \mathcal{M}_{n_2\gamma}^b \tag{4.147}$$

in other words we want to write explicit tensors for the images of  $e_\mu^b, \mu = n_1 + n_2, n_1 + n_2 - 2, \dots, -n_1 - n_2$ . Of course we send the the highest weight vector  $e_{n_1+n_2}^b \mapsto 'e_{n_1}^b \otimes ''e_{n_2}^b$ , this vector is the highest weight vector in the direct summand  $\mathcal{M}_{(n_1+n_2)\gamma, \mathbb{Q}}^b \subset \mathcal{M}_{(n_1+n_2)\gamma, \mathbb{Q}} \oplus \dots \oplus \mathcal{M}_{(n_1-n_2)\gamma, \mathbb{Q}}$ . In terms of the explicit realization of these modules we can say

$$X^{n_1+n_2} \mapsto 'X^{n_1} \otimes ''X^{n_2} \tag{4.148}$$

Now we apply the matrix  $\begin{pmatrix} 1 & 0 \\ t & 1 \end{pmatrix}$  to it, here we may think of  $t$  as an in determinant. Then we see

$$(X + tY)^{n_1+n_2} \mapsto ('X + t'Y)^{n_1} \otimes (''X + t''Y)^{n_2} \tag{4.149}$$

We expand on both sides and find

$$\sum_{\mu=0}^{n_1+n_2} \binom{n_1+n_2}{\mu} t^\mu X^{n_1+n_2-\mu} Y^\mu \mapsto \sum_{\mu=0}^{n_1+n_2} t^\mu \left( \sum_{\mu_1, \mu_2: \mu_1+\mu_2=\mu} \binom{n_1}{\mu_1} 'X^{n_1-\mu_1} Y^{\mu_1} \otimes \binom{n_2}{\mu_2} ''X^{n_2-\mu_2} \otimes ''X^{n_2-\mu_2} Y^{\mu_2} \right) \quad (4.150)$$

We remember the definition of the basis elements  $e_\mu^b$ , the formula above gives us

$$j_{n_1, n_2} : e_\mu^b \mapsto \sum_{\mu_1+\mu_2=\mu} 'e_{\mu_1}^b \otimes ''e_{\mu_2}^b \quad (4.151)$$

We apply this to the  $SU(2)$ -module

$$(\mathfrak{g}/\mathfrak{k})_{\mathbb{F}}^{\vee} \otimes \mathcal{M}_{n\gamma} \otimes_F \mathcal{M}_{n\bar{\gamma}},$$

this module contains a unique copy of  $\mathcal{M}_{2n+2}^b$ . We write

$$\mathfrak{g}/\mathfrak{k}_{\mathbb{F}}^{\vee} = F^0 e_2^b \oplus F^0 e_0^b \oplus F^0 e_{-2}^b, \quad \mathcal{M}_{n\gamma, F} = \bigoplus_{\mu} F e_{\mu}^b, \quad \mathcal{M}_{n\bar{\gamma}, F} = \bigoplus_{\mu} F \bar{e}_{\mu}^b \quad (4.152)$$

where of course  $\mu$  runs from  $n$  to  $-n$  and  $\mu \equiv n \pmod{2}$ . Then our copy of  $\mathcal{M}_{2n+2}^b$  comes with the basis

$$\bar{e}_{\mu}^b = \sum_{\mu_0+\mu_1+\mu_2=\mu} {}^0 e_{\mu_0}^b \otimes e_{\mu_1}^b \otimes \bar{e}_{\mu_2}^b$$

We have the invariant pairing (4.8) and this tells us that we can choose as our generator cangen

$$\omega^{\dagger, \bullet} = \sum_{\mu=0}^{2n+2} \Phi_{\lambda, \mu} \otimes \left( \sum_{\mu_0+\mu_1+\mu_2=n+1-\mu} {}^0 e_{\mu_0}^b \otimes e_{\mu_1}^b \otimes \bar{e}_{\mu_2}^b \right) \quad (4.153)$$

This generator is only determined up to a scalar, it is fixed once we choose a generator  $\Phi_{\lambda, n+1}$ .

### The "canonical" choice of the generator

Again we can fix the generator by requiring that certain Mellin transforms have a prescribed value at certain prescribed arguments.

We do essentially the same as in the case A). We can interpret  $\omega^{\dagger, 1}$  as a differential 1-form on  $G(\mathbb{R})$  with values in  $\mathcal{M}_{\lambda}^b \otimes \mathbb{C}$ . We can restrict this 1-form to the torus  $T^{\text{ad}}(\mathbb{R})_{>0} = \left\{ \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \mid t > 0 \right\}$ . We have the "cycles"  $e_{\mu_1} \otimes e_{\mu_2} \in \mathcal{M}_{\lambda}^{\vee}$ .

We evaluate  $\omega^{\dagger, 1}(X_0)$  on these "cycles" and get

$$\langle \omega^{\dagger, \bullet}(X_0), e_{\mu_1} \otimes e_{\mu_2} \rangle \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) = \Phi_{\lambda, n-\mu_1-\mu_2} \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) t^{\mu_1+\mu_2} = c_n t^{n+2+\mu_1+\mu_2} K_{n-\mu_1-\mu_2}(2\pi t) \quad (4.154)$$

We observe that the first factor in front does not depend on  $\mu_1, \mu_2$ . So we modify our generator and for  $\nu = -n - 1, -n, \dots, n + 1$  we now put  $\boxed{\text{Phi}}$

$$\Phi_{\lambda, \mu}(t) = \frac{2\pi^n}{\Gamma(n+1)} t^{n+2} K_{n-\mu}(t) \quad (4.155)$$

and with this choice of  $\Phi_{\lambda, \nu}$  the  $\omega^{\dagger, 1}$  (8.197) is our canonical generator.

Hence we may just choose  $\mu_1 = \mu_2 = 0$  to nail down  $\omega^{\dagger, \bullet}$ , it is not clear to me whether or not it is a "miracle" that the above relation holds for all values of  $\mu_1, \mu_2$ .

### The definition of the periods

The inner cohomology with rational coefficients is a semi-simple module under the action of the Hecke algebra (See Theorem 3.2.1). We find a finite Galois-extension  $F/\mathbb{Q}$  such that

$$H_!^{\bullet}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{\lambda} \otimes F) = \bigoplus_{\pi_f} H_!^{\bullet}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{\lambda} \otimes F)(\pi_f) \quad (4.156)$$

We assume that  $\Gamma = \text{Gl}_2(\mathbb{Z})$ , hence the  $\pi_f$  are homomorphisms  $\pi_f : \mathcal{H} \rightarrow \mathcal{O}_F$ . (See ???) In the case A) such an isotypical piece is a direct sum

$$H_!^{\bullet}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{\lambda} \otimes F)(\pi_f) = H_!^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{\lambda} \otimes F)(\pi_f)_+ \oplus H_!^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{\lambda} \otimes F)(\pi_f)_- \quad (4.157)$$

where both summands are of dimension one over  $F$ .

In case B) we get

$$H_!^{\bullet}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{\lambda} \otimes F)(\pi_f) = H_!^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{\lambda} \otimes F)(\pi_f) \oplus H_!^2(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{\lambda} \otimes F)(\pi_f) \quad (4.158)$$

and again the summands are one dimensional.

We have defined the module of integral classes  $H_{!, \text{int}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{\lambda}^b \otimes R) \subset H_!^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{\lambda}^b \otimes F)$  (See 2.54) and we consider the intersection

$$H_{!, \text{int}}^{\bullet}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{\lambda}^b \otimes \mathcal{O}_F)(\pi_f)_{\epsilon} = H_!^{\bullet}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{\lambda} \otimes F)(\pi_f)_{\epsilon} \cap H_{!, \text{int}}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{\lambda}^b \otimes \mathcal{O}_F)$$

is a locally free  $\mathcal{O}_F$ -module of rank 1, here  $\epsilon = \pm, \bullet = 1$  ( resp.  $\epsilon = 1, \bullet \in \{1, 2\}$ ) We assume for simplicity that it is actually free, otherwise the formulation of the following becomes slightly more complicated. (See below). On the set of  $\pi_f$  which occur in this decomposition we have an action of the Galois group (See (Theorem 3.2.1)) and the Galois action yields canonical isomorphisms

$$\Phi_{\sigma, \tau} : H_{!, \text{int}}^{\bullet}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{\lambda}^b \otimes \mathcal{O}_F)(\sigma \pi_f)_{\epsilon} \xrightarrow{\sim} H_{!, \text{int}}^{\bullet}(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{\lambda}^b \otimes \mathcal{O}_F)(\tau \pi_f)_{\epsilon} \quad (4.159)$$

We choose generators  $\sigma e_{\epsilon}^{\bullet}(\pi_f)$  and a simple argument using Hilbert theorem 90 shows that we can assume the consistency condition  $\boxed{\text{H90}}$

$$\Phi_{\sigma, \tau}(e_{\epsilon}^{\bullet}(\sigma \pi_f)) = e_{\epsilon}^{\bullet}(\tau \pi_f) \quad (4.160)$$

We get isomorphisms

$$\mathcal{F}_1^\bullet(\omega_\epsilon^\dagger) : \mathcal{W}(\sigma\pi_f) \otimes_F \mathbb{C} \xrightarrow{\sim} H_\epsilon^\bullet(\Gamma \backslash \mathbb{H}, \mathcal{M}_{\lambda^\vee})(\sigma\pi_f) \otimes_{F,\iota} \mathbb{C} \quad (4.161)$$

which is defined by Armand1

$$\mathcal{F}_1^\bullet(\omega_\epsilon^\dagger) : h_{\sigma\pi_f} \mapsto [\mathcal{F}_1(\omega_\epsilon^\dagger \otimes h_{\sigma\pi_f})], \quad (4.162)$$

here  $\mathcal{F}_1(\omega_\epsilon^\dagger \otimes h_{\sigma\pi_f})$  the (closed)  $\mathcal{M}_\lambda \otimes \mathbb{C}$  valued differential form corresponding to  $\mathcal{F}_1(\omega_\epsilon^\dagger \otimes h_{\sigma\pi_f})$  under the identification 8.3 and  $[\ ]$  is its class in cohomology.

Since we assume that  $\pi_f$  is unramified everywhere  $\mathcal{W}(\pi_f)$  we have the canonical basis element  $h_f^{(0)} = \prod_p h_{\sigma\pi_p}^{(0)}$  where  $h_{\sigma\pi_p}^{(0)}$  is defined by the equality 4.134.

Then we have obviously  $\sigma(h_{\pi_p}^{(0)}) = h_{\sigma\pi_p}^{(0)}$ .

Then we define the *periods* by the relation

$$\mathcal{F}_1(\omega_\epsilon^\dagger)(h_{\sigma\pi_f}^{(\dagger,0)}) = \Omega^\bullet(\sigma\pi_f, \epsilon) e_\epsilon^\bullet(\sigma\pi_f) \quad (4.163)$$

These periods depend of course on our choice of the "canonical" generator  $\omega_\epsilon^\dagger$ . We see that the numbers  $\Omega^\bullet(\sigma\pi_f, \epsilon)$  are well defined up to an element in  $\mathcal{O}_F^\times$ .

If  $H_{!, \text{int}}^\bullet(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_\lambda^b \otimes \mathcal{O}_F)(\pi_f)_\epsilon$  is not a free  $\mathcal{O}_F$  module, then we can find a covering by two open subsets  $U_1, U_2$  of  $\text{Spec}(\mathcal{O}_F)$  such that  $H_{!, \text{int}}^\bullet(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_\lambda^b \otimes \mathcal{O}_F(U_i))(\pi_f)_\epsilon$  is free. We can apply the above procedure and we get periods  $\Omega_1(\pi_f, \epsilon), \Omega_2(\pi_f, \epsilon)$ , they are well defined up to an element in  $\mathcal{O}_F(U_1)^\times, \mathcal{O}_F(U_2)^\times$  respectively. The ratio of these periods is an element in  $\mathcal{O}_F(U_1 \cap U_2)^\times$ . In the following we always pretend that the respective modules are free, just to simplify the formulation.

### Some little subtleties

We should notice that these periods are defined with respect to the "small" sheaves  $\tilde{\mathcal{M}}_\lambda^b$ . We have  $\tilde{\mathcal{M}}_\lambda^b \subset \tilde{\mathcal{M}}_\lambda$  and therefore the map

$$H_{!, \text{int}}^\bullet(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_\lambda^b \otimes \mathcal{O}_F)(\pi_f)_\epsilon \rightarrow H_{!, \text{int}}^\bullet(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_\lambda \otimes \mathcal{O}_F)(\pi_f)_\epsilon \quad (4.164)$$

may not be surjective. (The reader should not be puzzled by the fact that  $\tilde{\mathcal{M}}_\lambda^b \otimes F = \tilde{\mathcal{M}}_\lambda \otimes F$ .) Therefore, if we would work with  $\tilde{\mathcal{M}}_\lambda$  instead and define the periods  $\Omega^{\bullet, \#}(\sigma\pi_f, \epsilon)$  by the same procedure. Then we will get a relation

$$\Omega^{\bullet, \#}(\sigma\pi_f, \epsilon) = d(\pi_f, \epsilon) \Omega^\bullet(\sigma\pi_f, \epsilon)$$

where  $d(\pi_f, \epsilon)$  is a non zero factor in  $\mathcal{O}_F$ . The primes in these factors are the divisors of the binomial coefficients.

But we could also with the module  $H^\bullet(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_\lambda^b \otimes \mathcal{O}_F)_{\text{int}, !}(\pi_f)_\epsilon$  and define the periods with respect to this module. Again these periods will integral multiples of the periods  $\Omega^\bullet(\pi_f, \epsilon)$ .

In the following Chapter V we will discuss the rationality results (Manin and Shimura) which relate these periods to special values of the  $L$ -function (see section 5.6). But we also want to discuss this method not only for cuspidal classes but also for the Eisenstein cohomology classes, therefore we close this Chapter with a brief account of these Eisenstein classes.

### 4.1.9 The Eisenstein cohomology class

In section 3.3.3 we claimed the existence of the specific cohomology class  $\text{Eis}_n \in H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n)$ . In this section we give a construction of this class on transcendental level, i.e. we construct a cohomology class  $\text{Eis}(\omega_n) \in H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathbb{C})$  whose restriction to the boundary  $H^1(\partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}}_n \otimes \mathbb{C})$  is a given class  $\omega_n$ . For the general theory of Eisenstein cohomology we refer to Chapter 9.

We start from our highest weight module  $\mathcal{M}_\lambda$  and we observe that by definition we have an inclusion

$$i_0 : \mathfrak{J}_B^G \lambda_{\mathbb{R}}^{w_0} \hookrightarrow \mathcal{C}_\infty(\Gamma_\infty^+ \backslash G^+(\mathbb{R}))$$

where

$$\Gamma_\infty^+ = \left\{ \begin{pmatrix} t_1 & m \\ 0 & t_1 \end{pmatrix} \mid m \in \mathbb{Z} ; t_1 = \pm 1 \right\}.$$

Therefore we get an isomorphism

$$H^1(\mathfrak{g}, K_\infty, \mathfrak{J}_B^G \lambda_{\mathbb{R}}^{w_0} \otimes \mathcal{M}_\lambda \otimes \mathbb{C}) \xrightarrow{\sim} H^1(\Gamma_\infty^+ \backslash \mathcal{M}_\lambda \otimes \mathbb{C}) = H^1(\partial(\Gamma \backslash \mathbb{H}), \mathcal{M}_\lambda \otimes \mathbb{C})$$

The inclusion  $i_0$  sends the module  $\mathfrak{J}_B^G \lambda_{\mathbb{R}}^{w_0}$  into a space of functions which are  $\Gamma_\infty^+$  invariant under left translations. Therefore we get a homomorphism

$$\text{Eis} : \mathfrak{J}_B^G \lambda_{\mathbb{R}}^{w_0} \rightarrow \mathcal{C}_\infty(\Gamma \backslash \text{Sl}_2(\mathbb{R}))$$

if we make it invariant by summation, i.e. for  $f \in \mathfrak{J}_B^G \lambda_{\mathbb{R}}^{w_0}$  we define ESeries

$$\text{Eis}(f)(x) = \sum_{\Gamma_\infty^\pm \backslash \text{Sl}_2(\mathbb{Z})} f(\gamma x) \tag{4.165}$$

Of course we have to discuss the convergence of this infinite series. We could quote H. Jacquet: "Let us speak about convergence later", but here is a short interlude discussing this issue.

*Interlude:* Here is the point: We twist our module, for any complex number  $z \in \mathbb{C}$  we consider the induced module

$$\mathfrak{J}_B^G \lambda_{\mathbb{R}}^{w_0} |\rho|^z \subset \mathcal{C}_\infty(\Gamma_\infty^+ \backslash \text{Sl}_2(\mathbb{R}))$$

and again we write down the Eisenstein series. Now it is an elementary exercise to show that the map

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto (c, d)$$

provides a bijection

$$\Gamma_\infty^+ \backslash \text{Sl}_2(\mathbb{Z}) \xrightarrow{\sim} \{(c, d) \in \mathbb{Z} \times \mathbb{Z} \mid (c, d) \text{ coprime}\} / \{\pm 1\} = \mathbb{P}^1(\mathbb{Q}).$$

An element  $x \in \text{Sl}_2(\mathbb{R})$  can be written as  $x = \begin{pmatrix} t & u \\ 0 & t^{-1} \end{pmatrix} k$  with  $k \in K_\infty$ . Then

for  $f \in \mathfrak{I}_B^G \lambda_{\mathbb{R}}^{w_0} |\rho|^z$

$$\begin{aligned} f(\gamma x, z) &= \\ f\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} t & u \\ 0 & t^{-1} \end{pmatrix} k, z\right) &= \\ f\left(\begin{pmatrix} (c^2 t^2 + (cv + dt^{-1})^2)^{-1/2} & * \\ 0 & (c^2 t^2 + (cv + dt^{-1})^2)^{1/2} \end{pmatrix}\right) f(k(\gamma g)k, z) &= \\ (c^2 t^2 + (cv + dt^{-1})^2)^{-n-2-z} f(k(\gamma g)k). \end{aligned}$$

Since  $|f(k(\gamma)k)|$  is bounded the series

$$\text{Eis}(f, z)(x) = \sum_{\Gamma_{\infty}^+ \backslash \text{Sl}_2(\mathbb{Z})} f(\gamma x, z)$$

is converging if  $\Re(z) \gg 0$  and then it is also holomorphic in  $z$ . Selberg and others showed that it can be extended to a meromorphic function in the entire complex plane, it is now a special case of a theorem of Langlands [54]. If now the function  $x \mapsto \text{Eis}(f, z)(x)$  is holomorphic at  $z = 0$  then we do not care about convergence and we simply define

$$\text{Eis}(f)(x) = \sum_{\Gamma_{\infty}^+ \backslash \text{Sl}_2(\mathbb{Z})} f(\gamma x) = \text{Eis}(f, 0)(x).$$

In our special case it is easy to see that the series is convergent at  $z = 0$  provided we have  $n > 0$  and this is the only case where we will apply this construction.  
*End interlude*

This provides a homomorphism

$$\text{Eis}^{\bullet} : H^1(\mathfrak{g}, K_{\infty}, \mathfrak{I}_B^G \lambda_{\mathbb{R}}^{w_0} \otimes \mathcal{M}_{\lambda} \otimes \mathbb{C}) \rightarrow H^1(\Gamma \backslash \mathbb{H}, \mathcal{M}_{\lambda} \otimes \mathbb{C}) \quad (4.166)$$

In ??? we wrote down a distinguished generator  $\omega_n \in H^1(\mathfrak{g}, K_{\infty}, \mathfrak{I}_B^G \lambda_{\mathbb{R}}^{w_0} \otimes \mathcal{M}_{\lambda} \otimes \mathbb{C})$  and we define

$$\text{Eis}_n = \text{Eis}(\omega_n)$$

**Proposition 4.1.8.** *The restriction of  $\text{Eis}_n$  to  $H^1(\partial(\Gamma \backslash \mathbb{H}); \mathcal{M}_{\lambda} \otimes \mathbb{C})$  is the class  $[Y^n]$*

## Chapter 5

# Application to Number Theory

### 5.1 Modular symbols, $L$ - values and denominators of Eisenstein classes.

In this chapter we want to restrict to the case  $\Gamma = \mathrm{Sl}_2(\mathbb{Z})$  or  $\Gamma = \mathrm{Sl}_2(\mathcal{O})$  where  $\mathcal{O}$  is the ring of integers of an imaginary quadratic extension. We refer to section 4.1.1 then this means that  $\Gamma = \mathcal{G}(\mathbb{Z})$ . Our coefficient systems will be obtained from the modules  $\mathcal{M}_\lambda$ . We assume that we have  $d = 0$  and hence  $n \equiv 0 \pmod{2}$  in case A), and  $d_1 = d_2 = 0$ ,  $n_1 = n_1$  in case B). This has the effect that  $\lambda^\vee = \lambda$ .

We want to study the pairing

$$H_c^1(\Gamma \backslash X, \tilde{\mathcal{M}}_\lambda^b) \times H_1(\Gamma \backslash X, \partial(\Gamma \backslash X), \underline{\mathcal{M}}_\lambda) \rightarrow \mathbb{Z}, \quad (5.1)$$

#### 5.1.1 Modular symbols attached to a torus in $\mathrm{Gl}_2$ .

In a first step we construct (relative) cycles in  $C_1(\Gamma \backslash X, \underline{\mathcal{M}}_\lambda)$ ,  $C_1(\Gamma \backslash X, \partial(\Gamma \backslash X), \underline{\mathcal{M}}_\lambda)$ . Our starting point is a maximal torus  $T/\mathbb{Q} \subset G/\mathbb{Q}$  and we assume that it is split over a real quadratic extension  $F/\mathbb{Q}$ . Then the group of real points

$$T(\mathbb{R}) = \mathbb{R}^\times \times \mathbb{R}^\times$$

act on  $\mathbb{H}$  and  $\bar{\mathbb{H}}$  and it has two fixed points  $r, s \in \mathbb{P}^1(F)$ . There is a unique geodesic (half) circle  $\bar{C}_{r,s} \subset \bar{\mathbb{H}}$  joining these two points. Then  $T(\mathbb{R})$  acts transitively on  $C_{r,s} = \bar{C}_{r,s} \setminus \{r, s\}$ . We have two cases:

a) The torus  $T/\mathbb{Q}$  is split. Then the two points  $r, s \in \mathbb{P}^1(\mathbb{Q})$ . Here for instance we can take  $r = 0, s = \infty$ , then the geodesic circle is the line  $\{iy, y > 0\}$  and the torus is the standard diagonal split torus.

b) Here  $\{r, s\} \in \mathbb{P}^1(F) \setminus \mathbb{P}^1(\mathbb{Q})$ , then  $r, s$  are Galois-conjugates of each other. Our torus  $T/\mathbb{Q}$  is given by a suitable embedding

$$j : R_{F/\mathbb{Q}}(\mathbb{G}_m/F) = T \hookrightarrow \mathrm{Gl}_2/\mathbb{Q}.$$

In case a) we can choose any reasonable homeomorphism  $[0, 1] \xrightarrow{\sim} [0, \infty]$  - for instance  $x \mapsto x/(1-x)$  - and then we get a one chain

$$\sigma : [0, 1] \xrightarrow{\sim} \bar{C}_{r,s} = \mathbb{R}_{>0} \cup \{0\} \cup \{\infty\}, \sigma(0) = r, \sigma(1) = s \in \partial(\bar{\mathbb{H}}),$$

and for any  $m \in \mathcal{M}$  we can consider the image of  $\sigma \otimes m \in C_1(\bar{\mathbb{H}}) \otimes \mathcal{M}$  in  $C_1(\Gamma \backslash \bar{\mathbb{H}}, \partial(\Gamma \backslash \bar{\mathbb{H}}), \underline{\mathcal{M}})$ . By definition this is a cycle and hence we get a homology class

$$[\bar{C}_{r,s} \otimes m] \in H_1(\Gamma \backslash \bar{\mathbb{H}}, \partial(\Gamma \backslash \bar{\mathbb{H}}), \underline{\mathcal{M}}_\lambda), \quad (5.2)$$

it is easy to see that it does not depend on the choice of  $\sigma$ .

In case b) we have  $T(\mathbb{Q}) \xrightarrow{\sim} F^\times$ . Then the group  $T(\mathbb{Q}) \cap \Gamma$  is a subgroup of finite index in the group of units  $\mathcal{O}_F^\times = \{\epsilon_0\} \times \{\pm 1\}$ , where  $\epsilon_0$  is a fundamental unit. Hence

$$\Gamma_T = T(\mathbb{Q}) \cap \Gamma = \{\epsilon_T\} \times \mu_T \quad (5.3)$$

where  $\epsilon_T$  is an element of infinite order and  $\mu_T$  is trivial or  $\{\pm 1\}$ . This element  $\epsilon_T$  induces a translation on  $C_{r,s}$ . The quotient  $C_{r,s}/\Gamma_T$  is a circle. If we pick any point  $x \in C_{r,s}$  then  $[x, \epsilon_T x] \subset C_{r,s}$  is an interval and as above we can find a  $\sigma : [0, 1] \xrightarrow{\sim} [x, \epsilon_T x], \sigma(0) = x, \sigma(1) = \epsilon_T x$ . As before we can consider the 1-chain  $\sigma \otimes m \in C_1(\mathbb{H}) \otimes \mathcal{M}$ . Its boundary boundary is the zero chain  $\{x\} \otimes m - \{\epsilon_T x\} \otimes m$ . If we look at the images in  $C_\bullet(\Gamma \backslash \mathbb{H}, \underline{\mathcal{M}}_\lambda)$  then

$$\partial_1(\sigma \otimes m) = \sigma(0) \otimes (m - \epsilon_T m) = r \otimes (m - \epsilon_T m) \quad (5.4)$$

Hence we see that  $\sigma \otimes m$  is a 1-cycle if and only if  $m = \epsilon_T m$  and hence  $m \in \mathcal{M}^T$ . Hence we have constructed homology classes

$$[C_{r,s} \otimes m] \in H_1(\Gamma \backslash \mathbb{H}, \underline{\mathcal{M}}_\lambda) \text{ for all } m \in \mathcal{M}_\lambda^{<\epsilon_T>} = \mathcal{M}_\lambda^T \quad (5.5)$$

PDual

### 5.1.2 Evaluation of cuspidal classes on modular symbols

The following issue will also be discussed in greater generality and more systematically in chapter 8.2.1.

We start from a highest weight  $\lambda = n\gamma$  for simplicity we assume  $n$  to be even and  $d = 0$ . Then  $\lambda = \lambda^\vee$ , we consider the two modules  $\mathcal{M}_\lambda$  and  $\mathcal{M}_\lambda^b$ . Then we have the pairings

$$\begin{aligned} H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_\lambda^b) \times H_1(\Gamma \backslash \mathbb{H}, \underline{\mathcal{M}}_\lambda) &\rightarrow \mathbb{Z} \\ H_c^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_\lambda^b) \times H_1(\Gamma \backslash \mathbb{H}, \partial(\Gamma \backslash \mathbb{H}), \underline{\mathcal{M}}_\lambda) &\rightarrow \mathbb{Z} \end{aligned} \quad (5.6)$$

These two pairings are non degenerate if we invert 6 and divide by the torsion on both sides. (See [book]).

We have the surjective homomorphism  $H_c^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_\lambda^b) \rightarrow H_1^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_\lambda^b)$  and over a suitably large finite extension  $F/\mathbb{Q}$  we have the isotypical decomposition

$$H_1^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_\lambda^b \otimes F) = \bigoplus_{\pi_f} H_1^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_\lambda^b \otimes F)(\pi_f) \quad (5.7)$$

where the  $\pi_f$  are absolutely irreducible. (See Theorem 3.2.1, of course here it does not matter whether we work with  $\mathcal{M}_\lambda$  or  $\mathcal{M}_\lambda^b$ ). We choose an embedding  $\iota : K \hookrightarrow \mathbb{C}$ , in section 4.1.8 we constructed the isomorphism

$$\mathcal{F}_1^1(\omega_\epsilon^\dagger) : \mathcal{W}(\pi_f) \otimes_{F, \iota} \mathbb{C} \rightarrow H_{\epsilon, 1}^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_\lambda^b \otimes F)(\pi_f) \quad (5.8)$$

The space  $\mathcal{W}(\pi_f)$  is a very explicit space. Since we want to stick to the case  $K_f = K_f^{(0)}$  it is of dimension one and is generated by the element

$$h_{\pi_f}^{\dagger, 0} = \prod_p h_p^{\dagger, 0} \in \prod_p \mathcal{W}(\pi_p) \text{ where } h_p^{\dagger, 0}(\epsilon) = 1 \quad (5.9)$$

Now we want to compute the value

$$\langle \mathcal{F}_1^1(\omega_\epsilon^\dagger \times h_{\pi_f}^{\dagger, 0}), \bar{C}_{r,s} \otimes m \rangle. \quad (5.10)$$

This expression is not completely unproblematic. The argument  $C_{r,s}$  on the left lives in the relative homology group, hence the argument on the right should be in  $H_c^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_\lambda^b \otimes \mathbb{C})$ . Of course we can lift the class  $\mathcal{F}_1^1(\omega_\epsilon^\dagger)(h_{\pi_f}^{\dagger, 0})$  to a class

$$\mathcal{F}_1^1(\widetilde{\omega_\epsilon^\dagger \times h_{\pi_f}^{\dagger, 0}}) \in H_c^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_\lambda^b \otimes \mathbb{C}).$$

Then

$$\langle \mathcal{F}_1^1(\widetilde{\omega_\epsilon^\dagger \times h_{\pi_f}^{\dagger, 0}}), C_{r,s} \otimes m \rangle$$

makes sense, but the result may depend on the lift. We have paircusp

**Proposition 5.1.1.** *If  $\partial(C_{r,s} \otimes m)$  gives the trivial class in  $H_0(\partial(\Gamma \backslash \bar{\mathbb{H}}), \tilde{\mathcal{M}}_\lambda \otimes \mathbb{C})$  then  $\langle \mathcal{F}_1^1(\widetilde{\omega_\epsilon^\dagger})(h_{\pi_f}^{\dagger, 0}), C_{r,s} \otimes m \rangle$  does not depend on the lift, i.e. the value  $\langle \mathcal{F}_1^1(\omega_\epsilon^\dagger)(h_{\pi_f}^{\dagger, 0}), C_{r,s} \otimes m \rangle$  is well defined.*

*Proof.* This is rather clear, we refer to the systematic discussion in 6.3.11.  $\square$

Now we compute the value of the pairing. We realized the relative homology class by a  $\mathcal{M}_\lambda$  valued 1-chain  $\sigma \otimes m$ , the cohomology class  $\mathcal{F}_1^1(\omega_\epsilon^\dagger)(h_{\pi_f}^{\dagger, 0})$  is represented by  $\mathcal{F}_1^1(\widetilde{\omega_\epsilon^\dagger \times h_{\pi_f}^{\dagger, 0}})$ . (See 4.81, 8.3). We consider the pullback  $\sigma^*(\mathcal{F}_1^1(\widetilde{\omega_\epsilon^\dagger \times h_{\pi_f}^{\dagger, 0}}))$ , since  $\mathcal{F}_1^1(\omega_\epsilon^\dagger \times h_{\pi_f}^{\dagger, 0})$  is rapidly decaying if  $x \rightarrow 0$  or  $x \rightarrow 1$  this gives us a 1-form with values in  $\mathcal{M}_\lambda \otimes \mathbb{C}$  on the closed interval  $[0, 1]$ .

We claim - under the assumption  $[\partial(C_{r,s} \otimes m)] = 0$  - that

$$\langle \mathcal{F}_1^1(\omega_\epsilon^\dagger)(h_{\pi_f}^{\dagger, 0}), C_{r,s} \otimes m \rangle = \int_0^1 \langle \sigma^*(\mathcal{F}_1^1(\widetilde{\omega_\epsilon^\dagger \times h_{\pi_f}^{\dagger, 0}})), m \rangle. \quad (5.11)$$

**Needs a little argument**

We consider the special case that  $T$  is the standard split diagonal torus, this means that  $\{r, s\} = \{0, \infty\}$ . We use the above identification  $[0, 1] = [0, \infty]$  and our 1-chain is given by the map

$$\sigma : [0, \infty] \rightarrow \bar{\mathbb{H}} : t \mapsto \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} i = ti \in \bar{\mathbb{H}}, \quad (5.12)$$

especially  $\sigma(0) = 0$  and  $\sigma(\infty) = i\infty$ . The group  $T(\mathbb{R})$  acts transitively on the open part  $C_{0, i\infty}$ . This action can be used to trivialize the tangent bundle. The tangent space at  $i \in \mathbb{H}$  is identified to the subspace  $\mathfrak{p} \subset \mathfrak{g}$  (see 4.1.8) and  $\frac{H}{2}$  is a generator of the tangent space of  $C_{0, i\infty}$  at one. Using the translations by  $T(\mathbb{R})$  we get an invariant vector field on  $C_{0, i\infty}$ . If we identify  $C_{0, i\infty} = \mathbb{R}_{>0}$ , an easy calculation shows that this vector field is  $t \frac{d}{dt} = D^*$ .

Now an easy calculation (See 8.3) shows that

$$\mathcal{F}^1(\widetilde{\omega_\epsilon^\dagger \times h_{\pi_f}^{\dagger, 0}})(D^*)\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} e_f\right) = \rho_\lambda\left(\begin{pmatrix} t^{-1} & 0 \\ 0 & 1 \end{pmatrix}\right) \mathcal{F}^1(\omega_\epsilon^\dagger \times h_{\pi_f}^{\dagger, 0})\left(\frac{H}{2}\right)\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}, e_f\right)$$

and our integral in the formula above becomes

$$\int_0^\infty \langle \rho_\lambda\left(\begin{pmatrix} t^{-1} & 0 \\ 0 & 1 \end{pmatrix}\right) \mathcal{F}^1(\omega_\epsilon^\dagger\left(\frac{H}{2}\right) \times h_{\pi_f}^{\dagger, 0})\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}, e_f\right), m \rangle \frac{dt}{t}. \quad (5.13)$$

Our formulas in 4.1.8 give

$$\omega_\pm^\dagger\left(\frac{H}{2}\right) = \frac{1}{8}(\tilde{\psi}_{n+2} \otimes (X - Y \otimes i)^n \pm \tilde{\psi}_{-n-2} \otimes (X + Y \otimes i)^n) \quad (5.14)$$

this is an element in  $\tilde{\mathcal{D}}_\lambda^\pm \otimes \mathcal{M}_\lambda$ . We apply  $\mathcal{F}^1$  to  $\omega_\pm^\dagger\left(\frac{H}{2}\right) \times h_{\pi_f}^{\dagger, 0}$  and evaluate at  $\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}, e_f\right)$ . Applying  $\mathcal{F}^1$  means that we have to sum over  $a \in \mathbb{Q}^\times$  but since  $h_{\pi_f}^{\dagger, 0}$  is the Whittaker function attached to the unramified spherical function only the terms with  $a \in \mathbb{Z}$  can be non zero. Hence get

$$\begin{aligned} & \mathcal{F}^1(\omega_\pm^\dagger\left(\frac{H}{2}\right) \times h_{\pi_f}^{\dagger, 0})\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}, e_f\right) = \\ & \frac{1}{8} \sum_{a \in \mathbb{Z}; a \neq 0} (\tilde{\psi}_{n+2}\left(\begin{pmatrix} at & 0 \\ 0 & 1 \end{pmatrix}\right) \otimes (X - Y \otimes i)^n \pm \tilde{\psi}_{-n-2}\left(\begin{pmatrix} at & 0 \\ 0 & 1 \end{pmatrix}\right) \otimes (X + Y \otimes i)^n) h_{\pi_f}^{\dagger, 0}(a) \end{aligned} \quad (5.15)$$

We have seen that  $\tilde{\psi}_{n+2}\left(\begin{pmatrix} at & 0 \\ 0 & 1 \end{pmatrix}\right) = 0$  if  $at < 0$  and  $\tilde{\psi}_{n+2}\left(\begin{pmatrix} -at & 0 \\ 0 & 1 \end{pmatrix}\right) = \tilde{\psi}_{-n-2}\left(\begin{pmatrix} at & 0 \\ 0 & 1 \end{pmatrix}\right)$  and therefore our Fourier expansion becomes

$$\frac{1}{8} \sum_{a=1}^\infty \tilde{\psi}_{n+2}\left(\begin{pmatrix} at & 0 \\ 0 & 1 \end{pmatrix}\right) \otimes ((X - Y \otimes i)^n \pm i^n (X + Y \otimes i)^n) h_{\pi_f}^{\dagger, 0}(a) \quad (5.16)$$

We have

$$\begin{aligned} \rho_\lambda\left(\begin{pmatrix} t^{-1} & 0 \\ 0 & 1 \end{pmatrix}\right)((X - Y \otimes i)^n \pm i^n(X + Y \otimes i)^n) = \\ \sum_{\nu=0}^n \binom{n}{\nu} t^{\frac{n}{2}-\nu} X^\nu Y^{n-\nu} (i^{n+\nu} \pm i^{-\nu}), \end{aligned} \quad (5.17)$$

we remember that  $n$  is even, then the last factor is equal to  $i^{-\nu}((-1)^{\frac{n}{2}+\nu} \pm 1)$ . and this is  $i^{-\nu}$  times 2 or 0 or -2, depending on the choices of signs and the parity of  $\frac{n}{2}$  and  $\nu$ . The elements  $e_\nu = X^\nu Y^{n-\nu}$  form the dual basis to the basis  $\binom{n}{n-\nu} X^{n-\nu} Y^\nu$  of  $\mathcal{M}_\lambda^b$ , this implies: If we choose in our expression above  $m = e_{n-\nu}$  then pairinf

$$\langle \rho_\lambda\left(\begin{pmatrix} t^{-1} & 0 \\ 0 & 1 \end{pmatrix}\right)((X - Y \otimes i)^n \pm i^n(X + Y \otimes i)^n), m \rangle = t^{\frac{n}{2}-\nu} (i^{n-\nu} \pm i^{-\nu}) \quad (5.18)$$

and hence we have to compute

$$\frac{i^{n+\nu} \pm i^{-\nu}}{8} \int_0^\infty \sum_{a=1}^\infty \tilde{\psi}_{n+2}\left(\begin{pmatrix} at & 0 \\ 0 & 1 \end{pmatrix}\right) t^{\frac{n}{2}-\nu} h_{\pi_f}^{\dagger,0}(a) \frac{dt}{t}. \quad (5.19)$$

We remember  $\tilde{\psi}_{n+2}\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right) = t^{\frac{n}{2}+1} e^{-2\pi t}$ , we exchange summation and integration and after some innocent substitutions we get

$$\frac{i^{n+\nu} \pm i^{-\nu}}{8} \int_0^\infty \frac{t^{n-\nu+1}}{(2\pi)^{n-\nu+1}} e^{-t} \frac{dt}{t} \sum_{a=1}^\infty \frac{h_{\pi_f}^{\dagger}(a) a^{\frac{n}{2}}}{a^\nu} \quad (5.20)$$

We refer to the discussion of the  $L$ -function attached to  $\pi_f$  and get

$$\frac{i^{n+\nu} \pm i^{-\nu}}{8} \int_0^\infty \frac{t^{n-\nu+1}}{(2\pi)^{n-\nu+1}} e^{-t} \frac{dt}{t} \sum_{a=1}^\infty \frac{h_{\pi_f}^{\dagger}(a) a^{\frac{n}{2}}}{a^\nu} = \Lambda^{\text{coh}}(\pi, n+1-\nu) \quad (5.21)$$

**Of course some question concerning convergence has to be discussed**

In the case that  $\nu \neq 0, n$  we know that  $\partial(C_{0,\infty} \otimes X^{n-\nu} Y^\nu)$  is a torsion element in  $H^0(\partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}})$  and therefore the value of the integral is also the evaluation of the cohomology class  $\mathcal{F}_1^1(\omega_\epsilon^\dagger)(h_{\pi_f}^{\dagger,0})$  on a integral homology class. we get

$$\langle C_{0,\infty} \otimes X^\nu Y^{n-\nu}, \mathcal{F}_1^1(\omega_\epsilon^\dagger)(h_{\pi_f}^{\dagger,0}) \rangle = \Lambda^{\text{coh}}(\pi, n+1-\nu) \quad (5.22)$$

In section 4.1.8 we defined the periods  $\Omega_{U_\nu}^\epsilon(\pi_f)$ , we then know that

$$\frac{1}{\Omega_{U_\nu}^\epsilon(\pi_f)} \mathcal{F}_1^1(\omega_\epsilon^\dagger)(h_{\pi_f}^{\dagger,0}) \in H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}} \otimes \mathcal{O}_F(U_\nu)) \quad (5.23)$$

and hence we can conclude for  $\nu \neq 0, n$  ratint

$$\frac{1}{\Omega^\epsilon(\pi_f)} \Lambda^{\text{coh}}(\pi, n+1-\nu) \in \mathcal{O}_F \quad (5.24)$$

This argument fails for  $\nu = 0, n$  because  $\partial(C_{0,\infty} \otimes X^n) = \infty \otimes (X^n - Y^n)$  is not a torsion class in  $H_0(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_\lambda)$  (See section 3.2.1). We apply the Manin-Drinfeld principle to show that the rationality statement also holds for  $\nu = 0, n$  but we will get a denominator.

We pick a prime  $p$  then we know that the class  $[\partial(C_{0,\infty} \otimes X^n)]$  is an eigenclass modulo torsion for  $T_p$ , i.e.

$$T_p([\partial(C_{0,\infty} \otimes X^n)]) = (p^{n+1} + 1)[\partial(C_{0,\infty} \otimes X^n)] \quad (5.25)$$

This implies that  $\partial(T_p([\mathbb{C}_{0,\infty} \otimes X^n]) - (p^{n+1} + 1)[(C_{0,\infty} \otimes X^n)])$  is a torsion class, hence we can apply proposition 5.1.1 and get that the value of the pairing is equal to the integral against the modular symbol. If we exploit the adjointness formula for the Hecke operator then get

$$\begin{aligned} & \langle T_p([\mathbb{C}_{0,\infty} \otimes X^n]) - (p^{n+1} + 1)[(C_{0,\infty} \otimes X^n)], \mathcal{F}_1^1(\omega_\epsilon^\dagger \otimes h_{\pi_f}^{\dagger,0}) \rangle \\ &= \int_0^\infty \langle C_{0,\infty} \otimes X^n, \mathcal{F}_1^1(\omega_\epsilon^\dagger \otimes T_p(h_{\pi_f}^{\dagger,0})^\dagger) \rangle - (p^{n+1} + 1) \langle C_{0,\infty} \otimes X^n, \mathcal{F}_1^1(\omega_\epsilon^\dagger \otimes ((h_{\pi_f}^{\dagger,0})^\dagger)) \rangle \frac{dt}{t} \end{aligned} \quad (5.26)$$

We have  $T_p(h_{\pi_f}^{\dagger,0}) = a_p h_{\pi_f}^{\dagger,0}$  where  $a_p \in \mathcal{O}_F$  and hence we get

$$\begin{aligned} & \langle T_p([\mathbb{C}_{0,\infty} \otimes X^n]) - (p^{n+1} + 1)[(C_{0,\infty} \otimes X^n)], \mathcal{F}_1^1(\omega_\epsilon^\dagger \otimes h_{\pi_f}^{\dagger,0}) \rangle \\ &= (a_p - (p^{n+1} + 1)) \Lambda^{\text{coh}}(\pi_f, n + 1) \end{aligned} \quad (5.27)$$

It is again the Manin-Drinfeld principle that tells us that for almost all primes  $p$  the number  $a_p - (p^{n+1} + 1) \neq 0$ . More precisely we know that the greatest common divisor of these numbers is the numerator

$$Z(n) = \text{numerator}(\zeta(-1 - n)) \quad (5.28)$$

This gives us a modified rationality-integrality assertion: For  $\nu = n + 1, \dots, 0$  we have ratintE

$$\frac{1}{\Omega_{U_\nu}^\epsilon(\pi_f)} \Lambda^{\text{coh}}(\pi, \nu) \in \frac{1}{Z(n)} \mathcal{O}_F(U_\nu) \quad (5.29)$$

These rationality results go back to Manin and Shimura, In principle we may say that also the integrality assertion goes back to these authors, but here we have to take into account the fine tuning of the periods. (Deligne conjecture?)

### 5.1.3 Evaluation of Eisenstein classes on capped modular symbols

We have seen that MDEis

$$H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathbb{Q}) = H_!^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n \otimes \mathbb{Q}) \oplus \mathbb{Q}\text{Eis}_n \quad (5.30)$$

where  $\text{Eis}_n$  is defined by the two conditions

$$r(\text{Eis}_n) = [Y^n] \text{ and } T_p(\text{Eis}_n) = (p^{n+1} + 1)\text{Eis}_n, \quad (5.31)$$

for all Hecke operators  $T_p$ , in our special situation it suffices to check the second condition for  $p = 2$ . In (???) we raised the question to determine the denominator of the class  $\text{Eis}_n$ , i.e. we want to determine the smallest integer  $\Delta(n) > 0$  such that  $\Delta(n)\text{Eis}_n$  becomes an integral class.

To achieve this goal we compute the evaluation of  $\text{Eis}_n$  on the first homology group, i.e we compute the value  $\langle c, \text{Eis}_n \rangle$  for  $c \in H_1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n^\vee)$ . We have the exact sequence

$$H_1(\partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}}_n^\vee) \xrightarrow{j} H_1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n^\vee) \rightarrow H_1(\Gamma \backslash \mathbb{H}, \partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}}_n^\vee) \xrightarrow{\delta} H_0(\partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}}_n^\vee) \quad (5.32)$$

It follows from the construction of  $\text{Eis}_n$  that  $\langle c, \text{Eis}_n \rangle \in \mathbb{Z}$  for all the elements the image of  $j$ . Therefore we only have to compute the values  $\langle \tilde{c}_\nu, \text{Eis}_n \rangle$ , where  $\tilde{c}_\mu$  are lifts of a system of generators  $\{c_\mu\}$  of  $\ker(\delta)$ .

In our special case the elements  $C_{0,\infty} \otimes e_\nu^\vee$ , where  $\nu = 0, 1, \dots, n$  form a set of generators of  $H_1(\Gamma \backslash \mathbb{H}, \partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}}_n^\vee)$ . (Diploma thesis Gebertz). We observe:

The boundary of the element  $C_{0,\infty} \otimes e_n^\vee (= \pm C_{0,\infty} \otimes e_0^\vee)$  is an element of infinite order in  $H^0(\partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}}_n^\vee)$ ,

The boundary of an elements  $C_{0,\infty} \otimes e_\nu^\vee$  with  $0 < \nu < n$  are torsion elements in  $H^0(\partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}}_n^\vee)$ , This implies

**Proposition 5.1.2.** *The elements  $C_{0,\infty} \otimes m \in H_1(\Gamma \backslash \mathbb{H}, \partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}}_n^\vee)$  with  $\partial(C_{0,\infty} \otimes m) = 0$  are of the form*

$$c = C_{0,\infty} \otimes \left( \sum_{\nu=1}^{\nu=n-1} a_\nu e_\nu^\vee \right); \quad \text{with } a_\nu \in \mathbb{Z}$$

Now it seems to be tempting to choose for our our generators above the  $C_{0,\infty} \otimes e_\nu^\vee$ , but this is not possible because for  $\delta(C_{0,\infty} \otimes e_\nu^\vee)$  is not necessarily zero, it is only a torsion element. So we see that it is not clear how to find a suitable system of generators.

To overcome this difficulty we use the Hecke operators. If we want to determine the denominator  $\Delta(n)$  we can localize, i.e. for each prime  $p$  we have to determine the highest power  $p^{d(n,p)}$  which divides  $\Delta(n)$ . As usual we write  $d(n,p) = \text{ord}_p(\Delta(n))$ . We replace the ring  $\mathbb{Z}$  by its localization  $\mathbb{Z}_{(p)}$  and replace all our cohomology and homology groups by he localized groups. In other words we have to check we have to find a set of generators  $\{\tilde{c}_\nu\}_\mu \subset H_1(\partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}}_n^\vee \otimes \mathbb{Z}_{(p)})$  and compute the denominator  $\langle \tilde{c}_\nu, \text{Eis}_n \rangle \in \mathbb{Z}_{(p)}$ .

It follows from proposition 3.3.1 that for  $0 < \nu < n$  the torsion element  $\partial(c) = \partial(C_{0,\infty} \otimes (\sum_{\nu=1}^{\nu=n-1} a_\nu e_\nu^\vee))$  is annihilated by a sufficiently high power of the Hecke operator  $T_p^m$  and hence we see that  $T_p^m(c)$  can be lifted to an element  $\widetilde{T_p^m(c)} \in H_1(\partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}}_n^\vee \otimes \mathbb{Z}_{(p)})$ . Now

$$\langle \widetilde{T_p^m(c)}, \text{Eis}_n \rangle = \langle c, T_p^m(\text{Eis}_n) \rangle = (p^{n+1} + 1)^m \langle c, \text{Eis}_n \rangle \quad (5.33)$$

and hence  $\text{ord}_p(\langle \widetilde{T_p^m(c)}, \text{Eis}_n \rangle) = \text{ord}_p(\langle c, \text{Eis}_n \rangle)$ . Hence we get

**Proposition 5.1.3.** *If  $\nu$  runs from 1 to  $n - 1$  and if  $T_p^m(\widetilde{C_{0,\infty} \otimes e_\nu^\vee})$  is any lift of  $T_p^m(e_\nu^\vee)$  then*

$$d(n,p) = -\min(\min_\nu(\text{ord}_p(\langle \widetilde{T_p^m(C_{0,\infty} \otimes e_\nu^\vee)}, \text{Eis}_n \rangle)), 0)$$

*Proof.* This is now obvious.  $\square$

### 5.1.4 The capped modular symbol

Therefore we have to compute  $\langle T_p^m(\widetilde{C_{0,\infty} \otimes e_\nu^\vee}), \text{Eis}_n \rangle$ . At this point some meditation is in order. Our cohomology class  $\text{Eis}_n$  is represented by a closed differential form  $\text{Eis}(\omega_n)$  (See (??)) and this differential form lives on  $\Gamma \backslash \mathbb{H}$  a hence provides a cohomology class in  $\Gamma \backslash \mathbb{H}$ . But we know that the inclusion provides an isomorphism

$$H^1(\Gamma \backslash \mathbb{H}, \widetilde{\mathcal{M}}_n) \xrightarrow{\sim} H^1(\Gamma \backslash \bar{\mathbb{H}}, \widetilde{\mathcal{M}}_n)$$

and since  $T_p^m(\widetilde{C_{0,\infty} \otimes e_\nu^\vee}) \in H_1(\Gamma \backslash \bar{\mathbb{H}}, \widetilde{\mathcal{M}}_n)$  we can evaluate the cohomology class  $\text{Eis}_n$  on the cycle. But we want get this value  $\langle T_p^m(\widetilde{C_{0,\infty} \otimes e_\nu^\vee}), \text{Eis}_n \rangle$  by integration of the differential form against the cycle. This is a little bit problematic because the cycle has non trivial support in  $\partial(\Gamma \backslash \mathbb{H})$ , and on this circle at infinity the differential form is not really defined.

There are certainly several ways out of this dilemma. One possibility is to deform the cycle  $T_p^m(\widetilde{C_{0,\infty} \otimes e_\nu^\vee})$  and "pull" it into the interior  $\Gamma \backslash \mathbb{H}$ . The cycle is the sum of two 1-chains:

$$T_p^m(\widetilde{C_{0,\infty} \otimes e_\nu^\vee}) = C_{0,\infty} \otimes m_\nu + [\infty, T\infty] \otimes P_\nu$$

(recall definition of Borel-Serre construction from earlier chapters) where

$$\partial(C_{0,\infty} \otimes m_\nu) = \infty \otimes (m_\nu - wm_\nu) + \infty \otimes (1 - T)P_\nu = 0$$

Recall that  $C_{0,\infty}$  is the continuous extension of  $t \mapsto \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} i$  from  $\mathbb{R}_{>0}^\times$  to  $\mathbb{H}$  to a map from  $[0, \infty] \rightarrow \bar{\mathbb{H}}$ . We choose a sufficiently large  $t_0 \in \mathbb{R}_{>0}^\times$  and restrict  $C_{0,\infty}$  to  $[t_0^{-1}, t_0]$  we get the one chain  $C_{0,\infty}(t_0) \otimes m_\nu$ . The boundary of this 1-chain is  $\partial(C_{0,\infty}(t_0) \otimes m_\nu) = t_0 \otimes (m_\nu - wm_\nu)$ . Now we can do at this level the same as at infinity we get a 1-cycle

$$C_{0,\infty}(t_0) \otimes m_\nu = C_{0,\infty}(t_0) \otimes m_\nu + [t_0, Tt_0] \otimes P_\nu$$

This 1-cycle clearly defines the same class as  $T_p^m(\widetilde{C_{0,\infty} \otimes e_\nu^\vee})$  and since it is a cycle in  $C_1(\Gamma \backslash \mathbb{H}, \widetilde{\mathcal{M}})$  we get

$$\langle T_p^m(\widetilde{C_{0,\infty} \otimes e_\nu^\vee}), \text{Eis}_n \rangle = \int_{C_{0,\infty}(t_0) \otimes m_\nu + [t_0, Tt_0] \otimes P_\nu} \text{Eis}_n \quad (5.34)$$

The value of this integral does not depend on  $t_0$  and we check easily that for both summands the limit for  $t_0 \rightarrow \infty$  exists. We find that Nenner1

$$\begin{aligned} & \langle T_p^m(\widetilde{C_{0,\infty} \otimes e_\nu^\vee}), \text{Eis}_n \rangle = \\ & \int_0^\infty \langle T_p^m(C_{0,\infty} \otimes e_\nu^\vee), \text{Eis}_n \rangle \frac{dt}{t} + \lim_{t_0 \rightarrow \infty} \int_0^1 \langle [it_0, it_0 + x] \otimes P_\nu, \text{Eis}_n \rangle dx \end{aligned} \quad (5.35)$$

and For the first integral we have

$$\int_0^\infty \langle T_p^m(C_{0,\infty} \otimes e_\nu^\vee), \text{Eis}_n \rangle \frac{dt}{t} = (1+p^{n+1})^m \int_0^\infty \langle C_{0,\infty} \otimes e_\nu^\vee, \text{Eis}_n \rangle \frac{dt}{t}$$

and (handwritten notes page 49)

$$\int_0^\infty \langle C_{0,\infty} \otimes e_\nu^\vee, \text{Eis}_n \rangle \frac{dt}{t} = \frac{\zeta(-\nu)\zeta(\nu-n)}{\zeta(-1-n)} \quad (5.36)$$

remember this holds for  $0 < \nu < n$ .

For the second term we have to observe that it depends on the choice of  $P_\nu$ . We can replace  $P_\nu$  by  $P_\nu + V$  where  $V^T = V$ . (This means of course that  $V = aX^n$ ) Then  $[V] \in H^0(\partial(\Gamma \backslash \mathbb{H}), \tilde{\mathcal{M}}_\lambda)$  and

$$\lim_{t_0 \rightarrow \infty} \int_0^1 \langle [it_0, it_0+x] \otimes (P_\nu + V), \text{Eis}_n \rangle dx = \lim_{t_0 \rightarrow \infty} \int_0^1 \langle [it_0, it_0+x] \otimes P_\nu, \text{Eis}_n \rangle dx + \langle V, \omega_n \rangle .$$

Therefore the second term is only defined up to a number in  $\mathbb{Z}_{(p)}$  but this is ok because we are interested in the  $p$ -denominator in (5.35).

We have to evaluate the expression  $\langle [it_0, it_0+x] \otimes (P_\nu + V), \text{Eis}_n \rangle$ . Using the formula (8.3) we find

$$\langle [it_0, it_0+x] \otimes (P_\nu + V), \text{Eis}_n \rangle = \langle \begin{pmatrix} t_0 & x \\ 0 & 1 \end{pmatrix} P_\nu, \text{Eis}(\omega_n)(E_+) \left( \begin{pmatrix} t_0 & x \\ 0 & 1 \end{pmatrix} \right) \rangle \quad (5.37)$$

We know that for  $t_0 \gg 1$  the Eisenstein series is approximated by its constant term, i.e.

$$\text{Eis}(\omega_n)(E_+) \left( \begin{pmatrix} t_0 & x \\ 0 & 1 \end{pmatrix} \right) = t_0^{-n} Y^n + O(e^{-t_0}) \quad (5.38)$$

On the other hand we can write  $P_\nu(X, Y) = \sum p_\mu^{(\nu)} X^{n-\mu} Y^\mu$  with  $p_{\nu,\mu} \in \mathbb{Z}$ . Then

$$\begin{pmatrix} t_0 & x \\ 0 & 1 \end{pmatrix} P_\nu = t_0^n p_0^{(\nu)} X^n + \dots \quad (5.39)$$

and

$$\langle \begin{pmatrix} t_0 & x \\ 0 & 1 \end{pmatrix} P_\nu, \text{Eis}(\omega_n)(E_+) \left( \begin{pmatrix} t_0 & x \\ 0 & 1 \end{pmatrix} \right) \rangle = p_0^{(\nu)} + O(e^{-t_0}) \quad (5.40)$$

and hence we see that the limit exists and we get

$$\lim_{t_0 \rightarrow \infty} \int_0^1 \langle [it_0, it_0+x] \otimes (P_\nu + V), \text{Eis}_n \rangle dx = p_0^{(\nu)} = P_\nu(1, 0) \quad (5.41)$$

and hence we have the final formula

$$\langle T_p^m(\widetilde{C_{0,\infty}} \otimes e_\nu^\vee), \text{Eis}_n \rangle = \frac{\zeta(-\nu)\zeta(\nu-n)}{\zeta(-1-n)} + P_\nu(1, 0) \pmod{\mathbb{Z}_{(p)}}. \quad (5.42)$$

Therefore we have to compute  $P_\nu(1, 0) \pmod{\mathbb{Z}_{(p)}}$ . Recall that for any  $\nu, \nu \neq 0, n$  we have to choose a very large  $m > 0$  such that the zero chain  $T_p^m(e_\nu^\vee)$  is homologous to

$$T_p^m(e_\nu^\vee) \sim \{\infty\} \otimes L_\nu = \{\infty\} \otimes (1 - T)Q_\nu \quad (5.43)$$

with  $Q_\nu \in \mathcal{M}_n^\vee$ . Then we find  $P_\nu = Q_\nu \pm Q_{n+1-\nu}$ .

Hence we have to compute  $T_p^m(e_\nu^\vee)$ . A straightforward but lengthy computation yields

$$Q_\nu(1, 0) = \begin{cases} 0 & \text{if } (p-1) \nmid \nu+1 \\ \frac{1}{p^{\frac{\nu+1}{p-1}}} & \text{else} \end{cases} \quad (5.44)$$

Now we are ready to compute  $d(n, p)$ , it is the maximum over all  $\nu$  denomest

$$- \text{ord}_p\left(\frac{\zeta(-\nu)\zeta(\nu-n)}{\zeta(-1-n)} - (Q_\nu(1, 0) + Q_{n-\nu}(1, 0))\right) \pmod{\mathbb{Z}_{(p)}}. \quad (5.45)$$

We have to distinguish cases

I) We have  $(p-1) \nmid \nu+1$  and  $(p-1) \nmid n+1-\nu$ . In this case  $Q_\nu(1, 0) = Q_{n+1-\nu}(1, 0) = 0$  and

$$- \text{ord}_p\left(\frac{\zeta(-\nu)\zeta(\nu-n)}{\zeta(-1-n)}\right) = - \text{ord}_p((\zeta(-\nu)\zeta(\nu-n)) + \text{ord}_p(\zeta(-1-n))) \quad (5.46)$$

II) The number  $p-1$  divides exactly on of the numbers  $\nu+1$  or  $n+1-\nu$ . In this case let us assume that it divides  $\nu+1$  and let us write  $\nu+1 = p^{\alpha-1}\nu_0$ , with  $p^{\alpha-1} \mid \nu+1$ . Then the  $p$ -denominator of  $\zeta(-\nu)$  is  $p^\alpha$ . Then  $\nu-n-1 \equiv -n-1 \pmod{(p-1)p^{\alpha-1}}$  and hence it follows from the Kummer congruences that we can write

$$\zeta(\nu-n) = \zeta(-n-1) + p^\alpha Z(\nu, n); \text{ where } Z(\nu, n) \in \mathbb{Z}_{(p)} \quad (5.47)$$

and then

$$\frac{\zeta(-\nu)\zeta(\nu-n)}{\zeta(-1-n)} = \zeta(-\nu)\left(1 + p^\alpha \frac{Z(\nu, n)}{\zeta(-1-n)}\right) \quad (5.48)$$

The theorem of -von Staudt-Clausen tells us that

$$\zeta(-\nu) = \frac{-1}{p^{\frac{\nu}{p-1}}} + v \text{ with } v \in \mathbb{Z}_{(p)} \quad (5.49)$$

and hence the left hand side in the above equation becomes

$$\frac{-1}{p^{\frac{\nu}{p-1}}} + v - \frac{-p^\alpha}{p^{\frac{\nu}{p-1}}} \frac{Z(\nu, n)}{\zeta(-1-n)} + vp^\alpha \frac{Z(\nu, n)}{\zeta(-1-n)} \quad (5.50)$$

We have to subtract  $(Q_\nu(1, 0) + Q_{n-\nu}(1, 0))$  from this expression. Then  $Q_\nu(1, 0)$  cancels against the first term in the above expression, and  $Q_{n-\nu}(1, 0) \in \mathbb{Z}_{(p)}$ . Hence we see that in equation (5.45) we have to compute

$$- \text{ord}_p\left(-\frac{-p^\alpha}{p^{\frac{\nu}{p-1}}} \frac{Z(\nu, n)}{\zeta(-1-n)} + vp^\alpha \frac{Z(\nu, n)}{\zeta(-1-n)}\right) \quad (5.51)$$

By definition we have  $\alpha > 0$  and by definition the factor in front of the first term is a unit we see that for this  $\nu$  the expression in (5.45) is

$$- \operatorname{ord}_p(\zeta(-1-n)) + \operatorname{ord}_p(Z(\nu, n)) = - \operatorname{ord}_p(\zeta(-1-n)) + \operatorname{ord}_p(\zeta(\nu-n))$$

III) We have  $p-1|\nu+1$  and  $p-1|n+1-\nu$ . In this case an elementary computation shows that expression in (5.45) is  $\mathbb{Z}_{(p)}$ , i.e. it is a  $p$ -integer. To see this we write  $\nu+1 = (p-1)xp^{a-1}$ ,  $n+1-\nu = (p-1)yp^{b-1}$  with  $a > 0, b > 0$  and  $x, y$  prime to  $p$ . We assume  $a \leq b$  and compute

$$\frac{\zeta(1-(p-1)xp^{a-1})\zeta(1-(p-1)yp^{b-1})}{\zeta(1-(p-1)p^{a-1}(x+yp^{b-a}))} \pmod{\mathbb{Z}_{(p)}} \quad (5.52)$$

For a value  $\zeta(1-m)$  with  $p-1|m$  we write  $m = (p-1)xp^{k-1}$  with  $(x, p) = 1$ . We apply again the von Staudt-Clausen theorem

$$\zeta(1-m) = \zeta(1-(p-1)xp^{k-1}) = -\frac{1}{xp^k} + Z(x) \text{ where } Z(x) \in \mathbb{Z}_{(p)}$$

In our case this gives -let us assume  $a < b$  - for our expression above

$$\frac{-\frac{1}{(xp^a)} + Z(x))(-\frac{1}{(yp^b)} + Z(y))}{-\frac{1}{(x+yp^{b-a})p^a} + Z(x+yp^{b-a})} = -\frac{(x+yp^{b-a})(\frac{1}{x} + p^a Z(x))(\frac{1}{yp^b} + Z(y))}{1 + p^a(x+yp^{b-a})Z(x+yp^{b-a}y)} \quad (5.53)$$

The denominator is a unit, we need to know it modulo  $p^b$ , the numerator is a sum of eight terms we can forget all the terms in  $\mathbb{Z}_{(p)}$ . Then the above expression simplifies

$$\frac{\frac{1}{yp^b} + \frac{1}{xp^a} + \frac{p^{a-b}xZ(x)}{y}}{1 + p^axZ(x+yp^{b-a})} \quad (5.54)$$

We want this to be equal to  $\frac{1}{yp^b} + \frac{1}{xp^a}$ . Hence we have to verify the equality

$$\frac{1}{yp^b} + \frac{1}{xp^a} + \frac{p^{a-b}xZ(x)}{y} = \left(\frac{1}{yp^b} + \frac{1}{xp^a}\right)(1 + p^axZ(x+yp^{b-a})) \quad (5.55)$$

and this comes down to

$$p^{a-b}\frac{xZ(x)}{y} \equiv p^{a-b}\frac{xZ(x+yp^{b-a})}{y} \pmod{\mathbb{Z}_{(p)}} \quad (5.56)$$

and this means

$$Z(x) \equiv Z(x+yp^{b-a}) \pmod{p^{b-a}}$$

and this congruence is easy to verify.

Basically the same argument works if  $a = b$ . Then it can happen that  $x+y \equiv 0 \pmod{p}$ . Then we have to write  $x+y = p^cz$ . Then (5.53) changes into

$$\frac{(-\frac{1}{xp^a} + Z(x))(-\frac{1}{yp^a} + Z(y))}{-\frac{1}{zp^{a+c}} + Z(z)} = -\frac{zp^c(\frac{1}{x} + p^a Z(x))(\frac{1}{yp^a} + Z(y))}{1 + p^{a+c}zZ(z)}. \quad (5.57)$$

We ignore the denominator then the only non integral term is

$$(x+y) \frac{1}{x} \frac{1}{yp^a} = \frac{1}{xp^a} + \frac{1}{yp^a}$$

- This is now essentially the proof of (??), i.e.

**Theorem 5.1.1.** *If  $\Gamma = \text{Sl}_2(\mathbb{Z})$  then the denominator of the Eisenstein class in  $H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_n)$  is the numerator of  $\zeta(-1-n)$ .*

### The Deligne-Eichler-Shimura theorem

In this section the material is not presented in a satisfactory form. One reason is that at this point we should start using the language of adèles, but there are also other drawbacks. So in a final version of these notes this section probably be removed.

#### *Begin of probably removed section*

In this section I try to explain very briefly some results which are specific for  $\text{Gl}_2$  and a few other low dimensional algebraic groups. These results concern representations of the Galois group  $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$  which can be attached to irreducible constituents  $\Pi_f$  in the cohomology. These results are very deep and reaching a better understanding and more general versions of these results is a fundamental task of the subject treated in these notes. The first cases have been tackled by Eichler and Shimura, then Ihara made some contributions and finally Deligne proved a general result for  $\text{Gl}_2/\mathbb{Q}$ .

We start from the group  $G = \text{Gl}_2/\mathbb{Q}$ , this is now only a reductive group and its centre is isomorphic to  $\mathbb{G}_m/\mathbb{Q}$ . Its group of real points is  $\text{Gl}_2(\mathbb{R})$  and the centre  $\mathbb{G}_m(\mathbb{R})$  considered as a topological group has two components, the connected component of the identity is  $\mathbb{G}_m(\mathbb{R})^{(0)} = \mathbb{R}_{>0}^\times$ . Now we enlarge the maximal compact connected subgroup  $SO(2) \subset \text{Gl}_2(\mathbb{R})$  to the group  $K_\infty = SO(2) \cdot \mathbb{G}_m(\mathbb{R})^{(0)}$ . The resulting symmetric space  $X = \text{Gl}_2(\mathbb{R})/K_\infty$  is now a union of an upper and a lower half plane: We write  $X = \mathbb{H}_+ \cup \mathbb{H}_-$ .

We choose a positive integer  $N > 2$  and consider the congruence subgroup  $\Gamma(N) \subset \text{Gl}_2(\mathbb{Q})$ . We modify our symmetric space: This modification may look a little bit artificial at this point, it will be justified in the next chapter and is in fact very natural. (At this point I want to avoid to use the language of adèles.)

We replace the symmetric space by

$$X = (\mathbb{H}_+ \cup \mathbb{H}_-) \times \text{Gl}_2(\mathbb{Z}/N\mathbb{Z}).$$

On this space we have an action of  $\Gamma = \text{Gl}_2(\mathbb{Z})$ , on the second factor it acts via the homomorphism  $\text{Gl}_2(\mathbb{Z}) \rightarrow \text{Gl}_2(\mathbb{Z}/N\mathbb{Z})$  by translations from the left. Again we look at the quotient of this space by the action of  $\text{Gl}_2(\mathbb{Z})$ . This quotient space will have several connected components. The group  $\text{Gl}_2(\mathbb{Z})$  contains the group  $\text{Sl}_2(\mathbb{Z})$  as a subgroup of index two, because the determinant of an element is  $\pm 1$ . The element  $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$  interchanges the upper and the lower half plane and hence we see

$$\text{Gl}_2(\mathbb{Z}) \backslash X = \text{Gl}_2(\mathbb{Z}) \backslash ((\mathbb{H}_+ \cup \mathbb{H}_-) \times \text{Gl}_2(\mathbb{Z}/N\mathbb{Z})) = \text{Sl}_2(\mathbb{Z}) \backslash (\mathbb{H}_+ \times \text{Gl}_2(\mathbb{Z}/N\mathbb{Z})),$$

the connected components of  $(\mathbb{H}_+ \times \mathrm{Gl}_2(\mathbb{Z}/N\mathbb{Z}))$  are indexed by elements  $g \in \mathrm{Gl}_2(\mathbb{Z}/N\mathbb{Z})$ . The stabilizer of such a component is the full congruence subgroup

$$\Gamma(N) = \left\{ \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, d \equiv 1 \pmod{N}, b, c \equiv 0 \pmod{N} \right\}$$

this group is torsion free because we assumed  $N > 2$ .

The image of the natural homomorphism  $\mathrm{Sl}_2(\mathbb{Z}) \rightarrow \mathrm{Gl}_2(\mathbb{Z}/N\mathbb{Z})$  is the subgroup  $\mathrm{Sl}_2(\mathbb{Z}/N\mathbb{Z})$  (strong approximation), therefore the quotient is by this subgroup is  $(\mathbb{Z}/N\mathbb{Z})^\times$ .

We choose as system of representatives for the determinant the matrices  $t_a = \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix}$ ,  $a \in (\mathbb{Z}/N\mathbb{Z})^\times$ . The stabilizer of then we get an isomorphism

$$S_N = \mathrm{Gl}_2(\mathbb{Z}) \backslash (\mathbb{H} \times \mathrm{Gl}_2(\mathbb{Z}/N\mathbb{Z})) \xrightarrow{\sim} (\Gamma(N) \backslash \mathbb{H}) \times (\mathbb{Z}/N\mathbb{Z})^\times.$$

To any prime  $p$ , which does not divide  $N$  we can again attach Hecke operators. Again we can attach Hecke operators

$$T_{p^r} = T \left( \begin{pmatrix} p^r & 0 \\ 0 & 1 \end{pmatrix}, u \begin{pmatrix} p^r & 0 \\ 0 & 1 \end{pmatrix} \right)$$

to the double cosets and using strong approximation we can prove the recursion formulae.

We consider the cohomology groups  $H_c^\bullet(S_N, \tilde{\mathcal{M}}_n), H^\bullet(S_N, \tilde{\mathcal{M}}_n)$  and define  $H_!^\bullet(S_N, \tilde{\mathcal{M}}_n)$  as before. This is a semi simple module for the cohomology.

The theorem 3 extends to this situation without change. We have a small addendum: If denote by  $Z^{(N, \times)} \in \mathbb{Q}^\times$  the subgroup of those numbers which are units at the primes dividing  $N$ . We have the homomorphism  $r : Z^{(N, \times)} \rightarrow (\mathbb{Z}/N\mathbb{Z})^\times$

*On each absolutely irreducible component  $\Pi_f$  the Hecke operators  $T(z, u_z)$  act by a scalar  $\omega(z) \in \mathcal{O}_L$  and the map  $z \mapsto \omega(z)$  factors over  $r$  and induces a character  $\omega(\Pi_f) : (\mathbb{Z}/N\mathbb{Z})^\times \rightarrow (\mathcal{O}_L)^\times$ . This character is called the central character of  $\Pi_f$ .*

The following things will be explained in greater detail in the class

Now we exploit the fact, that the Riemann surface  $\Gamma(N) \backslash X$  is in fact the space of complex points of the moduli scheme  $M_N \rightarrow \mathrm{Spec}(\mathbb{Z}[1/N])$ . On this moduli scheme we have the universal elliptic curve with  $N$  level structure

$$\begin{array}{c} \mathcal{E} \\ \downarrow \pi \\ M_N \end{array}$$

On  $\mathcal{E}$  we have the constant  $\ell$ -adic sheaf  $\mathbb{Z}_\ell$ . For  $i = 0, 1, 2$  we can consider the  $\ell$ -adic sheaves  $R^i \pi_* (\mathbb{Z}_\ell)$  on  $M_N$ . We have the spectral sequence

$$H^p(M_N \times \bar{\mathbb{Q}}, R^q \pi_* (\mathbb{Z}_\ell)) \Rightarrow H^n(\mathcal{E} \times \bar{\mathbb{Q}}, \mathbb{Z}_\ell).$$

We can take the fibered product of the universal elliptic curve

$$\mathcal{E}^{(n)} = \mathcal{E} \times_{M_N} \mathcal{E} \times \cdots \times_{M_N} \mathcal{E} \xrightarrow{\pi_N} M_N$$

where  $n$  is the number of factors. This gives us a more general spectral sequence

$$H^p(M_N \times \bar{\mathbb{Q}}, R^q \pi_{N,*}(\mathbb{Z}_\ell)) \Rightarrow H^n(\mathcal{E}^{(n)} \times \bar{\mathbb{Q}}, \mathbb{Z}_\ell).$$

The stalk  $R^q \pi_{N,*}(\mathbb{Z}_\ell)_y$  of the sheaf  $R^q \pi_{N,*}(\mathbb{Z}_\ell)$  in a geometric point  $y$  of  $M_N$  is the  $q$ -th cohomology  $H^q(\mathcal{E}_y^{(n)}, \mathbb{Z}_\ell)$  and this can be computed using the Kuenneth formula

$$H^q(\mathcal{E}_y^{(n)}, \mathbb{Z}_\ell) \xrightarrow{\sim} \bigoplus_{a_1, a_2, \dots, a_n} H^{a_1}(\mathcal{E}_y, \mathbb{Z}_\ell) \otimes H^{a_2}(\mathcal{E}_y, \mathbb{Z}_\ell) \cdots \otimes H^{a_n}(\mathcal{E}_y, \mathbb{Z}_\ell),$$

where the  $a_i = 0, 1, 2$  and sum up to  $q$ . We have  $H^0(\mathcal{E}_y, \mathbb{Z}_\ell) = \mathbb{Z}_\ell(0)$ ,  $H^2(\mathcal{E}_y, \mathbb{Z}_\ell) = \mathbb{Z}_\ell(-1)$  and the most interesting factor is  $H^1(\mathcal{E}_y, \mathbb{Z}_\ell)$  which is a free  $\mathbb{Z}_\ell$  module of rank 2.

This tells us that the sheaf decomposes into a direct sum according to the type of Kuenneth summands. We also have an action of the symmetric group  $S_q$  which is obtained from the permutations of the factors in  $\mathcal{E}^{(n)}$  which also permutes the types. We are mainly interested in the case  $q = n$  and then we have the special summand where  $a_1 = a_2 = \cdots = a_n = 1$ . This summand is invariant under  $S_n$  and contains a summand on which  $S_n$  acts by the signature character  $\sigma : S_n \rightarrow \{\pm 1\}$ . This defines a unique subsheaf  $R^n \pi_{*,n}(\mathbb{Z}_\ell)(\sigma) \subset R^n \pi_{*,n}(\mathbb{Z}_\ell)$  and hence we get an inclusion

$$H^1(M_N \times \bar{\mathbb{Q}}, R^n \pi_{*,n}(\mathbb{Z}_\ell)(\sigma)) \hookrightarrow H^{n+1}(\mathcal{E}^{(n)} \times \bar{\mathbb{Q}}, \mathbb{Z}_\ell)$$

and we can do the same thing for the cohomology with compact supports.

Now I will explain:

A) If we extend the scalars from  $\mathbb{Q}$  to  $\mathbb{C}$  then the extension of  $R^n \pi_{*,n}(\mathbb{Q}_\ell)(\sigma)$  is isomorphic to the restriction of  $\mathcal{M}_n \otimes \mathbb{Q}_\ell$  to the étale topology.

B) The Hecke operators  $T_p$  for  $p \nmid N$  are coming from algebraic correspondences  $T_p \subset M_N \times M_N$  and induce endomorphisms  $T_p : H^1(M_N \times \bar{\mathbb{Q}}, R^n \pi_{*,n}(\mathbb{Z}_\ell)(\sigma)) \rightarrow H^1(M_N \times \bar{\mathbb{Q}}, R^n \pi_{*,n}(\mathbb{Z}_\ell)(\sigma))$  which commute with the action of  $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$  on the cohomology.

C) This tells us that after extension of the scalars of the coefficient system we get

$$H^1(M_N(\mathbb{C}), \mathcal{M}_n \otimes \mathbb{Q}_\ell) \xrightarrow{\sim} H^1(M_N \times \bar{\mathbb{Q}}, R^n \pi_{*,n}(\mathbb{Q}_\ell)(\sigma))$$

and this gives us the structure of a  $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \times \mathcal{H}_\Gamma$  on  $H^1(M_N(\mathbb{C}), \mathcal{M}_n \otimes \mathbb{Q}_\ell)$ .

D) The operation of the Galois group on  $H^1(M_N(\mathbb{C}), \mathcal{M}_n \otimes \mathbb{Q}_\ell)$  is unramified outside  $N$ , therefore we have the conjugacy class  $\Phi_p^{-1}$  for all  $p \nmid N$  as endomorphism of  $H^1(M_N(\mathbb{C}), \mathcal{M}_n \otimes \mathbb{Q}_\ell)$ .

Now we use another fact, which will be explained in Chapter III. We also can define a Hecke algebra  $\mathcal{H}_p$  for the primes  $p \mid N$ , and hence we get an action of a larger Hecke algebra

$$\mathcal{H}_N^{\text{large}} = \bigotimes_p' \mathcal{H}_p$$

and this algebra commutes with the action of the Galois group.

We now apply our theorem 2 to the cohomology  $H_1^1(M_N(\mathbb{C}), \mathcal{M}_n \otimes \mathbb{Q}_\ell)$ , as a module under this large Hecke algebra. Then the isotypical summands will be invariant under the Galois group.

**Theorem 4:** a) *The multiplicity of an irreducible representation  $\Pi_f \in \text{Coh}(M_N(\mathbb{C}), \tilde{\mathcal{M}}_{n, L_1})$  is two.*  
 b) *This gives a product decomposition*

$$H_1^1(M_N(\mathbb{C}), \mathcal{M}_n \otimes L_1) \xrightarrow{\sim} H_{\Pi_f} \otimes W(\Pi_f),,$$

where  $H_{\Pi_f}$  is irreducible of type  $\Pi_f$  and where  $W(\Pi_f)$  is a two dimensional  $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$  module.

The module  $W(\Pi_f)$  is unramified outside  $N$  and

$$\text{tr}(\Phi_p^{-1}|W(\Pi_f)) = \lambda(\pi_p), \det(\Phi_p^{-1}|W(\Pi_f)) = p^{n+1}\omega(\Pi_f)(p)$$

This theorem is much deeper than the previous ones. The assertion a) follows from the theory of automorphic forms on  $\text{GL}_2$  and b) requires some tools from algebraic geometry. We have to consider the reduction  $M_N \times \text{Spec}(\mathbb{F}_p)$  and to look at the reduction of the Hecke operator  $T_p$  modulo  $p$ . I will resume this discussion in Chap. V.

I want to discuss some applications.

A) To any isotypical component  $\Pi_f$  we can attach an ( so called automorphic)  $L$  function

$$L(\Pi_f, s) = \prod_p L(\pi_p, s)$$

where for  $p \nmid N$  we define

$$L(\pi_p, s) = \frac{1}{1 - \lambda(\pi_p)p^{-s} + p^{n+1}\omega(\Pi_f)(p)p^{-2s}}$$

and for  $p|N$  we have

$$L(\pi_p, s) = \begin{cases} \frac{1}{1 - p^{n+1}\omega(\Pi_f)(p)p^{-s}} & \text{if } \pi_p \text{ is a Steinberg module} \\ 1 & \text{else} \end{cases}$$

This  $L$ -function, which is defined as an infinite product is holomorphic for  $\Re(s) \gg 0$  it can be written as the Mellin transform of a holomorphic cusp form  $F$  of weight  $n + 2$  and this implies that

$$\Lambda(\Pi, s) = \frac{\Gamma(s)}{2\pi^s} L(\Pi_f, s)$$

has a holomorphic continuation into the entire complex plane and satisfies a functional equation

$$\Lambda(\Pi_f, s) = W(\Pi_f)(N(\Pi_f))^{s-1-n/2} \Lambda(\Pi_f, n + 2 - s)$$

Here  $W(\Pi_f)$  is the so called root number, it can be computed from the  $\pi_p$  where  $p|N$ , its value is  $\pm 1$ , the number  $N(\Pi_f)$  is the conductor of  $\Pi_f$  it is a positive integer, whose prime factors are contained in the set of prime divisors of  $N$ .

B) But we also can interpret an isotypic component as a submotive in  $H^{n+1}(\mathcal{E}^{(n)} \times \mathbb{Q}, \mathbb{Z})$ , this is the so called Scholl motive.

If we apply the results of Deligne in Weil II, which have been proved in the winter term 2003/4, we get the estimate

$$|\iota(\lambda(\pi_p))| \leq 2p^{(n+1)/2}$$

for any embedding  $\iota$  of  $L$  into  $\mathbb{C}$ .

*End of probably removed section*

### 2.2.5 The $\ell$ -adic Galois representation in the first non trivial case

Again we consider the module  $\mathcal{M} = \mathcal{M}_{10}[-10]$ . We choose a prime  $\ell$  and for some reason let us assume  $\ell > 7$ . Then we can consider the cohomology groups

$$H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}/\ell^n \tilde{\mathcal{M}})$$

and the projective limit

$$H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}} \otimes \mathbb{Z}_\ell) = \lim_{\leftarrow} H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}/\ell^n \tilde{\mathcal{M}}).$$

Now it is known that the quotient space is the "moduli space" of elliptic curves, this is an imprecise and even incorrect statement, but it contains a lot of truth. What is true is that we can define the moduli stack  $S/\text{Spec}(\mathbb{Z})$  of elliptic curves, this is a smooth stack and it has the universal elliptic curve  $\mathcal{E} \xrightarrow{\pi} S$  over it.

We can define etale torsion sheaves  $(\mathcal{M}/\ell^n \tilde{\mathcal{M}})_{et}$  on this stack and we know that

$$H_{et}^1(S \times_{\text{Spec}(\mathbb{Z})} \bar{\mathbb{Q}}, (\mathcal{M}/\ell^n \tilde{\mathcal{M}})_{et}) \xrightarrow{\sim} H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{10}/\ell^n \tilde{\mathcal{M}}_{10}).$$

On these etale cohomology groups we have an action of the Galois group. Using correspondences we can define Hecke operators  $T_p$  for all  $p \neq \ell$ , they induce endomorphism on the etale cohomology and they commute with the action of the Galois group.

We denote this action of the Galois group as a representation

$$\rho_n : \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{Gl}(H_{et}^1(S \times_{\text{Spec}(\mathbb{Z})} \bar{\mathbb{Q}}, (\mathcal{M}/\ell^n \tilde{\mathcal{M}})_{et})).$$

This representation is unramified outside  $\ell$ , and this means:

The finite extension  $K_\ell^{(n)}/\mathbb{Q}$  for which  $\text{Gal}(\bar{\mathbb{Q}}/K_\ell^{(n)})$  is the kernel of  $\rho_n$  is unramified outside  $\ell$ .

By transport of structure we have the same projective system of Hecke  $\times$  Galois modules on the right hand side.

We recall our fundamental exact sequence, the Galois groups acts on the individual terms of this sequence, we get projective systems of Galois-modules and passing to the limit yields

$$\rho_l : \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{Gl}(H_1^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}} \otimes \mathbb{Z}_\ell))$$

and

$$\rho_\partial : \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{Gl}(\mathbb{Z}_\ell e_{10}).$$

The field  $K_\ell = \bigcup_n K_\ell^{(n)}$  defines the kernel  $\text{Gal}(\bar{\mathbb{Q}}/K_\ell)$ , the extension  $K_\ell/\mathbb{Q}$  is unramified at all primes  $p \neq \ell$ . If  $\mathfrak{p}$  is a prime in  $\mathcal{O}_{K_\ell}$  which lies above then the geometric Frobenius  $\Phi_{\mathfrak{p}}$  is the unique element in  $\text{Gal}(K_\ell/\mathbb{Q})$  which fixes  $\mathfrak{p}$  and induces  $x \mapsto x^{-p}$  on the residue field  $\mathcal{O}_{K_\ell}/\mathfrak{p}$ . This element defines a unique conjugacy class  $\Phi_p$  in  $\text{Gal}(K_\ell/\mathbb{Q})$ .

**Theorem**(Deligne) *For any prime  $p \neq \ell$  we have*

$$\rho_\partial(\Phi_p) = p^{11} Id$$

and

$$\det(Id - \rho(\Phi_p)t | H_1^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}} \otimes \mathbb{Z}_\ell)) = 1 - \tau(p)t + p^{11}t^2$$

This is a special case of the general theorem stated in the previous section and it one of the aims of the subject treated in this book to generalize this theorem to larger groups.

We conclude by giving a few applications.

A) The function  $z \mapsto \Delta(z)$  is a function on the upper half plane  $\mathbb{H} = \{z | \Im(z) > 0\}$  and it satisfies

$$\Delta\left(\frac{az+b}{cz+d}\right) = (cz+d)^{12} \Delta(z)$$

and this means that it is a modular form of weight 12. Since it goes to zero if  $z = iy \rightarrow \infty$  it is even a modular cusp form.

For such a modular cusp form we can define the Hecke  $L$ -function

$$L(\Delta, s) = \int_0^\infty \Delta(iy)y^s \frac{dy}{y} = \frac{\Gamma(s)}{(2\pi)^s} \sum_{n=1}^\infty \frac{\tau(n)}{n^s} = \frac{\Gamma(s)}{(2\pi)^s} \prod_p \frac{1}{1 - \tau(p)p^{-s} + p^{11-2s}}$$

the product expansion has been discovered by Ramanujan and has been proved by Mordell and Hecke.

Now it is in any textbook on modular forms that the transformation rule

$$\Delta\left(-\frac{1}{z}\right) = z^{12} \Delta(z)$$

implies that  $L(\Delta, s)$  defines a holomorphic function in the entire  $s$  plane and satisfies the functional equation

$$L(\Delta, s) = (-1)^{12/2} L(\Delta, 12 - s) = L(\Delta, 12 - s).$$

This function  $L(\Delta, s)$  is the prototype of an automorphic  $L$ -function. The above theorem shows that it is equal to a "motivic"  $L$ -function. We gave some vague explanations of what this possibly means: We can interpret the projective system  $(\mathcal{M}/\ell^n \tilde{\mathcal{M}})_{et}$  as the  $\ell$ -adic realization of a motive:

$$\mathcal{M} = \text{Sym}^{10}(R^1(\pi : \mathcal{E} \rightarrow S))$$

(All this is a translation of Deligne's reasoning into a more sophisticated language.)

It is a general hope that "motivic"  $L$ -functions  $L(M, s)$  have nice properties as functions in the variable  $s$  (meromorphicity, control of the poles, functional equation). So far the only cases, in which one could prove such nice properties are cases where one could identify the "motivic"  $L$ -function to an automorphic  $L$  function. The greatest success of this strategy is Wiles' proof of the Shimura-Taniyama-Weil conjecture, but also the Riemann  $\zeta$ -function is a motivic  $L$ -function and Riemann's proof of the functional equation follows exactly this strategy.

B) But we also have a flow of information in the opposite direction. In 1973 Deligne proved the Weil conjectures, which in this case say that the two roots of the quadratic equation

$$x^2 - \tau(p)x + p^{11} = 0$$

have absolute value  $p^{11/2}$ , i.e. they have the same absolute value. This implies the famous Ramanujan-conjecture

$$\tau(p) \leq 2p^{11/2}$$

and for more than 50 years this has been a brain-teaser for mathematicians working in the field of modular forms.

C) We consider the Galois representation

$$\rho : \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{Gl}(H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}} \otimes \mathbb{Z}_\ell))$$

and its sub and quotient representations

$$\rho_l : \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{Gl}(H_l^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}} \otimes \mathbb{Z}_\ell)), \rho_\partial : \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{Gl}(\mathbb{Z}_\ell e_{10}).$$

The representation  $\rho_\partial$  is the  $\ell$ -adic realization of the Tate-motive  $\mathbb{Z}(-11)$  (For a slightly more precise explanation I refer to MixMot.pdf on my homepage). On  $\mathbb{Z}_\ell(-1) = H^2(\mathbb{P}^1 \times \bar{\mathbb{Q}}, \mathbb{Z}_\ell)$  the Galois group acts by the Tate-character

$$\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{Gal}(\mathbb{Q}(\zeta_{\ell^\infty})/\mathbb{Q}) \xrightarrow{\alpha} \mathbb{Z}_\ell^\times$$

where  $\mathbb{Q}(\zeta_{\ell^\infty})$  is the cyclotomic field of all  $\ell^n$ -th roots of unity ( $n \rightarrow \infty$ ). We identify  $\text{Gal}(\mathbb{Q}(\zeta_{\ell^\infty})/\mathbb{Q}) = \mathbb{Z}_\ell^\times$ , the identification is given by the map  $x \mapsto (\zeta \mapsto \zeta^x)$  and then  $\alpha(x) = x^{-1}$ . Hence the first assertion in Delignes theorem simply says:

$$\rho_\partial = \alpha^{11}.$$

We say a few words concerning

$$\rho_l : \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{Gl}(H_l^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}} \otimes \mathbb{Z}_\ell)).$$

It is easy to see that the cup product provides a non degenerate alternating pairing

$$\langle , \rangle : H_1^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}} \otimes \mathbb{Z}_\ell) \times H_1^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}} \otimes \mathbb{Z}_\ell) \rightarrow \mathbb{Z}_\ell(-11)$$

and clearly for any  $\sigma \in \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$  we must have

$$\langle \rho(\sigma)u, \rho(\sigma)v \rangle = \alpha^{11}(\sigma) \langle u, v \rangle .$$

This means we have  $\det(\rho(\sigma)) = \alpha^{11}(\sigma)$  and we can ask what is the image of  $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$  in  $\text{Gl}(H_1^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}} \otimes \mathbb{Z}_\ell) = \text{Gl}_2(\mathbb{Z}_\ell)$ . We ask a seemingly simpler question and we want to understand the image of

$$\rho!, \text{ mod } \ell \text{ Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{Gl}(H_1^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}} \otimes \mathbb{F}_\ell) = \text{Gl}_2(\mathbb{F}_\ell).$$

This question is discussed in the paper " On  $\ell$ -adic representations and congruences for coefficients of modular forms," Springer lecture Notes 350, Modular Functions of one Variable III by H.P.F. Swinnerton-Dyer.

Here we can say that the image of this homomorphism composed with the determinant will be  $(\mathbb{F}_\ell^\times)^{11} \subset \mathbb{F}_\ell^\times$ . It is shown in the above paper that for  $\ell \neq 2, 3, 5, 7, 23, 691$  the image of the Galois group will simply be as large as possible, namely it will be the inverse image of  $(\mathbb{F}_\ell^\times)^{11}$ .

We can apply the Manin-Drinfeld principle and conclude that after tensorization by  $\mathbb{Q}_\ell$  the representation  $\rho \otimes \mathbb{Q}_\ell$  splits

$$\rho \otimes \mathbb{Q}_\ell = \rho_1 \otimes \mathbb{Q}_\ell \oplus \mathbb{Q}_\ell e_{10}(-11).$$

In section 2.2.3 we have seen that we have such a splitting also for the integral cohomology, i.e. for the module  $H^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}} \otimes \mathbb{Z}_\ell)$  provided  $\ell$  is not one of the small primes, which have been inverted and  $\ell \neq 691$ .

But if  $\ell = 691$  then we have seen in 2.2.3 that we have a homomorphism

$$j : \mathbb{Z}/(691)(-11) \hookrightarrow H_1^1(\Gamma \backslash \mathbb{H}, \tilde{\mathcal{M}}_{\mathbb{Z}/(691)}),$$

this is a homomorphism of Galois-modules. This means that the representation of the of the Galois group modulo  $\ell = 691$  is of the form

$$\begin{aligned} \rho!, \text{ mod } 691 : \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) &\rightarrow \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \\ \rho!, \text{ mod } 691(\sigma) &\mapsto \begin{pmatrix} \alpha(\sigma)^{11} & u(\sigma) \\ 0 & 1 \end{pmatrix} \end{aligned}$$

The field  $K_{691}^{(1)}$  contains the 691- th roots of unity and is an unramified extension of degree 691, in a sense this extension is now obtained by an explicit construction.



## Chapter 6

# Cohomology in the adelic language

### 6.1 The spaces

#### 6.1.1 The (generalized) symmetric spaces

Our basic datum is a connected reductive group  $G/\mathbb{Q}$ . Let  $G^{(1)}/\mathbb{Q}$  be its derived group and let  $C/\mathbb{Q}$  its centre. Then  $G^{(1)}/\mathbb{Q}$  is semi simple and  $C/\mathbb{Q}$  is a torus. The multiplication provides a canonical map

$$m : C \times G^{(1)} \rightarrow G, \quad (6.1)$$

it is an isogeny, this means that the kernel  $\mu_C = C \cap G^{(1)}$  of this map is a finite group scheme of multiplicative type. A finite group scheme of multiplicative type is simply an abelian group together with an action of the Galois group  $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$  on it. If we have such an isogeny as in (6.1) we write  $G = C \cdot G^{(1)}$ .

Let  $S/\mathbb{Q}$  be the maximal  $\mathbb{Q}$ -split torus in  $C/\mathbb{Q}$ . Up to isogeny we have  $C = C_1 \cdot S$  where  $C_1$  is the maximal anisotropic subtorus of  $C/\mathbb{Q}$ . We also introduce the group  $G_1 = G^{(1)} \cdot C_1$ . We have an exact sequence

$$1 \rightarrow G^{(1)} \rightarrow G \xrightarrow{d_C} C' \rightarrow 1,$$

the quotient  $C'$  is a torus and the restricted map  $d_C : C \rightarrow C'$  is an isogeny.

If  $\tilde{G}^{(1)}/\mathbb{Q}$  is the simply connected covering of  $G^{(1)}$ , then we get an isogeny

$$m_1 : \tilde{G} = \tilde{G}^{(1)} \times C_1 \times S \rightarrow G \quad (6.2)$$

Let  $\mathfrak{g}, \mathfrak{g}^{(1)}, \mathfrak{c}, \mathfrak{c}_1, \mathfrak{z}$  be the Lie algebras of  $G/\mathbb{Q}, G^{(1)}/\mathbb{Q}, C/\mathbb{Q}, C_1/\mathbb{Q}, S/\mathbb{Q}$ , then the differential of  $m_1$  induces an isomorphism

$$D_{m_1} : \mathfrak{g} \rightarrow \mathfrak{g}^{(1)} \oplus \mathfrak{c}_1 \oplus \mathfrak{z} \quad (6.3)$$

On  $\mathfrak{g}$  we have the Killing form  $B : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{Q}$  be the Killing form, it is defined by the rule

$$(T_1, T_2) \mapsto \text{trace}(\text{ad}(T_1) \circ \text{ad}(T_2)) \quad (6.4)$$

(See [chap2] 1.2.2) The Killing form is actually a bilinear form on  $\mathfrak{g}^{(1)} = \mathfrak{g}/(\mathfrak{c}_1 \oplus \mathfrak{z})$  and the restriction  $B : \mathfrak{g}^{(1)} \times \mathfrak{g}^{(1)} \rightarrow \mathbb{Q}$  is nondegenerate (see chap2 and chap4).

An automorphism  $\Theta : \tilde{G}^{(1)} \times_{\mathbb{Q}} \mathbb{R} \rightarrow \tilde{G}^{(1)} \times_{\mathbb{Q}} \mathbb{R}$  is called a Cartan involution if  $\Theta^2 = \text{Id}$  and if the bilinear form

$$B_{\Theta}(T_1, T_2) = B(T_1, \Theta(T_2)) \quad (6.5)$$

on  $\mathfrak{g} \otimes \mathbb{R}$  is negative definite.

If  $\Theta$  is a Cartan involution then it induces an automorphism -also called  $\Theta$ - on the Lie algebra  $\mathfrak{g}_{\mathbb{R}} = \mathfrak{g} \otimes \mathbb{R}$  and decomposes it into a + and a - eigenspace

$$\mathfrak{g}_{\mathbb{R}} = \mathfrak{k} \oplus \mathfrak{p} \quad (6.6)$$

and then clearly the + eigenspace  $\mathfrak{k}$  is a Lie subalgebra and  $[\mathfrak{p}, \mathfrak{p}] \subset \mathfrak{k}$ . The Killing form is negative definite on  $\mathfrak{k}$  and positive definite on  $\mathfrak{p}$ . This explains the above assertion on  $B_{\Theta}$ .

The topological group of real points  $\tilde{G}^{(1)}(\mathbb{R})$  is connected (see ref?). Then we have the classical theorem

**Theorem 6.1.1.** *The fixed group  $K_{\infty}^{(1)} = \tilde{G}^{(1)}(\mathbb{R})^{\Theta}$  is a maximal compact subgroup and it is also connected. The Cartan involutions are conjugate under the action of  $\tilde{G}^{(1)}(\mathbb{R})$ , and therefore the maximal compact subgroups of  $\tilde{G}^{(1)}(\mathbb{R})$  are conjugate.*

We introduce the space  $\tilde{X}^{(1)}$  of Cartan involutions on  $\tilde{G}^{(1)} \times_{\mathbb{Q}} \mathbb{R}$ , it is a homogenous space under the action of  $\tilde{G}^{(1)}(\mathbb{R})$  by conjugation and if we choose a  $\Theta$  or  $K_{\infty}^{(1)}$  then

$$\tilde{X}^{(1)} = \tilde{G}^{(1)}(\mathbb{R})/K_{\infty}^{(1)} \quad (6.7)$$

This is the symmetric space attached to  $\tilde{G}^{(1)} \times_{\mathbb{Q}} \mathbb{R}$ .

**Proposition 6.1.1.** *The symmetric space  $\tilde{X}^{(1)} = \tilde{G}^{(1)}(\mathbb{R})/K_{\infty}^{(1)}$  is diffeomorphic to  $\mathbb{R}^d$ , where  $d = \dim \mathfrak{p}$ , it carries a Riemannian metric which is  $\tilde{G}^{(1)}(\mathbb{R})$  invariant.*

We have to be aware that it may happen that  $\Theta$  is the identity. Then  $\tilde{G}^{(1)}(\mathbb{R}) = K_{\infty}^{(1)}$  and our symmetric space is a point.

We extend  $\Theta$  to an involution on  $\tilde{G} \times \mathbb{R}$  it will be simply the identity on the other two factors. Then it also induces an involution, again called  $\Theta$  on  $G \times \mathbb{R}$ .

We return to our reductive group  $G/\mathbb{Q}$ . We compare it to  $\tilde{G}$  via the homomorphism  $m_1$  in (6.2). Let  $K_{\infty}^C$  be the connected component of the identity of the maximal compact subgroup in  $C_1(\mathbb{R})$  and let  $Z'(\mathbb{R})^0$  be the connected component of the identity of the group of real points a subtorus  $Z' \subset S$ . Then we put

$$K_{\infty} = m_1(K_{\infty}^{(1)} \times K_{\infty}^C \times Z'(\mathbb{R})^0)$$

This group  $K_{\infty}$  is connected and if we divide by  $Z'(\mathbb{R})^0$  it is compact, more precisely we can say that  $K_{\infty}/Z'(\mathbb{R})^0$  is the connected component of a maximal

compact subgroup in  $G(\mathbb{R})/Z'(\mathbb{R})^0$ . The choice of the subtorus  $Z'$  is arbitrary and in a certain sense irrelevant. We could choose  $Z' = Z$  then we call  $K_\infty$  *saturated*, this choice is very convenient but in certain situations it is better to make a different choice, for instance we may choose  $Z' = 1$ .

To such a pair  $(G, K_\infty)$  we attach the (*generalized*) *symmetric space*

$$X = G(\mathbb{R})/K_\infty.$$

Here are a few comments concerning the structure of this space. (see also Chap II. 1.3) We observe that by construction  $K_\infty$  is connected, hence we have that  $K_\infty \subset G(\mathbb{R})^0$ . So if as usual  $\pi_0(G(\mathbb{R}))$  denotes the set of connected components, then we see that

$$\pi_0(X) = \pi_0(G(\mathbb{R})).$$

The connected component of the identity of  $\tilde{G}(\mathbb{R})$  maps under  $m_1$  to the connected component of the identity of  $G(\mathbb{R})$ , i.e.

$$\tilde{G}(\mathbb{R}) = \tilde{G}^{(1)}(\mathbb{R}) \times C_1(\mathbb{R})^0 \times S(\mathbb{R})^0 \rightarrow G(\mathbb{R})^0$$

and if we divide by  $K_\infty^{(1)} \times K_\infty^C \times Z'(\mathbb{R})^0$ , resp.  $K_\infty$  we get a diffeomorphism with the connected component corresponding to the identity

$$\tilde{G}^{(1)}(\mathbb{R})/K_\infty^{(1)} \times C_1(\mathbb{R})^0/K_\infty^C \times S(\mathbb{R})^0/Z'(\mathbb{R}) \xrightarrow{\sim} X_1 \subset X.$$

We want to describe the other connected components of  $X$ . It is well known that we can find a maximal split torus  $\tilde{S}_1 \subset \tilde{G}^{(1)} \times \mathbb{R}$  which is invariant under our given Cartan involution  $\Theta$ . The homomorphism  $m_1$  maps  $\tilde{G}^{(1)}(\mathbb{R}) \rightarrow G^{(1)}(\mathbb{R})$ . The fixed group  $G^{(1)}(\mathbb{R})^\Theta$  is a compact subgroup whose connected component of the identity is the image of  $K_\infty^{(1)}$  under  $m_1$ . Our torus  $\tilde{S}_1$  sits as the first component in the maximal split torus

$$\tilde{S}_2 = \tilde{S}_1 \times C_1^{\text{split}} \times S$$

Then it is clear that  $\Theta$  induces the involution  $t \mapsto t^{-1}$  on  $\tilde{S}_1$ . Let  $S_2$  be the image of  $\tilde{S}_2$  under  $m_1$ . We have the following proposition

**Proposition 6.1.2.** *a) The group of 2-division points  $S_2[2]$  normalizes  $K_\infty$ .*

*b) We have an exact sequence*

$$\rightarrow \tilde{S}_2[2] \rightarrow S_2[2] \xrightarrow{r} \pi_0(G(\mathbb{R})) \rightarrow 0$$

*c) If  $K_\infty^0$  is the image of  $K_\infty^{(1)} \times K_\infty^C$  then  $K_\infty^0 \cdot S_2[2]$  is a maximal compact subgroup of  $G(\mathbb{R})$ .*

*Proof.* Rather obvious, the surjectivity of  $r$  requires an argument in Galois cohomology. (Details later)  $\square$

Now we can write down all the connected components. We choose a system  $\Xi$  of representatives for  $S_2[2]/\tilde{S}_2[2]$  and for any  $\xi \in \Xi$  we get a diffeomorphism

$$\tilde{G}^{(1)}(\mathbb{R})/K_\infty^{(1)} \times C_1(\mathbb{R})^0/K_\infty^C \times S(\mathbb{R})^0/Z'(\mathbb{R}) \rightarrow X_\xi \subset X \quad (6.8)$$

$$g \mapsto g\xi$$

We may formulate this differently

**Proposition 6.1.3.** *The multiplication from the left by  $S_2[2]$  on  $G(\mathbb{R})$  induces an action of  $S_2[2]/\tilde{S}_2[2]$  on  $X$  and this action is simple transitive on the set of connected components.*

Let  $x_0 = K_\infty \in X$ . For any other point  $x \in X$  we find an element  $g \in X$  which translates  $x_0$  to  $x$ . Then the derivative of the translation provides an isomorphism between the tangent spaces

$$D_g : T_{x_0} = \mathfrak{p} \xrightarrow{\sim} T_x.$$

This isomorphism depends of course on the choice of  $g$ . ( This will play a role in section (8.1)). But we apply this to the highest exterior power and get an isomorphism

$$D_g : \Lambda^d(\mathfrak{p}) \xrightarrow{\sim} \Lambda^d(T_x)$$

which does not depend on the choice of  $g$  because the connected group  $K_\infty$  acts trivially on  $\Lambda^d(\mathfrak{p})$ . Hence we can say that we can find a *consistent* orientation on  $X$  : We chose a generator in  $\Lambda^d(\mathfrak{p})$  the  $D_g$  yields a generator in  $\Lambda^d(T_x)$ .

If our reductive group is an anisotropic torus  $T/\mathbb{Q}$ , then we have for the connected component of the identity

$$T(\mathbb{R})^{(0)} \xrightarrow{\sim} (\mathbb{R}_{>0}^\times)^a \times (S^1)^b.$$

Then our maximal compact subgroup  $K_\infty^T$  is simply the product of the circles and

$$X_T = T(\mathbb{R})/K_\infty^T$$

is nothing else than as disjoint union of copies of  $\mathbb{R}^a$ . The situation is similar for a split torus but then we have the freedom, to divide out the connected component of a subtorus.

As a standard example we can take  $G/\mathbb{Q} = \mathrm{GL}_2/\mathbb{Q}$ , then the connected component of the real points of the centre is  $\mathbb{R}_{>0}^\times$  and in this case we can take  $K_\infty = \mathrm{SO}(2) \cdot \mathbb{R}_{>0}^\times \subset \mathrm{GL}_2(\mathbb{R})$ . In this case the symmetric space is the union of an upper and a lower half plane. If we choose for our split torus  $S_1/\mathbb{R}$  the standard diagonal torus, then  $S_1[2]$  is the group of diagonal matrices with entries  $\pm 1$  and this normalizes  $K_\infty$ .

### 6.1.2 The locally symmetric spaces

Let  $\mathbb{A}$  be the ring of adeles, we decompose it into its finite and its infinite part:  $\mathbb{A} = \mathbb{R} \times \mathbb{A}_f$ . We have the group of adeles  $G(\mathbb{A}) = G(\mathbb{R}) \times G(\mathbb{A}_f)$ . We denote elements in the adèle group by underlined letters  $\underline{g}, \underline{h} \dots$  and so on. If we decompose an element  $\underline{g}$  into its finite and its infinite part then we denote this by  $g_\infty \times \underline{g}_f$ . Let  $K_f$  be a (variable) open compact subgroup of  $G(\mathbb{A}_f)$ . We always assume that this group is a product of local groups  $K_f = \prod_p K_p$ .

To get such subgroups we choose an integral structure (explain at some other place)  $\mathcal{G}/\mathrm{Spec}(\mathbb{Z})$ . Then we know that we have  $K_p = \mathcal{G}(\mathbb{Z}_p)$  for almost all  $p$ . Furthermore we know that  $\mathcal{G} \times \mathrm{Spec}(\mathbb{Z}_p)/\mathrm{Spec}(\mathbb{Z}_p)$  is a reductive group scheme for almost all primes  $p$ .

If  $\mathcal{G}/\text{Spec}(\mathbb{Z})$  and  $K_f$  are given, then we select a finite set  $\Sigma$  of finite primes which contains the primes  $p$  where  $\mathcal{G}/\mathbb{Z}_p$  is not reductive and those where  $K_p$  is not equal to  $\mathcal{G}(\mathbb{Z}_p)$ . This set  $\Sigma$  will be called the set of *ramified* primes.

The general agreement will be that we use letters  $\mathcal{G}, \mathcal{T}, \mathcal{U}, \dots$  for group schemes over the integers, or over  $\mathbb{Z}_p$  and then their general fiber will be  $G, T, U, \dots$ .

Readers who are not so familiar with this language may think of the simple example where  $G/\mathbb{Q} = GSp_n/\mathbb{Q}$  is the group of symplectic similitudes on  $V = \mathbb{Q}^{2n} = \mathbb{Q}e_1 \oplus \dots \oplus \mathbb{Q}e_n \oplus \mathbb{Q}f_1 \oplus \dots \oplus \mathbb{Q}f_n$  with the standard symplectic form which is given by  $\langle e_i, f_i \rangle = 1$  for all  $i$  and where all other products zero. The vector space contains the lattice  $L = \mathbb{Z}^{2n} = \mathbb{Z}e_1 \oplus \dots \oplus \mathbb{Z}e_n \oplus \mathbb{Z}f_1 \oplus \dots \oplus \mathbb{Z}f_n$ . This lattice defines a unique integral structure  $\mathcal{G}/\mathbb{Z}$  on  $G/\mathbb{Q}$  for which  $\mathcal{G}(\mathbb{Z}_p) = \{g \in G(\mathbb{Q}_p) | g(L \otimes \mathbb{Z}_p) = (L \otimes \mathbb{Z}_p)\}$ . In this case the group scheme is reductive over  $\text{Spec}(\mathbb{Z})$ . This integral structure gives us a privileged choice of an open maximal compact subgroup: Within the ring  $\mathbb{A}_f$  of finite adeles we have the ring  $\hat{\mathbb{Z}} = \varprojlim \mathbb{Z}/m\mathbb{Z}$  of integral finite adeles and we can consider  $K_f^0 = \mathcal{G}(\hat{\mathbb{Z}}) = \prod_p \mathcal{G}(\mathbb{Z}_p)$ . This is a very specific choice. In this case the set  $\Sigma = \emptyset$ , we say that  $K_f = K_f^0$  is unramified.

Starting from there we can define new subgroups  $K_f$  by imposing some congruence conditions at a finite set  $\Sigma$  of primes. These congruence conditions then define congruence subgroups  $K_p \subset K_p^0$ . This set  $\Sigma$  of places where we impose congruence condition will then be the set of ramified primes. (See the example further down.) Then we define the level subgroup

$$K_f = \prod_{p \in \Sigma} K_p \times \prod_{p \notin \Sigma} \mathcal{G}(\mathbb{Z}_p). \tag{6.9}$$

The space  $(G(\mathbb{R})/K_\infty) \times (G(\mathbb{A}_f)/K_f)$  can be seen as a product of the symmetric space and an infinite discrete set, on this space  $G(\mathbb{Q})$  acts properly discontinuously (see below) and the quotients

$$\mathcal{S}_{K_f}^G = G(\mathbb{Q}) \backslash (G(\mathbb{R})/K_\infty \times G(\mathbb{A}_f)/K_f)$$

are the locally symmetric spaces whose topological properties we want to study. We denote by

$$\pi : G(\mathbb{R})/K_\infty \times G(\mathbb{A}_f)/K_f \rightarrow \mathcal{S}_{K_f}^G = G(\mathbb{Q}) \backslash (G(\mathbb{R})/K_\infty \times G(\mathbb{A}_f)/K_f),$$

the projection map.

To get an idea of how this space looks like we consider the action of  $G(\mathbb{Q})$  on the discrete space  $G(\mathbb{A}_f)/K_f$ . It follows from classical finiteness results that this quotient is finite, let us pick representatives  $\{\underline{g}_f^{(i)}\}_{i=1..m}$ . We look at the stabilizer of the coset  $\underline{g}_f^{(i)} K_f/K_f$  in  $G(\mathbb{Q})$ . This stabilizer is obviously equal to  $\Gamma_{\underline{g}_f^{(i)}} = G(\mathbb{Q}) \cap \underline{g}_f^{(i)} K_f (\underline{g}_f^{(i)})^{-1}$  which is an arithmetic subgroup of  $G(\mathbb{Q})$ . This subgroup acts properly discontinuously on  $X$  (See Chap. II, 1.6).

Now we call the level subgroup  $K_f$  neat, if all the subgroups  $\Gamma_{\underline{g}_f^{(i)}}$  are torsion free. It is not hard to see, that for any choice of  $K_f$  we can pass to a subgroup of finite index  $K'_f$ , which is neat. Then we have

**Proposition 6.1.4.** *For any subgroup  $K_f$  the space  $\mathcal{S}_{K_f}^G$  is a finite union of quotient spaces  $\Gamma_i^{g_f^{(i)}} \backslash X$  where  $X = G(\mathbb{R})/K_\infty$  and the  $\Gamma_i = \Gamma_i^{g_f^{(i)}}$  are varying arithmetic congruence subgroups. If  $K_f$  is neat, these spaces are locally symmetric spaces. If  $K_f$  is not neat then we may pass to a neat subgroup  $K'_f$  which is even normal in  $K_f$ : We get a covering  $\mathcal{S}_{K'_f}^G \rightarrow \mathcal{S}_{K_f}^G$  which induces coverings  $\Gamma'_j \backslash X \rightarrow \Gamma_i \backslash X$ , where the  $\Gamma'_j$  are torsion free and normal in  $\Gamma_i$ . So we see that in general the quotients are orbifold locally symmetric spaces. For any point  $y \in \mathcal{S}_{K_f}^G$  we can find a neighborhood  $V_y$  such that  $\pi^{-1}(V_y)$  is the disjoint union of connected components  $W_{\underline{x}}, \underline{x} = (x_\infty, \underline{g}_f) \in \pi^{-1}(y)$ , and  $V_y = \Gamma_{x_\infty} \backslash W_{\underline{g}_f}$ , where  $\Gamma_{x_\infty}$  is the stabilizer of  $x_\infty$  intersected with  $\Gamma^{g_f}$ .*

We will consider the special case where  $G/\mathbb{Q}$  is the generic fibre of a split reductive scheme  $\mathcal{G}/\mathbb{Z}$ . In that case we can choose  $K_f = \prod_p \mathcal{G}(\mathbb{Z}_p)$ , this is then a maximal compact subgroup in  $G(\mathbb{A}_f)$ . Then  $K_f$  is unramified we will also say that the space  $\mathcal{S}_{K_f}^G$  is unramified. If in addition the derived group  $G^{(1)}/\mathbb{Q}$  is simply connected, then it is not difficult to see, that  $G(\mathbb{Q})$  acts transitively on  $G(\mathbb{A}_f)/K_f$  and hence we get

$$\mathcal{S}_{K_f}^G \xrightarrow{\sim} \mathcal{G}(\mathbb{Z}) \backslash X.$$

The homomorphism  $\mathbb{G}(\mathbb{Z}) \rightarrow \pi_0(C'(\mathbb{R}))$  is surjective we can conclude that  $\mathbb{G}(\mathbb{Z})$  acts transitively on  $\pi_0(X)$  and if  $\Gamma_0$  is the stabilizer of a connected component  $X^0$  of  $X$  then we find

$$\mathcal{S}_{K_f}^G \xrightarrow{\sim} \Gamma_0 \backslash X^0$$

especially we see that the quotient is connected. We discuss an example.

We start from the group  $G/\text{Spec}(\mathbb{Z}) = \text{Gl}_n/\text{Spec}(\mathbb{Z})$  then we may choose  $K_\infty = \text{SO}(n) \times \mathbb{R}_{>0}^\times \subset \text{Gl}_n(\mathbb{R})$ . and  $X = \text{Gl}_n(\mathbb{R})/K_\infty$  is the disjoint union of two copies of the space  $X$  of positive definite symmetric  $(n \times n)$  matrices up to homothetic by a positive scalar (or what amounts to the same with determinant one). If we choose  $K_f$  as above then we find

$$\mathcal{S}_{K_f}^G = \text{Sl}_n(\mathbb{Z}) \backslash X.$$

We have another special case. Let us assume that  $G/\mathbb{Q}$  is semi simple and simply connected. The group  $G \times \mathbb{R}$  is a product of simple groups over  $\mathbb{R}$  and we assume in addition that there is at least one non compact factor. Then we have the strong approximation theorem ([51],[63]) which says that for any choice of  $K_f$  the map from  $G(\mathbb{Q})$  to  $G(\mathbb{A}_f)/K_f$  is surjective, i.e. any  $\underline{g}_f \in G(\mathbb{A}_f)$  can be written as  $\underline{g}_f = a\underline{k}_f, a \in G(\mathbb{Q}), \underline{k}_f \in K_f$ . This clearly implies that then

$$\mathcal{S}_{K_f}^G = \Gamma \backslash G(\mathbb{R})/K_\infty \tag{6.10}$$

where  $\Gamma = K_f \cap G(\mathbb{Q})$ .

There is a contrasting case, this is the case when  $G/\mathbb{Q}$  is still semi simple and simply connected, but where  $G(\mathbb{R})$  is compact. In this case our symmetric space  $X$  is simply a point  $*$  and

$$\mathcal{S}_{K_f}^G = G(\mathbb{Q}) \backslash (* \times G(\mathbb{A}_f) / K_f).$$

This means that our topological space is simply a discrete set of points, hence it looks as if this is an entirely uninteresting and trivial case. But this is not so. To determine the finite set and the stabilizers is a highly non trivial task. Later we will construct sheaves and discuss the action of the Hecke algebra on the cohomology of these sheaves. Then it turns out that that it is not only the set of points and the stabilizers that is of interest but also the "interaction" among these points is of interest. Then it turns out that this case is as difficult as the case where  $\Gamma \backslash X$  becomes an honest space.

We give a few examples of such spaces

In the choice of our group  $K_\infty$  a subtorus  $Z' \subset S$  enters. The choice of this subtorus has very little influence on the structure of our locally symmetric space  $\mathcal{S}_{K_f}^G$ . Remember that the isogeny  $m$  in (6.1) induces an isogeny  $C \rightarrow C'$  and this isogeny yields an isogeny from  $S$  to the maximal split subtorus  $S' \subset C'$ . This homomorphism induces an isomorphism  $S(\mathbb{R})^0 \rightarrow S'(\mathbb{R})^0$ . If  $G_1(\mathbb{R})$  is the inverse image of the the group of 2-division points  $S'[2]$  then we get from this isomorphism that  $G(\mathbb{R}) = G_1(\mathbb{R}) \times S(\mathbb{R})^0$ . If we now consider the two spaces  $\mathcal{S}_{K_f}^G$  and  $(\mathcal{S}_{K_f}^G)^\dagger$ , the first one defined with an arbitrary torus  $Z'$  the second one with  $Z' = S$  then the arguments above imply that

$$\mathcal{S}_{K_f}^G = (\mathcal{S}_{K_f}^G)^\dagger \times (S(\mathbb{R})^0 / Z'(\mathbb{R})^0) \tag{6.11}$$

the second factor on the right hand side is isomorphic to  $\mathbb{R}^b$  and since we are interested in the cohomology group of this space, the second factor is irrelevant.

In certain situations we encounter cases where it is natural to choose a subgroup  $K_\infty$  which is slightly larger and not connected. If this is the case we denote the connected component  $K_\infty^{(1)}$  and we get two locally symmetric spaces and a finite map

$$G(\mathbb{Q}) \backslash \left( G(\mathbb{R}) / K_\infty^{(1)} \times G(\mathbb{A}_f) / K_f \right) \rightarrow G(\mathbb{Q}) \backslash (G(\mathbb{R}) / K_\infty \times G(\mathbb{A}_f) / K_f) \tag{6.12}$$

This map is a covering if  $K_f$  is neat and the space on the right is a quotient of the space on the left by an action of the finite elementary abelian [2]-group  $K_\infty / K_\infty^{(1)}$ .

In accordance with the terminology in number theory we call the space  $\mathcal{S}_{K_f}^G$  *narrow* if  $K_\infty^{(1)} = K_\infty$  and in general we call the space on the left the *narrow cover* of  $G(\mathbb{R}) / K_\infty \times G(\mathbb{A}_f) / K_f$ .

Ccomp

### 6.1.3 The group of connected components, the structure of $\pi_0(\mathcal{S}_{K_f}^G)$ .

If we keep our assumptions that  $G/\mathbb{Q}$  is reductive and  $G^{(1)}/\mathbb{Q}$  is simply connected and satisfies strong approximation. We choose a level subgroup  $K_f \subset$

$G(\mathbb{A}_f)$  and we put  $d_{C'}(K_\infty \times K_f) = K_\infty^{C'} \times K_f^{C'}$ . Then we claim that under these conditions  $\boxed{\text{conncomp}}$

$$\pi_0(\mathcal{S}_{K_f}^G) \xrightarrow{\sim} \pi_0(\mathcal{S}_{K_\infty^{C'} \times K_f^{C'}}^{C'}). \quad (6.13)$$

To see this we need a theorem of Tate which says that the map  $C'(\mathbb{Q}) \rightarrow \pi_0(C'(\mathbb{R}))$  is surjective. This implies that  $\pi_0(\mathcal{S}_{K_\infty^{C'} \times K_f^{C'}}^{C'}) = C'(\mathbb{Q})^{(0)} \backslash C'(\mathbb{A}_f) / K_f^{C'}$ , where  $C'(\mathbb{Q})^{(0)} \subset C'(\mathbb{Q})$  are the elements whose image lies in  $C'(\mathbb{R})^{(0)}$ . Now we need a little argument from Galois cohomology. The map  $G(\mathbb{A}_f) \rightarrow C'(\mathbb{A}_f)$  is surjective because for all primes  $p$   $H^1(\mathbb{Q}_p, G^{(1)})$  consist of the trivial class only. (Kneser and Bruhat-Tits ([?]). ) This implies the surjectivity: For the injectivity assume  $\underline{x}, \underline{y} \in C'(\mathbb{A}_f)$  and there is an element  $a \in C'(\mathbb{Q})^{(0)}$  with  $a\underline{x} = \underline{y}$ . Then we need to find a lift of  $a$  to an element  $b \in G(\mathbb{Q})$ . Again we invoke the standard argument from Galois cohomology. We have the exact sequence

$$G(\mathbb{Q}) \rightarrow C'(\mathbb{Q}) \xrightarrow{\delta} H^1(\mathbb{Q}, G^{(1)})$$

the obstruction to find  $b$  is an element  $\delta(a) \in H^1(\mathbb{Q}, G^{(1)})$ . We have the Hasse principle  $H^1(\mathbb{Q}, G^{(1)}) \xrightarrow{\sim} H^1(\mathbb{R}, G^{(1)})$  ([?]) but since  $a \in C'(\mathbb{Q})^{(0)}$  it follows that the image of  $\delta(a) \in H^1(\mathbb{R}, G^{(1)})$  is trivial, hence  $\delta(a)$  is trivial.

We have seen in the previous section that we can choose a consistent orientation on  $X = G(\mathbb{R})/K_\infty$  provided  $K_\infty$  is narrow. Then it clear this induces also a consistent orientation on  $\mathcal{S}_{K_f}^G$ .

$\boxed{\text{BSC1}}$

### 6.1.4 The Borel-Serre compactification

In general the space  $\mathcal{S}_{K_f}^G$  is not compact. Recall that in the definition of this quotient the choice of a subtorus  $Z'/\mathbb{Q}$  of  $S/\mathbb{Q}$  enters. This If  $Z' \neq S$  then the quotient will never be compact. But this kind of non compactness is "uninteresting". In the following we assume that  $Z' = S$ .

In this case we have the criterion of Borel - Harish-Chandra which says

*The quotient space  $\mathcal{S}_{K_f}^G$  is compact if and only if the group  $G/\mathbb{Q}$  has no proper parabolic subgroup over  $\mathbb{Q}$ .*

If we have a non trivial parabolic subgroup  $P/\mathbb{Q}$  then we add a boundary part  $\partial_P \mathcal{S}_{K_f}^G$  to  $\mathcal{S}_{K_f}^G$  it will depend only the  $G(\mathbb{Q})$ -conjugacy class of  $P$ . We will describe this boundary piece later. We define the Borel-Serre boundary

$$\partial(\mathcal{S}_{K_f}^G) = \bigcup_P \partial_P \mathcal{S}_{K_f}^G,$$

where  $P$  runs over the set of  $G(\mathbb{Q})$  conjugacy classes of parabolic subgroups. We will put a topology on this space and if  $Q \subset P$  then  $\partial_Q \mathcal{S}_{K_f}^G$  will be in the closure of  $\partial_P \mathcal{S}_{K_f}^G$ . Then

$$\mathcal{S}_{K_f}^{\bar{G}} = \mathcal{S}_{K_f}^G \cup \partial(\mathcal{S}_{K_f}^G)$$

will be a compact Hausdorff-space.

We describe the construction of this compactification in more detail. In chap4.pdf 2.7.1 **Ref korrigieren** we studied the group  $\text{Hom}(P, \mathbb{G}_m)$  and have seen that

$$\text{Hom}(P, \mathbb{G}_m) \otimes \mathbb{Q} = \text{Hom}(S_P, \mathbb{G}_m) \otimes \mathbb{Q}.$$

For any character  $\gamma \in \text{Hom}(P, \mathbb{G}_m)$  we get a homomorphism  $\gamma_A : P(\mathbb{A}) \rightarrow \mathbb{G}_m(\mathbb{A}) = I_{\mathbb{Q}}$ , the group of ideles. We have the idele norm  $|\cdot| : \underline{x} \mapsto |\underline{x}|$  from the idele group to  $\mathbb{R}_{>0}^{\times}$  and then we get by composing

$$|\gamma| : P(\mathbb{A}) \rightarrow \mathbb{R}_{>0}^{\times}.$$

It is obvious that we can extend this definition to characters  $\gamma \in \text{Hom}(P, \mathbb{G}_m) \otimes \mathbb{Q}$ , for such a  $\gamma$  we find a positive non zero integer  $m$  such that  $m\gamma \in \text{Hom}(P, \mathbb{G}_m)$  and then we define

$$|\gamma| = (|m\gamma|)^{\frac{1}{m}}$$

Later we will even extend this to a homomorphism  $\text{Hom}(P, \mathbb{G}_m) \otimes \mathbb{C} \rightarrow \text{Hom}(P(\mathbb{A}), \mathbb{C}^{\times})$  by the rule XtimesC

$$\gamma \otimes z \mapsto |\gamma|^z \tag{6.14}$$

If we have a parabolic subgroup  $P/\mathbb{Q}$  and a point  $(x, \underline{g}_f) \in X \times G(\mathbb{A}_f)/K_f$  then we attach to it a (strictly positive) number

$$p(P, (x, \underline{g}_f)) = \text{vol}_{d_x u}(U(\mathbb{Q}) \cap \underline{g}_f K_f \underline{g}_f^{-1} \backslash U(\mathbb{R})). \tag{6.15}$$

This needs explanation. The group  $U(\mathbb{Q}) \cap \underline{g}_f K_f \underline{g}_f^{-1} = \Gamma_{U, \underline{g}_f}$  is a cocompact discrete lattice in  $U(\mathbb{R})$ , we can describe it as the group of elements  $\gamma \in U(\mathbb{Q})$  which fix  $\underline{g}_f K_f$ , so it can be viewed as a lattice of integral elements where integrality is determined by  $\underline{g}_f$ . The point  $x$  defines a positive definite bilinear form  $B_{\Theta_x}$  on the Lie algebra  $\mathfrak{g} \otimes \mathbb{R}$ , and this bilinear form can be restricted to the Lie-algebra  $\mathfrak{u}_P \otimes \mathbb{R}$  and this provides a volume form  $d_x u$  on  $U(\mathbb{R})$  the above number is the volume of the nilmanifold  $\Gamma_{U, \underline{g}_f} \backslash U(\mathbb{R})$  with respect to this measure.

If we are in the special case that  $G = \text{Sl}_2/\mathbb{Q}$  and  $K_f = \text{Sl}_2(\hat{\mathbb{Z}})$  then a parabolic subgroup  $P$  is a point  $r = \frac{p}{q} \in \mathbb{P}^1(\mathbb{Q})$  (or  $\infty$ ) and then  $p(P, (z, 1))$  is small if  $z$  lies in a small Farey circle, i.e. it is close to  $r$ .

These numbers have some obvious properties

a) They are invariant under conjugation by an element  $a \in G(\mathbb{Q})$ , this means we have

$$p(a^{-1}Pa, (x, \underline{g}_f)) = p(P, a(x, \underline{g}_f))$$

b) If  $\underline{p} \in P(\mathbb{A})$  then we have

$$p(P, \underline{p}(x, \underline{g}_f)) = p(P, (x, \underline{g}_f)) |\rho_P|^2$$

The  $G(\mathbb{Q})$  conjugacy classes of parabolic are in one to one correspondence with the subsets  $\pi'$  of the set relative simple roots  $\pi_G$ : The minimal parabolic corresponds to the empty set, the non proper parabolic subgroup  $G/\mathbb{Q}$  corresponds to  $\pi_G$  itself. In general  $\pi'$  is the set of relative simple roots of the semi

simple part of the reductive quotient of the parabolic subgroup. For a parabolic subgroup  $P'$  corresponding to  $\pi'$  we put  $d(P') = \#(\pi_G \setminus \pi')$ . For any  $i \in \pi_G \setminus \pi'$  we have a fundamental character

$$\gamma_i : P \rightarrow \mathbb{G}_m.$$

We have the Borel-Serre compactification

$$i : \mathcal{S}_{K_f}^G \rightarrow \bar{\mathcal{S}}_{K_f}^G$$

The compactification is a manifold with corners, the boundary is stratified

$$\partial(\bar{\mathcal{S}}_{K_f}^G) = \bigcup_P \partial_P(\bar{\mathcal{S}}_{K_f}^G)$$

where  $P$  runs over the  $G(\mathbb{Q})$  conjugacy classes of parabolic subgroups. If  $P \subset Q$  then the stratum  $\partial_Q(\bar{\mathcal{S}}_{K_f}^G) \subset \overline{\partial_P(\bar{\mathcal{S}}_{K_f}^G)}$ .

Locally at a point  $x \in \partial_P(\bar{\mathcal{S}}_{K_f}^G)$  we find neighborhoods of  $x$  in  $\bar{\mathcal{S}}_{K_f}^G$  which are of the form

$$U_x = W_x \times \{\dots, u_i, \dots\}_{i \in \pi_G \setminus \pi'; 0 \leq u_i < \epsilon} \quad (6.16)$$

where  $W_x$  is a neighborhood of  $x$  in the orbifold  $\partial_P(\bar{\mathcal{S}}_{K_f}^G)$ . The intersection  $\overset{\circ}{U}_x = U_x \cap \mathcal{S}_{K_f}^G$  consists of those elements where all the  $u_i > 0$ .

### 6.1.5 The easiest but very important example

If we take for instance  $\mathbb{G}/\mathbb{Z} = \mathrm{Gl}_2/\mathbb{Z}$  and if we pick an integer  $N$  then we can define the congruence subgroup  $K_f(N) = \prod_p K_p(N) \subset \mathbb{G}(\hat{\mathbb{Z}})$ . It is defined by the condition that at all primes  $p$  dividing  $N$  the subgroup

$$K_p(N) = \{\gamma \in \mathbb{G}(\hat{\mathbb{Z}}) \mid \gamma \equiv \mathrm{Id} \pmod{p^{n_p}}\}$$

where of course  $p^{n_p}$  is the exact power of  $p$  dividing  $N$ . At the other primes we take the full group of integral points. For the discussion of the example we put  $K_f(N) = K_f$ .

If we consider the action of  $G(\mathbb{Q})$  on  $G(\mathbb{A}_f)/K_f$  then the determinant gives us a map

$$\mathrm{Gl}_2(\mathbb{Q}) \backslash \mathrm{Gl}_2(\mathbb{A}_f) / K_f \rightarrow \mathbb{G}_m(\mathbb{A}_f) / \mathbb{Q}^* \mathfrak{U}_N$$

where  $\mathfrak{U}_N$  is the group of unit ideles in  $I_{\mathbb{Q},f} = \mathbb{G}_m(\mathbb{A}_f)$  which satisfy  $u_p \equiv 1 \pmod{p^{n_p}}$ . This map is a bijection as one can easily see from strong approximation in  $Sl_2$ , and the right hand side is equal to  $(\mathbb{Z}/N\mathbb{Z})^* / \{\pm 1\}$ . At the infinite place we have that our symmetric space has two connected components, we have

$$X = \mathrm{Gl}_2(\mathbb{R}) / SO(2) = \mathbb{C} \setminus \mathbb{R} = \mathbb{H}_+ \cup \mathbb{H}_-$$

where  $\mathbb{H}_{\pm}$  are the upper and lower half plane, respectively. We have a complex structure on  $X$  which is invariant under the action of  $\mathrm{Gl}_2(\mathbb{R})$ . The connected components of this quotient correspond (one to one) to the elements in

$$\mathbb{G}_m(\mathbb{A})/\mathbb{G}_m(\mathbb{Q})(\mathbb{G}_m(\mathbb{R})^0 \times \mathfrak{U}_N) = I_{\mathbb{Q}}/\mathbb{Q}^*\mathbb{R}_{>0}^*\mathfrak{U}_N = (\mathbb{Z}/N\mathbb{Z})^*.$$

We put  $\Gamma(N) = G(\mathbb{Q}) \cap K_f$  and then the components are

$$\Gamma(N) \backslash \begin{pmatrix} t_{\infty} & 0 \\ 0 & 1 \end{pmatrix} H_+ \times \begin{pmatrix} t_f & 0 \\ 0 & 1 \end{pmatrix} K_f/K_f$$

where  $t$  runs through a set of representatives of  $I_{\mathbb{Q}}/\mathbb{Q}^*\mathbb{R}_{>0}^*\mathfrak{U}_N = (\mathbb{Z}/N\mathbb{Z})^*$ .

These connected components are Riemann surfaces which are not compact. They can be compactified by adding a finite number of points, the so called *cusps*. These are in one to one correspondence with the orbits of  $\Gamma(N)$  on  $\mathbb{P}^1(\mathbb{Q})$  (see reduction theory).

(Compare to Borel-Serre)

## 6.2 The sheaves and their cohomology

### 6.2.1 Basic data and simple properties

Let  $\mathcal{M}_{\mathbb{Q}}$  be a finite dimensional  $\mathbb{Q}$ -vector space, let

$$r : G/\mathbb{Q} \rightarrow \text{Gl}(\mathcal{M}_{\mathbb{Q}})$$

a rational representation. This representation  $r$  provides a sheaf  $\tilde{\mathcal{M}}$  on  $\mathcal{S}_{K_f}^G$  whose sections on an open subset  $V \subset \mathcal{S}_{K_f}^G$  are given by

$$\tilde{\mathcal{M}}_{\mathbb{Q}}(V) = \{s : \pi^{-1}(V) \rightarrow \tilde{\mathcal{M}}_{\mathbb{Q}} | s \text{ locally constant and } s(\gamma v) = r(\gamma)s(v), \gamma \in G(\mathbb{Q})\}.$$

We call this the *right module description* of  $\tilde{\mathcal{M}}_{\mathbb{Q}}$ .

We can describe the stalk of the sheaf in a point  $y \in \mathcal{S}_{K_f}^G$ , we choose a point  $\underline{x} = (x_{\infty}, \underline{g}_f)$  in  $\pi^{-1}(y)$  and we choose a neighborhood  $V_y$  as in 1.2.1. Then we can evaluate an element  $s \in \tilde{\mathcal{M}}_{\mathbb{Q}}(V_y)$  at  $\underline{x}$  and this must be an element in  $\mathcal{M}_{\mathbb{Q}}^{\Gamma_{x_{\infty}}}$ , this means we get an isomorphism

$$e_{\underline{x}} : (\tilde{\mathcal{M}}_{\mathbb{Q}})_y \xrightarrow{\sim} \mathcal{M}_{\mathbb{Q}}^{\Gamma_{x_{\infty}}}.$$

By definition we have  $e_{\gamma \underline{x}} = \gamma e_{\underline{x}}$ .

In our previous example such a representation  $r$  is of the following form: We take the homogeneous polynomials  $P(X, Y)$  of degree  $n$  in two variables and with coefficients in  $\mathbb{Q}$ . This is a  $\mathbb{Q}$ -vector space of dimension  $n + 1$ , we choose another integer  $m$  and now we define an action of  $\text{Gl}_2/\mathbb{Q}$  on this vector space

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} P(X, Y) = P(aX + cY, bX + dY) \det\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}\right)^m.$$

This  $\text{Gl}_2$  module will be called  $\mathcal{M}_n[m]_{\mathbb{Q}}$  and it yields sheaves  $\tilde{\mathcal{M}}_n[m]_{\mathbb{Q}}$  on our space  $\mathcal{S}_{K_f}^G$ .

It is sometimes reasonable to start from an absolutely irreducible representation and therefore we consider representations defined after a base change

$r : G \times_{\mathbb{Q}} F \rightarrow \mathrm{Gl}(\mathcal{M}_F)$  where  $\mathcal{M}_F$  is a finite dimensional  $F$  vector space and the action is absolutely irreducible. Since  $G(\mathbb{Q})$  acts on  $\mathcal{M}_F$  we can define a sheaf  $\tilde{\mathcal{M}}_F$  of  $F$  vector spaces. In our notation we stick to  $\tilde{\mathcal{M}}_{\mathbb{Q}}$ , and keep in mind that  $\mathcal{M}_{\mathbb{Q}}$  may also be a vector space over an algebraic extension  $F/\mathbb{Q}$ .

If our group is a torus  $T/\mathbb{Q}$ , then we can find a finite normal extension  $E/\mathbb{Q}$  such that  $T \times_{\mathbb{Q}} E$  is split and then we denote by

$$X^*(T) = \mathrm{Hom}(T \times E, \mathbb{G}_m) \quad \text{resp} \quad X_*(T) = \mathrm{Hom}(\mathbb{G}_m, T \times_{\mathbb{Q}} E) \quad (6.17)$$

the character (resp. ) cocharacter module of  $T/\mathbb{Q}$ . Both modules come with an action of the Galois group  $\mathrm{Gal}(E/\mathbb{Q})$ . In this case an absolutely irreducible representation is simply a character  $\gamma \in X^*(T)$  and we denote by  $E[\gamma]$  a one dimensional  $E$ -vector space on which  $T/\mathbb{Q}$  acts by  $\gamma$ . Then  $E[\tilde{\gamma}]$  is a sheaf of  $F$ -vector spaces on  $S_{K_f}^T$ .

### Integral coefficient systems

We assume again that we have a rational representation of our group  $G/\mathbb{Q}$ , the following considerations easily generalize to the case of an arbitrary number field as base field. We want to define a subsheaf  $\tilde{\mathcal{M}}_{\mathbb{Z}} \subset \tilde{\mathcal{M}}_{\mathbb{Q}}$ . To do this we embed the field  $\mathbb{Q} \hookrightarrow \mathbb{A}_f$  and we consider the resulting sheaf of  $\mathbb{A}_f$ -modules  $\tilde{\mathcal{M}} \otimes \mathbb{A}_f$ . We consider the diagram

$$\begin{array}{ccc}
 & G(\mathbb{R})/K_{\infty} \times (G(\mathbb{A}_f)/K_f) & \\
 \nearrow \pi' & & \searrow \pi \\
 G(\mathbb{R})/K_{\infty} \times G(\mathbb{A}_f) & \xrightarrow{\quad \Pi \quad} & S_{K_f}^G \\
 \searrow \Pi_1 & & \nearrow \Pi_2 \\
 & G(\mathbb{Q}) \backslash G(\mathbb{R})/K_{\infty} \times G(\mathbb{A}_f) & 
 \end{array} \quad (6.18)$$

this means that the division by the action by  $K_f$  on the right and by  $G(\mathbb{Q})$  on the left (this gives  $\Pi$ ) is divided into two steps: In the lower diagram the projection  $\Pi_1$  is division by the action of  $G(\mathbb{Q})$  and then  $\Pi_2$  gives the division by the action of  $K_f$  on the right.

The sheaf  $\tilde{\mathcal{M}}_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{A}_f$  can be rewritten. For any open subset  $V \subset S_{K_f}^G$  we consider  $W = \Pi^{-1}(V)$  and by definition

$$\tilde{\mathcal{M}}_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{A}_f(V) = \{s : \Pi^{-1}(W) \rightarrow \mathcal{M}_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{A}_f \mid s(\gamma(x_{\infty}, \underline{g}_f k_f)) = \gamma(s(x_{\infty}, \underline{g}_f))\},$$

where these sections  $s$  are locally constant in the variable  $x_{\infty}$ . For any  $s \in \mathcal{M} \otimes \mathbb{A}_f(V)$  we define a map  $\tilde{s} : W \rightarrow \mathcal{M} \otimes \mathbb{A}_f$  by the formula

$$\tilde{s}(x_{\infty}, \underline{g}_f) = \underline{g}_f^{-1} s(x_{\infty}, \underline{g}_f K_f),$$

this makes sense because  $\mathcal{M} \otimes \mathbb{A}_f$  is a  $G(\mathbb{A}_f)$ -module. For  $\gamma \in G(\mathbb{Q})$  we have  $\tilde{s}(\gamma(x_\infty, \underline{g}_f)) = \tilde{s}((x_\infty, \underline{g}_f))$  hence we can view  $\tilde{s}$  as a map

$$\tilde{s} : G(\mathbb{Q}) \backslash G(\mathbb{R}) / K_\infty \times G(\mathbb{A}_f) \rightarrow \mathcal{M} \otimes_{\mathbb{Q}} \mathbb{A}_f.$$

We consider the projection

$$\Pi_2 : G(\mathbb{Q}) \backslash G(\mathbb{R}) / K_\infty \times G(\mathbb{A}_f) \rightarrow G(\mathbb{Q}) \backslash G(\mathbb{R}) / K_\infty \times G(\mathbb{A}_f) / K_f = \mathcal{S}_{K_f}^G$$

and then it becomes clear that  $\widetilde{\mathcal{M}} \otimes \mathbb{A}_f$  can be described as

$$\begin{aligned} \widetilde{\mathcal{M}} \otimes \mathbb{A}_f(V) &= \{ \tilde{s} : (\Pi_1^{-1}(V) \rightarrow \mathcal{M} \otimes_{\mathbb{Q}} \mathbb{A}_f | \\ &\tilde{s} \text{ locally constant in } x_\infty \text{ and } \tilde{s}((x_\infty, \underline{g}_f k_f)) = \underline{k}_f^{-1} \tilde{s}((x_\infty, \underline{g}_f)) \}. \end{aligned}$$

Hence we have identified the sheaf  $\widetilde{\mathcal{M}} \otimes_{\mathbb{Q}} \mathbb{A}_f$  which is defined in terms of the action of  $G(\mathbb{Q})$  on  $\mathcal{M}$  to the sheaf  $\widetilde{\mathcal{M}} \otimes_{\mathbb{Q}} \mathbb{A}_f$  which is defined in terms of the action of  $K_f$  on  $\mathcal{M} \otimes_{\mathbb{Q}} \mathbb{A}_f$ .

Now we assume that our group scheme  $G/\mathbb{Q}$  is the generic fiber of a flat group scheme  $\mathcal{G}/\text{Spec}(\mathbb{Z})$  (See 1.2). We choose our maximal compact subgroup  $K_f = \prod_p K_p$  such that  $K_p \subset \mathbb{G}(\mathbb{Z}_p)$  and with equality for all primes outside a finite set  $\Sigma$ . We can extend the vector space  $\mathcal{M}$  to a free  $\mathbb{Z}$  module  $\tilde{\mathcal{M}}_{\mathbb{Z}}$  of the same rank which provides a representation  $\mathbb{G}/\text{Spec}(\mathbb{Z}) \rightarrow \text{Gl}(\tilde{\mathcal{M}}_{\mathbb{Z}})$ .

As usual  $\hat{\mathbb{Z}}$  will be the ring of integral adeles. Then it is clear that  $\mathcal{M}_{\mathbb{Z}} \otimes \hat{\mathbb{Z}} \subset \mathcal{M} \otimes_{\mathbb{Q}} \mathbb{A}_f$  is invariant under  $K_f$  and hence we can define the sub sheaf

$$\widetilde{\mathcal{M}}_{\mathbb{Z}} \otimes \hat{\mathbb{Z}} \subset \widetilde{\mathcal{M}} \otimes_{\mathbb{Q}} \mathbb{A}_f,$$

this is the sheave where the sections  $\tilde{s}$  have values in  $\mathcal{M}_{\mathbb{Z}} \otimes \hat{\mathbb{Z}}$ . We put

$$\tilde{\mathcal{M}}_{\mathbb{Z}} = \widetilde{\mathcal{M}}_{\mathbb{Z}} \otimes \hat{\mathbb{Z}} \cap \widetilde{\mathcal{M}},$$

of course it depends on our choice of  $\mathcal{M}_{\mathbb{Z}} \subset \mathcal{M}$ . We get two exact sequences of sheaves

$$\begin{array}{ccccccc} 0 & \rightarrow & \tilde{\mathcal{M}}_{\mathbb{Z}} & \rightarrow & \tilde{\mathcal{M}} & \rightarrow & \widetilde{\mathcal{M}} \otimes (\mathbb{Q}/\mathbb{Z}) \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \rightarrow & \widetilde{\mathcal{M}}_{\mathbb{Z}} \otimes \hat{\mathbb{Z}} & \rightarrow & \widetilde{\mathcal{M}} \otimes_{\mathbb{Q}} \mathbb{A}_f & \rightarrow & \widetilde{\mathcal{M}} \otimes (\mathbb{A}_f/\hat{\mathbb{Z}}) \rightarrow 0 \end{array}$$

The far most vertical arrow to the right is an isomorphism, the inclusions  $\mathbb{Z} \hookrightarrow \hat{\mathbb{Z}}$  and  $\mathbb{Q} \hookrightarrow \mathbb{A}_f$  are flat. Writing down the resulting long exact sequences provides a diagram

$$\begin{array}{ccccc} \rightarrow & H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Z}}) & \xrightarrow{j_{\mathbb{Q}}} & H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}) & \rightarrow \\ & \downarrow i_{\mathbb{Z}} & & \downarrow i_{\mathbb{Q}} & \\ \rightarrow & H^\bullet(\mathcal{S}_{K_f}^G, \widetilde{\mathcal{M}}_{\mathbb{Z}} \otimes \hat{\mathbb{Z}}) & \xrightarrow{j_{\mathbb{A}}} & H^\bullet(\mathcal{S}_{K_f}^G, \widetilde{\mathcal{M}} \otimes_{\mathbb{Q}} \mathbb{A}_f) & \rightarrow \end{array}$$

The above remarks imply that the vertical arrows are injective, the horizontal arrows in the middle have the same kernel and cokernel. This implies

**Proposition 6.2.1.** *The integral cohomology*

$$H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Z}})$$

consists of those elements in  $H^\bullet(\mathcal{S}_{K_f}^G, \widetilde{\mathcal{M} \otimes \hat{\mathbb{Z}}})$  which under  $j_{\mathbb{A}}$  go to an element in the image under  $i_{\mathbb{Q}}$  or in brief

$$H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Z}}) = j_{\mathbb{A}}^{-1}(\text{im}(i_{\mathbb{Q}}))$$

This generalizes to the case where we have a representation  $r : G \times F \rightarrow \text{Gl}(\mathcal{M})$  where  $\mathcal{M}$  is a vector space over  $F$ . If our group scheme is an extension of a flat group scheme  $\mathcal{G}/\text{Spec}(\mathcal{O}_F)$  then can find a lattice  $\mathcal{M}_{\mathcal{O}_F}$  which yields a representation of  $\mathcal{G} \rightarrow \text{Gl}(\mathcal{M}_{\mathcal{O}_F})$ . Then we can define the sheaf  $\tilde{\mathcal{M}}_{\mathcal{O}_F}$  and define the cohomology groups

$$H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathcal{O}_F})$$

### Sheaves with support conditions

We can extend the sheaves to the Borel-Serre compactification. We have the inclusion

$$i : \mathcal{S}_{K_f}^G \rightarrow \bar{\mathcal{S}}_{K_f}^G$$

and we can extend the sheaf by the direct image functor  $i_*(\tilde{\mathcal{M}})$ . It follows easily from the description of the neighborhood of a point in the boundary (see 6.16) that  $R^q i_*(\tilde{\mathcal{M}}) = 0$  for  $q = 0$  and hence we get that the restriction map

$$H^\bullet(\bar{\mathcal{S}}_{K_f}^G, i_*(\tilde{\mathcal{M}})) \rightarrow H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})$$

is an isomorphism.

We may also extend the sheaf by zero (See [Vol I], 4.7.1), this yields the sheaf  $i_!(\tilde{\mathcal{M}})$  whose stalk at  $x \in \mathcal{S}_{K_f}^G$  is equal to  $\tilde{\mathcal{M}}_x$  and whose stalk is zero in points  $x \in \partial\mathcal{S}_{K_f}^G$ . Then we have by definition

$$H_c^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}) = H^\bullet(\bar{\mathcal{S}}_{K_f}^G, i_!(\tilde{\mathcal{M}}))$$

this is the cohomology with compact supports.

We are interested in the *integral* cohomology modules  $H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Z}})$ ,  $H_c^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Z}})$ . We introduced the boundary  $\partial\mathcal{S}_{K_f}^G$  of the Borel-Serre compactification then we have a first general theorem, which is due to Raghunathan.

**Theorem 6.2.1.** (i) *The cohomology groups  $H^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Z}})$ ,  $H^i(\partial\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Z}})$  and  $H_c^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Z}})$  are finitely generated.*

(ii) *We have the well known fundamental long exact sequence in cohomology*

$$\rightarrow H^{i-1}(\partial\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Z}}) \rightarrow H_c^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Z}}) \rightarrow H^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Z}}) \xrightarrow{r} H^i(\partial\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Z}}) \rightarrow .$$

We introduce the notation  $H_?( \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Z}})$  meaning that for  $? = \text{blank}$  we take the cohomology without support, for  $? = c$  we take the cohomology with compact support and for  $? = \partial$  we take cohomology of the boundary of the

Borel-Serre compactification. Later on we will also allow  $? = !$  this denotes the inner cohomology. The above proposition 6.2.1 holds for all choices of  $?$ .

Let  $\Sigma = \{P_1, \dots, P_s\}$  be a finite set of parabolic subgroups, we assume that none of them is a subgroup of another parabolic subgroup in this set. The union of the closures of the strata

$$\bigcup_i \bigcup_{Q \subset P_i} \partial_Q(\mathcal{S}_{K_f}^G) = \partial_\Sigma(\mathcal{S}_{K_f}^G)$$

is closed . We have the inclusions

$$j_\Sigma : \mathcal{S}_{K_f}^G \rightarrow \bar{\mathcal{S}}_{K_f}^G \setminus \partial_\Sigma(\bar{\mathcal{S}}_{K_f}^G), j^\Sigma : \bar{\mathcal{S}}_{K_f}^G \setminus \partial_\Sigma(\bar{\mathcal{S}}_{K_f}^G) \rightarrow \bar{\mathcal{S}}_{K_f}^G.$$

The inclusion  $i : \mathcal{S}_{K_f}^G \rightarrow \bar{\mathcal{S}}_{K_f}^G$  is the composition  $i = j^\Sigma \circ j_\Sigma$  we define the intermediate extension suppcond

$$i_{\Sigma, *, !}(\tilde{\mathcal{M}}) = j_{!, \Sigma}^\Sigma \circ j_{\Sigma, *}(\tilde{\mathcal{M}}), \tag{6.19}$$

this means that the stalk  $i_{\Sigma, *, !}(\tilde{\mathcal{M}})_y$  at a point  $y \in \partial_\Sigma(\bar{\mathcal{S}}_{K_f}^G)$  is zero. Now we can define the cohomology with supports  $H^\bullet(\mathcal{S}_{K_f}^G, i_{\Sigma, *, !}(\tilde{\mathcal{M}}))$ . If  $\Sigma = \emptyset$  then  $H^\bullet(\Sigma, *, !(\tilde{\mathcal{M}})_y) = H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})$  and if  $\Sigma$  is the set of all maximal parabolic subgroups then  $H^\bullet(\Sigma, *, !(\tilde{\mathcal{M}})_y) = H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})$ .

For these cohomology groups coefficients in sheaves with intermediate support conditions we can also formulate assertion like the one in the above theorem. Hence we get filtrations on the cohomology

$$W_0 H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}) = H_{!, \Sigma}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}) \subset W_1 H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}) \subset \dots \subset H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}) \tag{6.20}$$

on the cohomology, the bottom of this filtration will be the inner cohomology and the filtration steps will be the images cohomology with intermediate supports.

**Functorial properties**

The groups have some functorial properties if we vary the level subgroup  $K_f$ . If we pass to a smaller open subgroup  $K'_f \subset K_f$  then we get a surjective map

$$\pi_{K_f, K'_f} : \mathcal{S}_{K'_f}^G \rightarrow \mathcal{S}_{K_f}^G,$$

whose fibers are finite. This induces maps between cohomology groups

$$\pi_{K'_f, K_f ?}^\bullet : H_{?, \Sigma}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Z}}) \rightarrow H_{?, \Sigma}^\bullet(\mathcal{S}_{K'_f}^G, \tilde{\mathcal{M}}_{\mathbb{Z}}),$$

for  $? = c$  we exploit the fact that the fibers are finite.

We construct homomorphisms in the opposite direction. We exploit the finiteness a second time and find that the direct image functor  $(\pi_{K'_f, K_f}^G)_*$  is exact and hence

$$H_{?, \Sigma}^\bullet(\mathcal{S}_{K'_f}^G, \tilde{\mathcal{M}}_{\mathbb{Z}}) = H_{?, \Sigma}^\bullet(\mathcal{S}_{K_f}^G, (\pi_{K'_f, K_f}^G)_*(\tilde{\mathcal{M}}_{\mathbb{Z}})).$$

We define a trace homomorphism  $(\pi_{K'_f, K_f})_*(\tilde{\mathcal{M}}_{\mathbb{Z}}) \rightarrow \tilde{\mathcal{M}}_{\mathbb{Z}}$ : A section  $s \in (\pi_{K'_f, K_f})_*(\tilde{\mathcal{M}}_{\mathbb{Z}})(V)$  is a map  $\tilde{s} : \Pi^{-1}(V) \rightarrow \tilde{\mathcal{M}}_{\lambda} \otimes \hat{\mathbb{Z}}$  such that

$$\tilde{s}(\gamma(x_{\infty}, \underline{g}_f k'_f)) = (k'_f)^{-1} \tilde{s}((x_{\infty}, \underline{g}_f)) \text{ for all } k'_f \in K'_f.$$

This is a section of  $\tilde{\mathcal{M}}_{\mathbb{Z}}$  if and only if the corresponding section  $s$  takes values in  $\mathcal{M}$ . Then we define

$$\text{tr}(\tilde{s})(x_{\infty}, \underline{g}_f) = \sum_{\xi_f \in K_f/K'_f} \xi_f^{-1} \tilde{s}(x_{\infty}, \underline{g}_f)$$

and this now satisfies

$$\text{tr}(\tilde{s})(\gamma(x_{\infty}, \underline{g}_f k_f)) = k_f^{-1} \text{tr}(\tilde{s})(x_{\infty}, \underline{g}_f) \text{ for all } k_f \in K_f.$$

and since the corresponding section  $\text{tr}(s)$  takes values in  $\mathcal{M}$  we see that  $\text{tr}(\tilde{s}) \in \tilde{\mathcal{M}}_{\mathbb{Z}}(V)$ .

Remark: It may happen that this trace map is not the optimal choice, it can be the integral multiple of another homomorphism between these two sheaves. This happens the intersection  $C(\mathbb{Q}) \cap K_f$  is non trivial.

Then the homomorphism between the sheaves induces

$$H_{\mathbb{Z}}^{\bullet}(\mathcal{S}_{K'_f}^G, \tilde{\mathcal{M}}_{\mathbb{Z}}) = H_{\mathbb{Z}}^{\bullet}(\mathcal{S}_{K'_f}^G, (\pi_{K'_f, K_f})_*(\tilde{\mathcal{M}}_{\mathbb{Z}})) \xrightarrow{\pi_{K'_f, K_f}} H_{\mathbb{Z}}^{\bullet}(\mathcal{S}_{K_f}^G, (\tilde{\mathcal{M}}_{\mathbb{Z}})).$$

Later on our maps between the spaces will be denoted  $\pi, \pi_1, \dots$  and the notation simplifies accordingly. Heckalg

## 6.3 The action of the Hecke-algebra

### 6.3.1 The action on rational cohomology

In this section we assume that our coefficient systems are obtained from rational representations of a reductive group scheme  $G/\mathbb{Q}$  hence they are  $\mathbb{Q}$  vector spaces. We consider the *rational* cohomology groups

$$H_{\mathbb{Z}}^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Q}}) = H_c^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Q}}), H^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Q}}), H^i(\partial(\mathcal{S}_{K_f}^G), \tilde{\mathcal{M}}_{\mathbb{Q}}),$$

These cohomology groups are finite dimensional  $\mathbb{Q}$ -vector spaces and they are related exact fundamental sequence. We can pass to the direct limit

$$H_{\mathbb{Z}}^i(\mathcal{S}^G, \tilde{\mathcal{M}}_{\mathbb{Q}}) = \lim_{K_f} H_{\mathbb{Z}}^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Q}}).$$

GAF

**Proposition 6.3.1.** *On these limits we have an action of the group  $\pi_0(G(\mathbb{R})) \times G(\mathbb{A}_f)$ . We recover the cohomology with fixed level  $K_f$  by taking the inva, under this action, i.e. we have*

$$H_{\mathbb{Z}}^i(\mathcal{S}^G, \tilde{\mathcal{M}}_{\mathbb{Q}})^{K_f} = H_{\mathbb{Z}}^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Q}})$$

To define this action we represent an element in  $\pi_0(G(\mathbb{R}))$  by an element  $k_\infty$  in the normalizer of  $K_\infty$  in  $G(\mathbb{R})$ . An element  $\underline{x} = (k_\infty, \underline{x}_f) \in G(\mathbb{R}) \times G(\mathbb{A}_f)$  defines by multiplication from the right an isomorphism of spaces

$$m_{\underline{x}} : G(\mathbb{Q}) \backslash X \times G(\mathbb{A}_f) / K_f \xrightarrow{\sim} G(\mathbb{Q}) \backslash X \times G(\mathbb{A}_f) / \underline{x}_f^{-1} K_f \underline{x}_f.$$

It is clear from the definition that  $m_{\underline{x}}$  yields a bijection between the fibers  $\pi^{-1}(\underline{g}), \underline{g} \in G(\mathbb{Q}) \backslash X \times G(\mathbb{A}_f) / K_f$  and  $\pi^{-1}(m_{\underline{x}}(\underline{g}))$  and since the sheaf is described in terms of the left action by  $G(\mathbb{Q})$  we get  $m_{\underline{x},*}(\tilde{\mathcal{M}}) = \tilde{\mathcal{M}}$ . Passing to the limit gives us the action on  $H^i_\mathbb{Q}(\mathcal{S}^G, \tilde{\mathcal{M}}_\mathbb{Q})$ . The second assertion is obvious, but here we need that our coefficients are  $\mathbb{Q}$  vector spaces, we need to take averages.

We introduce the notation  $\pi_0(G(\mathbb{R})) \times (\mathbb{A}_f) := \tilde{G}(\mathbb{A})$  and then we denote this action by

$$\rho_{\tilde{\mathcal{M}}_\mathbb{Q}} : \tilde{G}(\mathbb{A}) \times H^i_\mathbb{Q}(\mathcal{S}^G_{K_f}, \tilde{\mathcal{M}}_\mathbb{Q}) \rightarrow H^i_\mathbb{Q}(\mathcal{S}^G_{K_f}, \tilde{\mathcal{M}}_\mathbb{Q}).$$

The interesting component of this representation is of course the action of the finite component  $G(\mathbb{A}_f)$ , it is simply the action which is induced by the right translation action of  $G(\mathbb{A}_f)$  on  $\mathcal{S}^G$ .

Now we fix a level  $K_f \subset G(\mathbb{A}_f)$  The Hecke algebra  $\mathcal{H}_{K_f}$  consists of the compactly supported functions  $h : G(\mathbb{A}_f) \rightarrow \mathbb{Q}$ , which are biinvariant under the action of  $K_f$ , we also write  $\mathcal{H}_{K_f} = \mathcal{C}_c(G(\mathbb{A}_f) // K_f, \mathbb{Q})$ . An element  $h \in \mathcal{H}_{K_f}$  is simply a finite linear combination of characteristic functions  $h = \sum c_{\underline{a}_f} \chi_{K_f \underline{a}_f K_f}$  with rational coefficients  $c_{\underline{a}_f}$ . The algebra structure is given by convolution with respect to the Haar measure on  $G(\mathbb{A}_f)$  which gives volume 1 to  $K_f$ . This convolution is given by

$$h_1 * h_2(\underline{g}_f) = \int_{G(\mathbb{A}_f)} h_1(\underline{x}_f) h_2(\underline{x}_f^{-1} \underline{g}_f) d\underline{x}_f.$$

With this choice of the measure it is clear that the characteristic function of  $K_f$  is the identity element of this algebra.

The action of the group  $G(\mathbb{A}_f)$  induces an action of  $\mathcal{H}_{K_f}$  on the cohomology with fixed level  $H^i_c(\mathcal{S}^G_{K_f}, \tilde{\mathcal{M}}), H^i(\mathcal{S}^G_{K_f}, \tilde{\mathcal{M}}), \dots$ : For an element  $v \in H^i_\mathbb{Q}(\mathcal{S}^G, \tilde{\mathcal{M}})$  we define

$$T_h(v) = \int_{G(\mathbb{A}_f)} h(\underline{x}_f) \underline{x}_f v d\underline{x}_f,$$

where the measure is still the one that gives volume 1 to  $K_f$ . Clearly we have  $T_{h_1 * h_2} = T_{h_1} T_{h_2}$ .

(Actually the integral is a finite sum: We find an open subgroup  $K'_f \subset K_f$  such that  $v$  is fixed by  $K'_f$  and then it is clear that

$$T_h(v) = \int_{G(\mathbb{A}_f)} h(\underline{x}_f) \underline{x}_f v d\underline{x}_f = \frac{1}{[K_f : K'_f]} \sum_{\underline{a}_f} \sum_{\underline{\xi}_f \in G(\mathbb{A}_f) / K'_f} c_{\underline{a}_f} \chi_{K_f \underline{a}_f K_f}(\underline{\xi}_f) \underline{\xi}_f v.$$

This makes it clear why we need rational coefficients .)

It is clear that  $T_h(v) \in H^i_\mathbb{Q}(\mathcal{S}^G_{K_f}, \tilde{\mathcal{M}})$  and hence  $T_h$  gives us an endomorphism of  $H^i_\mathbb{Q}(\mathcal{S}^G_{K_f}, \tilde{\mathcal{M}})$ . We will show later that we also get endomorphisms on the

cohomology of the boundary and therefore  $\mathcal{H}$  also acts on the fundamental long exact sequence (Seq).

If our function  $h$  is the characteristic function of a double coset  $K_f \underline{x}_f K_f$  then we change notation and write  $T_h = \mathbf{ch}(\underline{x}_f)$ . We give another definition of the Hecke operator  $\mathbf{ch}(\underline{x}_f)$  in terms of sheaf cohomology. We imitate the construction of the Hecke operators in Chapter 3, 3.1. We put  $K_f^{(\underline{x}_f)} = K_f \cap \underline{x}_f K_f \underline{x}_f^{-1}$  and consider the diagram

$$\begin{array}{ccc} S_{K_f^{(\underline{x}_f)}}^G & \xrightarrow{m_{\underline{x}_f}} & S_{K_f^{(\underline{x}_f^{-1})}}^G \\ \searrow \pi_1 & & \swarrow \pi_2 \\ & S_{K_f}^G & \end{array} \quad \text{Hop1}$$

where  $m_{\underline{x}_f}$  is induced by the multiplication by  $\underline{x}_f$  from the right. This yields in cohomology

$$H^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}) \xrightarrow{\pi_1^\bullet} H^\bullet(S_{K_f^{(\underline{x}_f)}}^G, \tilde{\mathcal{M}}) \xrightarrow{m_{\underline{x}_f, *}} H^\bullet(S_{K_f^{(\underline{x}_f^{-1})}}^G, m_{\underline{x}_f, *}(\tilde{\mathcal{M}})) \quad (\text{Hop2}).$$

Since we described the sheaf by the action of  $G(\mathbb{Q})$  and the map  $m_{\underline{x}_f}$  by multiplication from the right we have  $m_{\underline{x}_f, *}(\tilde{\mathcal{M}}) = \tilde{\mathcal{M}}$ , this yields an isomorphism  $i_{\underline{x}_f}$ . Since  $\pi_2$  is finite we have the trace homomorphism

$$\pi_{2, \bullet} : H^\bullet(S_{K_f^{(\underline{x}_f^{-1})}}^G, \tilde{\mathcal{M}}) \rightarrow H^\bullet(S_{K_f}^G, \tilde{\mathcal{M}})$$

and the composition is our Hecke operator

$$\pi_{2, \bullet} \circ i_{\underline{x}_f} \circ m_{\underline{x}_f, *} \circ \pi_1^\bullet = \mathbf{ch}(\underline{x}_f) : H^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}) \rightarrow H^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}).$$

This is simpler than the construction Chap.II 2.2. because we do not need the intermediate homomorphism  $u_\alpha$ . But we do not get Hecke operators on the integral cohomology.

### 6.3.2 The integral cohomology as a module under the Hecke algebra

We resume the discussion of the integral Hecke algebra acting on  $H^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Z}})$  from Chapter 3. Inside the Hecke algebra we may also look at the sub algebra of  $\mathbb{Z}$ -valued functions. This is in principle the algebra which is generated by the characteristic functions  $\mathbf{ch}(\underline{x}_f)$  of double cosets  $K_f \underline{x}_f K_f$ . These characteristic functions act by convolution on the cohomology  $H^\bullet(S_{K_f}^G, \mathcal{M})$  but this does not induce an action on the integral cohomology. Our next aim is to define a fractional ideal  $\mathfrak{n}(\underline{x}_f) \subset \mathbb{Q}$  or more generally  $\mathfrak{n}(\underline{x}_f) \subset F$  such that for any  $a \in \mathfrak{n}(\underline{x}_f)$  we can define an endomorphism

$$a \cdot \mathbf{ch}(\underline{x}_f) : H^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \rightarrow H^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$$

and if we send this to the rational cohomology then on  $H^\bullet(S_{K_f}^G, \mathcal{M})$  this will be the convolution endomorphism induced by  $\mathbf{ch}(\underline{x}_f)$  multiplied by  $a$ .

This ideal will depend on  $\underline{x}_f$  and on  $\lambda$  and further down we compute it in special cases.

(iv) These endomorphisms  $a \cdot \mathbf{ch}(\underline{x}_f)$  generate an algebra  $\mathcal{H}_{\mathbb{Z}}^{(\lambda)}$  acting on the integral cohomology and the arrows in the fundamental exact sequence above commute with this action.

v) Moreover, we have an action of  $\pi_0(G(\mathbb{R}))$  on the above sequence and this action also commutes with the action of the Hecke algebra. Hence we know that our above sequence is long exact sequence of  $\pi_0(G(\mathbb{R})) \times \mathcal{H}_{\mathbb{Z}}^{(\lambda)}$ .

We come to the definition of the ideal.

If we are in the special case that our group has strong approximation then we have

$$\Gamma \backslash X \xrightarrow{\sim} G(\mathbb{Q}) \backslash X \times G(\mathbb{A}_f) / K_f$$

(See (6.10)). We pick an element  $\alpha \in G(\mathbb{Q})$ . In Chap. 3, 3.1. we defined the Hecke operator  $T(\alpha, u_\alpha)$  where  $u_\alpha : \mathcal{M}^{(\alpha)} \rightarrow \mathcal{M}$  is the canonical choice. Let us denote the image of  $\alpha$  in  $G(\mathbb{A}_f)$  by  $\underline{\alpha}_f$ . We just attached a Hecke operator to the double coset  $K_f \underline{\alpha}_f \cdot K_f$ . We have the diagram of spaces

$$\begin{array}{ccc} \Gamma(\alpha) \backslash X & \xrightarrow{\quad\quad\quad} & G(\mathbb{Q}) \backslash G(\mathbb{R}) / K_\infty \times G(\mathbb{A}_f) / K_f^{\underline{\alpha}_f} & (6.21) \\ \downarrow l(\alpha^{-1}) & & \downarrow r(\underline{\alpha}_f) & \\ \Gamma(\alpha^{-1}) \backslash X & \xrightarrow{\quad\quad\quad} & G(\mathbb{Q}) \backslash G(\mathbb{R}) / K_\infty \times G(\mathbb{A}_f) / K_f^{\underline{\alpha}_f^{-1}} & \end{array}$$

Here the horizontal arrows are the isomorphisms provided by strong approximation, the arrow  $l(\alpha^{-1})$  is the isomorphism induced by left multiplication by  $\alpha^{-1}$  and  $r(\underline{\alpha}_f)$  by right multiplication by  $\underline{\alpha}_f$ . These two maps enter in the definition of the Hecke operators  $T(\alpha^{-1}, u_{\alpha^{-1}})$  and  $\mathbf{ch}(\underline{\alpha}_f)$  and a straightforward inspection of the sheaves yields

$$\mathbf{ch}(\underline{\alpha}_f) = T(\alpha^{-1}, u_{\alpha^{-1}}).$$

Hence we can conclude that under this assumption our newly defined Hecke operators coincide with the Hecke operators defined in Chap.3. This also tells us what we have to do if we want to define Hecke operators on integral cohomology.

To define the action of the Hecke algebra on the integral cohomology without the assumption of simple connectedness we have to translate their definition into the right module description. Then our sheaf  $\widehat{\mathcal{M}} \otimes \widehat{\mathbb{A}}_f$  is described by the action of  $K_f$  on  $\mathcal{M} \otimes \mathbb{A}_f$  and this allows us to define the sub sheaf  $\widehat{\mathcal{M}}_{\mathbb{Z}} \otimes \widehat{\mathbb{Z}}$ . We look at the same diagram. But now the sheaf  $m_{\underline{x}_f, *}(\widehat{\mathcal{M}} \otimes \widehat{\mathbb{A}}_f)$  is the sheaf described by the the  $K_f^{(\underline{x}_f)^{-1}}$  module  $(\mathcal{M} \otimes \mathbb{A}_f)^{(\underline{x}_f)}$ . This module is  $\mathcal{M} \otimes \mathbb{A}_f$  as abelian group, but  $\underline{g}_f \in K_f^{(\underline{x}_f)^{-1}}$  acts by  $\underline{m}_f \mapsto \underline{x}_f \underline{g}_f \underline{x}_f^{-1} \underline{m}_f$ . The map  $\underline{m}_f \rightarrow \underline{x}_f \underline{m}_f$

induces an isomorphism  $[\underline{x}_f]$  between the two  $K_f^{(\underline{x}_f)^{-1}}$  modules  $(\mathcal{M} \otimes \mathbb{A}_f)^{(\underline{x}_f)}$  and  $(\mathcal{M} \otimes \mathbb{A}_f)$ . We now consider the diagram *Hop1.* and replace in the sequence of maps the homomorphism  $i_{\underline{x}_f}$  by the map  $[\underline{x}_f]$  induced by the isomorphism  $[\underline{x}_f]$  between the sheaves. Then we can proceed as before and get an operator

$$p_{1,*} \circ [\underline{x}_f]^\bullet \circ m_{\underline{x}_f,*} \circ p_2^* = \mathbf{ch}(\underline{x}_f).$$

It is straightforward to check that this operator is an extension  $\pi_{2,\bullet} \circ i_{\underline{x}_f} \circ m_{\underline{x}_f,*} \circ \pi_1^\bullet$  to  $H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}} \otimes \mathbb{A}_f)$ .

Our right module sheaf contains the submodule sheaf  $\tilde{\mathcal{M}}_\lambda \otimes \hat{\mathbb{Z}}$ , we can write the same diagram but now it can happen that  $[\underline{x}_f]$  does not map  $\mathcal{M}_{\mathbb{Z}} \otimes \hat{\mathbb{Z}}$  into itself. This forces us to make the following definition

$$\mathfrak{n}(\underline{x}_f) = \{a \in \mathbb{Q} \mid [a\underline{x}_f] : \mathcal{M}_{\mathbb{Z}} \otimes \hat{\mathbb{Z}} \subset \mathcal{M}_{\mathbb{Z}} \otimes \hat{\mathbb{Z}}\}$$

Then we can again go back to our above diagram and it becomes clear that we can define Hecke operators

$$a \cdot \mathbf{ch}(\underline{x}_f) : H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Z}}) \rightarrow H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Z}}) \text{ for all } a \in \mathfrak{n}(\underline{x}_f).$$

### The case of a split group

We want to discuss this in the special case that  $\mathcal{G}/\mathrm{Spec}(\mathbb{Z})$  is split reductive, we assume that the derived group  $\mathcal{G}^{(1)}/\mathrm{Spec}(\mathbb{Z})$  is simply connected, we assume that the center  $\mathcal{C}/\mathrm{Spec}(\mathbb{Z})$  is a (split)-torus and that  $\mathcal{C} \cap \mathcal{G}^{(1)}$  is equal to the center  $Z^{(1)}$  of  $\mathcal{G}^{(1)}$ . This center is a finite multiplicative group scheme (See 6.1.1).

Accordingly we get decompositions up to isogeny of the character and cocharacter modules of the torus

$$X^*(\mathcal{T}) \hookrightarrow X^*(\mathcal{T}^{(1)}) \oplus X^*(\mathcal{C}) \quad X_*(\mathcal{T}^{(1)}) \oplus X_*(\mathcal{C}) \hookrightarrow X_*(\mathcal{T}) \quad (6.22)$$

they become isomorphisms after taking the tensor product by  $\mathbb{Q}$ . We numerate the simple positive roots  $I = \{1, 2, \dots, r\} = \{\alpha_1, \alpha_2, \dots, \alpha_r\} \subset X^*(\mathcal{T})$  and we define dominant fundamental weights  $\gamma_i \in X^*(\mathcal{T})_{\mathbb{Q}}$  which restricted to  $\mathcal{T}^{(1)}$  are the usual fundamental dominant weights and restricted to  $\mathcal{C}$  are trivial. Then a dominant weight can be written as

$$\lambda = \sum_{i \in I} a_i \gamma_i + \delta = \lambda^{(1)} + \delta, \quad (6.23)$$

where  $\delta \in X^*(\mathcal{C})$  and we must have the congruence condition

$$(\lambda^{(1)} + \delta)|_{Z^{(1)}} = 1 \quad (6.24)$$

We can construct a highest weight module  $\mathcal{M}_{\lambda, \mathbb{Z}}$ . We pick a prime  $p$ , we assume that is unramified (with respect to  $K_f$ ), this means that  $K_p = \mathcal{G}(\mathbb{Z}_p)$ . Any element  $t_p \in T(\mathbb{Q}_p)$  defines a double coset  $K_p t_p K_p$ . Of course only the image of  $t_p$  in  $T(\mathbb{Q}_p)/\mathcal{T}(\mathbb{Z}_p)$  matters and

$$T(\mathbb{Q}_p)/\mathcal{T}(\mathbb{Z}_p) = X_*(T)$$

we find  $\chi \in X_*(T)$  such that  $\chi(p) = t_p$ . We take a  $\chi$  in the positive chamber, i.e. we assume  $\langle \chi, \alpha \rangle \geq 0$  for all  $\alpha$ . We can produce the element

$$\underline{\chi}_p = (1 \dots, 1, \dots, \chi(p), 1 \dots, 1, \dots) \in T(\mathbb{A}_f)$$

and the Hecke operator

$$\text{ch}(\underline{\chi}_p) : H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}} \otimes \mathbb{Q}) \rightarrow H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}} \otimes \mathbb{Q})$$

We have to look at the ideal of those integers  $a$  for which

$$a \text{ch}(\underline{\chi}_p)(\mathcal{M}_{\lambda, \mathbb{Z}} \otimes \mathbb{Z}_p) \subset (\mathcal{M}_{\lambda, \mathbb{Z}} \otimes \mathbb{Z}_p).$$

This is easy: We have the decomposition into weight spaces

$$\mathcal{M}_{\lambda, \mathbb{Z}} = \bigoplus_{\mu} \mathcal{M}_{\lambda, \mathbb{Z}}(\mu)$$

and on a weight space the torus element  $\text{ch}(\underline{\chi}_p)$  acts by

$$\text{ch}(\underline{\chi}_p)x_{\mu} = p^{\langle \chi, \mu \rangle} x_{\mu}.$$

We get the smallest exponent if we choose for  $\mu$ , the lowest weight vector. We denote by  $w_0$  the longest element in the Weyl group, which sends all the positive roots into negative roots. The the element  $-w_0$  induces an involution  $i \rightarrow i'$  on the set of simple roots. Then

$$\mu = w_0(\lambda) = - \sum a_{i'} \gamma_i + \delta. \tag{6.25}$$

We say that our weight is (essentially) *self dual* if we have  $a_i = a_{i'}$ . If our weight is self dual then  $\mu = -\lambda^{(1)} + \delta$

Hence we see that our ideal is the principal ideal is given by

$$(p^{-\langle \chi, w_0 \lambda^{(1)} \rangle - \langle \chi, \delta \rangle}) \text{ or if } \lambda \text{ self dual } (p^{\langle \chi, \lambda^{(1)} \rangle - \langle \chi, \delta \rangle}) \tag{6.26}$$

and therefore, we have the Hecke operator

$$T_{p, \chi}^{\text{coh}, \lambda} = p^{-\langle \chi, w_0 \lambda^{(1)} \rangle - \langle \chi, \delta \rangle} \cdot \text{ch}(\underline{\chi}_p) : H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda, \mathbb{Z}}) \rightarrow H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda, \mathbb{Z}}) \tag{6.27}$$

The number  $-\langle \chi, w_0 \lambda^{(1)} \rangle$  is the relevant contribution in the exponent (let us call this the semi-simple term), the second term  $-\langle \chi, \delta \rangle$  is a correction term ( the abelian contribution) and it takes care of the central character. We come back to this in section 7.0.1.

GenHeck

### 6.3.3 Excursion: Finite dimensional $\mathcal{H}$ -modules and representations.

We fix a level  $K_f = \prod_p K_p$  and drop it in the notations. It follows from the theorem 6.2.1 that we have a finite Jordan-Hölder series on our cohomology groups such that the subquotients are irreducible Hecke-modules. If we take the tensor product with a suitable finite extension  $F/\mathbb{Q}$  then we can refine

the Jordan-Hölder series such that the quotients become absolutely irreducible modules for the Hecke algebra, we say a few words concerning the absolutely irreducible Hecke-modules.

We have a decomposition

$$\mathcal{H} = \bigotimes_p' \mathcal{H}_p = \bigotimes_p' \mathcal{C}_c(G(\mathbb{Q}_p)//K_p).$$

As the notation indicates we take the tensor product over all finite primes. This tensor product has to be taken in a restricted sense: for an element of the form  $h_f = \otimes h_p$  the local factor  $h_p$  is equal to the identity element  $e_p$  for almost all primes  $p$  (here  $e_p$  is the characteristic function of  $K_p$ ). All other elements are finite linear combinations of elements of the form above. We have the obvious embedding

$$\mathcal{H}_p \hookrightarrow \mathcal{H} \text{ we simply send } h_p \mapsto \otimes \dots e_p' \otimes h_p \otimes 1 \dots \quad (6.28)$$

The subalgebras  $\mathcal{H}_p$  commute with each other.

We are interested in categories of modules for the Hecke algebra, which will be finite dimensional  $\mathbb{Q}$ -vector spaces  $V$  together with a homomorphism  $\mathcal{H} \rightarrow \text{End}_{\mathbb{Q}}(V)$ . Let us call this category  $\mathbf{Mod}_{\mathcal{H}}$ . For any extension  $L/\mathbb{Q}$  we may consider the extension  $\mathcal{H}_L = \mathcal{H} \otimes L$  and the resulting category  $\mathbf{Mod}_{\mathcal{H}_L}$ . If we have an extension  $L \hookrightarrow K$  the tensor product yields a functor  $\mathbf{Mod}_{\mathcal{H}_L} \rightarrow \mathbf{Mod}_{\mathcal{H}_K}$ .

We briefly recall the theory of modules over a finite dimensional  $\mathbb{Q}$ -algebra  $\mathcal{A}$  more precisely for any extension  $L/\mathbb{Q}$  we consider the category  $\mathbf{Mod}_{\mathcal{A}_L}$  of finite dimensional  $L$ -vector spaces  $V$  together with a homomorphism  $\mathcal{A}_L \rightarrow \text{End}_L(V)$ .

We say that a finite dimensional  $\mathcal{A}_L$  module  $V$  is irreducible, if  $V$  does not contain a non trivial  $\mathcal{A}_L$  submodule. We say that  $V$  is absolutely irreducible if  $V \otimes \bar{L}$  is irreducible. We say that  $V$  is indecomposable if it can not be written as the direct sum of two non zero submodules.

We call such an algebra  $\mathcal{A}$  semi-simple if it does not contain a non trivial two sided ideal  $\mathcal{N}$  consisting of nilpotent elements. It is well known that this is equivalent to the semi simplicity of the category  $\mathbf{Mod}_{\mathcal{A}}$ , this means that for any  $\mathcal{A}$ -module  $V$  (finite dimensional over  $\mathbb{Q}$ ) and any submodule  $W \subset V$  we can find a  $\mathcal{A}$  submodule  $W'$  such that  $V = W \oplus W'$ . It is also well known that  $\mathcal{A}$  is semi simple if it has a faithful semi-simple (finite dimensional) module  $V \in \mathbf{Ob}(\mathbf{Mod}_{\mathcal{A}})$ , where faithful means that  $\mathcal{A} \rightarrow \text{End}_{\mathbb{Q}}(V)$  is injective and semi simple means of course that any  $\mathcal{A}$ -submodule  $W \subset V$  admits a complement.

It follows from a simple Galois-theoretic argument, that  $\mathcal{A}$  is semi simple if and only if  $\mathcal{A} \otimes_{\mathbb{Q}} L$  is semi simple for any extension  $L/\mathbb{Q}$ .

If we have two modules  $V_1, V_2$  in  $\mathbf{Mod}_{\mathcal{A}_L}$  and these modules become isomorphic after some extension  $L \hookrightarrow K$ , then they are already isomorphic over  $L$ . The isomorphism classes of irreducible modules for  $\mathcal{A}_L$  form a set which is called  $\text{Spec}(\mathcal{A}_L)$ . It is a standard fact from the theory of semi simple algebras that this spectrum can be identified to the set of two sided maximal ideals. We also know that we can write the identity element as a sum of commuting idempotents

$$1 = \sum_{\phi \in \text{Spec}(\mathcal{A}_L)} e_{\phi}; e_{\phi}^2 = e_{\phi}; e_{\phi}e_{\psi} = 0 \text{ for } \phi \neq \psi.$$

Then  $\mathcal{A}_L e_\psi$  is simple, i.e. has no non trivial two sided ideal. The maximal ideal corresponding to  $\phi$  is  $\bigoplus_{\psi:\psi \neq \phi} \mathcal{A} e_\psi$ . We have the decomposition

$$\mathcal{A}_L = \sum_{\phi \in \text{Spec}(\mathcal{A}_L)} \mathcal{A}_L e_\phi \tag{6.29}$$

Our algebra  $\mathcal{A}_L$  has a center  $\mathfrak{Z}_L$ , which is a commutative algebra over  $L$  and since it does not have nilpotent elements it is a direct sum of fields. The idempotents  $e_\phi \in \mathfrak{Z}_L$  and clearly

$$\mathfrak{Z}_L = \bigoplus_{\phi \in \text{Spec}(\mathcal{A}_L)} \mathfrak{Z} e_\phi$$

where  $\mathfrak{Z} e_\phi$  is a field. Hence we get an identification  $\text{Spec}(\mathcal{A}_L) = \text{Spec}(\mathfrak{Z}_L)$ .

We conclude that a semi-simple algebra  $\mathcal{A}_L$  whose center  $\mathfrak{Z}_L$  is a field is actually simple and then the structure theorem of Wedderburn implies

$$\mathcal{A}_L \xrightarrow{\sim} M_n(\mathcal{D})$$

where the right hand side is a matrix algebra of a central division algebra  $\mathcal{D}/\mathfrak{Z}_L$ . This algebra has only one irreducible non zero module: It acts by multiplication from the left on itself, any non zero minimal left ideal yields an irreducible module. These modules (minimal left ideals) are isomorphic to the ideal given by  $\mathfrak{c}_i$  where  $\mathfrak{c}_i$  consists of those matrices which have zero entries outside the  $i$ -th column. In this case  $\text{Spec}(\mathcal{A}_L) = (0)$  is the zero ideal. The unique irreducible module is not absolutely irreducible if  $\mathcal{D} \neq \mathfrak{Z}_L$ . We may choose an extension  $K/L$  which splits the division algebra, then  $\mathcal{A}_F = M_{nd}(K)$  where  $[\mathcal{D} : L] = d^2$ . If this is the case we call the algebra  $\mathcal{A}_K$  absolutely simple. The spectrum does not change.

This tells us that in general the set of isomorphism classes of irreducible  $\mathcal{A}_L$  is canonically isomorphic to  $\text{Spec}(\mathcal{A}_L)$  for any irreducible  $\mathcal{A}_L$  module  $Y_\phi$  we have exactly one  $\phi$  such that  $e_\phi Y = Y$ , and for all  $\psi \neq \phi$   $e_\psi Y = 0$ . On the other hand our construction above yields exactly one module irreducible module  $Y_\phi$  for a given  $\phi$ . For any  $\mathcal{A}_L$  -module  $X$  we get the isotypical decomposition

$$X = \sum_{\phi \in \text{Spec}(\mathcal{A})} e_\phi X,$$

The isotypical component where the isotypical component  $e_\phi X = Y_\phi^{m(X,\phi)}$ , and where  $m(X, \phi)$  is the multiplicity of this component. If we extend our ground field further  $Y_\phi \otimes_L K$  may become reducible, but if our extension  $L/\mathbb{Q}$  is large enough then  $Y_\phi$  will be absolutely irreducible.

Let us start from a semi simple algebra  $\mathcal{A}/\mathbb{Q}$ . Then its center  $\mathfrak{Z}$  is a direct sum of fields,  $\mathfrak{Z} = \bigoplus \mathfrak{Z}_i$ . We say that a finite extension  $F/\mathbb{Q}$  is a *splitting field* for  $\mathcal{A}$  if it is normal and if any summand  $\mathfrak{Z}_i$  can be embedded into  $F$ . Then we get

$$\mathcal{A}_F = \mathcal{A} \otimes_{\mathbb{Q}} F = \bigoplus_{\iota \in \text{Hom}(\mathfrak{Z}, F)} \mathcal{A} \otimes_{\mathfrak{Z}, \iota} F$$

Clearly the center  $\mathcal{A} \otimes_{\mathfrak{Z}, \iota} F = F$  and hence we see that this decomposition is the same as the above decomposition (6.29), we get

**Proposition 6.3.2.** *If  $F/\mathbb{Q}$  is a splitting field of  $\mathcal{A}/\mathbb{Q}$  then we get an action of the Galois group on  $\text{Spec}(\mathcal{A}_F)$ . The orbits of this actions are in one to one correspondence with the elements in  $\text{Spec}(\mathcal{A})$  in this is the set of summands of the decomposition of  $\mathfrak{Z}_{\mathbb{Q}}$  into a direct sum of fields.*

A summand  $\mathcal{A}e_{\phi}F$  has only one non zero irreducible module (up to isomorphism). This module  $Y_{\phi}$  is not necessarily absolutely irreducible because  $\mathcal{A}e_{\phi} \xrightarrow{\sim} M_n(\mathcal{D})$  where  $\mathcal{D}/F$  may be non trivial (we have a non trivial Schur multiplier).

We say that  $\mathcal{A}/\mathbb{Q}$  has trivial Schur multiplier if for all  $\phi \in \text{Spec}(\mathcal{A})$  the division algebra  $\mathcal{D}$  is trivial, i.e. equal to the center.

We apply these general principles to our Hecke -algebra and its action on the cohomology  $H_1^{\bullet}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda})$ . We define the ideal  $I_{K_f}^1$  to be the kernel of this action, then  $\mathcal{H}/I_{K_f}^1 = \mathcal{A}$  is a finite dimensional algebra. It is known- and will be proved later - that  $H_1^{\bullet}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})$  is a semi simple module and hence we see that  $\mathcal{A}$  is semi simple. Then we define the scheme

$$\text{Coh}(H_1^{\bullet}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda})) = \text{Spec}(\mathcal{A}).$$

We will denote the set of geometric points of this scheme, or more simple minded the set of isomorphism classes occurring in this cohomology, by  $\text{Coh}_1(G, K_f, \lambda)$ .

More generally we may consider the set of isomorphism classes of absolutely irreducible Hecke modules occurring in the Jordan-Hölder filtration of any of our cohomology modules  $H_?^{\bullet}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda})$  and denote this set by  $\text{Coh}_?(G, K_f, \lambda)$ . Since we have a fixed level  $K_f$  they are all defined over a suitable finite extension  $F/\mathbb{Q}$ .

### A central subalgebra

We still consider the action of  $\mathcal{H}/I_{K_f}^1 = \mathcal{A}$  on  $H_1^{\bullet}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}) = \bigoplus_q H_1^q(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})$ . For all  $p$  outside the finite set  $\Sigma$  we have  $K_p = \mathcal{G}(\mathbb{Z}_p)$ . In this case the algebra  $\mathcal{H}_p$  is finitely generated, integral and commutative. We say that  $\mathcal{H}_p$  is *unramified* if  $K_p = \mathcal{G}(\mathbb{Z}_p)$ . For an unramified Hecke-algebra  $\mathcal{H}_p$  its maximal spectrum

$\text{Hom}_{\text{alg}}(\mathcal{H}_p, \mathbb{C})$ , - i.e. the set of isomorphism classes of absolutely irreducible modules over  $\mathbb{C}$ -, is described by a theorem of Satake which we will recall in the next section.

The subalgebra

$$\mathcal{H}^{(\Sigma)} = \bigotimes'_{p \notin \Sigma} \mathcal{H}_p \tag{6.30}$$

is commutative and its image in  $\mathcal{H}/I_{K_f}^1$  lies in the center and hence also in the center of  $\mathcal{A}$ . Hence we can conclude that for a splitting field  $F$  for  $\mathcal{A}$  and any irreducible module  $Y_{\phi}$  for  $\mathcal{A}_F$  the restriction of the action to  $\mathcal{H}^{(\Sigma)}$  is given by a homomorphism

$$\phi^{(\Sigma)} : \mathcal{H}^{(\Sigma)} \rightarrow F.$$

Hence the module  $Y_{\phi}$  is determined by the action of  $\mathcal{H}_{\Sigma} = \prod_{p \in \Sigma} \mathcal{H}_p$  in  $\mathcal{A}_F$ . If we assume that  $Y_{\phi}$  is absolutely irreducible, then it follows from a standard argument that  $Y_{\phi} \xrightarrow{\sim} \bigotimes_{p \in \Sigma} Y_{\phi_p}$  where  $Y_{\phi_p}$  is an absolutely irreducible  $\mathcal{H}_p$ -module. For  $p \notin \Sigma$  let  $V_{\phi_p}$  be the one dimensional  $F$  vector space  $F$  with

canonical basis element  $1 \in F$  and an  $\mathcal{H}_p$  action given by the homomorphism  $\phi_p : \mathcal{H}_p \rightarrow F$ . Then we get an isomorphism

$$Y_\phi \xrightarrow{\sim} \bigotimes_p^{\prime} Y_{\phi_p}, \tag{Fl}$$

where we take the restricted tensor product in the usual sense, i.e. at almost all primes the factor in a tensor is equal to 1. Under our assumptions the homomorphism

$$\mathcal{H}_p \rightarrow \text{End}_F(Y_{\phi_p})$$

is surjective.

We get a map from the isomorphism classes of irreducible modules  $[Y_\phi]$  for  $\mathcal{A}_F$  to  $\phi^\sigma \in \text{Hom}(\mathcal{H}^{(\Sigma)}, F)$ . We say that  $\mathcal{H}^{(\Sigma)}$  acts *distinctively* on  $H_1^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}} \otimes F)$  if this map is injective, i.e. the isomorphism type  $[Y_\phi]$  is determined by its restriction to  $\mathcal{H}^{(\Sigma)}$ .

On the cohomology  $H_1^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})$  we still have the action of the group  $\pi_0(G(\mathbb{R}))$ , this action commutes with the action of the Hecke algebra. (See (6.3.7) This is an elementary abelian 2- group and we may decompose further according to characters  $\epsilon : \pi_0(G(\mathbb{R})) \rightarrow \{\pm 1\}$ .

We say that the  $\mathcal{H}$  module  $H_1^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})$  has *strong multiplicity one* (with respect to  $\Sigma$ ) if  $\mathcal{H}^{(\Sigma)}$  acts distinctively and for any splitting field  $F$  and any  $\phi^\Sigma : \mathcal{H}^{(\Sigma)} \rightarrow F$  we can find a degree  $q$  and an  $\epsilon$  such that

$$H_1^q(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})(\epsilon) \otimes_{\mathcal{H}^{(\Sigma)}, \phi^\Sigma} F$$

is an absolutely irreducible  $\mathcal{H}$ - module.

If this is so then the homomorphism

$$\mathcal{H}_\Sigma \rightarrow \text{End}_F(H_1^q(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})(\epsilon) \otimes_{\mathcal{H}^{(\Sigma)}, \phi^\Sigma} F)$$

is surjective and the Hecke module  $H_1^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})$  has trivial Schur multiplier.

### Representations and Hecke modules

For  $p \in \Sigma$  the category of finite dimensional modules is complicated, since the Hecke algebra will not be commutative in general.

Let  $F$  be a field of characteristic zero, let  $V$  be an  $F$ -vector space. An admissible representation of the group  $G(\mathbb{Q}_p)$  is an action of  $G(\mathbb{Q}_p)$  on  $V$  which has the following two properties

- (i) For any open compact subgroup  $K_p \subset G(\mathbb{Q}_p)$  the space  $V^{K_p}$  of  $K_p$  invariant vectors is finite dimensional.
- (ii) For any vector  $v \in V$  we can find an open compact subgroup  $K_p$  so that  $v \in V^{K_p}$  in other words  $V = \lim_{K_p} V^{K_p}$ .

An admissible  $G(\mathbb{Q}_p)$  -module  $V$  is irreducible if it does not contain an invariant proper submodule.

It is clear that the vector spaces  $V^{K_p}$  are modules for the Hecke algebra

$$\mathcal{H}_{K_p} \cdot \boxed{\text{VKirr}}$$

**Proposition 6.3.3.** *If  $V \neq (0)$  is a irreducible  $G(\mathbb{Q}_p)$  modules, and if  $K_p$  is an open compact subgroup with  $V^{K_p} \neq (0)$ . Then  $V^{K_p}$  is an irreducible  $\mathcal{H}_{K_p}$ -module.*

*Proof.* To see this we take the identity element  $e_{K_p}$  in our Hecke algebra, it induces a projector on  $V$  and a decomposition

$$V = V^{K_p} \oplus V' = e_{K_p} V \oplus (1 - e_{K_p})V.$$

Let assume we have a proper  $\mathcal{H}_{K_p}$ -invariant submodule  $W \subset V^{K_p}$  Now we convince ourselves that the  $G(\mathbb{Q}_p)$ -invariant subspace  $\tilde{W}$  generated by the elements  $gw$  is a proper subspace. We compute the integral

$$\int_{K_p} kgwk = \int_{K_p \times K_p} k_1 g k_2 w dk_2 dk_1.$$

The first integral gives us the projection to  $V^{K_p}$ , the second integral is the Hecke operator, hence the result is in  $W$ . We conclude that  $e_{K_p} \tilde{W} \subset W$  and tis shows that  $(0) \neq \tilde{W} \neq V$ .  $\square$

Now it is not hard to see, that the assignment

$$V \rightarrow V^{K_p}$$

from irreducible admissble  $G(\mathbb{Q}_p)$ -modules with  $V^{K_p} \neq (0)$  to finite dimensional irreducible  $\mathcal{H}_{K_p}$ -modules induces an bijection between the isomorphism classes of the respective types of modules. If we start from  $V^{K_p}$  we can reconstruct  $V$  by an appropriate form of induction.

### The dual module

Let us assume that  $V$  is a finite dimensional  $F$ -vector space with an action of the Hecke algebra  $\mathcal{H}$  (we fix the level). We have an involution on the Hecke algebra which is defined by

$${}^t h(\underline{x}_f) = h(\underline{x}_f^{-1})$$

a simple calculation shows that  ${}^t h_1 * {}^t h_2 = {}^t (h_2 * h_1)$ .

This allows us to introduce a Hecke-module structure on  $V^\vee = \text{Hom}_F(V, F)$  we for  $\phi \in V^\vee$  we simply put

$$T_h(\phi)(v) = \phi(T_{{}^t h}(v))$$

for all  $v \in V$ .

### Unitary and essentially unitary representations

Here it seems to be a good moment to recall the notion of unitary Hecke modules and unitary representations. In this book we make the convention that a character is a continuous homomorphism from a topological group  $H \rightarrow \mathbb{C}^\times$ , we do not require that its values have absolute value one. If this is the case we call the character unitary. Our ground field will now be  $F = \mathbb{C}$ , let  $V$  be a  $\mathbb{C}$  vector

space. We pick a prime  $p$ . We call a representation  $\rho : G(\mathbb{Q}_p) \rightarrow \text{Gl}(V)$  unitary if there is given a positive definite hermitian scalar product  $\langle , \rangle : V \times V \rightarrow \mathbb{C}$  which is invariant under the action of  $G(\mathbb{Q}_p)$ .

If our representation is irreducible then it has a central character  $\zeta_\rho : C(\mathbb{Q}_p) \rightarrow \mathbb{C}^\times$ . In this case the scalar product is unique up to a scalar. A necessary condition for the existence of such a scalar product is that  $|\zeta_\rho| = 1$ , in other words  $\zeta_\rho$  is unitary.

If this is not the case then our representation may still be *essentially unitary*: We have a unique homomorphism  $|\zeta_\rho^*| : C'(\mathbb{Q}_p) \rightarrow \mathbb{R}_{>0}^\times$  whose restriction to  $C(\mathbb{Q}_p)$  under  $d_C$  (see 1.1) is equal to  $|\zeta_\rho|$ . Then we may form the twisted representation  $\rho^* = \rho \otimes |\zeta_\rho^*|^{-1}$ . Then the central character of  $\rho^*$  is unitary. We say that  $\rho$  is called essentially unitary if  $\rho^*$  is unitary.

If our representation is not irreducible we still can define the notion of being essential unitary. This means that there exists a homomorphism  $|\zeta_\rho^*| : C'(\mathbb{Q}_p) \rightarrow \mathbb{R}_{>0}^\times$ , such that the twisted representation  $\rho^* = \rho \otimes |\zeta_\rho^*|^{-1}$  is unitary.

The same notions apply to modules for the Hecke algebra. A (finite dimensional)  $\mathbb{C}$  vector space  $V$  with an action  $\pi_p : \mathcal{H}_p \rightarrow \text{End}(V)$  is called unitary, if there is given a positive definite scalar product  $\langle , \rangle : V \times V \rightarrow \mathbb{C}$  such that

$$\langle T_h(v), w \rangle = \langle v, (T_h(w)) \rangle. \tag{6.31}$$

Recall that we always assume that our functions  $h \in \mathcal{H}_p$  take values in  $\mathbb{Q}$ , hence we do not need a complex conjugation bar in the expression on the right.

The restriction of  $\pi_p$  to  $C(\mathbb{Q}_p)$  induces a homomorphism  $\zeta_{\pi_p} : C(\mathbb{Q}_p) \rightarrow \mathbb{C}^\times$ . We call  $\pi_p$  isobaric if this action of the center is semi simple - and therefore a direct sum of characters  $\zeta_{\pi_p} = \sum \zeta_{\pi_p}^\nu$  - and if all these characters have the same absolute values  $|\zeta_{\pi_p}^\nu| = |\zeta_{\pi_p}|$ . This means that we can find  $|\zeta_{\pi_p}^*|$  as above. Then we call  $\pi_p$  essentially unitary if the Hecke module  $\pi_p^* = \pi_p \otimes |\zeta_{\pi_p}^*|^{-1}$  is unitary.

These boring considerations will be needed later, we will see that for an irreducible coefficient system  $\mathcal{M}$  the  $H_1^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}) \otimes \mathbb{C}$  is essentially unitary (see 8.1.6).

### 6.3.4 The Satake isomorphism

**Noch mal ein wenig überarbeiten** In the formulation of this theorem I will use the language of group schemes, the reader not so familiar with this language may think of  $\text{Gl}_n$  or the group of symplectic similitudes  $\text{GSp}_n$ . Since we assumed that for  $p \notin \Sigma$  the integral structure  $\mathcal{G}/\text{Spec}(\mathbb{Z}_p)$  is reductive it is also quasisplit. We can find a Borel subgroup  $\mathcal{B}/\text{Spec}(\mathbb{Z}_p) \subset \mathcal{G}/\text{Spec}(\mathbb{Z}_p)$  and a maximal torus  $\mathcal{T}/\text{Spec}(\mathbb{Z}_p) \subset \mathcal{B}/\text{Spec}(\mathbb{Z}_p)$ . Then our torus  $\mathcal{T}/\text{Spec}(\mathbb{Z}_p)$  splits over an unramified extension  $E_p/\mathbb{Q}_p$  and the Galois group  $\text{Gal}(E_p/\mathbb{Q}_p)$  acts on the character module  $X^*(\mathcal{T} \times E_p) = \text{Hom}(\mathcal{T} \times E_p, \mathbb{G}_m)$ . Let  $\{\alpha_1, \alpha_2, \dots, \alpha_r\} \subset X^*(\mathcal{T} \times E_p)$  be the set of positive simple roots, it is invariant under the action of the Galois group. Let  $W(\mathbb{Z}_p)$  be the centralizer of the Galois action in the absolute Weyl group  $W$ . We introduce the module of unramified characters on the torus this is

$$\text{Hom}_{\text{unram}}(\mathcal{T}(\mathbb{Q}_p), \mathbb{C}^\times) = \text{Hom}(\mathcal{T}(\mathbb{Q}_p)/\mathcal{T}(\mathbb{Z}_p), \mathbb{C}^\times) = \Lambda(\mathcal{T}).$$

Since we have  $T(\mathbb{Q}_p) = B(\mathbb{Q}_p)/U(\mathbb{Q}_p)$  the character  $\omega_p \in \Lambda(\mathcal{T})$  yields a character  $\omega_p : B(\mathbb{Q}_p) \rightarrow \mathbb{C}^\times$ . We write the module structure additively, i.e.  $(\omega_{1,p} + \omega_{2,p})(x) = \omega_{1,p}(x)\omega_{2,p}(x)$

The group of (rational) characters  $\text{Hom}(\mathcal{T}, \mathbb{G}_m) = X^*(T)^{\text{Gal}(E_p/\mathbb{Q}_p)}$  is a subgroup of  $\Lambda(\mathcal{T})$ : An element  $\gamma \in X^*(\mathcal{T})^{\text{Gal}(E_p/\mathbb{Q}_p)}$  defines a homomorphism  $\mathcal{T}(\mathbb{Q}_p) \rightarrow \mathbb{Q}_p^\times$  and this gives us the following element  $x \mapsto |\gamma(x)|_p \in \Lambda(\mathcal{T})$  which we denote by  $|\gamma|_p$ . We can even do this for elements  $\gamma \otimes \frac{1}{n} \in X^*(T) \otimes \mathbb{Q}$ , then  $\gamma \otimes \frac{1}{n}(x) = |\gamma(x)|_p^{1/n} \in \mathbb{R}_{>0}^\times$ .

Our open compact subgroup  $K_p = \mathcal{G}(\mathbb{Z}_p)$ . Since we have the Iwasawa decomposition  $G(\mathbb{Q}_p) = B(\mathbb{Q}_p)\mathcal{G}(\mathbb{Z}_p) = B(\mathbb{Q}_p)K_p$  we can attach to any  $\omega_p \in \Lambda(\mathcal{T})$  a *spherical function*

$$\phi_{\omega_p}(g) = \phi_{\omega_p}(b_p k_p) = (\omega_p + |\rho|_p)(b_p) \quad (6.32)$$

here  $\rho \in \Lambda(\mathcal{T}) \otimes \mathbb{Q}$  is the half sum of positive roots. This spherical function is of course an eigenfunction for  $\mathcal{H}_p$  under convolution, i.e. for  $h_p \in \mathcal{H}_p$

$$\int_{G(\mathbb{Q}_p)} \phi_{\omega_p}(gx)h_p(x)dx = \hat{h}_p(\omega_p)\phi_{\omega_p}(g) \quad (6.33)$$

and  $\mathfrak{s}(\omega_p) : h_p \mapsto \hat{h}_p(\omega_p)$  is an algebra homomorphism from  $\mathcal{H}_p$  to  $\mathbb{C}$ . Of course the measure  $dx$  gives volume 1 to  $\mathcal{G}(\mathbb{Z}_p) = K_p$ .

The theorem of Satake asserts:

**Theorem 6.3.1.** *The group  $W(\mathbb{Z}_p)$  acts on  $\Lambda(\mathcal{T})$ , we have  $\mathfrak{s}(w\omega_p) = \mathfrak{s}(\omega_p)$  and*

$$\Lambda(\mathcal{T})/W(\mathbb{Q}_p) \xrightarrow{\mathfrak{s}} \text{Hom}_{\text{alg}}(\mathcal{H}_p, \mathbb{C})$$

*is an isomorphism.*

We will write irreducible modules in this case as  $\pi_p = \pi_p(\omega_p)$  and  $\omega_p \in \Lambda(\mathcal{T})/W(\mathbb{Q}_p)$  is the so called *Satake parameter* of  $\pi_p$ .

The Hecke algebra is generated by the characteristic functions of double cosets  $K_p t_p K_p$  where  $t_p \in T(\mathbb{Q}_p)$  and where for all simple roots  $\alpha \in \pi$  we have  $|\alpha(t_p)|_p \leq 1$ , i.e.  $t_p \in T_+(\mathbb{Q}_p)$ . Then the evaluation in (6.33) comes down to the computation the integrals

$$\int_{K_p t_p K_p} \phi_{\omega_p}(gx)dx = \hat{t}_p(\omega_p)\phi_{\omega_p}(g) \quad (6.34)$$

We discuss this evaluation in (7.0.1) (????????)

### Spherical representations

Now we assume that Let  $F \subset \mathbb{C}$  be a finite extension of  $\mathbb{Q}$  and let  $V/F$  be a vector space. We choose  $K_p = \mathcal{G}(\mathbb{Z}_p)$ , i.e.  $p$  is unramified. An admissible representation

$$\tilde{\pi}_p : G(\mathbb{Q}_p) \rightarrow \text{Gl}(V)$$

is called *spherical* if  $V^{K_p} \neq 0$ , and this space is a module for the Hecke algebra. If the representation is absolutely irreducible, then it is well known (**Reference**) that  $\dim_F V^{K_p} = 1$ , this is a one dimensional module for  $\mathcal{H}_{K_p}$ , i.e. a homomorphism  $\pi_p : \mathcal{H}_{K_p} \rightarrow F$ . Let  $\omega_p \in \Lambda(\mathcal{T})$  the corresponding Satake parameter, it is well defined modulo the action of the group  $W(\mathbb{Q}_p)$ . We introduce the element  $\chi_p = \omega_p + |\rho|_p \in \Lambda(T)$  we observe that the values  $\chi_p(x) \in F_1$  where  $F_1$  is a finite normal extension of  $F$ .

Then our representation  $\tilde{\pi}_p \otimes_F F_1$  can be realised as a subquotient of the induced representation  $\square$

$$\text{Ind}_{B(\mathbb{Q}_p)}^{G(\mathbb{Q}_p)} \chi_p = \{f : G(\mathbb{Q}_p) \rightarrow F \mid f(bg) = \chi_p(b)f(g)\} \tag{6.35}$$

where  $f$  satisfies the (obvious) condition that there exists a finite index subgroup  $K'_p \subset K_p$  such that  $f$  is invariant under right translations by elements  $k' \in K'_p$ . In general the induced representation will be irreducible and then it is isomorphic to the representation  $\tilde{\pi}_p \otimes_F F_1$ .

If  $\tilde{\pi}_p^\vee$  is the spherical representation attached to the Satake parameter  $\omega_p^{-1}$  then we have a pairing  $\square$  dualSat

$$\begin{aligned} H_{\tilde{\pi}_p} \times H_{\tilde{\pi}_p^\vee} &\rightarrow \mathbb{C} \\ f_1 \times f_2 &\mapsto \int_{K_p} f_1(k_p) f_2(k_p) dk_p \end{aligned} \tag{6.36}$$

This tells us that the dual module to  $H_{\pi_p} = H_{\tilde{\pi}_p}^{K_p}$  has the Satake parameter  $\omega_p^{-1}$ . The representations  $H_{\tilde{\pi}_p}$  are called the representations of the unramified principal series.

We may consider the case that  $\omega_p$  is a unitary character, this means that  $\omega_p : \mathcal{T}(\mathbb{Q}_p)/\mathcal{T}(\mathbb{Z}_p) \rightarrow \mathbb{S}^1$ . Then we have  $\omega_p^{-1}(t) = \overline{\omega_p}(t)$  and our above pairing defines a positive definite hermitian scalar product

$$\langle , \rangle : H_{\tilde{\pi}_p} \times H_{\tilde{\pi}_p} \rightarrow \mathbb{C} \tag{6.37}$$

which is given by

$$\langle f_1, f_2 \rangle = \int_{K_p} f_1(k_p) \overline{f_2(k_p)} dk_p \tag{6.38}$$

If we allow for  $f \in H_{\tilde{\pi}_p}$  all the functions whose restriction to  $K_p$  lies in  $L^2(K_p)$  then  $H_{\tilde{\pi}_p}$  becomes a Hilbert space and the representation of  $G(\mathbb{Q}_p)$  on  $H_{\tilde{\pi}_p}$  is a unitary representation.

These representations are called the unitary principal series representations. It is not the case that these representations are the only unramified principal series representations which carry an invariant positive definite scalar product. (See [Sat]).

In the following section we discuss the classical case.  $\square$  SatakeGl

### 6.3.5 The case $\mathrm{Gl}_2$ .

In the case of  $\mathrm{Gl}_2$  the maximal torus is given by

$$T(\mathbb{Q}_p) = \left\{ \begin{pmatrix} t_1 & 0 \\ 0 & t_2 \end{pmatrix} \right\}.$$

It is contained in the two Borel subgroups  $B/\mathbb{Q}$  of upper and  $B_-/\mathbb{Q}$  of lower triangular matrices. Let  $U/\mathbb{Q}$  be the unipotent radical of  $B$ .

If we represent an element  $\bar{\omega}_p \in \Lambda(\mathcal{T})/W$  by  $\omega_p : T(\mathbb{Q}_p)/T(\mathbb{Z}_p) \rightarrow \mathbb{C}^\times$  then we get two numbers

$$\begin{aligned} \omega_p \left( \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} \right) &= \alpha'_p \\ \omega_p \left( \begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix} \right) &= \beta'_p \end{aligned}.$$

The local algebra is generated by two operators  $T_p, T(p, p)$  for which

$$\begin{aligned} \mathfrak{s}(\bar{\omega}_p)(T_p) &= p^{1/2}(\alpha'_p + \beta'_p) = \alpha_p + \beta_p \\ \mathfrak{s}(\bar{\omega}_p)(T(p, p)) &= p\alpha'_p\beta'_p = \alpha_p\beta_p \end{aligned}.$$

These two Hecke operators are -up to a normalizing factor - defined as the characteristic functions of the double cosets

$$\mathrm{Gl}_2(\mathbb{Z}_p) \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} \mathrm{Gl}_2(\mathbb{Z}_p) \text{ and } \mathrm{Gl}_2(\mathbb{Z}_p) \begin{pmatrix} p & 0 \\ 0 & p \end{pmatrix} \mathrm{Gl}_2(\mathbb{Z}_p).$$

The two numbers  $\alpha_p + \beta_p, \alpha_p\beta_p$  determine  $\omega_p$ . They are called the *arithmetic Satake parameters*. We also define the *semi-simple* component of the Satake parameter Satparmss

$$\omega^{(1)}(\pi_p) = \omega_p \left( \begin{pmatrix} p & 0 \\ 0 & p^{-1} \end{pmatrix} \right) = \frac{\alpha_p}{\beta_p} = \frac{\alpha'_p}{\beta'_p} \quad (6.39)$$

It is not difficult to prove Satake's theorem for  $\mathrm{Gl}_2/\mathbb{Q}_p$ . We write  $\mathrm{Gl}_2(\mathbb{Z}_p) = K_p$ . It is the theorem for elementary divisors that all the double cosets  $K_p \backslash G(\mathbb{Q}_p) / K_p$  are of the form

$$K_p \begin{pmatrix} p^a & 0 \\ 0 & p^b \end{pmatrix} K_p \text{ with } a \geq b.$$

If we want to understand the function  $h \mapsto \hat{h}(\lambda)$  it clearly suffices to compute its value on the characteristic function  $t_{p^m}$  of the double coset

$$K_p ([p^m]) K_p$$

For  $\lambda \in \Lambda(T)$  we have to evaluate the integral

$$\int_{G(\mathbb{Q}_p)} \phi_\lambda(x) t_{p^m}(x) dx = t_{p^m}^\wedge(\lambda).$$

We abbreviate  $[p^m] = \begin{pmatrix} p^m & 0 \\ 0 & 1 \end{pmatrix}$  and write our double coset as a union of right  $K_p$  cosets, i.e.

$$K_p[p^m]K_p = \bigcup_{\xi \in K_p/K_p \cap [p^m]K_p[p^m]^{-1}} \xi[p^m]K_p.$$

The volume of such a coset is one hence we get

$$\int_{G(\mathbb{Q}_p)} \phi_\lambda(x)t_{p^m}(x)dx = \sum_{\xi} \phi_\lambda(\xi[p^m])$$

The group

$$K_p \cap [p^m]K_p[p^m]^{-1} = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in K_p \mid b \equiv 0 \pmod{p^m} \right\},$$

this is the group of points  $B_-(\mathbb{Z}/p^m)$  of lower triangular matrices. Hence the coset space

$$\mathrm{Gl}_2(\mathbb{Z}/p^m)/B_-(\mathbb{Z}/p^m) = K_p/K_p \cap [p^m]K_p[p^m]^{-1} = \mathbb{P}^1(\mathbb{Z}/p^m).$$

The points in  $\mathbb{P}^1(\mathbb{Z}/p^m)$  are arrays  $\begin{pmatrix} a \\ b \end{pmatrix}$ ,  $a, b \in \mathbb{Z}/p^m$ ,  $a$  or  $b \in (\mathbb{Z}/p^m\mathbb{Z})^\times$ , modulo  $(\mathbb{Z}/p^m)^\times$ . Then  $K_p$  acts by multiplication from the left on this coset space and  $K_p \cap [p^m]K_p[p^m]^{-1}$  is the stabilizer of  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ . We still have an action of  $B(\mathbb{Z}/p^m)$  from the left on  $\mathbb{P}^1(\mathbb{Z}/p^m)$  and the orbits under this action from the left can be represented by

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix} \text{ and } \begin{pmatrix} p^\nu \\ 1 \end{pmatrix} \text{ for } \nu = 0, 1, \dots, m$$

On these orbits (under  $B(\mathbb{Z}/p^m)$ ) the function  $\xi \mapsto \phi_\lambda(\xi[p^m])$  is constant. We can take the representatives

$$\xi = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \text{ and } \begin{pmatrix} 1 & p^\nu \\ 0 & 1 \end{pmatrix}$$

and get the values

$$\begin{aligned} \phi_\lambda(w[p^m]) &= \beta_p^m \\ \phi_\lambda\left(\begin{pmatrix} 1 & p^\nu \\ 0 & 1 \end{pmatrix}[p^m]\right) &= (\alpha'_p)^m p^{-\frac{m}{2}} = \alpha_p^m p^{-m}. \end{aligned}$$

The total length of the of the orbits under  $B(\mathbb{Z}/p^m\mathbb{Z})$  is  $p^m$  and hence tpm

$$t_{p^m}(\lambda) = \alpha_p^m + \beta_p^m \tag{6.40}$$

We want to put on record that for an (irreducible) induced representation  $\mathrm{Ind}_{B(\mathbb{Q}_p)}^{\mathrm{Gl}_2(\mathbb{Q}_p)} \chi_p$  where  $\chi\left(\begin{pmatrix} t_1 & 0 \\ 0 & t_2 \end{pmatrix}\right) = \chi_1(t_1)\chi_2(t_2)$  the arithmetic Satake parameters are given by aritSat

$$p\chi_1(p) + \chi_2(p) ; p\chi_1(p)\chi_2(p). \tag{6.41}$$

### 6.3.6 Back to cohomology

#### The case of a torus and the central character

We consider the case that our group  $G/\mathbb{Q}$  is a torus  $T/\mathbb{Q}$ . This case is already discussed in [Ha-Gl2]. Our torus splits over a finite extension  $F/\mathbb{Q}$  and our absolutely irreducible representation is simply a character  $\gamma : T \times_{\mathbb{Q}} F \rightarrow \mathbb{G}_m$ , it defines a one dimensional  $T \times_{\mathbb{Q}} F$ - module  $F[\gamma]$ . Here  $F[\gamma]$  is simply the one dimensional vector space  $F$  over  $F$  with  $T \times_{\mathbb{Q}} F$  acting by the character  $\gamma$ .

We recall the notion of an algebraic Hecke character of type  $\gamma$ . We choose an embedding  $\iota : F \hookrightarrow \mathbb{Q}$  then  $\gamma$  induces a homomorphism  $T(\mathbb{C}) \rightarrow \mathbb{C}^{\times}$ . The restriction of this homomorphism to  $T(\mathbb{R})$  is called  $\gamma_{\infty} : T(\mathbb{R}) \rightarrow \mathbb{C}^{\times}$ .

A continuous homomorphism

$$\phi = \phi_{\infty} \times \prod_p \phi_p = \phi_{\infty} \times \phi_f : T(\mathbb{A})/T(\mathbb{Q}) \rightarrow \mathbb{C}^{\times}$$

is called an *algebraic Hecke character of type  $\gamma$*  if the restrictions to the connected component of the identity satisfy

$$\phi_{\infty}|_{T^{(0)}(\mathbb{R})} = \gamma_{\infty}^{-1}|_{T^{(0)}(\mathbb{R})}.$$

The finite part  $\phi_f : T(\mathbb{A}_f) \rightarrow \bar{\mathbb{Q}}^{\times}$  is trivial on some open compact subgroup  $K_f^T \subset T(\mathbb{A}_f)$ . We also say that a homomorphism  $\phi_1 : T(\mathbb{A}_f)/K_f^T \rightarrow \bar{\mathbb{Q}}^{\times}$  is an algebraic Hecke-character, if it is the finite part of an algebraic Hecke character, which is then uniquely defined.

In [Ha-Gl2], 2.5.5 we explain that the cohomology vanishes ( for any choice of  $K_f^C$  ) if  $\gamma$  is not the type of an algebraic Hecke character. In this case we give the complete description of the cohomology in [Ha-Gl2], 2.6: If we choose  $Z' = Z$  (see 1.1) then

$$H^0(S_{K_f^C}^C, F[\gamma] \otimes_{F,\iota} \bar{\mathbb{Q}}) = \bigoplus_{\phi_f : C(\mathbb{A}_f)/K_f^C \rightarrow \bar{\mathbb{Q}}^{\times} : \text{type}(\phi_f) = \gamma} \bar{\mathbb{Q}}\phi_f. \quad (6.42)$$

The property of  $\gamma$  to be the type of an algebraic Hecke character does not depend on the choice of  $\iota$ . If we fix the level then it is easy to see that the values of the characters  $\phi_f$  lie in a finite extension  $F_1$  of  $\iota(F)$  so we may replace in our formula above the algebraic closure  $\bar{\mathbb{Q}}$  by  $F_1$ .

If we return to our group  $G/\mathbb{Q}$  and if we start from an absolutely irreducible representation  $G \times_{\mathbb{Q}} F \rightarrow \text{Gl}(\mathcal{M})$  then its restriction to the center  $C/\mathbb{Q}$  is a character  $\zeta_{\mathcal{M}}$ . Our remark above implies that this character must be the type of an algebraic Hecke character if we want the cohomology  $H_?^{\bullet}(S_{K_f}^G, \tilde{\mathcal{M}})$  to be non trivial. (Look at a suitable spectral sequence).

In any case we can consider the sub algebra  $C_{K_f} \subset \mathcal{H}_{K_f}$  generated by central double cosets  $K_f z_f K_f = K_f \bar{z}_f$  with  $z_f \in C(\mathbb{A}_f)$  This provides an action of the group  $C(\mathbb{A}_f)/K_f^C$  on the cohomology  $H_?^{\bullet}(S_{K_f}^G, \tilde{\mathcal{M}})$ . Then the following proposition is obvious

**Proposition 6.3.4.** *Let  $H_{\pi_f}$  be an absolutely irreducible subquotient in the Jordan Hölder series in any of our cohomology groups. Then  $C(\mathbb{A}_f)/K_f^C$  acts by a character  $\zeta_{\pi_f}$  on  $H_{\pi_f}$  and  $\zeta_{\pi_f}$  is an algebraic Hecke character of type  $\zeta_{\mathcal{M}}$ .*

Note that  $\zeta_{\mathcal{M}}$  is the restriction of the abelian component  $\delta$  in  $\lambda = \lambda^{(1)} + \delta$  to the center.

### The cohomology in degree zero

Let us start from an absolutely irreducible representation  $r : G \times F \rightarrow \mathrm{Gl}(\mathcal{M})$ , we want to understand  $H^0(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})$ : To do this we have to understand the connected components of the space and the spaces of invariants in  $\tilde{\mathcal{M}}$  under the discrete subgroups  $\Gamma_f^g$  in 1.2.1. We assume that the groups  $\Gamma_f^g \cap G^{(1)}(\mathbb{Q})$  are Zariski dense in  $G^{(1)}$ . Then it is clear that we can have non trivial cohomology in degree zero if  $\mathcal{M}$  is one dimensional and  $G^{(1)}$  acts trivially. Hence  $\mathcal{M}$  is given by a character  $\delta : C' \times F \rightarrow \mathbb{G}_m \times F$ .

To simplify the situation we assume that the assumptions in (6.1.3) are fulfilled and we have a bijection

$$\pi_0(\mathcal{S}_{K_f}^G) \xrightarrow{\sim} \pi_0(\mathcal{S}_{K_\infty^{C'} \times K_f^{C'}}^{C'}) \quad (6.43)$$

where  $K_\infty^{C'}$  and  $K_f^{C'}$  are the images of the chosen compact subgroups respectively. With these data we define  $\mathcal{S}_{K_f^{C'}}^{C'}$  and we can view  $\mathcal{M}$  as a sheaf on  $\mathcal{S}_{K_f^{C'}}^{C'}$ , in our previous notation it is the sheaf  $\tilde{F}[\delta]$ .

Then we get for an absolutely irreducible  $G \times F$  module  $\mathcal{M}$  -and under the assumption that the  $\Gamma_f^g \cap G^{(1)}(\mathbb{Q})$  are Zariski dense in  $G^{(1)}$ - that (See 6.3.6)

$$H^0(\mathcal{S}_{K_f}^G, \mathcal{M} \otimes F_1) = \begin{cases} 0 & \text{if } \dim(\mathcal{M}) > 1 \\ \bigoplus_{\phi_f: \text{type}(\phi_f)=\delta} F_1 \phi & \text{if } \mathcal{M} = F[\delta] \end{cases} \quad (6.44)$$

The density assumption is fulfilled if  $G^{(1)}/\mathbb{Q}$  is quasisplit. We also observe that we have the isogeny  $d_C : C \rightarrow C'$  (See (1.1)). Then it is clear that the composition  $d_C \circ \delta$  is the character  $\zeta_{\mathcal{M}}$  in section 6.3.6.

### 6.3.7 The Manin-Drinfeld principle

For a moment we assume that our coefficient systems are rational vector spaces. This means that we start from a rational (preferably absolutely irreducible) representation  $\rho : G \times_{\mathbb{Q}} F_0 \rightarrow \mathrm{Gl}(\mathcal{M})$  where  $\mathcal{M}$  is a finite dimensional  $F_0$  vector spaces. We have an action of  $\mathcal{H}_{F_0}$  on our cohomology groups and we defined the spectra  $\mathrm{Coh}(H_?^*(\mathcal{S}_{K_f}^G, \mathcal{M}))$  which now will be a finite scheme over  $F_0$ . We will show that the modules  $H_?^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_L)$  are semi simple and if we pass to a splitting field  $F/F_0$  we can decompose

$$H_?^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})(\Pi_f) \otimes F = \bigoplus_{\pi_f} H_?^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})(\pi_f) = \bigoplus_{\pi_f} e_{\pi_f} H_?^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}) \quad (6.45)$$

Here we changed our notation slightly, we replaced the  $\phi$  by  $\pi_f$ . The isomorphism types  $\pi_f$  are not necessarily absolutely irreducible, but if we extend our field further then they decompose in a direct sum of modules of exactly one isomorphism type. We call the above decomposition the isotypical decomposition and under our assumption on  $F$  the summands are absolutely isotypical.

We say that for a cohomology group  $H^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F)$  (resp.  $H_c^*(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F)$ ) satisfies the *Manin-Drinfeld principle*, if  $\mathrm{Coh}(H_?^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F) \cap \mathrm{Coh}(H^i(\partial(\mathcal{S}_{K_f}^G), \tilde{\mathcal{M}}_F)) = \emptyset$  (resp  $\mathrm{Coh}(H_?^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F) \cap \mathrm{Coh}(H^{i-1}(\partial(\mathcal{S}_{K_f}^G), \tilde{\mathcal{M}}_F)) = \emptyset$ ).

We have defined  $\text{Coh}(X)(= \text{Spec}(\mathcal{H}/I(X)))$  for any Hecke-module  $X$  and if  $X$  is a submodule of another Hecke module  $Y$  then we say that  $X$  satisfies the Manin-Drinfeld principle with respect to  $Y$  if  $\text{Coh}(X) \cap \text{Coh}(Y/X) = \emptyset$ .

If the Manin-Drinfeld principle is valid we get canonical decompositions

$$H^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F) = \text{Im}(H^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F) \longrightarrow H^i(\partial(\mathcal{S}_{K_f}^G), \tilde{\mathcal{M}}_F)) \oplus H_1^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F), \quad (6.46)$$

$$H_c^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F) = \text{Im}(H^{i-1}(\partial\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F) \longrightarrow H_c^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F)) \oplus H_1^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F),$$

which is invariant under the action of the Hecke algebra and no irreducible representation  $\bar{\pi}_\infty \times \pi_f$  which occurs in  $H_1^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F)$  can occur as a sub quotient in  $\text{Im}(H^{i-1}(\partial\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F) \rightarrow H_c^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F))$ .

In the second case we will call the above image of the boundary cohomology the Eisenstein subspace or compactly supported Eisenstein cohomology and denote it by

$$\text{Im}(H^{i-1}(\partial\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F) \longrightarrow H_c^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})) = H_{c, \text{Eis}}^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}).$$

In the first case we can consider the module  $H_{\text{Eis}}^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F) \subset \text{Im}(H^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F) \longrightarrow H^i(\partial(\mathcal{S}_{K_f}^G), \tilde{\mathcal{M}}_F))$  as a submodule in  $H^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F)$  and this submodule is called the Eisenstein cohomology. Under the assumption of the Manin-Drinfeld principle we have a canonical section  $s : H_{\text{Eis}}^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F) \rightarrow H^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F)$ .

If we know the Manin-Drinfeld principle we can ask new questions. We return to the the integral cohomology  $H_?^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathcal{O}_F})$  and map it into the rational cohomology then the image is called  $H_?^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})_{\text{int}} \subset H_?^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F)$  this is also the module which we get if we divide  $H_?^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathcal{O}_F})$  by the torsion. (This may be not true for ? = !.)

Our decompositions above do not induce decomposition on the groups  $H_?^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})_{\text{int}}$ . Whenever we have a decomposition  $H_?^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F) = X \oplus Y$  we can take the intersections  $X_{\text{int}} \cap H_?^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})_{\text{int}}$  and the same for  $Y$  and get a decomposition *up to isogeny*

$$X_{\text{int}} \oplus Y_{\text{int}} \subset H_?^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})_{\text{int}},$$

where up to isogeny means that the left hand side is of finite index in the right hand side.

For instance the Manin-Drinfeld decomposition above yields ( if it exists ) a decomposition up to isogeny

$$H_{c, \text{Eis}}^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})_{\text{int}} \oplus H_1^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})_{\text{int}} \subset H_c^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})_{\text{int}},$$

it is canonical but the direct sum is only of finite index in the right hand side module. The primes dividing the order of the index are called *Eisenstein primes*.

These Eisenstein primes have been studied in the case  $G = \text{Gl}_2/\mathbb{Q}$  but they also seem to play a role in more general situation. The general philosophy is that they are related to the arithmetic of special values of  $L$ -functions. (See [31])

**Here general discussion of a decomposition into saturated submodules**

The same applies to the decomposition of  $H_1^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})_{\text{int}}$  in isotypical summands. We put

$$H_1^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})(\pi_f) \cap H_1^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathcal{O}_F})_{\text{int}} = H_1^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathcal{O}_F})_{\text{int}}(\pi_f).$$

Then we get an decomposition up to isogeny

$$\bigoplus_{\pi_f} H_1^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathcal{O}_F})_{\text{int}}(\pi_f) \subset H_1^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathcal{O}_F})_{\text{int}}. \tag{6.47}$$

It is a very interesting question to learn something about the the structure of the quotient of the right hand side by the left hand side. The structure of this quotient should be related to the arithmetic of special values of  $L$ -functions. (See [Hi]).

**The action of  $\pi_0(G(\mathbb{R}))$**

We have seen that we can choose a maximal torus  $T/\mathbb{Q}$  such that  $T(\mathbb{R})[2]$  normalizes  $K_\infty$ . We know that  $T(\mathbb{R})[2] \rightarrow \pi_0(G(\mathbb{R}))$  is surjective and that  $T(\mathbb{R})[2] \cap G^{(1)}(\mathbb{R}) \subset K_\infty$ . This allows us to define an action of  $\pi_0(G(\mathbb{R}))$  on the various cohomology groups and this action commutes with the action of the Hecke-algebra. Therefore we can decompose any isotypical subspace in a cohomology group into eigenspaces under this action

$$H_?^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F)(\pi_f) = \bigoplus_{\epsilon_\infty} H_?^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F)(\pi_f \times \epsilon_\infty) \tag{6.48}$$

and for the integral lattices we get a decomposition up to isogeny

$$\bigoplus_{\pi_f \times \epsilon_\infty} H_1^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathcal{O}_F})_{\text{int}}(\pi_f \times \epsilon_\infty) \subset H_1^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathcal{O}_F})_{\text{int}} \tag{6.49}$$

**6.3.8 The Gauss-Bonnet formula**

Of course we can be more modest we may only ask for the dimension of the cohomology groups  $H^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Q}})$ . This question can be answered in some cases, for instance we gave the answer  $\text{Sl}_2(\mathbb{Z})$  in section 2.1.3, and we will give the answer in some more cases further down.

If we are even more modest we can ask for the Euler characteristic

$$\chi(H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Q}})) = \sum_i (-1)^i \dim(H^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Q}}))$$

This question has an answer. We assume for the beginning that the subgroup  $K_f$  is neat (See 1.1.2.1) and we also assume that  $K_\infty$  is narrow. Then  $\mathcal{S}_{K_f}^G$  is a disjoint union of locally symmetric spaces on we can choose- in a consistent way- an orientation on  $\mathcal{S}_{K_f}^G$ . On these spaces exists a differential form of highest degree, which is obtained from differential geometric data, this is the Gauss-Bonnet form  $\omega^{GB}$ . Then the Gauss-Bonnet theorem yields that

$$\chi(H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Q}})) = \dim(\mathcal{M}_{\mathbb{Q}}) \int_{\mathcal{S}_{K_f}^G} \omega^{GB} \tag{6.50}$$

(See [22], [79],[68]).

We have a closer look at this formula. We can compute the differential form  $\omega^{GB}$  explicitly. The connected components of  $\mathcal{S}_{K_f}^G$  are of the form  $\Gamma_i \backslash X$  where  $X = G(\mathbb{R})/K_\infty$ . Then the top degree form  $\omega^{GB}$  is a  $G(\mathbb{R})$  invariant form on the symmetric space  $X$ . Since this form is  $G(\mathbb{R})$  invariant it is determined by its value on  $\Lambda^d(\mathfrak{p})$ . On  $\mathfrak{p}$  we have the euclidian metric given by the Killing form and we chose an orientation. These two data provide a second top degree form  $\omega^{Kill}$  on  $\Lambda^d(\mathfrak{p})$ , and hence an invariant form also called  $\omega^{Kill}$  on  $X$ . These two forms are proportional, i.e. we have

$$\omega^{GB} = \kappa_\infty(G)\omega^{Kill}, \quad (6.51)$$

the proportionality factor can be computed from the curvature tensor ( See [22], 2.2 Kobayashi-Nomizu) We have the following

**Proposition 6.3.5.** *The factor  $\kappa_\infty(G)$  is non zero if and only if  $G \times_{\mathbb{Q}} \mathbb{R}$  is an inner form of its compact dual  $G_c/\mathbb{R}$  or in other words  $G \times_{\mathbb{Q}} \mathbb{R}$  has a compact maximal torus. If it is non zero then it is a real number and its sign is  $(-1)^{\frac{d}{2}}$*

We remember that  $G/\mathbb{R} = G \times_{\mathbb{Q}} \mathbb{R}$ , the Lie-algebra  $\mathfrak{g}$  is a  $\mathbb{Q}$ -vector space. The Killing form  $B : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{Q}$  is non degenerate and hence defines a top degree form  $\omega_B$  on  $\Lambda^{d_G} \mathfrak{g}$ . If we use the decomposition  $\mathfrak{g} \otimes \mathbb{R} = \text{Lie}(K_\infty) \oplus \mathfrak{p}$  then we find  $\omega_B = \omega_B^{K_\infty} \wedge \omega^{Kill}$  and hence

$$\int_{K_\infty} \omega_B = \text{vol}_{\omega_B^{K_\infty}}(K_\infty)\omega^{Kill} \quad (6.52)$$

The top degree form  $\omega_B$  which is defined over  $\mathbb{Q}$  also provides invariant measures  $\omega_{B,p}$  on all the groups  $G(\mathbb{Q}_p)$  and also an invariant measure  $\omega_{B,\infty}$  on  $G(\mathbb{R})$ . We can multiply these measures and get the Tamagawa measure

$$\omega_G^{Tam} = \omega_{B,\infty} \times \prod_p \omega_{B,p} = \omega_{B,\infty} \times \omega_{G,f}^{Tam} \quad (6.53)$$

this product is absolutely convergent and provides an  $G(\mathbb{A})$ -invariant measure on  $G(\mathbb{A})$ . This is the *Tamagawa measure* on  $G(\mathbb{A})$ . It is an important fact, that this measure does not depend on the choice of the top degree form  $\omega_B$ . If we multiply  $\omega_B$  by a non zero number  $a \in \mathbb{Q}^\times$  then the local measures get multiplied by  $|a|_\infty$  at infinity and  $|a|_p$  at the finite places. Hence we see that  $\omega_B$  and  $a\omega_B$  yield the same Tamagawa measure. The number

$$\int_{G(\mathbb{Q}) \backslash G(\mathbb{A})} \omega_G^{Tam} = \tau(G)$$

is called the Tamagawa number.

If we have written the Tamagawa measure as a product as in (6.53) we say that the Tamagawa measure is represented by  $\omega_B$ . The remark above tells us that we may replace  $\omega_B$  by any non zero invariant top degree form.

*Now a miracle occurs*

**Theorem 6.3.2.** *If  $G/\mathbb{Q}$  is simply connected then*

$$\int_{G(\mathbb{Q}) \backslash G(\mathbb{A})} \omega_G^{Tam} = \tau(G) = 1$$

This theorem was conjectured by Weil, he ..... For a general semi simple  $G/\mathbb{Q}$  we can consider the universal covering by a simply connected  $\pi : G^{sc}/\mathbb{Q} \rightarrow G/\mathbb{Q}$ , then  $\tau(G)$  is a rational number which can be expressed in terms of Galois cohomology data of the finite kernel of  $\pi$ .

This gives us a way to compute the integral in (6.50). We recall that  $\mathcal{S}_{K_f}^G = \bigcup_i \Gamma_i \backslash G(\mathbb{R})/K_\infty = \bigcup_i \Gamma_i \backslash G(\mathbb{R})/K_\infty \times \underline{x}_i K_f/K_f$  (prop. 8.62) and hence

$$\int_{\mathcal{S}_{K_f}^G} \omega^{GB} = \sum_i \int_{\Gamma_i \backslash X} \omega^{GB} = \kappa_\infty(G) \sum_i \int_{\Gamma_i \backslash X} \omega^{Kill} = \frac{\kappa_\infty(G)}{\text{vol}_{\omega_{K_\infty}^{K_\infty}}(K_\infty)} \sum_i \int_{\Gamma_i \backslash X} \omega_B = \frac{\kappa_\infty(G)}{\text{vol}_{\omega_{K_\infty}^{K_\infty}}(K_\infty) \text{vol}_{\omega_{G,f}^{Tam}}(K_f)} \int_{G(\mathbb{Q}) \backslash G(\mathbb{A})} \omega_G^{Tam} \tag{6.54}$$

Hence we see that for a neat  $K_f$

$$\chi(H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\mathbb{Q})) = \frac{\kappa_\infty(G)}{\text{vol}_{\omega_{K_\infty}^{K_\infty}}(K_\infty) \text{vol}_{\omega_{G,f}^{Tam}}(K_f)} \dim(\mathcal{M}_\mathbb{Q}) \tau(G) \tag{6.55}$$

We study the factor in front. We assume that the level group  $K_f$  is a product over local factors. i.e. we assume  $K_f = \prod_{p:p \text{ prime}} K_p$ . Then clearly

$$\frac{1}{\text{vol}_{\omega_{G,f}^{Tam}}(K_f)} = \prod_p \frac{1}{\text{vol}_{\omega_{B,p}}(K_p)},$$

the message is that the factor in front is a product over local contributions at the places of  $\mathbb{Q}$ , i.e. a local contribution at infinity and a local contribution at each prime.

We study the local factors  $\text{vol}_{\omega_{B,p}}(K_p)$ . Of course we have to tell what  $K_p$  should be. To define such a subgroup  $K_f$  we choose a flat integral structure  $\mathcal{G}/\mathbb{Z}$  (See section 1.2.1) of  $G/\mathbb{Q}$  and define  $K_p$  a congruence subgroup of  $\mathcal{G}(\mathbb{Z}_p)$ . We know that there is a finite set  $\Sigma$  of primes such that for  $p \notin \Sigma$  the following is true

- a) The group scheme  $\mathcal{G} \times \mathbb{Z}_p/\mathbb{Z}_p$  is semi simple and  $K_p = \mathcal{G}(\mathbb{Z}_p)$
- b) The top degree form  $\omega_{B,p}$  on  $\Lambda^{d_G}(\mathfrak{g} \otimes \mathbb{Z}_p)$  is non zero mod  $p$ .

We think that it was Tamagawa who pointed out that under these conditions, i.e.  $p \notin \Sigma$  we have

$$\text{vol}_{\omega_{B,p}}(K_p) = p^{-d_G} \#\mathcal{G}(\mathbb{F}_p) \tag{6.56}$$

For the following we refer to the lectures of R. Steinberg [81]. We recall the well known formula for  $\#\mathcal{G}(\mathbb{F}_p)$ . For the moment we assume that  $G/\mathbb{Q}$  is an inner form of the split  $\mathbb{Q}$ -form, hence  $\mathcal{G} \times \mathbb{F}_p$  is a split Chevalley group. Then it is well known that

$$p^{-d_G} \#\mathcal{G}(\mathbb{F}_p) = (1 - \frac{1}{p})^r \prod_{i=1}^r ((1 + \frac{1}{p} + \dots + \frac{1}{p^{m_i}})) = \prod_{i=1}^r (1 - \frac{1}{p^{m_i+1}}) \tag{6.57}$$

where  $r$  is the rank of  $G/\mathbb{Q}$  ( the dimension of a maximal torus) and the  $m_i$  are so called exponents (See [?], [?] ,,). The expression  $(1 - \frac{1}{p^{m_i+1}}) = \zeta_p(m_i + 1)^{-1}$  where  $\zeta_p(s)$  is the local Euler factor of the Riemann-zeta function at  $p$ .

Hence we get

$$\frac{1}{\text{vol}_{\omega_{G,f}^{\text{Tam}}}(\mathcal{K}_f)} = \left( \prod_{p \in \Sigma} \frac{\prod_{i=1}^r \zeta_p(m_i + 1)^{-1}}{\text{vol}_{\omega_{B,p}}(\mathcal{K}_p)} \right) \prod_{i=1}^r \zeta(m_i + 1) \quad (6.58)$$

Hence we are left with the computation of  $\text{vol}_{\omega_{B,p}}(\mathcal{K}_p)$  at the finitely many ramified places  $p \in \Sigma$ . Of course  $\text{vol}_{\omega_{B,p}}(\mathcal{K}_p) = \frac{\text{vol}_{\omega_{B,p}}(\mathcal{G}(\mathbb{Z}_p))}{[\mathcal{G}(\mathbb{Z}_p) : \mathcal{K}_p]}$ . The computation of  $\text{vol}_{\omega_{B,p}}(\mathcal{G}(\mathbb{Z}_p))$  may become tedious depending on how badly ramified the group scheme  $\mathcal{G} \times \mathbb{Z}_p$  at  $p \in \Sigma$  will be. We know that for  $r \gg 0$

$$\text{vol}_{\omega_{B,p}}(\mathcal{G}(\mathbb{Z}_p)) = \#\mathcal{G}(\mathbb{Z}_p/p^r) p^{a(\omega_{B,p}) - \dim(G)r} \quad (6.59)$$

where  $a(\omega_{B,p})$  is an integer depending on the choice of  $\omega_G^{\text{Tam}}$ . Therefore it is clear that  $\text{vol}_{\omega_{B,p}}(\mathcal{K}_p)$  is a non zero rational number. For a very special case we discuss this computation further down

Therefore we can sum up and get for a split and simply connected  $G/\mathbb{Q}$  the final formula

$$\chi(H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{Q}})) = \frac{\kappa_\infty(G)}{\text{vol}_{\omega_B^{\text{K}\infty}}(\mathcal{K}_\infty)} \left( \prod_{p \in \Sigma} \frac{\prod_{i=1}^r \zeta_p(m_i + 1)^{-1}}{\text{vol}_{\omega_{B,p}}(\mathcal{K}_p)} \right) \dim(\mathcal{M}_{\mathbb{Q}}) \prod_{i=1}^r \zeta(m_i + 1) \quad (6.60)$$

Formulas of this kind have been proved by C. L. Siegel, I. Satake ([79] [68]) and others. For the case of general semi simple groups this is in [22]. The numbers  $m_i$  in [22] are the numbers  $m_i + 1$  here.

We also discuss the case that  $G/\mathbb{Q}$  is not an inner form of the split form  $G_0/\mathbb{Q}$ , we assume that  $G_0/\mathbb{Q}$  is simple, let  $\Phi$  be its Dynkin diagrams. It is one of the form  $A_n, n \geq 2, D_n, n \geq 4$  or  $E_6$ . In this case there is a unique normal extension  $L/\mathbb{Q}$  and a faithful action of  $\text{Gal}(L/\mathbb{Q})$  on  $\Phi$ , in other words an inclusion  $j : \text{Gal}(L/\mathbb{Q}) \hookrightarrow \text{Aut}(\Phi)$ . For these above diagrams the group of automorphisms is of order 2 except we are in the case  $D_4$ , in this case it is the symmetric group in three letters ( See for instance [81] , Chap. 10, p.85).

Again we find a finite set  $\Sigma$  of primes such that for  $p \notin \Sigma$  the group  $\mathcal{G} \times \mathbb{F}_p$  is semi simple, but possibly only quasisplit, assume we are in the case that  $[L : \mathbb{Q}] = 2$ , this is certainly the case if  $G_0/\mathbb{Q}$  is not of type  $D_4$ . If now  $p \notin \Sigma$  and  $p$  splits, then the formula (6.57) is still valid. If  $p$  is inert in  $L$ , i.e.  $\mathcal{O}_L/p\mathcal{O}_L = \mathbb{F}_{p^2}$  then we get from [81] the recipe how to modify the right hand side in (6.57).

a) In case  $G_0/\mathbb{Q}$  is of type  $A_n, D_n$  and  $n$  odd or  $E_6$  then we have to replace the factor  $(1 - \frac{1}{p^{m_i+1}})$  by  $(1 + \frac{1}{p^{m_i+1}})$  in case  $m_i$  is even.

b) In the case  $D_n$  and  $n$  even then  $m_i = n - 1$  occurs twice and we have to replace the factor

$$(1 - \frac{1}{p^n})^2 \text{ by } (1 - \frac{1}{p^n})(1 + \frac{1}{p^n}).$$

Finally we come to the case  $D_4$  and  $\text{Gal}(L/\mathbb{Q})$  is cyclic of order 3 or the symmetric group in three letters. Let  $\mathfrak{P}$  be a prime ideal in  $\mathcal{O}_L$  which lies over

$p$ . Then  $\mathcal{O}_L/\mathfrak{P} = \mathbb{F}_p, \mathbb{F}_{p^2}$  or  $\mathbb{F}_{p^3}$ . The first two cases are handled by b) . In the third case we have to replace (See[81], Table on p. 105)

$$\left(1 - \frac{1}{p^4}\right)^2 \text{ by } 1 + \frac{1}{p^4} + \frac{1}{p^8} = (1 - \zeta_3 p^{-4})(1 - \zeta_3^2 p^{-4})$$

where  $\zeta_3$  is a third root of unity ( $\neq 1$ ).

Now it is clear how we have to modify the formula (6.60). If  $[L : \mathbb{Q}] = 2$  we have the character  $\chi_{L/\mathbb{Q}}$  corresponding to this extension For  $p \notin \Sigma$  we have  $\chi_{L/\mathbb{Q}}(p) = -1$  if and only if  $p$  does not split in  $L$ . Attached to this character we have the Dirichlet  $L$ - function

$$L(\chi_{L/\mathbb{Q}}, s) = \prod_p \frac{1}{1 - \chi_{L/\mathbb{Q}}(p)p^{-s}}.$$

This means that in formula (6.60) we have to replace the factors  $\zeta(m_1 + 1)$  by  $L(\chi_{L/\mathbb{Q}}^{m_i+1}, m_i + 1)$ . If we agree that for  $m_i + 1$  even  $L(\chi_{L/\mathbb{Q}}^{m_i+1}, s) = \zeta(s)$  then (6.60) in case  $[L : \mathbb{Q}] = 2$  becomes

$$\begin{aligned} & \chi(H^\bullet(\mathcal{S}_{K'_f}^G, \tilde{\mathcal{M}}_{\mathbb{Q}})) = \\ & \frac{\kappa_\infty(G)}{\text{vol}_{\omega_{K_\infty}}(K_\infty)} \left( \prod_{p \in \Sigma} \frac{\prod_{i=1}^r L_p(\chi_{L/\mathbb{Q}}^{\epsilon(m_i)}, m_i+1)^{-1}}{\text{vol}_{\omega_{B,p}}(K_p)} \right) \dim(\mathcal{M}_{\mathbb{Q}}) \prod_{i=1}^r L(\chi_{L/\mathbb{Q}}^{\epsilon(m_i)}, m_i + 1) \end{aligned} \tag{6.61}$$

here we have to say what  $\epsilon(m_i)$  is.

a) If  $G_0/\mathbb{Q}$  is of type  $A_n, E_6$  or  $D_n$  with  $n$  odd then  $\epsilon(m_i) = 0$  if  $m_i$  is odd and  $\epsilon(m_i) = 1$  if  $m_i$  is even.

b) If  $G_0/\mathbb{Q}$  is of type  $D_n$  with  $n$  even then the exponent  $n - 1 = m_{n/2} = m_{n/2+1}$  occurs twice. In this case  $\epsilon(m_i) = 0$  for  $i \neq \frac{n}{2}$  or  $\frac{n}{2} + 1$  and  $\epsilon(m_{n/2}) = 0, \epsilon(m_{n/2+1}) = 1$  (Hence we see that in the case  $D_n$  we have exactly one genuine Dirichlet  $L$ - function in the product.)

c) Finally we look at the case  $D_4$  and assume  $\text{Gal}(L/\mathbb{Q})$  is the symmetric group in three letters or cyclic of order three. In this case we have an irreducible representation  $\rho_2 : \text{Gal}(L/\mathbb{Q}) \rightarrow \text{GL}_2(\mathbb{Q})$ ,. to this representation we attach the Artin- $L$ - function. Then it is clear that we have to replace the factor  $\zeta(4)^2$  by the factor  $L(\rho_2, 4)$ .

This implies of course, that for a covering  $\mathcal{S}_{K'_f}^G \rightarrow \mathcal{S}_{K_f}^G$ , where  $K'_f \subset K_f$  and both groups are neat, we get

$$\chi(\mathcal{S}_{K'_f}^G, \tilde{\mathcal{M}}) = \chi(H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})[K_f : K'_f]),$$

a fact which also follows easily from topological considerations.

This leads us following C.T.C. Wall- to introduce the orbifold Euler characteristic for a not necessarily neat  $K_f$  by

$$\chi_{\text{orb}}(H^\bullet(\mathcal{S}_{K'_f}^G, \tilde{\mathcal{M}})) = \frac{1}{[K'_f : K_f]} \chi(\mathcal{S}_{K'_f}^G, \tilde{\mathcal{M}}) \tag{6.62}$$

where  $K'_f \subset K_f$  is a neat subgroup of finite index. The orbifold Euler characteristic may differ from the Euler characteristic  $\chi(H^\bullet(\mathcal{S}_{K'_f}^G, \tilde{\mathcal{M}}))$  by a sum of contributions coming from the set of fixed points of the  $\Gamma_i$  on  $X$  (See 1.1.2.1). Hence the formulae (6.60) and (6.61) remain valid without the assumption  $K_f$  neat once we replace  $\chi(\mathcal{S}_{K'_f}^G, \tilde{\mathcal{M}})$  by  $\chi_{\text{orb}}(H^\bullet(\mathcal{S}_{K'_f}^G, \tilde{\mathcal{M}}))$  on the left hand side.

The Gauss-Bonnet formula implies that the orbifold Euler characteristic is linear in  $\dim(\mathcal{M}_{\mathbb{Q}})$ . But this is an obvious consequence of our considerations in section 2.1.2. We compute the cohomology from the Čech complex given by an orbiconvex covering. If our group  $K_f$  is neat then all the terms  $\mathcal{M}(U_i)$  in (??) are of the form  $\tilde{\mathbb{Q}}(U_i) \otimes \mathcal{M}$  where of course  $\tilde{\mathbb{Q}}$  is the sheaf obtained from the trivial one-dimensional representation. The differentials in the complex are only acting on the first factor and hence

$$C^\bullet(\mathfrak{U}, \tilde{\mathcal{M}}) \xrightarrow{\sim} C^\bullet(\mathfrak{U}, \tilde{\mathbb{Q}}) \otimes \mathcal{M}.$$

Since  $\chi(\mathcal{S}_{K'_f}^G, \tilde{\mathcal{M}}) = \chi(C^\bullet(\mathfrak{U}, \tilde{\mathcal{M}})) = \chi(C^\bullet(\mathfrak{U}, \tilde{\mathbb{Q}})) \times \dim \mathcal{M}$  the linearity follows.

### Gauss-Bonnet and the special values

We discuss some arithmetic consequences of the Gauss-Bonnet formula. By definition  $\chi_{\text{orb}}(H^\bullet(\mathcal{S}_{K'_f}^G, \tilde{\mathcal{M}})) \in \mathbb{Q}$ , hence we can conclude that the right side must be a rational number. This argument gives us non trivial consequences for the special values  $\zeta(m_i + 1)$ , but only if the curvature factor  $\kappa_\infty(G) \neq 0$ . We analyse this condition.

Remember that we want to assume that  $G/\mathbb{Q}$  is absolutely simple, We consider the base extension  $G \times_{\mathbb{Q}} \mathbb{R}$ , then the complex conjugation  $\mathbf{c}$  induces an involution on  $\Phi$ . Now it is known

**Proposition 6.3.6.** *The Dynkin diagrams is of the form  $A_1, B_n, C_n, E_7, E_8, F_4, G_2$  have trivial automorphism groups and  $G \times_{\mathbb{Q}} \mathbb{R}$  is an inner form of its compact dual. The Dynkin diagram  $D_n$  has non trivial automorphisms. In this case  $G \times_{\mathbb{Q}} \mathbb{R}$  is an inner form of its compact dual if*

- a) *the complex conjugation  $\mathbf{c}$  acts trivially on  $\Phi$  if  $n$  is even.*
- b) *the complex conjugation  $\mathbf{c}$  acts non trivially on  $\Phi$  if  $n$  is odd.*

*In the remaining cases  $G \times_{\mathbb{Q}} \mathbb{R}$  is an inner form of its compact dual if  $\mathbf{c}$  acts non trivially.*

*In the cases where  $\mathbf{c}$  acts trivially all the  $m_i$  are odd.*

Now we have a look at the numbers  $\zeta(m_i + 1)$  and  $L(\chi_{L/\mathbb{Q}}^{\epsilon(m_i)}, m_i + 1)$  on the right hand side. Euler has shown that

$$\text{For even integers } m \geq 2 \text{ the numbers } \frac{\zeta(m)}{\pi^m} \in \mathbb{Q}. \quad (6.63)$$

We also know that the matching answer for  $L(\chi_{L/\mathbb{Q}}, m)$  depends on the parity  $p(\chi_{L/\mathbb{Q}})$  of  $\chi_{L/\mathbb{Q}}$ , where  $p(\chi_{L/\mathbb{Q}}) = 1$  if  $L_\infty = \mathbb{C}$  and 0 else. If we write  $L = \mathbb{Q}[\sqrt{d}]$  we have  $p(\chi_{L/\mathbb{Q}}) = 1$  if and only if  $d < 0$ . Then we have the more general result

$$\text{For integers } m > 0 \text{ and } m + p(\chi_{L/\mathbb{Q}}) \text{ even the numbers } \frac{L(\chi_{L/\mathbb{Q}}, m)}{\pi^m \sqrt{|d|}} \in \mathbb{Q} \quad (6.64)$$

The values  $\zeta(m)$  for  $m = 2, 4, \dots$  or  $L(\chi_{L/\mathbb{Q}}, m)$  for  $m > 0$ ;  $m + p(\chi_{L/\mathbb{Q}}) \equiv 0 \pmod{2}$  are the so called *special values* of the Riemann  $\zeta$  function or more generally Dirichlet  $-L$  function.

For the following we refer to [60] Chapter VII. We still have the functional equation. We introduce the Euler-factor at infinity

$$L_\infty(\chi_{L/\mathbb{Q}}, s) := \left(\frac{|d|}{\pi}\right)^{\frac{s}{2}} \Gamma\left(\frac{s + p(\chi_{L/\mathbb{Q}})}{2}\right)$$

where  $\Gamma$  is the Gamma-function and here we also assume that  $d$  is squarefree. We define the completed  $L$ -function

$$\Lambda(\chi_{L/\mathbb{Q}}, s) = L_\infty(\chi_{L/\mathbb{Q}}, s)L(\chi_{L/\mathbb{Q}}, s) \tag{6.65}$$

If  $\chi_{L/\mathbb{Q}}$  is not trivial then  $\Lambda(\chi_{L/\mathbb{Q}}, s)$  is holomorphic in the entire complex plane, if  $\chi_{L/\mathbb{Q}}$  is the trivial character then we get the completed Riemann  $\zeta$  function  $\Lambda(s) = \frac{\Gamma(s/2)}{(2\pi)^{s/2}}\zeta(s)$ . It is meromorphic function in the entire complex plane and has two simple poles at  $s = 1$  and  $s = 0$ . For this completed  $L$  function we have the functional equation (See [60], Chap. VII, Theorem 2.8)

$$\Lambda(\chi_{L/\mathbb{Q}}, s) = W(\chi)\Lambda(\chi_{L/\mathbb{Q}}, 1 - s) \tag{6.66}$$

where in this special case  $W(\chi)$  is an integral power of  $i = \sqrt{-1}$  and we observe that  $\chi_{L/\mathbb{Q}}$  is real.

This tells us something about the special values at negative integers: For  $m > 0$ ;  $m + p(\chi_{L/\mathbb{Q}}) \equiv 0 \pmod{2}$  we get

$$L(\chi_{L/\mathbb{Q}}, 1 - m) = \frac{L_\infty(\chi_{L/\mathbb{Q}}, m)}{L_\infty(\chi_{L/\mathbb{Q}}, 1 - m)}L(\chi_{L/\mathbb{Q}}, m). \tag{6.67}$$

Using the functional equation for the  $\Gamma$ -function and  $\Gamma(\frac{1}{2}) = \sqrt{\pi}$  we get for ratio of the two Euler factors at infinity

$$\left|\frac{d}{\pi}\right|^{m-\frac{1}{2}} \frac{\Gamma\left(\frac{m+p(\chi_{L/\mathbb{Q}})}{2}\right)}{\Gamma\left(\frac{1-m-p(\chi_{L/\mathbb{Q}})}{2}\right)} = \left(\frac{|d|}{\pi}\right)^m \frac{\sqrt{\pi}}{\sqrt{|d|}} \frac{\Gamma(m)}{\sqrt{\pi}2^{m-1}}$$

and if we insert this in (6.67) we get

$$L(\chi_{L/\mathbb{Q}}, 1 - m) = |d|^m \frac{\Gamma(m)}{2^{m-1}} \frac{L(\chi_{L/\mathbb{Q}}, m)}{\sqrt{|d|\pi^m}} \in \mathbb{Q}^\times \tag{6.68}$$

Finally we have understand the contribution from the infinite place, i.e. the term  $\frac{\kappa_\infty(G)}{\text{vol}_{\omega_B^{K_\infty}}(K_\infty)}$ . This term has been computed in [22] in the case of split groups  $G/F$ , where  $F$  is a totally real number field. In our case  $F = \mathbb{Q}$  this gives

$$\frac{\kappa_\infty(G)}{\text{vol}_{\omega_B^{K_\infty}}(K_\infty)} = (-1)^{\frac{g}{2}} \frac{\prod_{i=1}^r (m_i + 1)!}{\#W_{K_\infty} \pi^{r+\sum_{i=1}^r m_i}} = \frac{a_\infty(G)}{\pi^{r+\sum_{i=1}^r m_i}} \tag{6.69}$$

here  $W_{K_\infty}$  is the Weyl group of  $K_\infty$  and  $a_\infty(G)$  is an explicitly computable rational number. Hence equation (6.60) becomes

$$\chi_{\text{orb}}(H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})) = a_\infty(G)a(G, K_f) \dim(\mathcal{M}_\mathbb{Q}) \prod_{i=1}^r \frac{\zeta(m_i + 1)}{\pi^{m_i + 1}} \text{ with } a(G, K_f) \in \mathbb{Q}^\times, \quad (6.70)$$

the non zero rational number  $a(G, K_f)$  can be explicitly computed.

The formula becomes much nicer if we apply the functional equation for the  $\zeta$ -function and look at the special values at the negative arguments.

If  $\mathcal{G}_0/\mathbb{Z}$  is split semi simple and simply connected and if we choose  $K_f = \prod_p \mathcal{G}_0(\mathbb{Z}_p)$  then the computation in [22] p. 452-453 gives

$$\chi_{\text{orb}}(H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})) = \frac{\#W_G}{2^r \#W_{K_\infty}} \dim(\mathcal{M}_\mathbb{Q}) \prod_{i=1}^r \zeta(-m_i) \quad (6.71)$$

Now we see that the Gauss-Bonnet formula reproves Euler's rationality results. We consider the split groups schemes of type  $A_1$  and  $B_n$  or  $C_n$ . In these cases the exponents are the odd numbers  $1, 3, \dots, 2n - 1$ . (See [8] , Planche II, III) If we apply the formula for  $A_1$  we get  $\zeta(2)/\pi^2 \in \mathbb{Q}^\times$ . Applying it for  $B_2$  or  $C_2$  gives  $\zeta(2)/\pi^2 \times \zeta(4)/\pi^4 \in \mathbb{Q}^\times$  hence  $\zeta(4)/\pi^4 \in \mathbb{Q}^\times$ . Clearly we get Euler's result by induction.

We also get the corresponding rationality results for the Dirichlet- $L$  functions attached to quadratic characters  $\chi_{L/\mathbb{Q}}$ . We consider absolutely simple groups  $G/\mathbb{Q}$  with non trivial action of  $\text{Gal}(L/\mathbb{Q})$  on their Dynkin diagram  $\Phi$ . If  $L/\mathbb{Q}$  is real quadratic, i.e.  $p(\chi_{L/\mathbb{Q}}) = 0$  then the situation is the essentially same as in the case of a trivial character.

If  $L = \mathbb{Q}[\sqrt{d}]$  is imaginary quadratic we choose a group  $G/\mathbb{Q}$  of type  $A_n$  with  $n \geq 2$ , or  $D_n$  with  $n$  odd, or  $E_6$ . In these cases the Gauss-Bonnet formula becomes

$$\chi_{\text{orb}}(H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})) = a_\infty(G)a(G, K_f) \dim(\mathcal{M}_\mathbb{Q}) \prod_{i=1}^r L(\chi_{L/\mathbb{Q}}^{\epsilon(m_i)}, -m_i) \quad (6.72)$$

Hence see that for an imaginary quadratic extension  $L/\mathbb{Q}$  we can start from groups  $G/\mathbb{Q}$  of type  $A_n$  for  $n = 2, 3, 4, \dots$  to prove (??).

Of course we can also consider the case that  $G(\mathbb{R})$  is compact. (See ??) in this case the Gauss-Bonnet theorem is a tautology, the quotient

$$\mathcal{S}_{K_f}^G = G(\mathbb{Q}) \backslash * \times G(\mathbb{A}_f) / K_f$$

is a finite set. If  $K_f$  is neat then

$$\chi(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}) = \dim H^0(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}) = \#(G(\mathbb{Q}) \backslash * \times G(\mathbb{A}_f) / K_f) \dim(\mathcal{M}_\mathbb{Q}).$$

If  $K_f^0 \subset G(\mathbb{A}_f)$  is not then we choose a normal neat subgroup  $K_f \subset K_f^0$ , and we consider the diagram

$$\begin{array}{ccc} G(\mathbb{A}_f) / K_f & \xrightarrow{\pi_{K_f}} & G(\mathbb{Q}) \backslash G(\mathbb{A}_f) / K_f = \mathcal{S}_{K_f}^G \\ q_0 \downarrow & & q_1 \downarrow \\ G(\mathbb{A}_f) / K_f^0 & \xrightarrow{\pi_{K_f^0}} & G(\mathbb{Q}) \backslash G(\mathbb{A}_f) / K_f^0 = \mathcal{S}_{K_f^0}^G \end{array} \quad (6.73)$$

We want to compute the orbifold Euler characteristic

$$\chi_{\text{orb}}(\mathcal{S}_{K_f^0}^G, \tilde{\mathcal{M}}) = \frac{1}{[K_f^0 : K_f]} \chi(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})$$

to do this we have to understand the fibers of  $q_1$ . For a point  $\underline{x} \in \mathcal{S}_{K_f^0}^G$  we pick a point  $\underline{y} \in \pi_{K_f^0}^{-1}(\underline{x})$ . Then we choose a point  $\underline{y}_1 \in q_0^{-1}(\underline{y})$ , now we can identify  $q_0^{-1}(\underline{y}_1) = K_f^0/K_f$  by  $\underline{k}_f \mapsto \underline{y}_1 \underline{k}_f$ . If we apply  $\pi_{K_f}$  to the fiber  $q_0^{-1}(\underline{y}_1)$  we get the fiber  $q_1^{-1}(\underline{y})$ . Now two points  $\underline{y}_1 \underline{k}_f, \underline{y}_1 \underline{k}'_f$  map to the same point in  $q_1^{-1}(\underline{y})$  if there is a  $\gamma \in G(\mathbb{Q})$  such that  $\gamma \underline{y}_1 \underline{k}_f K_f^0 = \underline{y}_1 \underline{k}'_f K_f^0$ . Since  $K_f$  was a normal subgroup this means that  $\gamma \underline{y}_1 \underline{k}'_f \underline{k}_f^{-1} \in \underline{y}_1 K_f^0$  and hence  $\gamma \in \underline{y}_1 K_f^0 \underline{y}_1^{-1}$ . Since  $K_f$  is neat we get an injection

$$\Gamma_{\underline{y}_1} := G(\mathbb{Q}) \cap \underline{y}_1 K_f^0 \underline{y}_1^{-1} / \underline{y}_1 K_f \underline{y}_1^{-1} \hookrightarrow \underline{y}_1 K_f^0 \underline{y}_1^{-1} / \underline{y}_1 K_f \underline{y}_1^{-1} \xrightarrow{\sim} K_f^0 / K_f$$

Now it is easy to see that the conjugacy class of the finite subgroup  $\Gamma_{\underline{y}_1} \subset K_f^0 / K_f$  only depends on  $\underline{x} \in \mathcal{S}_{K_f^0}^G$  and not on the choices  $\underline{y}$  or  $\underline{y}_1$ . Therefore all the  $\Gamma_{\underline{y}_1}$  are isomorphic and we put  $\#\Gamma_{\underline{x}} := \#\Gamma_{\underline{y}_1}$ .

Now we see

$$\chi_{\text{orb}}(\mathcal{S}_{K_f^0}^G, \tilde{\mathcal{M}}) = \frac{1}{[K_f^0 : K_f]} \left( \sum_{\underline{x} \in \mathcal{S}_{K_f^0}^G} \sum_{\underline{x}_1 \in q_1^{-1}(\underline{x})} \dim_{\mathbb{Q}}(\mathcal{M}) \right) = \left( \sum_{\underline{x} \in \mathcal{S}_{K_f^0}^G} \frac{1}{\#\Gamma_{\underline{x}}} \right) \dim(\mathcal{M}_{\mathbb{Q}}) \tag{6.74}$$

We see that the orbifold system  $\tilde{\mathcal{M}}$  enters only by its dimension. This changes if we look at the Euler characteristic itself. Then we get obviously

$$\chi(\mathcal{S}_{K_f^0}^G, \tilde{\mathcal{M}}) = \sum_{\underline{x} \in \mathcal{S}_{K_f^0}^G} \dim(\mathcal{M}_{\mathbb{Q}}^{\Gamma_{\underline{x}}}) \tag{6.75}$$

and we notice that  $G$ -module structure of  $\mathcal{M}$  matters. Since in general  $\frac{\dim(\mathcal{M}_{\mathbb{Q}})}{\#\Gamma_{\underline{x}}} \neq \dim(\mathcal{M}^{\Gamma_{\underline{x}}})$  we see that we should expect  $\chi_{\text{orb}}(\mathcal{S}_{K_f^0}^G, \tilde{\mathcal{M}}) \neq \chi(\mathcal{S}_{K_f^0}^G, \tilde{\mathcal{M}})$  in general, for instance if  $\mathcal{M} = \mathbb{Q}$ . Of course the formulas (6.60), (6.61) also apply in this situation. But here we have the advantage that the curvature factor  $\kappa_{\infty}(G) = 1$ .

There are cases where  $\mathcal{G}/\mathbb{Z}$  is semi-simple and simply connected and  $\mathcal{G}(\mathbb{R})$  is compact. In this case the computation in [22] gives us a variant of equation (6.71)

$$\chi_{\text{orb}}(\mathcal{S}_{K_f}^G, \mathbb{Q}) = \frac{1}{2^r} \prod_{i=1}^r \zeta(-m_i) \tag{6.76}$$

To give an example we consider  $n$ -dimensional unimodular lattices. An unimodular lattice  $L$  is a free  $\mathbb{Z}$ -module, which is equipped with a symmetric bilinear form  $F : L \times L \rightarrow \mathbb{Z}$  which has the following properties

- a) It is positive definite, i.e.  $F(x, x) > 0$  for all  $x \neq 0$
- b) For any saturated (See (??))  $x \in L$  we find a  $y \in L$  such that  $F(x, y) = 1$ .

c) The values  $F(x, x)$  are even.

Then we know that  $n \equiv 0 \pmod{8}$  (See ???) and the group  $\mathrm{SO}(F)/\mathbb{Z}$  is indeed a semi simple. (If we consider the base extension  $\mathrm{SO}(F) \times_{\mathbb{Z}} \mathbb{Z}_p$  for any prime  $p$  the group scheme is isomorphic to  $\mathrm{SO}(\frac{n}{2}, \frac{n}{2})/\mathbb{Z}_p$ ). This group scheme is not simply connected we have a degree 2 covering  $\mathcal{G}/\mathbb{Z} = \mathrm{Spin}(F)/\mathbb{Z} \rightarrow \mathrm{SO}(F)/\mathbb{Z}$ . Then equation (6.71) yields

$$\frac{1}{2^{\frac{n}{2}}} \prod_i \zeta(-m_i) = \sum_{\mathbf{x} \in \mathcal{S}_{K_f}^G} \frac{1}{\#\Gamma_{\mathbf{x}}} \quad (6.77)$$

We do this for  $n = 8$ . In this case the root lattice  $(L_8, F_8)$  of  $E_8$  (See [?], or infinitely many other references) satisfies the conditions a), b), c) above and we get

$$\frac{1}{2^4} \zeta(-1)\zeta(-3)\zeta(-3)\zeta(-5) = \frac{1}{696729600} = \sum_{\mathbf{x} \in \mathcal{S}_{K_f}^G} \frac{1}{\#\Gamma_{\mathbf{x}}} \quad (6.78)$$

This implies that  $\mathcal{S}_{K_f}^G = \mathcal{G}(\mathbb{Q}) \setminus * \times \mathcal{G}(\mathbb{A}_f) / \mathcal{G}(\hat{\mathbb{Z}})$  consists of one element, namely the identity  $\mathbf{x}_1$  and

$$\Gamma_{\mathbf{x}_1} = \mathcal{G}(\mathbb{Z}) = \mathcal{G}(\mathbb{Q}) \cap \mathcal{G}(\hat{\mathbb{Z}}).$$

We have the homomorphism  $\mathcal{G}(\mathbb{Z}) \rightarrow \mathrm{SO}(F)(\mathbb{Z})$  its kernel is the group  $\mu_2 = \{\pm 1\}$ , and a simple calculation shows that the cokernel is also  $\{\pm 1\}$ , Since  $(L_8, F_8)$  is the root lattice of  $E_8$  we get that  $\mathrm{SO}(F_8)(\mathbb{Z})$  is the Weyl  $W(E_8)$  group of  $E_8$ . The order of the Weyl group of  $E_8$  is 696729600, hence we have verified equation (6.71) in this particular case.

If we play the same game for  $n = 16$  then we start from the lattice  $L_8 \oplus L_8$ . The automorphism group of this lattice is  $\Gamma_{\mathbf{x}_1} \times \Gamma_{\mathbf{x}_1} \times \mathbb{Z}/2\mathbb{Z}$ , we may flip the two summands. Then

$$\frac{1}{2^8} \zeta(-1)\zeta(-3)\zeta(-5)\zeta(-7)^2\zeta(-9)\zeta(-11)\zeta(-13) = \sum_{\mathbf{x}} \frac{1}{\#\Gamma_{\mathbf{x}}} \quad (6.79)$$

On the right hand side we have the summand  $\frac{1}{2 * 696729600^2}$  we subtract it and get

$$\frac{1}{2^8} \zeta(-1) \dots \zeta(-13) - \frac{1}{2 * 696729600^2} = \frac{1}{685597979049984000}$$

Hence we see that  $\mathcal{S}_{K_f}^G$  consists of exactly two elements, we have the lattice  $L_8 \oplus L_8$  and still another one. This has been discovered by E. Witt in [88]. In the same paper Witt mentions that he has found more then 10 different lattices for  $n = 24$ .

The case  $n = 24$  was solved by Niemeier in [61], he showed that there exactly 24 different lattices, one of them is the famous Leech lattice (See editors note to the paper [88]).

Unimodular lattices are studied intensively in the the book of G. Chenevier-J. Lannes [10]

We get also get (semi)- simple group schemes  $\mathcal{G}/\mathbb{Z}$  with  $\mathcal{G}(\mathbb{R})$  compact if start from a Dynkin diagram for which the simply connected group has trivial center. This follows from the Hasse principle. Hence we can find such a  $\mathcal{G}/\mathbb{Z}$  of type  $E_8$ . Then we get

$$\begin{aligned} \frac{1}{2^8} \zeta(-1)\zeta(-7)\zeta(-11)\zeta(-13)\zeta(-17)\zeta(-19)\zeta(-23)\zeta(-29) = \\ \frac{2155741910416889170788798426985697}{154705492508859411569049600000} = \sum_{x \in S_{K_f}^G} \frac{1}{\#\Gamma_x} \end{aligned} \tag{6.80}$$

**The comparison**

If we have two semi simple groups  $G/\mathbb{Q}, G'/\mathbb{Q}$  which are inner forms of each other, then for both groups the product of  $L$ - values in (6.72) is the same. Hence we see that the ratio  $\chi_{\text{orb}}(S_{K_f}^G, \tilde{\mathcal{M}})/\chi_{\text{orb}}(S_{K'_f}^{G'}, \tilde{\mathcal{M}}')$  of the Euler characteristics is a number which can be computed from comparing local data at a finite number of primes, this is sometimes called the Hirzebruch proportionality principle.

We want to be a little more precise. We need to find a way to compare the groups  $K_f$  and  $K'_f$ . To make such a comparison possible, we use the ideas of Bruhat-Tits.

We start from any extension of  $G/\mathbb{Q}, G'/\mathbb{Q}$  to flat group schemes over  $\mathbb{Z}$ . (See section 1.2.1). For all primes  $p$  outside a finite set  $\Sigma$  these two extensions will be semi simple at  $p$ . Then we get semi simple extensions  ${}^*\mathcal{G}/(\text{Spec}(\mathbb{Z}) \setminus \Sigma)$  and  ${}^*\mathcal{G}'/(\text{Spec}(\mathbb{Z}) \setminus \Sigma)$ . At the primes  $p \in \Sigma$  we choose extensions  $G/\mathbb{Q}, G'/\mathbb{Q}$  to flat, smooth Bruhat-Tits group schemes  $\mathcal{G} \times_{\mathbb{Z}} \mathbb{Z}_p$  and  $\mathcal{G}' \times_{\mathbb{Z}} \mathbb{Z}_p$ . We require that the two group schemes  $\mathcal{G} \times_{\mathbb{Z}} \mathbb{Z}_p$  and  $\mathcal{G}' \times_{\mathbb{Z}} \mathbb{Z}_p$  are locally isomorphic for the etale topology. In less educated language this means that we can find a finite unramified extension  $F_p/\mathbb{Q}_p$  such that  $\mathcal{G} \times_{\mathbb{Z}_p} \mathcal{O}_{F,p}$  and  $\mathcal{G}' \times_{\mathbb{Z}_p} \mathcal{O}_{F,p}$  become isomorphic. Then we can use these extensions to extend  ${}^*\mathcal{G}, {}^*\mathcal{G}'$  to flat, smooth group scheme  $\mathcal{G}/\mathbb{Z}, \mathcal{G}'/\mathbb{Z}$  (We will give an example further down).

Now we may choose  $K_p = \mathcal{G}(\mathbb{Z}_p), K'_p = \mathcal{G}'(\mathbb{Z}_p)$  and  $K_f = \prod_p K_p, K'_f = \prod_p K'_p$ . At a finite set of primes we may modify our choice and take full congruence subgroups  $K_p = \mathcal{G}(\mathbb{Z}_p)(p^r), K'_p = \mathcal{G}'(\mathbb{Z}_p)(p^r)$ . This makes it clear that -up to a power of  $p$  -the ratio

$$\frac{\text{vol}_{\omega_{B,p}}(K_p)}{\text{vol}_{\omega_{B,p}}(K'_p)} = p^{\delta_p(G,G')} \frac{\#\mathcal{G}(\mathbb{F}_p)}{\#\mathcal{G}'(\mathbb{F}_p)}, \tag{6.81}$$

the exponent  $\delta(G, G')$  is explicitly computable, and the orders of the finite groups follow from Bruhat-Tits. The computation is basically straightforward but not completely trivial.

We discuss an example. Let  $G/\mathbb{Q} = \text{Sl}_2/\mathbb{Q}$ , we choose a prime  $p \equiv 3 \pmod 4$  and we consider the division algebra  $D(-p, -1)$  (see section 1.1.3) and put  $G'/\mathbb{Q} = D^{(1)}(-p, -1)$  the norm one group of this division algebra. We put  $L = \mathbb{Q}[\sqrt{-1}]$  then we get (see (1.15))

$$G'(\mathbb{Q}) = \{x \in \text{Sl}_2(L) \mid x = \begin{pmatrix} 0 & -p \\ 1 & 0 \end{pmatrix} \sigma(x) \begin{pmatrix} 0 & -p \\ 1 & 0 \end{pmatrix}^{-1} \} \tag{6.82}$$

and with  $x = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  this means

$$G'(\mathbb{Q}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} \sigma(c) & -p\sigma(c) \\ p^{-1}\sigma(b) & \sigma(a) \end{pmatrix} \right\} \quad (6.83)$$

and hence

$$G'(\mathbb{Q}) = \left\{ \begin{pmatrix} a & pb \\ -\sigma(b) & \sigma(a) \end{pmatrix} \mid a\sigma(a) + pb\sigma(b) = 1; a, b \in L \right\} \quad (6.84)$$

We choose extensions  $\mathcal{G}'/\mathbb{Z}$  and  $\mathcal{G}/\mathbb{Z}$  of our groups, For any prime  $\ell$  we require that

$$\mathcal{G}'(\mathbb{Z}_\ell) = \left\{ \begin{pmatrix} a & pb \\ -\sigma(b) & \sigma(a) \end{pmatrix} \mid a\sigma(a) + pb\sigma(b) = 1; a, b \in \mathbb{Z}_\ell[i] \right\}$$

For  $\ell \neq 2$  or  $\ell \neq p$  it is easy to see that  $\mathcal{G}/\mathbb{Z}_\ell \xrightarrow{\sim} \mathrm{Sl}_2/\mathbb{Z}_\ell$  and hence semi simple. For  $\ell = 2$  we have to use  $p \equiv 3 \pmod{4}$  and hence  $-p \in N_{L/\mathbb{Q}}(\mathbb{Z}_2[i]^\times)$ . Then it is again easy to see that  $\mathcal{G}'/\mathbb{Z}_\ell$  must be semi-simple. It remains the case  $\ell = p$ . In this case  $p$  does not split in  $\mathbb{Z}[i]$  and hence  $\mathbb{Z}[i]/(p) = \mathbb{F}_{p^2}$ . Hence we see that the reduction  $\pmod{p}$  gives us

$$\mathcal{G}'(\mathbb{F}_p) = \left\{ \begin{pmatrix} a & 0 \\ -\sigma(b) & \sigma(a) \end{pmatrix} \mid a\sigma(a) = 1, a, b \in \mathbb{F}_{p^2} \right\}$$

Now we see that  $\mathcal{G}' \times_{\mathbb{Z}} \mathbb{F}_p$  is not semi-simple, it has a non trivial unipotent radical, which is isomorphic to  $R_{\mathbb{F}_{p^2}/\mathbb{F}_p}(\mathbb{G}_a)$ . We get  $\#\mathcal{G}'(\mathbb{F}_p) = (p+1)p^2$ .

It is clear that this extension of  $\mathcal{G}'/\mathbb{Q}$  to  $\mathcal{G}'/\mathbb{Z}$  is "optimal".

Now we extend  $\mathrm{Sl}_2/\mathbb{Z}$  to  $\mathcal{G}/\mathbb{Z}$ , for any prime  $\ell \neq p$  we choose the obvious extension  $\mathrm{Sl}_2/\mathbb{Z}_\ell$ . For  $p$  we choose an Iwahori -Bruhat-Tits group scheme  $\mathcal{G}/\mathbb{Z}_p$ , it is smooth and flat and

$$\mathcal{G}(\mathbb{Z}_p) = \left\{ \begin{pmatrix} a & b \\ pc & d \end{pmatrix} \mid a, b, c, d \in \mathbb{Z}_p, ad - pbc = 1 \right\}.$$

The reduction  $\pmod{p}$  gives  $\mathcal{G} \times_{\mathbb{Z}} \mathbb{F}_p = \mathbb{G}_m \ltimes (\mathbb{G}_a \times \mathbb{G}_a)$ . It is clear that  $\mathcal{G} \times \mathbb{Z}_p$  and  $\mathcal{G}' \times \mathbb{Z}_p$  are locally isomorphic in the etale topology.

Hence we see that we can choose for our set  $\Sigma = \{p\}$  and we get

$$\frac{\#\mathcal{G}(\mathbb{F}_p)}{\#\mathcal{G}'(\mathbb{F}_p)} = \frac{p-1}{p+1}$$

We still have to discuss the factor  $p^{\delta_p(G, G')}$  and the contribution from the infinite place. We must go back to the definition of the Tamagawa measure. Since  $\mathcal{G}/\mathbb{Z}$  and  $\mathcal{G}'/\mathbb{Z}$  are smooth the Lie algebras of these group schemes are free  $\mathbb{Z}$ -modules, they are given by

$$\mathrm{Lie}(\mathcal{G}) = \mathbb{Z}H \oplus \mathbb{Z}pE_+ \oplus \mathbb{Z}E_-; \quad \mathrm{Lie}(\mathcal{G}') = \mathbb{Z} \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \oplus \mathbb{Z} \begin{pmatrix} 0 & p \\ -1 & 0 \end{pmatrix} \oplus \mathbb{Z} \begin{pmatrix} 0 & pi \\ i & 0 \end{pmatrix}. \quad (6.85)$$

To define the Tamagawa measure we have to choose top degree non zero invariant differential forms, in this situation we gauge them by requiring

$$\omega_G(H \wedge pE_+ \wedge E_-) = 1; \quad \omega_{G'}\left(\begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \wedge \begin{pmatrix} 0 & p \\ -1 & 0 \end{pmatrix} \wedge \begin{pmatrix} 0 & pi \\ i & 0 \end{pmatrix}\right) = 1.$$

For any prime  $\ell$  these linear forms provide invariant measures  $\omega_{G,\ell}, \omega_{G',\ell}$  on  $G(\mathbb{Q}_\ell), G'(\mathbb{Q}_\ell)$  and  $\omega_{G,\infty}, \omega_{G',\infty}$  at the infinite place, the product of these local measures gives the Tamagawa -measures  $\omega_G^{\text{Tam}}, \omega_{G'}^{\text{Tam}}$ . We recall that these two measures do not depend on the choice of  $\omega_G, \omega_{G'}$ , but the local factors do. In a sense the choice of these two forms is optimal with respect to the " arithmetic aspects".

With this product realisation of the Tamagawa measure we get for the local factor at  $p$  in formula (See 6.60)

$$\frac{\zeta_p(2)}{\text{vol}_{\omega_G^{\text{Tam},f}}(K_p)} = \frac{1}{p}(p+1); \quad \frac{\zeta_p(2)}{\text{vol}_{\omega_{G'}^{\text{Tam},f}}(K'_p)} = \frac{1}{p}(p-1)$$

We have to consider the contributions at the infinite place. We have to compare the measures  $\omega_{G,\infty}, \omega_{G',\infty}$  to the measures defined by the Killing form, and we have to compute the factor  $\kappa_\infty(G)$ .

We consider the cases  $G = \text{Sl}_2/\mathbb{Q}$  first. Then we have the decomposition

$$\mathfrak{g} = \mathbb{Q}Y \oplus \mathbb{Q}H \oplus \mathbb{Q}V = \mathfrak{k} \oplus \mathfrak{p}$$

this is an orthonormal decomposition for the Killingform  $B$  and  $B(Y, Y) = -8, B(H, H) = 8, B(V, V) = 8$ . and now we replace  $B$  by  $\frac{1}{8}B$ . This normalised Killing form defines a top differential  $\omega_B$  which satisfies  $\omega_B(Y, H, V) = 1$  this differential is of the form  $\omega_B^{K_\infty} \wedge \omega_{|\mathfrak{p}B}$ , here  $\omega_{|\mathfrak{p}B}$  is a 2-form on  $\mathfrak{p}$ , it is normalised by  $\omega_{|\mathfrak{p}B}(H, V) = 1$ .

We have to compare  $\omega_B$  and  $\omega_G$ . We must express  $Y, H, V$  as linear combination of  $H, pE_+, E_-$  and then we get easily

$$\omega_B = \pm \frac{p}{2} \omega_G$$

Now it is well known that in this case  $\omega^{GB} = -\frac{1}{2\pi} \omega_{|\mathfrak{p},B}$  ( i.e.  $\kappa_\infty(G) = -\frac{1}{2\pi} \cdot$ ) (See ???) and hence finally

$$\omega^{GB} = \pm \frac{p}{4\pi} \omega_G \tag{6.86}$$

Now we represent the Tamagawa number by  $\omega_G$ . We get

$$1 = \int_{G(\mathbb{Q}) \backslash G(\mathbb{A})} \omega^{\text{Tam}} = \int_{G(\mathbb{Z}) \backslash G(\mathbb{R})} \omega_{B,\infty} \times \int_{K_f} \omega_{B,f} = \frac{p}{p+1} \zeta(2)^{-1} \times \int_{G(\mathbb{Z}) \backslash G(\mathbb{R})} \omega_{G,\infty} \tag{6.87}$$

For the last factor we have

$$\int_{G(\mathbb{Z}) \backslash G(\mathbb{R})} \omega_{G,\infty} = \pm \frac{2}{p} \int_{G(\mathbb{Z}) \backslash G(\mathbb{R})} \omega_{B,\infty} \tag{6.88}$$

Now  $\int_{K_\infty} \omega_B^{K_\infty} = 2\pi$  and hence

$$\int_{\mathcal{G}(\mathbb{Z}) \backslash \mathbb{G}(\mathbb{R})} \omega_{B,\infty} = \pm \frac{4\pi}{p} \int_{\mathcal{G}(\mathbb{Z}) \backslash X} \omega_{|\mathfrak{p},B} = \pm \frac{8\pi^2}{p} \int_{\mathcal{G}(\mathbb{Z}) \backslash X} \omega^{GB} \quad (6.89)$$

and this finally results in

$$\chi_{\text{orb}}(\mathcal{G}(\mathbb{Z}) \backslash X) = \pm \int_{\mathcal{G}(\mathbb{Z}) \backslash X} \omega^{GB} = -\frac{p+1}{8} \frac{\zeta(2)}{\pi^2} = -\frac{p+1}{48} \quad (6.90)$$

Here we recollect that  $\mathcal{G}(\mathbb{Z}) = \Gamma_0(p) := \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{Sl}_2(\mathbb{Z}) \mid c \equiv 0 \pmod{p} \right\}$  and  $X = \mathbb{H}$  the upper half plane.

We do the same calculation for  $\mathcal{G}'$ . The group

$$G'(\mathbb{R}) = \left\{ \begin{pmatrix} x & py \\ -\sigma(y) & \sigma(x) \end{pmatrix} \mid x\sigma(x) + py\sigma(y) = 1; x, y \in \mathbb{C} \right\}$$

and we get the well known isomorphism

$$\Phi: G'(\mathbb{R}) \xrightarrow{\sim} S^3 \subset \mathbb{C}^2; \Phi: \begin{pmatrix} x & py \\ -\sigma(y) & \sigma(x) \end{pmatrix} \mapsto (x, \sqrt{p}y) \quad (6.91)$$

The identity element  $e_{G'} \in G'(\mathbb{R})$  is mapped to  $(1, 0) \in \mathbb{C}^2$  (" = "(1, 0, 0, 0)  $\in \mathbb{R}^4$ ). The tangent space to the sphere at this point is  $i\mathbb{R} \oplus \mathbb{C}$ . The Lie algebra  $\mathfrak{g}' \otimes \mathbb{R}$  is the tangent space of  $G'(\mathbb{R})$  at  $e_{G'}$  and the derivative  $D_\Phi$  maps

$$\begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \mapsto (i, 0) \text{ " = "(0, 1, 0, 0); } \begin{pmatrix} 0 & p \\ -1 & 0 \end{pmatrix} \mapsto (0, \sqrt{p}); \begin{pmatrix} 0 & pi \\ i & 0 \end{pmatrix} \mapsto (0, \sqrt{p}i),$$

we see that the images of our three basis vectors are orthogonal to each other. Hence the euclidian volume form evaluated at the triple of these three vectors gives

$$\omega^{\text{eucl}}((i, 0), (0, \sqrt{p}), (0, \sqrt{p}i)) = p$$

The volume of the 3-sphere with respect to the euclidian volume form is  $4\pi^2$ . Hence we get  $\text{vol}_{\omega_{G',\infty}}(K'_\infty) = \frac{4\pi^2}{p}$ . If we perform the same computation as in the first case we end up with

$$\chi_{\text{orb}}(\mathcal{S}'_{K'_f}, \tilde{\mathcal{M}}') = \frac{p-1}{4\pi^2} \zeta(2) = \frac{p-1}{48} \quad (6.92)$$

### 6.3.9 Some (philosophical) remarks

Of course the Gauss-Bonnet theorem only gives an alternating sum of dimensions of cohomology groups, we do not have any control of the possible cancellation. This is especially bad in the case when we have  $\kappa_\infty(G) = 0$ . We have seen that in the case where  $G(\mathbb{R})$  is compact the space  $\mathcal{S}_{K_f}^G$  is of dimension zero and hence all the cohomology sits in degree zero and there is no cancellation. In this case the differential geometric subtleties also disappear, i.e. we have  $\kappa_\infty(G) = 1$ .

We still may ask: How do the cohomology groups behave once we vary the level  $K_f$  or the coefficient system  $\mathcal{M}$ . What happens if  $K_f$  gets smaller and smaller? If our coefficient system is a highest weight module  $\mathcal{M}_\lambda$  where  $\lambda = \sum n_i \gamma_i$ , what happens if all the  $n_i \rightarrow \infty$ ?

Let us fix a neat reference level  $K_f^{(0)}$ , and the reference coefficient system  $\mathbb{Q}$ . Then we have seen that for  $K_f \subset K_f^{(0)}$  and any  $\mathcal{M}_\lambda$  we have

$$\chi(\mathcal{S}_{K_f}^G, \mathcal{M}) = [K_f^{(0)} : K_f] \times \chi(\mathcal{S}_{K_f^{(0)}}^G, \mathbb{Q}) \times \dim(\mathcal{M})$$

We consider the case  $\kappa_\infty(G) \neq 0$ . In this case the dimension  $d = \dim X$  is even. In this one might expect that in a certain sense

$$\chi(\mathcal{S}_{K_f}^G, \mathcal{M}) \simeq (-1)^{\frac{d}{2}} \dim H^{\frac{d}{2}}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}). \tag{6.93}$$

This expectation can be verified and also made precise in a few cases and it is also supported by experimental data.

If the highest weight  $\lambda$  is regular, i.e. if  $n_i > 0 \forall i$  then it has been shown by J. S. Li -J. Schwermer in [?] and L. Saper in [?] that  $H^\nu(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) = 0$  for  $\nu < \frac{d}{2}$ . Moreover it can be shown that all the cohomology  $H^\nu(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$  with  $\nu > \frac{d}{2}$  is Eisenstein cohomology (see section 9.2), and this gives us some confirmation of the statement above.

We drop the assumption on  $\lambda$  and vary  $K_f$ , then we have results by Lück and ...that

$$\lim_{K_f \rightarrow \epsilon} \frac{1}{[K_f^{(0)} : K_f]} H^\nu(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}) = \begin{cases} 0 & \text{if } \nu \neq \frac{d}{2} \\ \chi_{orb}(\mathcal{S}_{K_f}^G, \mathcal{M}) & \text{if } \nu = \frac{d}{2} \end{cases} \tag{6.94}$$

This is another piece of evidence for the above principle.

This last formula makes also sense if  $\kappa_\infty(G) = 0$ .

### The topological trace formula

The Gauss-Bonnet formula is a special case of the topological trace formula (See [2],[19],[34]). The topological trace formula is a tool to compute the trace

$$\text{tr}(T_h | H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})) = \sum_\nu (-1)^\nu \text{tr}(T_h | H^\nu(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})) \tag{6.95}$$

of a Hecke operator  $T_h$  (See section 6.3). If we choose for  $h$  the characteristic function of  $K_f$  then this trace is equal to  $\chi_{orb}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})$ . The topological trace formula gives a formula for the traces of Hecke operators on the cohomology in terms of orbifold Euler characteristics of fixed point sets. These fixed point sets are again locally symmetric spaces and hence again can be computed using the Gauss-Bonnet theorem.

We come back to our two groups  $G/\mathbb{Q}, G'/\mathbb{Q}$  and we assume that we have chosen compatible extensions  $\mathcal{G}/\mathbb{Z}$  and  $\mathcal{G}'/\mathbb{Z}$  as above. Again  $\Sigma$  will be the set of places where  $\mathcal{G}, \mathcal{G}'$  are not semi simple. Then we can identify the central

sub algebras  $\mathcal{H}^{(\Sigma)} = \bigotimes_{p \notin \Sigma} \mathcal{H}_p = \mathcal{H}^{(\prime, \Sigma)} = \bigotimes_{p \notin \Sigma} \mathcal{H}'_p$ . This means that we can compare the  $\mathcal{H}^{(\Sigma)}$  - modules  $H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})$  and  $H^\bullet(\mathcal{S}_{K_f}^{G'}, \mathcal{M}')$ .

To get such a comparison we can invoke the topological trace formula, We have to make some clever choices of Hecke operators  $h = h_{(\Sigma)} \times h^{(\Sigma)} \in \prod_{p \in \Sigma} \mathcal{H}_p \times \mathcal{H}^{(\Sigma)}$  and  $h' = h'_{(\Sigma)} \times h^{(\Sigma)} \in \prod_{p \in \Sigma} \mathcal{H}'_p \times \mathcal{H}^{(\Sigma)}$ .

Now we compute the traces on the two cohomology groups using the topological trace formula

$$\begin{aligned} \text{tr}(T_h | H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})) &= \sum_{\underline{x}: \text{fixed point}} T_h \underline{x} \\ \text{tr}(T_{h'} | H^\bullet(\mathcal{S}_{K_f}^{G'}, \mathcal{M}')) &= \sum_{\underline{x}': \text{fixed point}} T_{h'} \underline{x}' \end{aligned} \tag{6.96}$$

where the numbers  $T_h \underline{x}, T_{h'} \underline{x}'$  are local contributions, they are Euler characteristics of fixed point sets times so called orbital integrals.

Now we try to establish a correspondence between the two sets of fixed points such that for two corresponding points  $\underline{x} \leftrightarrow \underline{x}'$  we have adapted our Hecke operators such that  $T_h \underline{x} = T_{h'} \underline{x}'$ . In case we do not find a corresponding point  $\underline{x}'$  to a given  $\underline{x}$  we must have  $T_h \underline{x} = 0$ . If we are lucky-and in fact we are in a few cases- we can show that the right hand sides in ( 6.96) are equal.

This comparison between the cohomology groups attached to two different groups has been executed in some detail in [34] for a pair  $G = \text{Gl}_2/\mathbb{Q}$  and the multiplicative group  $G'/\mathbb{Q}$  of a division algebra. We also discuss the much more subtle of comparing  $\text{Sl}_2/\mathbb{Q}$  and where  $G'/\mathbb{Q}$  is the norm one group of a division algebra. (See also [52],)

### Arthur-Selberg trace formula vs. topological trace formula

In Chapter 8 we will discuss the description of the cohomology  $H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathbb{C}})$  in terms of automorphic forms. In the theory of automorphic forms we also can compare the spaces of automorphic forms for a pair of groups which are inner forms of each other. This occurs the first time in the fundamental book of Jacquet and Langlands [43] for the two groups  $\text{Gl}_2$  and the multiplicative group of a division algebra. The result is the Jacquet- Langlands correspondence, which plays a predominant role in the theory of modular forms, The Jacquet -Langlands correspondence implies the above results on cohomology.

The main tool to prove the Jacquet-Langlands correspondence is the Arthur-Selberg trace formula, J. Arthur and many other people have developed this instrument to the case of general reductive groups. As an application they get results which allow a comparison of spaces of automorphic forms (See Arthur's papers) on different groups. The formulation and the proof of the Arthur-Selberg trace formula are peppered with enormous analytical difficulties, which make it difficult to apply it. The problem is the non compactness of  $\mathcal{S}_{K_f}^G$ , one encounters situations in which certain infinite sums or certain integrals are divergent and one has to renormalise them.

These subtle analytical problems disappear if we use the topological trace formula instead. In this context we encounter the problem how to treat the "fixed points at infinity", this is discussed and solved in [2] in the rank one case

and in [19] in greater generality. It should be possible to prove many of the relevant consequences of the Arthur-Selberg trace formula by using the topological trace formula, provided we restrict our attention to the "cohomological part" of the space of automorphic forms. This applies especially to the comparison of the cohomology of to different groups and to questions of endoscopy. The proofs would dramatically simplify. On the other hand the problems with the stabilisation of the trace formula remain the same.

In section 3.2.1 we gave a general strategy how to write an algorithm to compute -at least in principle- cohomology groups  $H^\bullet(\mathcal{S}_{K_f}^G, \mathcal{M})$  and in addition to compute the action of a Hecke operator  $T_{p, \chi}^{\text{coh}, \lambda}$  (??) on it. Together with H. Gangl we wrote such an algorithm in a baby case (3.3), and we used this to verify an assertion about denominators of Eisenstein classes experimentally.

We come back to this algorithm and discuss it in the case that  $G(\mathbb{R})$  is compact. In this case  $\mathcal{S}_{K_f}^G = * \times G(\mathbb{A}_f)/K_f$  is a finite set, let  $\mathcal{A}(* \times G(\mathbb{A}_f)/K_f, \mathbb{Z})$  be the module of  $\mathbb{Z}$ - valued functions on this finite set, it is of course equal to  $H^0(\mathcal{S}_{K_f}^G, \mathbb{Z})$ . The space  $\mathcal{A}(* \times G(\mathbb{A}_f)/K_f, \mathbb{C})$  is also called the space of algebraic modular forms. We have a basis given by the delta functions  $\delta_{\underline{x}}$ , where  $\underline{x}$  runs through the points in  $\mathcal{A}(* \times G(\mathbb{A}_f)/K_f, \mathbb{C})$ .

We recall the definition of Hecke operators in this special situation. We pick an element  $x_p \in G(\mathbb{Q}_p)$  we extend it to an adelic point  $\underline{x}_p = (1, 1, \dots, x_p, \dots 1, \dots)$  We consider the group  $K(\underline{x}_p)_f := K_f \cap \underline{x}_p K_f \underline{x}_p^{-1}$  and the projection map  $\pi_+ : G(\mathbb{Q}) \setminus * \times G(\mathbb{A}_f)/K(\underline{x}_p)_f \rightarrow G(\mathbb{Q}) \setminus * \times G(\mathbb{A}_f)/K_f$ . If we multiply by  $\underline{x}_p$  from the right we get a map

$$m(\underline{x}_p) : G(\mathbb{Q}) \setminus * \times G(\mathbb{A}_f)/K(\underline{x}_p)_f \rightarrow G(\mathbb{Q}) \setminus * \times G(\mathbb{A}_f)/K(\underline{x}_p^{-1})_f$$

$$m(\underline{x}_p) : \underline{y} \mapsto \underline{y} \underline{x}_p$$

Now the Hecke operator  $T(\underline{x}_p) : H^0(\mathcal{S}_{K_f}^G, \mathbb{Z}) \rightarrow H^0(\mathcal{S}_{K_f}^G, \mathbb{Z})$  does the following: We choose a set  $\underline{u}_1, \dots, \underline{u}_t$  for  $K_f/K(\underline{x}_p)_f$  We pick an  $\underline{x} \in G(\mathbb{Q}) \setminus * \times G(\mathbb{A}_f)/K_f$  and represent by  $\tilde{\underline{x}} \in G(\mathbb{A}_f)$ . We consider the fiber  $\pi_+^{-1}(\underline{x})$  the points in the fiber are represented by  $\underline{x} \underline{u}_i, i = 1 \dots t$ . A point  $\underline{y} \in \pi_+^{-1}(\underline{x})$  comes with a multiplicity  $m(\underline{y})$ , the number of times it is represented by a  $\tilde{\underline{x}} \underline{u}_i$ . The map  $m(\underline{x}_p)$  maps the points  $\tilde{\underline{x}} \underline{u}_i$  to  $G(\mathbb{Q}) \setminus * \times G(\mathbb{A}_f)/K(\underline{x}_p^{-1})_f$ . To this set of points (with multiplicities) we apply the projection  $\pi_- : G(\mathbb{Q}) \setminus * \times G(\mathbb{A}_f)/K(\underline{x}_p^{-1})_f \rightarrow G(\mathbb{Q}) \setminus * \times G(\mathbb{A}_f)/K_f$ . We get a finite set of points  $T(\underline{x}, \underline{x}_p) \subset G(\mathbb{Q}) \setminus * \times G(\mathbb{A}_f)/K_f$ , where each point  $\underline{z} \in T(\underline{x}, \underline{x}_p)$  comes with a finite multiplicity  $a_{\underline{x}, \underline{z}}(\underline{x}_p)$ , this is the number of times it is hit by a  $\pi_-(m(\underline{x}_p)(\tilde{\underline{x}} \underline{u}_i))$ . Hence  $a_{\underline{x}, \underline{z}}(\underline{x}_p)$  is an integer  $> 0$  for  $\underline{z} \in T(\underline{x}, \underline{x}_p)$ , we put  $a_{\underline{x}, \underline{z}}(\underline{x}_p) = 0$  for  $\underline{z} \notin T(\underline{x}, \underline{x}_p)$ . Then

$$T(\underline{x}_p)(\delta_{\underline{x}}) = \sum_{\underline{z} \in G(\mathbb{Q}) \setminus * \times G(\mathbb{A}_f)/K_f} a_{\underline{x}, \underline{z}}(\underline{x}_p) \delta_{\underline{z}} \tag{6.97}$$

The computation of this incidence matrix  $(a_{\underline{x}, \underline{z}}(\underline{x}_p))$  may become very difficult. In a slightly different context such computations are carried out in [59] and their results are presented in [10]. Even in the case of the group  $\text{Spin}(L_8 \oplus L_8)$

where we have only two elements in  $\mathcal{S}_{K_f}^G$  the computation of the incidence matrix is by no means trivial. In [10] the authors define a Hecke operator  $T_2$  using "Kneser Neighbors", this is essentially a  $T(x_2)$  as described above. And they give the resulting matrix

$$T_2 = \begin{pmatrix} 20025 & 18225 \\ 12870 & 14670 \end{pmatrix} \quad (6.98)$$

(One of the reasons we give this matrix here is the following observation: The difference of the two eigenvalues is divisible by 691 !) In [10] the authors also discuss  $T_2$  for the lattice  $L_8 \oplus L_8 \oplus L_8$  but they do not write the resulting  $(24 \times 24)$ -matrix.

On the other hand there are formulas in [10] which can not be obtained simple from a computer program, for instance **Theorem A** in section 1.2. on Kneser neighbors.

Of course we may also do this if we have a non trivial coefficient system  $\mathcal{M}$ . In case we need to know the incidence matrix but in addition we also have to keep track of the linear maps between the stalks of sheaves. Then even the case of the 8 dimensional lattice  $L_8$  becomes non trivial.

It seems to be an interesting exercise to consider the the two groups  $\mathcal{G}'/\mathbb{Z} = \text{Spin}(L_8)/\mathbb{Z}$  and  $\mathcal{G}/\mathbb{Z} =$  the split Chevalley group of type  $D_4$ . Now we look at the unramified cohomology, i.e. we take  $K_f = \mathcal{G}(\hat{\mathbb{Z}}), K'_f = \mathcal{G}'(\hat{\mathbb{Z}})$  and compare the Hecke modules

$$H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}) \text{ and } H^0(\mathcal{S}_{K'_f}^G, \tilde{\mathcal{M}}') \quad (6.99)$$

The result should be compared to the results of Arthur.

This is perhaps the right moment, to discuss another minor technical point. When we discuss the action of the Hecke algebra  $\mathcal{H}_{K_f} = C_c(G(\mathbb{A}_f)/K_f, \mathbb{Q})$  on  $H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})$  then we chose the same  $K_f$  for the space and for the Hecke algebra. We also normalized the measure on the group so that it gave volume 1 to  $K_f$ . But we have of course an inclusion of Hecke algebras  $\mathcal{H}_{K_f} \subset \mathcal{H}_{K'_f}$ . Therefore  $\mathcal{H}_{K_f}$  also acts on  $H^\bullet(\mathcal{S}_{K'_f}^G, \tilde{\mathcal{M}})$ . This contains  $H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})$  but then the inclusion is not compatible with the action of the Hecke algebra. We therefore choose a measure independently of the level, if we are in a situation where we vary the level. In such a case a measure provided by an invariant form  $\omega_G$  on  $G$  (See 2.1.3) is a good choice. If we now define the action of the Hecke operators by means of this measure. With this choice of a measure the inclusion  $\mathcal{H}_{K_f} \subset \mathcal{H}_{K'_f}$  is compatible with the inclusion of the cohomology groups.

Then we see the new Hecke operator  $T_h^{(\omega_G)}$ , and the old one are related by the formula

$$T_h = \frac{1}{\text{vol}_{|\omega_G|}(K_f)} T_h^{(\omega_G)}$$

The reader might raise the question, why we work with fixed levels and why we do not pass to the limit. The reason is that for some questions we need to work with the integral cohomology, and this does not behave so well under change of level.

### 6.3.10 Some questions and and some general facts

#### Homology

We may also define homology groups  $H_i(\mathcal{S}_{K_f}^G, \underline{\mathcal{M}}_\lambda)$  and  $H_i(\mathcal{S}_{K_f}^G, \partial\mathcal{S}_{K_f}^G, \underline{\mathcal{M}}_\lambda)$ , here  $\mathcal{M}_\lambda$  is a ‘‘cosheaf’’. The ‘‘costalk’’  $\underline{\mathcal{M}}_{\mathbb{Z},x}$  is obtained as follows: We consider  $\pi^{-1}(x)$  and

$$\bigoplus_{\underline{y}=y \times \underline{g}_f K_f / K_f} \underline{g}_f \mathcal{M}_\lambda,$$

and the action of  $G(\mathbb{Q})$  on this direct sum. Then  $\underline{\mathcal{M}}_{\lambda,x}$  is the module of coinvariants. If we pick a point  $\underline{y} = y \times \underline{g}_f K_f / K_f$ , which maps to  $x \in \mathcal{S}_{K_f}^G$  then we get an isomorphism

$$\underline{\mathcal{M}}_{\lambda,x} \simeq (g_f \mathcal{M}_\lambda)_{\Gamma_y^{(g_f)}}.$$

We define the chain complex

$$C_i(\mathcal{S}_{K_f}^G, \underline{\mathcal{M}}_\lambda)$$

and the above homology groups are given by the homology of this complex.

If we assume that  $\mathcal{S}_{K_f}^G$  is oriented (ref. to prop 1.3) then we know (Chap. II 2. 1. 5) that we have isomorphisms which are compatible with the fundamental exact sequence

$$\begin{array}{ccc} \downarrow & & \downarrow \\ H^{i-1}(\partial\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) & \xrightarrow{\sim} & H_{d-i}(\partial\mathcal{S}_{K_f}^G, \underline{\mathcal{M}}_\lambda) \\ \downarrow & & \downarrow \\ H_c^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) & \xrightarrow{\sim} & H_{d-i}(\mathcal{S}_{K_f}^G, \underline{\mathcal{M}}_\lambda) \\ \downarrow & & \downarrow \\ H^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) & \xrightarrow{\sim} & H_{d-i}(\mathcal{S}_{K_f}^G, \partial\mathcal{S}_{K_f}^G, \underline{\mathcal{M}}_\lambda) \\ \downarrow & & \downarrow \\ H^i(\partial\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) & \xrightarrow{\sim} & H_{d-i-1}(\partial\mathcal{S}_{K_f}^G, \underline{\mathcal{M}}_\lambda) \\ \downarrow & & \downarrow \end{array}$$

poincdu

### 6.3.11 Poincaré duality

We assume that  $\mathcal{S}_{K_f}^G$  is connected. If we denote the dual representation by  $\mathcal{M}_\lambda^\vee = \mathcal{M}_{w_0(\lambda)}$  ( we choose the right lattice  $\mathcal{M}_\mathbb{Z}^\vee \subset \mathcal{M}_\mathbb{Q}^\vee$ ) we have the canonical homomorphism  $\Phi_\lambda : \mathcal{M}_\lambda \otimes \mathcal{M}_\lambda^\vee \rightarrow \mathbb{Z}$  and the standard pairing between the homology and the cohomology groups yields pairings

$$\begin{array}{ccccc} H_c^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \times H_i(\mathcal{S}_{K_f}^G, \partial\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda^\vee) & \rightarrow & H^0(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \tilde{\mathcal{M}}_\lambda^\vee) & \rightarrow & H^0(\mathcal{S}_{K_f}^G, \mathbb{Z}) \\ \downarrow & & \downarrow & & \downarrow \\ H^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \times H_i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda^\vee) & \rightarrow & H^0(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \tilde{\mathcal{M}}_\lambda^\vee) & \rightarrow & H^0(\mathcal{S}_{K_f}^G, \mathbb{Z}) \end{array}$$

This pairing is of course compatible with the isomorphism between homology and cohomology and then the pairing becomes the cup product. We get the diagram

$$\begin{array}{ccccccc}
H_c^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \times H^{d-i}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda^\vee) & \rightarrow & H_c^d(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \tilde{\mathcal{M}}_\lambda^\vee) & \rightarrow & H_c^d(\mathcal{S}_{K_f}^G, \mathbb{Z}) \\
\downarrow & & \downarrow & & \downarrow \\
H^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \times H_c^{d-i}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda^\vee) & \rightarrow & H_c^d(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \tilde{\mathcal{M}}_\lambda^\vee) & \rightarrow & H_c^d(\mathcal{S}_{K_f}^G, \mathbb{Z}) \\
\uparrow & & \uparrow & & \uparrow
\end{array}$$

We know that the manifold with corners  $\partial\mathcal{S}_{K_f}^G$  "smoothable" it can be approximated by a  $\mathcal{C}$ -manifold and therefore we also have a pairing  $\langle \cdot, \cdot \rangle_\partial$  on the cohomology of the boundary. This pairing is consistent with the fundamental long exact sequence (Thm. 6.2.1). We write this sequence twice but the second time in the opposite direction and the pairing  $\langle \cdot, \cdot \rangle$  in vertical direction:

$$\begin{array}{ccccc}
\rightarrow & H^p(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) & \xrightarrow{r} & H^p(\partial\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) & \xrightarrow{\delta} \\
& \times & & \times & \\
\leftarrow & H_c^{d-p}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda^\vee) & \xleftarrow{\delta} & H^{d-p-1}(\partial\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda^\vee) & \leftarrow \\
& \downarrow \langle \cdot, \cdot \rangle & & \downarrow \langle \cdot, \cdot \rangle_\partial & \\
& H_c^d(\mathcal{S}_{K_f}^G, \mathbb{Z}) & \xleftarrow{\delta_d} & H_c^{d-1}(\partial\mathcal{S}_{K_f}^G, \mathbb{Z}) & 
\end{array} \quad (6.100)$$

then we have: For classes  $\xi \in H^p(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda), \eta \in H^{d-p-1}(\partial\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda^\vee)$  we have the equality

$$\langle \xi, \delta(\eta) \rangle = \delta_d(\langle r(\xi), \eta \rangle_\partial) \quad (6.101)$$

### Non degeneration of the pairing

The spaces  $\mathcal{S}_{K_f}^G$  and  $\partial\mathcal{S}_{K_f}^G$  are not connected in general. Let us assume that we have a consistent orientation on  $\mathcal{S}_{K_f}^G$ . Then each connected component  $M$  of  $\mathcal{S}_{K_f}^G$  is an oriented manifold which is natural embedded into its compactification  $\tilde{M}$ . It is obvious that the cohomology groups are the direct sums of the cohomology groups of the connected components and that we may restrict the pairing to the components

$$H^p(M, \tilde{\mathcal{M}}_\lambda) \times H_c^{d-p}(M, \tilde{\mathcal{M}}_\lambda^\vee) \rightarrow H_c^d(M, \mathbb{Z}) = \mathbb{Z}. \quad (6.102)$$

We recall the results which are explained in Vol. I 4.8.4. The fundamental group  $\pi_1(M)$  is an arithmetic subgroup  $\Gamma_M \subset G(\mathbb{Q})$  and  $\mathcal{M}_\lambda, \mathcal{M}_\lambda^\vee$  are  $\Gamma_M$  modules. For any commutative ring with identity  $\mathbb{Z} \rightarrow R$  the  $\Gamma_M$  modules  $\mathcal{M}_\lambda \otimes R, \mathcal{M}_\lambda^\vee \otimes R$  provide local systems  $\widetilde{\mathcal{M}_\lambda \otimes R}, \widetilde{\mathcal{M}_\lambda^\vee \otimes R}$ , and we have the extension of the cup product pairing

$$H^p(M, \widetilde{\mathcal{M}_\lambda \otimes R}) \times H_c^{d-p}(M, \widetilde{\mathcal{M}_\lambda^\vee \otimes R}) \rightarrow H_c^d(M, R) = R$$

**Proposition 6.3.7.** *If  $R = k$  is a field then the pairing is non degenerate. .*

*If  $R$  is a Dedekind ring then the pairing then the cohomology may contain some torsion submodules and*

$$H^p(M, \widetilde{\mathcal{M}_\lambda \otimes R})/\text{Tors} \times H_c^{d-p}(M, \widetilde{\mathcal{M}_\lambda^\vee \otimes R})/\text{Tors} \rightarrow H_c^d(M, R) = R$$

*is non degenerate.*

(See Vol. I 4.8.9)

We want to discuss the consequences of this result for the cohomology of  $H_c^*(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$ . Before we do this we want to recall some simple facts concerning the representations of the algebraic group  $G/\mathbb{Q}$ . We consider two highest weights  $\lambda, \lambda_1 \in X^*(T \times F)$  which are dual modulo the center. By this we mean that we have (See 6.22)

$$\lambda = \lambda^{(1)} + \delta, \lambda_1 = -w_0(\lambda^{(1)}) + \delta_1 \tag{6.103}$$

Then  $\delta + \delta_1$  is a character on  $X^*(C' \times F)$  and yields a one dimensional module

(?????)  $H_c^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \times H^{d-i}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda^\vee) \rightarrow H_c^d(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \tilde{\mathcal{M}}_\lambda^\vee) \rightarrow H_c^d(\mathcal{S}_{K_f}^G, \mathbb{Z})$   
 for  $G \times F$ , of course the action of  $G^{(1)}$  on this module is trivial. Then we get a  $G$  invariant non trivial pairing

$$\mathcal{M}_{\lambda, F} \times \mathcal{M}_{\lambda_1, F} \rightarrow \mathcal{N}_{\lambda \circ \lambda_1}$$

which induces a pairing

$$H_c^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda, F}) \times H^{d-i}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda_1, F}) \rightarrow H_c^d(\mathcal{S}_{K_f}^G, \mathcal{N}_{\lambda \circ \lambda_1}),$$

this only a slight generalization of the previous pairing.

Now we recall that (under certain assumptions) we have the inclusion  $\pi_0(\mathcal{S}_{K_f}^G) \hookrightarrow \pi_0(\mathcal{S}_{K_\infty^{C'} \times K_f^{C'}})$  and then we get

$$H_c^d(\mathcal{S}_{K_f}^G, \mathcal{N}_{\lambda \circ \lambda_1}) \subset H^0(\mathcal{S}_{K_\infty^{C'} \times K_f^{C'}}^G, \mathcal{N}_{\lambda \circ \lambda_1}) = \bigoplus_{\chi': \text{type}(\chi') = \lambda \circ \lambda_1} F\chi'$$

The character  $\chi'$  has a restriction to  $C(\mathbb{A})$  let us call this restriction  $\chi$ .

The group  $C(\mathbb{A}_f)$  acts on the cohomology groups and this action has an open kernel  $K_f^C$ . Hence we can decompose the cohomology groups on the left hand side according to characters

$$H_c^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda, F}) = \bigoplus_{\zeta_f: \text{type}(\zeta_f) = \delta} H_c^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda, F})(\zeta_f) \tag{6.104}$$

$$H^{d-i}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda_1, F}) = \bigoplus_{\zeta_{1, f}: \text{type}(\zeta_{1, f}) = \delta_1} H^{d-i}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda_1, F})(\zeta_{1, f}). \tag{6.105}$$

With these notations we get another formulation of Poincaré duality.

**Proposition 6.3.8.** *If we have three algebraic Hecke characters  $\zeta_f, \zeta_{1, f}, \chi'_f$  of the correct type and if we have the relation  $\zeta_f \cdot \zeta_{1, f} = \chi'_f$  then the cup product induces a non degenerate pairing*

$$H_c^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda, F})(\zeta_f) \times H^{d-i}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda_1, F})(\zeta_{1, f}) \rightarrow F\chi'_f$$

This is an obvious consequence of our considerations above. Fixing the central characters has the advantage that the target space of the pairing becomes one dimensional over  $F$ , The field  $F$  should contain the values of the characters.

We return to the diagram (6.100) and consider the images  $\text{Im}(r^q)(\zeta_f) = \text{Im}(H_c^q(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda, F})(\zeta_f) \rightarrow H_c^{d-q-1}(\partial\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda, F}^\vee)(\zeta_f)$  and  $\text{Im}(r^{\vee, d-q-1})$ . Then the following proposition is an obvious consequence of the non degeneration of the pairing and (6.101)

**Proposition 6.3.9.** *The images  $\text{Im}(r^p(\zeta_f))$  and  $\text{Im}(r^{\vee, d-p-1})(\zeta_{1,f})$  are mutual orthogonal complements of each other with respect to  $\langle, \rangle_{\partial}$ .*

*The pairing in proposition 6.3.8 induces a non degenerate pairing*

$$H_!^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda, F})(\zeta_f) \times H_!^{d-i}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda_1, F})(\zeta_{1,f}) \rightarrow F\chi'$$

*Proof.* Let  $\eta \in H^{d-p-1}(\zeta_{1,f})$ . Then we know from the exactness of the sequence that  $\eta \in \text{Im}(r^{\vee, d-p-1})(\zeta_{1,f}) \iff \delta(\eta) = 0 \iff \langle \delta(\eta), \xi \rangle = 0$  for all  $\xi \in H^p(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda})(\zeta_f) \iff \langle \eta, r(\xi) \rangle = 0$  for all  $\xi \in H^p(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda})(\zeta_f) \iff \langle \eta, \xi' \rangle_{\partial} = 0$  for all  $\xi' \in \text{Im}(r^q)(\zeta_f)$ .

The second assertion is rather obvious. If we have  $\xi \in H_!^p(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda})(\zeta_f)$ ,  $\xi_1 \in H_!^{d-p}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda^{\vee}})(\zeta_f)$  then we can lift either of these classes - say  $\xi_1$  - to a class  $\tilde{\xi}_1 \in H_c^p(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda})(\zeta_f)$  and then  $\langle \xi_1, \xi_2 \rangle = \langle \tilde{\xi}_1, \xi_2 \rangle$ . It is clear that the result does not depend on the choice of class which we lift. It is also obvious that the pairing is non degenerate.  $\square$

Of course we also have a version of proposition 6.3.9 for the integral cohomology. Since we fixed the level we have only a finite number of possible central characters  $\zeta_f, \zeta_{1,f}$  of the required type. The values of these characters evaluated on  $C(\mathbb{A}_f)$  lie in a finite extension  $F/\mathbb{Q}$  and of of course they are integral. If we now invert a few small primes and pass to a quotient ring  $R = \mathcal{O}_F[1/N]$  then we get the decomposition (6.104) but with coefficient systems which are  $R$ -modules:

$$H_c^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda, R}) = \bigoplus_{\zeta_f: \text{type}(\zeta_f) = \delta} H_c^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda, R})(\zeta_f) \quad (6.106)$$

$$H^{d-i}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda_1, R}) = \bigoplus_{\zeta_{1,f}: \text{type}(\zeta_{1,f}) = \delta_1} H^{d-i}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda_1, R})(\zeta_{1,f}) \quad (6.107)$$

Then it becomes clear that we get an integral version of proposition 6.3.8 where replace the  $F$ -vector space coefficient systems  $\tilde{\mathcal{M}}_{\lambda, F}$  by  $R$ -module coefficient systems. We get a pairing (See [28] 4.8.4)

$$H_c^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda, R})(\zeta_f)/\text{Tors} \times H^{d-i}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda_1, R})(\zeta_{1,f})/\text{Tors} \rightarrow R\chi' \quad (6.108)$$

and this pairing is non degenerate. (See [28] Thm. 4.8.9. The finiteness assumptions are easy consequences of reduction theory)

We recall the notion of non degenerate. Our ring  $R$  is a Dedekind ring and all our cohomology groups are finitely generated  $R$  modules. If we divide any finitely generated  $R$ -module by the subgroups of torsion elements then the result is a projective  $R$ -module and it is locally free for Zariski topology. An element  $\xi \in H_c^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda, R})(\zeta_f)/\text{Tors}$  is called *primitive* if the submodule  $R\xi$  is -locally for the Zariski topology- a direct summand or what amounts to the same if  $H_c^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda, R})(\zeta_f)/\text{Tors}/R\xi$  is torsion free. Then the assertion that the above pairing is non degenerate means:

*For any primitive element  $\eta \in H_c^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda, R})(\zeta_f)/\text{Tors}$  we find elements  $\xi_1, \xi_2, \dots, \xi_r \in H^{d-i}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda_1, R})(\zeta_{1,f})/\text{Tors}$  such that the ideal generated by  $\langle \xi_1, \eta \rangle, \langle \xi_2, \eta \rangle, \dots, \langle \xi_r, \eta \rangle$  is equal to  $R$ .*

We want to formulate an integral version of (6.101). Here the statement is not quite symmetric. It is clear from ??? that we get a pairing  $\boxed{\text{intint}}$

$$H_!^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda,R})(\zeta_f)_{\text{int}} \times H^{d-i}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda_1,R})(\zeta_{1,f})_{\text{int},!} \rightarrow R\chi'. \quad (6.109)$$

It is also clear from proposition (6.3.7)

**Proposition 6.3.10.** *This pairing is partially non degenerate. For any primitive element  $\eta \in H^{d-i}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda_1,R})(\zeta_{1,f})_{\text{int},!}$  we find elements*

$$\xi_1, \xi_2, \dots, \xi_r \in H_!^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda,R})(\zeta_f)_{\text{int}}$$

such that the ideal generated by  $\langle \eta, \xi_1 \rangle, \langle \eta, \xi_2 \rangle, \dots, \langle \eta, \xi_r \rangle$  is equal to  $R$ .

Here we see that the possibility that

$$H^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda_1,R})(\zeta_{1,f})_{\text{int},!} / H_!^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda,R})(\zeta_f)_{\text{int}} \neq (0)$$

plays a role.

### Inner Congruences

We choose a highest weight  $\lambda = \lambda^{(1)} + d\delta$  and the dual weight  $\lambda^\vee = -w_0(\lambda) - d\delta$ . Let us also fix a central character  $\zeta_f$  whose type is equal to the restriction of  $d\delta$  to the central torus  $C$ .

We look at the pairing in prop. 6.3.9 where we assume in addition that  $\zeta_{1,f} = \zeta_f^{-1}$  and we take the action of the Hecke algebra into account, i.e we look at the decomposition into eigenspaces (see(6.45)). Then we get a non degenerate pairing between isotypical subspaces

$$H_!^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda,F})(\pi_f) \times H_!^{d-i}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda^\vee,F})(\pi_f^\vee) \rightarrow F$$

where we assume that the central characters of the summands are  $\zeta_f, \zeta_f^{-1}$ .

If we try to extend this to the integral cohomology. In this case the above decomposition yields decompositions up to isogeny

$$\begin{aligned} H_!^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda,R})/\text{Tors} &\supset \bigoplus_{\pi_f} H_!^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda,R})/\text{Tors}(\pi_f) \\ H_!^{d-i}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda^\vee,R})/\text{Tors} &\supset \bigoplus_{\pi_f} H_!^{d-i}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda^\vee,R})/\text{Tors}(\pi_f^\vee) \end{aligned} \quad (6.110)$$

where we should fix the central characters as above. We choose a pair  $\pi_f, \pi_f^\vee$ . Then our non degenerate pairing from the above proposition induces a pairing

$$H_!^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda,R})/\text{Tors}(\pi_f) \times H_!^{d-i}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda^\vee,R})/\text{Tors}(\pi_f^\vee) \rightarrow R \quad (6.111)$$

and this pairing is non degenerate if and only if both modules are direct summands in the above decomposition up to isogeny.

But it may happen that the values of the pairing generate a proper ideal  $\Delta(\pi_f) \subset R$ , and in this case the above submodules will not be direct summands and this implies that we will have congruences between the Hecke-module  $\pi_f$  and some other module in the decomposition up to isogeny. This yields the *inner congruences*.

The ideal  $\Delta(\pi_f)$  should be expressed in terms of  $L$ -values, in the classical case this has been done by Hida [Hi].



# Chapter 7

## The fundamental question

Let  $\Sigma$  be a finite set. Of course any product  $V = \otimes H_{\pi_p}$  of finite dimensional absolutely irreducible modules for the  $\mathcal{H}_p$ , for which  $\mathcal{H}_p$  is spherical for all  $p \notin \Sigma$  gives us an absolutely irreducible module for the Hecke algebra.

*We may ask: Can we formulate non tautological conditions for the irreducible representation  $V$  or for the collection  $\{\pi_p\}_{p:\text{prime}}$ , which are necessary or (and) sufficient for the occurrence of  $\otimes'_p \pi_p$  in the cohomology*

This question can be formulated in the more general framework of the theory automorphic forms, but in this book we only consider "cohomological" (or certain limits of those) automorphic forms. This restricted question is difficult enough. A speculative answer is outlined in the following section

### 7.0.1 The Langlands philosophy

Let us start from a product  $V = \otimes H_{\pi_p}$ . For the primes outside the finite set  $\Sigma$  the module  $H_{\pi_p}$  is determined by its Satake parameter  $\omega_p$ .

#### The dual group

There is another way of looking at these Satake parameters  $\omega_p$ . We explain this in the case that  $\mathcal{G}/\mathbb{Z}_p$  is a split reductive group. We choose a maximal split torus  $\mathcal{T}$  over  $\mathbb{Z}$  and a Borel subgroup  $\mathcal{B}/\mathbb{Z}$ . For any commutative ring with identity ring  $R$  we have a canonical isomorphism  $X_*(\mathcal{T}) \otimes R^\times \xrightarrow{\sim} \mathcal{T}(R)$ , which is given by  $\chi \otimes a \mapsto \chi(a)$ . Then  $\mathcal{T}(\mathbb{Q}_p)/\mathcal{T}(\mathbb{Z}_p) = X_*(\mathcal{T}) \otimes \mathbb{Q}_p^\times/\mathbb{Z}_p^\times = X_*(\mathcal{T})$ . We apply this to the maximal split torus  $\mathcal{T}/\mathbb{Z}_p \subset \mathbb{G}/\mathbb{Z}_p$ . Then  $\Lambda(\mathcal{T}) = \text{Hom}(X_*(\mathcal{T}), \mathbb{C}) = X^*(\mathcal{T}) \otimes \mathbb{C}^\times = T^\vee(\mathbb{C})$  where  $T^\vee$  is the torus over  $\mathbb{Q}$  whose cocharacter module is  $X^*(\mathcal{T})$ . This torus over  $\mathbb{Q}$  is called the dual torus. There is a canonical construction of a dual group  ${}^L G/\mathbb{C}$ , this is a reductive group with maximal torus  $T^\vee$  such that the Weyl group of  $T^\vee$  in this dual group is equal to the Weyl group of  $\mathcal{T} \subset \mathbb{G}$  (See also (7.0.1)). This dual torus sits in a Borel subgroup  ${}^L B \subset {}^L G$ . Recall that we have a canonical pairing

$$\langle, \rangle: X_*(\mathcal{T}) \times X^*(\mathcal{T}) \rightarrow \mathbb{Z}, \quad \gamma \circ \chi(x) \mapsto x^{\langle \chi, \gamma \rangle}. \quad (7.1)$$

The positive simple roots in  $X^*(T^\vee)$  in the dual group  ${}^L G/\mathbb{C}$  are the cocharacters  $\alpha_i^\vee \in X_*(\mathcal{T}^{(1)})$  defined by

$$\langle \alpha_i^\vee, \gamma_j \rangle = \delta_{i,j}.$$

Hence we can interpret  $\omega_p \in \Lambda(T) = X^*(\mathcal{T}) \otimes \mathbb{C}^\times = T^\vee(\mathbb{C})$  as a semi simple conjugacy class in  ${}^L G(\mathbb{C})$ . Remember that  $\omega_p$  is only determined by the local component  $\pi_p$  up to an element in the Weyl group, hence we only get a conjugacy class.

We assume that  $\mathbb{G}/\mathbb{Z}$  is a split reductive group scheme. Then the dual group  ${}^L G$  is also split over  $\mathbb{Z}$  and the absolutely irreducible highest weight modules  $\mathcal{M}_\lambda$  for  $\mathbb{G}/\mathbb{Z}$  and the highest weight module attached to  $\chi$  are defined over  $\mathbb{Q}$ . Let  $\pi_f \in \text{Coh}_1(G, K_f, \lambda)$  be absolutely irreducible and defined over a finite extension  $E/\mathbb{Q}$ . Hence we see that our absolutely irreducible  $\pi_f$  provides a collection of conjugacy classes  $\{\omega(\pi_p) = \omega_p\}_{p \notin \Sigma}$  in the dual group  ${}^L G(E)$ .

A rather vague but also very bold formulation of the general Langlands philosophy predicts:

*The isotypical components under the action of the Hecke algebra, namely the  $H^i(\mathcal{S}_{K_f}^G, \mathcal{M})(\pi_f)$ , should correspond to a collection  $\{\mathbb{M}(\pi_f, r_\chi)\}_{r_\chi}$  of motives (with coefficients in  $E$ ). The correspondence should be defined via the equality of certain automorphic and motivic  $L$ -functions.*

This formulation is definitely somewhat cryptic, we will try to make it a little bit more precise in the following sections.

Such a motive could in principle be a "direct summand" the  $H^i(X)$  of a smooth projective scheme  $X/\mathbb{Q}$ , which in a certain sense is cut out by a projector. In some cases, where the space  $\mathcal{S}_{K_f}^G$  "is a Shimura variety", these motives have been constructed, we will discuss this issue in Chap. V.

### The cyclotomic case

We consider the special case that  $G = \mathbb{G}_m/\mathbb{Q}$  and our coefficient system  $\mathbb{Q}(n)$  is given by the character  $[n] : x \mapsto x^n$ . We fix a level  $K_f$  and we consider our isotypical decomposition over  $\mathbb{Q}$

$$H^0(\mathcal{S}_{K_f}^G, \mathbb{Q}(n)) = \bigoplus_{\Phi} \mathbb{Q}(\Phi_f).$$

In this case  $\mathbb{Q}(\Phi_f)$  is a field, and the action of the group is simply an irreducible action of the group of finite ideles  $G(\mathbb{A}_f) = I_{\mathbb{Q},f}$  on the  $\mathbb{Q}$ -vector space  $\mathbb{Q}(\Phi_f)$ . If we extend our field to  $\bar{\mathbb{Q}}$  we get a decomposition

$$H^0(\mathcal{S}_{K_f}^G, \bar{\mathbb{Q}}(n)) = \bigoplus_{\chi: \text{type}(\chi)=[n]} \bar{\mathbb{Q}}(\chi),$$

and the collection of conjugate characters  $\chi$  are in one to one correspondence with the  $\Phi_f$ . We can attach two different kinds of  $L$ -functions to our isotypical component  $\Phi_f$  namely an automorphic  $L$ -function and a motivic  $L$ -function.

Actually we get a collection of such  $L$ -functions which are labelled by the embeddings  $\iota : \mathbb{Q}(\Phi) \rightarrow \mathbb{Q} \subset \mathbb{C}$ . Such an embedding yields an algebraic Hecke character

$$\chi_f^{(\iota)} = \iota \circ \Phi_f : G(\mathbb{A}_f) = I_{\mathbb{Q},f} \rightarrow \bar{\mathbb{Q}}^\times$$

and

$$\chi^{(\iota)} = \iota \circ \Phi : G(\mathbb{Q}) \backslash G(\mathbb{A}) = \mathbb{Q}^\times \backslash I_{\mathbb{Q}} \rightarrow \mathbb{C}^\times$$

and to any of these Hecke characters we attach the (the automorphic  $L$ -function) namely

$$L(\chi^{(\iota)}, s) = \prod_p (1 - \chi^{(\iota)}(p)p^{-s})^{-1}$$

where  $\chi^{(\iota)}(p) = \chi^{(\iota)}(1, 1, \dots, p, \dots)$  and it is zero if the character is ramified.

Now we can attach a motive  $\mathbb{M}(\Phi)$  to our isotypical component. To do this we assume first that  $\mathbb{Q}(\Phi) = \mathbb{Q}$ , then we have only one embedding. Then we have  $\chi(x) = \alpha^n(x) = |x|^n$  for some integer  $n$ . This is an algebraic Hecke character of type  $[-n] : x \mapsto x^{-n}$ . Then we attach the motive  $\mathbb{Z}(-n)$  to this Hecke character. At this moment we do not need to know what a motive is, the only thing we need to know that it provides a compatible system of  $\ell$ -adic representations of the Galois group: For any prime  $\ell$  we define a module To this motive we attach a motivic  $L$  function using the compatible system of  $\ell$ -adic representations. For a prime  $\ell$  and a prime  $p \neq \ell$  we have the local Euler factor

$$L_p(\mathbb{Z}(-n), s) = \frac{1}{\det(1 - F_p^{-1} | \mathbb{Z}_\ell(-n)p^{-s})} = \frac{1}{1 - p^n p^{-s}},$$

where  $F_p$  is the Frobenius at  $p$ . The  $\ell$ -adic representation is unramified outside  $\ell$  and the Frobenius  $F_p$  corresponds to  $p$  under the reciprocity map  $r$ . Hence we see that the Frobenius  $F_p$  acts by the multiplication by  $\alpha^n(p) = |p|^n = p^{-n}$  on  $\mathbb{Z}_\ell(-n)$ . In the general case we start from the representation  $\Phi_f : I_{\mathbb{Q},f} \rightarrow \mathbb{Q}(\Phi_f)^\times$ , it is unramified outside a finite set  $\Sigma$  of primes. The reciprocity map from class field theory provides a homomorphism  $r : I_{\mathbb{Q},f} \rightarrow \text{Gal}_\Sigma(\bar{\mathbb{Q}}/\mathbb{Q})_{\text{abelian}}$ , this is the maximal abelian quotient of the Galois group which is unramified outside  $\Sigma$ , the image of the reciprocity map is dense. If we fix a prime  $\ell$  then we get an  $\ell$ -adic representation

$$\rho(\Phi) : \text{Gal}_\Sigma(\bar{\mathbb{Q}}/\mathbb{Q})_{\text{abelian}} \rightarrow (\mathbb{Q}(\Phi_f) \otimes \mathbb{Q}_\ell)^\times,$$

which is determined by the rule  $\rho(\Phi)(F_p) = \Phi_f(p)$ . If we now choose an embedding  $\iota : \mathbb{Q}(\Phi_f) \rightarrow \bar{\mathbb{Q}}$  and an extension  $\mathfrak{l}$  of  $\ell$  to a place of  $\bar{\mathbb{Q}}$  and we get a one dimensional  $\mathfrak{l}$  adic representation

$$\rho(\iota \circ \Phi) : \text{Gal}_\Sigma(\bar{\mathbb{Q}}/\mathbb{Q})_{\text{abelian}} \rightarrow \bar{\mathbb{Q}}_\mathfrak{l}^\times,$$

from which we get a motivic  $L$ -function  $(\mathbb{M}(\Phi) \circ \iota, s)$ , whose local factor at  $p$  is

$$L_p(\mathbb{M}(\Phi)^{(\iota)}, s) = \frac{1}{1 - \rho(\iota \circ \Phi)(F_p)^{-1} p^{-s}}$$

These are the collections of  $\ell$ -adic representations of our motives  $\mathbb{M}(\Phi)$ . Then the relation between the automorphic and the  $\ell$ -adic  $L$  functions is:

The collection of automorphic  $L$ -functions attached to  $\Phi$  is equal to the collection of motivic  $L$ -functions attached to  $\mathbb{M}(\Phi^{-1})$ .

We will sometimes modify the notation slightly. If  $\chi$  is an algebraic Hecke character then this datum corresponds to a pair  $(\Phi, \iota)$  and hence we can attach to it a character  $\chi_\iota : \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow \bar{\mathbb{Q}}_\iota$  and then we get the equality of local  $L$ -factors

$$L_p(\chi, s) = \frac{1}{1 - \chi(p)p^{-s}} = \frac{1}{1 - \chi_\iota^{-1}(F_p)^{-1}p^{-s}}$$

(Nochmal ein wenig besser schreiben!!!!!!!!!!!!!!!!!!!!)

### The $L$ -functions

Let us choose a cocharacter  $\chi : G_m \rightarrow T$ , we assume that it is in the positive chamber, i.e. we have  $\langle \chi, \alpha_i \rangle \geq 0$  for all positive simple roots. It yields an element  $\chi(p) \in T(\mathbb{Q}_p)$ . For  $\omega_p \in \Lambda(T)$  we put EVTchi

$$S_{\chi, \omega_p} = p^{\langle \chi, \rho \rangle} \sum_{w \in W/W_\chi} \omega_p(w(\chi(p))) \tag{7.2}$$

then we get a formula

$$\int_{\text{ch}(\chi(p))} \phi_{\omega_p}(xg)dg = (S_{\chi, \omega_p} + \sum_{\chi' < \chi} a(\chi, \chi') S_{\chi', \omega_p}) \phi_{\omega_p}(x) \tag{7.3}$$

where the  $\chi'$  are in the positive chamber,  $\chi' < \chi$  means that  $\chi - \chi' = \sum n_i \chi_i, n_i \geq 0$  and the coefficients  $a(\chi, \chi') \in \mathbb{Z}$ . The expression on the right hand side is invariant under  $W$  and hence only depends on  $\omega_p$  modulo  $W$ . Let (**Give reference!**)

The number  $\langle \chi, \rho \rangle$  is a half integer, hence  $p^{\langle \chi, \rho \rangle}$  may not lie in a fixed number field if  $p$  varies. But for those  $\chi'$  which may occur in the summation we have  $\langle \chi - \chi', \rho \rangle \in \mathbb{Z}$ .

We consider an unramified prime. The theorem of Satake yields that we can define a Hecke operator  $S_\chi \in \mathcal{H}_p$  such that  $S_\chi * \phi_{\omega_p} = S_{\chi, \omega_p} \phi_{\omega_p}$  and the formula (7.3) tells us that we get another recursion

$$S_\chi = \text{ch}(\chi) + \sum_{\chi' < \chi} b(\chi, \chi') \text{ch}(\chi') \tag{7.4}$$

where again  $b(\chi, \chi') \in \mathbb{Z}$ .

Since we assume that our absolutely irreducible module  $V_{\pi_f}, \pi_f = \otimes' \pi_p$  occurs in  $\text{Coh}(G, K_f, \lambda)$ , the Hecke module is a vector space over a finite extension  $F/\mathbb{Q}$ . We can conclude that the eigenvalue of the convolution operator  $\text{ch}(\chi)$  is in  $F$  and it follows that

$$S_{\chi, \omega_p} \in F$$

for any cocharacter  $\chi$ .

Since we can replace  $\chi$  by  $n\chi$  for any integer  $n \geq 1$  it follows that the numbers  $w(\chi(p))$  lie in a finite extension of  $F$  and the polynomial

$$\prod_{w \in W/W_\chi} (X \cdot \text{Id} - p^{\langle \chi, \rho \rangle} w(\chi(p))) \in F[X].$$

Our cocharacter  $\chi \in X_*(T)$  can also be interpreted as a character in  $X^*(T^\vee)$ , i.e it is a character on the dual torus. Since we assumed it to be in the positive chamber we can view  $\chi$  as the highest weight of an irreducible representation  $r_\chi : {}^L G \rightarrow \text{Gl}(\mathcal{E}_\chi)$ . (Since we assume that  $G$  is split the dual group is also split over  $\mathbb{Q}$  and hence  $r_\chi$  is defined over  $\mathbb{Q}$ .) The eigenvalues of the endomorphism  $r_\chi(\omega_p)$  are of the form  $\omega_p(w(\chi'(p)))$  where  $\chi' \leq \chi$  and this implies that the polynomial

$$\det(X \cdot \text{Id} - p^{\langle \chi, \rho \rangle} r_\chi(\omega_p)|\mathcal{E}_\chi) \in F[X].$$

We attach a local Euler factor to the data  $\pi_p, \omega_p = \omega(\pi_p), \chi$ :

$$L_p^{\text{rat}}(\pi_f, r_\chi, s) = \frac{1}{\det(\text{Id} - p^{\langle \chi, \rho \rangle} r_\chi(\omega_p) p^{-s} | \mathcal{E}_\chi)} \tag{7.5}$$

which is a formal power series in the variable  $p^{-s}$  with coefficients in  $F$ . We define

$$L^{\text{rat}}(\pi_f, r_\chi, s) = \prod_{p \in \Sigma} L_p(\pi_f, r_\chi, s) \left( \prod_{p \notin \Sigma} \frac{1}{\det(\text{Id} - p^{\langle \chi, \rho \rangle} r_\chi(\omega_p) p^{-s} | \mathcal{E}_\chi)} \right), \tag{7.6}$$

at the moment we do not say anything about the Euler factors at the bad primes.

At this moment  $L^{\text{rat}}(\pi_f, r_\chi, s)$  is a product of formal power series in infinitely many variables  $p^{-s}$  which in some sense encodes the collection of eigenvalues of the different Hecke eigenvalues.

We want to relate this  $L$ -function to some other  $L$ -functions which are defined in the theory of automorphic forms.

To define the automorphic  $L$ -function we start from an absolutely irreducible Hecke-module  $V_{\pi_f}$  over  $\mathbb{C}$ , its isomorphism type is still denoted by  $\pi_f$ . This  $\pi_f$  will be the first argument (in our notation) in the automorphic  $L$ -function. It has a central character  $\zeta_{\pi_f}$  and we assume that this central character is the finite component of a character  $\zeta_\pi : C(\mathbb{Q}) \backslash C(\mathbb{A}) \rightarrow \mathbb{C}^\times$ . (In the back of our mind of  $\pi_f$  to be the finite component of an automorphic form  $\pi$ , then this assumption is automatically fulfilled. But for the definition of the  $L$ -functions we do not need this.)

Then we define the unitary (automorphic)  $L$ -function: Here we require that the central character  $\zeta_{\pi_f}$  of  $\pi_f$  is unitary and put

$$L^{\text{unit}}(\pi_f, r_\chi, s) = \prod_{p \in \Sigma} L_p(\pi_f, r_\chi, s) \left( \prod_{p \notin \Sigma} \frac{1}{\det(\text{Id} - r_\chi(\omega_p) p^{-s} | \mathcal{E}_\chi)} \right) \tag{7.7}$$

If the central character is not unitary we define the automorphic  $L$ -function essentially by the same formula:

$$L^{\text{aut}}(\pi_f, r_\chi, s) = \prod_{p \in \Sigma} L_p(\pi_f, r_\chi, s) \left( \prod_{p \notin \Sigma} \frac{1}{\det(\text{Id} - r_\chi(\omega_p) p^{-s} | \mathcal{E}_\chi)} \right) \tag{7.8}$$

This  $L$ -function is related to an unitary  $L$ -function by a shift in the variable  $s$ . The isogeny  $d_C$  induces a homomorphism  $d' : C(\mathbb{Q}) \backslash C(\mathbb{A}) \rightarrow C'(\mathbb{Q}) \backslash C'(\mathbb{A})$  and it is well known that this map has a compact kernel. We compose  $\zeta_\pi$  with the norm  $|\cdot| : \mathbb{C}^\times \rightarrow \mathbb{R}_{>0}^\times$ , this composition is trivial on the kernel of  $d'$ . Therefore we find a homomorphism  $|\zeta_\pi|^* : C'(\mathbb{A}_f) \rightarrow \mathbb{R}_{>0}^\times$  which satisfies  $|\cdot| \circ \zeta_\pi = |\zeta_\pi|^* \circ d'$ . We look at the finite components of these characters and put as in (6.3.3)

$$\pi_f^* = \pi_f \otimes (|\zeta_\pi|^*)^{-1}. \quad (7.9)$$

This module has a unitary central character. It is easy to see how the Satake parameter changes under the twisting. We have the homomorphism  $T(\mathbb{A}) \rightarrow C'(\mathbb{A})$  and therefore  $(|\zeta_\pi|^*)^{-1}$  induces also a homomorphism from  $T(\mathbb{A}_f)$  to  $\mathbb{R}_{>0}^\times$ . Then it is clear that we get for the Satake parameters the equality

$$\omega(\pi_p \otimes (|\zeta_\pi|^*)^{-1}) = \omega(\pi_p)(|\zeta_\pi|^*)^{-1} \quad (7.10)$$

Let us assume that  $\pi_f$  occurs as an isotypical subspace in some  $H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{C})$ , where  $\lambda = \lambda^{(1)} + \delta$ . The element  $\delta$  is an element in  $X^*(C') \otimes \mathbb{Q}$ . To an element  $\eta \in X^*(C') \otimes \mathbb{R}$  we have attached an element  $|\eta|$  and since  $\zeta_{\pi_f}$  is of type  $\delta \circ d_C$  we have

$$(|\zeta_\pi|^*)^{-1} = |\delta|.$$

We also have the cocharacter  $\chi : \mathbb{G}_m \rightarrow T$  then it is clear that the composition  $(|\zeta_\pi|^*)^{-1} \circ \chi$  induces a homomorphism  $\mathbb{G}_m(\mathbb{Q}) \backslash \mathbb{G}_m(\mathbb{A}) \rightarrow \mathbb{R}_{>0}^\times$  which is of the form

$$((|\zeta_\pi|^*)^{-1} \circ \chi)_\mathbb{A} : \underline{x} \mapsto |\underline{x}|^{<\chi, \delta>}. \quad (7.11)$$

Then we have

$$L^{\text{unit}}(\pi_f^*, r_\chi, s) = L^{\text{aut}}(\pi_f, r_\chi, s + \langle \chi, \delta \rangle) \quad (7.12)$$

We now assume that  $\pi_f^*$  is the finite part of a cuspidal unitary representation (See 8.1.6), then the functions  $L^{\text{unit}}(\pi_f^*, r_\chi, s)$  are studied in the theory of automorphic forms. The Euler factors are now meromorphic functions in the variable  $s \in \mathbb{C}$ . Since  $\pi_f^*$  is unitary it follows that the Satake parameters satisfy some bounds and this implies that the infinite product converges if  $\Re(s) \gg 0$ . If for all  $p \notin \Sigma$  the representation  $\pi_p^*$  is in the unitary principal series, i.e.  $|\omega_{i,p}^*| = 1$  then it follows from standard arguments that the infinite product over  $p \notin \Sigma$  converges for  $\Re(s) > 1$ .

It is a conjecture (proved in some cases) that  $L^{\text{unit}}(\pi_f, r_\chi, s)$  has analytic continuation into the entire complex plane and that there is a functional equation relating  $L^{\text{unit}}(\pi_f, r_\chi, s)$  and  $L^{\text{unit}}(\pi_f^\vee, r_\chi, 1 - s)$ .

But of course any theorem proved for the  $L$ -functions  $L^{\text{unit}}(\pi_f^*, r_\chi, s)$  translates into a theorem for the automorphic  $L$  functions  $L^{\text{aut}}(\pi_f, r_\chi, s)$ .

Given a automorphic representation  $\pi$  which occurs in the cuspidal spectrum then we may twist it by any character  $\xi : C'(\mathbb{Q}) \backslash C'(\mathbb{A}) \rightarrow \mathbb{R}_{>0}^\times$ , this group of characters is equal to  $X^*(C') \otimes \mathbb{R}$ . We get a principal homogenous space (a torsor) of automorphic representations  $\{\pi \otimes \xi\}_{\xi \in \Xi}$ .

For the Euler factors  $p \notin \Sigma$  we have

$$\frac{1}{\det(\text{Id} - r_\chi((\omega_p)(\pi_p \otimes \xi_p))p^{-s}|\mathcal{E}_\chi)} = \frac{1}{\det(\text{Id} - r_\chi((\omega_p)(\pi_p))p^{-\langle \chi, \xi \rangle - s}|\mathcal{E}_\chi)} \tag{7.13}$$

and hence we get for our automorphic  $L$ -function

$$L^{\text{aut}}(\pi_f \otimes \xi_f, r_\chi, s) = L^{\text{aut}}(\pi_f, r_\chi, s + \langle \chi, \xi \rangle) \tag{7.14}$$

The representation  $\pi^*$  is then the unique cuspidal (in the above sense) representation in this principal homogeneous spaces  $\{\pi \otimes \xi\}_{\xi \in \Xi}$ , i.e. it is the unique representation which has a unitary central character. In other words  $\pi_f^*$  provides a trivialization of the torsor. Then we define for any  $\pi \otimes \xi$

$$L^{\text{unit}}(\pi_f \otimes \xi_f, r_\chi, s) = L^{\text{unit}}(\pi_f^*, r_\chi, s) \tag{7.15}$$

the unitary  $L$ -function is constant on the torsor, i.e. invariant under twisting.

We compare the automorphic  $L$ - function to the rational  $L$ - function. We start from an absolutely irreducible module  $\pi_f$  which occurs in  $\text{Coh}_1(G, K_f, \lambda)$  and which is defined over some finite extension  $F/\mathbb{Q}$ . As usual we write  $\lambda = \lambda^{(1)} + \delta$ , (See(6.22)). We know that the central character  $\zeta_{\pi_f}$  is an algebraic Hecke character of type  $\delta$ . Our Hecke module  $\pi_f$  is an absolutely irreducible module over  $F$ . If we want to compare its  $L$  functions to automorphic  $L$ -functions we need to choose an embedding  $\iota : F \hookrightarrow \mathbb{C}$  and consider the module  $V_{\pi_f} \otimes_{F, \iota} \mathbb{C} = V_{\iota \circ \pi_f}$ . The we will see in section 8.1.6 that  $\iota \circ \pi_f$  is the finite part of an automorphic representation occurring in the discrete (or the cuspidal) spectrum. Hence we have defined  $\tilde{L}^{\text{aut}}(\iota \circ \pi_f, r_\chi, s)$ . We can also consider the "extension" of the rational  $L$ -function

$$\iota \circ L^{\text{rat}}(\pi_f, r_\chi, s) = \prod_{p \in \Sigma} \iota \circ L_p^{\text{rat}}(\pi_f, r_\chi, s) \prod_{p \notin \Sigma} \frac{1}{\det(\text{Id} - \iota(p^{\langle \chi, \rho \rangle} r_\chi(\omega_p(\pi_p)))p^{-s}|\mathcal{E}_\chi)}$$

Then it is clear that

$$\iota \circ L^{\text{rat}}(\pi_f, r_\chi, s) = L^{\text{aut}}(\iota \circ \pi_f, r_\chi, s - \langle \chi, \rho \rangle). \tag{7.16}$$

The central character of  $\iota \circ \pi_f$  is of type  $\delta$ , it follows from (6.22) that some non zero multiple  $r\delta \in X^*(T)$ . Then we put  $\langle \chi, \delta \rangle = \frac{1}{r} \langle \chi, r\delta \rangle$ , this is a rational number. Then we get

$$\iota \circ L^{\text{rat}}(\pi_f, r_\chi, s) = L^{\text{aut}}(\iota \circ \pi_f, r_\chi, s - \langle \chi, \delta \rangle) \tag{7.17}$$

We still have another  $L$  function which is attached to a Hecke module  $\pi_f$  which occurs in the cohomology, this is the cohomological  $L$  function. Let us decompose the representation  $\mathcal{E}_\lambda$  into weight spaces

$$\mathcal{E}_\chi = \bigoplus_{\nu} \mathcal{E}_{\chi, \nu} = \bigoplus_{\nu \in X_{*,+}(T)} \bigoplus_{w \in W/W_\nu} \mathcal{E}_{\chi, w(\nu)}$$

then we get with  $m(\nu, \chi) = \dim(\mathcal{E}_{\chi, w(\nu)})$ . Such a weight vector space is zero unless we have  $\nu < \chi$ .

$$\det(\text{Id} - r_\chi(\omega_p)p^{-s}|\mathcal{E}_\chi) = \prod_{\nu \in X_{*,+}(T)} \prod_{w \in W/W_\nu} (1 - \omega_p(w(\nu))p^{-s})^{m(\nu, \chi)}$$

For a given  $\nu$  we expand the inner product

$$\prod_{w \in W/W_\nu} (1 - \omega_p(w(\nu))p^{-s}) = 1 - \left( \sum_{w \in W/W_\nu} \omega_p(w(\nu)) \right) p^{-s} \dots$$

Now we recall that

$$p^{\langle \chi, \lambda^{(1)} \rangle - \langle \chi, \delta \rangle} \text{ch}(\chi) = S_\chi^{(\lambda)}$$

is an operator on the integral cohomology (See (6.27)). Then our recursion formula (7.4) implies that

$$p^{\langle \chi, \lambda^{(1)} \rangle - \langle \chi, \delta \rangle} S_{\chi'}$$

is an operator on the integral cohomology, we simply have to observe that  $\langle \chi, \lambda^{(1)} \rangle \geq \langle \chi', \lambda^{(1)} \rangle$ . From this it follows directly that for  $\nu \in X_{*,+}(T)$  which occurs as a weight in  $r_\chi$  we have

$$p^{\langle \chi, \lambda^{(1)} + \rho \rangle - \langle \chi, \delta \rangle} \sum_{w \in W/W_\nu} \omega_p(w(\nu)) \in \mathcal{O}_F$$

because  $\langle \chi, \lambda^{(1)} \rangle > \langle \nu, \lambda^{(1)} \rangle$ . Then the right hand side in the above formula can be written

$$1 - p^{\langle \chi, \lambda^{(1)} + \rho \rangle - \langle \chi, \delta \rangle} \left( \sum_{w \in W/W_\nu} \omega_p(w(\nu)) \right) p^{-s - \langle \chi, \lambda^{(1)} + \rho \rangle + \langle \chi, \delta \rangle} \dots$$

We introduce the new variable  $s' = s + \langle \chi, \lambda^{(1)} + \rho \rangle - \langle \chi, \delta \rangle$  and put

$$c(\chi, \lambda) = \langle \chi, \lambda^{(1)} + \rho \rangle - \langle \chi, \delta \rangle \quad (7.18)$$

$$\prod_{w \in W/W_\nu} (1 - p^{c(\chi, \lambda)} \omega_p(w(\nu)) p^{-s'}) = 1 - p^{c(\chi, \lambda)} \left( \sum_{w \in W/W_\nu} \omega_p(w(\nu)) \right) p^{-s'} \dots \quad (7.19)$$

Hence we define the cohomological local Euler factor at  $p$

$$L_p^{\text{coh}}(\pi_f, r_\chi, s) = \frac{1}{\det(\text{Id} - p^{c(\chi, \lambda)} r_\chi(\omega_p) p^{-s})}. \quad (7.20)$$

(It seems to be reasonable and very adequate to define for any highest weight  $\lambda$  the modified weight  $\tilde{\lambda} = \lambda + \rho$ .)

We look at this local Euler factor from a slightly different point of view. Our  $\pi_f$  is an absolutely irreducible module which occurs in the cohomology  $H_i^*(\mathcal{S}_{K_f}^G, \mathcal{M}_\lambda \otimes F)$ , where  $F/\mathbb{Q}$  is an abstract (normal) finite extension of  $\mathbb{Q}$ . For an unramified prime  $p$  the local factor is simply a homomorphism  $\pi_p : \mathcal{H}_p \rightarrow E$ .

The previous computations show that the denominator is equal to a polynomial in the "variable"  $p^{-s}$  and with coefficients in  $\mathcal{O}_F$ , i.e.

$$\det(\text{Id} - p^{c(\chi, \lambda)} r_\chi(\omega_p) p^{-s}) = 1 - A_1(p, \lambda, \chi)(\pi_p) p^{-s} + A_2(p, \lambda, \chi)(\pi_p) p^{-2s} \dots \in \mathcal{O}_F[p^{-s}] \tag{7.21}$$

where the  $A_i(p, \lambda, \chi)$  are certain explicitly computable elements in  $\mathcal{H}_{\mathbb{Z}}^{(\lambda)}$ . (We showed this only for  $A_1(p, \lambda, \chi)$  but the same kind of reasoning gives it for the other  $A_i(p, \lambda, \chi)$ .) In the expression of the right hand side the Satake parameter does not enter.

The cohomological  $L$  function is defined as

$$L^{\text{coh}}(\pi_f, r_\chi, s) = \prod_{p \in \Sigma} L_p^{\text{coh}}(\pi_p, r_\chi, s) \prod_{p \notin \Sigma} \frac{1}{1 - A_1(p, \lambda, \chi)(\pi_p) p^{-s} + A_2(p, \lambda, \chi)(\pi_p) p^{-2s} \dots} \tag{7.22}$$

Again we do not discuss the factors at the primes in  $\Sigma$ .

In the definition of the automorphic  $L$  function the Satake parameter is an element in  ${}^L G(\mathbb{C})$  or in other words  $\omega_p(\nu) \in \mathbb{C}^\times$  and  $L_p^{\text{aut}}(\pi_f, r_\chi, s)$  is an honest analytic function in the complex variable  $s$  for  $\Re(s) \gg 0$ .

If we want to compare the cohomological  $L$ -function to the automorphic  $L$ -function we have to pick an element  $\iota \in I(F, \mathbb{C})$ , then  $\iota \circ \pi_f$  is an absolutely irreducible Hecke module over  $\mathbb{C}$ . To  $\iota \circ \pi_p$  belongs a Satake parameter  $\omega_p$  and then

$$\det(\text{Id} - r_\chi(\omega_p) p^{-s+c(\chi, \lambda)}) = 1 - \iota(A_1(p, \lambda, \chi)(\pi_p)) p^{-s} + \iota(A_2(p, \lambda, \chi)(\pi_p)) p^{-2s} \dots$$

and this tells us that we have

$$L^{\text{coh}}(\iota \circ \pi_f, r_\chi, s) = L^{\text{aut}}(\iota \circ \pi_f, r_\chi, s - c(\chi, \lambda)) \tag{7.23}$$

**Invariance under twisting**

We remember that we introduced the quotient  $\mathcal{C}' = \mathcal{T}/\mathcal{T}^{(1)}$  and the isogeny  $d_C : \mathcal{C} \rightarrow \mathcal{C}'$ . (See 6.1.1). The map  $d_C$  in 1.1 induces a map from our locally symmetric space

$$\mathcal{S}_{K_f}^G \xrightarrow{d_{C'}} \mathcal{S}_{K_\infty^{C'} \times K_f^{C'}}^{C'}$$

We assume that  $K_\infty$  is connected and then  $K_\infty^{C'}$  is also connected.

We can modify our system of coefficients if we replace  $\lambda$  by  $\lambda + \delta_1$  with  $\delta_1 \in X^*(\mathcal{C}')$ . Then  $\delta_1$  provides a local coefficient system  $\mathbb{Z}[\delta_1]$  on  $\mathcal{S}_{K_\infty^{C'} \times K_f^{C'}}^{C'}$  and since  $K_\infty^{C'}$  is connected we get a canonical class

$$e_{\delta_1} \in H^0(\mathcal{S}_{K_\infty^{C'} \times K_f^{C'}}^{C'}, \mathbb{Z}[\delta_1])$$

which generates the rank one submodule of type  $|\delta_f|^{-1}$  in the decomposition (6.42). We pull this back by  $d_C'$  and we get a class in

$$e_{\delta_1} \in H^0(\mathcal{S}_{K_f}^G, \mathbb{Z}[\delta_1]) \tag{7.24}$$

(see section (6.3.6)). We have the isomorphism  $\mathcal{M}_{\lambda, \mathbb{Z}} \otimes \mathbb{Z}[\delta_1] \xrightarrow{\sim} \mathcal{M}_{\lambda+\delta_1, \mathbb{Z}}$  and then the cup product with  $e_{\delta_1}$  yields an isomorphism

$$H_!^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda, \mathbb{Z}}) \cup e_{\delta_1} \xrightarrow{\sim} H_!^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda+\delta_1, \mathbb{Z}}) \quad (7.25)$$

This isomorphism is compatible with the action of the integral Hecke algebra provided we choose the right identification

$$\mathcal{H}_{\mathbb{Z}}^{(\lambda)} \rightarrow \mathcal{H}_{\mathbb{Z}}^{(\lambda+\delta_1)}$$

which is given by  $a \cdot \mathbf{ch}(\underline{x}_f) \mapsto p^{\langle \mathbf{ch}(\underline{x}_f), \delta_1 \rangle} a \cdot \mathbf{ch}(\underline{x}_f)$ .

If we extend the coefficients to  $F$  then this cup product yields an isomorphism

$$H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda, F})(\pi_f) \xrightarrow{\sim} H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda+\delta_1, F})(\pi_f \otimes |\delta_{1, f}|^{-1}) \quad (7.26)$$

Then our cohomological  $L$ -function has the property

$$L^{\text{coh}}(\pi_f \otimes |\delta_{1, f}|^{-1}, r_\chi, s) = L^{\text{coh}}(\pi_f, r_\chi, s) \quad (7.27)$$

This invariance under twists is of course also a consequence of the definition in terms of the automorphic  $L$ -function.

We may interpret this differently. Our  $\lambda$  is a sum of a semi-simple component  $\lambda^{(1)}$  plus an abelian part  $\delta$ . We can use the isomorphisms in (7.26) to define a vector space

$$H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda^{(1)+}, F})\{\pi_f\}, \quad (7.28)$$

this vector space has a distinguished isomorphism to any of the  $H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda+\delta_1, F})(\pi_f \otimes |\delta_{1, f}|^{-1})$ , we could say that it is the direct limit of all these spaces. By  $\{\sigma_f\}$  we understand the array

$$\{\sigma_f\} = \{\dots, \pi_f \otimes |\delta_{1, f}|^{-1}, \}_{\delta_1 \in X^*(C')}.$$

Using (7.27) we have now defined  $L^{\text{coh}}(\{\pi_f\}, r_\chi, s)$

For any pair  $\chi \in X_*(T)$ ,  $\lambda \in X^*(T)$ , where  $\chi$  is in the positive chamber and  $\lambda$  a dominant weight we define the weight

$$\mathbf{w}(\chi, \lambda) = \langle \chi, \lambda^{(1)} + \rho \rangle. \quad (7.29)$$

Here we observe that  $\chi$  provides a highest weight representation  $r = r_\chi$  of  ${}^L G$  and  $\lambda$  a highest weight representation of  $G$  so we could also write

$$\mathbf{w}(\chi, \lambda) = \mathbf{w}(r_\chi, \mathcal{M}_\lambda) = \mathbf{w}(r, \mathcal{M}). \quad (7.30)$$

This means that we may consider the weight as a number attached to a pair of irreducible rational representations of  ${}^L G$  and  $G$ . It also depends only on the semi simple part of  $\lambda$ .

**A different look**

We could look at the previous discussion from another point of view. Given our coefficient system  $\mathcal{M}_\lambda$  where  $\lambda = \lambda^{(1)} + \delta$  and an absolutely irreducible module  $\pi_f \in \text{Coh}_!(G, \lambda, K_f)$ . As explained above we get  $X^*(C')$  torsor  $(\lambda + \delta', \pi_f \otimes |\delta'_f|)$  of such objects. If we choose a  $\iota : F \hookrightarrow \mathbb{C}$  then we can think of  $\iota \circ \pi_f$  as the finite part of an automorphic representation  $\pi$ . Then we get a second torsor for the above group  $\Xi = X^*(C') \otimes \mathbb{R}$ . The inclusion  $X^*(C') \hookrightarrow \Xi$  yields an interpolation of the first torsor into the second one. To any element  $\pi \otimes \xi$  we defined the automorphic  $L$  function  $L^{\text{aut}}(\iota \circ \pi_f \otimes \xi_f, r_\chi, s)$ . Now the unitary and the cohomological  $L$ -function are defined as the automorphic  $L$  function of a specific point in the torsor, i.e. a specific trivialization.

To define the unitary  $L$  function we choose the specific point for which the central character is unitary, for the cohomological  $L$ -function we choose the "optimal" point  $\pi_f \otimes |\delta'_f|$  for which we have

$$L_p^{\text{coh}}(\pi_f \otimes |\delta'_f|, r_\chi, s)^{-1} \in \mathcal{O}_F[p^{-s}]. \tag{7.31}$$

If we are investigating analytic questions concerning automorphic forms the unitary  $L$  is the right object, but if we want to capture the integral structure of the cohomology we prefer to work with the cohomological  $L$  function.

**The motives**

We consider an isotypical submodule  $H_!^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda;F})(\pi_f)$  in the inner cohomology. The Langlands philosophy predicts the existence of a collection of pure motives over  $\mathbb{Q}$  with coefficients in  $F$ .

$$\{\mathbb{M}(\pi_f, r_\chi)\}_{r_\chi}$$

which has certain properties. We will not be absolutely precise in the following but we list certain properties this motive should have. We should assume that  $\pi_f$  is not some kind of exceptional Hecke module (for instance it should not be endoscopic), and I can not give a precise definition what that means. We will make it more precise later when we discuss the case that our group is  $\text{Gl}_n$ .

This motive should be invariant under twists, i.e. we want that

$$\mathbb{M}(\pi_f \otimes |\delta_f|, r_\chi) = \mathbb{M}(\pi_f, r_\chi)$$

First of all this motive has a Betti-realization  $\mathbb{M}(\pi_f, r_\chi)_B$ , which is simply an  $F$  vector space of dimension  $\dim(r_\chi)$ . Such a motive has a de-Rham realization  $\mathbb{M}(\pi_f, r_\chi)_{dRh}$ , this is another  $F$ -vector space of the same dimension. It has a descending filtration

$$\begin{aligned} \mathbb{M}(\pi_f, r_\chi)_{dRh} &= F^0(\mathbb{M}(\pi_f, r_\chi)_{de-Rh}) \supset F^1(\mathbb{M}(\pi_f, r_\chi)_{de-Rh}) \supset \dots \\ \dots &\supset F^{\mathbf{w}}(F^0(\mathbb{M}(\pi_f, r_\chi)_{dRh}) \supset F^{\mathbf{w}+1}(F^0(\mathbb{M}(\pi_f, r_\chi)_{dRh})) = 0. \end{aligned}$$

The number  $\mathbf{w} = \mathbf{w}(\pi_f, \chi)$  is the weight of the motive it is equal to  $\mathbf{w}(\chi, \lambda)$ .

Furthermore we have a comparison isomorphism

$$I_{B-dRh} : \mathbb{M}(\pi_f, r_\chi)_B \otimes \mathbb{C} \xrightarrow{\sim} \mathbb{M}(\pi_f, r_\chi)_{dRh} \otimes \mathbb{C},$$

this yields periods and these periods should be related to  $\pi_f$ , this is rather mysterious.

For any prime  $\ell$  and any prime  $l \nmid \ell$  in  $F$  we get a Galois representation

$$\rho(\pi_f, \chi) : \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow GL(\mathbb{M}(\pi_f, r_\chi)_B \otimes F_l)$$

which is unramified outside  $\Sigma \cup \{l\}$  and for any such prime we have

$$\det(\text{Id} - \rho(\pi_f, \chi)(\Phi_p^{-1})p^{-s}, \mathbb{M}(\pi_f, r_\chi)_B \otimes F_l) = L_p^{\text{coh}}(\pi_f, r_\chi, s)^{-1},$$

or in other words we expect that the semi-simple conjugacy classes

$$\rho(\pi_f, \chi)(\Phi_p^{-1}) \sim p^{c(\chi, \lambda)} r_\chi(\omega_p) \quad (7.32)$$

and hence we want

$$L^{\text{coh}}(\pi_f, r_\chi, s) = L(\mathbb{M}(\pi_f, r_\chi), s)$$

The existence of these hypothetical motives has a lot of consequences. Once we have established such a relation

$$L^{\text{coh}}(\pi_f, r_\chi, s) = L(\mathbb{M}(\pi_f, r_\chi), s)$$

then we can exploit this in both directions. We have a certain chance to prove the conjectural analytic properties and the conjectural functional equation for the  $L$ -function of the motive  $\mathbb{M}(\pi_f, r_\chi)$ , provided we can prove this for  $L^{\text{coh}}(\pi_f, r_\chi, s)$ . On the automorphic side we know many cases in which we can prove these properties of the  $L$ -function using the theory of automorphic forms.

In the other direction we have Deligne's theorem concerning the absolute values of the Frobenius. This implies Ramanujan (more details later)

We seem to be very far away from proving these conjectures, but there are many instances where some parts of this program have been established and there are also some very interesting cases where this correspondence has been verified experimentally.

### The case $G = \text{Gl}_n$

#### Notations for the dual group ${}^L G$

We want to verify formula (7.3) in the special case  $G = \text{Gl}_n/\mathbb{Z}$ . In this case we have the cocharacters  $\chi_i$  which send  $t$  to the diagonal matrix  $t \mapsto \text{diag}(t, \dots, t, 1, \dots, 1)$  where  $t$  is placed to the first  $i$  dots. They satisfy  $\langle \chi_i, \alpha_j \rangle = \delta_{i,j}$  for  $1 \leq i \leq n, 1 \leq j \leq n-1$ . They are uniquely determined by this condition modulo the cocharacter  $\chi_n$  which identifies  $\mathbb{G}_m$  with the center. For  $1 \leq \nu \leq n-1$  the cocharacter  $\chi_i$  determines a maximal parabolic subgroup  $P_i \supset T$  whose roots  $\Delta_{P_i} = \{\alpha \mid \langle \chi_i, \alpha \rangle \geq 0\}$ . The parabolic subgroup  $P_i^-$  will be the opposite parabolic subgroup.

Let  $\eta_i : \mathbb{G}_m \rightarrow T$  be the cocharacter which sends  $t$  to  $t$  on the  $i$ -th spot on the diagonal and to 1 at all others. If we identify the module of cocharacters with the character group of the dual torus  $T^\vee \subset {}^L G = \text{Gl}_n$  then the differences  $\eta_i - \eta_j$  will be the roots, the simple roots are  $\eta_i - \eta_{i+1}$  and the fundamental dominant weights are the semi simple components  $(\sum_{i=1}^i \eta_i)^{(1)}$ .

**Formulas for the Hecke operators**

We consider the homomorphism  $r : K_p = \text{Gl}_n(\mathbb{Z}_p) \rightarrow \text{Gl}_n(\mathbb{F}_p)$  then we check easily that the intersection  $K_p \cap \chi_i(p)K_p\chi_i(p)^{-1} = K_p^{(\chi_i(p))}$  is the inverse image of the parabolic subgroup  $P_i^-(\mathbb{F}_p)$  under  $r$ .

We want to evaluate the integral

$$\int_{K_p\chi_i(p)K_p} \phi_{\omega_p}(x)dx$$

We write choose representatives  $\xi$  for the cosets of  $K_p/K_p^{(\chi_i(p))}$  and write  $K_p = \cup_{\xi} \xi K_p^{(\chi_i(p))}$ . We observe that  $\phi_{\omega_p}$  is constant on the cosets  $\xi K_p^{(\chi_i(p))}$ . Hence we see that

$$\int_{K_p\chi_i(p)K_p} \phi_{\omega_p}(x)dx = \sum_{\xi} \phi_{\omega_p}(\xi\chi_i(p)) \tag{7.33}$$

The Bruhat decomposition gives us a nice system of representatives for  $K_p/K_p^{(\chi_i(p))} = \text{Gl}_n(\mathbb{F}_p)/P_i^-(\mathbb{F}_p)$ . Let  $W_{M_i}$  be the Weyl group of the standard Levi subgroup  $M_i = P_i \cap P_i^-$  and we choose a system of representatives  $W^{P_i}$  for  $W/W_{M_i}$ . Then we get a disjoint decomposition

$$\text{Gl}_n(\mathbb{F}_p) = \bigcup_{w \in W^{P_i}} U_B(\mathbb{F}_p)wP_i^-(\mathbb{F}_p),$$

here  $U_B$  is the unipotent radical of the standard Borel subgroup. The function  $\phi_{\omega_p}$  is constant on the double cosets. If we write a representative in the form  $\xi = uw$  then the factor  $w$  is determined by  $\xi$  but the factor  $u$  is not. This factor is only unique up to multiplication from the right by a factor  $u \in U_B^{(w,-)}(\mathbb{F}_p) = U_B(\mathbb{F}_p) \cap wP_i^-w^{-1}(\mathbb{F}_p)$ . Hence we may choose our  $u$  in the subgroup

$$U_B^{(w,+)}(\mathbb{F}_p) = \prod_{\alpha \in \Delta^+ | \langle \chi_i, w^{-1}\alpha \rangle > 0} U_{\alpha}(\mathbb{F}_p) \tag{7.34}$$

and our sum in (7.33) becomes

$$\sum_{w \in W^{P_i}} \sum_{u \in U_B^{(w,+)}(\mathbb{F}_p)} \phi_{\omega_p}(uw\chi_i(p)) = \sum_{w \in W^{P_i}} p^{l(w)} \phi_{\omega_p}(w\chi_i(p)w^{-1}) \tag{7.35}$$

where  $l(w)$  is the cardinality of the set  $\{\alpha \in \Delta^+ | \langle \chi_i, w^{-1}\alpha \rangle > 0\}$ . We recall

the definition of the spherical function and get for our integral

$$\sum_{w \in W/W_{M_i}} p^{l(w)} \omega_p(w\chi_i(p)w^{-1})|\rho|_p(w\chi_i(p)w^{-1}) = \sum_{w \in W/W_{M_i}} p^{l(w) - \langle \chi_i, w^{-1}\rho \rangle} \omega_p((w\chi_i)(p)) \tag{7.36}$$

Now one checks easily that  $p^{l(w) - \langle \chi_i, w^{-1}\rho \rangle} = p^{\langle \chi_i, \rho \rangle}$  and hence we get the desired formula

$$\int_{K_p\chi_i(p)K_p} \phi_{\omega_p}(x)dx = p^{\langle \chi_i, \rho \rangle} \sum_{w \in W/W_{M_i}} \omega_p((w\chi_i)(p)) \tag{7.37}$$

This is the formula (7.3) for the group  $\mathrm{Gl}_n$  and the special choice of the cocharacters  $\chi = \chi_i$ . The only cocharacter  $\chi' < \chi_i$  is the trivial cocharacter, in our situation its contribution to (7.3) is zero.

Let us have a brief look at an arbitrary reductive (split or may be only quasisplit) group  $G/\mathbb{Q}$ , let us assume that the center is a connected torus  $C/\mathbb{Q}$ . We choose a maximal torus  $T/\mathbb{Q}$  which is contained in a Borel subgroup  $B/\mathbb{Q}$ . We have the homomorphism to the adjoint group  $G \rightarrow G_{\mathrm{ad}}$  it maps  $T$  to  $T_{\mathrm{ad}} = T/C$ . Again we may also define the fundamental cocharacters  $\chi_i : \mathbb{G}_m \rightarrow T$  which satisfy  $\langle \chi_i, \alpha_j \rangle = \delta_{i,j}$ . They are only well defined modulo cocharacters  $\chi : \mathbb{G}_m \rightarrow C$  but this does not matter so much. Our above method to compute the eigenvalue of  $\mathrm{ch}(\chi_i)$  still works if the cocharacter  $\chi_i$  is "minuscule" which means that  $\langle \chi_i, \alpha_j \rangle \in \{-1, 0, 1\}$ . In this case the formula (7.37) is still valid, again there is no contribution from the trivial character.

We return to  $G = \mathrm{Gl}_n$  and to our speculations about motives. We choose a weight module  $\mathcal{M}_\lambda$  where  $\lambda = \sum_i a_i \gamma_i + d\delta$ , where the  $\gamma_i$  are the fundamental weights and  $\delta$  is the determinant. The  $a_i$  are integers and we have the consistency condition  $\sum a_i \equiv nd \pmod n$ . Let us pick an isotypical submodule  $H^\bullet(\mathcal{S}_{K_f}^G, \mathcal{M}_\lambda \otimes F)(\pi_f)$ . In section 6.3.2 we define the Hecke operators

$$T_\chi^{\mathrm{coh}, \lambda} : H_\sharp^\bullet(\mathcal{S}_{K_f}^G, \mathcal{M}_\lambda) \rightarrow H_\sharp^\bullet(\mathcal{S}_{K_f}^G, \mathcal{M}_\lambda)$$

and these endomorphisms induce endomorphisms

$$T_\chi^{\mathrm{coh}, \lambda} : H_{\sharp, \mathrm{int}}^\bullet(\mathcal{S}_{K_f}^G, \mathcal{M}_\lambda \otimes F)(\pi_f) \rightarrow H_{\sharp, \mathrm{int}}^\bullet(\mathcal{S}_{K_f}^G, \mathcal{M}_\lambda \otimes F)(\pi_f)$$

Let  $\pi_f = \otimes \pi_p$  be an irreducible Hecke module and at an unramified place  $p$  let  $\omega_p$  be the Satake parameter. Our Satake parameter is determined by the  $n$ -tuple of numbers

$$\omega_p(\eta_i(p)) = \omega_{i,p} \text{ for } i = 1, \dots, n$$

The cocharacter  $\chi_n : \mathbb{G}_m \rightarrow T$  identifies  $\mathbb{G}_m$  with the center of  $\mathrm{Gl}_n$ . Our Hecke-module  $\pi_f$  has a central character and this provides a Hecke character

$$\pi_f \circ \chi_n : \mathbb{G}_m(\mathbb{A}_f) = I_{\mathbb{Q},f} \rightarrow F^\times$$

The restriction of  $\mathcal{M}_\lambda$  to  $\mathbb{G}_m$  is the character  $\omega_\lambda : t \mapsto t^{nd}$  and the type of  $\pi_f \circ \chi_n$  is of course  $\omega_\lambda$ .

Our cocharacters  $\chi_i$  define representations of the dual group which is again  $\mathrm{Gl}_n$  and in fact  $\chi_1$  yields the tautological representation  $r_1 : \mathrm{Gl}_n \xrightarrow{\sim} \mathrm{Gl}(V)$ . Then  $\chi_i$  yields the representation  $r_i = \Lambda^i(r_1) : \mathrm{Gl}_n \rightarrow \mathrm{Gl}(\Lambda^i(V))$ . For any subset  $I \subset \{1, 2, \dots, n\}$  we define

$$\omega_{I,p} = \prod_{i \in I} \omega_{i,p}$$

and then our formula (7.37) in combination with the formula (6.27) in section 6.3.2 and the observation that  $\langle \chi_i, \delta \rangle = i$  yields

$$T_{\chi_i}^{\mathrm{coh}, \lambda}(\pi_p) = p^{\langle \chi_i, \lambda^{(1)+\rho} \rangle - id} \sum_{I: \#I=i} \omega_{I,p} \quad (7.38)$$

and by the same token we get for the cohomological  $L$ -function

$$L^{\text{coh}}(\pi_f, r_\nu, s) = \prod_{p \in S} L_p^{\text{coh}}(\pi_f, r_i, s) \prod_{p \notin S} \left( \prod_{I: \#I=i} \frac{1}{(1 - p^{\langle \chi_i, \lambda^{(1)} + \rho \rangle - id} \omega_{I,p} p^{-s})} \right) \tag{7.39}$$

Here we see in a very transparent way the independence of the twist: If we modify  $\lambda$  to  $\lambda + r\delta$  then we have to modify  $\pi_f$  to  $\pi_f \otimes |\delta_f|^{-r}$ . This means that the  $\omega_{I,p}$  get multiplied by  $p^{ir}$  and the modifications cancel out.

We assume that  $\pi_f \in \text{Coh}(H_1^*(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda))$ , then we will see in section 8.1.6 that  $\pi_f$  is essentially unitary. The central character of  $\mathcal{M}_\lambda$  is  $x \mapsto x^{nd}$  and hence we get that  $\pi_f^* = \pi_f \otimes |\delta_f|^d$  is unitary. Then the Satake parameter of  $\pi_f^*$  is given by

$$\omega_{i,p}^* = \omega_{i,p} p^{-d} \text{ for } i = 1, \dots, n \tag{7.40}$$

where the factor  $p^{-d} = |p|_p^d$  and we observe that these numbers are also invariant under twists by a power of  $|\delta_f|$ .

Since the operators  $T_{\chi_i}^{\text{coh}, \lambda}$  operate on the integral cohomology it follows that the numbers  $T_{\chi_i}^{\text{coh}, \lambda}(\pi_f)$  are algebraic integers. We easily check that for all  $i \leq n$

$$i(\langle \chi_1, \lambda^{(1)} + \rho \rangle - d) \geq \langle \chi_i, \lambda^{(1)} + \rho \rangle - id$$

and this implies that the numbers

$$\sum_{I: \#I=i} \prod_{\nu \in I} p^{\langle \chi_1, \lambda^{(1)} + \rho \rangle - d} \omega_{\nu,p}$$

are algebraic integers and hence we can conclude

*The numbers*

$$\tilde{\omega}_{i,p} = p^{\langle \chi_1, \lambda^{(1)} + \rho \rangle - d} \omega_{i,p} = p^{\langle \chi_1, \lambda^{(1)} + \rho \rangle} \omega_{i,p}^* \tag{7.41}$$

*are algebraic integers*

Observe that these numbers are invariant under twists by a power of  $|\delta_f|$ .

We want to make a few remarks about the relationship between the automorphic and the cohomological  $L$ -functions, especially we comment the shift in the variable  $s$ .

For the automorphic  $L$ -function we assume that we are over  $\mathbb{C}$ , we have chosen an embedding  $\iota : F \hookrightarrow \mathbb{C}$ . If our isotypical Hecke module  $\pi_f$  is cuspidal (see Thm. 8.1.1) then the considerations around this theorem show that  $\pi_f$  is essentially unitary. The center  $C = \mathbb{G}_m$ , the quotient  $C' = \mathbb{G}_m$  and the isogeny  $d_C : x \mapsto x^n$ .

We come back to the Langlands philosophy. It predicts that for our a "cuspidal"  $\pi_f$  and the cocharacter  $\chi_1$  we should be able to attach a motive  $\mathbb{M}(\pi_f, r_1) = \mathbb{M}(\pi_f, \chi_1)$  with coefficients in  $F$ . This motive provides a compatible system of  $\mathbb{L}$ -adic Galois representations

$$\rho_\iota(\pi_f, \chi_1) : \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{Gl}_n(F_\iota) = \text{Gl}(\mathbb{M}(\pi_f, \chi_1)_{\text{ét}, \iota}) \tag{7.42}$$

which are unramified outside  $\{l\} \cup S$  and for  $p \notin S \cup \{l\}$  we should have

$$\det(\text{Id} - \rho_\iota(\pi_f, \chi_1)(\Phi_p^{-1})p^{-s}) = \prod_i (1 - p^{\langle \chi_1, \lambda^{(1)} + \rho \rangle - d} \omega_{i,p} p^{-s}) \quad (7.43)$$

and this means that up to the local factors at the bad primes we should have

$$L^{\text{mot}}(\mathbb{M}(\pi_f, \chi_1), s) = L^{\text{coh}}(\pi_f, \chi_1, s) \quad (7.44)$$

The existence of the compatible system of Galois representation has been shown by Harris - Kai-Wen Lan -Taylor and Thorne and by P. Scholze.

Once we have the motive for the cocharacter  $\chi_1$  we easily get the other  $\chi_i$  we simply have to look at the exterior powers  $\Lambda^i(\mathbb{M}(\pi_f, \chi_1))$ .

Now we see that that numbers  $\tilde{\omega}_{\nu,p}$  can be interpreted as the eigenvalues of the Frobenius on  $\mathbb{M}_{\text{ét},\iota}(\pi_f, \chi_1)$ . Under the assumption that  $\pi_f$  is "cuspidal" we expect that the motive  $\mathbb{M}(\pi_f, \chi_1)$  is pure of weight  $\mathbf{w}(\chi_1, \lambda)$  we get

$$|\tilde{\omega}_{\nu,p}| = p^{\frac{\mathbf{w}(\chi_1, \lambda)}{2}}$$

and this is the Ramanujan conjecture. We will explain in the section on analytic aspects, that for cuspidal  $\pi_f$  the Ramanujan conjecture says that for any embedding  $\iota : F \hookrightarrow \mathbb{C}$  we have

$$|\iota \circ \omega_{\nu,p}^*| = 1$$

This suggests that we call the array  $\tilde{\omega}_p = \{\tilde{\omega}_{1,p}, \dots, \tilde{\omega}_{n,p}\}$  the *motivic* Satake parameter (with respect to the tautological representation  $r_1$ .) Of course it can always be defined, independently of the existence of the motive.

We will see in the next section that the inner cohomology is trivial unless our highest weight is essentially self dual, this means that  $\lambda^{(1)} = -w_0(\lambda^{(1)})$ . Let us assume that this is the case. If  $r_1^\vee$  is the dual of the tautological representation then the eigenvalues of  $r_1^\vee(\omega_p)$  are by

$$r_1^\vee(\omega_p) = \{\omega_{1,p}^{-1}, \dots, \omega_{n,p}^{-1}\}.$$

The highest weight of  $r_1^\vee$  is the cocharacter  $-\eta_n = \sum_{i=1}^{n-1} \eta_i - \det$  (This has to be read in  $X^*(T^\vee)$ ) Then

$$c(-\eta_n, \lambda) = \langle \chi_1, -w_0(\lambda^{(1)}) \rangle + d$$

and under our assumption that  $\lambda$  is essentially self dual we know

$$\langle \chi_1, -w_0(\lambda^{(1)}) \rangle = \langle \chi_1, \lambda^{(1)} \rangle = \frac{\mathbf{w}(\chi_1, \lambda)}{2}.$$

This implies that the motivic Satake parameters with respect to the dual representation  $r_1^\vee$  are the numbers

$$\{p^{\langle \chi_1, \lambda^{(1)} \rangle + d\delta} \omega_{1,p}^{-1}, \dots, p^{\langle \chi_1, \lambda^{(1)} \rangle + d\delta} \omega_{n,p}^{-1}\} \quad (7.45)$$

In the following section on Poincaré duality we will see that for any isotypical module  $H_1^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda, F})(\pi_f)$  the dual module  $\pi_f^\vee$  appears in  $H_1^{d-i}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda^\vee, F})$ . Then we get an equality of local Euler factors

$$L^{\text{coh}}(\pi_p, r_1^\vee, s) = L^{\text{coh}}(\pi_p^\vee, r_1, s) \quad (7.46)$$

The concept of motives allows us to define the dual motive. If our motive has weight  $\mathbf{w}(M)$  then Poincaré duality suggests that we define the motive

$$\mathbb{M}^\vee = \text{Hom}(\mathbb{M}, \mathbb{Z}(-\mathbf{w}(M))) \quad (7.47)$$

The  $\ell$  adic realization as Galois module gives us

$$\mathbb{M}_{\text{ét}, \ell}^\vee = \text{Hom}(\mathbb{M}_{\text{ét}, \ell}, \mathbb{Z}_\ell(-\mathbf{w}(M)))$$

If  $\{\alpha_1, \dots, \alpha_m\}$  are the eigenvalues of  $\Phi_p^{-1}$  on  $\mathbb{M}_{\text{ét}, \ell}$  then  $\{\alpha_1^{-1}p^{\mathbf{w}(M)}, \dots, \alpha_m^{-1}p^{\mathbf{w}(M)}\}$  are the eigenvalues of  $\Phi_p^{-1}$  on  $\mathbb{M}_{\text{ét}, \ell}^\vee$ .

Therefore we can say: If we find a motive  $\mathbb{M}(\pi_f, \chi_1)$  for  $\pi_f$  then we also find the motive for  $\pi_f^\vee$  and we have

$$\mathbb{M}(\pi_f^\vee, \chi_1) = \mathbb{M}(\pi_f, \chi_1)^\vee$$



# Chapter 8

## Analytic methods

### 8.1 The representation theoretic de-Rham complex

#### 8.1.1 Rational representations

We start from a reductive group  $G/\mathbb{Q}$  for simplicity we assume that the semi simple component  $G^{(1)}/\mathbb{Q}$  is quasisplit. There is a unique finite normal extension  $F/\mathbb{Q}, F \subset \mathbb{C}$  such that  $G^{(1)} \times_{\mathbb{Q}} F$  becomes split, if  $T^{(1)}/\mathbb{Q}$  is a maximal torus which is contained in a Borel subgroup  $B/\mathbb{Q}$  then the Galois group  $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$  acts on  $X^*(T^{(1)} \times_{\mathbb{Q}} F)$  and by permutations on the set of positive roots  $\pi_G \subset X^*(T^{(1)} \times_{\mathbb{Q}} F)$  corresponding to  $B/\mathbb{Q}$ . This action factors over the quotient  $\text{Gal}(F/\mathbb{Q})$ . Then it also acts on the set of highest weights. Since our group is quasi split we find for any highest weight an absolutely irreducible  $G \times_{\mathbb{Q}} F$ -module  $\mathcal{M}_\lambda$ .

$$r : G \times_{\mathbb{Q}} K \rightarrow \text{Gl}(\mathcal{M}_\lambda)$$

whose highest weight is  $\lambda$ . Since we assumed that  $\mathbb{Q} \subset F \subset \bar{\mathbb{Q}} \subset \mathbb{C}$  we get the extension

$$r_{\mathbb{C}} : (G \times_{\mathbb{Q}} K) \times_K \mathbb{C} \rightarrow \text{Gl}(\mathcal{M}_\lambda \otimes_F \mathbb{C}).$$

Given such an absolutely irreducible rational representation, we can construct two new representations. At first we can form the dual  $\mathcal{M}_{\lambda, \mathbb{C}}^\vee = \text{Hom}_{\mathbb{C}}(\mathcal{M}_\lambda, \mathbb{C})$  and the complex conjugate  $\bar{\mathcal{M}}_{\mathbb{C}}$  of our module  $\mathcal{M}_\lambda$ . On the dual module we have the contragredient representation  $r^\vee$ , which is defined by  $\phi(r_{\mathbb{C}}(g)(v)) = r_{\mathbb{C}}^\vee(g^{-1})(\phi)(v)$ .

To get the rational representation on the conjugate module  $\bar{\mathcal{M}} \otimes_F \mathbb{C}$ , we recall its definition: As abelian groups we have  $\mathcal{M} \otimes_F \mathbb{C} = \bar{\mathcal{M}} \otimes_F \mathbb{C}$  but the action of the scalars is conjugated, we write this as  $z \cdot_c m = \bar{z}m$ . Then the identity gives us an identification

$$\text{End}_{\mathbb{C}}(\mathcal{M} \otimes_F \mathbb{C}) = \text{End}_{\mathbb{C}}(\bar{\mathcal{M}}_\lambda \otimes_F \mathbb{C}).$$

Now we define an action  $\bar{r}_{\mathbb{C}}$  on  $\bar{\mathcal{M}}_\lambda \otimes_F \mathbb{C}$ : For  $g \in G(\mathbb{C})$  we put

$$\bar{r}_{\mathbb{C}}(g)m = r_{\mathbb{C}}(g) \cdot_c m.$$

This defines an action of the abstract group  $G(\mathbb{C})$ , but this is in fact obtained from a rational representation. Therefore  $\mathcal{M}_{\mathbb{C}}^{\vee}$  and  $\overline{\mathcal{M}}_C$  both are given by a highest weight.

The highest weight of  $\mathcal{M}_{\lambda}^{\vee}$  is  $-w_0(\lambda)$ . Here  $w_0$  is the unique element  $w_0 \in W$ , which sends the system of positive roots  $\Delta^+$  into the system  $\Delta^- = -\Delta^+$ .

The highest weight of  $\overline{\mathcal{M}}_{\lambda} \otimes_F \mathbb{C}$  is  $c(\lambda)$  where  $c \in \text{Gal}(\mathbb{C}/\mathbb{R}) \subset \text{Gal}(F/\mathbb{Q})$  is the complex conjugation acting on  $X^*(T \times_{\mathbb{Q}} F)$ . So we may say:  $\overline{\mathcal{M}}_{\lambda C} = \mathcal{M}_{\lambda}$ .

We will call the module  $\mathcal{M}_{\lambda}$ -conjugate-*autodual* or simply *c-autodual* if

$$c(\lambda) = -w_0(\lambda) \tag{8.1}$$

In the following few sections (until 8.1.7 we will always assume that our local system (resp. the corresponding representation) are local systems in  $\mathbb{C}$ -vector spaces (resp.  $\mathbb{C}$ -vector spaces  $\mathcal{M}_{\lambda}$ ). Therefore we will suppress the factor  $\otimes \mathbb{C}$ .

HCmod

### 8.1.2 Harish-Chandra modules and $(\mathfrak{g}, K_{\infty})$ -cohomology.

Now we consider the group of real points  $G(\mathbb{R})$ , it has the Lie algebra  $\mathfrak{g}$ , inside this Lie algebra we have the Lie algebra  $\mathfrak{k}$  of the group  $K_{\infty}$ . We have the notion of a  $(\mathfrak{g}, K_{\infty})$  module: This is a  $\mathbb{C}$ -vector space  $V$  together with an action of  $\mathfrak{g}$  and an action of the group  $K_{\infty}$ . We have certain assumptions of consistency:

i) The action of  $K_{\infty}$  is differentiable, this means it induces an action of  $\mathfrak{k}$ , the derivative of the group action.

ii) The action of  $\mathfrak{g}$  restricted to  $\mathfrak{k}$  is the derivative of the action of  $K_{\infty}$ .

iii) For  $k \in K_{\infty}, X \in \mathfrak{g}$  and  $v \in V$  we have

$$(\text{Ad}(k)X)v = k(X(k^{-1}v)).$$

Inside  $V$  we have the subspace of  $K_{\infty}$  finite vectors, a vector  $v$  is called  $K_{\infty}$  finite if the  $\mathbb{C}$ -subspace generated by all translates  $kv$  is finite dimensional, i.e.  $v$  lies in a finite dimensional  $K_{\infty}$  invariant subspace. The  $K_{\infty}$  finite vectors form a subspace  $V^{(K_{\infty})}$  and it is obvious that  $V^{(K_{\infty})}$  is invariant under the action of  $\mathfrak{g}$ , hence it is a  $(\mathfrak{g}, K_{\infty})$  sub module of  $V$ . We call a  $(\mathfrak{g}, K_{\infty})$  module a Harish-Chandra module if  $V = V^{(K_{\infty})}$ .

For such a  $(\mathfrak{g}, K_{\infty})$ -module we can write down a complex

$$\text{Hom}_{K_{\infty}}(\Lambda^{\bullet}(\mathfrak{g}/\mathfrak{k}), V) = \{0 \rightarrow V \rightarrow \text{Hom}_{K_{\infty}}(\Lambda^1(\mathfrak{g}/\mathfrak{k}), V) \rightarrow \text{Hom}_{K_{\infty}}(\Lambda^2(\mathfrak{g}/\mathfrak{k}), V) \rightarrow \dots\}$$

where the differential is given by liealgc

$$d\omega(X_0, X_1, \dots, X_p) = \sum_{i=0}^p (-1)^i X_i \omega(X_0, \dots, \hat{X}_i, \dots, X_p) + \sum_{0 \leq i < j \leq p} (-1)^{i+j} \omega([X_i, X_j], X_0, \dots, \hat{X}_i, \dots, \hat{X}_j, \dots, \tag{8.2}$$

A few comments are in order. We have inclusions

$$\text{Hom}_{K_{\infty}}(\Lambda^{\bullet}(\mathfrak{g}/\mathfrak{k}), V) \subset \text{Hom}(\Lambda^{\bullet}(\mathfrak{g}/\mathfrak{k}), V) \subset \text{Hom}(\Lambda^{\bullet}(\mathfrak{g}), V).$$

The above differential defines the structure of a complex for the rightmost term, we have to verify that the leftmost term is a subcomplex, this is not so difficult.

We define the  $(\mathfrak{g}, K_\infty)$  cohomology as the cohomology of this complex, i.e.

$$H^\bullet(\mathfrak{g}, K_\infty, V) = H^\bullet(\text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), V)).$$

It is clear that the map

$$H^\bullet(\mathfrak{g}, K_\infty, V^{(K_\infty)}) \rightarrow H^\bullet(\mathfrak{g}, K_\infty, V)$$

is an isomorphism.

If we have two  $(\mathfrak{g}, K_\infty)$  modules  $V_1, V_2$  and form the algebraic tensor product  $W = V_1 \otimes V_2$  then we have a natural structure of a  $(\mathfrak{g}, K_\infty)$  -module on  $W$  : The group  $K_\infty$  acts via the diagonal and  $U \in \mathfrak{g}$  acts by the Leibniz-rule  $U(v_1 \otimes v_2) = Uv_1 \otimes v_2 + v_1 \otimes Uv_2$ . If both modules are Harish-Chandra modules, then the tensor product is also a Harish-Chandra module. Of course any finite dimensional rational representation of the algebraic group also yields a Harish-Chandra module.

deRhamiso

### 8.1.3 The representation theoretic de-Rham isomorphism

For us the  $(\mathfrak{g}, K_\infty)$  module  $\mathcal{C}_\infty(G(\mathbb{Q})\backslash G(\mathbb{A})/K_f)$ , - this is the space of functions which are  $\mathcal{C}_\infty$  in the variable  $g_\infty$ - is one of the most important  $(\mathfrak{g}, K_\infty)$  -modules. We may also consider the limit over smaller and smaller levels  $K_f$  we get the space  $\mathcal{C}_\infty(G(\mathbb{Q})\backslash G(\mathbb{A}))$ , which consists of those functions on  $G(\mathbb{A})$ , which are left invariant under  $G(\mathbb{Q})$ , right invariant under a suitably small open subgroup  $K_f \subset G(\mathbb{A}_f)$  and which are  $\mathcal{C}_\infty$  in the variable  $g_\infty$ . On these functions the group  $G(\mathbb{A})$  acts by translations from the right, since our functions are  $\mathcal{C}_\infty$  we also get an action of the Lie algebra  $\mathfrak{g}$ . Hence this is also a  $(\mathfrak{g}, K_\infty) \times G(\mathbb{A}_f)$ -module.

If we fix the level see that  $\mathcal{C}_\infty(G(\mathbb{Q})\backslash G(\mathbb{A})/K_f)$  is a  $(\mathfrak{g}, K_\infty) \times \mathcal{H}_{K_f}$ , the Hecke algebra acts by convolution. We choose a highest weight module  $\mathcal{M}_\lambda$  and apply the previous considerations to the Harish-Chandra module

$$V = \mathcal{C}_\infty(G(\mathbb{Q})\backslash G(\mathbb{A})/K_f) \otimes \mathcal{M}_\lambda.$$

Notice that we can evaluate an element  $f \in \mathcal{C}_\infty(G(\mathbb{Q})\backslash G(\mathbb{A})/K_f) \otimes \mathcal{M}_\lambda$  in a point  $\underline{g} = (g_\infty, \underline{g}_f)$  and the result  $f(\underline{g}) \in \mathcal{M}_\lambda$ . The Hecke algebra acts via convolution on the first factor.

Let us assume that our compact subgroup  $K_f \subset G(\mathbb{A}_f)$  is neat, i.e. for any  $\underline{g} = (g_\infty, \underline{g}_f) \in G(\mathbb{A})$  we have  $\underline{g}^{-1}(K_\infty \times K_f)\underline{g} \cap G(\mathbb{Q}) = \{e\}$ . In this case we know that  $\tilde{\mathcal{M}}$  is a local system and we can form the de-Rham complex  $\Omega^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$ .

We have an action of the Hecke algebra on this complex and we have the following fundamental fact: Borel

**Proposition 8.1.1.** *We have a canonical isomorphism of complexes*

$$\text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \mathcal{C}_\infty(G(\mathbb{Q})\backslash G(\mathbb{A})/K_f) \otimes \mathcal{M}_\lambda) \xrightarrow{\sim} \Omega^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{C}),$$

*this isomorphism is compatible with the action of the Hecke algebra on both sides*

This is rather clear. We have the projection map

$$q : G(\mathbb{R}) \times G(\mathbb{A}_f) \rightarrow G(\mathbb{R})/K_\infty \times G(\mathbb{A}_f)/K_f = X \times G(\mathbb{A}_f)/K_f$$

let  $x_0 \in X \times G(\mathbb{A}_f)/K_f$  be the image of the identity  $e \in G(\mathbb{R})$ . The differential  $D_q(e)$  maps the Lie algebra  $\mathfrak{g}$  = tangent space of  $G(\mathbb{R})$  at  $e$  to the tangent space  $T_{X,x_0}$  at  $x_0 \times e_f$ . This provides the identification  $T_{X,x_0} \xrightarrow{\sim} \mathfrak{g}/\mathfrak{k}$ .

An element  $\omega \in \text{Hom}_{K_\infty}(\Lambda^p(\mathfrak{g}/\mathfrak{k}), \mathcal{C}_\infty(G(\mathbb{Q}) \backslash G(\mathbb{A})/K_f) \otimes \mathcal{M}_\lambda)$  can be evaluated on a  $p$ -tuple  $(X_0, X_1, \dots, X_{p-1})$  and the result

$$\omega(X_0, X_1, \dots, X_{p-1}) \in \mathcal{C}_\infty(G(\mathbb{Q}) \backslash G(\mathbb{A})/K_f) \otimes \mathcal{M}_\lambda.$$

We want to produce an element  $\tilde{\omega}$  in the de-Rham complex  $\Omega^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$ . Pick a point  $x \times \underline{g}_f \in X \times G(\mathbb{A}_f)/K_f$ , we find an element  $(g_\infty, \underline{g}_f) \in G(\mathbb{R}) \times G(\mathbb{A}_f)$  such that  $g_\infty x_0 = x$ . Our still to be defined form  $\tilde{\omega}$  can be evaluated at a  $p$ -tuple  $(Y_0, \dots, Y_{p-1})$  of tangent vectors in  $x \times \underline{g}_f$  and the result has to be an element in  $\mathcal{M}_{\mathbb{C},x}$ . We find a  $p$ -tuple  $(X_0, X_1, \dots, X_{p-1})$  of tangent vectors at  $x_0$  which are mapped to  $(Y_0, \dots, Y_{p-1})$  under the differential  $D_{g_\infty}$  of the left translation by  $D_{g_\infty}$ . We put Armand

$$\tilde{\omega}(Y_0, \dots, Y_{p-1})(x \times \underline{g}_f) = g_\infty^{-1}(\omega(X_0, \dots, X_{p-1})(g_\infty, \underline{g}_f)). \quad (8.3)$$

At this point I leave it as an exercise to the reader that this gives the isomorphism we want. (Ref ???)

We recall that the de-Rham complex (Reference Book Vol. !) computes the cohomology and therefore we can rewrite the de-Rham isomorphism BodeRh

$$H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \xrightarrow{\sim} H^\bullet(\text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \mathcal{C}_\infty(G(\mathbb{Q}) \backslash G(\mathbb{A})/K_f) \otimes \mathcal{M}_\lambda)) \quad (8.4)$$

From now on the complex  $\text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \mathcal{C}_\infty(G(\mathbb{Q}) \backslash G(\mathbb{A})/K_f) \otimes \mathcal{M}_\lambda)$  will also be called the de-Rham complex.

By the same token we can compute the cohomology with compact supports BodeRhcs

$$H_c^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \xrightarrow{\sim} H_c^\bullet(\text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \mathcal{C}_{c,\infty}(G(\mathbb{Q}) \backslash G(\mathbb{A})/K_f) \otimes \mathcal{M}_\lambda)) \quad (8.5)$$

where  $\mathcal{C}_{c,\infty}(G(\mathbb{Q}) \backslash G(\mathbb{A})/K_f)$  are the  $\mathcal{C}_\infty$  function with compact support. These isomorphisms are also valid if we drop the assumption that  $K_f$  is neat.

The Poincaré duality on the cohomology is induced by the pairing on the de-Rham complexes: PD

**Proposition 8.1.2.** *If  $\omega_1 \in \text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \mathcal{C}_\infty(G(\mathbb{Q}) \backslash G(\mathbb{A})/K_f) \otimes \tilde{\mathcal{M}})$  is a closed form and  $\omega_2 \in \text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \mathcal{C}_{\infty,c}(G(\mathbb{Q}) \backslash G(\mathbb{A})/K_f) \otimes \tilde{\mathcal{M}}^\vee)$  a closed form with compact support in complementary degree then the value of the cup product pairing of the classes  $[\omega_1] \in H^p(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$ ,  $[\omega_2] \in H_c^{d-p}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda^\vee)$  is given by*

$$\langle [\omega_1] \cup [\omega_2] \rangle = \int_{\mathcal{S}_{K_f}^G} \langle \omega_1 \wedge \omega_2 \rangle$$

(Reference Book Vol. !)

### 8.1.4 Input from representation theory of real reductive groups.

Let us consider an arbitrary irreducible  $(\mathfrak{g}, K_\infty)$ -module  $V$ . We also assume that for any  $\vartheta \in \hat{K}_\infty$  the multiplicity of  $\vartheta$  in  $V$  is finite (we say that  $V$  is admissible). Then we can extend the action of the Lie-algebra  $\mathfrak{g}$  to an action of the universal enveloping algebra  $\mathfrak{U}(\mathfrak{g})$  on  $V$  and we can restrict this action to an action of the centre  $\mathfrak{Z}(\mathfrak{g})$ . The structure of this centre is well known by a theorem of Harish-Chandra, it is a polynomial algebra in  $r = \text{rank}(G)$  variables, here the rank is the absolute rank, i.e. the dimension of a maximal torus in  $G/\mathbb{Q}$ . (See Chap. 4 sect. 4)

Clearly this centre respects the decomposition into  $K_\infty$  types, since these  $K_\infty$  types come with finite multiplicity we can apply the standard argument, which proves the Lemma of Schur. Hence  $\mathfrak{Z}(\mathfrak{g})$  has to act on  $V$  by scalars, we get a homomorphism  $\chi_V : \mathfrak{Z}(\mathfrak{g}) \rightarrow \mathbb{C}$ , which is defined by

$$zv = \chi_V(z)v.$$

This homomorphism is called the central character of  $V$ .

A fundamental theorem of Harish-Chandra asserts that for a given central character there exist only finitely many isomorphism classes of irreducible, admissible  $(\mathfrak{g}, K_\infty)$ -modules with this central character.

Of course for any rational finite dimensional representation  $r : G/\mathbb{Q} \rightarrow \text{Gl}(\mathcal{M}_\lambda)$  we can consider  $\mathcal{M}_\lambda \otimes \mathbb{C}$  as  $(\mathfrak{g}, K_\infty)$ -module. If  $\mathcal{M}_\lambda$  is absolutely irreducible with highest weight  $\lambda$  (See chap. IV) then it also has a central character  $\chi_{\mathcal{M}} = \chi_\lambda$ .

**Wigner's lemma:** *Let  $V$  be an irreducible, admissible  $(\mathfrak{g}, K_\infty)$ -module, let  $\mathcal{M} = \mathcal{M}_\lambda$ , a finite dimensional, absolutely irreducible rational representation. Then  $H^\bullet(\mathfrak{g}, K_\infty, V \otimes \mathcal{M}_\mathbb{C}) = 0$  unless we have*

$$\chi_V(z) = \chi_{\mathcal{M}^\vee}(z) = \chi_{\mathcal{M}_{\lambda^\vee}}(z) \text{ for all } z \in \mathfrak{Z}(\mathfrak{g})$$

Since we also know that the number of isomorphism classes of irreducible, admissible  $(\mathfrak{g}, K_\infty)$ -modules with a given central character is finite, we can conclude that for a given absolutely irreducible rational module  $\mathcal{M}_\lambda$  the number of isomorphism classes of irreducible, admissible  $(\mathfrak{g}, K_\infty)$ -modules  $V$  with  $H^\bullet(\mathfrak{g}, K_\infty, V \otimes \mathcal{M}_\mathbb{C}) \neq 0$  is finite.

The proof of Wigner's lemma is very elegant. We have  $\mathcal{M} \otimes V = \mathcal{M}^\vee \otimes V$  and hence we have  $H^0(\mathfrak{g}, K_\infty, \mathcal{M} \otimes V) = \text{Hom}(\mathcal{M}^\vee, V)^{(\mathfrak{g}, K_\infty)} = \text{Hom}_{\mathfrak{g}, K_\infty}(\mathcal{M}^\vee, V)$ . In [B-W], Chap.I 2.4 it is shown, that the category of  $\mathfrak{g}, K_\infty$ -modules has enough injective and projective elements (See [B-W], I. 2.5). If  $I$  is an injective  $\mathfrak{g}, K_\infty$ -module then  $\mathcal{M} \otimes I$  is also injective because for any  $\mathfrak{g}, K_\infty$ -module  $A$  we have  $\text{Hom}(A, \mathcal{M} \otimes I) = \text{Hom}(\mathcal{M}^\vee, I)$ . Hence an injective resolution  $0 \rightarrow V \rightarrow I^0 \rightarrow I^1 \dots$  yields an injective resolution  $0 \rightarrow \mathcal{M} \rightarrow \mathcal{M} \otimes I^0 \rightarrow \mathcal{M} \otimes I^1 \dots$  and from this we get

$$H^q(\mathfrak{g}, K_\infty, \mathcal{M} \otimes V) = \text{Ext}_{\mathfrak{g}, K_\infty}^q(\mathcal{M}^\vee, V).$$

Any  $z \in \mathfrak{Z}(\mathfrak{g})$  induces an endomorphism of  $\mathcal{M}_\lambda$  and  $V$ . Since  $\text{Ext}^\bullet$  is functorial in both variables, we see that  $z$  induces endomorphisms  $z_1$  (via the action on  $\mathcal{M}_\lambda$ ) and  $z_2$  (via the action on  $V$ ) on  $\text{Ext}_{\mathfrak{g}, K_\infty}^q(\mathcal{M}^\vee, V)$ . We show that  $z_1 = z_2$ . This is clear by definition for  $\text{Ext}_{\mathfrak{g}, K_\infty}^0(\mathcal{M}^\vee, V) = \text{Hom}_{\mathfrak{g}, K_\infty}(\mathcal{M}^\vee, V)$ : For  $z \in \mathfrak{Z}(\mathfrak{g})$  and  $\phi \in \text{Hom}_{\mathfrak{g}, K_\infty}(\mathcal{M}^\vee, V)$ ,  $m \in \mathcal{M}_\lambda$  we have  $z_1\phi(m) = \phi(zm) = z_2(\phi(m))$ . To prove it for an arbitrary  $q$  we use devissage and induction. We embed  $V$  into an injective  $\mathfrak{g}, K_\infty$  module  $I$  and get an exact sequence

$$0 \rightarrow V \rightarrow I \rightarrow I/V \rightarrow 0$$

and from this and  $\text{Ext}_{\mathfrak{g}, K_\infty}^q(\mathcal{M}_\lambda, I)$  for  $q > 0$  we get

$$\text{Ext}^{q-1}(\mathfrak{g}, K_\infty, \mathcal{M}_\lambda, I/V) = \text{Ext}^q(\mathfrak{g}, K_\infty, \mathcal{M}_\lambda, V) \text{ for } q > 0.$$

Now by induction we know  $z_1 = z_2$  on the left hand side, so it also holds on the right hand side.

If now  $\chi_V \neq \chi_{\mathcal{M}^\vee}$  then we can find a  $z \in \mathfrak{Z}(\mathfrak{g})$  such that  $\chi_{\mathcal{M}^\vee}(z) = 0, \chi_V(z) = 1$ . This implies that  $z_1 = 0$  and  $z_2 = 1$  on all  $\text{Ext}^q(\mathfrak{g}, K_\infty(\mathcal{M}_\lambda, V))$ . Since we know that  $z_1 = z_2$  we see that the identity on  $\text{Ext}^q(\mathfrak{g}, K_\infty(\mathcal{M}_\lambda, V))$  is equal to zero and this implies the assertion.

On the universal enveloping algebra  $\mathfrak{U}(\mathfrak{g})$  we have an antiautomorphism  $u \mapsto {}^t u$  which is induced by the antiautomorphism  $X \mapsto -X$  on the Lie algebra  $\mathfrak{g}$ . If  $V$  is an admissible  $(\mathfrak{g}, K_\infty)$ -module, then we can form the dual module  $V^\vee$  and if we denote the pairing between  $V, V^\vee$  by  $\langle \cdot, \cdot \rangle_V$  then

$$\langle Uv, \phi \rangle_V = \langle v, {}^t U\phi \rangle_V \text{ for all } U \in \mathfrak{U}(\mathfrak{g}), v \in V, \phi \in V^\vee.$$

If  $V$  is irreducible, then it has a central character and we get

$$\chi_{V^\vee}(z) = \chi_V({}^t z).$$

This applies to finite dimensional and infinite dimensional  $(\mathfrak{g}, K_\infty)$ -modules.

### 8.1.5 Representation theoretic Hodge-theory.

We consider irreducible unitary representations  $G(\mathbb{R}) \rightarrow U(H)$ . We know from the work of Harish-Chandra:

1) If we fix an isomorphism class  $\vartheta$  irreducible representations of  $K_\infty$  then the isotypical subspace  $\dim_{\mathbb{C}} H(\vartheta) \leq \dim(\vartheta)^2$ , i.e.  $\vartheta$  occurs at most with multiplicity  $\dim(\vartheta)$ .

2) The direct sum  $\sum_{\vartheta \subset K_\infty} H(\vartheta) = H^{(K)} \subset H$  is dense in  $H$  and it is an admissible irreducible Harish-Chandra -module.

We call an irreducible  $(\mathfrak{g}, K_\infty)$ -module unitary, if it is isomorphic to such an  $H^{(K)}$ .

For a given  $G/\mathbb{R}$  and any rational irreducible module  $\mathcal{M}_\lambda$  Vogan and Zuckerman give a finite list of certain irreducible, admissible  $(\mathfrak{g}, K_\infty)$ -modules  $A_q(\lambda)$ , for which  $H^\bullet(\mathfrak{g}, K_\infty, A_q(\lambda) \otimes \mathcal{M}_\lambda) \neq 0$  they compute these cohomology group. This list contains all unitary, irreducible  $(\mathfrak{g}, K_\infty)$ -modules, which have non trivial cohomology with coefficients in  $\mathcal{M}_\lambda$ .

For the following we refer to [B-W] Chap. II ,§ 1-2 . We want to apply the methods of Hodge-theory to compute the cohomology groups  $H^\bullet(\mathfrak{g}, K_\infty, V \otimes$

$\mathcal{M}_\lambda$ ) for an unitary  $(\mathfrak{g}, K_\infty)$ -module  $V$ . This means have a positive definite scalar product  $\langle \cdot, \cdot \rangle_V$  on  $V$ , for which the action of  $K_\infty$  is unitary and for  $U \in \mathfrak{g}$  and  $v_1, v_2 \in V$  we have  $\langle Uv_1, v_2 \rangle_V + \langle v_1, Uv_2 \rangle_V = 0$ .

In the next step we introduce for all  $p$  a hermitian form on  $\text{Hom}_{K_\infty}(\Lambda^p(\mathfrak{g}/\mathfrak{k}), V \otimes \mathcal{M}_\lambda)$ . To do this we construct a hermitian form on  $\mathcal{M}_\lambda$ .

(The following considerations are only true modulo the centre). We consider the Lie algebra and its complexification  $\mathfrak{g}_\mathbb{C} = \mathfrak{g} \otimes \mathbb{C}$ . On this complex vector space we have the complex conjugation  $\bar{\cdot} : U \mapsto \bar{U}$ . We rediscover  $\mathfrak{g}$  as the set of fixed points under  $\bar{\cdot}$ . We also have the Cartan involution  $\Theta$  which is the involution which has  $\mathfrak{k}$  as its fixed point set. Then we get the Cartan decomposition

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p} \text{ where } \mathfrak{p} \text{ is the } -1 \text{ eigenspace of } \Theta.$$

The Killing form is negative definite on  $\mathfrak{k}$  and positive definite on  $\mathfrak{p}$ , we have for the Lie bracket  $[\mathfrak{p}, \mathfrak{p}] \subset \mathfrak{k}$ . We consider the invariants under  $\bar{\cdot} \circ \Theta$ , this is the Lie algebra  $\mathfrak{g}_c = \mathfrak{k} \oplus \sqrt{-1} \otimes \mathfrak{p}$ . On this real Lie algebra the Killing form is negative definite and  $\mathfrak{g}_c$  is the Lie algebra of an algebraic group  $G_c/\mathbb{R}$  whose base extension  $G_c \otimes_{\mathbb{R}} \mathbb{C} \xrightarrow{\sim} G \otimes_{\mathbb{R}} \mathbb{C}$  and whose group  $G_c(\mathbb{R})$  of real points is compact (this is the so called compact form of  $G$ ). We still have the representation  $G_c/\mathbb{R} \rightarrow \text{Gl}(\mathcal{M}_\lambda)$  which is irreducible and hence we find a hermitian form  $\langle \cdot, \cdot \rangle_\lambda$  on  $\mathcal{M}_\lambda$ , which is invariant under  $G_c(\mathbb{R})$  and which is unique up to a scalar.

This form satisfies the equations

$$\langle Um_1, m_2 \rangle_{\mathcal{M}} + \langle m_1, Um_2 \rangle_\lambda = 0 \text{ for all } m_1, m_2 \in \mathcal{M}_\lambda, U \in \mathfrak{k}$$

this is the invariance under  $K_\infty$  and

$$\langle Um_1, m_2 \rangle_{\mathcal{M}} = \langle m_1, Um_2 \rangle_\lambda \text{ for all } m_1, m_2 \in \mathcal{M}_\lambda, U \in \mathfrak{p}$$

this is the invariance under  $\sqrt{-1} \otimes \mathfrak{p}$ .

Now we define a hermitian metric on  $V \otimes \mathcal{M}_\lambda$ , we simply take the tensor product  $\langle \cdot, \cdot \rangle_V \otimes \langle \cdot, \cdot \rangle_\lambda = \langle \cdot, \cdot \rangle_{V \otimes \lambda}$ . Finally we define the (hermitian) scalar product on  $\text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), V \otimes \mathcal{M}_\lambda)$ . We choose an orthonormal (with respect to the Killing form) basis  $E_1, E_2, \dots, E_d$  on  $\mathfrak{p}$ , we identify  $\mathfrak{g}/\mathfrak{k} \xrightarrow{\sim} \mathfrak{p}$ . Then a form  $\omega \in \text{Hom}_{K_\infty}(\Lambda^p(\mathfrak{g}/\mathfrak{k}), V \otimes \mathcal{M}_\lambda)$  is given by its values  $\omega(E_I) \in V \otimes \mathcal{M}_\lambda$ , where  $I = \{i_1, i_2, \dots, i_p\}$  runs through the ordered subsets of  $\{1, 2, \dots, d\}$  with  $p$  elements. For  $\omega_1, \omega_2 \in \text{Hom}_{K_\infty}(\Lambda^p(\mathfrak{g}/\mathfrak{k}), V \otimes \mathcal{M}_\lambda)$  we put

$$\langle \omega_1, \omega_2 \rangle = \sum_{I, |I|=p} \langle \omega_1(E_I), \omega_2(E_I) \rangle_{V \otimes \lambda} \tag{8.6}$$

Now we can define an adjoint operator

$$\delta : \text{Hom}_{K_\infty}(\Lambda^p(\mathfrak{g}/\mathfrak{k}), V \otimes \mathcal{M}_\lambda) \rightarrow \text{Hom}_{K_\infty}(\Lambda^{p-1}(\mathfrak{g}/\mathfrak{k}), V \otimes \mathcal{M}_\lambda), \tag{8.7}$$

which can be defined by a straightforward calculation. We simply write a formula for  $\delta$ : For an element  $E_i$  we define  $E_i^*(v \otimes m) = -E_i v \otimes m + v \otimes E_i m$ . Then we can define  $\delta$  by the following formula:

We have to evaluate  $\delta(\omega)$  on  $E_J = (E_{i_1}, \dots, E_{i_{p-1}})$  where  $J = \{i_1, \dots, i_{p-1}\}$ . We put

$$\delta(\omega)(E_J) = \sum_{i \notin J} (-1)^{p(i, J \cup \{i\})} E_i^* \omega_{J \cup \{i\}},$$

where  $p(i, J \cup \{i\})$  denotes the position of  $i$  in the ordered set  $J \cup \{i\}$ . With this definition we get for a pair of forms  $\omega_1 \in \text{Hom}_{K_\infty}(\Lambda^{p-1}(\mathfrak{g}/\mathfrak{k}), V \otimes \mathcal{M}_\lambda)$  and  $\omega_2 \in \text{Hom}_{K_\infty}(\Lambda^p(\mathfrak{g}/\mathfrak{k}), V \otimes \mathcal{M}_\lambda)$  (See [B-W], II, prop. 2.3)

$$\langle d\omega_1, \omega_2 \rangle = \langle \omega_1, \delta\omega_2 \rangle \tag{8.8}$$

We define the Laplacian  $\Delta = \delta d + d\delta$ . Then we have ([B-W], II, Thm.2.5)

$$\langle \Delta\omega, \omega \rangle \geq 0 \text{ and we have equality if and only if } d\omega = 0, \delta\omega = 0 \tag{8.9}$$

Inside  $\mathfrak{Z}(\mathfrak{g})$  we have the the Casimir operator  $C$  (See Chap. 4). An element  $z \in \mathfrak{Z}(\mathfrak{g})$  acts on  $V \otimes \mathcal{M}_\lambda$  by  $z \otimes \text{Id}$  via the action on the first factor and by the scalar  $\chi_\lambda(z)$  via the action on the second factor. Then we have

**Kuga's lemma :** *The action of the Casimir operator and the Laplace operator on  $\text{Hom}_{K_\infty}(\Lambda^p(\mathfrak{g}/\mathfrak{k}), V \otimes \mathcal{M}_\lambda)$  are related by the identity*

$$\Delta = C \otimes \text{Id} - \chi_\lambda(C).$$

*If the  $(\mathfrak{g}, K_\infty)$  module is irreducible, then  $\Delta$  acts by multiplication by the scalar  $\chi_V(C) - \chi_\lambda(C)$*

This has the following consequence

*If  $V$  is an irreducible unitary  $\mathfrak{g}, K_\infty$ - module and if  $\mathcal{M}_\lambda$  is an irreducible representation with highest weight  $\lambda$  then*

$$H^\bullet(\mathfrak{g}, K_\infty, V \otimes \mathcal{M}_\mathbb{C}) = \begin{cases} 0 & \text{if } \chi_V(C) - \chi_\lambda(C) \neq 0 \\ \text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), V \otimes \mathcal{M}_\lambda) & \text{if } \chi_V(C) - \chi_\lambda(C) = 0 \end{cases}.$$

This only applies for unitary  $\mathfrak{g}, K_\infty$ -modules, but for these it is much stronger: It says that under the assumption  $\chi_V(C) = \chi_\lambda(C)$  we have  $\chi_V = \chi_\lambda$  ( we only have to test the Casimir operator) and it says that all the differentials in the complex are zero.

### 8.1.6 Input from the theory of automorphic forms

We apply this to the spaces of square integrable functions on  $G(\mathbb{Q}) \backslash G(\mathbb{A}) / K_f$ . Because of the presence of a non trivial center, we have to consider functions which transform in a certain way under the action of the center. We may assume that coefficient system  $\mathcal{M}_\lambda$  has a central character and this central character defines a character  $\zeta_\lambda$  on the maximal  $\mathbb{Q}$ -split torus  $S \subset C$ . This character can be evaluated on the connected component of the identity of the real valued points and induces a (continuous) homomorphism  $\zeta_\infty : S^0(\mathbb{R}) \rightarrow \mathbb{R}_{>0}^\times$ . Then we define

$$\mathcal{C}_\infty(G(\mathbb{Q}) \backslash G(\mathbb{A}) / K_f, \zeta_\infty^{-1}) \tag{8.10}$$

to be the subspace of those  $\mathcal{C}_\infty$  functions which satisfy  $f(z_\infty g) = \zeta_\infty^{-1}(z_\infty)f(g)$  for all  $z_\infty \in S^0(\mathbb{R}) \backslash \underline{G}(\mathbb{A})$ . The isogeny  $d_C : C \rightarrow C'$  (see 6.1.1) induces an isomorphism  $S^0(\mathbb{R}) \xrightarrow{\sim} S'^0(\mathbb{R})$ , where  $S'$  is the maximal  $\mathbb{Q}$  split torus in  $C'$ . Therefore we get a character  $\zeta'_\infty : S'^0(\mathbb{R}) \rightarrow \mathbb{R}_{>0}^\times$  and this is also a character  $\zeta'_\infty : G(\mathbb{R}) \rightarrow \mathbb{R}_{>0}^\times$  and its restriction to  $S^0(\mathbb{R})$  is  $\zeta_\infty$ . If now  $f \in \mathcal{C}_\infty(G(\mathbb{Q}) \backslash G(\mathbb{A}) / K_f, \zeta_\infty^{-1})$  then

$$f(g)\zeta'_\infty(g) \in \mathcal{C}_\infty(G(\mathbb{Q})S^0(\mathbb{R}) \backslash G(\mathbb{A}) / K_f) \tag{8.11}$$

We say that  $f \in \mathcal{C}_\infty(G(\mathbb{Q}) \backslash G(\mathbb{A}) / K_f, \zeta_\infty^{-1})$  is square integrable if

$$\int_{(G(\mathbb{Q})S^0(\mathbb{R}) \backslash G(\mathbb{A}) / K_f)} |f(g)\zeta'_\infty(g)|^2 dg < \infty \tag{8.12}$$

and this allows us to define the Hilbert space  $L^2(G(\mathbb{Q}) \backslash G(\mathbb{A}) / K_f, \zeta_\infty^{-1})$ . Since the space  $(G(\mathbb{Q})S^0(\mathbb{R}) \backslash G(\mathbb{A}) / K_f)$  has finite volume we know that

$$\zeta'_\infty \in L^2(G(\mathbb{Q}) \backslash G(\mathbb{A}) / K_f, \zeta_\infty^{-1}).$$

The group  $G(\mathbb{R})$  acts on  $\mathcal{C}_\infty(G(\mathbb{Q}) \backslash G(\mathbb{A}) / K_f, \zeta_\infty^{-1})$  by right translations and hence we get by differentiating an action of the universal enveloping algebra  $\mathfrak{U}(\mathfrak{g})$  on it. We define by  $\mathcal{C}_\infty^{(2)}(G(\mathbb{Q}) \backslash G(\mathbb{A}) / K_f, \zeta_\infty^{-1})$  the subspace of functions  $f$  for which  $Uf$  is square integrable for all  $U \in \mathfrak{U}(\mathfrak{g})$ .

This allows us to define a sub complex of the de-Rham complex Ltwo

$$\text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \mathcal{C}_\infty^{(2)}(G(\mathbb{Q}) \backslash G(\mathbb{A}) / K_f, \zeta_\infty^{-1}) \otimes \mathcal{M}_\lambda). \tag{8.13}$$

We will not work with this complex because its cohomology may show some bad behavior. (See remark below).

We do something less sophisticated, we simply define  $H_{(2)}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \subset H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$  to be the image of the cohomology of the complex (8.13) in the cohomology. Hence  $H_{(2)}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$  is the space of cohomology classes which can be represented by square integrable forms.

Remark: Some authors also define  $L^2$  de-Rham complexes, using the above complex (8.13) and then they take suitable completions to get complexes of Hilbert spaces. These complexes also give cohomology groups which run under the name of  $L^2$ -cohomology. These  $L^2$ -cohomology groups are related but not necessarily equal to our  $H_{(2)}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$ . They can be infinite dimensional.

The Hilbert space  $L^2(G(\mathbb{Q}) \backslash G(\mathbb{A}) / K_f, \zeta_\infty^{-1})$  is a module for  $G(\mathbb{R}) \times \mathcal{H}_{K_f}$  the group  $G(\mathbb{R})$  acts by unitary transformations and the algebra  $\mathcal{H}_{K_f}$  is selfadjoint.

Let us assume that  $H = H_{\pi_\infty \times \pi_f}$  is an irreducible unitary module for  $G(\mathbb{R}) \times \mathcal{H} = \bigotimes'_p \mathcal{H}_p$  and assume that we have an inclusion of this  $G(\mathbb{R}) \times \mathcal{H}$ -module

$$j : H \hookrightarrow L^2(G(\mathbb{Q}) \backslash G(\mathbb{A}) / K_f, \zeta_\infty^{-1}).$$

It follows from the finiteness results in 8.1.5 that induces an inclusion into the space of square integrable  $\mathcal{C}_\infty$  functions

$$H^{(K_\infty)} \hookrightarrow \mathcal{C}_\infty^{(2)}(G(\mathbb{Q}) \backslash G(\mathbb{A}) / K_f, \zeta_\infty^{-1})^{(K_\infty)}.$$

We consider the  $(\mathfrak{g}, K_\infty)$ -cohomology of this module with coefficients in our irreducible module  $\mathcal{M}_\lambda$ , we assume  $\chi_V(C) = \chi_\lambda(C)$ . We have  $H^\bullet(\mathfrak{g}, K_\infty, H \otimes \mathcal{M}_\lambda) = \text{Hom}_{K_\infty}(\mathfrak{g}, K_\infty, H^{(K_\infty)} \otimes \mathcal{M}_\lambda)$  and get

$$H^\bullet(\mathfrak{g}, K_\infty, H^{(K_\infty)} \otimes \mathcal{M}_\mathbb{C}) \xrightarrow{j^\bullet} H^\bullet(\mathfrak{g}, K_\infty, \mathcal{C}_\infty(G(\mathbb{Q}) \backslash G(\mathbb{A})/K_f, \zeta_\infty^{-1})^{(K_\infty)} \otimes \mathcal{M}_\lambda).$$

This suggests that we try to "decompose"  $\mathcal{C}_\infty(G(\mathbb{Q}) \backslash G(\mathbb{A})/K_f, \zeta_\infty^{-1})^{(K_\infty)}$  into irreducibles and then investigate the contributions of the irreducible summands to the cohomology. Essentially we follow the strategy of [Bo-Ga] and [Bo-Ca] but instead of working with complexes of Hilbert spaces we work with complexes of  $\mathcal{C}_\infty$  forms and modify the arguments accordingly.

It has been shown by Langlands, that we have a decomposition into a discrete and a continuous spectrum

$$L^2(G(\mathbb{Q}) \backslash G(\mathbb{A})/K_f) = L^2_{\text{disc}}(G(\mathbb{Q}) \backslash G(\mathbb{A}_f)/K_f) \oplus L^2_{\text{cont}}(G(\mathbb{Q}) \backslash G(\mathbb{A}_f)/K_f),$$

where  $L^2_{\text{disc}}(G(\mathbb{Q}) \backslash G(\mathbb{A}_f)/K_f)$  is the closure of the sum of all irreducible closed subspaces occurring in  $L^2(G(\mathbb{Q}) \backslash G(\mathbb{A})/K_f)$  and where  $L^2_{\text{cont}}(G(\mathbb{Q}) \backslash G(\mathbb{A}_f)/K_f)$  is the complement.

The discrete spectrum  $L^2_{\text{disc}}(G(\mathbb{Q}) \backslash G(\mathbb{A}_f)/K_f)$  contains as a subspace the *cuspidal spectrum*  $L^2_{\text{cusp}}(G(\mathbb{Q}) \backslash G(\mathbb{A}_f)/K_f)$  :

A function  $f \in L^2(G(\mathbb{Q}) \backslash G(\mathbb{A}_f)/K_f)$  is called a *cuspidal form* if for all proper parabolic subgroups  $P/\mathbb{Q} \subset G/\mathbb{Q}$ , with unipotent radical  $U_P/\mathbb{Q}$  the integral

$$\mathcal{F}^P(f)(g) = \int_{U_P(\mathbb{Q}) \backslash U_P(\mathbb{A})} f(\underline{u}g) d\underline{u} = 0,$$

this means that the integral is defined for almost all  $\underline{g}$  and zero for almost all  $\underline{g}$ . The function  $\mathcal{F}^P(f)(\underline{g})$ , which is an almost everywhere defined function on  $P(\mathbb{Q}) \backslash G(\mathbb{A})/K_f$  is called the constant Fourier coefficient of  $f$  along  $P/\mathbb{Q}$ . The cuspidal spectrum is the intersection of all the kernels of the  $\mathcal{F}^P$ .

If our group is anisotropic, then it does not have any proper parabolic subgroup and in this case we have  $L^2_{\text{cusp}}(G(\mathbb{Q}) \backslash G(\mathbb{A}_f)/K_f) = L^2_{\text{disc}}(G(\mathbb{Q}) \backslash G(\mathbb{A}_f)/K_f) = L^2(G(\mathbb{Q}) \backslash G(\mathbb{A}_f)/K_f)$ .

For any unitary  $G(\mathbb{R}) \times \mathcal{H}$ -module  $H_\pi = H_{\pi_\infty} \otimes H_{\pi_f}$  we put  $W_{\pi, \text{cusp}} = \text{Hom}_{G(\mathbb{R}) \times \mathcal{H}}(H_\pi, L^2_{\text{cusp}}(G(\mathbb{Q}) \backslash G(\mathbb{A}_f)/K_f))$ . We can ignore the  $\mathcal{H}$ -module structure and define

$$W_{\pi_\infty, \text{cusp}} = \text{Hom}_{G(\mathbb{R})}(H_{\pi_\infty} \otimes H_{\pi_f}, L^2_{\text{cusp}}(G(\mathbb{Q}) \backslash G(\mathbb{A}_f)/K_f)).$$

It has been shown by Gelfand-Graev and Langlands that

$$m_{\text{cusp}}(\pi_\infty) = \sum_{\pi_f} \dim(W_{\pi_\infty, \text{cusp}}) < \infty.$$

We get a decomposition into isotypical subspaces

$$L^2_{\text{cusp}}(G(\mathbb{Q}) \backslash G(\mathbb{A}_f)/K_f) = \overline{\bigoplus_{\pi_\infty \otimes \pi_f} (L^2_{\text{cusp}}(G(\mathbb{Q}) \backslash G(\mathbb{A}_f)/K_f)(\pi_\infty \times \pi_f))},$$

where  $L^2_{\text{cusp}}(G(\mathbb{Q})\backslash G(\mathbb{A}_f)/K_f)(\pi_\infty \times \pi_f)$  is the image of  $W_{\pi, \text{cusp}} \otimes H_\pi$  in  $L^2_{\text{cusp}}(G(\mathbb{Q})\backslash G(\mathbb{A}_f)/K_f)$ .

The cuspidal spectrum has a complement in the discrete spectrum, this is the *residual spectrum*  $L^2_{\text{res}}((G(\mathbb{Q})\backslash G(\mathbb{A}_f)/K_f)$ . It is called residual spectrum, because the irreducible subspaces contained in it are obtained by residues of Eisenstein classes.

Again we define  $W_{\pi, \text{res}} = \text{Hom}_{G(\mathbb{R}) \times \mathcal{H}}(H_\pi, L^2_{\text{res}}(G(\mathbb{Q})\backslash G(\mathbb{A}_f)/K_f))$ , (resp.  $W_{\pi_\infty, \text{res}} = \text{Hom}_{G(\mathbb{R})}(H_{\pi_\infty}, L^2_{\text{cusp}}(G(\mathbb{Q})\backslash G(\mathbb{A}_f)/K_f))$ , and it is a deep theorem of Langlands that  $m_{\text{res}}(\pi_\infty) = \dim(W_{\pi_\infty, \text{res}}) < \infty$ . Hence we get a decomposition

$$L^2_{\text{res}}(G(\mathbb{Q})\backslash G(\mathbb{A}_f)/K_f) = \overline{\bigoplus_{\pi_\infty \otimes \pi_f} (L^2_{\text{res}}(G(\mathbb{Q})\backslash G(\mathbb{A}_f)/K_f)(\pi_\infty \times \pi_f))}.$$

If our group  $G/\mathbb{Q}$  is isotropic, then the one dimensional space of constants is in the residual (discrete) spectrum but not in the cuspidal spectrum.

Langlands has given a description of the continuous spectrum using the theory of Eisenstein series, we have a decomposition decomp-cont

$$L^2_{\text{cont}}(G(\mathbb{Q})\backslash G(\mathbb{A}_f)/K_f) = \overline{\bigoplus_{\Sigma} \tilde{H}_P^+(\pi_\Sigma)}, \tag{8.14}$$

we briefly explain this decomposition following [Bo-Ga]. The  $\Sigma$  are so called cuspidal data, this are pairs  $(P, \pi_\Sigma)$  where  $P$  is a proper parabolic subgroup and  $\pi_\Sigma$  is a representation of  $M(\mathbb{A}) = P(\mathbb{A})/U(\mathbb{A})$  occurring in the discrete spectrum  $L^2_{\text{cusp}}(M(\mathbb{Q})\backslash M(\mathbb{A}))$ .

Let  $M^{(1)}/\mathbb{Q}$  be the semi simple part of  $M$  and recall that  $C/\mathbb{Q}$  was the center of  $G/\mathbb{Q}$ . We consider the character module  $Y^*(P) = \text{Hom}(C \cdot M^{(1)}, \mathbb{G}_m)$ . The elements  $Y^*(P) \otimes \mathbb{C}$  provide homomorphisms  $\gamma \otimes z : M(\mathbb{A})/C(\mathbb{A})M^{(1)}(\mathbb{A}) \rightarrow \mathbb{C}^\times$ . (See (6.14)). The module  $Y^*(P) \otimes \mathbb{Q}$  comes with a canonical basis which is given by the dominant fundamental weights  $\gamma_\mu$  which are trivial on  $M^{(1)}$ . We define

$$\Lambda_\Sigma = Y^*(P) \otimes i\mathbb{R} = \left\{ \sum_{\mu} \gamma_\mu \otimes it_\mu \mid t_\mu \in \mathbb{R} \right\}$$

this is a group of unitary characters. For  $\sigma \in \Lambda_\Sigma$  we define the unitarily induced representation

$$\text{Ind}_{P(\mathbb{A})}^{G(\mathbb{A})} \pi_\Sigma \otimes (\sigma + \rho_P) = I_P^G \pi_\Sigma \otimes \sigma \tag{8.15}$$

$$\{f : G(\mathbb{A}) \rightarrow L^2_{\text{res}}(M(\mathbb{Q})\backslash M(\mathbb{A}))(\pi_\Sigma) \mid f(\underline{p}g) = (\sigma + |\rho_P|)(\underline{p})\pi_\Sigma(\underline{p})f(\underline{g})\}$$

where of course  $\underline{p} \in P(\mathbb{A}), \underline{g} \in G(\mathbb{A})$  and  $\rho_P \in Y^*(P) \otimes \mathbb{Q}$  is the half sum of the roots in the unipotent radical of  $P$ . This gives us a unitary representation of  $G(\mathbb{A})$ . Let  $d_\Sigma$  be the Lebesgue measure on  $\Lambda_\Sigma$  then we can form the direct integral unitary representations

$$H_P(\pi_\Sigma) = \int_{\Lambda_\Sigma} I_P^G \pi_\Sigma \otimes \sigma \, d_\Sigma \sigma \tag{8.16}$$

The theory of Eisenstein series gives us a homomorphism of  $G(\mathbb{R}) \times \mathcal{H}$ -modules

$$\text{Eis}_P(\pi_\Sigma) : H_P(\pi_\Sigma) \rightarrow L_{\text{cont}}^2(G(\mathbb{Q}) \backslash G(\mathbb{A}_f) / K_f). \quad (8.17)$$

Let us put

$$\Lambda_\Sigma^\pm = \left\{ \sum_{\mu} \gamma_{\mu} \otimes it_{\mu} \mid t_{\mu} \geq 0 \right\}$$

then the restriction

$$\text{Eis}_P(\pi_\Sigma) : H_P^\pm(\pi_\Sigma) = \int_{\Lambda_\Sigma^\pm} I_P^G \pi_\Sigma \otimes \sigma \, d_\Sigma \sigma \rightarrow L_{\text{cont}}^2(G(\mathbb{Q}) \backslash G(\mathbb{A}_f) / K_f). \quad (8.18)$$

is an isometric embedding. The image will be denoted by  $\tilde{H}_P^\pm(\pi_\Sigma)$  these spaces are the elementary subspaces in [B-G]. Two such elementary subspaces  $\tilde{H}_P^\pm(\pi_\Sigma)$ ,  $\tilde{H}_{P_1}^\pm(\pi_{\Sigma_1})$  are either orthogonal to each other or they are equal. We get the above decomposition if we sum over a suitable set of representatives of cuspidal data.

Now we are ready to discuss the contribution of the continuous spectrum to the cohomology. If we have a closed square integrable form

$$\omega \in \text{Hom}_{K_\infty}(\Lambda^p(\mathfrak{g}/\mathfrak{k}), \mathcal{C}_\infty^2(G(\mathbb{Q}) \backslash G(\mathbb{A}) / K_f) \otimes \mathcal{M}_\lambda),$$

then we can decompose it

$$\omega = \omega_{\text{res}} + \omega_{\text{cont}},$$

both summands are  $\mathcal{C}_\infty^2$  and closed.

**Proposition 8.1.3.** *The cohomology class  $[\omega_{\text{cont}}]$  is trivial.*

*Proof.* This now the standard argument in Hodge theory, but this time we apply it to a continuous spectrum instead of a discrete one. We follow Borel-Casselman and prove their Lemma 5.5 (See[B-C]) in our context. We may assume that  $\omega_\infty$  lies in one of the summands, i.e.  $\omega_{\text{cont}} = \text{Eis}(\int_{\Lambda_\Sigma} \omega^\vee(\sigma) d_\Sigma \sigma)$  where  $\omega^\vee(\sigma) \in \text{Hom}_{K_\infty}(\Lambda^p(\mathfrak{g}/\mathfrak{k}), I_P^G \pi_\Sigma \otimes \sigma \otimes \mathcal{M}_\lambda)$  is the Fourier transform of  $\omega_\infty$  in the  $L^2$ , (theorem of Plancherel). As it stands the expression  $\int_{\Lambda_\Sigma} \omega^\vee(\sigma) d_\Sigma \sigma$  does not make sense because the integrand is in  $L^2$  and not necessarily in  $L^1$ . If we choose a symmetric positive definite quadratic form  $h(\sigma) = \sum_{\nu, \mu} b_{\nu, \mu} t_\nu t_\mu$  and a positive real number  $\tau$  then the function

$$h_\tau(\sigma) = (1 + \tau h(\sigma))^m \in L^2(\Lambda_\Sigma)$$

and then  $\omega^\vee(\sigma) h_\tau(\sigma)$  is in  $L^1$  and by definition

$$\lim_{\tau \rightarrow 0} \int_{\Lambda_\Sigma} \omega^\vee(\sigma) h_\tau(\sigma) d_\Sigma \sigma = \int_{\Lambda_\Sigma} \omega^\vee(\sigma) d_\Sigma \sigma \quad (8.19)$$

where the convergence is in the  $L^2$  sense. Since  $\omega_\infty \in \text{Hom}_{K_\infty}(\Lambda^p(\mathfrak{g}/\mathfrak{k}), I_P^G \pi_\Sigma \otimes \sigma \otimes \mathcal{M}_\lambda)$  we get that  $\omega^\vee(\sigma)$  has the following property

For any polynomial  $P(\sigma) = \sum a_\mu t^\mu$  in the variables  $t_\mu$  and with real coefficients the section diffmult

$$\omega^\vee(\sigma) P(\sigma) \text{ is square integrable} \quad (8.20)$$

this follows from the well known rules that differentiating a function provides multiplication by the variables for the Fourier transform.

The Lemma of Kuga implies

$$\Delta(\omega^\vee(\sigma)) = (\chi_\sigma(C) - \chi_\lambda(C))\omega^\vee(\sigma)$$

and if  $\sigma = \sum \gamma_\mu \otimes it_\mu$  the eigenvalue is

$$\chi_\sigma(C) - \chi_\lambda(C) = \sum a_{\nu,\mu} t_\nu t_\mu + \sum b_\mu t_\mu + c_{\pi_\Sigma} - c_\lambda. \tag{8.21}$$

where  $c_{\pi_\Sigma}$  is the eigenvalue of the Casimir operator of  $M^{(1)}$  on  $\pi_\Sigma$  If the  $t_\mu \in \mathbb{R}$  then this expression is always  $\leq 0$  especially we see that the quadratic form on the right hand side is negative definite. This implies that for  $\sigma \in \Lambda_F$  the expression  $\chi_\sigma(C) - \chi_\lambda(C)$  assumes a finite number of maximal values all of them  $\leq 0$  and hence

$$V_\Sigma = \{\sigma | \chi_\sigma(C) - \chi_\lambda(C) = 0\} \tag{8.22}$$

is a finite set of point. This set has measure zero, since we assumed that  $P$  was a proper parabolic subgroup. The of  $\sigma$  for which  $H^\bullet(\mathfrak{g}, K_\infty, H_{\Lambda_\Sigma}(\sigma) \otimes \mathcal{M}_\mathbb{C}) \neq 0$  is finite. We choose a  $\mathcal{C}_\infty$  function  $h_\Sigma(\sigma)$  which is positive, which takes value 1 in a small neighborhood of  $V_\Sigma$ , which takes values  $\leq 1$  in a slightly larger neighborhood and which is zero outside this second neighborhood. Then we write

$$\omega_\infty = \text{Eis}(\int_{\Lambda_\Sigma^+} h_\Sigma(\sigma)\omega^\vee(\sigma)d_\Sigma\sigma) + \text{Eis}(\int_{\Lambda_\Sigma^+} (1 - h_\Sigma(\sigma))\omega^\vee(\sigma)d_\Sigma\sigma)$$

We have  $d\omega^\vee(\sigma) = 0$  and hence we get

$$\Delta((1 - h_\Sigma(\sigma))\omega^\vee(\sigma)) = d((\chi_\sigma(C) - \chi_\lambda(C))(1 - h_\Sigma(\sigma))\delta\omega^\vee(\sigma))$$

and this implies that

$$\text{Eis}(\int_{\Lambda_\Sigma^+} (1 - h_\Sigma(\sigma))\omega^\vee(\sigma)d_\Sigma\sigma) = d \text{Eis}(\int_{\Lambda_\Sigma^+} (1 - h_\Sigma(\sigma))(\chi_\sigma(C) - \chi_\lambda(C))^{-1}\delta\omega^\vee(\sigma)d_\Sigma\sigma)$$

It is clear that the integrand in the second term-  $\int_{\Lambda_\Sigma^+} (1 - h_\Sigma(\sigma))(\chi_\sigma(C) - \chi_\lambda(C))^{-1}\delta\omega^\vee(\sigma)$  still satisfies (8.20) and then our well known rules above imply that  $\psi = \text{Eis}(\int_{\Lambda_\Sigma^+} (1 - h_\Sigma(\sigma))(\chi_\sigma(C) - \chi_\lambda(C))^{-1}\delta\omega^\vee(\sigma)d_\Sigma\sigma)$  is  $\mathcal{C}_\infty^2$ . Therefore the second term in our above formula is a boundary.

$$\omega_{\text{cont}} = \int_{\Lambda_\Sigma} h_\Sigma(\sigma)\omega(\sigma)d_\Sigma\sigma + d\psi.$$

This is true for any choice of  $h_\Sigma$ . Hence the scalar product  $\langle \omega - d\psi, \omega - d\psi \rangle$  can be made arbitrarily small. Then we claim that the cohomology class  $[\omega] \in H^\bullet(\text{Hom}_{K_\infty}(\Lambda^p(\mathfrak{g}/\mathfrak{k}), \mathcal{C}_\infty(G(\mathbb{Q}) \backslash G(\mathbb{A})/K_f) \otimes \mathcal{M}_\lambda))$  must be zero. This needs a tiny final step.

We invoke Poincaré duality: A cohomology class in  $[\omega] \in H^p(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$  is zero if and only the value of the pairing with any class  $[\omega_2] \in H_c^{d-p}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda^\vee)$  is zero. But the (absolute) value  $[\omega] \cup [\omega_2]$  of the cup product can be given by an integral (See Prop.8.1.2). Therefore it can be estimated by the norm  $\langle \omega - d\psi, \omega - d\psi \rangle$  (Cauchy-Schwarz inequality) and hence must be zero.  $\square$

As usual we denote by  $\widehat{G(\mathbb{R})}$  the unitary spectrum, for us it is simply the set of unitary irreducible representations of  $G(\mathbb{R})$ . Given  $\tilde{\mathcal{M}}_\lambda$ , we define

$$\text{Coh}(\lambda) = \{\pi_\infty \in \widehat{G(\mathbb{R})} \mid H^\bullet(\mathfrak{g}, K_\infty, H_{\pi_\infty} \otimes \tilde{\mathcal{M}}_\lambda) \neq 0\}.$$

The theorem of Harish-Chandra says that this set is finite.

Let

$$H_{\text{Coh}(\lambda)} = \bigoplus_{\pi: \pi_\infty \in \text{Coh}(\lambda)} L^2_{\text{disc}}(G(\mathbb{Q}) \backslash G(\mathbb{A}_f) / K_f)(\pi_\infty \times \pi_f) = \bigoplus_{\pi: \pi_\infty \in \text{Coh}(\lambda)} H_{\pi_\infty}(\pi_f)$$

the theorem of Gelfand-Graev and Langlands assert that this is a finite sum of irreducible modules. This space decomposes again into  $H_{\text{Coh}(\lambda)}^{\text{cusp}} \oplus H_{\text{Coh}(\lambda)}^{\text{res}}$

Then we get the following theorem which due to Borel, Garland, Matsushima and Murakami Bo-Ga-Mu

**Theorem 8.1.1.** *a) The map*

$$H^\bullet(\mathfrak{g}, K_\infty, H_{\text{Coh}(\lambda)}^{(K_\infty)} \otimes \mathcal{M}_\lambda) = \text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), H_{\text{Coh}(\lambda)}^{(K_\infty)} \otimes \mathcal{M}_\lambda) \rightarrow H_{(2)}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$$

*surjective. Especially the image contains  $H_1^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$ .*

*b) (Borel) The homomorphism*

$$H^\bullet(\mathfrak{g}, K_\infty, H_{\text{Coh}(\lambda)}^{(\text{cusp}, K_\infty)} \otimes \mathcal{M}_\lambda) \rightarrow H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$$

*is injective.*

In [5] Prop.5.6, they do not consider the above space  $H_{(2)}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$  we added an  $\epsilon > 0$  to this proposition by claiming that this space is the image.

In general the homomorphism

$$H^\bullet(\mathfrak{g}, K_\infty, H_{\text{res}(\lambda)}^{\text{res}}, K_\infty) \otimes \mathcal{M}_\lambda \rightarrow H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$$

is not injective. We come to this issue in the next section.

If we denote by  $H_{\text{cusp}}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$  the image of the homomorphism in b), then we get a filtration of the cohomology by four subspaces four

$$H_{\text{cusp}}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \subset H_1^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \subset H_{(2)}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \subset H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda). \quad (8.23)$$

We want to point out that our space  $H_{(2)}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$  is not the space denoted by the same symbol in the paper [Bo-Ca]. They define  $L^2$  cohomology as the complex of square integrable forms, i.e.  $\omega$  and  $d\omega$  have to be square integrable. But then a closed form  $\omega$  which is in  $L^2$  gives the trivial class in their cohomology if we can write  $\omega = d\psi$  where  $\psi$  must also be square integrable. In our definition we do not have that restriction on  $\psi$ .

**Hier ein wenig zu Coh( $\lambda$ ) sagen, Vogon-Zuckerman und discrete series, Nochmal Schwermer -J.S. Li**

**A formula for the Poincaré duality pairing**

We assume that  $-w_0(\lambda) = c(\lambda)$ . We have the positive definite hermitian scalar product on  $\text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), H_{\text{Coh}(\lambda)}^{(K_\infty)} \otimes \mathcal{M}_\lambda)$  (See(ü8.6)). On the other hand we have the Poincaré duality pairing

$$H_!^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)(\omega_f) \times H_!^{d-i}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda^\vee})(\omega_{1,f}) \rightarrow \mathbb{C} \quad (8.24)$$

where  $\omega_f \cdot \omega_{1,f} = 1$ . To relate these two products we recall the Hodge  $*$ -operator. (See for instance Vol. I. 4.11) This operator yields an isomorphism

$$* : \text{Hom}_{K_\infty}(\Lambda^p(\mathfrak{g}/\mathfrak{k}), \mathcal{C}_\infty(G(\mathbb{Q}) \backslash G(\mathbb{A})/K_f) \otimes \mathcal{M}_\lambda) \xrightarrow{\sim} \text{Hom}_{K_\infty}(\Lambda^{d-p}(\mathfrak{g}/\mathfrak{k}), \mathcal{C}_\infty(G(\mathbb{Q}) \backslash G(\mathbb{A})/K_f) \otimes \mathcal{M}_{c\lambda}) \quad (8.25)$$

We can use the  $*$  operator to define the adjoint  $\delta = (-1)^{d(p+1)+1} * d*$  and hence the Laplacian  $\Delta$  (See (8.7)). Especially the  $*$  operator yields an identification between the  $\mathcal{C}_\infty$ -functions and the  $\mathcal{C}_\infty$  differential forms in top degree.

We consider two differential forms

$$\omega_1, \omega_2 \in \text{Hom}_{K_\infty}(\Lambda^p(\mathfrak{g}/\mathfrak{k}), \mathcal{C}_\infty^2(G(\mathbb{Q}) \backslash G(\mathbb{A})/K_f) \otimes \mathcal{M}_\lambda)$$

which are square integrable, then we defined the scalar product (See(8.6)  $\langle \omega_1, \omega_2 \rangle$ ) of these two forms. By definition this scalar product is an integral over a function

$$\langle \omega_1, \omega_2 \rangle = \int_{\mathcal{S}_{K_f}^G} \{\omega_1, \omega_2\}.$$

If we have two closed forms  $\omega_1 \in \text{Hom}_{K_\infty}(\Lambda^p(\mathfrak{g}/\mathfrak{k}), \mathcal{C}_\infty^2(G(\mathbb{Q}) \backslash G(\mathbb{A})/K_f) \otimes \mathcal{M}_\lambda)$ ,  $\omega_2 \in \text{Hom}_{K_\infty}(\Lambda^{d-p}(\mathfrak{g}/\mathfrak{k}), \mathcal{C}_\infty^2(G(\mathbb{Q}) \backslash G(\mathbb{A})/K_f) \otimes \mathcal{M}_{\lambda^\vee})$  and if one of these forms has compact support -say  $\omega_2$ -then they define cohomology classes  $[\omega_1] \in H^p(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$ ,  $[\omega_2] \in H_c^{d-p}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda^\vee})$  and the cup product  $[\omega_1] \cup [\omega_2]$  is defined and given by an integral (See proposition 8.1.2) over a form in top degree. Now we check easily - and this is the way how the  $*$  operator is designed that for  $\omega_1, \omega_2 \in \text{Hom}_{K_\infty}(\Lambda^p(\mathfrak{g}/\mathfrak{k}), \mathcal{C}_\infty^2(G(\mathbb{Q}) \backslash G(\mathbb{A})/K_f) \otimes \mathcal{M}_\lambda)$  the integrand

$$\{\omega_1, \omega_2\} = \langle \omega_1 \wedge * \omega_2 \rangle.$$

Now we can formulate the

**Proposition 8.1.4.** *If  $\omega_1, \omega_2 \in \text{Hom}_{K_\infty}(\Lambda^p(\mathfrak{g}/\mathfrak{k}), H_{\text{Coh}(\lambda)}^{(K_\infty)} \otimes \mathcal{M}_\lambda)$  and if both classes  $[\omega_1], [* \omega_2]$  are inner classes, i.e. can be represented by compactly supported forms, then*

$$\langle \omega_1, \omega_2 \rangle = [\omega_1] \cup [* \omega_2]$$

*Proof.* Here we give only a sketch of the proof, for some details we refer to section 8.1.9. Of course we have to recall that the right hand side is defined since we have proposition 6.3.9, we need that both classes are inner classes. We write  $\omega_1 = \tilde{\omega}_1 + d\psi$  where  $\tilde{\omega}_1$  has compact support. Then the value of this cup product is equal to cup12

$$[\omega_1] \cup [* \omega_2] = \int_{\mathcal{S}_{K_f}^G} \tilde{\omega}_1 \wedge * \omega_2. \quad (8.26)$$

We have compact subsets  $\mathcal{S}_{K_f}^G(c)$ , (See ??) we can choose a  $c > 0$  such that the support of  $\tilde{\omega}_1$  lies in  $\mathcal{S}_{K_f}^G(c)$ . Then we get

$$\int_{\mathcal{S}_{K_f}^G(c)} \omega_1 \wedge *\omega_2 = \int_{\mathcal{S}_{K_f}^G(c)} (\tilde{\omega}_1 + d\psi) \wedge *\omega_2 = [\omega_1] \cup [* \omega_2] + \int_{\mathcal{S}_{K_f}^G(c)} \omega_1 \wedge d\psi \tag{8.27}$$

For  $c \rightarrow 0$  the left hand side converges to  $\langle \omega_1, \omega_2 \rangle$ , hence we have to show that

$$\lim_{c \rightarrow 0} \int_{\mathcal{S}_{K_f}^G(c)} \omega_1 \wedge d\psi = \pm \lim_{c \rightarrow 0} \int_{\partial(\mathcal{S}_{K_f}^G(c))} \omega_1 \wedge \psi = 0 \tag{8.28}$$

We know of course that the limit at the right hand side exists. We invoke section 8.1.9, there we will show that we can take  $\psi$  to be square integrable and then it follows that the limit is zero. □

This proposition is delicate. If the quotient  $\mathcal{S}_{K_f}^G$  is compact, then it is of course a consequence of Hodge theory. But if this is not the case we really need that both classes are inner. In fact we have the standard example which shows that this assumption is needed. If take  $\omega_1 = \omega_2$  to be the form in degree zero given by the constant function 1. Then the left hand side is non zero but the class  $*1$  is the volume form which is trivial if  $\mathcal{S}_{K_f}^G$  is not compact, and therefore the right hand side is not zero.

The proposition has the following nice corollary

**Corollary 8.1.1.** *If  $\omega \in \text{Hom}_{K_\infty}(\Lambda^p(\mathfrak{g}/\mathfrak{k}), H_{\text{Coh}(\lambda)}^{(K_\infty)} \otimes \mathcal{M}_\lambda)$  is non zero and if the restriction of  $*\omega$  to the boundary is zero then  $[\omega] \neq 0$ .*

This last Corollary could be useful if we want to understand the kernel of the map

$$\text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), H_{\text{Coh}(\lambda)}^{(K_\infty)} \otimes \mathcal{M}_\lambda) \rightarrow H^\bullet(\mathcal{S}_{K_f}^G, \mathcal{M}_\lambda), \tag{8.29}$$

but a closer look tells us that this may not be so easy, because the restriction the cohomology to the boundary cohomology is not so easy to understand.

Now we remember that in the previous sections we made the convention (See end of (8.1.1)) that our coefficient systems  $\mathcal{M}_\lambda$  are  $\mathbb{C}$  vector spaces. We now revoke this convention and recall that the coefficient systems  $\mathcal{M}_\lambda$  should be replaced by  $\mathcal{M}_\lambda \otimes_F \mathbb{C}$ . Then in the above list (8.23) of four subspaces in the cohomology the second and the fourth subspace have a natural structure of  $F$ -vector spaces and they have a combinatorial definition, whereas the first and third subspace need some input from analysis in their definition. In other words if we replace  $\mathcal{M}_\lambda$  in (8.23) by  $\mathcal{M}_\lambda \otimes_f \mathbb{C}$  then the second and the fourth space can be written as

$$H_i^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \otimes_F \mathbb{C} \subset H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \otimes_F \mathbb{C}$$

We believe that also the third space has a combinatorial definition, for this we need the weighted cohomology groups: *Weighted cohomology*; G. Harder; R. MacPherson; M. Goresky *Inventiones mathematicae* (1994).

**8.1.7 Consequences.**

**Vanishing theorems**

If  $V$  is unitary and irreducible, then we have that  $\bar{V} \xrightarrow{\sim} V^\vee$  and this implies for the central character

$$\overline{\chi_V(z)} = \chi_{V^\vee}(z) \text{ for all } z \in \mathfrak{Z}(\mathfrak{g}).$$

Combining this with Wigner’s lemma we can conclude

*If  $V$  is an irreducible unitary  $(\mathfrak{g}, K_\infty)$ -module,  $\mathcal{M}_\lambda$  is an irreducible rational representation, and if*

$$H^\bullet(\mathfrak{g}, K_\infty, V \otimes \mathcal{M}_\lambda) \neq 0$$

*then  $\chi_{\mathcal{M}_\lambda^\vee}(z) = \chi_{\mathcal{M}_\lambda}(z) = \chi_{\tilde{\mathcal{M}}_\lambda}(z)$*

*In other words: For an unitary irreducible  $(\mathfrak{g}, K_\infty)$ -module  $V$  the cohomology with coefficients in an irreducible rational representation  $\mathcal{M}$  vanishes, unless we have  $\mathcal{M}_\lambda^\vee \xrightarrow{\sim} \tilde{\mathcal{M}}_\lambda$ , or in terms of highest weights unless  $-w_0(\lambda) = c(\lambda)$ . (See 3.1.1)*

If we combine this with the considerations following Wigner’s lemma we get

**Corollary** *If  $\mathcal{M}$  is an absolutely irreducible rational representation and if  $\mathcal{M}_\lambda^\vee$  is not isomorphic to  $\tilde{\mathcal{M}}_\lambda$  then*

$$H_{(2)}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) = 0.$$

Hence also

$$H_!^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) = 0.$$

We will discuss examples for this in section 8.1.7

**The group  $G/\mathbb{Q} = \mathrm{Sl}_2/\mathbb{Q}$**

Let us consider the group  $G/\mathbb{Q} = \mathrm{Sl}_2/\mathbb{Q}$ . We have tautological representation  $\mathrm{Sl}_2 \hookrightarrow \mathrm{Gl}(\mathbb{Q}^2) = \mathrm{Gl}(V)$  and we get all irreducible representations of we take the symmetric powers  $\mathcal{M}_n = \mathrm{Sym}^n(V)$  of  $V$ . (See 2, these are the  $\mathcal{M}_n[m]$  restricted to  $\mathrm{Sl}_2$ , then the  $m$  drops out.)

In this case the Vogan-Zuckerman list is very short. It is discussed in [Slzwei] for the groups  $\mathrm{Sl}_2(\mathbb{R})$  and  $\mathrm{Sl}_2(\mathbb{C})$ , where both groups are considered as real Lie-groups.

In the case  $\mathrm{Sl}_2(\mathbb{R})$  we have the trivial module  $\mathbb{C}$  and for any integer  $k \geq 2$  we have two irreducible unitarizable  $(\mathfrak{g}, K_\infty)$ -modules  $\mathcal{D}_k^\pm$  (the discrete series representations) (See [Slzwei], 4.1.5 ). These are the only  $(\mathfrak{g}, K_\infty)$ -modules which have non trivial cohomology with coefficients in a rational representation. If we now pick one of our rational representation  $\mathcal{M}_n$ , then the non vanishing cohomology groups are

$$\begin{aligned} H^q(\mathfrak{g}, K_\infty, \mathcal{M}_n \otimes \mathbb{C}) &= \mathbb{C} \text{ for } l = 0, q = 0, 2 \\ H^q(\mathfrak{g}, K_\infty, \mathcal{D}_k^\pm \otimes \mathcal{M}_n \otimes \mathbb{C}) &= \mathbb{C} \text{ for } l = k - 2, q = 1 \end{aligned}$$

The trivial  $(\mathfrak{g}, K_\infty)$ -module  $\mathbb{C}$  occurs with multiplicity one in  $L^2(G(\mathbb{Q})\backslash G(\mathbb{A})/K_f)$  hence we get for the trivial coefficient system a contribution

$$H^\bullet(\mathfrak{g}, K_\infty, \mathbb{C} \otimes \mathcal{M}_n \otimes \mathbb{C}) = H^0(\mathfrak{g}, K_\infty, \mathbb{C}) \oplus H^2(\mathfrak{g}, K_\infty, \mathbb{C}) = \mathbb{C} \oplus \mathbb{C} \rightarrow H_{(2)}^\bullet(\mathcal{S}_{K_f}^G, \mathbb{C}).$$

This map is injective in degree 0 and zero in degree 2.

For the modules  $\mathcal{D}_k^\pm$  we have to determine the multiplicities  $m^\pm(k)$  of these modules in the discrete spectrum of  $L^2(G(\mathbb{Q})\backslash G(\mathbb{A})/K_f)$ . A simple argument using complex conjugation tells us  $m^+(k) = m^-(k)$ . Now we have the fundamental observation made by Gelfand and Graev, which links representation theory to automorphic forms:

*We have an isomorphism*

$$\begin{aligned} \text{Hom}_{\mathfrak{g}, K_\infty}(\mathcal{D}_k^+, L_{\text{disc}}^2(G(\mathbb{Q})\backslash G(\mathbb{A})/K_f)) &\xrightarrow{\sim} S_k(G(\mathbb{Q})\backslash \mathbb{H} \times G(\mathbb{A}_f)/K_f) = \\ &\text{space of holomorphic cusp forms of weight } k \text{ and level } K_f \end{aligned}$$

This is also explained in [Slzwei] on the pages following 23. We explain how we get starting from a holomorphic cusp form  $f$  of weight  $k$  an inclusion

$$\Phi_f : \mathcal{D}_k^+ \hookrightarrow L_{\text{disc}}^2(G(\mathbb{Q})\backslash G(\mathbb{A})/K_f)$$

and that this map  $f \mapsto \Phi_f$  establishes the above isomorphism. This gives us the famous Eichler-Shimura isomorphism

$$S_k(G(\mathbb{Q})\backslash \mathbb{H} \times G(\mathbb{A}_f)/K_f) \oplus \overline{S_k(G(\mathbb{Q})\backslash \mathbb{H} \times G(\mathbb{A}_f)/K_f)} \xrightarrow{\sim} H_1^G(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{k-2}).$$

**The group**  $G/\mathbb{Q} = R_{F/\mathbb{Q}}(\text{Sl}_2/F)$ .

For any finite extension  $F/\mathbb{Q}$  we may consider the base restriction  $G/\mathbb{Q} = R_{F/\mathbb{Q}}(\text{Sl}_2/F)$ . (See Chap-II. 1.1.1). Here we want to consider the special case the  $F/\mathbb{Q}$  is imaginary quadratic. In this case we have  $G \otimes \mathbb{C} = \text{Sl}_2 \times \text{Sl}_2/\mathbb{C}$  the factors correspond to the two embeddings of  $F$  into  $\mathbb{C}$ . The rational irreducible representations are tensor products of irreducible representations of the two factors  $\mathcal{M}_\lambda = \mathcal{M}_{k_1} \otimes \mathcal{M}_{k_2}$  where again  $\mathcal{M}_k = \text{Sym}^k(\mathbb{C}^2)$ . These representations are defined over  $F$ .

In this case we discuss the Vogan-Zuckerman list in [Slzwei], here we want to discuss a particular aspect. We observe that

$$\mathcal{M}_\lambda^\vee \xrightarrow{\sim} \mathcal{M}_{k_1} \otimes \mathcal{M}_{k_2}, \bar{\mathcal{M}}_\lambda = \mathcal{M}_{k_2} \otimes \mathcal{M}_{k_1}$$

and hence our corollary above yields for any choice of  $K_f$

$$H_1^\bullet(\mathcal{S}_{K_f}^G, \mathcal{M}) = 0 \text{ if } k_1 \neq k_2.$$

In Chapter II we discuss the special examples in low dimensions. We take  $F = \mathbb{Q}[i]$  and  $\Gamma = \text{Sl}_2[\mathbb{Z}[i]]$  this amounts to taking the standard maximal compact subgroup  $K_f = \text{Sl}_2[\mathcal{O}_F]$ . If now for instance  $k_1 > 0$  and  $k_2 = 0$ , then we get  $H_1^\bullet(\mathcal{S}_{K_f}^G, \mathcal{M}_\lambda) = 0$ . Hence we have by definition  $H_1^\bullet(\mathcal{S}_{K_f}^G, \mathcal{M}) = H_{\text{Eis}}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})$

and we have complete control over the Eisenstein- cohomology in this case. Hence we know the cohomology in this case if we apply the analytic methods.

On the other hand in Chapter II we have written an explicit complex of finite dimensional vector spaces, which computes the cohomology. It is not clear to me how we can read off this complex the structure of the cohomology groups.

We get another example where this phenomenon happens, if we consider the group  $Sl_n/\mathbb{Q}$  if  $n > 2$ . In Chap. IV 1.2 we described the simple roots  $\alpha_1, \alpha_2, \dots, \alpha_{n-1}$ , accordingly we have the fundamental highest weights  $\omega_1, \dots, \omega_{n-1}$ . The element  $w_0$  (See 8.1.1) has the effect of reversing the order of the weights. Hence we see that for  $\lambda = \sum n_i \omega_i$  we have

$$H_!^\bullet(\mathcal{S}_{K_f}^G, \mathcal{M}_\lambda) = 0$$

unless we have  $-w_0(\lambda) = \lambda$  and this means  $n_i = n_{n-1-i}$ .

**The algebraic  $K$ -theory of number fields**

I briefly recall the definition of the  $K$ -groups of an algebraic number field  $F/\mathbb{Q}$ . We consider the group  $Gl_n(O_F)$ , it has a classifying space  $BG_n$ . We can pass to the limit  $\lim_{n \rightarrow \infty} Gl_n(O_F) = Gl(O_F) = G$  and let  $BG$  its classifying space. Quillen invented a procedure to modify this space to another space  $BG^+$ , whose fundamental group is now abelian, but which has the same homology and cohomology as  $BG$ . Then he defines the algebraic  $K$ -groups as

$$K_i(\mathcal{O}_F) = \pi_i(BG^+).$$

The space is an  $H$ -space, this means that we have a multiplication  $m : BG^+ \times BG^+ \rightarrow BG^+$  which has a two sided identity element. Then we get a homomorphism  $m^\bullet : H^\bullet(BG^+, \mathbb{Z}) \rightarrow H^\bullet(BG^+ \times BG^+, \mathbb{Z})$  and if we tensorize by  $\mathbb{Q}$  and apply the Künneth-formula then we get the structure of a Hopf algebra on the Cohomology

$$m^\bullet : H^\bullet(BG^+, \mathbb{Q}) \rightarrow H^\bullet(BG^+, \mathbb{Q}) \otimes H^\bullet(BG^+, \mathbb{Q})$$

Then a theorem of Milnor asserts that the rational homotopy groups

$$\pi_i(BG^+) \otimes \mathbb{Q} = \text{prim}(H^i(BG, \mathbb{Q})),$$

where  $\text{prim}$  are the primitive elements, i.e. those elements  $x \in H^i(BG, \mathbb{Q})$  for which

I sketch a second application. We discuss the group  $G = R_{F/\mathbb{Q}}(Gl_n/F)$ , where  $F/\mathbb{Q}$  is an algebraic number field. the coefficient system  $\tilde{\mathcal{M}}_\lambda = \mathbb{C}$  is trivial. In this case Borel, Garland and Hsiang have shown hat in low degrees  $q \leq n/4$

$$H^q(\mathcal{S}_{K_f}^G, \mathbb{C}) = H_{(2)}^q \mathcal{S}_{K_f}^G, \mathbb{C}.$$

On the other hand it follows from the Vogan-Zuckerman classification, that the only irreducible unitary  $(\mathfrak{g}, K_\infty)$  modules  $V$ , for which  $H^q(\mathfrak{g}, K_\infty, V) \neq 0$  and  $q \leq n/4$  are one dimensional.

Hence we see that in low degrees

$$H^q(\mathfrak{g}, K_\infty, \mathbb{C}) \rightarrow H^q(\mathcal{S}_{K_f}^G, \mathbb{C})$$

is an isomorphism (Injectivity requires some additional reasoning.)

On the other hand we have  $H^q(\mathfrak{g}, K_\infty, \mathbb{C}) = \text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \mathbb{C})$  and obviously this last complex is isomorphic to the complex  $\Omega^\bullet(X)^{G(\mathbb{R})}$  of  $G(\mathbb{R})$ -invariant forms on the symmetric space  $G(\mathbb{R})/K_\infty$ . Our field has different embeddings  $\tau : F \hookrightarrow \mathbb{C}$ , the real embeddings factor through  $\mathbb{R}$ , they form the set  $S_\infty^{\text{real}}$  and the pairs of may conjugate embeddings into  $\mathbb{C}$  form the set  $S_\infty^{\text{comp}}$ . Then

$$X = \prod_{v \in S_\infty^{\text{real}}} \text{Sl}_n(\mathbb{R})/SO(n) \times \prod_{S_\infty^{\text{comp}}} \text{Sl}_n(\mathbb{C})/SU(n).$$

Now the complex  $\Omega^\bullet(X)^{G(\mathbb{R})}$  of invariant differential forms (all differentials are zero) does not change if we replace the group

$$G(\mathbb{R}) = \prod_{v \in S_\infty^{\text{real}}} \text{Sl}_n(\mathbb{R}) \times \prod_{S_\infty^{\text{comp}}} \text{Sl}_n(\mathbb{C})$$

by its compact form  $G_c(\mathbb{R})$  and then we get the complex of invariant forms on the compact twin of our symmetric space

$$X_c = \prod_{v \in S_\infty^{\text{real}}} SU_n(\mathbb{R})/SO(n) \times \prod_{S_\infty^{\text{comp}}} (SU(n) \times SU(n))/SU(n),$$

but then

$$\Omega(X_c)^{G_c(\mathbb{R})} = H^\bullet(X_c, \mathbb{C}).$$

The cohomology of the topological spaces like the one on the right hand side has been computed by Borel in the early days of his career.

If we let  $n$  tend to infinity, we can consider the limit of these cohomology groups, then the limit becomes a Hopf algebra and we can consider the primitive elements

### The semi-simplicity of the inner cohomology

Now we assume again that our representation  $\tilde{\mathcal{M}}_\lambda$  is defined over some number field  $F$  we consider it as a subfield of  $\mathbb{C}$ . In other word we have a representation  $r : G \times F \rightarrow \text{Gl}(\mathcal{M}_\lambda)$ . We have defined  $H_1^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})$ , this is a finite dimensional  $F$ -vector space and Theorem 3.1.1 in Chapter 3 asserts that this is a semi simple module under the Hecke algebra. The following argument shows that this is an easy consequence of our results above.

The module  $H_1 \subset L_{\text{disc}}^2(G(\mathbb{Q}) \backslash G(\mathbb{A}_f)/K_f)$  can also be decomposed into a finite direct sum of irreducible  $G(\mathbb{R}) \times \mathcal{H}_{K_f}$  modules

$$H_1 = \bigoplus_{\pi_\infty \otimes \pi_f \in \hat{H}_1} (H_{\pi_\infty} \otimes H_{\pi_f})^{m_1(\pi_\infty \times \pi_f)},$$

this module is clearly semi-simple. Of course it is not a  $(\mathfrak{g}, K_\infty)$ -module, but we can restrict to the  $K_\infty$ -finite vectors and get

$$H^\bullet(\mathfrak{g}, K_\infty, H_1^{(K_\infty)} \otimes \mathcal{M}_\lambda \otimes \mathbb{C}) = \bigoplus_{\pi_\infty \otimes \pi_f \in \hat{H}_1} (\text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), H_{\pi_\infty} \otimes \mathcal{M}_\mathbb{C}) \otimes H_{\pi_f})^{m_1(\pi_\infty \times \pi_f)}$$

This is a decomposition of the left hand side into irreducible  $\mathcal{H}_{K_f}$  modules. Now we have the surjective map

$$H^\bullet(\mathfrak{g}, K_\infty, H_1^{(K_\infty)} \otimes \mathcal{M}_\lambda \otimes \mathbb{C}) \rightarrow H_{(2)}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{C})$$

hence it follows that  $H_{(2)}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{C})$  is a semi simple  $\mathcal{H}_{K_f}$  module and hence also  $H_!^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$  is a semi simple  $\mathcal{H}_{K_f}$  module.

At this point we encounter an interesting problem. We have the three subspaces (See end of 3.2)

$$H_{\text{cusp}}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{C}) \subset H_!^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \otimes \mathbb{C} \subset H_{(2)}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{C}) \subset H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \otimes \mathbb{C},$$

note the positions of the tensor symbol  $\otimes$ . The first and the third space are only defined after we tensorize the coefficient system by  $\mathbb{C}$ , whereas the second and the fourth cohomology groups by definition  $F$  vector spaces tensorized by  $\mathbb{C}$ .

Now the question is whether the first and the third space also have a natural  $F$ -vector space structure. Of course we get a positive answer, if the Manin-Drinfeld principle holds. All the vector spaces are of course modules under the Hecke algebra and we can look at their spectra

$$\begin{aligned} \Sigma(H_{\text{cusp}}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{C})) &= \Sigma_{\text{cusp}} & \Sigma(H_!^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{C})) &= \Sigma_! \\ \Sigma(H_{(2)}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{C})) &= \Sigma_{(2)} & \Sigma(H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{C})) &= \Sigma \end{aligned}$$

If now for instance  $\Sigma_{\text{cusp}} \cap (\Sigma_! \setminus \Sigma_{\text{cusp}}) = \emptyset$  then we can define  $H_{\text{cusp}}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \subset H_!^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$  as the subspace which is the sum of the isotypical components in  $\Sigma_{\text{cusp}}$ .

If this is the case we say that the cuspidal cohomology is *intrinsically definable* and we get a canonical decomposition

$$H_!^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) = H_{\text{cusp}}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \oplus H_{!, \text{noncusp}}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda).$$

The classical Manin-Drinfeld principle refers to the two spectra  $\Sigma_! \subset \Sigma$ , if it is true in this case we get a decomposition

$$H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) = H_!^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \oplus H_{\text{Eis}}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$$

the canonical complement is called the Eisenstein cohomology. (See Chap. II 2.2.3 and Chap III 5.)

### 8.1.8 Growth of cohomology classes

The fundamental exact sequence yields a short sequence fil0

$$0 \rightarrow H_!^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}) \rightarrow H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}) \rightarrow H^\bullet(\mathcal{N}(\mathcal{S}_{K_f}^G), \tilde{\mathcal{M}}) \quad (8.30)$$

and we gained some understanding of  $H_!^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})$  using analytic methods. We have seen that classes in  $[\omega] \in H_{(2)}^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})$  can be represented by harmonic forms  $\omega$ . Of course the condition that  $\omega$  is square integrable implies some

restriction on the growth of  $\omega$ . It is our goal in this section to find criteria which imply that a closed form  $\omega$  or a class  $[\omega]$  is square integrable.

To attack this kind of question we study the "asymptotic behavior" of the cohomology at infinity, this means that we have to study  $H^q(\dot{\mathcal{N}}(\mathcal{S}_{K_f}^G), \tilde{\mathcal{M}})$ . We apply reduction theory (See section 1.2.8) and start from the covering

$$\dot{\mathcal{N}}(\mathcal{S}_{K_f}^G) = \bigcup_{P:P\text{proper}} \Gamma_P \backslash X^P(c_{\pi'}, r(\pi')) \tag{8.31}$$

Of course we know: The form  $\omega$  is square integrable if and only if its restriction to the open sets is square integrable.

We start by describing the cohomology of the open sets in the covering, i.e. we consider the cohomology  $H^\bullet(X^P(c_{\pi'}, r(\pi')), \tilde{\mathcal{M}})$  we recall that we have the spectral sequence (2.39)

$$H^p(\Gamma_M \backslash X^M(r), H^q(\widetilde{(\Gamma_{U_P} \backslash U_P(\mathbb{R}))}, \tilde{\mathcal{M}})) \Rightarrow H^{p+q}(\Gamma_P \backslash X^P(C(\tilde{z})), \tilde{\mathcal{M}})$$

and the first step is to get more information on the  $M$ - module  $H^\bullet(\Gamma_{U_P} \backslash U_P(\mathbb{R}), \tilde{\mathcal{M}})$ .

**The cohomology of unipotent groups**

We drop the subscript  $P$ , we know that the group scheme  $U/\mathbb{Q}$  is a unipotent group scheme, this means that  $U/\mathbb{Q}$  has a filtration by subschemes  $U_0 = \{e\} \subset U_1 \subset U_2 \subset \dots \subset U_{m-1} \subset U_m$  such that  $U_i/U_{i-1} \xrightarrow{\sim} \mathbb{G}_a$ . The subgroup  $\Gamma_U \subset U(\mathbb{Q})$  is Zariski dense, more precisely we know the following: If  $\Gamma_i = U_i(\mathbb{Q}) \cap \Gamma$  then  $\Gamma_i/\Gamma_{i-1} \xrightarrow{\sim} \mathbb{Z} \subset U_i/U_{i-1}(\mathbb{R}) \xrightarrow{\sim} \mathbb{R}$ .

We consider the category of  $U/\mathbb{Q}$  modules  $\text{Mod}_U$  (see section 1.1.1). Then it is clear that the functor  $\mathcal{M} \rightarrow \mathcal{M}^U$  is equal to  $\mathcal{M} \rightarrow \mathcal{M}^{\Gamma_U}$ . ( Our  $\mathbb{Z}$  module  $\mathcal{M}$  above is now a  $\mathbb{Q}$ - vector space, i.e. we consider coefficient systems with rational coefficients.)

We choose the action of  $U$  on  $A$  by left translations on  $A$ . It follows from Frobenius reciprocity that the  $U/\mathbb{Q}$  module  $A$  is an injective module in  $\text{Mod}_U$ . (See ???) This implies that we get an injective resolution of the  $U/\mathbb{Q}$  -module  $\mathbb{Q}$  by

$$0 \rightarrow \mathbb{Q} \rightarrow A \rightarrow (A/\mathbb{Q}) \otimes A \rightarrow \dots = 0 \rightarrow \mathbb{Q} \rightarrow I^0 \rightarrow I^1 \rightarrow \tag{8.32}$$

and hence

$$H^q(U, \mathcal{M}) = H^q(\Gamma_U \backslash U(\mathbb{R}), \mathcal{M}) = H^q(0 \rightarrow (I^1 \otimes \mathcal{M})^U \rightarrow (I^2 \otimes \mathcal{M})^U \rightarrow \dots) = H^q((I^\bullet \otimes \mathcal{M})^U) \tag{8.33}$$

Since  $U/\mathbb{Q}$  is the unipotent radical of the parabolic group  $P/\mathbb{Q}$ , the parabolic group  $P/\mathbb{Q}$  acts via the adjoint action on the modules  $I^m$ . This action respects the submodules  $(I^m)^U$  and  $U/\mathbb{Q}$  acts trivially on  $(I^m)^U$ , this implies that the modules  $(I^m)^U$  are  $M/\mathbb{Q} = (P/U)/\mathbb{Q}$  modules. The group  $M/\mathbb{Q}$  is reductive and we know that the category of  $M/\mathbb{Q}$  modules is semi simple (???). This implies that we can decompose

$$(I^\bullet)^U = \mathbb{H}^\bullet(U, \mathcal{M}) \oplus ACI(I^\bullet)^U \tag{8.34}$$

where the first summand is a complex of  $M/\mathbb{Q}$ -modules in which all the differentials are zero and the second is an acyclic complex of  $M/\mathbb{Q}$ -modules. Hence

$$H^\bullet(U, \mathcal{M}) = H^\bullet(\Gamma_U, \mathcal{M}) \xrightarrow{\sim} \mathbb{H}^\bullet(U, \mathcal{M}) \tag{8.35}$$

We get a "smaller" resolution from the (algebraic) de-Rham complex of differential forms. On the smooth affine scheme  $U/\mathbb{Q}$  we have the sheaves of differential forms  $\Omega_U^p = \Lambda^p \Omega_U^1$  ([29], 7.5) and we have the de-Rham complex

$$\Omega(U)^\bullet = 0 \rightarrow \mathbb{Q} \rightarrow A \rightarrow \Omega^1(U) \rightarrow \Omega^2(U) \rightarrow \dots \tag{8.36}$$

where  $\Omega^p(U) = \Omega_U^p(U)$  is the module of global sections and  $A = \Omega^0(U)$ . These modules of differentials are free  $A$  modules, hence they are injective. Since our unipotent group scheme  $U/\mathbb{Q}$  is isomorphic to the affine space  $\mathbb{A}^d$  (as affine scheme) we see easily that this complex is exact, hence it provides an acyclic resolution. As before we get the cohomology by taking the complex  $(\Omega^p(U) \otimes \mathcal{M})^U$  of invariants under the action of  $U/\mathbb{Q}$ . Since an  $U/\mathbb{Q}$ -invariant differential form with values in  $\mathcal{M}$  is determined by its value at the identity  $e$  the complex of invariants under  $U/\mathbb{Q}$  becomes

$$0 \rightarrow \mathcal{M} \rightarrow \text{Hom}(\mathfrak{u}, \mathcal{M}) \rightarrow \text{Hom}(\Lambda^2 \mathfrak{u}, \mathcal{M}) \rightarrow \dots = 0 \rightarrow \text{Hom}(\Lambda^\bullet \mathfrak{u}, \mathcal{M}) \tag{8.37}$$

and the cohomology of this complex is the cohomology  $H^\bullet(\mathfrak{u}, \mathcal{M})$ . We still have the action of  $P/\mathbb{Q}$  on  $\mathfrak{u}$  by the adjoint action, hence we get an action of  $P$  on  $\text{Hom}(\Lambda^\bullet \mathfrak{u}, \mathcal{M})$  and we have

**Theorem 8.1.2.** (*van Est [?]*)

$$H^\bullet(\mathfrak{u}, \mathcal{M}) \xrightarrow{\sim} \mathbb{H}^\bullet(\mathfrak{u}, \mathcal{M}) = (\text{Hom}(\Lambda^\bullet \mathfrak{u}, \mathcal{M}))^U,$$

and therefore  $H^\bullet(\mathfrak{u}, \mathcal{M})$  is a  $M/\mathbb{Q}$  module.

*Proof.* later □

A theorem of Kostant yields a description of the  $M/\mathbb{Q}$  module  $(\text{Hom}(\Lambda^\bullet \mathfrak{u}, \mathcal{M}))^U$ , it gives us the decomposition into highest modules. Let  $\lambda \in X^*(T)$  be the highest weight of  $\mathcal{M}$ , i.e. we have  $\mathcal{M} = \mathcal{M}_\lambda$ . The set

$$W^P = \{w \in W \mid w^{-1}(\alpha) \in \Delta^+\} \tag{8.38}$$

is the set of Kostant representatives for  $W^M \backslash W$ . For any  $w \in W^P$  we define the element

$$\omega_w = \Lambda_{\alpha \in \Delta_U; w^{-1}\alpha < 0} u_\alpha^\vee \otimes e_{w\lambda} \tag{8.39}$$

**Proposition 8.1.5.** *This element  $\omega_w$  lies in  $\mathbb{H}^\bullet(\mathfrak{u}, \mathcal{M})$  and it is a highest weight vector for the action of  $M/\mathbb{Q}$ , the weight is  $w \cdot \lambda = w\lambda + w\rho - \rho = w(\lambda + \rho) - \rho$ .*

*Proof.* This is an easy computation. □

This highest weight vector provides an irreducible highest weight module  $\mathcal{M}_{w \cdot \lambda}$  for  $M/\mathbb{Q}$  and we have the famous theorem of Kostant

**Theorem 8.1.3.**

$$\mathbb{H}^\bullet(\mathbf{u}, \mathcal{M}) = \bigoplus_{w \in W^P} \mathcal{M}_{w \cdot \lambda}[l(w)]$$

where the summand  $\mathcal{M}_{w \cdot \lambda}$  sits in degree  $l(w) = ?$ .

*Proof.* Rather clear after the preparation.  $\square$

Since the differentials in the complex  $\mathbb{H}^\bullet(\mathbf{u}, \mathcal{M})$  are zero, the spectral sequence degenerates and we get cohbounstrat

$$H^n(\Gamma_P \backslash X^P(C(\tilde{\mathcal{C}})), \tilde{\mathcal{M}}) \xrightarrow{\sim} \bigoplus_{w \in W^P} H^{n-l(w)}(\Gamma_M \backslash X^M(c_P), \mathbb{H}^{l(w)}(\mathbf{u}, \mathcal{M})[w \cdot \lambda]), \quad (8.40)$$

this is the decomposition of the cohomology of the boundary stratum into weight spaces.

The cohomology groups  $H^\bullet(\Gamma_P \backslash X^P(C(\tilde{\mathcal{C}})), \tilde{\mathcal{M}} \otimes \mathbb{C})$  can be computed as the cohomology groups of the de-Rham complex

$$H^p(\Gamma_P \backslash X^P(C(\tilde{\mathcal{C}})), \tilde{\mathcal{M}} \otimes \mathbb{C}) \xrightarrow{\sim} H^p(\Omega^\bullet(\Gamma_P \backslash X^P(C(\tilde{\mathcal{C}})), \tilde{\mathcal{M}} \otimes \mathbb{C}) \quad (8.41)$$

here  $\Omega^\bullet(\Gamma_P \backslash X^P(C(\tilde{\mathcal{C}}))$  is the complex of those  $\mathcal{C}_\infty$  differential forms which extend to a  $\mathcal{C}_\infty$  form into a small open neighborhood of  $X^P(C(\tilde{\mathcal{C}}))$ . We want to use the decomposition of the cohomology into weight spaces to establish a "much smaller" sub-complex

$$\Omega_{\log}^\bullet(\Gamma_P \backslash X^P(C(\tilde{\mathcal{C}})), \tilde{\mathcal{M}} \otimes \mathbb{C}) \hookrightarrow \Omega^\bullet(\Gamma_P \backslash X^P(C(\tilde{\mathcal{C}})), \tilde{\mathcal{M}} \otimes \mathbb{C})$$

such that the inclusion induces an isomorphism in cohomology. We recall the map

$$q_{P,M} : \Gamma_P \backslash X^P(c_{\pi'}, r(c_{\pi'})) \rightarrow \Gamma_M \backslash X^M(r(c_P)) \times \prod_{\alpha \in \pi'} (0, c_\alpha] \quad (8.42)$$

it provides a map

$$q_{P,M}^\bullet : \Omega^\bullet(\Gamma_M \backslash X^M(r(c_P)) \times \prod_{\alpha \in \pi'} (0, c_\alpha]) \otimes \mathbb{H}^\bullet(\mathbf{u}, \mathcal{M}) \rightarrow \Omega^\bullet(\Gamma_P \backslash X^P(C(\tilde{\mathcal{C}})) \otimes \mathcal{M} \quad (8.43)$$

This map is defined as follows. Let

$$\omega^p \otimes \omega_U^q \in \Omega^p(\Gamma_M \backslash X^M(r(c_P)) \times \prod_{\alpha \in \pi'} (0, c_\alpha]) \otimes \mathbb{H}^q(\mathbf{u}, \mathcal{M})$$

For a point  $x \in \Gamma_P \backslash X^P(c_{\pi'}, r(c_{\pi'}))$  we have to give the value of  $q_{P,M}^{p+q}(\omega^p \otimes \omega_U^q)(x)$ . Hence we to determine the value of  $q_{P,M}^{p+q}(\omega^p \otimes \omega_U^q)(x)$  at a  $p+q$ -tuple of tangent vectors. We choose  $p$  tangent vectors  $t_1^M, \dots, t_p^M$  arbitrarily, they map to tangent vectors  $\bar{t}_1, \dots, \bar{t}_p$  under  $q_{P,M}$ . Then we choose  $q$  tangent vectors  $u_1, \dots, u_q$  which are tangent to the fiber. The fiber is identified to  $\Gamma_U \backslash U(\mathbb{R})$  and hence  $u_1, \dots, u_q \in \mathfrak{u}^\vee$ . With  $m = (m_1, a) = q_{P,M}$  we get

$$q_{P,M}^{p+q}(\omega^p \otimes \omega_U^q)(x) = \omega^p(\bar{t}_1, \dots, \bar{t}_p)((m_1, a)) \omega_U^q(u_1, \dots, u_q) \quad (8.44)$$

and (8.40) implies that  $q_{P,M}^\bullet$  induces an isomorphism in cohomology.

The image under this map is not yet what we want. Again we can consider the sub complex

$$\Omega^\bullet(\Gamma_M \backslash X^M(r(\underline{c}_P))) \otimes \Omega^\bullet\left(\prod_{\alpha \in \pi'} (0, c_\alpha)\right) \subset \Omega^\bullet(\Gamma_M \backslash X^M(r(\underline{c}_P) \times \prod_{\alpha \in \pi'} (0, c_\alpha))).$$

this inclusion induces an isomorphism in cohomology. We pick an element  $w \in W^P$  and consider the complex

$$\Omega^\bullet(\Gamma_M \backslash X^M(r(\underline{c}_P))) \otimes \Omega^\bullet\left(\prod_{\alpha \in \pi'} (0, c_\alpha)\right) \otimes \mathbb{H}^{l(w)}(\mathfrak{u}, \mathcal{M})[w \cdot \lambda], \quad (8.45)$$

this complex still computes the cohomology  $H^p(\Gamma_P \backslash X^P(C(\tilde{c})), \tilde{\mathcal{M}} \otimes \mathbb{C})[w \cdot \lambda]$ . We look for a suitable small sub complex of  $\Omega^\bullet(\prod_{\alpha \in \pi'} (0, c_\alpha)) \otimes \mathbb{H}^{l(w)}(\mathfrak{u}, \mathcal{M})[w \cdot \lambda]$ . We embed  $\prod_{\alpha \in \pi'} (0, c_\alpha) \subset \prod_{\alpha \in \pi'} \mathbb{R}_{>0}^\times = A_{\pi'}$ , we have the restriction

$$\Omega^\bullet(A_{\pi'}) \otimes \mathbb{H}^{l(w)}(\mathfrak{u}, \mathcal{M})[w \cdot \lambda] \xrightarrow{res} \Omega^\bullet\left(\prod_{\alpha \in \pi'} (0, c_\alpha)\right) \otimes \mathbb{H}^{l(w)}(\mathfrak{u}, \mathcal{M})[w \cdot \lambda].$$

Then  $\mathbb{H}^{l(w)}(\mathfrak{u}, \mathcal{M})[w \cdot \lambda]$  is a  $A_{\pi'}$ -module, the action is given by the restriction of  $w \cdot \lambda$  to  $A_{\pi'}$ . Let  $\mathfrak{a}_{\pi'}$  be the Lie-algebra of  $A_{\pi'}$  then

$$\Omega^\bullet(A_{\pi'}) \otimes \mathbb{H}^{l(w)}(\mathfrak{u}, \mathcal{M})[w \cdot \lambda] = \text{Hom}(\Lambda^\bullet(\mathfrak{a}_{\pi'}), \mathcal{C}_\infty(A_{\pi'}) \otimes \mathbb{H}^{l(w)}(\mathfrak{u}, \mathcal{M})[w \cdot \lambda]).$$

Of course we know that the cohomology of this complex sits in degree zero, our small sub complex hence our small sub complex must have the same property.

We look at the degree zero, for an element  $\omega = f \otimes u \in \mathcal{C}_\infty(A_{\pi'}) \otimes \mathbb{H}^{l(w)}(\mathfrak{u}, \mathcal{M})[w \cdot \lambda]$  and an element  $H \in \mathfrak{a}$  we have

$$d(f \otimes u)(H) = Hf \otimes u - d(w \cdot \lambda(H)) \otimes u$$

and we have  $d\omega = 0 \iff f = cw \cdot \lambda$ , this means that

$$H^0(\Omega^\bullet(A_{\pi'}) \otimes \mathbb{H}^{l(w)}(\mathfrak{u}, \mathcal{M})[w \cdot \lambda]) = \mathbb{H}^{l(w)}(\mathfrak{u}, \mathcal{M})[w \cdot \lambda] \otimes \mathbb{C}w \cdot \lambda$$

as it should be. We consider the subspace

$$P_{\log}(A_{\pi'}) = \{f \in \mathcal{C}_\infty(A_{\pi'}) \mid f \text{ is a polynomial in } \log(x_\alpha), \alpha \in \pi'\}$$

and define

$$\Omega_{\log}^p(A_{\pi'}) = \left\{ \sum f_I \frac{dx_{\alpha_1}}{x_{\alpha_1}} \wedge \cdots \wedge \frac{dx_{\alpha_p}}{x_{\alpha_p}} \right\} \quad (8.46)$$

where  $I = \{\alpha_1, \dots, \alpha_p\}$  and  $f_I \in P_{\log}(A_{\pi'})$ . Observe that  $d \log(x_\alpha) = \frac{dx_\alpha}{x_\alpha}$  and hence it is clear that the inclusion

$$\Omega_{\log}^\bullet(A_{\pi'}) \otimes \mathbb{H}^{l(w)}(\mathfrak{u}, \mathcal{M})[w \cdot \lambda] \hookrightarrow \Omega^\bullet(A_{\pi'}) \otimes \mathbb{H}^{l(w)}(\mathfrak{u}, \mathcal{M})[w \cdot \lambda]$$

induces an isomorphism in cohomology. If we now define  $\Omega_{\log}^\bullet(\prod_{\alpha \in \pi'} (0, c_\alpha)) \otimes \mathbb{H}^{l(w)}(\mathfrak{u}, \mathcal{M})[w \cdot \lambda]$  to be the image of  $\Omega_{\log}^\bullet(A_{\pi'}) \otimes \mathbb{H}^{l(w)}(\mathfrak{u}, \mathcal{M})[w \cdot \lambda]$  under the

restriction then it is clear that

$$\begin{aligned} & \Omega^\bullet(\Gamma_M \backslash X^M(r(\underline{c}_P))) \otimes \Omega_{\log}^\bullet(\prod_{\alpha \in \pi'}(0, c_\alpha)) \otimes \mathbb{H}^{l(w)}(\mathbf{u}, \mathcal{M})[w \cdot \lambda] \hookrightarrow \\ & \Omega^\bullet(\Gamma_M \backslash X^M(r(\underline{c}_P))) \otimes \Omega^\bullet(\prod_{\alpha \in \pi'}(0, c_\alpha)) \otimes \mathbb{H}^{l(w)}(\mathbf{u}, \mathcal{M})[w \cdot \lambda] \quad (8.47) \\ & \hookrightarrow \Omega^{\bullet+\bullet+l(w)}(\Gamma_P \backslash X^P(r(c_P), c_P), \tilde{\mathcal{M}} \otimes \mathbb{C}) \end{aligned}$$

induces an isomorphism in cohomology.

We define a global subcomplex  $\Omega_{\log}^\bullet(\Gamma \backslash X) \otimes \mathcal{M}_{\mathbb{C}}$ , it consists of those forms whose restriction to  $\Gamma_P \backslash X^P(r(c_P), c_P)$  lie asymptotically in the log sub complex, this means for a suitable choice  $c'_P < c_P$  the restriction to  $\Gamma_P \backslash X^P(r(c_P), c'_P)$  lies in  $\Omega_{\log}^\bullet(\Gamma_P \backslash X^P, (r(c_P), c'_P)) \otimes \mathcal{M}_{\mathbb{C}}$ . Then

**Proposition 8.1.6.** *The inclusion*

$$\Omega_{\log}^\bullet(\Gamma \backslash X) \otimes \mathcal{M}_{\mathbb{C}} \hookrightarrow \Omega^\bullet(\Gamma \backslash X) \otimes \mathcal{M}_{\mathbb{C}}$$

*induces an isomorphism in cohomology.*

*Proof.* We pick a closed form  $\omega \in \Omega^p(\Gamma \backslash X) \otimes \mathcal{M}_{\mathbb{C}}$ , we restrict this form to the sets  $X^B(r_B, c_\pi)$  where  $B$  runs through a set of representatives of Borel subgroups (or more generally minimal parabolic subgroups). These sets are contained in slightly larger subsets  $X^B(r'_B, c'_\pi)$ . They are disjoint for different  $B$ . We can find a  $\mathcal{C}_\infty$  function  $h_P \in \mathcal{C}(\Gamma \backslash X)$  which is constant equal to one on  $X^B(r_B, c_\pi)$  and zero outside of the larger set  $X^B(r'_B, c'_\pi)$ . Now we can find a form  $\psi_P \in \Omega^{p-1}(\Gamma_B \backslash X^B(r_B, c_\pi)) \otimes \mathcal{M}_{\mathbb{C}}$  such that

$$\omega|_{X^B(r'_B, c'_\pi)} - d\psi \in \Omega_{\log}^p(X^B(r'_B, c'_\pi))$$

The form  $\omega_1 = \omega - d(h_P \psi)$  extends to  $\Gamma \backslash X$  and satisfies the condition for being asymptotically in  $\Omega_{\log}$  with respect to the subgroups  $B$ .

For a given Borel subgroup  $B$  we look at the different parabolic subgroups  $P \supset B$  whose rank drops by one, i.e. the next to minimal ones. We apply the same procedure to the restriction of  $\omega_1$  to  $X^P(r_P, c_P)$  and we get a form  $\omega_2 = \omega_1 - \sum_P d\psi_P$  whose restriction to the  $X^P(r_P, c_P)$  is asymptotically in  $\Omega_{\log}^\bullet$ . We have to be a little bit careful since we have modified  $\omega_1$  also on the  $X^B(r_B, c_B)$  but it is clear that also  $\omega_2$  restricted to  $X^B(r_B, c_B)$  is asymptotically in  $\Omega_{\log}^\bullet$ . This goes on and stops if we have reached the maximal parabolic subgroup  $G$  and then being in  $\Omega_{\log}$  becomes an empty condition.

This proves at least that the map in proposition 8.1.6 induces a surjective map in cohomology.  $\square$

We introduced this sub complex because now we can say something about the growth of cohomology classes or the asymptotic behavior. We consider this behavior on the different sets  $X^P((r(c_P), c_P))$ . The restriction of a form  $\omega \in \Omega_{\log}(\Gamma \backslash X) \otimes \mathcal{M}$  to  $\Gamma_P \backslash X^P((r(c_P), c_P))$  is asymptotically of the form  $\sum_{w \in W^P} \omega_w$ .

Let  $\omega = \omega_w \in \Omega^p(\Gamma_M \backslash X^M(r(\underline{c}_P))) \otimes \Omega_{\log}^i(\prod_{\alpha \in \pi'}(0, c_\alpha)) \otimes \mathbb{H}^{l(w)}(\mathbf{u}, \mathcal{M})[w \cdot \lambda]$ , we study the "growth" of the value of this form. We evaluate it at points  $x = (x_1, a) \in \Gamma_M \backslash X^M(r(\underline{c}_P)) \times \prod_{\alpha \in \pi'}(0, c_\alpha)$ , this means we pick tangent

vectors  $t_1^M, \dots, t_p^M$  at  $m_1$ . The tangent bundle on  $A_{\pi'}$  is trivialized by translation invariant vector fields. An  $i$ -tuple  $t_1^A, \dots, t_i^A \in \text{Lie}A_{\pi'}$  gives an  $i$ -tuple  $t_1^A, \dots, t_i^A$  of tangent vectors in the point  $a$ . Now we consider the value

$$\omega_w(\underline{T})(x) := \omega_w^M(x_1)(t_1^M, \dots, t_p^M)\omega_w^A(a)(t_1^A, \dots, t_i^A) \in \mathbb{H}^{l(w)}(\mathfrak{u}, \mathcal{M})[w \cdot \lambda].$$

We have a hermitian scalar product  $\langle, \rangle$  on  $\mathbb{H}^{l(w)}(\mathfrak{u}, \mathcal{M})[w \cdot \lambda]$  and we are interested in the value

$$\langle \omega_w(\underline{T})(x), \omega_w(\underline{T})(x) \rangle = \|\omega_w^M(x_1)(t_1^M, \dots, t_p^M)\|^2 \|\omega_w^A(a)(t_1^A, \dots, t_i^A)\|^2$$

The variable  $x_1$  runs through a compact set, hence value the first factor is bounded. The term  $\omega_w^A(a)(t_1^A, \dots, t_i^A) \in \mathbb{C}(w \cdot \lambda)P_{\log}(A_{\pi'})$  and this implies

$$\langle \omega_w(\underline{T})(x), \omega_w(\underline{T})(x) \rangle < C(2w \cdot \lambda)(a)a^{-\epsilon}$$

where  $\epsilon > 0$  and  $a^{-\epsilon} = \prod_{\alpha \in \pi'} x_{\alpha}^{-\epsilon}$ .

Now we can formulate a criterion to decide whether  $\omega_w$  is square integrable. We have to evaluate the integral

$$\int_{\Gamma_P \backslash X^P(r(c_P), c'_P)} \|\omega_w(\underline{p})\|^2 d\underline{p} \tag{8.48}$$

The measure  $d\underline{p}$  is of course the restriction of the invariant measure on  $\Gamma \backslash X$ , it is of the form  $2\rho_P(a)dud\underline{a}dm$ . The differential form is invariant under left translations under  $U(\mathbb{R})$  and hence we have to evaluate

$$\int_{\Gamma_M \backslash X^M(r(c_P) \times (\prod_{\alpha \in \pi'} (0, c_{\alpha}]))} \|\omega_w^M(\underline{m})\|^2 \|\omega_w^A(a)\|^2 (2\rho)(a) d\underline{m} da \tag{8.49}$$

The integral over  $\Gamma_M \backslash X^M(r(c_P))$  is finite hence we are left with

$$\int_{\prod_{\alpha \in \pi'} (0, c_{\alpha}]} \|\omega_w^A(a)\|^2 2\rho_P(a) da \tag{8.50}$$

Of course we assume that  $\omega_w \neq 0$  and then we can find a constant  $C > 0$  and an  $\epsilon > 0$  such that

$$C(2(w(\lambda + \rho) - \rho) + 2\rho_P)(a)a^{\epsilon} \leq \|\omega_w^A(a)\|^2 2\rho_P(a) \leq C(2(w(\lambda + \rho) - \rho) + 2\rho_P)(a)a^{-\epsilon} \tag{8.51}$$

Since  $\rho|_{A_{\pi'}} = \rho_P|_{A_{\pi'}}$  we get

$$C(2(w(\lambda + \rho)))(a)a^{\epsilon} \leq \|\omega_w^A(a)\|^2 2\rho_P(a) \leq C(2(w(\lambda + \rho)))(a)a^{-\epsilon}. \tag{8.52}$$

The relative roots  $\alpha^P$  form a basis for  $X^*(S)$ , any character  $\mu \in X^*(S)$  can be written as linear combination  $\mu = \sum_{\alpha \in \pi'} r_{\mu, \alpha} \alpha^P$  with  $r_{\mu, \alpha} \in \mathbb{Q}$ . We say that  $\mu$  is in the positive cone (with respect to the roots) if  $r_{\mu, \alpha} > 0$  for all  $\alpha \in \pi'$ , we write  $\mu >_P 0$ . Then

$$w(\lambda + \rho)(a) = w(\lambda + \rho)(\{ \dots, x_{\alpha}, \dots \}) = \prod_{\alpha \in \pi'} x_{\alpha}^{r_{\lambda + \rho, \alpha}}.$$

and come to the conclusion

**Proposition 8.1.7.** *a) The integral (8.50) is finite  $\iff w(\lambda + \rho) >_P 0$ .*

*b) A closed differential form  $\omega \in \Omega_{\log}^p(\Gamma \backslash X) \otimes \mathcal{M}$  is square integrable if and only if for all parabolic subgroups  $P$  and the resulting decompositions*

$$\omega|_{X^P}(r(c_P), c_P) = \sum_{w \in W^P} \omega_w$$

*the components  $\omega_w = 0$  if  $w(\lambda + \rho) \not>_P 0$ .*

We are now able to show

**Proposition 8.1.8.** *If  $\omega \in \Omega^p(\Gamma \backslash X) \otimes \mathcal{M}$  be a closed square integrable form and if the class  $[\omega] \in H_1^q(\Gamma \backslash X, \mathcal{M})$  then we can find a square integrable  $\psi \in \Omega^{p-1}(\Gamma \backslash X) \otimes \mathcal{M}$  such that  $\omega - d\psi$  has compact support*

*Proof.* We know that we can find a form  $\omega_1 \in \Omega_{\log}^p(\Gamma \backslash X) \otimes \mathcal{M}$  which represents the same class. Our previous arguments show that  $\omega_1$  is again square integrable.  $\square$

### 8.1.9 Franke's Theorem

The theorem 8.1.1 tells us that we have the very small sub complex

$$\text{Hom}_{K_\infty}(\Lambda^p(\mathfrak{g}/\mathfrak{k}), H_{\text{Coh}(\lambda)}^{(K_\infty)} \otimes \mathcal{M}_\lambda) \subset \text{Hom}_{K_\infty}(\Lambda^p(\mathfrak{g}/\mathfrak{k}), \mathcal{C}_\infty(\Gamma \backslash G(\mathbb{R})) \otimes \mathcal{M}_\lambda)$$

such that this induces a surjective map in cohomology

$$H^\bullet(\mathfrak{g}, K_\infty, H_{\text{Coh}(\lambda)}^{(K_\infty)} \otimes \mathcal{M}_\lambda) \xrightarrow{j_{(2)}} H_{(2)}^\bullet(\Gamma \backslash X, \mathcal{M}_\lambda),$$

if  $\Gamma \backslash X$  is not compact the map  $j_c$  is not necessarily an isomorphism, the kernel can be computed in principle by using Proposition 8.1.4.

By definition

$$H_{\text{Coh}(\lambda)}^{(K_\infty)} = \{f \in L_{\text{disc}}^2(\Gamma \backslash G(\mathbb{R})) \mid zf = \chi_\lambda(z)f \ \forall z \in \mathfrak{Z}(\mathfrak{g})\}.$$

A. Borel proposed to replace  $H_{\text{Coh}(\lambda)}^{(K_\infty)}$  by a larger space

$$\mathcal{A}_\lambda(\Gamma \backslash G(\mathbb{R})) := \{f \in \mathcal{C}_\infty(\Gamma \backslash G(\mathbb{R})) \mid \exists N \text{ such that } (z - \chi_\lambda(z))^N f = 0\} \quad (8.53)$$

where  $f$  also satisfies a growth condition. Borel conjectured the following theorem which was proved by Franke

**Theorem 8.1.4.** (Franke [17]) *The inclusion  $\mathcal{A}_\lambda(\Gamma \backslash G(\mathbb{R})) \subset \mathcal{C}_\infty(\Gamma \backslash G(\mathbb{R}))$  induces an isomorphism in cohomology*

$$H^\bullet(\mathfrak{g}, K_\infty, \mathcal{A}_\lambda(\Gamma \backslash G(\mathbb{R})) \otimes \mathcal{M}_\lambda) \xrightarrow{\sim} H^\bullet(\mathcal{S}_{K_f}^G, \mathcal{M}_\lambda)$$

The main tool for proving this theorem is again the theory of Eisenstein series. As in section 4.1.8 we start from certain induced  $\mathfrak{I}_P^G \sigma \times |\rho_P|^z$  where  $P$  runs over the conjugacy classes of parabolic subgroups,  $\sigma$  is a (cuspidal) cohomology class on locally symmetric space attached to the reductive quotient  $M/\mathbb{Q}$  of  $P$ , and  $z \in \mathbb{C}^r$ . We can write down Eisenstein series which yield an embedding

$$\text{Eis}(\cdot, z) : \mathfrak{I}_P^G \sigma \times |\rho_P|^z \hookrightarrow \mathcal{C}_\infty(G(\mathbb{Q})\mathbb{G}(\mathbb{A}))$$

these series are absolutely and locally uniformly converging if  $\Re(z - i) \gg 0$ . In [54] Langlands proves that these Eisenstein series have a meromorphic continuation into the entire  $\mathbb{C}^r$ . If we now "evaluate at  $z = 0$ " then we get functions in  $\mathcal{A}_\lambda(\Gamma \backslash G(\mathbb{R}))$ . But now the process of evaluation may become delicate, because the Eisenstein series may be singular at  $z = 0$ . So we may have to take residues and derivatives of such Eisenstein series which will give us the space  $\mathcal{A}_\lambda(\Gamma \backslash G(\mathbb{R}))$ .

## 8.2 Modular symbols

GP

### 8.2.1 The general pattern

We start from a flat group scheme  $\mathcal{G}/\text{Spec}(\mathbb{Z})$  whose generic fiber  $G/\mathbb{Q} = \mathcal{G} \times \mathbb{Q}$  is reductive. Let  $F/\mathbb{Q}$  be a finite normal extension, let  $\mathcal{O}_F$  be its ring of integers. We choose a highest weight, which is defined over  $F$  and consider a representation  $\rho_\lambda : \mathcal{G} \times \mathcal{O}_F \rightarrow \text{Gl}(\mathcal{M}_{\mathcal{O}_F})$  which after tensorization by  $F$  becomes the highest weight representation  $\mathcal{M}_{F,\lambda}$ . In the following we write  $\mathcal{M} = \mathcal{M}_{\mathcal{O}_F}$ , if we change the ring of scalars we write  $\mathcal{M}_R := \mathcal{M} \otimes_{\mathcal{O}_F} R$ . Let  $K_f^{(0)} = \mathcal{G}(\hat{\mathbb{Z}})$  and  $K_f \subset K_f^{(0)}$  be an open subgroup.

We want to describe a general method to construct homology classes in

$$H_d(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}) \text{ resp. relative homology groups } H_d(\mathcal{S}_{K_f}^G, \partial\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}),$$

which are obtained from (reductive) subgroups  $H \subset G$ , these classes will be the modular symbols. We want to put the considerations in Chapter 5 into a general framework.

Let  $H/\mathbb{Q}$  be a (reductive) subgroup of our ambient group  $G/\mathbb{Q}$ , we also consider the flat closure  $\mathcal{H}/\mathbb{Z}$ . We assume that its derived subgroup  $H^{(1)}$  is simply connected and satisfies strong approximation. The quotient  $H/H^{(1)} = C'$  is a torus. Let  $K_\infty^{H,(1)}$  be the connected component of the identity of a maximal compact subgroup of  $H(\mathbb{R})$  we put  $X^H = H(\mathbb{R})/K_\infty^{H,(1)}$ . We have the two spaces

$$\mathcal{S}_{K_f}^G = G(\mathbb{Q}) \backslash X \times G(\mathbb{A}_f)/K_f, \quad \mathcal{S}_{K_f^H}^H = H(\mathbb{Q}) \backslash X^H \times H(\mathbb{A}_f)/K_f.$$

and it follows from the considerations in section 6.1.3 that

$$\pi_0(\mathcal{S}_{K_f^H}^H) \xrightarrow{\sim} \pi_0(\mathcal{S}_{K_f^{C'}}^{C'}). \tag{8.54}$$

From the inclusion  $i : H \rightarrow G$  we will get maps between these locally symmetric spaces

$$j(x, \underline{g}_f) : \mathcal{S}_{K_f^H}^H \rightarrow \mathcal{S}_{K_f}^G$$

which depend on the choice of "pin points"  $(x, \underline{g}_f) \in X \times G(\mathbb{A}_f)$ . These pin points have to be chosen with some care:

a) The point  $x \in X$  can be viewed as a Cartan involution  $\Theta_x$  on  $G(\mathbb{R})$  and  $\Theta_x$  should fix  $H(\mathbb{R})$ . Hence it is also a Cartan involution on  $H$  and we require that it is the identity on our chosen  $K_\infty^{H,(1)}$ . Let us denote this subset of  $X$  by

$$X^{(H, K_\infty^{H,(1)})} = \{x \in X \mid \Theta_x(H(\mathbb{R})) = H(\mathbb{R}); \Theta_x = \text{identity on } K_\infty^{H,(1)}\}.$$

Let  $N$  be the subgroup of the normalizer of  $H/\mathbb{Q}$  which also normalizes  $K_\infty^{H,(1)}$ . Then  $N(\mathbb{R})$  acts on  $X^{(H, K_\infty^{H,(1)})}$ . I think that this action is transitive and the orbits under the group  $N(\mathbb{R})^{(1)}$  are the connected components.

b) The element  $\underline{g}_f$  has to satisfy a similar condition:

$$K_f^H \underline{g}_f K_f = \underline{g}_f K_f \quad (8.55)$$

we say that  $\underline{g}_f$  is adapted.

(Recall that we always have to make careful choices of the level once we deal with integral cohomology.) Such a pin point  $(x, \underline{g}_f)$  provides a map pinpoint

$$j(x, \underline{g}_f) : H(\mathbb{Q}) \backslash H(\mathbb{R}) / K_\infty^H \times H(\mathbb{A}_f) / K_f^H \longrightarrow \mathcal{S}_{K_f}^G \quad (8.56)$$

which is defined by

$$(h_\infty, \underline{h}_f) \mapsto (h_\infty x, \underline{h}_f \underline{g}_f).$$

We restrict this representation to  $H/\mathbb{Q}$  then we can decompose the rational module  $\mathcal{M}_\lambda \otimes F = \mathcal{M}_{\lambda F}^{\geq 1} \oplus \mathcal{M}_{\lambda F}^{H^{(1)}}$  where the first summand is the direct sum of irreducible modules of dimension  $> 1$  and the second summand is the module of  $H^{(1)}$  invariants. We define the module of  $\mathcal{H}^{(1)}$  coinvariants

$$\mathcal{M}_{\lambda, H^{(1)}} = \mathcal{M}_\lambda / \mathcal{M}_\lambda \cap \mathcal{M}_{\lambda F}^{\geq 1}.$$

This module of coinvariants is now a module for  $C'$  we assume that our field is large enough so that we can assume that  $C' \times \mathcal{O}_F$  is a split torus. Then we get that  $\mathcal{M}_{\lambda, H^{(1)}} = \bigoplus_{\mu \in X^*(C' \times \mathcal{O}_F)} \mathcal{M}_{\lambda, H^{(1)}}[\mu]$ . Then  $\mathcal{M}_{\lambda, H^{(1)}}[\mu]$  is a projective  $\mathcal{O}_F$  module of finite rank on which  $C' \times \mathcal{O}_F$  acts by the character  $\mu$ . We assume for simplicity that  $\mathcal{M}_{\lambda, H^{(1)}}[\mu]$  is actually free, hence we can write it as a direct sum of modules  $\mathcal{O}_F e_{\mu, j}$  where we chose a generator for each summand (in our examples this module is always of rank one). Let  $\mathcal{O}_\mu$  be the  $\mathcal{O}_F$ -module  $\mathcal{O}_F$  (with canonical generator 1) and with the action of  $C'$  by the character  $\mu$ , then of course  $\mathcal{O}_F e_{\mu, j} \xrightarrow{\sim} \mathcal{O}_\mu$ . Any  $C'$  homomorphism  $\phi_\mu : \mathcal{M}_{\lambda, H^{(1)}} \rightarrow \mathcal{O}_\mu$  provides a homomorphism of  $\mathcal{H}$  modules

$$\phi_\mu : \mathcal{M}_\lambda \rightarrow \mathcal{O}_\mu. \quad (8.57)$$

we denote it by the same letter. This induces a homomorphism of sheaves

$$\phi_\mu^* : j(x, \underline{g}_f)^*(\tilde{\mathcal{M}}_\lambda) \rightarrow \tilde{\mathcal{O}}_\mu. \quad (8.58)$$

especially any of the  $e_{\mu, j}$  gives us such a homomorphism.

Then these data provide a homomorphism for the cohomology groups

$$\phi_\mu \circ j(x, \underline{g}_f)^\bullet : H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}) \rightarrow H^\bullet(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{O}}_\mu).$$

We are interested in this homomorphism in degree  $d_H = \dim \mathcal{S}_{K_f^H}^H$ .

Let us assume for a moment that  $\mathcal{S}_{K_f^H}^H$  is compact. We have an orientation on  $\mathcal{S}_{K_f^H}^H$ , because we chose the compact subgroup  $K_\infty^H$  to be narrow. Therefore we see that the cohomology group  $H^{d_H}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{O}}_\mu)$  is the sum of cohomology groups over the connected components, and hence ( See8.62)

$$H^{d_H}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{O}}_\mu) \supset \bigoplus_{\tilde{\mu}_f} H^{d_H}(\mathcal{S}_{K_f^H}^H, (\tilde{\mathcal{O}}_\mu)[\tilde{\mu}_f]). \tag{8.59}$$

where we sum over characters  $\tilde{\mu}_f$  of type  $\mu$  on  $\tilde{C}'(\mathbb{A})/K_f^{C'}$  (See (6.3.6)). The eigenspaces are projective  $\mathcal{O}$ -modules of rank one let us assume that they are free and that we have chosen generators  $c_{\tilde{\mu}_f}$ . We will call such generators modular symbols.

We still have the variable  $\underline{g}_f$ , it has to satisfy the above condition b). We have to fix the level because we want to work with integral cohomology groups. But once we tensorize our coefficient systems with  $F$  ( the quotient field of  $\mathcal{O}$  ) then we can consider the limit

$$\lim_{K_f} H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F) = H^\bullet(\mathcal{S}^G, \tilde{\mathcal{M}}_F),$$

and this limit is now a  $\tilde{G}(\mathbb{A})$ - module (Section 6.3). Doing this also with  $\mathcal{S}_{K_f^H}^H$  we can forget the constraint on  $\underline{g}_f$ , the condition b) is certainly fulfilled for some choice of levels.

We recall the definition of an induced representation, we have

$$\text{Ind}_{\tilde{H}(\mathbb{A})}^{\tilde{G}(\mathbb{A})} H^{d_H}(\mathcal{S}^H, \tilde{F}_\mu) = \{ \Phi : \tilde{G}(\mathbb{A}) \rightarrow H^{d_H}(\mathcal{S}^H, \tilde{F}_\mu) \}$$

where  $\Phi$  satisfies  $\Phi(\underline{h}g) = \rho_{F_\mu}(\underline{h})\Phi(g)$  for all  $\underline{h} \in \tilde{H}(\mathbb{A}), g \in \tilde{G}(\mathbb{A})$  and where  $\Phi$  is right invariant under some open compact subgroup  $K_f^I$ . The map

$$J(\phi_\mu) : \xi \mapsto r_{\phi_\mu} \circ j(x, \underline{g}_f)(\xi) \tag{8.60}$$

yields an intertwining operator between  $\tilde{G}(\mathbb{A})$  modules Jphi

$$J(\phi_\mu) : H^{d_H}(\mathcal{S}^G, \tilde{\mathcal{M}}_F) \rightarrow \text{Ind}_{\tilde{H}(\mathbb{A})}^{\tilde{G}(\mathbb{A})} H^{d_H}(\mathcal{S}^H, \tilde{F}_\mu) \tag{8.61}$$

On the right hand side we can decompose further. We have seen that

$$H^{d_H}(\mathcal{S}^H, \tilde{F}_\mu) = \bigoplus_{\tilde{\mu}_f : \text{type}(\tilde{\mu}_f) = \mu} \text{Ind}_{\tilde{H}(\mathbb{A})}^{\tilde{G}(\mathbb{A})} H^{d_H}(\mathcal{S}^H, \tilde{F}_\mu)[\tilde{\mu}_f] = \bigoplus_{\tilde{\mu}_f : \text{type}(\tilde{\mu}_f) = \mu} F[\tilde{\mu}_f] \tag{8.62}$$

here we have to take into account that we have to enlarge our field  $F$  so that it contains the values of  $\tilde{\mu}_f(C'(\mathbb{A}))$ .

We project to the  $\tilde{\mu}_f$  component and get intertwining operators prchi

$$J(\phi_\mu, \tilde{\mu}_f) : H^{d_H}(\mathcal{S}^G, \tilde{\mathcal{M}}_F) \rightarrow \text{Ind}_{\tilde{H}(\mathbb{A})}^{\tilde{G}(\mathbb{A})} H^{d_H}(\mathcal{S}^H, \tilde{F}_\mu)(\tilde{\mu}_f) = \text{Ind}_{\tilde{H}(\mathbb{A})}^{\tilde{G}(\mathbb{A})} F \otimes \tilde{\mu}_f \quad (8.63)$$

Again the question arises to compute this intertwining operator. We have to explain what this means. At this point we only give a first approximation of what it means to compute this operator, better approximations come later.

Assume we have an absolutely irreducible  $\tilde{G}(\mathbb{A})$  "explicitly given" module  $V(\epsilon \times \pi_f)/F$ , here  $\epsilon \times \pi_f$  is a isomorphism type of an absolutely irreducible representation of  $\tilde{G}(\mathbb{A})$  on a  $F$ -vector space. The infinite component of such a representation is simply a character  $\epsilon : \pi_\infty(G(\mathbb{R})) \rightarrow \{\pm 1\}$ .

Now we also assume that we have an embedding  $\Phi(\pi_f) : V(\epsilon \times \pi_f) \hookrightarrow H^{d_H}(\mathcal{S}^G, \tilde{\mathcal{M}}_F)$ , i.e.  $V(\pi_f)$  is a  $F$ -model space (see further down). We also choose a character  $\tilde{\mu}_f$  of type  $\mu$ , we assume that the values of  $\tilde{\mu}_f$  are in  $F$ . Then  $H^{d_H}(\mathcal{S}^H, \tilde{F}_\mu)[\tilde{\mu}_f]$  is of rank one and our intertwining operator gives us an  $\tilde{G}(\mathbb{A})$ -module homomorphism

$$J(\phi_\mu, \tilde{\mu}_f) \circ \Phi(\pi_f) : V(\pi_\infty \times \pi_f) \rightarrow \text{Ind}_{\tilde{H}(\mathbb{A})}^{\tilde{G}(\mathbb{A})} \tilde{\mu}_f. \quad (8.64)$$

Now we will see further down that we encounter situations where the space of these intertwining operators is of dimension one. Moreover we will be able to identify an explicit non zero such operator  $I^{\text{loc}} : V(\epsilon \times \pi_f) \rightarrow \text{Ind}_{\tilde{H}(\mathbb{A})}^{\tilde{G}(\mathbb{A})} \tilde{\mu}_f$ . (See (8.162)) Then we get

$$J(\phi_\mu, \tilde{\mu}_f) \circ \Phi(\pi_f) = \mathcal{L}(\pi_f, \tilde{\mu}_f) I^{\text{loc}} \quad (8.65)$$

and computing the intertwining operator means to compute the number  $\mathcal{L}(\pi_f, \tilde{\mu}_f)$ . Since all our vector spaces on stage are defined over  $F$  we get the rationality result

$$\mathcal{L}(\pi_f, \tilde{\mu}_f) \in F \quad (8.66)$$

We will make this more precise later.

The passage to the to the limit has the technical advantage that we are dealing with representations of  $G(\mathbb{A}_f)$  instead of Hecke-modules, for the representations certain issues are easier to handle. Especially it is easier to compute dimensions of spaces of intertwining operators.

But now we going back to the case of a fixed level. We want to extend our considerations to the case that  $\mathcal{S}_{K_f^H}^H$  is not compact. In this case we study the extension of  $j(x, \underline{g}_f)$  to the compactification

$$\bar{j}(x, \underline{g}_f) : \bar{\mathcal{S}}_{K_f^H}^H \rightarrow \bar{\mathcal{S}}_{K_f}^G$$

We recall the construction of sheaves with intermediate support conditions (See(6.19)). Let us assume that we can find a  $\Sigma$  such that the image of  $\partial(\bar{\mathcal{S}}_{K_f^H}^H)$

factors through  $\partial_\Sigma(\mathcal{S}_{K_f}^G)$ . Then our homomorphism  $r$  together with a choice of a  $\phi_\mu$  yields a homomorphism between sheaves (see ( 6.19))

$$r_{\phi_\mu}^! : \bar{j}(x, \underline{g}_f)^*(i_{\Sigma,*,!}(\tilde{\mathcal{M}})) \rightarrow i_!(\tilde{\mathcal{O}}_\mu). \quad (8.67)$$

and hence we get a homomorphism in cohomology  $\boxed{\text{jxr}}$

$$(r_{\phi_\mu}^! \circ \bar{j}((x, \underline{g}_f)))^{d_H} : H^{d_H}(\mathcal{S}_{K_f}^G, i_{\Sigma,*,!}(\tilde{\mathcal{M}})) \rightarrow H^{d_H}(\mathcal{S}_{K_f}^H, i_!(\tilde{\mathcal{O}}_\mu)) \quad (8.68)$$

and now the left hand side is again of the form the composition  $J(\phi_\mu, \tilde{\mu}_f) \circ \Phi(\pi_f)$ . On the right hand side we can decompose again

$$H^{d_H}(\mathcal{S}_{K_f}^H, i_!(\tilde{\mathcal{O}}_\mu)) = H_c^{d_H}(\mathcal{S}_{K_f}^H, \tilde{\mathcal{O}}_\mu) \supset \bigoplus_{\tilde{\mu}_f} H_c^{d_H}(\mathcal{S}_{K_f}^H, \tilde{\mathcal{O}}_\mu)[\tilde{\mu}_f], \quad (8.69)$$

and the summands are locally free  $\mathcal{O}$  modules of rank one. If we project to the  $\tilde{\mu}_f$  component we get an operator

$$J(\phi_\mu, \tilde{\mu}_f) := P_{\tilde{\mu}_f} \circ (\phi_\mu \circ \bar{j}((x, \underline{g}_f)))^{d_H} : H^{d_H}(\mathcal{S}_{K_f}^G, i_{\Sigma,*,!}(\tilde{\mathcal{M}})) \rightarrow \text{Ind}_{\tilde{H}(\mathbb{A})}^{\tilde{G}(\mathbb{A})} \tilde{\mu}_f. \quad (8.70)$$

We are in the same situation as before, we have to find absolutely irreducible modules defined over  $F$  and an embedding

$$\Phi(\pi_f) : V(\epsilon \times \pi_f) \hookrightarrow H^{d_H}(\mathcal{S}_{K_f}^G, i_{\Sigma,*,!}(\tilde{\mathcal{M}}_F)).$$

and then we can try again to investigate the operator  $J(\phi_\mu, \tilde{\mu}_f) \circ \Phi(\pi_f)$

Here we have to discuss a subtle point. Let us consider the case that  $\Sigma$  is the set of all maximal parabolic subgroups, then  $H^{d_H}(\mathcal{S}_{K_f}^G, i_{\Sigma,*,!}(\tilde{\mathcal{M}}_F)) = H_c^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F)$ . We have the exact sequence

$$H^{d_H-1}(\partial(\mathcal{S}_{K_f}^G), \tilde{\mathcal{M}}_F) \rightarrow H_c^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F) \rightarrow H_1^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F) \rightarrow 0.$$

Sometimes it is easier to construct homomorphisms  $\Phi : V(\epsilon \times \pi_f) \hookrightarrow H_1^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F)$ . and we would like to form again in a canonical way the composition  $(r_{\phi_\mu}^! \circ \bar{j}((x, \underline{g}_f)))^{d_H} \circ \Phi$ .

We have two different instances, when this is possible. We look at the homology, then we have the boundary map

$$H_{d_H}(\mathcal{S}_{K_f}^H, \partial(\mathcal{S}_{K_f}^H), \underline{F}_\mu) \xrightarrow{\partial} H_{d_H-1}(\partial(\mathcal{S}_{K_f}^H), \underline{F}_\mu) \quad (8.71)$$

and from the target we have map

$$H_{d_H-1}(\partial(\mathcal{S}_{K_f}^H), \underline{F}_\mu) \xrightarrow{j} H_{d_H-1}(\partial(\mathcal{S}_{K_f}^G), \underline{\mathcal{M}}_F) \quad (8.72)$$

Hence we get a map

$$j \circ \partial_{\tilde{\mu}_f} : H_{d_H}(\mathcal{S}_{K_f}^H, \partial(\mathcal{S}_{K_f}^H), \underline{F}_\mu)[\tilde{\mu}_f] \rightarrow H_{d_H-1}(\partial(\mathcal{S}_{K_f}^G), \underline{\mathcal{M}}_F) \quad (8.73)$$

Now we have the following

**Proposition 8.2.1.** *If the map  $j \circ \partial_{\tilde{\mu}_f} = 0$  then the homomorphism  $J(\phi_\mu, \pi_f, \tilde{\mu}_f)$  vanishes on  $H^{d_H-1}(\partial(\mathcal{S}_{K_f}^G), \tilde{\mathcal{M}}_F)$  and hence it factors over the quotient*

$$J(\phi_\mu, \tilde{\mu}_f) : H_!^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F) \rightarrow H_!^{d_H}(\mathcal{S}_{K_f}^H, \tilde{F}_\mu)[\tilde{\mu}_f]$$

The second instance that that may be satisfied is the Manin-Drinfeld principle applies to the exact sequence above, i.e. we have an isotypical decomposition

$$H_{\text{Eis}}^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F) \oplus H_!^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F). \quad (M_2)$$

Then we may restrict  $J(\phi_\mu, \tilde{\mu}_f)$  to the second summand. We get

$$J_!(\phi_\mu, \tilde{\mu}_f) : H_!^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F) \longrightarrow \text{Ind}_{\tilde{H}(\mathbb{A}_f)}^{\tilde{G}(\mathbb{A}_f)} \tilde{\mu}_f,$$

and this intertwining operator is defined over  $F$ .

We used both arguments already in section 5.6 in a very special case.

## 8.2.2 Model spaces

We want to find the modules  $V(\pi_\infty \times \pi_f)$  and operators  $\Phi(\pi_f)$ . We introduce some abstract concept of the production of cohomology classes and the evaluation of these intertwining operators on these classes. For this purpose we introduce model spaces.

We assume that we have a family of smooth and admissible representations  $\{X_{\pi_v}\}$  of  $G(\mathbb{Q}_v)$  where  $v$  runs over all places. At this moment the  $X_{\pi_v}$  are  $\mathbb{C}$ -vector spaces. For almost all finite places  $p$  the representation  $\{X_{\pi_p}\}$  should be an unramified irreducible principal series representation. We assume that  $X_{\pi_\infty}$  is an irreducible Harish-Chandra module with non trivial cohomology  $H^\bullet(\mathfrak{g}, K_\infty, X_{\pi_\infty} \otimes \mathcal{M}_{\mathbb{C}}) \neq 0$ . We denote  $\pi = \pi_\infty \times \pi_f$ . Furthermore we assume that we have an intertwining operator of  $(\mathfrak{g}, K_\infty) \times G(\mathbb{A}_f)$ -modules

$$\Psi(\pi) : X_{\pi_\infty} \otimes \bigotimes_p X_{\pi_p} \longrightarrow \mathcal{C}_\infty(G(\mathbb{Q}) \backslash G(\mathbb{A})). \quad (8.74)$$

At this point a comment is in order. We should think of the spaces  $X_{\pi_v}$  as very specific spaces of  $\mathbb{C}$  valued functions on  $G(\mathbb{Q}_v)$  on which  $G(\mathbb{Q}_v)$  acts by right translations. In all cases known to the author the operator  $\Psi(\pi)$  is given by an infinite summation, i.e if  $f = \prod f_v \in X_{\pi_\infty} \otimes \bigotimes_p X_{\pi_p}$  then

$$\Psi(\pi)(f)(g) : \sum_{a \in H(\mathbb{Q})} f(ag) \quad (8.75)$$

where for instance  $H/\mathbb{Q}$  is a subgroup or a quotient of a subgroup by another subgroup. In any case it is clear that the construction of these  $\Psi(\pi)$  will be a transcendental process.

This induces of course an intertwining operator  $\Psi(\pi)$

$$\begin{aligned} H^\bullet(\mathfrak{g}, K_\infty, X_{\pi_\infty} \otimes \mathcal{M}_{\mathbb{C}}) \otimes \bigotimes_p X_{\pi_p} &\xrightarrow{\Psi^\bullet(\pi)} H^\bullet(\mathfrak{g}, K_\infty, \mathcal{C}_\infty(G(\mathbb{Q}) \backslash G(\mathbb{A})) \otimes \mathcal{M}_{\mathbb{C}}) \\ &= H^\bullet(\mathcal{S}^G, \tilde{\mathcal{M}}_{\mathbb{C}}) \end{aligned}$$

We introduce a subspace of  $\mathcal{C}_\infty(G(\mathbb{Q})\backslash G(\mathbb{A}))$ . We consider the subspace of functions of moderate growth and inside this space we consider the space of functions which are cuspidal along the strata  $\partial_P(\mathcal{S}^G)$  for the parabolic subgroups  $P \in \Sigma$ , i.e. which satisfy

$$\int_{U_P(\mathbb{Q})\backslash U_P(\mathbb{A})} f(\underline{u}g) d\underline{u} \equiv 0$$

for these parabolic subgroups. Let us call this subspace  $\mathcal{C}_\infty^{(\Sigma)}(G(\mathbb{Q})\backslash G(\mathbb{A}))$ . We assume that our intertwining operator factors through the subspace of  $\Sigma$  cuspidal functions

$$\Psi(\pi) : X_{\pi_\infty} \otimes \bigotimes_p X_{\pi_p} \longrightarrow \mathcal{C}_\infty^{(\Sigma)}(G(\mathbb{Q})\backslash G(\mathbb{A})). \tag{8.76}$$

We have an action of  $\pi_0(G(\mathbb{R}))$  on  $H^\bullet(\mathfrak{g}, K_\infty, X_{\pi_\infty} \otimes \mathcal{M}_\mathbb{C})$  let  $\epsilon : \pi_0(G(\mathbb{R})) \rightarrow \{\pm 1\}$  be a character and let  $\omega_\epsilon$  be a differential form representing an eigenclass  $[\omega_\epsilon]$ . In [Ha-Gl2] we explain how a Hecke character  $\tilde{\mu}_f$  extends uniquely to a character  $\tilde{\mu}_f^{-1} = \epsilon \times \tilde{\mu}_f : \pi_0(H(\mathbb{R}))H(\mathbb{A}_f) \rightarrow \{\pm 1\}$ . We have the homomorphism  $\pi_0(H(\mathbb{R})) \rightarrow \pi_0(G(\mathbb{R}))$  and we require that  $\chi_\infty = \epsilon$ .

We get a diagram

$$\begin{array}{ccc} H^{d_H}(\mathfrak{g}, K_\infty, X_{\pi_\infty} \otimes \mathcal{M}_\mathbb{C})(\epsilon) \otimes \bigotimes_p X_{\pi_p} & & \\ \downarrow \Psi(\pi)^{d_H} & & \\ H^{d_H}(\mathfrak{g}, K_\infty, \mathcal{C}_\infty^{(\Sigma)}(G(\mathbb{Q})\backslash G(\mathbb{A})) \otimes \mathcal{M}_\mathbb{C}) & \xrightarrow{\text{dRh}} & H^{d_H}(\mathcal{S}^G, \tilde{\mathcal{M}} \otimes \mathbb{C}) \\ & & \uparrow i_\Sigma^{d_H} \otimes \mathbb{C} \\ \text{Ind}_{\tilde{H}(\mathbb{A})}^{\tilde{G}(\mathbb{A})} \tilde{\mu}_f^{-1} \otimes \mathbb{C} & \xrightarrow{J(\phi_\mu, \tilde{\mu}_f)} & H^{d_H}(\mathcal{S}^G, i_{\Sigma, *, !}(\tilde{\mathcal{M}})) \otimes \mathbb{C} \end{array}$$

**Proposition 8.2.2.** *The image of dRh is contained in the image of  $i_\Sigma^{d_H} \otimes \mathbb{C}$*

*Proof.* We do not give the proof of this general assertion here, it is a careful analysis using reduction theory and the considerations in ???. We simply mention the case of a compact  $\mathcal{S}_{K_f^H}^H$  then we may choose  $\Sigma = \emptyset$  to be the set of all maximal parabolic subgroups and  $i_\Sigma^{d_H} \otimes \mathbb{C}$  is the identity and hence the proposition is obvious in this case. On the other hand if  $\Sigma$  is the set of all maximal parabolic subgroups then the image of  $i_\Sigma^{d_H} \otimes \mathbb{C}$  is the inner cohomology and since in this the functions in  $\mathcal{C}_\infty^{(\Sigma)}(G(\mathbb{Q})\backslash G(\mathbb{A}))$  are cuspidal the assertion follows from Theorem 8.1.1. □

We put  $H_{\Sigma, !}^{d_H}(\mathcal{S}^G, \tilde{\mathcal{M}}) = i_\Sigma^{d_H}(H^{d_H}(\mathcal{S}^G, i_{\Sigma, *, !}(\tilde{\mathcal{M}})))$  and we assume that one of the conditions above is satisfied, i.e. either we can apply proposition 8.2.1 or we have Manin-Drinfeld. We have the action of the  $\pi_0(G(\mathbb{R}))$  on  $H^\bullet(\mathfrak{g}, K_\infty, X_{\pi_\infty} \otimes \tilde{\mathcal{M}} \otimes \mathbb{C})$ , we decompose into eigenspaces according to characters  $\epsilon$ .

We get an arrow

$$H^\bullet(\mathfrak{g}, K_\infty, X_{\pi_\infty} \otimes \mathcal{M}_\mathbb{C})(\epsilon) \otimes \bigotimes_p X_{\pi_p} \xrightarrow{J(\phi_\mu, \tilde{\mu}_f) \circ \text{dRh} \circ \Psi^{d_H}(\epsilon \times \pi)} \text{Ind}_{\tilde{H}(\mathbb{A})}^{\tilde{G}(\mathbb{A})} \tilde{\mu}_f^{-1} \otimes \mathbb{C}. \tag{8.77}$$

We choose an element  $\omega_\epsilon \in \text{Hom}_{K_\infty}(\Lambda^{d_H}(\mathfrak{g}/\mathfrak{k}), X_{\pi_\infty} \otimes \mathcal{M}_{\mathbb{C}})[\epsilon]$  then this provides a homomorphism of  $G(\mathbb{A}_f)$ -modules

$$J(\phi_\mu, \tilde{\mu}_f) \circ \Psi^{d_H}(\omega_\epsilon \times \pi_f) : \bigotimes_p X_{\pi_p} \rightarrow \text{Ind}_{H(\mathbb{A})}^{\tilde{G}(\mathbb{A})} \tilde{\mu}_f^{-1} \otimes \mathbb{C} \quad (8.78)$$

For an element  $\psi_f \in \bigotimes_p X_{\pi_p}$  this map is given by the formula

intcyc0

$$J(\phi_\mu, \tilde{\mu}_f) \circ \Psi^{d_H}(\omega_\epsilon \times \pi_f)(\psi_f)(\underline{g}_f) = \int_{S_{K_f^H}^H} \phi_\mu(j^*(x, \underline{h}_f \underline{g}_f)(\omega_\epsilon \times \psi_f)) \tilde{\mu}_f(\underline{h}_f) d\underline{h}_f, \quad (8.79)$$

here  $d\underline{h}_f$  is the invariant measure on  $H(\mathbb{A}_f)$  which has value one on  $K_f^H$ .

We still have the problem to compute this operator. But now the situation has changed, we can be a little bit more precise in formulating what we mean by computing this operator. The source and the target of the operator

$$J(\phi_\mu, \pi_f, \tilde{\mu}_f, \omega_\epsilon) := J(\phi_\mu, \tilde{\mu}_f) \circ \Psi^{d_H}(\omega_\epsilon \times \pi_f) \quad (8.80)$$

are restricted tensor products of local representations. So a necessary condition for  $J(\phi_\mu, \pi_f, \tilde{\mu}_f, \omega_\epsilon) \neq 0$  is that for all primes  $p$  the vector space

$$\text{Hom}_{G(\mathbb{Q}_p)}(X_{\pi_p}, \text{Ind}_{H(\mathbb{Q}_p)}^{G(\mathbb{Q}_p)} \tilde{\mu}_p^{-1}) \neq 0. \quad (I_p)$$

Therefore we assume that this condition is fulfilled.

local condition  $(I_p)$  is satisfied for all primes  $p$ , then there are many interesting special cases where

$$\dim \text{Hom}_{G(\mathbb{Q}_p)}(X_{\pi_p}, \text{Ind}_{H(\mathbb{Q}_p)}^{G(\mathbb{Q}_p)} \tilde{\mu}_p^{-1}) = 1 \quad (I_{pp})$$

Therefore we assume that the representations  $X_{\pi_p}$  are somehow given to us as very concrete representations and  $(I_{pp})$  is true for all primes  $p$ . Moreover we assume at each prime  $p$  we see some natural choice of a generator

$$I_{\tilde{\mu}_p}^{\text{loc}} \in \text{Hom}_{G(\mathbb{Q}_p)}(X_{\pi_p}, \text{Ind}_{H(\mathbb{Q}_p)}^{G(\mathbb{Q}_p)} \tilde{\mu}_p^{-1})$$

(This will be discussed in our examples.) We can define a local intertwining operator

$$I_{\tilde{\mu}_f}^{\text{loc}} = \bigotimes_p I_{\tilde{\mu}_p}^{\text{loc}} \in \text{Hom}_{G(\mathbb{A}_f)}(\bigotimes_p X_{\pi_p}, \text{Ind}_{H(\mathbb{A}_f)}^{G(\mathbb{A}_f)} \tilde{\mu}_f^{-1}) \quad (8.81)$$

and now we can formulate the following basic question:

*The operator  $J(\phi_\mu, \pi_f, \tilde{\mu}_f, \omega_\epsilon)$  is a multiple of  $I_{\tilde{\mu}_f}^{\text{loc}}$  and the problem of computing this intertwining operator comes down to compute a number namely the proportionality factor in*

$$J(\phi_\mu, \pi_f, \tilde{\mu}_f, \omega_\epsilon) = \mathcal{L}(\pi_f, \tilde{\mu}) \cdot I_{\tilde{\mu}_f}^{\text{loc}}. \quad \text{bquest}$$

The general philosophy says that this proportionality factor should be obtained from the data  $\pi_f, \tilde{\mu}_f$  for instance it should be essentially a special value of an  $L$ -function attached to  $\pi_f = \bigotimes_p \pi_p$ . We will see in the examples in the section below that this is indeed sometimes the case.under

### 8.2.3 Rationality and integrality results

Now go back to the situation where we fix a finite level  $K_f$ , we also assume that  $\Sigma$  is the set of all maximal parabolic subgroups, and we assume that proposition ??? applies (this depends on the choice of  $\mu$ . see (??)). Hence we have the map

$$\phi_\mu \circ \bar{j}((x, \underline{g}_f))^{d_H} : H_!^{d_H} \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F \rightarrow H_!^{d_H} (\mathcal{S}_{K_f}^H, \tilde{F}_\mu) \quad (8.82)$$

We assume that our finite extension  $F/\mathbb{Q}$  is large enough so that we get an isotypical decomposition

$$H_!^{d_H} (\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F) = \bigoplus_{\pi_f} H_!^{d_H} \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F(\pi_f)$$

where the  $\pi_f$  are isomorphism types of absolutely irreducible modules for the Hecke algebra. Of course we may also require that  $F/\mathbb{Q}$  is normal.

We intersect  $H_!^{d_H} (\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})_F(\pi_f)$  with the integral cohomology  $H^{d_H} (\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathcal{O}_F})$  and get the submodule  $H_!^{d_H} (\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F)(\pi_f)_{\text{int}} \subset H^{d_H} (\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})_F(\pi_f)_{\text{int}}$ . We have seen in ??? that we may alternatively define the submodule

$$H^{d_H} (\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathcal{O}_F})(\pi_f)_{\text{int},!} \subset H^{d_H} (\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathcal{O}_F})_{\text{int}} \quad (8.83)$$

and we recall that the quotient

$$H^{d_H} (\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathcal{O}_F})(\pi_f)_{\text{int},!} / H_!^{d_H} (\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathcal{O}_F})(\pi_f)_{\text{int}} = \mathcal{T}(\pi_f) \quad (8.84)$$

is a torsion module which is isomorphic to a sub quotient of the torsion module of  $H^{d_H} (\partial(\mathcal{S}_{K_f}^G), \tilde{\mathcal{M}}_{\mathcal{O}_F})$ .

On our isotypical subspace we still have the action of  $\pi_0(G(\mathbb{R}))$  which commutes with the action of the Hecke algebra. Since this group of connected components is an elementary abelian 2 group we get a decomposition

$$H_!^{d_H} (\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F)(\pi_f) = \bigoplus_{\epsilon} H_!^{d_H} (\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F)(\epsilon \times \pi_f)$$

where  $\epsilon$  runs over the characters of  $\epsilon : \pi_0(G(\mathbb{R})) \rightarrow \{\pm 1\}$

We now make the assumption that  $(\epsilon \times \pi_f)$  occurs with multiplicity one, or in other words  $H_!^{d_H} (\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F)(\epsilon, \pi_f) \neq 0$  ( resp.  $H_!^{d_H} (\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_F)(\epsilon, \pi_f) \neq 0$ ) and it is an irreducible module for the Hecke algebra (resp. for  $G(\mathbb{A}_F)$ .)

We also assume that all the local components of our model space  $X_{\pi_p}$  are also defined over  $F$ . For a finite place  $p$  this means that  $X_{\pi_p}$  is a vector space over  $F$  with an action of  $G(\mathbb{Q}_p)$ . If we choose an open compact subgroup  $K_p \subset G(\mathbb{Q}_p)$  then  $X_{\pi_p}^{K_p}$  is a finite dimensional  $F$  vector space with an action of the Hecke algebra  $\mathcal{H}_{K_p}$  on it. This action is absolutely irreducible if  $\pi_p$  is absolutely irreducible. If our underlying flat group scheme  $\mathcal{G}/\mathbb{Z}$  is reductive at the prime  $p$  and  $K_p = \mathcal{G}(\mathbb{Z}_p)$  (some people say that  $K_p$  is hyperspecial) then  $X_{\pi_p}^{K_p}$  is of dimension one (or zero). The Hecke algebra module is given by a homomorphism  $h \mapsto \pi_p(h)$  from  $\mathcal{H}_{K_p} \rightarrow F$ .

We discuss the concept of rationality for  $X_{\pi_\infty}$ . Our group  $G/\mathbb{Q}$  is defined over  $\mathbb{Q}$  we choose the Cartan-involution  $\Theta$  which provides  $K_\infty$  also defined over

$\mathbb{Q}$ . Hence the Lie algebras  $\mathfrak{g}$  and  $\mathfrak{k}$  are defined over  $\mathbb{Q}$ , i.e. they are  $\mathbb{Q}$ -vector spaces. Our module  $\mathcal{M}$  is obtained from an absolutely irreducible highest weight representation  $\rho_\lambda : G \times F \rightarrow \mathrm{Gl}(\mathcal{M}_F)$ , If we choose a basis  $\{\dots, f_i, \dots\}$  and  $a \in G(\mathbb{Q})$  then  $\rho_\lambda(a)f_i = \sum a_{i,j}f_j$  with  $a_{i,j} \in F$ . The same applies for the action of  $\mathfrak{g}$  on  $X_{\pi_\infty}$ , this module has a countable basis  $\{\dots, g_i, \dots\}$  and for  $X \in \mathfrak{g}$  we get again  $Xf_i = \sum a_{i,j}f_j$ ,  $a_{i,j} \in F$  and where the sum is finite, i.e. only finitely many  $a_{i,j} \neq 0$ .

Then it is clear that for any  $\sigma \in \mathrm{Gal}(F/\mathbb{Q})$  we can define the conjugate  $(\mathfrak{g}, K_\infty)$ -module  $X_{\pi_\infty}$  and the conjugate  $\mathcal{G}$  module  $\mathcal{M}$ . In this sense we can say that the system  $(X_{\pi_\infty}, \mathcal{M})$  is defined over  $\mathbb{Q}$ . We also get the system of conjugate cohomology groups  $\{\dots, H_!^{d_H}(\mathcal{S}^G, \tilde{\mathcal{M}}_F)(\epsilon \times \pi_f), \dots\}_{\sigma \in \mathrm{Gal}(F/\mathbb{Q})}$

We assume that  $F \subset \mathbb{C}$ , and assume that we have the intertwining operator

$$\Psi^{d_H}(\omega_\epsilon) : \left( \bigotimes_p X_{\pi_p} \right) \otimes \mathbb{C} \rightarrow H_!^{d_H}(\mathcal{S}^G, \tilde{\mathcal{M}}_F)(\epsilon \times \pi_f) \otimes \mathbb{C} \quad (8.85)$$

these are isomorphisms over  $\mathbb{C}$  between absolutely irreducible  $G(\mathbb{A}_f)$  modules which are defined over  $F$ . Hence we can find numbers (the periods)  $\Omega(\epsilon \times \pi_f) \in \mathbb{C}^\times$  such that

periodrat

$$\Phi^{d_H}(\omega_\epsilon \times \pi_f) = \frac{\Psi^{d_H}(\omega_\epsilon \times \pi_f)}{\Omega(\epsilon \times \pi_f)} : \bigotimes_p X_{\pi_p} \xrightarrow{\sim} H_!^{d_H}(\mathcal{S}^G, \tilde{\mathcal{M}}_F)(\epsilon \times \pi_f) \quad (8.86)$$

is an isomorphism over  $F$ .

But we can do better. We may also assume that after fixing a level we have an integral structure on our model space, i.e we have chosen lattices  $X_{\pi_p, \mathcal{O}_F}^{K_p} \subset X_{\pi_p}^{K_p}$ . For almost all  $p$  this lattice is of rank one and  $X_{\pi_f, \mathcal{O}_F}^{K_f} = \bigotimes_p X_{\pi_p, \mathcal{O}_F}^{K_p}$  is a (locally) free module of finite rank. We require that our periods satisfy

$$\frac{\Psi^{d_H}(\omega_\epsilon \times \pi_f)}{\Omega(\epsilon \times \pi_f)} : \left( \bigotimes_p X_{\pi_p, \mathcal{O}_F}^{K_p} \right) \rightarrow H_!^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathcal{O}_F})(\epsilon \times \pi_f)_{\mathrm{int}} \quad (8.87)$$

and that this choice of periods is optimal, i.e. if  $r \in F$  and  $r \frac{\Psi^{d_H}(\omega_\epsilon \times \pi_f)}{\Omega(\epsilon \times \pi_f)} \left( \bigotimes_p X_{\pi_p, \mathcal{O}_F}^{K_p} \right) \subset H_!^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathcal{O}_F})(\epsilon \times \pi_f)_{\mathrm{int}}$  then  $r \in \mathcal{O}_F$ . This pins down the periods up to an element in  $\mathcal{O}_F^\times$ .

If now proposition 8.2.1 applies we can then we can form the composition

$$J(\phi_\mu, \chi_f) \circ \frac{\Psi^{d_H}(\omega_\epsilon \times \pi_f)}{\Omega(\epsilon \times \pi_f)} : \bigotimes_p X_{\pi_p, \mathcal{O}_F}^{K_p} \rightarrow \mathrm{Ind}_{\tilde{H}(\mathbb{A})}^{\tilde{G}(\mathbb{A})} \tilde{\mu}_f^{-1}.$$

Now we assume that the condition  $(I_{pp})$  is satisfied and we assume that our local intertwining operators  $I_p^{\mathrm{loc}}$  are defined over  $F$ . We define as above  $I_{\tilde{\mu}_f}^{\mathrm{loc}} = \otimes I_{\tilde{\mu}_p}$  and again we get a formula

$$J(\phi_\mu, \tilde{\mu}_f) \circ \frac{\Psi^{d_H}(\omega_\epsilon \times \pi_f)}{\Omega(\epsilon \times \pi_f)} = \mathcal{L}(\pi \otimes \chi, \mu) I_{\tilde{\mu}_f}^{\mathrm{loc}} \quad (8.88)$$

Of course there is still the unknown quantity  $\mathcal{L}(\pi \otimes \chi, \tilde{\mu})$  but we can say

**Proposition 8.2.3.** *If proposition 8.2.1 applies or if we have Manin-Drinfeld then  $\mathcal{L}(\pi \otimes \chi, \tilde{\mu}) \in F$*

But we want to do better. On the left hand side we have the integral structure and if we evaluate at an adapted argument  $\underline{g}_f$ , i.e.  $\underline{g}_f$  satisfies (8.55)

then we get for  $\psi_f \in X_{\pi_f, \mathcal{O}_F}^{K_f}$  Spv1

$$P_{\chi_f} \circ (r_{\phi_\mu}^! \circ \bar{j}((x, \underline{g}_f)))^{d_H} \circ \frac{\Psi^{d_H}(\omega_\epsilon \times \pi_f)}{\Omega(\epsilon \times \pi_f)}(\psi_f) = \mathcal{L}(\pi \otimes \chi, \mu) I_{\tilde{\mu}_f}^{\text{loc}}(\psi_f)(\underline{g}_f) \tag{8.89}$$

If we can apply proposition 8.2.1 then the left hand side is an integer in  $\mathcal{O}_F$  hence we know that the right hand side  $\mathcal{L}(\pi \otimes \chi, \mu) I_{\chi_f}^{\text{loc}}(\psi_f)(\underline{g}_f)$  is also an integer. To get information about the denominator of  $\mathcal{L}(\pi \otimes \tilde{\mu}_f, \mu)$  we have to optimize the numerator of  $I_{\tilde{\mu}_f}^{\text{loc}}(\psi_f)(\underline{g}_f)$ .

We have to choose  $\psi_f \in \bigotimes_p X_{\mathcal{O}_F}^{K_p}$ , and we choose  $\underline{g}_f$  such that  $K_f^H \underline{g}_f K_f = \underline{g}_f K_f$ ). The first choice provides an integral cohomology class in  $H^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathcal{O}_F})(\pi_f)$ . But this class is not necessarily the image of an integral class under  $r_{\Sigma}$ , this will be the case if we multiply it with  $\Delta(\pi_f)$ . Once we have done this we get that Spv2

$$j((x, \underline{g}_f), r_{\lambda, \mu})(\Phi^{d_H}(\frac{[\omega_\epsilon]}{\Omega(\pi_f, \omega_\epsilon)} \times \Delta(\pi_f)\psi_f)) = \Delta(\pi_f)\mathcal{L}(\pi \otimes \tilde{\mu}) I_{\tilde{\mu}_f}^{\text{loc}}(\psi_f)(\underline{g}_f) c_{\tilde{\mu}_f} \tag{8.90}$$

is a number in  $\mathcal{O}_F$ .

Then we have to optimize the choice of  $\underline{g}_f$ , this means that we have to keep the numerator of  $I_{\tilde{\mu}_f}^{\text{loc}}(\psi_f)(\underline{g}_f)$  small. Then we get an integrality result for the  $L$ -value.

### 8.2.4 The special case $\text{Gl}_2/F_0$

Let  $F_0/\mathbb{Q}$  be an algebraic number field, we consider the algebraic group  $G/\mathbb{Q} = R_{F_0/\mathbb{Q}}(\text{Gl}_2/F_0)$ . We embed  $\mathbb{G}_m \times \mathbb{G}_m = T_0$  into  $\text{Gl}_2/F_0$  by

$$(t_1, t_2) \mapsto \begin{pmatrix} t_1 & 0 \\ 0 & t_2 \end{pmatrix}$$

the diagonal  $\mathbb{G}_m \subset \mathbb{G}_m \times \mathbb{G}_m$  maps to the center  $C_0/F$ . Let  $B_0 \supset T_0$  be the standard Borel subgroup of upper triangular matrices and  $U_0$  its unipotent radical. Then  $T/\mathbb{Q} = R_{F/\mathbb{Q}}(T_0)$ ,  $B/\mathbb{Q} = R_{F/\mathbb{Q}}(B_0)U/\mathbb{Q} = R_{F/\mathbb{Q}}(U_0)B/\mathbb{Q} = R_{F/\mathbb{Q}}(U_0)$  and  $C/\mathbb{Q} = R_{F/\mathbb{Q}}(C_0)$

We want to apply our above considerations to the case  $H = T$  and to the case  $H = \text{Gl}_2 \subset \text{Gl}_2 \times \text{Gl}_2$ , diagonally embedded.

#### The spaces

Let  $\Sigma$  be the set of embeddings  $\iota : F_0 \hookrightarrow \mathbb{C}$ , on this set we have the action of complex conjugation  $\mathbf{c}$ . The set of embeddings  $\iota : F_0 \rightarrow \mathbb{R}$  is the set of

elements the fixed under conjugation, this is also the set of real places. The other embeddings come in pairs  $\iota, c\iota$ . Let  $S_\infty$  be the set of equivalence classes under this action, then  $S_\infty$  is the set of archimedean places of  $F_0$  and

$$F_0 \otimes \mathbb{R} = \prod_{v \in S_\infty} F_v = \mathbb{R}^{r_1} \times \mathbb{C}^{r_2}$$

of course  $F_v = \mathbb{R}$  resp.  $\mathbb{C}$ , if  $v$  is a real (resp. complex) place, Hence  $G(\mathbb{R}) = \text{Gl}_2(F \otimes \mathbb{R}) = \prod_{v \in S_\infty} \text{Gl}_2(F_v)$ . Let

$$K_v = (\text{SO}(2) \text{ ( resp. } U(2) ) \times C(F_v)^{(0)}) \subset \text{Gl}_2(F_v)$$

be the ess. maximal compact subgroups (See (4.1.2)). Our symmetric space

$$X = \prod_{v \in S_\infty} \text{Gl}_2(K_v)/K_v = \prod_{v \in S_\infty} X_v. \tag{8.91}$$

Let  $\mathcal{O}_{F,0}$  be the ring of integers in  $F_0$ , let  $\hat{\mathcal{O}}_{F,0} \subset \mathbb{A}_{F_0}$  be the ring of integral adeles, we consider the group scheme  $\mathcal{G}/\mathbb{Z} = R_{\mathcal{O}_{F,0}/\mathbb{Z}}(\text{Gl}_2/c\mathcal{O}_{F,0})$ . We choose an open compact subgroup  $K_f \subset \mathcal{G}(\hat{\mathcal{O}}_{F,0})$ . With these choices we define again

$$S_{K_f}^G = G(\mathbb{Q}) \backslash X \times G(\mathbb{A}_Q)/K_f \tag{8.92}$$

The Lie-algebra  $\mathfrak{g} \otimes \mathbb{R}$  of  $G \times_{\mathbb{Q}} \mathbb{R}$  is the direct sum  $\mathfrak{g} \otimes \mathbb{R} = \bigoplus_{v \in S_\infty} \mathfrak{g}_v$ , our standard Cartan involution is the product of the involution  $\Theta_v$ . (See 4.1.2). Then we get the corresponding Cartan decomposition

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p} \text{ where } \mathfrak{k} = \bigoplus_{v \in S_\infty} \mathfrak{k}_v, \mathfrak{p} = \bigoplus_{v \in S_\infty} \mathfrak{p}_v. \tag{8.93}$$

**The highest weight modules, the sheaves and their cohomology groups.**

Let  $\bar{\mathbb{Q}}$  the subfield of algebraic numbers of  $\mathbb{C}$ , then the maps  $\iota \in \Sigma$  factor through  $\bar{\mathbb{Q}}$ . Let  $F \in \bar{\mathbb{Q}}$  be the subfield generated by the  $\iota(F_0)$ , this is of course the normal closure of  $F_0$ . In (4.1.1) we gave a description of the character module of  $T_0/F$  : is an array The character module

$$X^*(T \times \bar{\mathbb{Q}}) = \text{Hom}(T \times \bar{\mathbb{Q}}, \mathbb{G}_m) = \prod_{\iota_0: F \rightarrow \bar{\mathbb{Q}}} X^*(T_0 \times_{F_0, \iota} \bar{\mathbb{Q}}) \tag{8.94}$$

and hence an element  $\underline{\lambda} \in X^*(T \times \bar{\mathbb{Q}})$  is an array

$$\underline{\lambda} = \{ \dots, n_\iota \gamma + d_\iota \det, \dots \}_{\iota \in \Sigma}. \tag{8.95}$$

We call  $\underline{\lambda}$  dominant if  $n_\iota \geq 0$  for all  $\iota \in \Sigma$ . The Galois group  $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$  acts on  $X^*(T \times \bar{\mathbb{Q}}, \mathbb{G}_m)$ , let  $F_\lambda \subset \bar{\mathbb{Q}}$  be the stabilizer field, i.e.  $\text{Gal}(\mathbb{Q}/F_\lambda)$  is the stabilizer of  $\underline{\lambda}$ . We can obviously extend our constructions in (4.1.1) to this situation and construct a  $\mathcal{G}$  module  $\mathcal{M}_\lambda^b$  of highest weight  $\underline{\lambda}$ . This is a free  $\mathcal{O}_{F_\lambda}$  module, for the extension  $\mathcal{M}_\lambda^b \otimes \mathcal{O}_{\bar{F}}$  we have a canonical isomorphism KueM

$$\mathcal{M}_\lambda^b \otimes \mathcal{O}_{\bar{F}} \xrightarrow{\sim} \bigotimes_{\iota \in \Sigma} \mathcal{M}_{n_\iota \gamma + d_\iota \det}^b \tag{8.96}$$

The tensor product  $\mathcal{M}_\lambda^b \otimes_{F_\lambda} F$  is of course isomorphic to the standard highest weight module  $\mathcal{M}_{\lambda,F}$ . We recall the explicit realization

$$\mathcal{M}_{n_\iota \gamma + d_\iota \det}^b = \{P(X_\iota, Y_\iota) = \sum_{m=0}^{n_\iota} a_\iota \binom{n_\iota}{m} X_\iota^{n_\iota-m} Y_\iota^m \mid a_\iota \in \mathcal{O}_F\} \quad (8.97)$$

We apply the considerations of section (6.2) to these modules. we get sheaves  $\tilde{\mathcal{M}}_\lambda^b$  on  $S_{K_f}^G$ . If the field  $F_0$  has at least one real place the sheaves  $\tilde{\mathcal{M}}_\lambda^b$  are zero unless all the coefficients  $d_\iota$  are all equal, i.e.  $d_\iota = d$ . (See [?]). Therefore we require that this is always so. The parameter  $d$  is actually rather irrelevant it only serves to fulfill the parity condition. We also require that  $\lambda$  is unitary, this means that for all the complex embeddings we have  $n_\iota = n_{\text{co}\iota}$  (See Thm. 4.1.2.). Of course, if  $F$  is totally real  $\lambda$  is always unitary.

We want to investigate the cohomology groups and the fundamental exact sequence

$$\rightarrow H_c^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}_\lambda^b) \xrightarrow{j_c} H^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}_\lambda^b) \xrightarrow{r} H^\bullet(\mathcal{N} S_{K_f}^G, \tilde{\mathcal{M}}_\lambda^b) \rightarrow \quad (8.98)$$

as modules for the Hecke-algebra in this special case. As usual we also introduce the inner "cohomology"  $H_!^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}_\lambda^b) = \ker(r) = \text{Im}(j_c)$ .

We apply theorem 8.1.1. The first step is to determine  $\text{Coh}(\lambda)$ . The Künneth-formula implies that any  $\pi_\infty \in \text{Coh}(\lambda)$  is a tensor product  $\pi_\infty = \bigotimes_{v \in S_\infty} \pi_v$  where  $\pi_v \in \text{Coh}(\lambda_v) = \text{Coh}(\lambda_v)_{\text{res}} \cup \text{Coh}(\lambda_v)_{\text{cusp}}$ . (See section 4.1.4 ). It is well known that  $H_{\pi_\infty}^{K_\infty} \neq \{0\}$  can occur if either all  $\pi_v \in \text{Coh}(\lambda_v)_{\text{res}}$  or all  $\pi_v \in \text{Coh}(\lambda_v)_{\text{cusp}}$ . The first case can only happen if  $n_\iota = 0$  for all  $\iota \in \Sigma$ . Then we get a decomposition

$$H_!^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{C}) = H_{!,\text{res}}^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{C}) \oplus H_{\text{cusp}}^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{C})$$

and the Manin-Drinfeld principle implies that this decomposition is defined over  $F_\lambda$ . (See [?]).

Therefore we get two decompositions deco-eis-cusp

$$\begin{aligned} H_c^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F_\lambda) &= H_{\text{cusp}}^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F_\lambda) \oplus H_{c,\text{Eis}}^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F_\lambda) \\ H^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F_\lambda) &= H_{\text{cusp}}^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F_\lambda) \oplus H_{\text{Eis}}^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F_\lambda) \end{aligned} \quad (8.99)$$

Of course this may not give a decomposition over  $\mathcal{O}_{F_\lambda}$ , the reason for this are the "denominators of the Eisenstein-classes". We only have decompositions up to isogeny

$$\begin{aligned} H_c^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}_\lambda^b \otimes \mathcal{O}_{F_\lambda}) &\supset H_{\text{cusp}}^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}_\lambda^b \otimes \mathcal{O}_{F_\lambda}) \oplus H_{c,\text{Eis}}^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}_\lambda^b \otimes \mathcal{O}_{F_\lambda}) \\ H^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}_\lambda^b \otimes \mathcal{O}_{F_\lambda}) &\supset H_{\text{cusp}}^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}_\lambda^b \otimes \mathcal{O}_{F_\lambda}) \oplus H_{\text{Eis}}^\bullet(S_{K_f}^G, \tilde{\mathcal{M}}_\lambda^b \otimes \mathcal{O}_{F_\lambda}) \end{aligned} \quad (8.100)$$

These denominators are actually very interesting, we have studied them in a special case in chapter V.

### The modular symbols

We apply our consideration in (8.2.1) to this special situation. We have two cases: In the first case our group  $G/\mathbb{Q}$  is  $G/\mathbb{Q} = \mathrm{Gl}_2/\mathbb{Q}$  and  $H/\mathbb{Q} = T/\mathbb{Q}$ , the standard maximal torus. In the second case we choose our group  $G/\mathbb{Q} = \mathrm{Gl}_2 \times \mathrm{Gl}_2/\mathbb{Q}$  and  $H/\mathbb{Q} = \mathrm{Gl}_2/\mathbb{Q}$  and we embed  $\mathrm{Gl}_2/\mathbb{Q}$  diagonally into  $\mathrm{Gl}_2 \times \mathrm{Gl}_2/\mathbb{Q}$ . For the archimedean component of our pin point we choose  $x = \prod_v K_v$  (resp.  $x = \prod_v (K_v \times K_v)$ ).

In the next step we compute the restriction of  $\mathcal{M}_\lambda$  to  $H$ , we do it separately for the two cases. The action of the torus  $H/\mathbb{Q} = T/\mathbb{Q}$  on  $\mathcal{M}_\lambda$  is semi simple and it is clear from (8.96) that the restriction to  $T/\mathbb{Q}$  decomposes into free rank one modules restr

$$\mathcal{M}_\lambda^b = \bigoplus_{\underline{\mu}} \mathcal{O}_{F_\mu} e_\mu^b \quad (8.101)$$

where  $\underline{\mu} = \{\dots, m_\iota \gamma_\iota + d \det, \dots\}$  where  $-n_\iota \leq m_\iota \leq n_\iota, n_\iota \equiv m_\iota \pmod{2}$  and where

$$e_\mu^b = \prod_{\iota} \binom{n_\iota}{m_\iota} X_\iota^{n_\iota - m_\iota} Y_\iota^{m_\iota}$$

is a generator. The homomorphism  $\phi_\mu$  will be the projection to the summand  $\mathcal{O}e_\mu^b$ . The space

$$S_{K_f^T}^T = T(\mathbb{Q}) \backslash (T(\mathbb{R})/K_\infty^T \times C(\mathbb{R})^{(0)}) \times T(\mathbb{A}_f)/K_f^T,$$

it has several connected components, each of these components is isomorphic to  $(S^1)^{r_1+r_2-1} \times R_{>0}^\times$ , the dimension is  $d_T = r_1 + r_2$ . Our data so far provide a homomorphism

$$\phi_\mu^! \circ j(x, \underline{g}_f) : H_c^{r_1+r_2}(\mathcal{S}_{K_f^G}^G, \tilde{\mathcal{M}}_\lambda^b) \rightarrow H_c^{r_1+r_2}(\mathcal{S}_{K_f^T}^T, \tilde{\mathcal{O}}_{F_\lambda} e_\mu) \quad (8.102)$$

This homomorphism factors over the quotient  $H_c^{r_1+r_2}(\mathcal{S}_{K_f^G}^G, \tilde{\mathcal{M}}_\lambda^b)$  if proposition 8.2.1 applies. If this proposition does not apply, then we have Manin-Drinfeld instead then we only get a homomorphism

$$\phi_\mu^! \circ j(x, \underline{g}_f) : H_c^{r_1+r_2}(\mathcal{S}_{K_f^G}^G, \tilde{\mathcal{M}}_\lambda \otimes F) \rightarrow H_c^{r_1+r_2}(\mathcal{S}_{K_f^T}^T, F_\lambda e_\mu) \quad (8.103)$$

Of course we want that the cohomology  $H_c^\bullet(\mathcal{S}_{K_f^T}^T, \tilde{\mathcal{O}}_{F_\lambda} e_\mu) \neq 0$ . A necessary condition for this to be the case, is that  $\underline{\mu}$  is *pure of weight*  $w(\underline{\mu})$ . This means that for the real embeddings all the numbers  $m_\iota = w(\underline{\mu})$  and for the pairs of complex embeddings we have  $m_\iota + m_{c\iota} = 2w(\underline{\mu})$ .

This is now the situation where we can try the strategy outlined in section 8.2.2. The module  $H_{\mathrm{cusp}}^{r_1+r_2}(\mathcal{S}_{K_f^G}^G, \tilde{\mathcal{M}}_\lambda)$  is semi simple, we can find a finite (normal over  $\mathbb{Q}$ ) extension  $F/F_\lambda$  such that we get an isotypical decomposition

$$H_{\mathrm{cusp}}^{r_1+r_2}(\mathcal{S}_{K_f^G}^G, \tilde{\mathcal{M}}_\lambda \otimes F) = \bigoplus_{\epsilon \times \pi_f} H_{\mathrm{cusp}}^{r_1+r_2}(\mathcal{S}_{K_f^G}^G, \tilde{\mathcal{M}}_\lambda \otimes F)(\epsilon \times \pi_f)$$

where  $\epsilon \times \pi_f$  is an isomorphism class of a (finite dimensional)  $F$ -vector space with an irreducible action of  $\pi_0(G(\mathbb{R})) \times \mathcal{H}$  on it.

For the integral cohomology we get a decomposition up to isogeny

$$H_{\text{cusp}}^{r_1+r_2}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda^b \otimes \mathcal{O}_F) \supset \bigoplus_{\epsilon \times \pi_f} H_{\text{cusp}}^{r_1+r_2}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda^b \otimes \mathcal{O}_F)(\epsilon \times \pi_f).$$

We can also decompose the right hand side. We have seen in ?? : We get a decomposition

$$H_!^{r_1+r_2}(\mathcal{S}_{K_f^T}^T, \tilde{F}_\lambda e_\mu) = \bigoplus_{\tilde{\mu}_f: \text{type}(\tilde{\mu}_f)=\mu} F e_{\tilde{\mu}_f}$$

and over the integers this gives us

$$H_!^{r_1+r_2}(\mathcal{S}_{K_f^T}^T, \tilde{\mathcal{O}}_{F_\lambda} e_\mu) \supset \bigoplus_{\phi: \text{type}(\phi)=\mu} \mathcal{O}_F e_{\tilde{\mu}_f}$$

and this gives us the projection map  $P_{\tilde{\mu}_f} : H_!^{r_1+r_2}(\mathcal{S}_{K_f^T}^T, \tilde{\mathcal{O}}_{F_\lambda} e_\mu) \rightarrow \mathcal{O}_F \tilde{e}_{\tilde{\mu}_f}$ . Hence we see: If we restrict to the  $\epsilon \times \pi_f$  component on the left hand side and project to the  $\tilde{\mu}_f$  component on the right hand side, then we get a homomorphism PiChi

$$P_{\tilde{\mu}_f} \phi_\mu \circ j(x, \underline{g}_f)(\epsilon \times \pi_f) : H_{\text{cusp}}^{r_1+r_2}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda^b \otimes \mathcal{O}_F)(\epsilon \times \pi_f) \rightarrow \mathcal{O}_F \tilde{e}_{\tilde{\mu}_f} \quad (8.104)$$

Recall that this only works if proposition 8.2.1 applies. But if we tensor our coefficient system with  $F$  then we can invoke the Manin-Drinfeld principle. We can change the level and go to the limit over smaller and smaller  $K_f$ . We get a  $\pi_0(G(\mathbb{R})) \times G(\mathbb{A}_f)$ - module homomorphism

$$J(\phi_\mu, \epsilon \times \pi_f, \tilde{\mathcal{H}}_f) : H_!^{r_1+r_2}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F)(\epsilon \times \pi_f) \rightarrow \text{Ind}_{T(\mathbb{A}_f)}^{G(\mathbb{A}_f)} \tilde{\mu}_f^{-1} \quad (8.105)$$

which is defined by

$$\xi \mapsto \{ \underline{g}_f \mapsto P_\phi \circ \phi_\mu \circ j(x, \underline{g}_f)(\epsilon \times \pi_f) \}. \quad (8.106)$$

This is now the situation where we can try the strategy outlined in section 8.2.2. In the following section we find a model space  $X_\epsilon(\pi_f) = \prod_{\mathfrak{p}} X(\pi_{\mathfrak{p}})$  together with an isomorphism  $\Phi : X_\epsilon(\pi_f) \xrightarrow{\sim} H_!^{r_1+r_2}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F)(\epsilon \times \pi_f)$ . We get the composite

$$J(\phi_\mu, \epsilon \times \pi_f, \tilde{\mu}_f \circ \Phi : X_\epsilon(\pi_f) \rightarrow \text{Ind}_{T(\mathbb{A}_f)}^{G(\mathbb{A}_f)} \tilde{\mu}_f^{-1}. \quad (8.107)$$

Now the  $G(\mathbb{A}_f)$ - modules are product of local  $G(F_{0,\mathfrak{p}})$ - modules and we find some "natural" local operators  $I_{\mathfrak{p}}^{\text{loc}} : X(\pi_{\mathfrak{p}}) \hookrightarrow \text{Ind}_{T_0(F_{0,\mathfrak{p}})}^{\text{Gl}_2(\mathbb{F}_{0,\mathfrak{p}})} \tilde{\mu}_{\mathfrak{p}}$ . Our final goal is to find an explicit formula for the factor in the comparison

$$J(\phi_\mu, \epsilon \times \pi_f, \tilde{\mu}_f) \circ \Phi = \mathcal{L}(\epsilon \times \pi_f, \tilde{\mu}_f) \prod_{\mathfrak{p}} I_{\mathfrak{p}}^{\text{loc}}. \quad (8.108)$$

### The Whittaker models

We assume that  $\pi_f$  is a representation which occurs in the decomposition of  $H_{\text{cusp}}^{r_1+r_2}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathcal{O}_F)$ . Let  $\pi_\infty = \otimes_{v \in S_\infty} \pi_v$  an isomorphism class of Harish-Chandra modules with  $\pi_v \in \text{Coh}_{\text{cusp}}(\lambda_v)$ . Then the isomorphism type  $\pi_\infty \otimes \pi_f$  occurs in  $L_{\text{cusp}}^2(G(\mathbb{Q}) \backslash G(\mathbb{A}))$ , we have to find model spaces  $X_{\mathfrak{p}}, X_{\pi_\infty}$ . We proceed as in section 4.1.6.

The adèle ring  $\mathbb{A} = \mathbb{A}_{F_0}$  is the restricted product

$$\mathbb{A} = (F_0 \otimes \mathbb{R}) \times \prod'_p (F_0 \otimes \mathbb{Q}_p) = \prod_{v \in S_\infty} F_v \times \prod_{\mathfrak{p}}' F_{\mathfrak{p}}.$$

The trace map  $\text{tr}_{F_0/\mathbb{Q}} : F_0 \rightarrow \mathbb{Q}$  induces a homomorphism  $\text{tr}_{F_0/\mathbb{Q}} : \mathbb{A}/F_0 \rightarrow \mathbb{A}_{\mathbb{Q}}/\mathbb{Q}$  which is of course the product  $\prod_v \text{tr}_{F_v/\mathbb{Q}_v}$ . We compose  $\text{tr}_{F_0/\mathbb{Q}}$  with the character  $\psi_1 : \mathbb{A}_{\mathbb{Q}}/\mathbb{Q} \rightarrow \mathbb{C}^\times$  in (4.107) and get a character  $\psi = \psi_1 \circ \text{tr}_{F_0/\mathbb{Q}} : \mathbb{A} \rightarrow S^1$ . (See ??). This character is of course a product  $\prod_{v \in S_\infty} \psi_v \times \prod_{\mathfrak{p}} \psi_{\mathfrak{p}}$ .

The character  $\psi_{1,p} : \mathbb{Q}_p \rightarrow S^1$  has a trivial "additive" conductor, this means that  $\psi_{1,p}(\mathbb{Z}_p) = 1$  and  $\psi_{1,p}(\frac{1}{p}\mathbb{Z}_p) \neq 1$ , or in other words  $\psi_{1,p} : \frac{1}{p}\mathbb{Z}_p \rightarrow S^1$  is a non trivial character. For any prime  $\mathfrak{p}$  which lies above  $p$  there is a largest integer  $d_{\mathfrak{p}} \geq 0$  such that  $\text{tr}_{F_0, \mathfrak{p}/\mathbb{Q}_p}(\mathfrak{p}^{-d_{\mathfrak{p}}}\mathcal{O}_{F_0, \mathfrak{p}}) \subset \mathbb{Z}_p$ . Then it is clear that  $d_{\mathfrak{p}}$  is the largest integer such that  $\psi_{\mathfrak{p}} : \mathfrak{p}^{-d_{\mathfrak{p}}}\mathcal{O}_{F_0, \mathfrak{p}} \rightarrow S^1$  is the trivial character. The ideal  $\mathfrak{p}^{d_{\mathfrak{p}}}$  and sometimes also simply  $d_{\mathfrak{p}}$  is the "additive" conductor of  $\psi$ . Of course we have  $d_{\mathfrak{p}} = 0$  for almost all  $\mathfrak{p}$ , the ideal  $\vartheta_{F_0} = \prod_{\mathfrak{p}} \mathfrak{p}^{d_{\mathfrak{p}}}$  is the different of  $F_0$ . The character  $\psi$  is trivial on  $F_0 \subset \mathbb{A}$  hence  $\psi \in \text{Hom}(\mathbb{A}/F_0, S^1)$ . It is in some sense a distinguished element: It is obtained by a canonical construction from  $\psi_1$ . All other characters in  $\text{Hom}(\mathbb{A}/F_0, S^1)$  are of the form

$$\psi^{[a]} := \underline{x} \mapsto \psi(a\underline{x}) \text{ with some } a \in F_0,$$

hence we can say  $\text{Hom}(\mathbb{A}/F_0, S^1) = F_0$ .

Now we know that for any place  $v$  (finite or archimedean) and any -not one dimensional- irreducible representation  $\pi_{\mathfrak{p}}$  we have a Whittaker model, this means that we have an unique subspace

$$\mathcal{W}(\pi_v, \psi_v)_{\mathbb{C}} \subset \{f : \text{Gl}_2(F_v) \rightarrow \mathbb{C} \mid f\left(\begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix} g\right) = \psi_v(u)f(g)\} \quad (8.109)$$

which invariant under right translations and isomorphic to  $\pi_v$ . This Whittaker model depends of course the choice of  $\psi_v$  which in the following always is the  $v$ -local component of the distinguished  $\psi$ . At some instances we have to use the fact that we have an isomorphism shift

$$R_{a_v} : \mathcal{W}(\pi_v, \psi_v) \xrightarrow{\sim} \mathcal{W}(\pi_v, \psi_v^{[a_v]})$$

$$f(g) \mapsto f\left(\begin{pmatrix} a_v & 0 \\ 0 & 1 \end{pmatrix} g\right) \quad (8.110)$$

At the archimedean places these are the modules  $\tilde{\mathcal{D}}_{\lambda_v}$ . We can form the tensor product

$$\mathcal{W}(\pi, \psi)_{\mathbb{C}} := \bigotimes_v \mathcal{W}(\pi_v, \psi_v)_{\mathbb{C}} \quad (8.111)$$

and these spaces will be our  $X_{\mathfrak{p}}, X_{\pi_{\infty}}$ .

Our level subgroup should be a product  $K_f = \prod_{\mathfrak{p}} K_{\mathfrak{p}}$ , then

$$\mathcal{W}(\pi_f^{K_f}, \psi_f)_{\mathbb{C}} = \prod_{\mathfrak{p}} \mathcal{W}(\pi_{\mathfrak{p}}^{K_{\mathfrak{p}}}, \psi_{\mathfrak{p}})_{\mathbb{C}}$$

is a module for the Hecke algebra  $\mathcal{H}_{K_f}^G$ .

The Fourier expansion gives us an isomorphism

$$\begin{aligned} \mathcal{F}_1 : \mathcal{W}(\pi, \psi) &\rightarrow \mathcal{A}(\mathrm{Gl}_2(F) \backslash \mathrm{Gl}_2(\mathbb{A}))(\pi) \\ \underline{f}(\underline{g}) &= \{\dots, f_v, \dots\} \mapsto \sum_{a \in T_1(\mathbb{Q})} \underline{f}(a\underline{g}) \end{aligned} \tag{8.112}$$

this will be our intertwining operator in (8.76). We get an isomorphism

$$\mathcal{F}_1^{d_T} : \bigotimes_{v \in S_{\infty}} H^1(\mathfrak{g}_v, K_v, \mathcal{D}_v \otimes \mathcal{M}_{\lambda_v}) \otimes \bigotimes_{\mathfrak{p}} \mathcal{W}(\pi_{\mathfrak{p}}^{K_{\mathfrak{p}}}, \psi_{\mathfrak{p}}) \xrightarrow{\sim} H_{\mathrm{cusp}}^{r_1+r_2}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda} \otimes F)(\pi_f) \otimes \mathbb{C} \tag{8.113}$$

A well known consequence is that  $H_{\mathrm{cusp}}^{r_1+r_2}(\mathcal{S}^G, \tilde{\mathcal{M}}_{\lambda} \otimes F)(\epsilon \times \pi_f) \otimes \mathbb{C}$  occurs with multiplicity one. Therefore  $H_{\mathrm{cusp}}^{r_1+r_2}(\mathcal{S}^G, \tilde{\mathcal{M}}_{\lambda} \otimes F)(\epsilon \times \pi_f)$  is a  $F$ -vector space with an absolutely irreducible  $G(\mathbb{A}_f)$  module structure, hence we can also say that we realized  $\pi_f$  over  $F$ , we abbreviate  $H(\pi_f) := H_{\mathrm{cusp}}^{r_1+r_2}(\mathcal{S}^G, \tilde{\mathcal{M}}_{\lambda} \otimes F)(\epsilon \times \pi_f)$

### Rational and integral structures on the Whittaker model.

Of course we have  $H(\pi_f) = \bigotimes' H(\pi_{\mathfrak{p}})$  At a finite place  $\mathfrak{p}$  we can realize our local representation  $\pi_{\mathfrak{p}}$  as a subspace  $\mathcal{W}_{\overline{\mathbb{Q}}}(\pi_{\mathfrak{p}}, \psi_{\mathfrak{p}})_{\overline{\mathbb{Q}}}$  in the space of  $\overline{\mathbb{Q}}$ -valued functions

$$\mathcal{W}_{\overline{\mathbb{Q}}}(\psi_{\mathfrak{p}}) = \left\{ f : G(F_{\mathfrak{p}}) \rightarrow \overline{\mathbb{Q}} \mid f \left( \begin{pmatrix} 1 & u_{\mathfrak{p}} \\ 0 & 1 \end{pmatrix} g_{\mathfrak{p}} \right) = \psi_{\mathfrak{p}}(u_{\mathfrak{p}}) f(g_{\mathfrak{p}}) \right\}.$$

We briefly sketch how we get this realisation. We choose a no zero linear form  $L_0 : H(\pi_f) \rightarrow F$  and then we define a second linear form  $L : H(\pi_f) \rightarrow F$  by the integral

$$L(h) := \int_{U_0(F_{0,\mathfrak{p}})} (\rho_{\pi_{\mathfrak{p}}} \left( \begin{pmatrix} 1 & u_{\mathfrak{p}} \\ 0 & 1 \end{pmatrix} \right) h) \overline{\psi_{\mathfrak{p}}(u)} du$$

here is a minor issue with convergence, this will be discussed later. Then it is clear that  $L(\left( \begin{pmatrix} 1 & u_{\mathfrak{p}} \\ 0 & 1 \end{pmatrix} \right) h) = L(h) \psi_{\mathfrak{p}}(u)$ . and

$$h \mapsto \{g_{\mathfrak{p}} \mapsto L(\rho_{\pi_{\mathfrak{p}}}(g_{\mathfrak{p}})(h))\} \tag{8.114}$$

is either zero or a  $\mathrm{Gl}_2(F_{0,\mathfrak{p}})$  - isomorphism of this  $H(\pi_{\mathfrak{p}})$  with a subspace in  $\mathcal{W}_{\overline{\mathbb{Q}}}(\psi_{\mathfrak{p}})$ .

On this space we define an action of the Galois group: The values  $\psi_{\mathfrak{p}}(u_{\mathfrak{p}})$  are  $p^m$ -th roots of unity, we have the reciprocity homomorphism

$$\alpha : \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \mathrm{Gal}(\mathbb{Q}(\zeta_{p^{\infty}})/\mathbb{Q}) \xrightarrow{\sim} \mathbb{Z}_p^{\times}.$$

For  $f \in \mathcal{W}_{\overline{\mathbb{Q}}}(\psi_{\mathfrak{p}})$  and  $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$  we put ([23])

$$f^\sigma(g) = \left( f \left( \begin{pmatrix} \alpha(\sigma)^{-1} & 0 \\ 0 & 1 \end{pmatrix} g \right) \right)^\sigma,$$

If we take an element  $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$  then it conjugates the representation  $\pi_{\mathfrak{p}}$  into  $\pi_{\mathfrak{p}}^\sigma$  and we get a map

$$\begin{array}{ccc} \mathcal{W}_{\overline{\mathbb{Q}}}(\pi_{\mathfrak{p}}, \psi_{\mathfrak{p}}) & \xrightarrow{\tilde{\sigma}} & \mathcal{W}_{\overline{\mathbb{Q}}}(\pi_{\mathfrak{p}}^\sigma, \psi_{\mathfrak{p}}) \\ f & \mapsto & f^\sigma \end{array}$$

This map is a semilinear isomorphism and  $\mathbb{Q}(\pi_{\mathfrak{p}})$  is the number field for which  $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\pi_{\mathfrak{p}}))$  is the stabilizer of  $\mathcal{W}_{\overline{\mathbb{Q}}}(\pi_{\mathfrak{p}}, \psi_{\mathfrak{p}})$ . The space  $\mathcal{W}_{\overline{\mathbb{Q}}}(\pi_{\mathfrak{p}}, \psi_{\mathfrak{p}})$  is the union of finite dimensional  $\overline{\mathbb{Q}}$ , modules  $\mathcal{W}_{\overline{\mathbb{Q}}}(\pi_{\mathfrak{p}}^{K_{\mathfrak{p}}}, \psi_{\mathfrak{p}})$  where  $K_{\mathfrak{p}}$  runs over the open compact subgroups. The space of functions in  $\mathcal{W}(\pi_{\mathfrak{p}}^{K_{\mathfrak{p}}}, \psi_{\mathfrak{p}})$  which are invariant under  $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\pi_{\mathfrak{p}}))$  is a  $\mathbb{Q}(\pi_{\mathfrak{p}})$  vector space  $\mathcal{W}(\pi_{\mathfrak{p}}^{K_{\mathfrak{p}}}, \psi_{\mathfrak{p}})$  on which  $\mathcal{H}(G(F_{\mathfrak{p}})/K_{\mathfrak{p}})$  acts absolutely irreducibly. Of course  $\mathcal{W}(\pi_{\mathfrak{p}}, \psi_{\mathfrak{p}})$  is the union of the  $\mathcal{W}(\pi_{\mathfrak{p}}^{K_{\mathfrak{p}}}, \psi_{\mathfrak{p}})$  and clearly  $\mathcal{W}(\pi_{\mathfrak{p}}, \psi_{\mathfrak{p}}) \otimes_{\mathbb{Q}(\pi_{\mathfrak{p}})} \overline{\mathbb{Q}} = \mathcal{W}_{\overline{\mathbb{Q}}}(\pi_{\mathfrak{p}}, \psi_{\mathfrak{p}})$ .

Of course  $\mathbb{Q}(\pi_{\mathfrak{p}}) \subset \mathbb{Q}(\pi_f)$ , and we define a subring  $\mathcal{O}(\pi_f) \subset \mathbb{Q}(\pi_f)$ . We have the action of  $\mathcal{H}_{\mathbb{Z}}^{\text{coh}}$  (See 1.2.1.(ii)) on the cohomology and hence we get an action of the algebra  $\mathcal{H}(G(F_{\mathfrak{p}})/K_{\mathfrak{p}})_{\mathbb{Z}}$  on  $\mathcal{W}(\pi_{\mathfrak{p}}, \psi_{\mathfrak{p}})$  and this gives us a finitely generated  $\mathcal{O}(\pi_{\mathfrak{p}})$ -module of endomorphisms. Hence we can find invariant lattices  $\mathcal{W}_{\mathcal{O}(\pi_{\mathfrak{p}})}(\pi_{\mathfrak{p}}^{K_{\mathfrak{p}}}, \psi_{\mathfrak{p}})_{\mathcal{O}(\pi_{\mathfrak{p}})}$ . If we invert a few more primes then we can achieve that two such choices just differ by an element  $a \in \mathcal{O}(\pi_{\mathfrak{p}})$ . We assume that such a choice of lattices has been made at all primes  $\mathfrak{p}$ . If we are in the unramified case then we will make a very particular choice later. We put  $\mathcal{W}_{\mathcal{O}(\pi_f)}(\pi_f, \psi_{\mathfrak{p}}) = \bigotimes_{\mathfrak{p}} \mathcal{W}_{\mathcal{O}(\pi_f)}(\pi_{\mathfrak{p}}, \psi_{\mathfrak{p}})$  ( See 2.2.7 ).

### The Newvector

For any integer  $n \geq 0$  we define the congruence subgroups

$$\begin{aligned} K_{\mathfrak{p},0}(\mathfrak{p}^{f(\pi_{\mathfrak{p}})}) &= \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid c \equiv 0 \pmod{\mathfrak{p}^{f(\pi_{\mathfrak{p}})}} \right\} \subset \text{Gl}_2(\mathcal{O}_{\mathfrak{p}}) \\ K_{\mathfrak{p},1}(\mathfrak{p}^{f(\pi_{\mathfrak{p}})}) &= \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid c \equiv 0 \pmod{\mathfrak{p}^{f(\pi_{\mathfrak{p}})}} \text{ and } a \equiv 1 \pmod{\mathfrak{p}^{f(\pi_{\mathfrak{p}})}} \right\} \subset \text{Gl}_2(\mathcal{O}_{\mathfrak{p}}). \end{aligned}$$

Clearly we have  $K_{\mathfrak{p},0}(\mathfrak{p}^{f(\pi_{\mathfrak{p}})}) = K_{\mathfrak{p},1}(\mathfrak{p}^{f(\pi_{\mathfrak{p}})}) \times \left\{ \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix} \right\}_{a \in (\mathcal{O}_{\mathfrak{p}})/\mathfrak{p}^{f(\pi_{\mathfrak{p}})}} = C(\mathcal{O}_{\mathfrak{p}}/\mathfrak{p}^{f(\pi_{\mathfrak{p}})})$ .

A theorem of Casselman and Novikovskii([?]) implies that there is smallest integer  $f(\pi_{\mathfrak{p}}) \geq 0$  such that  $H(\pi_{\mathfrak{p}})^{K_{\mathfrak{p},1}(\mathfrak{p}^{f(\pi_{\mathfrak{p}})})} \neq 0$  and it also says that the dimension of this space is one. An element  $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in K_{\mathfrak{p},0}(\mathfrak{p}^{f(\pi_{\mathfrak{p}})})$  acts by multiplication the central character  $\zeta_{\mathfrak{p}}(a)$  on this one dimensional space. The ideal  $\mathfrak{p}^{f(\pi_{\mathfrak{p}})}$  is the *conductor* of  $\pi_{\mathfrak{p}}$ .

We assume that  $K_{\mathfrak{p}} = K_{\mathfrak{p},1}(\mathfrak{p}^{f(\pi_{\mathfrak{p}})})$  at all finite primes. Then  $\mathcal{W}(\pi_{\mathfrak{p}}^{K_{\mathfrak{p}}}, \psi_{\mathfrak{p}})$  is of dimension one over  $\mathbb{Q}(\pi_f)$ , we can choose any non zero element  $h_{\mathfrak{p}}$  a a

generator of such a lattice. In the following we give a rule to choose a specific generator. At first we consider

**The Principal Series**

To see what is going on we consider the special case that  $\pi_{\mathfrak{p}}$  is a principal series representation. This means that

$$\chi_{\mathfrak{p}} : \begin{pmatrix} t_1 & * \\ 0 & t_2 \end{pmatrix} \longrightarrow \chi_{\mathfrak{p},1}(t_1) \cdot \chi_{\mathfrak{p},2}(t_2)$$

is an unramified character and  $\pi_{\mathfrak{p}}$  is the induced representation from  $\chi_{\mathfrak{p}}$ , i.e. we consider the space of functions

$$\text{Ind}_{B(F_{\mathfrak{p}})}^{G(F_{\mathfrak{p}})} \chi_{\mathfrak{p}} = \left\{ f : G(F_{\mathfrak{p}}) \rightarrow \mathbb{C} \mid f \left( \begin{pmatrix} t_1 & * \\ 0 & t_2 \end{pmatrix} g \right) = \chi_{\mathfrak{p},1}(t_1) \chi_{\mathfrak{p},2}(t_2) f(g) \right\},$$

since we want the representation to be admissible the function  $f$  must be right invariant under some open subgroup  $K'_{\mathfrak{p}}$ .

Let us denote the restriction of  $\chi_{\mathfrak{p}}$  to the subtorus  $T^{(1)}(F_{\mathfrak{p}}) = \left\{ \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix} \mid t \in F_{\mathfrak{p}}^{\times} \right\} = F_{\mathfrak{p}}^{\times}$  by  $\chi_{\mathfrak{p}}^{(1)}$ .

**Proposition 8.2.4.** *i) The induced representation  $\text{Ind}_{B(F_{\mathfrak{p}})}^{G(F_{\mathfrak{p}})} \chi_{\mathfrak{p}}$  is irreducible unless  $\chi_{\mathfrak{p}}^{(1)} = \mathbf{1}$  (the trivial character) or  $|\mathfrak{p}$ .*

*ii) If  $\chi_{\mathfrak{p}}^{(1)} = \mathbf{1}$ , the one dimensional space of functions  $g \mapsto a \chi_{1,\mathfrak{p}}(\det(g))$  form an invariant subspace the quotient by this subspace is irreducible. This quotient is called the Steinberg module  $\text{St}(\chi_{\mathfrak{p}})$ .*

*iii) If  $\chi_{\mathfrak{p}}^{(1)} = |\mathfrak{p}$  the integral*

$$f \mapsto \int_{\text{Sl}_2(\mathcal{O}_{\mathfrak{p}})} f(k) dk.$$

*defines an invariant linear form, the kernel is irreducible and isomorphic to  $\text{St}(\chi_{\mathfrak{p}})$*

See [18]

In this case we have an obvious option for an intertwining operator to the Whittaker model:

$$R_{\mathfrak{p}} : \text{Ind}_{B(F_{\mathfrak{p}})}^{G(F_{\mathfrak{p}})} \chi_{\mathfrak{p}} \longrightarrow \mathcal{W}(\pi_{\mathfrak{p}}(\chi_{\mathfrak{p}}), \psi_{\mathfrak{p}}),$$

it is given by

$$R_{\mathfrak{p}}(f) \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) = \int_{U(F_{\mathfrak{p}})} f \left( w \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix} \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) \overline{\psi_{\mathfrak{p}}(u)} du$$

where  $w = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ , and  $du$  is the additively invariant measure that gives volume one to  $\mathcal{O}_{F_{\mathfrak{p}}}$ . After a small substitution the integral becomes

$$|t|_{\mathfrak{p}} \chi_{2,\mathfrak{p}}(t) \int_{U(F_{\mathfrak{p}})} f \left( \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix} \right) \overline{\psi_{\mathfrak{p}}(tu)} du \tag{8.115}$$

where of course  $|t|_{\mathfrak{p}} = N(\mathfrak{p})^{-\text{ord}_{\mathfrak{p}}(t)}$  is the normalised  $\mathfrak{p}$ -adic absolute value.

Our integral can be written as an infinite sum RP

$$\int_{\mathcal{O}_{F_0, \mathfrak{p}}} f\left(w \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix}\right) \overline{\psi_{\mathfrak{p}}(tu)} du + \sum_{\nu=1}^{\infty} \int_{\varpi_{\mathfrak{p}}^{-\nu} \mathcal{O}_{F_0, \mathfrak{p}} \setminus \varpi_{\mathfrak{p}}^{-\nu+1} \mathcal{O}_{F_0, \mathfrak{p}}} f\left(w \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix}\right) \overline{\psi_{\mathfrak{p}}(tu)} du \quad (8.116)$$

We choose a local uniformizing element  $\varpi_{\mathfrak{p}}$  and write any element  $u \in F_{0, \mathfrak{p}}^{\times}$  as  $u = \varpi_{\mathfrak{p}}^{-\nu} \varepsilon$  with  $\varepsilon$  a unit and of course  $\nu = \text{ord}_{\mathfrak{p}}(u)$ . Then the  $\nu$ -th summand becomes term1

$$N(\mathfrak{p})^{\nu} \int_{\mathcal{O}_{F_0, \mathfrak{p}} \setminus \varpi_{\mathfrak{p}} \mathcal{O}_{F_0, \mathfrak{p}}} f\left(w \begin{pmatrix} 1 & \varpi_{\mathfrak{p}}^{-\nu} u \\ 0 & 1 \end{pmatrix}\right) \overline{\psi_{\mathfrak{p}}(t \varpi_{\mathfrak{p}}^{-\nu} u)} du \quad (8.117)$$

We have to compute the value of the function  $f$ , we apply the Iwasawa decomposition and for  $u \in \mathcal{O}_{F_0, \mathfrak{p}} \setminus \varpi_{\mathfrak{p}} \mathcal{O}_{F_0, \mathfrak{p}}$  we have

$$\underline{f}\left(w \begin{pmatrix} 1 & \varpi_{\mathfrak{p}}^{-\nu} u \\ 0 & 1 \end{pmatrix}\right) = f\left(\begin{pmatrix} \varpi_{\mathfrak{p}}^{\nu} u^{-1} & -1 \\ 0 & \varpi_{\mathfrak{p}}^{-\nu} u \end{pmatrix} \begin{pmatrix} 1 & 0 \\ \varpi_{\mathfrak{p}}^{\nu} u^{-1} & 1 \end{pmatrix}\right) = \chi_{\mathfrak{p}}^{(1)}(\varpi_{\mathfrak{p}})^{\nu} \chi_{\mathfrak{p}}^{(1)}(u)^{-1} f\left(\begin{pmatrix} 1 & 0 \\ \varpi_{\mathfrak{p}}^{\nu} u^{-1} & 1 \end{pmatrix}\right) \quad (8.118)$$

Our first concern is the convergence of this integral, we have to show that almost all summands are zero, the integral is actually a finite sum. We look at an individual term. We observe that for  $\nu$  large enough we have

$$f\left(\begin{pmatrix} 1 & 0 \\ \varpi_{\mathfrak{p}}^{\nu} u^{-1} & 1 \end{pmatrix}\right) = f\left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}\right),$$

hence we have to study the expression Gsum

$$G(\varpi_{\mathfrak{p}}^{-\nu}, \chi_{\mathfrak{p}}, \psi_{\mathfrak{p}}) = \int_{\mathcal{O}_{F_0, \mathfrak{p}} \setminus \varpi_{\mathfrak{p}} \mathcal{O}_{F_0, \mathfrak{p}}} \chi_{\mathfrak{p}}^{(1)}(u)^{-1} \overline{\psi_{\mathfrak{p}}(t \varpi_{\mathfrak{p}}^{-\nu} u)} du. \quad (8.119)$$

We remind the reader of the notion of the "additive" conductor of  $\psi_{\mathfrak{p}}$ . This is the smallest integer  $d_{\mathfrak{p}} = d(\psi_{\mathfrak{p}})$  for which the character  $\psi_{\mathfrak{p}}$  is trivial on  $\varpi_{\mathfrak{p}}^{-d_{\mathfrak{p}}} \mathcal{O}_{F_0, \mathfrak{p}}$ . We also have the multiplicative conductor  $\mathfrak{f}(\chi_{\mathfrak{p}}) = \mathfrak{f}(\chi_{\mathfrak{p}}^{(1)})$ . This is the smallest integer  $\mathfrak{f}_0$  such that  $\chi_{\mathfrak{p}}^{(1)}$  is trivial on the subgroup  $\mathcal{O}_{F_0, \mathfrak{p}}^{(\mathfrak{f}_0)} = \{u \in \mathcal{O}_{F_0, \mathfrak{p}}^{\times} \mid u \equiv 1 \pmod{\varpi_{\mathfrak{p}}^{\mathfrak{f}_0}}\}$ . Then the last integral gives

$$\int_{\mathcal{O}_{F_0, \mathfrak{p}}^{\times}} \chi_{\mathfrak{p}}^{(1)}(\varepsilon^{-1}) \overline{\psi_{\mathfrak{p}}(t \varpi_{\mathfrak{p}}^{-\nu} \varepsilon^{-1})} d\varepsilon = \sum_{\eta \in \mathcal{O}_{F_0, \mathfrak{p}}^{\times} / \mathcal{O}_{F_0, \mathfrak{p}}^{(\mathfrak{f}_0)}} \chi_{\mathfrak{p}}^{(1)}(\eta^{-1}) \int_{\mathcal{O}_{F_0, \mathfrak{p}}^{(\mathfrak{f}_0)}} \psi_{\mathfrak{p}}(t \varpi_{\mathfrak{p}}^{-\nu} \eta u) du \quad (8.120)$$

We claim that the integral in this expression has value zero if  $\nu \gg 0$ . To see this we look at the case  $\mathfrak{f}_0 = 0$  first. Then

$$\int_{\mathcal{O}_{F_0, \mathfrak{p}}^{\times}} \psi_{\mathfrak{p}}(t \varpi_{\mathfrak{p}}^{-\nu} \eta u) du = \int_{\mathcal{O}_{F_0, \mathfrak{p}}} \psi_{\mathfrak{p}}(t \varpi_{\mathfrak{p}}^{-\nu} u) du - \int_{\varpi_{\mathfrak{p}} \mathcal{O}_{F_0, \mathfrak{p}}} \psi_{\mathfrak{p}}(t \varpi_{\mathfrak{p}}^{-\nu} u) du$$

For any  $t_1 = \varpi_{\mathfrak{p}}^m \varepsilon \in F_{0,\mathfrak{p}}^\times$  we know the character  $\mathcal{O}_{F_{0,\mathfrak{p}}} \rightarrow \mathbb{C}^\times, u \mapsto \psi_{\mathfrak{p}}(t_1 u)$  is trivial  $\iff d_{\mathfrak{p}} + m \geq 0$  and therefore

$$\int_{\mathcal{O}_{F_{0,\mathfrak{p}}}} \psi_{\mathfrak{p}}(t_1 u) du = \begin{cases} 1 & \text{if } d_{\mathfrak{p}} + m \geq 0 \\ 0 & \text{else} \end{cases}. \tag{8.121}$$

Therefore

$$\int_{\mathcal{O}_{F_{0,\mathfrak{p}}} \setminus \varpi_{\mathfrak{p}} \mathcal{O}_{F_{0,\mathfrak{p}}}} \psi_{\mathfrak{p}}(t \varpi_{\mathfrak{p}}^{-\nu} u) du = \begin{cases} 0 & \text{if } -\nu + \text{ord}_{\mathfrak{p}}(t) < -d_{\mathfrak{p}} - 1 \\ -\frac{1}{N(\mathfrak{p})} & \text{if } -\nu + \text{ord}_{\mathfrak{p}}(t) = -d_{\mathfrak{p}} - 1 \\ 1 & \text{if } -\nu + \text{ord}_{\mathfrak{p}}(t) \geq -d_{\mathfrak{p}} \end{cases} \tag{8.122}$$

and this implies the claim in the case  $f_0 = 0$ . If  $f_0 > 0$  then we write  $\varepsilon = 1 + \varpi_{\mathfrak{p}}^{f_0} v$  with  $v \in \mathcal{O}_{F_{0,\mathfrak{p}}}$  and then  $\psi_{\mathfrak{p}}(t \varpi_{\mathfrak{p}}^{-\nu} \eta \varepsilon) = \psi_{\mathfrak{p}}(t \varpi_{\mathfrak{p}}^{-\nu} \eta) \psi_{\mathfrak{p}}(t \varpi_{\mathfrak{p}}^{-\nu + f_0} \eta v)$  and then  $v \rightarrow \psi_{\mathfrak{p}}(t \varpi_{\mathfrak{p}}^{-\nu + f_0} \eta v)$  is a non trivial character on  $\mathcal{O}_{F_{0,\mathfrak{p}}}$  provided  $\nu \gg 0$  and hence the integral is zero. This proves the claim and this implies that the sum over  $\nu$  is finite, hence there is no problem with convergence.

We recall the definition of the Schwartz-spaces  $\mathcal{S}(F_{0,\mathfrak{p}})$ - this are the locally constant  $\mathbb{Q}$  valued functions with compact support-, and  $\mathcal{S}(F_{0,\mathfrak{p}}^\times)$ , this is the space of those functions in  $\mathcal{S}(F_{0,\mathfrak{p}})$  which vanish at 0. This is of course equal to the locally constant  $\mathbb{Q}$  valued functions on  $F_{0,\mathfrak{p}}^\times$  with compact support.

a) *Covergence: The terms with  $\nu \gg 0$  vanish, hence the sum is finite.*

b) *If  $\text{ord}_{\mathfrak{p}}(t) \ll 0$ , i.e.  $t \rightarrow \infty$  we get  $R_{\mathfrak{p}}(f)\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right) = 0$  this means that  $R_{\mathfrak{p}}(f)\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right)$  lies in  $\mathcal{S}(F_{0,\mathfrak{p}})$ .*

c) *The space of all restriction  $R_{\mathfrak{p}}(f)$  to  $\left\{\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right\} = F_{0,\mathfrak{p}}^\times$  contains  $\mathcal{S}(F_{0,\mathfrak{p}}^\times)$  and the quotient  $\text{Im}(R_{\mathfrak{p}})/\mathcal{S}(F_{0,\mathfrak{p}}^\times)$  is of rank 2 or 1. (See further down)*

The composition of  $R_{\mathfrak{p}}$  with the restriction to  $F_{0,\mathfrak{p}}^\times$  is injective, the image is called the Kirillov-model. (See further down).

We apply this computation to a generator in  $f_0 \in (\text{Ind}_{B(F_{\mathfrak{p}})}^{G(F_{\mathfrak{p}})} \chi_{\mathfrak{p}}))^{K_{\mathfrak{p},1}(\mathfrak{p}^{f(\pi_{\mathfrak{p}})})}$ .

We go back to equation (8.118) and for the last factor on the right we get

$$f_0\left(\begin{pmatrix} 1 & 0 \\ \varpi_{\mathfrak{p}}^\nu u^{-1} & 1 \end{pmatrix}\right) = f_0\left(\begin{pmatrix} 1 & 0 \\ 0 & u^{-1} \end{pmatrix}\right) \left(\begin{pmatrix} 1 & 0 \\ \varpi_{\mathfrak{p}}^\nu & 1 \end{pmatrix}\right) \left(\begin{pmatrix} 1 & 0 \\ 0 & u \end{pmatrix}\right) = \chi_{2,\mathfrak{p}}(u^{-1}) f_0\left(\begin{pmatrix} 1 & 0 \\ \varpi_{\mathfrak{p}}^\nu & 1 \end{pmatrix}\right)$$

because  $\begin{pmatrix} 1 & 0 \\ 0 & u \end{pmatrix} \in K_{\mathfrak{p},1}(\mathfrak{p}^{f(\pi_{\mathfrak{p}})})$ . Therefore the term on the right in (8.118) becomes

$$N(\mathfrak{p})^\nu \chi^{(1)}(\varpi_{\mathfrak{p}}^\nu) \chi_{1,\mathfrak{p}}(u)^{-1} f_0\left(\begin{pmatrix} 1 & 0 \\ \varpi_{\mathfrak{p}}^\nu & 1 \end{pmatrix}\right)$$

hence we get finally

$$R_{\mathfrak{p}}(f_0)\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right) = |t|_{\mathfrak{p}} \chi_{2,\mathfrak{p}}(t) \times \left[ \int_{\mathcal{O}_{F_0,\mathfrak{p}}} f_0(w) \overline{\psi_{\mathfrak{p}}(tu)} du + \right. \\ \left. + \sum_{\nu=1}^{\infty} N(\mathfrak{p})^{\nu} \chi_{\mathfrak{p}}^{(1)}(\varpi_{\mathfrak{p}}^{\nu}) f_0\left(\begin{pmatrix} 1 & 0 \\ \varpi_{\mathfrak{p}}^{\nu} & 1 \end{pmatrix}\right) \int_{\mathcal{O}_{F_0,\mathfrak{p}}^{\times}} \chi_{1,\mathfrak{p}}(u)^{-1} \overline{\psi_{\mathfrak{p}}(t\varpi_{\mathfrak{p}}^{-\nu} u)} du \right] \quad (8.123)$$

We need to fix a specific generator  $f_0 = h_{\pi_{\mathfrak{p}}}^{(0)} \in (\text{Ind}_{B(F_{\mathfrak{p}})}^{G(F_{\mathfrak{p}})} \chi_{\mathfrak{p}})^{K_{\mathfrak{p},1}(\mathfrak{p}^{f(\pi_{\mathfrak{p}})})}$  ( resp.  $\text{St}(\chi_{\mathfrak{p}})^{K_{\mathfrak{p},1}(\mathfrak{p}^{f(\pi_{\mathfrak{p}})})}$ ). We look at the double coset decomposition

$$\text{Gl}_2(\mathcal{O}_{\mathfrak{p}}) = \bigcup_{\xi} B(\mathcal{O}_{\mathfrak{p}}) \xi K_{\mathfrak{p},0}(\mathfrak{p}^{f(\pi_{\mathfrak{p}})}) = \bigcup_{\xi} B(\mathcal{O}_{\mathfrak{p}}) \xi K_{\mathfrak{p},1}(\mathfrak{p}^{f(\pi_{\mathfrak{p}})}) \quad (8.124)$$

It is easy to see that a system of representatives for these double cosets is given by the matrices

$$\left\{ \begin{pmatrix} 1 & 0 \\ \varpi_{\mathfrak{p}}^{\nu} & 1 \end{pmatrix} \right\}_{\nu=1, \dots, f(\pi_{\mathfrak{p}})} \cup \left\{ \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \right\} \quad (8.125)$$

The space  $(\text{Ind}_{B(F_{\mathfrak{p}})}^{G(F_{\mathfrak{p}})} \chi_{\mathfrak{p}})^{K_{\mathfrak{p},1}(\mathfrak{p}^{f(\pi_{\mathfrak{p}})})}$  is spanned by functions

$$f_{\xi} : B(\mathcal{O}_{\mathfrak{p}}) \xi K_{\mathfrak{p},1}(\mathfrak{p}^{f(\pi_{\mathfrak{p}})}) \rightarrow \bar{\mathbb{Q}}$$

which are supported on  $B(\mathcal{O}_{\mathfrak{p}}) \xi K_{\mathfrak{p},1}(\mathfrak{p}^{f(\pi_{\mathfrak{p}})})$  and satisfy

$$f_{\xi}(b_1 \xi k) = \chi_{\mathfrak{p}}(b_1) f_{\xi}(\xi) \quad \forall b_1 \in B(\mathcal{O}_{\mathfrak{p}}), k \in K_{\mathfrak{p},1}(\mathfrak{p}^{f(\pi_{\mathfrak{p}})})$$

This is a very restrictive condition if we want  $f_{\xi} \neq 0$ , actually it follows from the definition of  $f(\pi_{\mathfrak{p}})$  and the above theorem of Casselmann and Novovorskii( that there is exactly one double coset  $\xi_0$  for which we find a function  $f_{\xi_0} = f_0 \neq 0$ . We have several cases.

a) If  $f(\pi_{\mathfrak{p}}) = 0$  then  $K_{\mathfrak{p},1}(\mathfrak{p}^{f(\pi_{\mathfrak{p}})}) = \text{Gl}_2(\mathcal{O}_{F_0,\mathfrak{p}})$  and the character  $\chi_{\mathfrak{p}}$  is unramified. In this case we choose for the function  $f_0$  the spherical function which has value one at the identity element.

b) We have  $f(\pi_{\mathfrak{p}}) > 0$ . Assume we have double coset  $\xi$  and a function  $f_{\xi} \neq 0$ . Assume this coset is of the form  $\xi = \begin{pmatrix} 1 & 0 \\ \varpi_{\mathfrak{p}}^{\nu} & 1 \end{pmatrix}$ . For any  $\varepsilon \in \mathcal{O}_{F_0,\mathfrak{p}}^{\times}$  we have

$$f_{\xi}\left(\begin{pmatrix} 1 & 0 \\ \varpi_{\mathfrak{p}}^{\nu} \varepsilon & 1 \end{pmatrix}\right) = f_{\xi}\left(\begin{pmatrix} 1 & 0 \\ 0 & \varepsilon \end{pmatrix} \begin{pmatrix} 1 & 0 \\ \varpi_{\mathfrak{p}}^{\nu} & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 & 0 \\ 0 & \varepsilon^{-1} \end{pmatrix}\right) = \chi_{2,\mathfrak{p}}(\varepsilon) f_{\xi}\left(\begin{pmatrix} 1 & 0 \\ \varpi_{\mathfrak{p}}^{\nu} & 1 \end{pmatrix}\right). \quad (8.126)$$

For  $\varepsilon \equiv 1 \pmod{\mathfrak{p}^{f(\pi_{\mathfrak{p}}) - \nu}}$  the far most left term does not depend on  $\varepsilon$  hence we can conclude that  $\chi_{2,\mathfrak{p}}(\varepsilon) = 1$  or what amounts to the same

$$f(\chi_{2,\mathfrak{p}}) \geq f(\pi_{\mathfrak{p}}) - \nu.$$

On the other hand for any  $v \in \mathcal{O}_{F_0, \mathfrak{p}}$  we have the equality

$$f_\xi \left( \begin{pmatrix} 1 & 0 \\ \varpi_\mathfrak{p}^\nu & 1 \end{pmatrix} \right) = f_\xi \left( \begin{pmatrix} 1 & 0 \\ \varpi_\mathfrak{p}^\nu & 1 \end{pmatrix} \begin{pmatrix} 1 & v \\ 0 & 1 \end{pmatrix} \right). \quad (8.127)$$

and

$$\begin{pmatrix} 1 & 0 \\ \varpi_\mathfrak{p}^\nu & 1 \end{pmatrix} \begin{pmatrix} 1 & v \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} (1 + v\varpi_\mathfrak{p}^\nu)^{-1} & * \\ 0 & (1 + v\varpi_\mathfrak{p}^\nu) \end{pmatrix} \begin{pmatrix} 1 & 0 \\ \frac{\varpi_\mathfrak{p}^\nu}{1 + v\varpi_\mathfrak{p}^\nu} & 1 \end{pmatrix}$$

Therefore we get

$$f_\xi \left( \begin{pmatrix} 1 & 0 \\ \varpi_\mathfrak{p}^\nu & 1 \end{pmatrix} \right) = \chi_{1, \mathfrak{p}}^{-1}(1 + v\varpi_\mathfrak{p}^\nu) f_\xi \left( \begin{pmatrix} 1 & 0 \\ \varpi_\mathfrak{p}^\nu & 1 \end{pmatrix} \right) \quad (8.128)$$

and this implies  $d_\mathfrak{p}(\chi_{1, \mathfrak{p}}) \leq \nu$ . Hence we see that  $\nu$  must satisfy

$$\mathfrak{f}(\chi_{1, \mathfrak{p}}) \geq \nu \geq \mathfrak{f}(\pi_\mathfrak{p}) - \mathfrak{f}(\chi_{2, \mathfrak{p}})$$

If on the other hand  $\nu$  satisfies this inequality we can write down a non zero function  $f_\xi$ . But since the  $\nu$  is unique we find that actually nufone

$$\mathfrak{f}(\pi_\mathfrak{p}) = \mathfrak{f}(\chi_{2, \mathfrak{p}}) + \mathfrak{f}(\chi_{1, \mathfrak{p}}) \text{ and } \nu = \mathfrak{f}(\chi_{1, \mathfrak{p}}). \quad (8.129)$$

Therefore we normalise the generator  $f_0$  so that  $f_0 \left( \begin{pmatrix} 1 & 0 \\ \varpi_\mathfrak{p}^{\nu_0} & 1 \end{pmatrix} \right) = 1$

We still have the double coset  $\xi_0 = w$ . Then

$$f_w(w) = f_{\xi_0} \left( w \begin{pmatrix} 1 & 0 \\ 0 & \varepsilon \end{pmatrix} \right) = f_w \left( \begin{pmatrix} \varepsilon & 0 \\ 0 & 1 \end{pmatrix} w \right) = \chi_{1, \mathfrak{p}}(\varepsilon) f_w(w)$$

and this implies  $\chi_{1, \mathfrak{p}}(\varepsilon) = 1$ . We normalise  $f_w(w) = 1$ . We are still in the case b) i.e.  $\mathfrak{f}(\chi_{2, \mathfrak{p}}) = \mathfrak{f}(\pi_\mathfrak{p}) > 0$ .

Now we leave case b), we consider the case that  $\chi_\mathfrak{p}$  is unramified. Here we have to be alert in the exceptional case that  $\chi_\mathfrak{p}^{(1)} = \mathbf{1}$  (resp.  $= | \cdot |_\mathfrak{p}$ ). In this case  $\text{Ind}_{B(F_\mathfrak{p})}^{G(F_\mathfrak{p})} \chi_\mathfrak{p}$  is not irreducible and has the Steinberg module  $\pi_\mathfrak{p} = \text{St}(\chi_\mathfrak{p})$  as quotient (resp. submodule). Then  $\mathfrak{f}(\pi_\mathfrak{p}) = 1$  and the Bruhat decomposition  $\text{Gl}_2(\mathcal{O}_\mathfrak{p}) = B(\mathcal{O}_\mathfrak{p}) \cup B(\mathcal{O}_\mathfrak{p})wK_{\mathfrak{p}, 0}(\mathfrak{p})$  gives us

$$\left( \text{Ind}_{B(F_\mathfrak{p})}^{G(F_\mathfrak{p})} \chi_\mathfrak{p} \right)^{K_{\mathfrak{p}, 0}(\mathfrak{p})} = \bar{\mathbb{Q}}f_e + \bar{\mathbb{Q}}f_w, \quad (8.130)$$

here  $f_e$  resp.  $f_w$  are the characteristic functions of  $B(\mathcal{O}_\mathfrak{p})$  resp.  $B(\mathcal{O}_\mathfrak{p})wK_{\mathfrak{p}, 0}(\varpi_\mathfrak{p})$ . The element  $f_e + f_w$  spans the one dimensional kernel of  $\text{Ind}_{B(F_\mathfrak{p})}^{G(F_\mathfrak{p})} \chi_\mathfrak{p} \rightarrow \text{St}(\chi_\mathfrak{p})$ , hence the image of  $f_e$  spans  $\text{St}(\chi_\mathfrak{p})^{K_{\mathfrak{p}, 0}(\mathfrak{p})}$ . If we realise  $\text{St}(\chi_\mathfrak{p})$  as kernel of  $\text{Ind}_{B(F_\mathfrak{p})}^{G(F_\mathfrak{p})} | \cdot |_\mathfrak{p} \chi_\mathfrak{p}^{-1} \rightarrow \bar{\mathbb{Q}} | \cdot |_\mathfrak{p} \chi_\mathfrak{p}^{-1}$  then  $f_0 := f_e - \frac{1}{q} f_w$  can be chosen as generator of  $\text{St}(\chi_\mathfrak{p})^{K_{\mathfrak{p}, 0}(\mathfrak{p})}$ .

We resume the computation of the  $R_\mathfrak{p}(f_0)$  after equation (8.123) we start with the case  $\chi_\mathfrak{p}$  is unramified. Let  $f$  be any linear combination of these two

functions, for  $\nu \geq 1$  we have  $f\left(\begin{pmatrix} 1 & 0 \\ \varpi_p^\nu \varepsilon^{-1} & 1 \end{pmatrix}\right) = f\left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}\right)$  and (8.123) becomes

$$\begin{aligned} R_{\mathfrak{p}}(f)\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right) &= |t|_{\mathfrak{p}} \chi_{2,\mathfrak{p}}(t) \times [f(w) \int_{\mathcal{O}_{F_0,\mathfrak{p}}} \overline{\psi_{\mathfrak{p}}(tu)} du + \\ & f\left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}\right) \sum_{\nu=1}^{\infty} N(\mathfrak{p})^{\nu} \chi_{\mathfrak{p}}^{(1)}(\varpi_{\mathfrak{p}}^{\nu}) \int_{\mathcal{O}_{F_0,\mathfrak{p}}^{\times}} \overline{\psi_{\mathfrak{p}}(t\varpi_{\mathfrak{p}}^{-\nu}u)} du] \end{aligned} \quad (8.131)$$

We have our above formulas for the integrals and get: If we write  $t = \varpi_{\mathfrak{p}}^{\text{ord}_{\mathfrak{p}}(t)}$  then

$$\begin{aligned} R_{\mathfrak{p}}(f)\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right) &= |t|_{\mathfrak{p}} \chi_{2,\mathfrak{p}}(\varpi_{\mathfrak{p}})^{\text{ord}_{\mathfrak{p}}(t)} \times [f(w) \int_{\mathcal{O}_{F_0,\mathfrak{p}}} \overline{\psi_{\mathfrak{p}}(tu)} du \\ & + f(e)(1 - \frac{1}{N(\mathfrak{p})}) \left( \sum_{\nu=1}^{\text{ord}_{\mathfrak{p}}(t)-d_{\mathfrak{p}}} N(\mathfrak{p})^{\nu} \chi_{\mathfrak{p}}^{(1)}(\varpi_{\mathfrak{p}})^{\nu} - N(\mathfrak{p})^{\text{ord}_{\mathfrak{p}}(t)} \chi_{\mathfrak{p}}^{(1)}(\varpi_{\mathfrak{p}})^{\text{ord}_{\mathfrak{p}}(t)+1} \right)] \end{aligned} \quad (8.132)$$

We consider the special case  $f_0 = f_e + f_w$  then a simple manipulation gives us

Wsph

$$\begin{aligned} R_{\mathfrak{p}}(f_0)\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right) &= \\ & (1 - \chi_{\mathfrak{p}}^{(1)}(\varpi_{\mathfrak{p}})) N(\mathfrak{p})^{-\text{ord}_{\mathfrak{p}}(t)} \left( \sum_{\nu=0}^{\text{ord}_{\mathfrak{p}}(t)-d_{\mathfrak{p}}} (N(\mathfrak{p}) \chi_{1,\mathfrak{p}}(\varpi_{\mathfrak{p}}))^{\nu} \chi_{2,\mathfrak{p}}(\varpi_{\mathfrak{p}})^{\text{ord}_{\mathfrak{p}}(t)-\nu} = \right. \\ & \left. (1 - \chi_{\mathfrak{p}}^{(1)}(\varpi_{\mathfrak{p}})) (N(\mathfrak{p}) \chi_{2,\mathfrak{p}}(\varpi_{\mathfrak{p}}))^{d_{\mathfrak{p}}} \left( \sum_{\nu=0}^{\text{ord}_{\mathfrak{p}}(t)-d_{\mathfrak{p}}} \chi_{1,\mathfrak{p}}(\varpi_{\mathfrak{p}})^{\nu} (N(\mathfrak{p})^{-1} \chi_{2,\mathfrak{p}}(\varpi_{\mathfrak{p}}))^{\text{ord}_{\mathfrak{p}}(t)-d_{\mathfrak{p}}-\nu} \right) \right) \end{aligned} \quad (8.133)$$

We look back to section 4.1.6. There  $\mathfrak{p}$  was a rational prime  $p$  and we introduced the spherical Whittaker function  $h_{\pi_p}^{(0)} \in \mathcal{W}(\pi_p, \psi_p)^{\text{G}1_2(\mathbb{Z}_p)}$  we normalised it so it takes value one at one, its values on the torus  $T_1(\mathbb{Q}_p)$  were encoded in formula (4.134).

Essentially the same is true in this more general situation. In the above special case we had  $d_p = 0$ , here we may have  $d_{\mathfrak{p}} \neq 0$ .

Assume  $\chi_{\mathfrak{p}}$  is unramified and  $\text{Ind}_{B(F_{\mathfrak{p}})}^{G(F_{\mathfrak{p}})} \chi_{\mathfrak{p}}$  is irreducible, we define  $h_{\pi_{\mathfrak{p}}}^{(0)}$  by the equation (recall that  $\text{Ind}_{B(F_{\mathfrak{p}})}^{G(F_{\mathfrak{p}})} \chi_{\mathfrak{p}}$  irreducible  $\iff \chi^{(1)}(\varpi_{\mathfrak{p}}) \neq \mathbf{1}$  or  $|\cdot|_{\mathfrak{p}}$ )

$$R_{\mathfrak{p}}(f_0)\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right) := (1 - \chi_{\mathfrak{p}}^{(1)}(\varpi_{\mathfrak{p}})) (N(\mathfrak{p}) \chi_{2,\mathfrak{p}}(\varpi_{\mathfrak{p}}))^{d_{\mathfrak{p}}} h_{\pi_{\mathfrak{p}}}^{(0)}\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right) \quad (8.134)$$

We get the following identity of formal power series PSI

$$\sum_{\nu=0}^{\infty} h_{\pi_{\mathfrak{p}}}^{(0)}\left(\begin{pmatrix} \varpi_{\mathfrak{p}}^{\nu-d_{\mathfrak{p}}} & 0 \\ 0 & 1 \end{pmatrix}\right) q^{\nu} = \frac{1}{(1 - \chi_{1,\mathfrak{p}}(\varpi_{\mathfrak{p}})q)(1 - N(\mathfrak{p})^{-1} \chi_{2,\mathfrak{p}}(\varpi_{\mathfrak{p}})q)} \quad (8.135)$$

If  $\chi_{\mathfrak{p}}^{(1)} = \mathbf{1}$  then  $\text{Ind}_{B(F_{\mathfrak{p}})}^{G(F_{\mathfrak{p}})}\chi_{\mathfrak{p}}$  is reducible and  $R_{\mathfrak{p}}(f_0) = 0$ , the function  $f_0$  generates the kernel of  $R_{\mathfrak{p}}$ . The quotient  $\text{Ind}_{B(F_{\mathfrak{p}})}^{G(F_{\mathfrak{p}})}\chi_{\mathfrak{p}}/Ff_0$  is the Steinberg module. Now we go back to (8.131) and we evaluate  $R_{\mathfrak{p}}$  at the function  $f_w$ . Then  $f_w\left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}\right) = 0$  and the terms in the summation over  $\nu$  are zero. Therefore we get PsSt

$$R_{\mathfrak{p}}(f_w)\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right) = N(\mathfrak{p})^{-\text{ord}_{\mathfrak{p}}(t)}\chi_{2,\mathfrak{p}}(\varpi_{\mathfrak{p}})^{\text{ord}_{\mathfrak{p}}(t)}\int_{\mathcal{O}_{F_0,\mathfrak{p}}} \overline{\psi_{\mathfrak{p}}(tu)}du \quad (8.136)$$

We choose as canonical generator

$$h_{\pi_{\mathfrak{p}}}^{(0)} = N(\mathfrak{p})^{d_{\mathfrak{p}}}\chi_{2,\mathfrak{p}}(\varpi_{\mathfrak{p}})^{-d_{\mathfrak{p}}}R_{\mathfrak{p}}(f_w). \quad (8.137)$$

and again we get an identity for power series

$$\sum_{\nu=0}^{\infty} h_{\pi_{\mathfrak{p}}}^{(0)}\left(\begin{pmatrix} \varpi_{\mathfrak{p}}^{\nu-d_{\mathfrak{p}}} & 0 \\ 0 & 1 \end{pmatrix}\right)q^{\nu} = \frac{1}{1 - N(\mathfrak{p})^{-1}\chi_{2,\mathfrak{p}}(\varpi_{\mathfrak{p}})q} \quad (8.138)$$

Now we consider the case that  $\chi_{\mathfrak{p}}$  is ramified. We compute the value  $R_{\mathfrak{p}}(f_{\xi})$ , for  $\xi$  running through the elements in (8.125). We begin with the case that  $\chi_{1,\mathfrak{p}}$  is unramified, we can take  $\xi = w$ . Then all the terms in the summation over  $\nu$  are equal to zero (8.123), we normalise  $f_w(w) = 1$  and get

$$R_{\mathfrak{p}}(f_w)\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right) = |t|_{\mathfrak{p}}\chi_{2,\mathfrak{p}}(t)\int_{\mathcal{O}_{F_0,\mathfrak{p}}} \psi_{\mathfrak{p}}(ut)du \quad (8.139)$$

Consequently we define in this case

$$h_{\pi_{\mathfrak{p}}}^{(0)}\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right) = N(\mathfrak{p})^{d_{\mathfrak{p}}}\chi_{2,\mathfrak{p}}(t)R_{\mathfrak{p}}(f_w)\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right) \quad (8.140)$$

We see that in all cases the support of the restriction of  $h_{\pi_{\mathfrak{p}}}^{(0)}$  to  $T_1(F_{0,\mathfrak{p}}) = F_{0,\mathfrak{p}}^{\times}$  in the set  $\{t \mid \text{ord}_{\mathfrak{p}}(t) \geq -d_{\mathfrak{p}}\}$  and the value of  $h_{\pi_{\mathfrak{p}}}^{(0)}$  on  $\varpi_{\mathfrak{p}}^{-d_{\mathfrak{p}}}$  is 1 (or at least a unit).

We look at the case  $\xi_{\nu_0} = \begin{pmatrix} 1 & 0 \\ \varpi_{\mathfrak{p}}^{\nu_0} & 1 \end{pmatrix}$ , we normalise  $f_{\xi_{\nu_0}}(\xi_{\nu_0}) = 1$ . In the summation in formula (8.116) the only (possibly) non zero term is  $\nu = \nu_0$ . Hence the value of the sum is xinu

$$R_{\mathfrak{p}}(f_{\xi_{\nu_0}})\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right) = |t|_{\mathfrak{p}}\chi_{2,\mathfrak{p}}(t)N(\mathfrak{p})^{\nu_0}\chi_{\mathfrak{p}}^{(1)}(\varpi_{\mathfrak{p}})^{\nu_0}\int_{\mathcal{O}_{F_0,\mathfrak{p}} \setminus \varpi_{\mathfrak{p}}\mathcal{O}_{F_0,\mathfrak{p}}} \chi_{1,\mathfrak{p}}(u)^{-1}\overline{\psi_{\mathfrak{p}}(t\varpi_{\mathfrak{p}}^{-\nu_0}u)}du. \quad (8.141)$$

The last integral is essentially a Gaussian sum, we write

$$G(t\varpi_{\mathfrak{p}}^{-\nu_0}, \chi_{1,\mathfrak{p}}, \psi_{\mathfrak{p}}) = N(\mathfrak{p})^{f(\chi_{1,\mathfrak{p}})}\int_{\mathcal{O}_{F_0,\mathfrak{p}}^{\times}} \chi_{1,\mathfrak{p}}(u)^{-1}\overline{\psi_{\mathfrak{p}}(t\varpi_{\mathfrak{p}}^{-\nu_0}u)}du \quad (8.142)$$

These Gaussian sums are computed in any textbook on algebraic number theory. we refer to [60]. The Gaussian sum is zero unless the "additive" conductor of  $u \mapsto \psi_{\mathfrak{p}}(t\varpi_{\mathfrak{p}}^{-\nu_0}u)$  is equal to the conductor  $f(\chi_{1,\mathfrak{p}})$ , i.e. the Gaussian sum is zero unless

$$\text{ord}_{\mathfrak{p}}(t) + d_{\mathfrak{p}} = 0 \quad (8.143)$$

where we used (8.129). If this equality holds then a well known computation yields

$$G(t\varpi_{\mathfrak{p}}^{-\nu_0}, \chi_{1,\mathfrak{p}}, \psi_{\mathfrak{p}}) \overline{G(t\varpi_{\mathfrak{p}}^{-\nu_0}, \chi_{1,\mathfrak{p}}, \psi_{\mathfrak{p}})} = N(\mathfrak{p})^{f(\chi_{1,\mathfrak{p}})}$$

Here a short discussion about normalisations of measures is in order. Our measure  $du$  is additively invariant on  $F_{0,\mathfrak{p}}$  and  $\text{vol}_{du}(\mathcal{O}_{F_{0,\mathfrak{p}}}) = 1$ . The measure  $\frac{du}{|u|_{\mathfrak{p}}}$  is a multiplicatively invariant measure on  $F_{0,\mathfrak{p}}^{\times}$ , for the volume of the group of units we have

$$\text{vol}_{\frac{du}{|u|_{\mathfrak{p}}}}(\mathcal{O}_{F_{0,\mathfrak{p}}}^{\times}) = \left(1 - \frac{1}{N(\mathfrak{p})}\right).$$

We define the local Tamagawa measure on  $T(F_{0,\mathfrak{p}})$  by  $(1 - \frac{1}{N(\mathfrak{p})})^{-1} \frac{dt_{\mathfrak{p}}}{|t_{\mathfrak{p}}|_{\mathfrak{p}}}$  it gives volume one to  $\mathcal{O}_{F_{0,\mathfrak{p}}}^{\times}$ , and we define the Tamagawa measure

$$d_{\text{Tam}}t_f = \prod_{\mathfrak{p}} \left(1 - \frac{1}{N(\mathfrak{p})}\right)^{-1} \frac{dt}{|t|_{\mathfrak{p}}}.$$

Then we have by definition  $\text{vol}_{d_{\text{Tam}}t_f} T(\hat{\mathcal{O}}_{F_{0,\mathfrak{p}}}^{\times}) = 1$ .

Any residue class  $x + \varpi_{\mathfrak{p}}^{f(\chi_{1,\mathfrak{p}})}$  has volume  $(\frac{1}{N(\mathfrak{p})})^{f(\chi_{1,\mathfrak{p}})}$  for the measure  $du$  and therefore we can say that the Gaussian sum is equal to

$$G(t\varpi_{\mathfrak{p}}^{-\nu_0}, \chi_{1,\mathfrak{p}}, \psi_{\mathfrak{p}}) = \sum_{\varepsilon \in (\mathcal{O}_{F_{0,\mathfrak{p}}}/(\varpi_{\mathfrak{p}}^{f(\chi_{1,\mathfrak{p}})}))^{\times}} \chi_{1,\mathfrak{p}}(\varepsilon)^{-1} \overline{\psi_{\mathfrak{p}}(t\varpi_{\mathfrak{p}}^{-\nu_0}\varepsilon)}. \quad (8.144)$$

if (8.143) holds and zero otherwise. Then it is clear that the Gaussian sum is an algebraic integer. If we replace  $t$  by  $t\eta$  with  $\eta \in F_{0,\mathfrak{p}}^{\times}$  then clearly  $\overline{\text{teta}}$

$$G(t\eta\varpi_{\mathfrak{p}}^{-\nu_0}, \chi_{1,\mathfrak{p}}, \psi_{\mathfrak{p}}) = \chi_{1,\mathfrak{p}}(\eta) G(t\varpi_{\mathfrak{p}}^{-\nu_0}, \chi_{1,\mathfrak{p}}, \psi_{\mathfrak{p}}) \quad (8.145)$$

This tells us that in this case  $h_{\pi_{\mathfrak{p}}}^{(0)}\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right)$  is supported on the annulus  $\varpi_{\mathfrak{p}}^{\nu_0+d_{\mathfrak{p}}-f\chi_{1,\mathfrak{p}}}\mathcal{O}_{F_{0,\mathfrak{p}}}^{\times}$  and again we can normalise by requiring  $h_{\pi_{\mathfrak{p}}}^{(0)}\left(\begin{pmatrix} \varpi_{\mathfrak{p}}^{\nu_0+d_{\mathfrak{p}}-f\chi_{1,\mathfrak{p}}} & 0 \\ 0 & 1 \end{pmatrix}\right) = 1$ .

Looking at these computation we easily see that the functions  $t \mapsto R_{\mathfrak{p}}(f_0)\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right)$  have a simple asymptotic behaviour. For  $|t| \rightarrow 0$ , the computations yield

$$R_{\mathfrak{p}}(f_0)\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right) \sim a|t|\chi_{2,\mathfrak{p}}(t) + b\chi_{1,\mathfrak{p}}(t) \quad (8.146)$$

We still have the supercuspidal representations  $\pi_{\mathfrak{p}}$ , these are the representations for which the Kirillov-model is  $\mathcal{S}(F_{0,\mathfrak{p}}^{\times})$ , we follow the argumentation in

[Cas]. Let  $n_0 = f(\pi_p)$  and let  $h_1$  be a generator of the one dimensional vector space  $H(\pi_p)^{K_{p,1}(\mathfrak{p}^{n_0})}$ . We consider the element  $W(\mathfrak{p}^{n_0}) = \begin{pmatrix} 0 & \varpi_p^{-n_0} \\ -1 & 0 \end{pmatrix}$  (the Atkin-Lehner involution), it conjugates  $K_{p,1}(\mathfrak{p}^{n_0})$  into  $K'_{p,1}(\mathfrak{p}^{n_0})$  where the condition  $a \equiv 1 \pmod{\mathfrak{p}^{n_0}}$  is replaced by  $d \equiv 1 \pmod{\mathfrak{p}^{n_0}}$ . Therefore  $\pi_p(W(\mathfrak{p}^{n_0})h_1) = h_2$  will be a generator of  $H(\pi_p)^{K'_{p,1}(\mathfrak{p}^{n_0})}$ . We look at the restriction of  $h_1$  to the annuli  $\varpi_p^\nu \mathcal{O}_{F_0,p}^\times$ . It is clear that  $h_1(\varpi_p^\nu \varepsilon) = h_1(\varpi_p^\nu) \zeta_p(\varepsilon)$ . Let us assume that  $d_p = 0$  then it is also clear that  $h_1(\varpi_p^\nu) = 0$  if  $\nu < 0$ . The computations in [Cas] yield

$$W(\mathfrak{p}^{n_0})h_1\left(\begin{pmatrix} \varpi_p^\nu \varepsilon & 0 \\ 0 & 1 \end{pmatrix}\right) = C(\psi_p)h_1\left(\begin{pmatrix} \varpi_p^{-\nu} \varepsilon^{-1} & 0 \\ 0 & 1 \end{pmatrix}\right)$$

The right hand side is zero if  $-\nu < 0$  hence we see that the values of  $h_1, h_2$  on an annulus  $\varpi_p^\nu \mathcal{O}_{F_0,p}^\times$  is zero unless  $\nu = 0$ . Hence we define the generator  $h_{\pi_p}^{(0)}$  by the requirement that it assumes the value 1 at the identity element. If  $d_p >= 0$  we use the isomorphism in equation 8.110. It tells us that

$$g \mapsto h_{\pi_p}^{(0)}\left(\begin{pmatrix} \varpi_p^{-d_p t} & 0 \\ 0 & 1 \end{pmatrix} g\right) \in \mathcal{W}(\pi_p, \psi_v^{[-d_p]}) \tag{8.147}$$

and since  $\psi^{[-d_p f p]}$  has additive character 0 we conclude that  $t \mapsto h_{\pi_p}^{(0)}\left(\begin{pmatrix} \varpi_p^{-d_p t} & 0 \\ 0 & 1 \end{pmatrix} t\right)$  is supported on the annulus  $\mathcal{O}_{F_0,v}^\times$ . Therefore we can normalise  $h_{\pi_p}^{(0)}$  by requiring

$$h_{\pi_p}^{(0)}\left(\begin{pmatrix} \varpi_p^{-d_p} & 0 \\ 0 & 1 \end{pmatrix}\right) = 1.$$

For any irreducible  $\pi_p$  we introduce the number  $\epsilon(e_p)$ , this is the smallest integer for which the value  $h_{\pi_p}^{(0)}\left(\begin{pmatrix} \varpi_p^{\epsilon(e_p)} & 0 \\ 0 & 1 \end{pmatrix}\right) \neq 0$  and we normalise such that this value is actually equal to 1. It does not depend on the choice of the generator

**Periods again**

Now that we have chosen a generator  $h_{\pi_p}^{(0)}$  at all finite places we choose generators in  $\bigotimes_{v \in S_\infty} H^1(\mathfrak{g}_v, K_v, \tilde{\mathcal{D}}_{\lambda_v} \otimes \mathcal{M}_{\lambda_v})$  and of course these generators will be tensor product of the generators in the factors.

If  $v \in S_\infty$  is real, then the maximal compact subgroup  $K_v^*$  containing  $K_v$  is not connected. As before we denote the Whittaker realisation of  $\mathcal{D}_{\lambda_v}$  by  $\tilde{\mathcal{D}}_{\lambda_v}$ . The module  $\tilde{\mathcal{D}}_{\lambda_v}$  is a sum of two copies which are switched under the action of  $K_v^*/K_v$ . The space

$$H^1(\mathfrak{g}_v, K_v, \tilde{\mathcal{D}}_{\lambda_v} \otimes \mathcal{M}_{\lambda_v}) = \text{Hom}_{K_v}(\Lambda^1(\mathfrak{g}/\mathfrak{k}), \tilde{\mathcal{D}}_{\lambda_v} \otimes \mathcal{M}_{\lambda_v})$$

is the direct sum of a + and a - eigenspace, both of dimension 1. In (4.133) we wrote down generators of these one dimensional spaces

$$\omega_+^\dagger = \frac{1}{2}(\omega^\dagger + i^n \bar{\omega}^\dagger); \quad \omega_-^\dagger = \frac{1}{2}(\omega^\dagger - i^n \bar{\omega}^\dagger) \tag{8.148}$$

The choice of these generators depends on the choice of several specific basis elements. To justify the selection of these basis elements we can put a  $\mathbb{Z}$ -module structures on  $\mathcal{D}_v, \mathfrak{g}/\mathfrak{k}$ , (See [33]). The module  $\mathcal{M}_{\lambda^v}$  has a  $\mathbb{Z}$ -module structure by definition and the choices become natural.

We do essentially the same for a complex place  $v$ , we have seen that for  $i = 1, 2$  the space  $\text{Hom}_{K_\infty}(\Lambda^i(\mathfrak{g}/\mathfrak{k})_R, \tilde{\mathcal{D}}_v \otimes \mathcal{M}_{\lambda^v})$  is of dimension one. For the elements  $e_\mu$  in (??) we can choose tensor product of monomials  $X^{n-\nu}Y^\nu \otimes \bar{X}^{n-\bar{\nu}}\bar{Y}^{\bar{\nu}}$ . Then we require that our generator  $\omega^{1,\dagger}$  in degree one satisfies

$$\int_0^\infty \langle \omega_\epsilon^{1,\dagger}, H \otimes X^n \otimes \bar{X}^n \rangle = \frac{\Gamma(2n+2)}{(2\pi)^{2n+2}} \quad (8.149)$$

In degree two we use the isomorphism  $\kappa : \Lambda^1(\mathfrak{g}/\mathfrak{k}) \xrightarrow{\sim} \Lambda^2(\mathfrak{g}/\mathfrak{k})$  we define  $\omega^{2,\dagger} = {}^t \kappa^{-1}(\omega^{1,\dagger})$ .

Let  $\nu = 1$  or  $2$  and  $\underline{\epsilon} = \prod_{v \in S_\infty, \text{real}} \epsilon_v$  be a character

$$\underline{\epsilon} : \pi_0(G(\mathbb{R})) = \prod_{v \in S_\infty, \text{real}} K_v^*/K_v \rightarrow \mathbb{C}^\times$$

then

$$\omega_{(\underline{\epsilon})}^{\nu,\dagger} \in \text{Hom}_{K_\infty}(\Lambda^{r_1+\nu r_2}(\mathfrak{g}/\mathfrak{k}), \mathcal{D}_\lambda^\underline{\epsilon} \otimes \mathcal{M}_\lambda)$$

will be the product over the  $v \in S_\infty$  where the local factor is  $\omega_v^\dagger$  for a real place and  $\omega^{(\nu),\dagger}$  for a complex place.

Now we have constructed an isomorphism between  $G(\mathbb{A}_f)$  modules

$$\mathcal{F}_1^{(\bullet)}(\omega_\epsilon^{\nu,\dagger}) : \bigotimes_{\mathfrak{p}} \mathcal{W}(\pi_{\mathfrak{p}}, \psi_{\mathfrak{p}}) \otimes \mathbb{C} \longrightarrow H^{r_1+\nu r_2}(\mathcal{S}^G, \mathcal{M}_\lambda)(\epsilon \times \pi_f) \otimes \mathbb{C},$$

The two vector spaces  $\mathcal{W}(\pi_{\mathfrak{p}}, \psi_{\mathfrak{p}}), H^{r_1+\nu r_2}(\mathcal{S}^G, \mathcal{M}_\lambda)(\epsilon \times \pi_f)$  are  $\mathbb{Q}(\pi_f)$  vector spaces, and hence we can define a complex numbers  $\Omega^{(\nu)}(\underline{\epsilon} \times \pi_f)$  such that

$$\frac{1}{\Omega^{(\nu)}(\underline{\epsilon} \times \pi_f)} \cdot \mathcal{F}_1^{(1)}(\omega_\epsilon^\dagger) : \bigotimes_{\mathfrak{p}} \mathcal{W}_{\mathcal{O}(\pi_f)}(\pi_f, \psi_{\mathfrak{p}}) \xrightarrow{\sim} H^{r_1+\nu r_2}(\mathcal{S}^G, \mathcal{M})(\epsilon \times \pi_f), \quad (8.150)$$

these numbers are well defined modulo an element in  $\mathbb{Q}(\pi_f)^\times$ .

But we can do better. We choose a level subgroup, actually we choose  $K_f = \prod_{\mathfrak{p}} K_{\mathfrak{p},1}(\mathfrak{p}^{f(\pi_{\mathfrak{p}})})$ , this is in a sense the optimal level for  $\pi_f$ . Then

$$H^{r_1+\nu r_2}(\mathcal{S}_{K_f}^G, \mathcal{M} \otimes \mathcal{O}_F)(\epsilon \times \pi_f)_{\text{int}} \subset H^{r_1+\nu r_2}(\mathcal{S}_{K_f}^G, \mathcal{M} \otimes F^\dagger)(\epsilon \times \pi_f)$$

is a locally free  $\mathcal{O}_F$  module of rank 1. Hence we can find a covering  $\cup_\mu U_\mu = \text{Spec}(\mathcal{O}_F)$  by Zariski open subset such that

$$H^{r_1+\nu r_2}(\mathcal{S}_{K_f}^G, \mathcal{M} \otimes \mathcal{O}_F(U_\mu))(\pi_f) = \mathcal{O}_F(U_\mu)e_\mu$$

is actually free of rank one.

On the other hand we have chosen specific generators  $h_{\pi_{\mathfrak{p}}}^{(0)} \in \mathcal{W}(\pi_{\mathfrak{p}}^{K_{\mathfrak{p}}}, \psi_{\mathfrak{p}})$  and hence  $h_{\pi_f}^{(0)} \in \mathcal{W}(\pi_f^{K_f}, \psi_f)$ . Then we can define periods locally over  $U_{\mu}$  by requiring

$$\frac{1}{\Omega_{\underline{\epsilon}}^{(\nu)}(\pi_f, \mu)} \mathcal{F}_1^{(\nu)}(\omega_{\underline{\epsilon}}^{(\nu)}(h_f)) = e_{\mu}$$

these periods are now well defined up to an element of  $\mathcal{O}_F(U_{\mu})^{\times}$ . Moreover it is clear that for two indices  $\mu_1, \mu_2$  we have

$$\Omega_{\underline{\epsilon}}^{(\nu)}(\pi_f, \mu_1) = \Omega_{\underline{\epsilon}}^{(\nu)}(\pi_f, \mu_2)c_{1,2}, \text{ with } c_{1,2} \in \mathcal{O}_F(U_{\mu_1} \cap U_{\mu_2})^{\times}.$$

For the following we assume for simplicity that  $\mathcal{O}_F$  has class number one, then we can remove the argument  $\mu$  in the periods and we denote them as  $\Omega_{\underline{\epsilon}}^{(\nu)}(\pi_f)$

These numbers are the periods attached to the data  $\pi_f, \underline{\epsilon}, \nu$ .

We may also look at the conjugates of  $\dots \pi_f^{\sigma} \dots$  of  $\pi_f$ . We can choose these periods consistently (see [Ha-Mod]) and hence we even get a period vector

$$\Omega_{\epsilon}(\Pi_f)^{-1} = (\dots \Omega_{\epsilon}(\pi_f^{\sigma})^{-1} \dots)_{\sigma: \mathbb{Q}(\pi_f) \rightarrow \mathbb{C}}.$$

Let us assume that we have an isotopical component  $H_1^{r_1+r_2}(\mathcal{S}^G, \tilde{\mathcal{M}}_{\mathbb{Q}(\pi_f)}^{\vee})(\pi_f)$ , then we can consider the composition

$$J(\phi_{\mu}, \epsilon \times \pi_f, \tilde{\mu}_f) \circ \frac{1}{\Omega^{(\nu)}(\underline{\epsilon} \times \pi_f)} \mathcal{F}_1^{(1)}(\omega_{\epsilon}) : \bigotimes_{\mathfrak{p}} \mathcal{W}_{\mathcal{O}(\pi_f)}(\pi_{\mathfrak{p}}, \psi_{\mathfrak{p}}) \longrightarrow \text{Ind}_{\tilde{H}(A_f)}^{\tilde{G}(A_f)} \tilde{\mu}_f^{-1}.$$

Since we will see in the following subsection that the condition  $(I_{pp})$  is satisfied and since we have some natural choices for the local intertwining operators, this comes down to the computation of a number and this number is expressible in terms of  $L$ -values, this is our ultimate goal.

**The local intertwining operators**

Next issue is to investigate the space of intertwining operators, i.e. we have to check  $(I_p)$  and  $(I_{pp})$  and to study the space

$$\text{Hom}_{G(F_{\mathfrak{p}})}(\mathcal{W}(\pi_{\mathfrak{p}}, \psi_{\mathfrak{p}}), \text{Ind}_{T(F_{\mathfrak{p}})}^{G(F_{\mathfrak{p}})} \tilde{\mu}_{\mathfrak{p}}^{-1}).$$

Of course we need to assume that the central character  $\zeta_{\mathfrak{p}}$  is equal to the character  $\tilde{\mu}_{\mathfrak{p}}$  restricted to the centre. We restrict the functions in  $\mathcal{W}(\pi_{\mathfrak{p}}, \psi_{\mathfrak{p}})$  to  $T_1(F_{\mathfrak{p}})$  the space of these restrictions is the Kirillov model  $\mathcal{K}(\pi_{\mathfrak{p}})$  this restriction map is injective ([18]) We introduce the subtorus

$$T_1(F_{\mathfrak{p}}) = \left\{ \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right\}$$

of  $T(F_{\mathfrak{p}})$ . Of course a function in  $\mathcal{K}(\pi_{\mathfrak{p}})$  is determined by its restriction to  $T_1(F_{\mathfrak{p}})$ , hence we may consider  $\mathcal{K}(\pi_{\mathfrak{p}})$  as a space of functions on  $T_1(F_{\mathfrak{p}}) = F_{\mathfrak{p}}^{\times}$ . The restriction  $\tilde{\mu}_{\mathfrak{p}}$  to this subgroup  $T_1(F_{\mathfrak{p}})$  is also called  $\tilde{\mu}_{\mathfrak{p}}$ , since the central character is given this restriction determines  $\tilde{\mu}_{\mathfrak{p}}$ . For  $t \in F_{\mathfrak{p}}^{\times}$ , we denote by  $[t]$  the matrix  $[t] = \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}$ . The space of Schwartz-functions  $\mathcal{S}(F_{\mathfrak{p}}^{\times})$  is

of codimension 0, 1 or 2 in  $\mathcal{K}(\pi_{\mathfrak{p}})$ , The action of  $T(F_{\mathfrak{p}})$  on  $\mathcal{K}(\pi_{\mathfrak{p}})$ , is given by  $\pi_{\mathfrak{p}}([t])(f)(x) = f(tx)$ , hence it is clear that the space  $\mathcal{S}(F_{\mathfrak{p}}^{\times})$  invariant under this action. Therefore we have an intertwining operator  $I_{\mathfrak{p}}$  from  $\mathcal{S}(F_{\mathfrak{p}}^{\times})$  to  $F[\tilde{\mu}_{\mathfrak{p}}]$  which given by

$$I_{\mathfrak{p}}(f)([t_0]) = \int_{T_1(F_{\mathfrak{p}})} f([t_0 t] \tilde{\mu}_{\mathfrak{p}}([t]) d^{\times} t,$$

this operator is unique up to a scalar. Here  $d^{\times} t$  is (momentarily ) the multiplicatively invariant measure which gives value one to  $\mathcal{O}_{\mathfrak{p}}^{\times}$ . We apply Frobenius and see

$$\mathrm{Hom}_{G(F_{\mathfrak{p}})}(\mathcal{W}(\pi_{\mathfrak{p}}, \psi_{\mathfrak{p}}), \mathrm{Ind}_{T(F_{\mathfrak{p}})}^{G(F_{\mathfrak{p}})} \tilde{\mu}_{\mathfrak{p}}^{-1}) = \mathrm{Hom}_{T(F_{\mathfrak{p}})}(\mathcal{W}(\pi_{\mathfrak{p}}, \psi_{\mathfrak{p}}), \tilde{\mu}_{\mathfrak{p}}^{-1}).$$

If  $\pi_{\mathfrak{p}}$  is supercuspidal this is our choice of a local intertwining operator  $I_{\mathfrak{p}}^{\mathrm{loc}}$  at  $\mathfrak{p}$  up to a normalisation of the measure

If  $\pi_{\mathfrak{p}}$  is not supercuspidal we have to discuss the question whether this operator has a (unique) extension to  $\mathcal{K}(\pi_{\mathfrak{p}}, \psi_{\mathfrak{p}})$ . Our representation  $\pi_{\mathfrak{p}}$  is either a induced representation  $\mathrm{Ind}_{B(F_{\mathfrak{p}})}^{G(F_{\mathfrak{p}})} \chi_{\mathfrak{p}}$  or it is a Steinberg representation. In both cases we can consider the quotient (See (8.146))

$$\mathcal{K}(\pi_{\mathfrak{p}})/\mathcal{S}(F_{\mathfrak{p}}) \xrightarrow{\sim} \begin{cases} F_{\mathfrak{p}} |t| \chi_{2,\mathfrak{p}}(t) \oplus F_{\mathfrak{p}} \chi_{1,\mathfrak{p}}(t) & \text{if } \pi_{\mathfrak{p}} \text{ is induced} \\ |t| & \text{if } \pi_{\mathfrak{p}} \text{ is Steinberg} \end{cases} \quad (8.151)$$

We consider the exact sequence

$$\begin{aligned} 0 \rightarrow \mathrm{Hom}_{T_1(F_{\mathfrak{p}})}(\mathcal{K}(\pi_{\mathfrak{p}})/\mathcal{S}(F_{\mathfrak{p}}), F[\tilde{\mu}_{\mathfrak{p}}^{-1}]) &\rightarrow \mathrm{Hom}_{T_1(F_{\mathfrak{p}}^{\times})}(\mathrm{Ind}_{B(F_{\mathfrak{p}})}^{G(F_{\mathfrak{p}})} F \chi_{\mathfrak{p}}, F[\tilde{\mu}_{\mathfrak{p}}^{-1}]) \rightarrow \\ &\rightarrow \mathrm{Hom}_{T_1(F_{\mathfrak{p}}^{\times})}(\mathcal{S}(F_{\mathfrak{p}}^{\times}), F[\tilde{\mu}_{\mathfrak{p}}^{-1}]) \rightarrow \mathrm{Ext}_{T_1(F_{\mathfrak{p}}^{\times})}^1(\mathcal{K}(\pi_{\mathfrak{p}})/\mathcal{S}(F_{\mathfrak{p}}), F[\tilde{\mu}_{\mathfrak{p}}^{-1}]) \end{aligned} \quad (8.152)$$

Now it is easy to see that the abelian groups

$$\mathrm{Hom}_{T_1(F_{\mathfrak{p}})}(\mathcal{K}(\pi_{\mathfrak{p}})/\mathcal{S}(F_{\mathfrak{p}}), F \tilde{\mu}_{\mathfrak{p}}^{-1}) \text{ and } \mathrm{Ext}_{T_1(F_{\mathfrak{p}}^{\times})}^1(\mathcal{K}(\pi_{\mathfrak{p}})/\mathcal{S}(F_{\mathfrak{p}}), \tilde{\mu}_{\mathfrak{p}}^{-1}) = 0$$

are both trivial unless we have

$$\chi_{1,\mathfrak{p}} |t| = \tilde{\mu}_{\mathfrak{p}}^{-1} \text{ or } \chi_{2,\mathfrak{p}} |t|^{-1} = \tilde{\mu}_{\mathfrak{p}}^{-1}. \quad \text{pole}$$

Hence we see that in case that (pole) is not true ( we do not have a pole ) the map

$$\mathrm{Hom}_{T_1(F_{\mathfrak{p}}^{\times})}(\mathrm{Ind}_{B(F_{\mathfrak{p}})}^{G(F_{\mathfrak{p}})} \chi_{\mathfrak{p}}, F[\tilde{\mu}_{\mathfrak{p}}^{-1}]) \rightarrow \mathrm{Hom}_{T_1(F_{\mathfrak{p}}^{\times})}(\mathcal{S}(F_{\mathfrak{p}}^{\times}), F[\tilde{\mu}_{\mathfrak{p}}^{-1}]) \quad (8.153)$$

is an isomorphism, our intertwining operator has a unique extension. We want a formula for this extension and consider the expression

$$\int_{T_1(F_{\mathfrak{p}})} f([t_0] \tilde{\mu}_{\mathfrak{p}}([t]) d^{\times} t \quad (8.154)$$

for all functions in  $f \in \mathcal{K}(\pi_{\mathfrak{p}})$ . Of course as it stands it does not always make sense. But given  $f$  we can find an integer  $N_0 > 0$  such that

$$f|\{[t] \in T_1(F_{\mathfrak{p}}) \mid |t|_{\mathfrak{p}} \leq N(\mathfrak{p})^{-N_0}\} = a|t|_{\mathfrak{p}}\chi_{2,\mathfrak{p}}(t) + b\chi_{1,\mathfrak{p}}(t). \tag{8.155}$$

Let us put  $c_0 = N(\mathfrak{p})^{-N_0}$  and let  $T_1(F_{\mathfrak{p}})(\leq c_0)$  be the above neighbourhood of 0. Then our above expression becomes

$$\begin{aligned} \int_{T_1(F_{\mathfrak{p}})} f([tt_0])\tilde{\mu}_{\mathfrak{p}}([t])d^{\times}t &= \int_{T_1(F_{\mathfrak{p}})(>c_0)} f([tt_0])\tilde{\mu}_{\mathfrak{p}}([t])d^{\times}t \\ &+ \int_{T_1(F_{\mathfrak{p}})(\leq c_0)} f([tt_0])\tilde{\mu}_{\mathfrak{p}}([t])d^{\times}t \end{aligned}$$

The first term is well defined, the second is equal to

$$\int_{T_1(F_{\mathfrak{p}})(\leq c_0)} (a|t|_{\mathfrak{p}}\chi_{2,\mathfrak{p}}(t) + b\chi_{1,\mathfrak{p}}(t))\tilde{\mu}_{\mathfrak{p}}([t])d^{\times}t \tag{8.156}$$

hence we have to "compute"

$$\int_{T_1(F_{\mathfrak{p}})(\leq c_0)} (|t|^{\pm}\chi_{j,\mathfrak{p}}(t)\tilde{\mu}_{\mathfrak{p}}([t])d^{\times}t \tag{8.157}$$

We identify  $F_{\mathfrak{p}}^{\times}$  to  $T_1(F_{\mathfrak{p}})$  via the map  $h$ . It is clear that the integral over an annulus

$$\int_{F_{\mathfrak{p}}^{\times}(|t|=N(\mathfrak{p})^N)} (|t|^m\chi_{j,\mathfrak{p}}(t)\tilde{\mu}_{\mathfrak{p}}(t)d^{\times}t = 0$$

if the character  $\chi_{j,\mathfrak{p}}(t)\tilde{\mu}_{\mathfrak{p}}(t)$  is ramified. Hence the expression in (8.157) simply has value zero in this case.

Let  $\eta_{\mathfrak{p}} : F_{\mathfrak{p}}^{\times} \rightarrow F^{\times}$  be any character ( which is the local component of an algebraic Hecke character). Then at least formally

$$\int_{F_{\mathfrak{p}}^{\times}(\leq c_0)} \eta_{\mathfrak{p}}(t)d^{\times}t = \sum_{\nu \geq N_0} \int_{F_{\mathfrak{p}}^{\times}(|t|=N(\mathfrak{p})^{-\nu})} \eta_{\mathfrak{p}}(t)d^{\times}t = \begin{cases} 0 & \text{if } \eta_{\mathfrak{p}} \text{ is ramified} \\ \frac{\eta_{\mathfrak{p}}(\varpi_{\mathfrak{p}})^{N_0}}{1-\eta_{\mathfrak{p}}(\varpi_{\mathfrak{p}})} & \text{if } \eta_{\mathfrak{p}} \text{ is unramified} \end{cases} \tag{8.158}$$

We introduce the local Euler factor

$$L(\eta_{\mathfrak{p}}, s) := \begin{cases} 1 & \text{if } \eta_{\mathfrak{p}} \text{ is ramified} \\ \frac{1}{1-\eta_{\mathfrak{p}}(\varpi_{\mathfrak{p}})N(\mathfrak{p})^{-s}} & \text{if } \eta_{\mathfrak{p}} \text{ is unramified} \end{cases}$$

here  $s$  is a complex variable, but we could as well interpret  $N(\mathfrak{p})^{-s}$  as a formal variable. Therefore we get

$$L(\pi_{\mathfrak{p}}, 0)^{-1} \int_{F_{\mathfrak{p}}^{\times}(\leq c_0)} \eta_{\mathfrak{p}}(t)d^{\times}t \tag{8.159}$$

is a finite sum and has a well defined value.

Let  $\chi_{\mathfrak{p}}$  be an unramified character and  $\pi_{\mathfrak{p}} = \text{Ind}_{B(F_{\mathfrak{p}})}^{G(F_{\mathfrak{p}})} \chi_{\mathfrak{p}}$  be an irreducible induced representation or  $\pi_{\mathfrak{p}} = \text{St}(\chi_{\mathfrak{p}})$ . For any  $\eta_{\mathfrak{p}} : T(F_{\mathfrak{p}}) \rightarrow E^{\times}$  as above we define localL

$$L(\pi_{\mathfrak{p}} \times \eta_{\mathfrak{p}}, s) = \begin{cases} L(|\cdot|_{\mathfrak{p}}^{-1} \chi_{1,\mathfrak{p}} \eta_{1,\mathfrak{p}}, N(\mathfrak{p})^{-s}) (L(\chi_{2,\mathfrak{p}} \eta_{2,\mathfrak{p}}, N(\mathfrak{p})^{-s}) & \text{if } \pi_{\mathfrak{p}} \text{ is induced} \\ L(|\cdot|_{\mathfrak{p}}^{-1} \chi_{1,\mathfrak{p}} \eta_{1,\mathfrak{p}}, N(\mathfrak{p})^{-s}) & \text{if } \pi_{\mathfrak{p}} \text{ is Steinberg} \end{cases} \quad (8.160)$$

If  $\chi_{\mathfrak{p}}$  is ramified but  $\chi_{1,\mathfrak{p}}$  is ramified and  $\pi_{\mathfrak{p}} = \text{Ind}_{B(F_{\mathfrak{p}})}^{G(F_{\mathfrak{p}})} \chi_{\mathfrak{p}}$  then we put

$$L(\pi_{\mathfrak{p}} \times \eta_{\mathfrak{p}}, s) = L(|\cdot|_{\mathfrak{p}}^{-1} \chi_{1,\mathfrak{p}} \eta_{1,\mathfrak{p}}, N(\mathfrak{p})^{-s}) \quad (8.161)$$

The above  $\pi_{\mathfrak{p}}$  are those irreducible representations for which  $h_{\pi_{\mathfrak{p}}}^{(0)}$  does not have compact support. For the other representations and especially for the cuspidal then we put  $L(\pi_{\mathfrak{p}} \otimes \eta_{\mathfrak{p}}, s) = 1$ .

Hence we define the local intertwining operator

localintop

$$\begin{aligned} I_{\mathfrak{p}}^{\text{loc}} : \mathcal{K}(\pi_{\mathfrak{p}}, \psi_{\mathfrak{p}}) &\rightarrow \text{Ind}_{T(F_{\mathfrak{p}})}^{G(F_{\mathfrak{p}})}(\tilde{\mu}_{\mathfrak{p}}) \\ f(\cdot) &\mapsto L(\pi_{\mathfrak{p}} \times \tilde{\mu}_{\mathfrak{p}}, 1)^{-1} \int_{T_1(F_{\mathfrak{p}})} f([t]\cdot) \tilde{\mu}_{\mathfrak{p}}([t]) d^{\times} t \end{aligned} \quad (8.162)$$

We remember that we want to study the integral cohomology, therefore a level subgroup  $K_{\mathfrak{p}}^T$  is given to ust, it has to satisfy the condition b) above. In this case our function  $f$  and  $\tilde{\mu}_{\mathfrak{p}}$  are invariant under  $K_f^T$  and then we renormalise our operator and define

$$I_{\mathfrak{p}}^{\text{loc}}(f)(g) = \frac{[T_1(\mathcal{O}_{F_0,\mathfrak{p}}) : K_{\mathfrak{p}}^{T_1}]}{L(\pi_{\mathfrak{p}} \times \tilde{\mu}_{\mathfrak{p}}, 1)} \int_{T_1(F_{\mathfrak{p}})} f([t]g) \tilde{\mu}_{\mathfrak{p}}([t]) d^{\times} t, \quad (8.163)$$

here the measure  $d^{\times} t = d_{\text{Tam}} t$ . Then the right hand side is actually a finite sum

$$I_{\mathfrak{p}}^{\text{loc}}(f)(g) = L(\pi_{\mathfrak{p}} \times \tilde{\mu}_{\mathfrak{p}}, 1)^{-1} \sum_{\varepsilon \in T_1(\mathcal{O}_{F_0,\mathfrak{p}})/K_{\mathfrak{p}}^{T_1}} f(\varepsilon g) \tilde{\mu}_{\mathfrak{p}}(\varepsilon) \quad (8.164)$$

From this we conclude that the local intertwining operator  $I^{\text{loc}}(\pi_{\mathfrak{p}}, \chi_{\tilde{\mu}_{\mathfrak{p}}^{-1}})$  is defined over  $\mathbb{Q}(\pi_{\mathfrak{p}}, \chi^{(1)})$  we get

$$I^{\text{loc}}(\pi_{\mathfrak{p}}, \tilde{\mu}_{\mathfrak{p}}^{-1}) : \mathcal{W}(\pi_{\mathfrak{p}}, \psi_{\mathfrak{p}})_{\mathbb{Q}(\pi_{\mathfrak{p}}, \chi^{(1)})} \rightarrow (\text{Ind}_{T(\mathbb{Q}_{\mathfrak{p}})}^{G(\mathbb{Q}_{\mathfrak{p}})} \tilde{\mu}_{\mathfrak{p}}^{-1})_{\mathbb{Q}(\pi_{\mathfrak{p}}, \chi^{(1)})}$$

If the conductor  $d_{\mathfrak{p}}(\psi_{\mathfrak{p}}) = 0$  it transforms the spherical function  $h_{\pi_{\mathfrak{p}}}^{(0)}$  into the spherical function in the induced module which also takes value one at the identity element.

We come to the final computation, for our pin point  $\underline{g}_f = \prod_{\mathfrak{p}} g_{\mathfrak{p}}$  and we have a level subgroup  $K_f^T = \prod_{\mathfrak{p}} K_{\mathfrak{p}}^T$  which satisfies b) with respect to this pin point. We know that the Manin-Drinfeld principle is valid in this case. Hence ( see also the argument further down)

$$\begin{aligned}
 & \langle j((x, \underline{g}_{\mathfrak{p}}), r_{\lambda, \mu})(\mathcal{F}^\bullet(\frac{[\omega_\epsilon]}{\Omega(\pi_f, \epsilon)} \times h_{\pi_f}^{(0)})), \tilde{\mu}_{\mathfrak{p}} \rangle = \\
 [T_1(\hat{\mathcal{O}}_{F_0}) : K_f^{T_1}] & \int_{T_1(\mathbb{Q}) \backslash T_1(\mathbb{A})} \langle \mathcal{F}(\frac{\omega_\epsilon}{\Omega(\underline{\epsilon} \times \pi_f)} \times h_{\pi_f}^{(0)})(\underline{t}g_{\mathfrak{p}}), \tilde{\mu}_{\mathfrak{p}} \rangle d_\infty^\times t_\infty \times d_{\text{Tam}} t_f = \\
 [T_1(\hat{\mathcal{O}}_{F_0}) : K_f^{T_1}] & \int_{T_1(\mathbb{Q}) \backslash T_1(\mathbb{A})} (\sum_{a \in T(\mathbb{Q})} \frac{\omega_\epsilon}{\Omega(\underline{\epsilon} \times \pi_f)} \times h_{\pi_f}^{(0)}(atg_{\mathfrak{p}}), \tilde{\mu}_{\mathfrak{p}} \rangle d_{\text{Tam}} t.
 \end{aligned} \tag{8.165}$$

Now we allow ourselves to interchange summation and integration then the last expression becomes

$$\begin{aligned}
 [T_1(\hat{\mathcal{O}}_{F_0}) : K_f^{T_1}] & \int_{T_1(\mathbb{A})} (\langle \frac{\omega_\epsilon}{\Omega(\underline{\epsilon} \times \pi_f)} \times h_{\pi_f}^{(0)}(\underline{t}g_{\mathfrak{p}}), \tilde{\mu}_{\mathfrak{p}} \rangle d_{\text{Tam}} t = \\
 \frac{1}{\Omega(\underline{\epsilon} \times \pi_f, \cdot)} & \prod_{v \in S_\infty} \int_{T_1(F_v)} \langle \omega_v^\dagger, e_{\mu_v} \rangle d_\infty^\times t_v \prod_{\mathfrak{p}} L(\pi_{\mathfrak{p}} \times \tilde{\mu}_{\mathfrak{p}}, 1) I^{\text{loc}}(h_{\mathfrak{p}}^{(0)})(g_{\mathfrak{p}})
 \end{aligned} \tag{8.166}$$

For the archimedean places  $v$  we denoted the Whittaker model of the representation  $\pi_v$  by  $\tilde{\mathcal{D}}_{\lambda_v}^\pm$ , for real places,  $\tilde{\mathcal{D}}_{\lambda_v}$  for complex places. We have the local Euler factor

$$L_v(\pi_v, s) = \frac{\Gamma(s)}{(2\pi)^s} \text{ and } L_\infty(\pi_\infty, s) = \prod_{v \in S_\infty} L_v(\pi_v, s) \tag{8.167}$$

We define the completed  $L$ -function

$$\Lambda(\pi, s) = L_\infty(\pi_\infty, s) L(\pi_f, s) \tag{8.168}$$

Now we see from the definition of the generators  $\omega_v^\dagger$  we get

$$\int_{T_1(F_v)} \langle \omega_v^\dagger, e_{\mu_v} \rangle d^* t_v = \frac{\Gamma(1 + w(\mu))}{(2\pi)^{1+w(\mu)}} \tag{8.169}$$

and hence we get the final formula

$$\begin{aligned}
 & \langle j((x, \underline{g}_{\mathfrak{p}}), r_{\lambda, \mu})(\mathcal{F}^\bullet(\frac{[\omega_\epsilon]}{\Omega(\pi_f, \epsilon)} \times h_{\pi_f}^{(0)})) = \\
 & \frac{1}{\Omega(\pi_f, \epsilon)} \Lambda(\pi \otimes \tilde{\mu}_{\mathfrak{p}}, 1) I^{\text{loc}}(h_{\pi_f}^{(0)})(g_{\mathfrak{p}})
 \end{aligned} \tag{8.170}$$

This is now exactly the expression we want to see. We have identified the factor  $\mathcal{L}(\pi \otimes \chi, \mu)$  in (8.89),(8.90) a special values of an  $L$ -function. We have to interpret this formula.

It is clear from our computations above that for all places  $\mathfrak{p}$  where  $\chi_{\mathfrak{p}}, \tilde{\mu}_{\mathfrak{p}}$  are unramified and where  $t_{0, \mathfrak{p}} \in T_1(\mathcal{O}_{F, \mathfrak{p}})$  we have  $I_{\mathfrak{p}}^{\text{loc}}(h_{\pi_{\mathfrak{p}}}^{(0)})(t_{0, \mathfrak{p}}) = 1$ . Hence it

is clear that in the infinite product  $I^{\text{loc}}(h_{\pi_f^{(0)}})([t_{0,f}]) = \prod_{\mathfrak{p}} I_{\mathfrak{p}}^{\text{loc}}(h_{\pi_{\mathfrak{p}}}^{(0)}(t_{0,\mathfrak{p}}))$  almost all factors are equal to one, hence there is no issue with convergence.

It is also clear that  $I_{\mathfrak{p}}^{\text{loc}}(h_{\pi_{\mathfrak{p}}}([t_{0,\mathfrak{p}}])) \in F(\tilde{\mu}_f)$ , this was the field extension of  $F$  which was generated by the values of  $\tilde{\mu}_f$ . We may evaluate the local intertwining operator at any function  $h_{\pi_f} = \prod' h_{\pi_{\mathfrak{p}}}$  where  $h_{\pi_{\mathfrak{p}}} = h_{\pi_{\mathfrak{p}}}^{(0)}$  for almost all  $\mathfrak{p}$  we can find an element  $h_{\pi_f}$  such that  $I_{\mathfrak{p}}^{\text{loc}}(h_{\pi_{\mathfrak{p}}}([t_{0,\mathfrak{p}}])) \neq 0, ?$  for all  $\mathfrak{p}$ . We apply our formula above to this function then we can conclude

**Corollary 8.2.1.** *The number  $\frac{1}{\Omega(\pi_f, \epsilon)} \Lambda(\pi \otimes \tilde{\mu}_{\mathfrak{p}}, 1) \in F[\tilde{\mu}_{\mathfrak{p}}^{-1}]$ , For any element  $\sigma \in \text{Gal}(F(\tilde{\mu}_f))/\mathbb{Q}$  we have*

$$\left(\frac{1}{\Omega(\pi_f, \epsilon^\sigma)} \Lambda(\pi^\sigma \otimes \tilde{\mu}_{\mathfrak{p}}, 1)\right)^\sigma = \frac{1}{\Omega(\pi_f, \epsilon)} \Lambda(\pi \otimes (\tilde{\mu}_{\mathfrak{p}})^\sigma, 1)$$

Of course we also get an integrality statement. We are still working with our level subgroup  $K_f = \prod_{\mathfrak{p}} K_{\mathfrak{p},1}(\mathfrak{p}^{f(\pi_{\mathfrak{p}})})$ , but we choose an arbitrary pin point  $(x, \underline{g}_f)$  and a level subgroup  $K_f^T \in T(\mathbb{A}_f)$  such that  $K_f^T \underline{g}_f K_f = \underline{g}_f K_f$  -this is our condition b) above. We get the maps

$$j(x, \underline{t}_f) : \mathcal{S}_{K_f^T}^T \rightarrow \mathcal{S}_{K_f}^G, j(x, \underline{t}_f)^\bullet : H_c^{r_1+r_2}(\mathcal{S}_{K_f}^G, \mathcal{M}^b \otimes \mathcal{O}_F) \rightarrow H_c^{r_1+r_2}(\mathcal{S}_{K_f^T}^T, \mathcal{O}_F \otimes \tilde{\mu}_f),$$

it follows from the definition of the periods that the cohomology class  $\mathcal{F}^\bullet\left(\frac{[\omega_\epsilon]}{\Omega(\pi_f, \epsilon)} \times h_{\pi_f}^{(0)}\right) \in H_c^{r_1+r_2}(\mathcal{S}_{K_f}^G, \mathcal{M}^b \otimes \mathcal{O}_F)(\epsilon \times \pi_f)_{!, \text{int}}$ . Hence we can lift this class to a class  $\mathcal{F}^\bullet\left(\frac{[\omega_\epsilon]}{\Omega(\pi_f, \epsilon)} \times h_{\pi_f}^{(0)}\right)^*$  in  $H_c^{r_1+r_2}(\mathcal{S}_{K_f}^G, \mathcal{M}^b \otimes \mathcal{O}_F)$  and the restriction

$$j(x, \underline{t}_f)^\bullet \left(\mathcal{F}^\bullet\left(\frac{[\omega_\epsilon]}{\Omega(\pi_f, \epsilon)} \times h_{\pi_f}^{(0)}\right)^*\right) \in H_c^{r_1+r_2}(\mathcal{S}_{K_f^T}^T, \mathcal{O}_F \otimes \tilde{\mu}_f)$$

gives us the number

$$\langle j(x, \underline{t}_f)^\bullet \left(\mathcal{F}^\bullet\left(\frac{[\omega_\epsilon]}{\Omega(\pi_f, \epsilon)} \times h_{\pi_f}^{(0)}\right)^*\right), e_{\tilde{\mu}_f} \rangle$$

which is the number in the top line in (8.165). But this number may depend on the lift. We still have proposition 8.2.1 which says that this number does not depend on the lift if  $j \circ \partial_{\chi_{\tilde{\mu}}} = 0$ . and hence we can say

**Theorem 8.2.1.** *If  $j \circ \partial_{\tilde{\mu}_f} = 0$ . then*

$$\langle j(x, \underline{g}_f)^\bullet \left(\mathcal{F}^\bullet\left(\frac{[\omega_\epsilon]}{\Omega(\pi_f, \epsilon)} \times h_{\pi_f}^{(0)}\right), \tilde{\mu}_f \right) \rangle := \langle j(x, \underline{g}_f)^\bullet \left(\mathcal{F}^\bullet\left(\frac{[\omega_\epsilon]}{\Omega(\pi_f, \epsilon)} \times h_{\pi_f}^{(0)}\right)^*, e_{\tilde{\mu}_f} \right) \rangle =$$

$$\int_{T_1(\mathbb{Q}) \backslash T_1(\mathbb{A})} \langle \mathcal{F}\left(\frac{\omega_\epsilon}{\Omega(\underline{\epsilon} \times \pi_f)} \times h_{\pi_f}^{(0)}\right)(\underline{g}_f), e_{\tilde{\mu}_f} \rangle d^* \underline{t} =$$

$$\frac{1}{\Omega(\pi_f, \epsilon)} \Lambda(\pi \otimes \tilde{\mu}_{\mathfrak{p}}, 1) I^{\text{loc}}(h_{\pi_f}^{(0)})(\underline{g}_f)$$

where  $d^* \underline{t}_f$  has volume 1 on  $K_f^T$ . As far as  $j \circ \partial_{\chi_{\tilde{\mu}}} = 0$  is concerned we have

**Proposition 8.2.5.** *The class  $j \circ \partial_{\chi_{\tilde{\mu}}} = 0$  unless  $\underline{\lambda}$  is of parallel weight, i.e. all the  $n_\lambda$  are equal to the same number  $m$ .*

*If  $\underline{\lambda}$  is of parallel weight we still have  $j \circ \partial_{\chi_{\tilde{\mu}}} = 0$  unless  $\underline{\mu} = \underline{\lambda}, w_{0\underline{\lambda}}$ .*

*Proof.* Postponed □

If we are in the exceptional case that  $j \circ \partial_{\chi_{\tilde{\mu}}} \neq 0$  we use the Manin-Drinfeld argument. We can find Hecke operators  $C'$  in ( the central subalgebra ) of the Hecke algebra which annihilate  $j \circ \partial_{\chi_{\tilde{\mu}}} \neq 0$  and act by multiplication by a non zero algebraic integer  $\pi_f(T_h) \in \mathcal{O}_F$  on  $H_c^{r_1+r_2}(\mathcal{S}_{K_f}^G, \mathcal{M}^b \otimes \mathcal{O}_F)(\pi_f)$  (See [?]). We consider the ideal generated by all these numbers  $\pi_f(T_h) \in \mathcal{O}_F$  and we assume for simplicity that it is a principal ideal  $\Delta(\pi_f, \mu)$ . Then we can apply proposition 8.2.1 and get

$$\begin{aligned} < (j(x, \underline{g}_f) \bullet (\mathcal{F} \bullet (\frac{[\omega_\epsilon]}{\Omega(\pi_f, \epsilon)} \times h_{\pi_f}^{(0)}), T_h \chi^{[\mu, 1]} > := < (j(x, \underline{g}_f) \bullet (\mathcal{F} \bullet (\frac{[\omega_\epsilon]}{\Omega(\pi_f, \epsilon)} \times h_{\pi_f}^{(0)})^*, T_h \chi^{[\mu, 1]} > = \\ & \int_{T_1(\mathbb{Q}) \backslash T_1(\mathbb{A})} < \mathcal{F}(\frac{\omega_\epsilon}{\Omega(\epsilon \times \pi_f)} \times \pi_f(T_h) h_{\pi_f}^{(0)})(\underline{g}_f), \tilde{\mu}_p > d^x t = \\ & \frac{\pi_f(T_h)}{\Omega(\pi_f, \epsilon)} \Lambda(\pi \otimes \tilde{\mu}_p, 1) I^{\text{loc}}(h_{\pi_f}^{(0)})(\underline{g}_f) \end{aligned} \tag{8.171}$$

This formula is a supplement to the theorem above if the proposition 8.2.1 does not apply directly. We have used this argument already before in section 5.6.

**Corollary 8.2.2.** *If  $j \circ \partial_{\chi_{\tilde{\mu}}} = 0$  then*

$$\frac{\Lambda(\pi \otimes \tilde{\mu}_p, 1)}{\Omega(\pi_f, \epsilon)} I^{\text{loc}}(h_{\pi_f}^{(0)})(\underline{g}_f) \in \mathcal{O}_{F[\tilde{\mu}_p]}$$

and if this is not the case then we get the weaker result

$$\Delta(\pi_f, \mu) \frac{\Lambda(\pi \otimes \tilde{\mu}_p, 1)}{\Omega(\pi_f, \epsilon)} I^{\text{loc}}(h_{\pi_f}^{(0)})(\underline{g}_f) \in \mathcal{O}_{F[\tilde{\mu}_p]}$$

The numbers  $\frac{\Lambda(\pi \otimes \tilde{\mu}_p, 1)}{\Omega(\pi_f, \epsilon)}$  are of arithmetic interest, for instance the factorisation into primes contains information about the structure of the cohomology (see further down). For instance we can ask whether they are integers themselves, or if not what are the denominators. This amounts to the study of the numbers  $\Delta(\pi_f, \mu)$  and  $I^{\text{loc}}(h_{\pi_f}^{(0)})(\underline{g}_f) = \prod_p I^{\text{loc}}(h_{\pi_p}^{(0)})(\underline{g}_p)$ .

The number  $\Delta(\pi_f, \mu)$  is of global nature, it should be a denominator of the Eisenstein class. We determined this denominator in a very special case in Chapter V Theorem 5.1.1, in this case  $\mathcal{G}/\mathbb{Z} = \text{GL}_2/\text{Spec}(\mathbb{Z})$  and the level was  $K_f = \mathcal{G}(\hat{\mathbb{Z}})$ . It is certainly not too difficult to extend this result to the case of congruence subgroups.

I believe that it is very interesting problem to study the numbers  $\Delta(\pi_f, \mu)$  if  $F_0/\mathbb{Q}$  is a non trivial extension of  $\mathbb{Q}$ , for instance simply a real quadratic extension.

We look at the numbers  $I^{\text{loc}}(h_{\pi_f}^{(0)})(\underline{g}_f) = \prod_p I_p^{\text{loc}}(h_{\pi_p}^{(0)})(\underline{g}_p)$ . They are algebraic integers, of course our goal must be to arrange the data such that the

number  $I^{\text{loc}}(h_{\pi_{\mathfrak{p}}}^{(0)})(g_{\mathfrak{p}}) \neq 0$  and to keep the ( the number of) prime divisors "small" . Given our pin point  $g_{\mathfrak{p}}$  we compute

$$\begin{aligned} & [T(\mathcal{O}_{F_0, \mathfrak{p}}) : K_{\mathfrak{p}}^T] \int_{T_1(F_0, \mathfrak{p})} h_{\pi_{\mathfrak{p}}}^{(0)}([t_{\mathfrak{p}}]g_{\mathfrak{p}})\tilde{\mu}_{1, \mathfrak{p}}([t_{\mathfrak{p}}])d_{\text{Tam}}t_{\mathfrak{p}} = \\ & [T(\mathcal{O}_{F_0, \mathfrak{p}}) : K_{\mathfrak{p}}^T] \sum_{n \in \mathbb{Z}} \tilde{\mu}_{1, \mathfrak{p}}(\varpi_{\mathfrak{p}})^n \int_{\mathcal{O}_{F_0, \mathfrak{p}}^{\times}} h_{\pi_{\mathfrak{p}}}^{(0)}\left(\begin{pmatrix} \varpi_{\mathfrak{p}}^n \varepsilon & 0 \\ 0 & 1 \end{pmatrix} g_{\mathfrak{p}}\right)\tilde{\mu}_{1, \mathfrak{p}}(\varepsilon)d_{\text{Tam}}\varepsilon \end{aligned} \quad (8.172)$$

We have the freedom to choose  $g_{\mathfrak{p}}$ , of course we always want condition b).

We start with the choice  $g_{\mathfrak{p}} = [t_0] = \begin{pmatrix} \varpi_{\mathfrak{p}}^{m_0} & 0 \\ 0 & 1 \end{pmatrix}$ , since  $h_{\pi_{\mathfrak{p}}}^{(0)}$  is the new vector we know  $h_{\pi_{\mathfrak{p}}}^{(0)}\left(\begin{pmatrix} \varpi_{\mathfrak{p}}^{n+m_0} \varepsilon & 0 \\ 0 & 1 \end{pmatrix}\right) = \zeta_{\mathfrak{p}}(\varepsilon)h_{\pi_{\mathfrak{p}}}^{(0)}\left(\begin{pmatrix} \varpi_{\mathfrak{p}}^{n+m_0} & 0 \\ 0 & 1 \end{pmatrix}\right)$  and hence the right hand side in (8.172) becomes

$$\sum_{n \in \mathbb{Z}} \tilde{\mu}_{1, \mathfrak{p}}(\varpi_{\mathfrak{p}})^n h_{\pi_{\mathfrak{p}}}^{(0)}\left(\begin{pmatrix} \varpi_{\mathfrak{p}}^{n+m_0} & 0 \\ 0 & 1 \end{pmatrix}\right) \int_{\mathcal{O}_{F_0, \mathfrak{p}}^{\times}} \zeta_{\mathfrak{p}}(\varepsilon)\tilde{\mu}_{1, \mathfrak{p}}(\varepsilon)d^{\times}\varepsilon \quad (8.173)$$

This is of course only useful if  $\zeta_{\mathfrak{p}}\tilde{\mu}_{1, \mathfrak{p}}$  is unramified, i.e.  $\zeta_{\mathfrak{p}}\tilde{\mu}_{1, \mathfrak{p}}(\varepsilon) = 1$ . We assume that this is the case, then our pin point  $g_{\mathfrak{p}}$  does not put any constraint on  $K_{\mathfrak{p}}^T$ . Hence we may assume that  $K_{\mathfrak{p}}^T \cap T_1(F_{0, \mathfrak{p}}^{\times}) = \mathcal{O}_{F_0, \mathfrak{p}}^{\times}$  and the integral is simply equal to 1.

Again we have to discuss different cases. We have seen that  $h_{\pi_{\mathfrak{p}}}^{(0)}\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right)$  is supported on an annulus if  $\pi_{\mathfrak{p}}$  is a principal series representation and  $\chi_{1, \mathfrak{p}}$  is ramified, or if  $\pi_{\mathfrak{p}}$  is a discrete series representation. Hence we can choose  $m_0$  so that in the summation the only surviving term is the term  $n = 0$  and hence under these conditions we can achieve

$$I_{\mathfrak{p}}^{\text{loc}}(h_{\pi_{\mathfrak{p}}}^{(0)})(g_{\mathfrak{p}}) = 1.$$

If the function  $h_{\pi_{\mathfrak{p}}}^{(0)}\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}\right)$  is not supported on an annulus, then our computations above show that we can find a  $m_0$  such that

$$I_{\mathfrak{p}}^{\text{loc}}(h_{\pi_{\mathfrak{p}}}^{(0)})\left(\begin{pmatrix} \varpi_{\mathfrak{p}}^{m_0} & 0 \\ 0 & 1 \end{pmatrix}\right) = 1.$$

We come to the case where  $\zeta_{\mathfrak{p}}\tilde{\mu}_{1, \mathfrak{p}}$  is ramified, we have to choose a different pin point. We take

$$g_{\mathfrak{p}} = \begin{pmatrix} \varpi_{\mathfrak{p}}^{m_0} & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & \varpi_{\mathfrak{p}}^{-\nu_0} \\ 0 & 1 \end{pmatrix}$$

where  $\nu_0 > 0$ . This imposes some restriction on the choice of our level  $K_{\mathfrak{p}}^T$ , if we want condition b) ( for  $K_{\mathfrak{p}} = K_{\mathfrak{p}, 1}(\mathfrak{p}^{n_0})$ ) we must have:

$$\begin{pmatrix} t_1 & 0 \\ 0 & t_2 \end{pmatrix} \in K_{\mathfrak{p}}^T \implies \frac{t_1}{t_2} \equiv 1 \pmod{\mathfrak{p}^{\nu_0}}$$

We compute the value

$$h_{\pi_{\mathfrak{p}}}^{(0)}\left(\begin{pmatrix} \varpi_{\mathfrak{p}}^{n+m_0} \varepsilon & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & \varpi_{\mathfrak{p}}^{-\nu_0} \\ 0 & 1 \end{pmatrix}\right) = h_{\pi_{\mathfrak{p}}}^{(0)}\left(\begin{pmatrix} 1 & \varepsilon \varpi_{\mathfrak{p}}^{-\nu_0+n+m_0} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \varpi_{\mathfrak{p}}^{n+m_0} \varepsilon & 0 \\ 0 & 1 \end{pmatrix}\right)$$

$$h_{\pi_{\mathfrak{p}}}^{(0)}\left(\begin{pmatrix} \varpi_{\mathfrak{p}}^{n+m_0} & 0 \\ 0 & 1 \end{pmatrix}\right) \zeta(\varepsilon) \psi_{\mathfrak{p}}(\varepsilon \varpi_{\mathfrak{p}}^{-\nu_0+n+m_0})$$

and hence our expression in (8.172) becomes

$$[T_1(\mathcal{O}_{F_0, \mathfrak{p}}) : K_{\mathfrak{p}}^T] \sum_{n \in \mathbb{Z}} h_{\pi_{\mathfrak{p}}}^{(0)}\left(\begin{pmatrix} \varpi_{\mathfrak{p}}^n & 0 \\ 0 & 1 \end{pmatrix}\right) \int_{\mathcal{O}_{F_0, \mathfrak{p}}^\times} \psi_{\mathfrak{p}}(\varpi_{\mathfrak{p}}^{-\nu_0+n} \varepsilon) \zeta_{\mathfrak{p}}(\varepsilon) \tilde{\mu}_{\mathfrak{p}}(\varepsilon) d^\times \varepsilon$$

The integral is a Gaussian sum, the conductor  $f(\zeta_{\mathfrak{p}} \tilde{\mu}_{1, \mathfrak{p}}) > 0$  and we know that the Gaussian sum is zero unless the additive character  $u \mapsto \psi_{\mathfrak{p}}(\varpi_{\mathfrak{p}}^{-\nu_0+n+m_0} u)$  restricted to  $\mathcal{O}_{F_0, \mathfrak{p}}$  has conductor  $f(\zeta_{\mathfrak{p}} \tilde{\mu}_{1, \mathfrak{p}})$ . Hence the Gaussian sum is only non zero if

$$-\nu_0 + n + f(\zeta_{\mathfrak{p}} \tilde{\mu}_{1, \mathfrak{p}}) = -d_{\mathfrak{p}}.$$

We have normalised  $h_{\pi_{\mathfrak{p}}}^{(0)}\left(\begin{pmatrix} \varpi_{\mathfrak{p}}^{-d_{\mathfrak{p}}} & 0 \\ 0 & 1 \end{pmatrix}\right) = 1$  hence we have to choose  $\nu_0 = f(\zeta_{\mathfrak{p}} \tilde{\mu}_{1, \mathfrak{p}})$  and we find

$$I_{\mathfrak{p}}^{\text{loc}}(h_{\pi_{\mathfrak{p}}}^{(0)})(\underline{g}_{\mathfrak{p}}) = G(\zeta_{\mathfrak{p}} \tilde{\mu}_{1, \mathfrak{p}}, \psi_{\mathfrak{p}}) \tag{8.174}$$

The Gaussian sum only depends on the restriction of  $\zeta_{\mathfrak{p}} \tilde{\mu}_{1, \mathfrak{p}}$  to the units  $T(\mathcal{O}_{F_0, \mathfrak{p}})$ . The values of  $\mathfrak{p} \tilde{\mu}_{1, \mathfrak{p}}$  on the units are  $(N(\mathfrak{p}) - 1) \times N(\mathfrak{p})^{f(\zeta_{\mathfrak{p}} \tilde{\mu}_{1, \mathfrak{p}}) - 1}$  th roots of unity, the values of  $\psi_{\mathfrak{p}}$  are also  $N(\mathfrak{p})^{f(\zeta_{\mathfrak{p}} \tilde{\mu}_{1, \mathfrak{p}}) - 1}$  th roots of unity. Hence the number  $G(\zeta_{\mathfrak{p}} \tilde{\mu}_{1, \mathfrak{p}}, \psi_{\mathfrak{p}})$  is an algebraic integer, it lies in the field  $\mathbb{Q}[\zeta_{N(\mathfrak{p})-1}, \zeta_{N(\mathfrak{p})^{f(\zeta_{\mathfrak{p}} \tilde{\mu}_{1, \mathfrak{p}})}}$ . In factorisation of this integer into prime ideals only primes lying above  $p$  occur. Hence we have control over the numbers  $I_{\mathfrak{p}}^{\text{loc}}(h_{\pi_{\mathfrak{p}}}^{(0)})(\underline{g}_{\mathfrak{p}})$ .

### 8.2.5 Poincare duality and modular symbols.

We consider the second example where can apply the strategy which we outlined in section 8.2.1. We start from an arbitrary quasi split group  $H/\mathbb{Q}$ . Let  $T/\mathbb{Q} \subset B/\mathbb{Q}$  a maximal torus in a Borel sub group  $B/\mathbb{Q}$ . Let us denote the center of  $H/\mathbb{Q}$  by  $C_H/\mathbb{Q} \subset T/\mathbb{Q}$ , for any  $\lambda \in X^*(T \times_{\mathbb{Q}} \bar{\mathbb{Q}})$  the restriction of  $\lambda$  to  $C_H$  is denoted by  $\lambda_{C_H}$ .

We take for our ambient group  $G/\mathbb{Q} = H \times H/\mathbb{Q}$  and we embed  $H/\mathbb{Q} \hookrightarrow G/\mathbb{Q}$  diagonally. We follow the steps in 8.2.1. We choose a level subgroup  $K_f^H \subset H(\mathbb{A}_f)$  and put  $K_f^H \times K_f^H = K_f^G$ . Then we have  $\mathcal{S}_{K_f^H}^H \times \mathcal{S}_{K_f^H}^H = \mathcal{S}_{K_f^G}^G$ . We choose the base point  $e_0 \in H(\mathbb{R})/K_{\infty}^H = X$  and  $x_0 = (e_0, e_0)$ . From this we get the map

$$j(x_0, e_f) : \mathcal{S}_{K_f^H}^H = H(\mathbb{Q}) \backslash H(\mathbb{R})/K_{\infty}^H \times H(\mathbb{A}_f)/K_f^H \rightarrow \mathcal{S}_{K_f^G}^G. \tag{8.175}$$

An irreducible representation of  $G/\mathbb{Q}$  is of the form  $\mathcal{M}_{\lambda} = \mathcal{M}_{\lambda_1} \otimes \mathcal{M}_{\lambda_2}$ , where  $\lambda_1, \lambda_2$  are dominant weights. We choose a one dimensional representation  $\mu : H/\mathbb{Z} \rightarrow \mathcal{O}^{\times}$ . We have to understand the module of  $H$ -homomorphisms  $\text{Hom}_H(\mathcal{M}_{\lambda}, \mathcal{O}^{\times})$ . We know

**Proposition 8.2.6.** *The module  $\text{Hom}_H(\mathcal{M}_\lambda, \mathcal{O}\mu)$  is free of rank  $d_{\lambda, \mu}$ . We have*

$$d_{\lambda, \mu} = \begin{cases} 1 & \text{if } \lambda_1^{(1)} = -w_0(\lambda_1^{(2)}) \text{ and } \lambda_{1, C_H}^{(1)} + \lambda_{2, C_H}^{(1)} = \mu_{C_H} \\ 0 & \text{else} \end{cases}$$

*Proof.* This is obvious from the theory of representations of algebraic groups.  $\square$

We assume now that  $d_{\lambda, \mu} = 1$  and choose a generator  $r_{\lambda, \mu} \in \text{Hom}_H(\mathcal{M}_\lambda, \mathcal{O}\mu)$ . We get a homomorphism (See 8.82)

$$j(x, e_f, r_{\lambda, \mu})^{d_H} : H_c^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \rightarrow H^{d_H}(\mathcal{S}_{K_f}^H, \mathcal{O}\mu).$$

We know that the Manin-Drinfeld principle is valid, this means that we get a canonical splitting for the Hecke modules

$$H_c^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F) = \text{Im}(H^{d_H-1}(\partial(\mathcal{S}_{K_f}^G), \tilde{\mathcal{M}}_\lambda \otimes F)) \oplus H_!^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F)$$

here  $F$  is the quotient field of  $\mathcal{O}$ . Hence we can restrict our homomorphism

$$j(x, e_f, r_{\lambda, \mu})^{d_H} : H_!^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F) \rightarrow H^{d_H}(\mathcal{S}_{K_f}^H, F\mu).$$

We want to discuss the integral cohomology. We start from the exact sequence and get a diagram

$$\begin{array}{ccccccc} H^{d_H-1}(\partial(\mathcal{S}_{K_f}^G), \tilde{\mathcal{M}}_\lambda) & \xrightarrow{\delta} & H_c^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) & \rightarrow & H_!^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) & \rightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow & & \\ H^{d_H-1}(\partial(\mathcal{S}_{K_f}^G), \tilde{\mathcal{M}}_\lambda)_{\text{int}}^{\text{sat}} & \xrightarrow{\delta} & H_c^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)_{\text{int}} & \rightarrow & H_!^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)_{\text{int}} & \rightarrow & 0 \end{array} \quad (8.176)$$

and the Manin-Drinfeld principle gives us a splitting up to isogeny

$$H_c^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)_{\text{int}} \supset H^{d_H-1}(\partial(\mathcal{S}_{K_f}^G), \tilde{\mathcal{M}}_\lambda)_{\text{int}}^{\text{sat}} \oplus H_!^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)_{\text{int}} \quad (8.177)$$

the reader should pay attention to the difference between the subscripts  $_{\text{int}}$  and  $_{\text{int}}$  we have an inclusion  $H_!^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)_{\text{int}} \subset H_!^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)_{\text{int}}$  the quotient is a finite module, which may be difficult to understand. There is a non zero number  $\Delta_\lambda \in \mathcal{O}$  such that  $\Delta_\lambda H_!^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)_{\text{int}} \subset H_!^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)_{\text{int}}$ . Then it is clear that  $j(x, e_f, r_{\lambda, \mu})^{d_H}$  induces Hecke invariant homomorphisms

$$\begin{aligned} j(x, e_f, r_{\lambda, \mu})^{d_H} : H_!^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)_{\text{int}} &\rightarrow H_!^{d_H}(\mathcal{S}_{K_f}^G, \mathcal{O}\mu) \\ j(x, e_f, r_{\lambda, \mu})^{d_H} : H_!^{d_H}(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)_{\text{int}} &\rightarrow \frac{1}{\Delta_\lambda} H_!^{d_H}(\mathcal{S}_{K_f}^G, \mathcal{O}\mu) \end{aligned} \quad (8.178)$$

Assume that the argument, which I have in my mind, is correct, then we may even apply proposition 8.2.1 and then we can take  $\Delta_\lambda = 1$ .

We can produce classes in  $H_!^{d_H}((\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda))$ . For any  $0 \leq r \leq d_H$  we have the Künneth homomorphism

$$H_!^r(\mathcal{S}_{K_f}^H, \tilde{\mathcal{M}}_{\lambda_1}) \times H_!^{d_H-r}(\mathcal{S}_{K_f}^H, \tilde{\mathcal{M}}_{\lambda_2}) \rightarrow H_!^{d_H}((\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda))$$

and taking the composition with  $j(x, e_f, r_{\lambda, \mu})_!^{d_H}$  we get

$$J(r) : H_!^r(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1}) \times H_!^{d_H-r}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_2}) \rightarrow \frac{1}{\Delta_\lambda} H_!^{d_H}(\mathcal{S}_{K_f^H}^H, \mathcal{O}_\mu) \quad (8.179)$$

It is clear from the definitions that this homomorphism is the cup product.

If necessary we enlarge our field such that we get decompositions ( up to isogeny) into absolutely irreducible Hecke modules

$$\begin{aligned} H_!^r(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1}) &\supset \bigoplus_{\pi_{1,f}} H_!^r(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1})[\pi_{1,f}] \\ H_!^{d_H-r}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_2}) &\supset \bigoplus_{\pi_{2,f}} H_!^{d_H-r}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_2})[\pi_{2,f}] \\ H_!^{d_H}(\mathcal{S}_{K_f^H}^H, \mathcal{O}_\mu) &\supset \bigoplus_{\tilde{\mu}: \text{type}(\tilde{\mu})=\mu} \mathcal{O}_{\tilde{\mu}}. \end{aligned}$$

Hence we have to compute the pairing

$$H_!^r(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1})[\pi_{1,f}] \times H_!^{d_H-r}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_2})[\pi_{2,f}] \rightarrow \frac{1}{\Delta_\lambda} \mathcal{O}_{\tilde{\mu}} \quad (8.180)$$

This pairing is zero unless  $\pi_{1,f}, \pi_{2,f}$  are essentially dual, i.e. dual up to a twist. scussion in ??)

But we still have to go one step further, we to take into account the action of  $\pi_0(G(\mathbb{R}))$  on the different cohomology groups, our pairing becomes

$$H^r(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1})[\epsilon_1 \times \pi_{1,f}] \times H^{d_H-r}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_2})[\epsilon_2 \times \pi_{2,f}] \rightarrow \frac{1}{\Delta_\lambda} \mathcal{O}_{\tilde{\mu}} \quad (8.181)$$

and the consistency rule  $\epsilon_1 \epsilon_2 = \tilde{\mu}_\infty$  should be satisfied.

Here we described a very general situation, it seems to be a very difficult problem to compute this pairing, at the end of section 8.2.2 we formulated the expectation that the value of this pairing should be expressible in terms of  $L$ -functions. attached to  $\pi_{1,f}$ , I have no idea how to do this in general.

### A special example

We stop our general reasoning and consider a very special case, we choose a finite extension  $F/\mathbb{Q}$  and choose  $H/\mathbb{Q} = R_{F/\mathbb{Q}}(\text{Gl}_2/F)$  and  $G/\mathbb{Q} = R_{F/\mathbb{Q}}((\text{Gl}_2 \times \text{Gl}_2)/F)$ . In this situation we can work with the Whittaker model. In this case  $d_H = 2r_1 + 3r_2$ , we pick two isomorphism types  $\pi_{1,f}, \pi_{2,f}$  which occur in the cuspidal cohomology  $H^\bullet(\mathcal{S}_{K_f^G}^G, \tilde{\mathcal{M}}_{\lambda_1}), H^\bullet(\mathcal{S}_{K_f^G}^G, \tilde{\mathcal{M}}_{\lambda_2})$ .

We want to compute the value of the pairing

$$H^{r_1+r_2}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1})[\epsilon_2 \times \pi_{2,f}] \times H^{r_1+2r_2}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1})[\epsilon_2 \times \pi_{2,f}] \rightarrow \frac{1}{\Delta(\pi_{1,f}, \pi_{2,f})} \mathcal{O}_{\tilde{\mu}} \quad (8.182)$$

here  $\epsilon_1, \epsilon_2$  and  $\eta$  are characters on  $(\mathbb{Z}/2\mathbb{Z})^{r_1} = \pi_0(H(\mathbb{R}))$ . Of course this pairing is zero unless we have  $\pi_{2,f}^\vee = \pi_{1,f}$  and  $\epsilon_1 \epsilon_2 \eta = 1$ .

We start from the Whittaker model  $\tilde{\mathcal{D}}_\lambda \otimes \mathcal{W}(\pi_f, \psi_f)$  and we choose generators  $\omega_\epsilon^{(\dagger, \nu)} \in \text{Hom}_{K_\infty}(\Lambda^{r_1+\nu r_2}(\mathfrak{g}/\mathfrak{k}),$

$\tilde{t}ilde{\mathcal{D}}_\lambda^\epsilon, \psi_\infty) \otimes \mathcal{M}_\lambda)$  and  $h_f^{(i)} = \prod_{\mathfrak{p}} h_{\mathfrak{p}}^{(i)} \in \prod_{\mathfrak{p}} \mathcal{W}(\pi_{i,\mathfrak{p}}, \psi_{\mathfrak{p}})$ , where we even choose  $h_{\mathfrak{p}}^{(i)} = h_{\pi_{i,\mathfrak{p}}}^{(0)}$  our previously chosen generators. Then the cup product of the two integral(!) cohomology classes  $\boxed{\text{cupff}}$

$$\left[ \frac{1}{\Omega_\epsilon^{(1)}(\pi_f)} \mathcal{F}_1^{(1)}(\omega_\epsilon^{(\dagger,1)} \times h_f^{(1)}) \right] \cup \left[ \frac{1}{\Omega_\epsilon^{(2)}(\pi_f)} \mathcal{F}_1^{(2)}(\omega_\epsilon^{(\dagger,2)} \times h_f^{(2)}) \right] \quad (8.183)$$

is given by the integral (see section 6.3.11)

$$\frac{1}{\Omega_\epsilon^{(1)}(\pi_f) \Omega_\epsilon^{(2)}(\pi_f)} \int_{S_{\kappa^H}^H} \mathcal{F}_1^{(1)}(\omega_\epsilon^{(\dagger,1)} \times h_f^{(1)}) \wedge \mathcal{F}_1^{(1)}(\omega_\epsilon^{(\dagger,2)} \times h_f^{(2)}),$$

by construction the expression under the integral is a differential form in top degree.

We choose a specific invariant volume form  $dy = dy_\infty \times dy_f$  on  $H(\mathbb{A})$ . We normalize  $\text{vol}_{dy_f}(K_f^H) = 1$ , and we write  $dy_\infty = dx_\infty \times dk_\infty$ , we require  $\text{vol}_{dk_\infty}(K_\infty) = 1$  and  $dx_\infty$  will be the volume form given by the Riemannian metric. To write down the integral explicitly we choose an orthonormal basis of  $\mathfrak{p} \otimes \mathbb{R}$ . This basis will consist of bases of the  $\mathfrak{p}_v$ . For the  $\mathfrak{p}_v$  we choose the following basis

a) If  $v$  is real our basis will be  $X_{v,+} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, X_{v,-} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$

b) For  $v$  complex we have the basis  $X_{v,0}, X_{v,1}, X_{v,-1}$

Hence we get a basis

$$\{\dots, X_{v,+}, X_{v,-}, \dots\}_{S_\infty^{\text{real}}} \cup \{\dots, X_{v,0}, X_{v,1}, X_{v,-1}, \dots\}_{S_\infty^{\text{comp}}} \quad (8.184)$$

To evaluate the integral we have to look at the value

$$\omega_\epsilon^{(\dagger,1)} \wedge \omega_{\epsilon'}^{(\dagger,2)} (\Lambda^{2r_1+3r_2} X_\nu) \quad (8.185)$$

where the  $X_\nu$  run through the above basis. The result is an element in  $\mathcal{W}(\mathcal{D}_\lambda) \otimes \mathcal{M}_\lambda$ . To compute this value we have to divide the above basis into a subset  $A = \{\dots, X_\nu, \dots\}$  consisting of  $r_1 + r_2$  elements and a complement  $B = \{\dots, X_\mu^*, \dots\}$  consisting of  $r_1 + 2r_2$  elements of our basis, we have to multiply  $\omega_\epsilon^{(\dagger,1)}(\Lambda X_\nu) \omega_{\epsilon'}^{(\dagger,2)}(\Lambda X_\mu)$ . Then we have to sum over all these divisions into two subsets.

But looking at the definition of our  $\omega_\epsilon^{(\dagger,1)}, \omega_{\epsilon'}^{(\dagger,2)}$  we see that only one division into two disjoint sets can give a non zero contribution. We describe this division. Recall that  $\epsilon = \{\dots, \epsilon_v, \dots\}_{v \in S_\infty^{\text{real}}}$  is an array of signs, and  $\epsilon'$  is the opposite array. Then the first  $r_1$  elements in  $A$  will be the  $X_{v,\epsilon_v}$  with  $v \in S_\infty^{\text{real}}$  and this will be supplemented by the  $X_{v,0}^*$  with  $v \in S_\infty^{\text{comp}}$ , this is the set  $A_1$ . The set  $B_1$  is the complement, but we also give the explicit description. The first  $r_1$  elements will be the  $X_{v,\epsilon'_v}$  and the second part consists of the elements  $\{\dots, X_{v,1}, X_{v,-1}, \dots\}$ . Hence we see that

$$\omega_\epsilon^{(\dagger,1)}(\wedge \omega_{\epsilon'}^{(\dagger,2)} (\Lambda^{2r_1+3r_2} X_\nu)) = \omega_\epsilon^{(\dagger,1)}((\dots \wedge X_\mu \wedge \dots)_{X_\mu \in A_1}) \omega_{\epsilon'}^{(\dagger,1)}((\dots \wedge X_\mu \wedge \dots)_{X_\mu \in B_1}) \quad (8.186)$$

To evaluate the above integral we apply a method which goes back to Asai(?). We write the constant function 1 on  $H(\mathbb{Q}) \backslash H(\mathbb{A})$  as the residue of an Eisenstein series. More precisely for any complex number  $s$  we define a "height" function  $H_s(\underline{y}) = H_s(\underline{bk}) = \rho(\underline{b})^{2+s}$  where  $\underline{y} \in H(\mathbb{A}), \underline{b} \in B_H(\mathbb{A})$  and  $\underline{k} \in K_\infty \times K_f$ . This function is invariant under  $B_H(\mathbb{Q})$  and we define

$$\text{Eis}(s, \underline{y}) = \text{Res}_{s=0} \sum_{\gamma \in B_H(\mathbb{Q}) \backslash H(\mathbb{Q})} H_s(\gamma \underline{y})$$

It is well known that this series converges for  $\Re(s) \gg 0$ , hence it defines an analytic function in  $\Re(s) \gg 0$  and it has a meromorphic continuation into the entire  $s$ - plane. It is known that this function in  $s$  has a simple pole at  $s = 0$  and

$$\text{Res}_{s=0} \text{Eis}(s, \underline{y}) = \text{Res}_{s=0} \frac{\zeta_F(s+1)}{\zeta_F(s+2)}$$

especially we see that this residue considered as function in  $\underline{y}$  is a constant  $c_F$ . Therefore we compute the integral IE

$$\int_0^\infty (\text{content}) \text{Eis}(s, \underline{y}) d\underline{y} \tag{8.187}$$

and compute its residue at  $s = 0$ .

Let us denote the elements  $(\cdots \wedge X_\mu \wedge \cdots)_{X_\mu \in A_1}$  resp.  $(\cdots \wedge X_\mu \wedge \cdots)_{X_\mu \in B_1}$  by  $\mathcal{X}_{A_1}$  resp.  $\mathcal{X}_{B_1}$ , then our integral becomes

$$\begin{aligned} & \int_{H(\mathbb{Q}) \backslash H(\mathbb{A})} \mathcal{F}(\omega_\underline{\epsilon}^{(\dagger,1)} \times h_f^{(1)})(\mathcal{X}_{A_1})(y_\infty, \underline{y}_f) \mathcal{F}(\omega_\underline{\epsilon}^{(\dagger,1)} \times h_f^{(2)})(\mathcal{X}_{B_1})(y_\infty, \underline{y}_f) \text{Eis}(s, \underline{y}) d\underline{y} = \\ & \int_{H(\mathbb{Q}) \backslash H(\mathbb{A})} \mathcal{F}(\omega_\underline{\epsilon}^{(\dagger,1)} \times h_f^{(1)})(\mathcal{X}_{A_1})(y_\infty, \underline{y}_f) \mathcal{F}(\omega_\underline{\epsilon}^{(\dagger,1)} \times h_f^{(2)})(\mathcal{X}_{B_1})(y_\infty, \underline{y}_f) (\sum_{\gamma \in B_H(\mathbb{Q}) \backslash H(\mathbb{Q})} H_s(\gamma \underline{y})) d\underline{y} \\ & \int_{B_H(\mathbb{Q}) \backslash H(\mathbb{A})} \mathcal{F}(\omega_\underline{\epsilon}^{(\dagger,1)} \times h_f^{(1)})(\mathcal{X}_{A_1})(y_\infty, \underline{y}_f) \mathcal{F}(\omega_\underline{\epsilon}^{(\dagger,1)} \times h_f^{(2)})(\mathcal{X}_{B_1})(y_\infty, \underline{y}_f) H_s(\underline{y}) d\underline{y} \end{aligned}$$

We recall the decomposition  $H(\mathbb{A}) = B_H(\mathbb{A}) K_\infty^H K_f^{H,0}$  then our measure  $d\underline{y} = d\underline{b} \times (dk_\infty \times dk_f)$ . Then  $H_s(\underline{bk}_f) = H_s(\underline{b})$ , the expression under the integral is invariant under the action of  $K_\infty^H$  from the right and hence our integral becomes

$$\int_{B_H(\mathbb{Q}) \backslash B_H(\mathbb{A})} \int_{K_f^{H,0}} \mathcal{F}(\omega_\underline{\epsilon}^{(\dagger,1)} \times h_f^{(1)})(\mathcal{X}_{A_1})(y_\infty, \underline{y}_f) \mathcal{F}(\omega_\underline{\epsilon}^{(\dagger,1)} \times h_f^{(2)})(\mathcal{X}_{B_1})(y_\infty, \underline{y}_f) dk_f (H_s(\underline{b})) d\underline{b}$$

This integral converges for  $\Re(s) \gg 0$  and the value of the residue at  $s = 0$  is equal to the value of our integral. Now  $(\omega_\underline{\epsilon}^{(\dagger,1)}(\mathcal{X}_{A_1}) \times h_f^{(1)}, \omega_{\underline{\epsilon}'}^{(\dagger,2)}(\mathcal{X}_{B_1}) \times h_f^{(2)})$  are Whittaker functions (with values in  $\mathcal{M}_\lambda^b \otimes \mathbb{C}, \mathcal{M}_\lambda \otimes \mathbb{C}$  respectively) and we have the Fourier-expansions

$$\mathcal{F}(\omega_\underline{\epsilon}^{(\dagger,1)}(\mathcal{X}_{A_1}) \times h_f^{(1)}(\underline{u}\underline{t})) = \sum_{a \in F^\times} \omega_\underline{\epsilon}^{(\dagger,1)}(\mathcal{X}_{A_1}) \times h_f^{(1)} \left( \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & \underline{u} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \underline{t} & 0 \\ 0 & 1 \end{pmatrix} \right)$$

and

$$\mathcal{F}(\omega_{\underline{\epsilon}'}^{(\dagger,2)}(\mathcal{X}_{B_1}) \times h_f^{(2)})(\underline{ut}) = \sum_{a \in F^\times} \omega_{\underline{\epsilon}'}^{(\dagger,2)}(\mathcal{X}_{B_1}) \times h_f^{(2)} \left( \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & \underline{u} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \underline{t} & 0 \\ 0 & 1 \end{pmatrix} \right)$$

Since the functions  $\omega_{\underline{\epsilon}}^{(\dagger,1)}(\mathcal{X}_{A_1}) \times h_f^{(1)}$ ,  $\omega_{\underline{\epsilon}'}^{(\dagger,2)}(\mathcal{X}_{B_1}) \times h_f^{(2)}$  are Whittaker functions i.e. they satisfy

$$\omega_{\underline{\epsilon}}^{(\dagger,1)}(\mathcal{X}_{A_1}) \times h_f^{(1)} \left( \begin{pmatrix} 1 & \underline{u} \\ 0 & 1 \end{pmatrix} \underline{y} \right) = \psi(\underline{u}) \omega_{\underline{\epsilon}}^{(\dagger,1)}(\mathcal{X}_{A_1}) \times h_f^{(1)}(\underline{y})$$

$$\omega_{\underline{\epsilon}'}^{(\dagger,2)}(\mathcal{X}_{B_1}) \times h_f^{(2)} \left( \begin{pmatrix} 1 & \underline{u} \\ 0 & 1 \end{pmatrix} \underline{y} \right) = \psi(\underline{u}) \omega_{\underline{\epsilon}'}^{(\dagger,2)}(\mathcal{X}_{B_1}) \times h_f^{(2)}(\underline{y})$$

Our volume form is  $d\underline{y} = c_F |\underline{t}|^{-1} d\underline{u} \times d^\times \underline{t} d\underline{k}$  where all these measures are product over local measures, we require  $\text{vol}_{dk_v}(K_v) = 1$  and  $\text{vol}_{\underline{u}} U(\mathbb{Q}) \backslash U(\mathbb{A}) = 1$  the constant  $c_F$  is essentially the inverse of the discriminant.

Then

$$\begin{aligned} & \int_{U(\mathbb{A})} \omega_{\underline{\epsilon}}^{(\dagger,1)}(\mathcal{X}_{A_1}) \times h_f^{(1)} \left( \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & \underline{u} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \underline{t} & 0 \\ 0 & 1 \end{pmatrix} \underline{k} \right) \omega_{\underline{\epsilon}'}^{(\dagger,2)}(\mathcal{X}_{B_1}) \times h_f^{(2)} \left( \begin{pmatrix} b & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & \underline{u} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \underline{t} & 0 \\ 0 & 1 \end{pmatrix} \underline{k} \right) d\underline{u} \\ &= \begin{cases} \omega_{\underline{\epsilon}'}^{(\dagger,2)}(\mathcal{X}_{B_1}) \times h_f^{(1)} \left( \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \underline{t} & 0 \\ 0 & 1 \end{pmatrix} \right) \omega_{\underline{\epsilon}'}^{(\dagger,2)}(\mathcal{X}_{B_1}) \times h_f^{(2)} \left( \begin{pmatrix} b & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \underline{t} & 0 \\ 0 & 1 \end{pmatrix} \right) \underline{k} & \text{if } a + b = 0 \\ 0 & \text{else} \end{cases} \end{aligned} \quad (8.188)$$

and therefore our integral becomes

$$\int_{T(\mathbb{Q}) \backslash T(\mathbb{A})} \int_{K_f^{H,0}} \sum_{a \in F^\times} \omega_{\underline{\epsilon}}^{(\dagger,1)}(\mathcal{X}_{A_1}) \times h_f^{(1)} \left( \begin{pmatrix} at & 0 \\ 0 & 1 \end{pmatrix} \underline{k}_f \right) \omega_{\underline{\epsilon}'}^{(\dagger,2)}(\mathcal{X}_{B_1}) \times h_f^{(2)} \left( \begin{pmatrix} -at & 0 \\ 0 & 1 \end{pmatrix} \underline{k}_f \right) |\underline{t}|^s d\underline{k}_f d^\times \underline{t} \quad (8.189)$$

and since  $T(\mathbb{Q}) = F^\times$  for the value of the integral

$$\int_{T(\mathbb{A})} \int_{K_f^{H,0}} \omega_{\underline{\epsilon}}^{(\dagger,1)}(\mathcal{X}_{A_1}) \times h_f^{(1)} \left( \begin{pmatrix} \underline{t} & 0 \\ 0 & 1 \end{pmatrix} \underline{k}_f \right) \omega_{\underline{\epsilon}'}^{(\dagger,2)}(\mathcal{X}_{B_1}) \times h_f^{(2)} \left( \begin{pmatrix} -\underline{t} & 0 \\ 0 & 1 \end{pmatrix} \underline{k}_f \right) |\underline{t}|^s d\underline{k}_f d^\times \underline{t} \quad (8.190)$$

In the variable  $\underline{k}_f$  our functions are right invariant under  $K_f^H$  hence the integral over  $\underline{k}_f$  is actually a finite sum. Then for a fixed value of  $\underline{k}_f$  our functions are products of local Whittaker functions, i.e.

$$\omega_{\underline{\epsilon}}^{(\dagger,1)}(\mathcal{X}_{A_1}) \times h_f^{(1)} \left( \begin{pmatrix} \underline{t} & 0 \\ 0 & 1 \end{pmatrix} \underline{k}_f \right) = \omega_{\underline{\epsilon}}^{(\dagger,1)}(\mathcal{X}_{A_1}) \left( \begin{pmatrix} t_\infty & 0 \\ 0 & 1 \end{pmatrix} \right) \prod_{\mathfrak{p}} h_{\mathfrak{p}}^{(1)} \left( \begin{pmatrix} t_{\mathfrak{p}} & 0 \\ 0 & 1 \end{pmatrix} k_{\mathfrak{p}} \right)$$

$$\omega_{\underline{\epsilon}'}^{(\dagger,2)}(\mathcal{X}_{B_1}) \times h_f^{(2)} \left( \begin{pmatrix} -\underline{t} & 0 \\ 0 & 1 \end{pmatrix} \underline{k}_f \right) = \omega_{\underline{\epsilon}'}^{(\dagger,2)}(\mathcal{X}_{B_1}) \left( \begin{pmatrix} t_\infty & 0 \\ 0 & 1 \end{pmatrix} \right) \prod_{\mathfrak{p}} h_{\mathfrak{p}}^{(2)} \left( \begin{pmatrix} t_{\mathfrak{p}} & 0 \\ 0 & 1 \end{pmatrix} k_{\mathfrak{p}} \right)$$

and hence our integral becomes

$$\text{vol}_{d\mathbf{k}}(K_f^H) \sum_{\mathbf{k}_f \in K_f^{H,0}/K_f} \prod_v \int_{T(F_v)} h_{\mathbf{p}}^{(1)}\left(\begin{pmatrix} t_v & 0 \\ 0 & 1 \end{pmatrix} k_v\right) h_{\mathbf{p}}^{(2)}\left(\begin{pmatrix} t_v & 0 \\ 0 & 1 \end{pmatrix} k_v\right) |t_v|_v^s d^\times t_v$$

The local Whittaker functions are explicitly given to us. We look at the different places. We begin with a finite place  $\mathfrak{p}$  and if  $\pi_{\mathfrak{p}}$  is unramified, i.e.  $K_{\mathfrak{p}}^H$  is maximal. We have to compute

$$\int_{T(F_{\mathfrak{p}})} h_{\mathbf{p}}^{(1)}\left(\begin{pmatrix} t_{\mathfrak{p}} & 0 \\ 0 & 1 \end{pmatrix}\right) h_{\mathbf{p}}^{(2)}\left(\begin{pmatrix} t_{\mathfrak{p}} & 0 \\ 0 & 1 \end{pmatrix}\right) |t_{\mathfrak{p}}|_{\mathfrak{p}}^s d^\times t_{\mathfrak{p}}$$

We recall the explicit formulas for the values of  $h_{\mathbf{p}}^{(1)}\left(\begin{pmatrix} t_{\mathfrak{p}} & 0 \\ 0 & 1 \end{pmatrix}\right)$  and  $h_{\mathbf{p}}^{(2)}\left(\begin{pmatrix} t_{\mathfrak{p}} & 0 \\ 0 & 1 \end{pmatrix}\right)$ . Let  $\omega(\pi_{1,\mathfrak{p}})$  be the Satake parameter of  $\pi_{1,\mathfrak{p}}$  then - as usual - we define

$$\alpha_{\mathfrak{p}} = N(\mathfrak{p})^{\frac{1}{2}} \omega(\pi_{1,\mathfrak{p}})\left(\begin{pmatrix} \varpi_{\mathfrak{p}} & 0 \\ 0 & 1 \end{pmatrix}\right), \beta_{\mathfrak{p}} = N(\mathfrak{p})^{\frac{1}{2}} \omega(\pi_{1,\mathfrak{p}})\left(\begin{pmatrix} 1 & 0 \\ 0 & \varpi_{\mathfrak{p}} \end{pmatrix}\right)$$

we introduced the Euler factor in (4.100)

$$L(\pi_{1,\mathfrak{p}}, s) = \frac{1}{(1 - \alpha_{\mathfrak{p}} N(\mathfrak{p})^{-s})(1 - \beta_{\mathfrak{p}} N(\mathfrak{p})^{-s})}.$$

After expanding we get

$$L(\pi_{1,\mathfrak{p}}, s) = \sum_{n=0}^{\infty} \left( \sum_{\nu=0}^n \alpha_{\mathfrak{p}}^{n-\nu} \beta_{\mathfrak{p}}^{\nu} \right) N(\mathfrak{p})^{-ns} = \sum_{n=0}^{\infty} h_{\mathbf{p}}^{(1)}\left(\begin{pmatrix} \varpi_{\mathfrak{p}}^{f(\pi_{\mathfrak{p}})} & 0 \\ 0 & 1 \end{pmatrix}\right) N(\mathfrak{p})^{n(1-s)} \quad (8.191)$$

For the second factor we have  $\pi_{2,\mathfrak{p}} = \pi_{1,\mathfrak{p}}^{\vee}$  hence the Satake parameter is  $\omega(\pi_{2,\mathfrak{p}}) = \omega(\pi_{1,\mathfrak{p}})^{-1}$ . If we now define

$$\beta'_{\mathfrak{p}} = N(\mathfrak{p})^{\frac{1}{2}} \omega(\pi_{2,\mathfrak{p}})\left(\begin{pmatrix} \varpi_{\mathfrak{p}} & 0 \\ 0 & 1 \end{pmatrix}\right), \alpha'_{\mathfrak{p}} = N(\mathfrak{p})^{\frac{1}{2}} \omega(\pi_{2,\mathfrak{p}})\left(\begin{pmatrix} 1 & 0 \\ 0 & \varpi_{\mathfrak{p}} \end{pmatrix}\right),$$

then we have  $\alpha_{\mathfrak{p}} \beta'_{\mathfrak{p}} = \alpha'_{\mathfrak{p}} \beta_{\mathfrak{p}} = N(\mathfrak{p})$  and hence  $\alpha_{\mathfrak{p}} \alpha'_{\mathfrak{p}} \beta_{\mathfrak{p}} \beta'_{\mathfrak{p}} = N(\mathfrak{p})^2$ .

We get

$$L(\pi_{2,\mathfrak{p}}, s) = \sum_{n=0}^{\infty} \left( \sum_{\nu=0}^n \alpha_{\mathfrak{p}}^{m-\nu} \beta_{\mathfrak{p}}^{\nu} \right) N(\mathfrak{p})^{-s} = \sum_{n=0}^{\infty} h_{\mathbf{p}}^{(2)}\left(\begin{pmatrix} \varpi_{\mathfrak{p}}^{f(\pi_{\mathfrak{p}})} & 0 \\ 0 & 1 \end{pmatrix}\right) N(\mathfrak{p})^{n(1-s)} \quad (8.192)$$

We express the inner sums in terms of the semi-simple Satake parameters (See (??) and remark after it), we have

$$\alpha_{\mathfrak{p}}^{n-\nu} \beta_{\mathfrak{p}}^{\nu} = \left(\frac{\alpha_{\mathfrak{p}}}{\beta_{\mathfrak{p}}}\right)^{\frac{n}{2}-\nu} (\alpha_{\mathfrak{p}} \beta_{\mathfrak{p}})^{\frac{n}{2}} = \omega^{(1)}(\pi_{1,\mathfrak{p}})^{\frac{n}{2}-\nu} (\alpha_{\mathfrak{p}} \beta_{\mathfrak{p}})^{\frac{n}{2}},$$

the same holds for  $\pi_{2,\mathfrak{p}}$ . and therefore

$$h_{\mathbf{p}}^{(1)}\left(\begin{pmatrix} t_{\mathfrak{p}} & 0 \\ 0 & 1 \end{pmatrix}\right) h_{\mathbf{p}}^{(2)}\left(\begin{pmatrix} t_{\mathfrak{p}} & 0 \\ 0 & 1 \end{pmatrix}\right) = \left( \sum_{\nu=0}^n \omega^{(1)}(\pi_{\mathfrak{p}})^{\frac{n}{2}-\nu} \right)^2$$

Now we have the following identity in power series ring  $\mathbb{Z}[u, 1/u][[t]]$  :

$$\frac{1-t^2}{(1-u^2t)(1-t)^2(1-u^{-2}t)} = \sum_{n=0}^{\infty} \left( \sum_{\nu=0}^n u^{n-2\nu} \right)^2 t^n$$

(According to Jacquet ([?], ??) the proof is a refreshing exercise.) We put  $t = N(\mathfrak{p})^{-1-s}$  and  $u = \omega^{(1)}(\pi_{\mathfrak{p}})$  then this identity gives us

$$\begin{aligned} & \frac{1 - N(\mathfrak{p})^{-2-s}}{(1 - \omega^{(1)}(\pi_{\mathfrak{p}})N(\mathfrak{p})^{-1-s})(1 - N(\mathfrak{p})^{-1-s})^2(1 - \omega^{(1)}(\pi_{\mathfrak{p}})^{-1}(N(\mathfrak{p})^{-1-s}))} = \\ & \sum_{n=0}^{\infty} \Phi_{\mathfrak{p}}^{(1)} \left( \begin{pmatrix} \varpi_{\mathfrak{p}}^{f(\pi_{\mathfrak{p}})} & 0 \\ 0 & 1 \end{pmatrix} \right) \Phi_{\mathfrak{p}}^{(2)} \left( \begin{pmatrix} \varpi_{\mathfrak{p}}^{f(\pi_{\mathfrak{p}})} & 0 \\ 0 & 1 \end{pmatrix} \right) N(\mathfrak{p})^{n(-1-s)} \end{aligned} \quad (8.193)$$

The factor in the numerator is the inverse of the local factor of the Dedekind  $\zeta_{\mathfrak{p}}(\cdot)$  function at  $s+2$ , the factor  $(1 - N(\mathfrak{p})^{-1-s})$  in the denominator gives us the local zeta factor  $\zeta_{\mathfrak{p}}(1+s)$ . The remaining expression gives us local factor of the adjoint  $L$ -function, i.e.

$$\frac{1}{(1 - \omega^{(1)}(\pi_{\mathfrak{p}})N(\mathfrak{p})^{-1-s})(1 - N(\mathfrak{p})^{-1-s})(1 - \omega^{(1)}(\pi_{\mathfrak{p}})^{-1}(N(\mathfrak{p})^{-1-s}))} = L(\pi_{\mathfrak{p}}, \text{Ad}, s+1) \quad (8.194)$$

Therefore we get for an unramified  $\pi_{\mathfrak{p}}$

$$\frac{\zeta_{\mathfrak{p}}(1+s)}{\zeta_{\mathfrak{p}}(2+s)} L(\pi_{\mathfrak{p}}, \text{Ad}, s+1) = \int_{T(F_{\mathfrak{p}})} h_{\mathfrak{p}}^{(1)} \left( \begin{pmatrix} t_{\mathfrak{p}} & 0 \\ 0 & 1 \end{pmatrix} \right) h_{\mathfrak{p}}^{(2)} \left( \begin{pmatrix} t_{\mathfrak{p}} & 0 \\ 0 & 1 \end{pmatrix} \right) |t_{\mathfrak{p}}|_{\mathfrak{p}}^s d^{\times} t_{\mathfrak{p}} \quad (8.195)$$

In this book we try to avoid the discussions of the subtle phenomena at ramified  $\pi_{\mathfrak{p}}$ , therefore we assume that a similar formula also holds at the finite number of ramified places, we may take this as definition of the local Euler factor at these places.

### The integral at the archimedean places

We treat the cases of a real and a complex place separately.

A) The place  $v$  is real. We have the two generators  $\omega_{v,\pm}^{\dagger}$  (4.133) and the factor at our place  $v$  becomes

$$\int_0^{\infty} \langle \omega_{v,+}^{\dagger}(H) \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right), \omega_{v,-}^{\dagger}(V) \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) \rangle > t^s \frac{dt}{t}$$

We recall the definition of the generators and then our integral becomes (up to some small power of 2 (to be fixed later))

$$\langle (X-Y \otimes i)^n, (X+Y \otimes i)^n \rangle > \int_0^{\infty} t^{n+2} e^{-4\pi t} t^s \frac{dt}{t} = \langle (X-Y \otimes i)^n, (X+Y \otimes i)^n \rangle > \frac{\Gamma(n+2+s)}{(4\pi)^{n+2+s}}$$

The factor in front is

$$\langle (X - Y \otimes i)^n, (X + Y \otimes i)^n \rangle = \sum_{\nu, \mu} i^{\mu - \nu} \binom{n}{\nu} \binom{n}{\mu} \langle X^{n-\nu} Y^\nu, X^{n-\mu} Y^\mu \rangle$$

and by definition we have (See ??)  $\langle X^{n-\nu} Y^\nu, X^{n-\mu} Y^\mu \rangle = 0$  unless we have  $\nu + \mu = n$  and then

$$\langle X^{n-\nu} Y^\nu, X^\nu Y^{n-\mu} \rangle = \binom{n}{\nu}^{-1}.$$

Hence we see that one of the binomial factor cancels and we find  $\langle (X - Y \otimes i)^n, (X + Y \otimes i)^n \rangle = (2i)^n$ . So we finally get

$$\int_0^\infty \langle \omega_{v,+}(H) \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right), \omega_{v,-}(V) \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) \rangle t^s \frac{dt}{t} = (2i)^n \frac{\Gamma(n+2+s)}{(4\pi)^{n+2+s}} \tag{8.196}$$

Let us call this last expression  $\mathfrak{G}_v(n, s)$

B) The place  $v$  is a complex place, this case is more difficult (interesting, amusing). In this case we have to evaluate

$$\int_0^\infty \langle \omega_v^{\dagger,1}(X_{0,v}) \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right), \omega_v^{\dagger,2}(X_{1,v}, X_{-1,v}) \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) \rangle t^s \frac{dt}{t} =$$

We have the explicit formula (8.197) for these these factors for  $Z = X_{v,0}$  or  $Z = (X_{1,v}, X_{-1,v})$  we have

$$\omega^{\dagger, \bullet}(Z) = \sum_{\mu=-n}^n \Phi_{\lambda, \mu} \otimes \left( \sum_{\mu_1 + \mu_2 = |\mu|} e_{\mu_1}^b \otimes \bar{e}_{\mu_2}^b \right) \tag{8.197}$$

Hence we multiply and get a sum

$$\sum_{\mu, \mu'} \Phi_{\lambda, \mu} \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) \Phi_{\lambda, \mu'} \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) \otimes \left( \sum_{\mu_1, \mu'_1} T(\mu_1, \mu'_1) \right)$$

where

$$T(\mu_1, \mu'_1) = \langle \rho_\lambda \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) (e_{\mu_1}^b \otimes \bar{e}_{\mu_2}^b), \rho_{\lambda'} \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) (e_{\mu'_1}^b \otimes \bar{e}_{\mu'_2}^b) \rangle.$$

But since our pairing is invariant under the action of  $G(\mathbb{R})$  we can ignore the  $\rho_\lambda \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right)$  and we find that the value

$$\langle e_{\mu_1}^b \otimes \bar{e}_{\mu_2}^b, e_{\mu'_1}^b \otimes \bar{e}_{\mu'_2}^b \rangle = \begin{cases} \binom{|\mu|}{\mu_1} \binom{|\mu|}{|\mu| - \mu_1} & \text{if } \mu_1 = -\mu'_1, \mu_2 = -\mu'_2 \\ 0 & \text{else} \end{cases} \tag{8.198}$$

and taking into account the formulas for the pairing we get for the integrand

$$\sum_{\mu=n}^{-n} \Phi_{\lambda,\mu} \Phi_{\lambda,-\mu} \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) \left( \sum_{\mu_1} \binom{|\mu|}{\mu_1} \binom{|\mu|}{|\mu| - \mu_1} \right) t^s$$

We have our explicit expressions for the  $\Phi_{\lambda,\mu}$  we have to compute the Mellin transform

$$\begin{aligned} & \frac{4\pi^{2n}}{\Gamma(n+1)^2} \int_0^\infty K_\mu(2\pi t) K_{-\mu}(2\pi t) t^{2n+4+s} \frac{dt}{t} = \\ & \frac{4\pi^{2n}}{\Gamma(n+1)^2} \frac{\Gamma(n+1+\mu+s/2)\Gamma(n+1-\mu+s/2)\Gamma(n+2+s/2)^2}{(2\pi)^{2n+4+s}} = \quad (8.199) \\ & \frac{\Gamma(n+2+s/2)^2}{4\pi^{4+s}\Gamma(n+1)^2} \Gamma(n+1+\mu+s/2)\Gamma(n+1-\mu+s/2) \end{aligned}$$

To get the value of the above integral we have to sum over the  $\mu$ . Hence finally we get

$$\begin{aligned} \mathfrak{G}_v(n, s) & := \int_0^\infty \langle \omega_v^{\dagger,1}(X_{0,v}) \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right), \omega_v^{\dagger,2}(X_{1,v}, X_{-1,v}) \left( \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) \rangle t^s \frac{dt}{t} = \\ & \frac{1}{4\pi^{s+4}} \frac{\Gamma(n+2+s/2)^2}{\Gamma(n+1)} \left( \sum_{\mu=-n}^{\mu=n} \Gamma(n+1+\mu+s/2)\Gamma(n+1-\mu+s/2) \left( \sum_{\mu_1} \binom{|\mu|}{\mu_1} \binom{|\mu|}{|\mu| - \mu_1} \right) \right) \quad (8.200) \end{aligned}$$

Eventually we are interested in the value at  $s = 0$ , then we have the following identity, which I checked experimentally

$$\sum_{\mu=-n}^{\mu=n} \Gamma(n+1+\mu)\Gamma(n+1-\mu) \left( \sum_{\mu_1} \binom{|\mu|}{\mu_1} \binom{|\mu|}{|\mu| - \mu_1} \right) = \Gamma(2n+2) \quad (8.201)$$

so that  $\mathfrak{G}_v(n, 0) = \frac{(n+1)^2}{4\pi^4} (2n+1)!$  We put  $\mathfrak{G}_\infty(\lambda, s) = \prod_{v \in S_\infty} \mathfrak{G}(n_v, s)$   
Then we see that the value our integral in (8.187) eventually is given by

$$c_F \frac{\zeta_F(1+s)}{\zeta_F(s+2)} \mathfrak{G}_\infty(\lambda, s) L(\pi_f, \text{Ad}, s+1) \quad (8.202)$$

We have to take the residue at  $s = 0$ , we know that all the factors except  $\zeta_F(s+1)$  are holomorphic at  $s = 0$  and hence we get for the cup product of the two cohomology classes

$$\begin{aligned} & \frac{1}{\Omega_\epsilon^{(1)}(\pi_f)} \mathcal{F}_1^{(1)}(\omega_\epsilon^{(1)} \times h_f^{(1)}) \cup \frac{1}{\Omega_\epsilon^{(2)}(\pi_f)} \mathcal{F}_1^{(2)}(\omega_\epsilon^{(2)} \times h_f^{(1)}) = \\ & \frac{1}{\Omega_\epsilon^{(1)}(\pi_f)\Omega_\epsilon^{(2)}(\pi_f)} \frac{\text{Res}_{s=0} \zeta_F(1+s)}{\zeta_F(2)} \mathfrak{G}_\infty(\lambda, 0) L(\pi_f, \text{Ad}, 1) \quad (8.203) \end{aligned}$$

We know that this number is in  $\frac{1}{\Delta_\lambda} \mathcal{O}_F$ . Let  $\mathfrak{p}$  be a prime in  $\mathcal{O}_F[\frac{1}{\Delta_\lambda}]$ , which divides this number, i.e.

$$\mathfrak{p}^{\delta(\pi_f)} \parallel \frac{1}{\Omega_\epsilon^{(1)}(\pi_f)\Omega_\epsilon^{(2)}(\pi_f)} \frac{\text{Res}_{s=0} \zeta_F(1+s)}{\zeta_F(2)} \mathfrak{G}_\infty(\lambda, 0) L(\pi_f, \text{Ad}, 1).$$

We have the non degenerate pairing

$$H^{r_1+r_2}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1} \otimes \mathcal{O}_F[\frac{1}{\Delta_\lambda}])_{\text{int},!} \times H^{r_1+2r_2}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_2} \otimes \mathcal{O}_F[\frac{1}{\Delta_\lambda}])_{\text{int},!} \rightarrow \mathcal{O}_F[\frac{1}{\Delta_\lambda}]$$

and the decomposition into saturated Hecke submodules

$$\begin{aligned} & H^{r_1+r_2}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1} \otimes \mathcal{O}_F[\frac{1}{\Delta_\lambda}])_{\text{int},!} \supset \\ & H^{r_1+r_2}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1} \otimes \mathcal{O}_F[\frac{1}{\Delta_\lambda}])_{\text{int},!(\pi_{1,f})} \oplus H^{r_1+r_2}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1} \otimes \mathcal{O}_F[\frac{1}{\Delta_\lambda}])_{\text{int},!(\pi_{1,f})}^\perp \end{aligned} \quad (8.204)$$

where  $^\perp$  means that we take the saturated direct sum over the  $\pi'_f \neq \pi_f$ . We introduce the quotient

$$\begin{aligned} & \tilde{H}^{r_1+r_2}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1} \otimes \mathcal{O}_F[\frac{1}{\Delta_\lambda}])_{\text{int},!(\pi_{1,f})} = \\ & H^{r_1+r_2}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1} \otimes \mathcal{O}_F[\frac{1}{\Delta_\lambda}])_{\text{int},!(\pi_{1,f})} / H^{r_1+r_2}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1} \otimes \mathcal{O}_F[\frac{1}{\Delta_\lambda}])_{\text{int},!(\pi_{1,f})}^\perp \end{aligned} \quad (8.205)$$

and the above pairing induces a non degenerate pairing

$$\tilde{H}^{r_1+r_2}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1} \otimes \mathcal{O}_F[\frac{1}{\Delta_\lambda}])_{\text{int},!(\pi_{1,f})} \times H^{r_1+2r_2}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_2} \otimes \mathcal{O}_F[\frac{1}{\Delta_\lambda}])_{\text{int},!(\pi_{2,f})} \rightarrow \mathcal{O}_F[\frac{1}{\Delta_\lambda}] \quad (8.206)$$

We choose a character  $\underline{\epsilon}'$  then  $H^{r_1+2r_2}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_2} \otimes \mathcal{O}_F[\frac{1}{\Delta_\lambda}])_{\text{int},!(\underline{\epsilon}' \times \pi_{2,f})}$  is a free  $\mathcal{O}_F[\frac{1}{\Delta_\lambda}]$  module of rank one, a generator is  $y_0 = [\frac{1}{\Omega_{\underline{\epsilon}'}}(\pi_f)^{(2)} \mathcal{F}_1^{(2)}(\omega_{\underline{\epsilon}}^{(2)} \times h_f^{(1)})]$ . Let  $x_0$  be the corresponding generator in  $H^{r_1+r_2}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1} \otimes \mathcal{O}_F[\frac{1}{\Delta_\lambda}])_{\text{int},!(\underline{\epsilon} \times \pi_{1,f})}$ . We can find an element  $\tilde{x}_0 \in \tilde{H}^{r_1+r_2}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1} \otimes \mathcal{O}_F[\frac{1}{\Delta_\lambda}])_{\text{int},!(\underline{\epsilon} \times \pi_{1,f})}$  such that  $\langle x_0, y_0 \rangle = 1$ . Let  $\varpi_{\mathfrak{p}}$  be a uniformizer for  $\mathfrak{p}$  we lift  $x_0$  to an element  $\tilde{x}_0^* \in H^{r_1+r_2}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1} \otimes \mathcal{O}_F[\frac{1}{\Delta_\lambda}])_{\text{int},!}$  and we can write ( we localize at  $\mathfrak{p}$ )

$$\tilde{x}_0^* = \frac{x_0 + z}{\varpi_{\mathfrak{p}}^m} \text{ with } z \in H^{r_1+r_2}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1} \otimes \mathcal{O}_F[\frac{1}{\Delta_\lambda}])_{\text{int},!(\pi_{1,f})}^\perp. \quad (8.207)$$

Then

$$1 = \langle \tilde{x}_0, y_0 \rangle = \frac{\langle x_0, y_0 \rangle}{\varpi_{\mathfrak{p}}^m} \quad (8.208)$$

and this implies  $m = \delta(\pi_f)$ .

We can slightly modify this argument. Any element  $\tilde{x} \in \tilde{H}^{r_1+r_2}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1} \otimes \mathcal{O}_F[\frac{1}{\Delta_\lambda}])_{\text{int},!}$  can be written as above in the form

$$\tilde{x} = \frac{x + y}{\varpi_{\mathfrak{p}}^{\delta(\pi_f)}}$$

and then the map  $\tilde{x} \mapsto y \pmod{\varpi_{\mathfrak{p}}^{\delta(\pi_f)}} H^{r_1+r_2}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1} \otimes \mathcal{O}_F[\frac{1}{\Delta_\lambda}])_{\text{int},!(\pi_{1,f})}^\perp$  yields an inclusion

$$\tilde{H}^{r_1+r_2}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1})_{\text{int},!(\pi_{1,f})} \otimes \mathcal{O}_F[\frac{1}{\Delta_\lambda}]/\mathfrak{p}^{\delta(\pi_{\mathfrak{p}})} \hookrightarrow H^{r_1+r_2}(\mathcal{S}_{K_f^H}^H, \tilde{\mathcal{M}}_{\lambda_1})_{\text{int},!(\pi_{1,f})}^\perp \otimes \mathcal{O}_F[\frac{1}{\Delta_\lambda}]/\mathfrak{p}^{\delta(\pi_{\mathfrak{p}})} \quad (8.209)$$

This has consequences for congruences, it is clear how to formulate a theorem corresponding to Theorem (3.3.2).

At this place references to Urban, Dimitroff and Namikawa will be added.

### Fixing the period

The actual of computation the period may be a highly non trivial. Actually this may even not be so important. But it is indeed of interest to compute the factorization of the  $L$ -values, this means we have to compute the numbers

$$\text{ord}_{\mathfrak{p}}\left(\frac{L(\pi_f \otimes \chi, \mu)}{\Omega(\pi_f, \omega_\epsilon)}\right) \quad (8.210)$$

for as many  $\mathfrak{p} \subset \mathcal{O}_F$  as possible.

Of course we have problems to fix the period if the class number of  $\mathcal{O}_F$  is not one, but this does not matter for the above question, we have to fix a prime  $p$  and then we have to choose a good period locally at  $p$ . This means we solve the problem alluded to in (8.87) only locally at  $p$ .

We discuss this problem in a very special case where our group  $G = \text{Gl}_2$ , the maximal compact subgroup  $K_f = \prod_p \text{Gl}_2(\mathbb{Z}_p)$  and our coefficient system  $\mathcal{M}$  is the module of homogenous polynomials  $P(X, Y)$  of degree  $n$  and coefficients in  $\mathbb{Z}$ . Hence the Hecke algebra  $\mathcal{H}_{K_f} = \otimes'_p \mathcal{H}_{K_p}$  is unramified at all primes  $p$  it is commutative. Our isotypical component  $\pi_f$  defines an ideal  $\mathcal{I}(\Pi_f) \subset \mathcal{H}_{K_f}$  and the quotient  $\mathcal{H}_{K_f}/\mathcal{I}(\Pi_f)$  is an order in the field  $\mathbb{Q}(\mathcal{I}(\Pi_f)) = \mathcal{H}_{K_f}/\mathcal{I}(\Pi_f) \otimes \mathbb{Q}$ , which is finite extension of  $\mathbb{Q}$ . (I replaced  $\pi_f$  by  $\Pi_f$  because the ideal does not change if we conjugate  $\pi_f$  the ideal  $\mathcal{I}(\Pi_f)$  is associated to the Galois orbit of  $\pi_f$ . I prefer to view  $\mathbb{Q}(\Pi_f)$  as an abstract extension of  $\mathbb{Q}$ .) This ideal  $\mathcal{I}(\Pi_f)$  defines a submodule  $H_!^1(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})_{\text{int}}(\mathcal{I}(\Pi_f)) = \text{Ann}(\mathcal{I}(\Pi_f))$ , this is the submodule annihilated by  $\mathcal{I}(\Pi_f)$ .

We can think of  $\pi_f$  as simply being a modular cusp form  $f$  of weight  $k = n+2$ . To get our isotypical module  $H_!^1(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathcal{O}_F})_{\text{int}}$  we have to find a homomorphism  $\sigma : \mathcal{H}_{K_f}/\mathcal{I}(\Pi_f) \rightarrow \mathcal{O}_F$  and then

$$H_!^1(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathcal{O}_F})_{\text{int}}(\pi_f) = H_!^1(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})_{\text{int}}(\mathcal{I}(\Pi_f)) \otimes_{\mathcal{H}_{K_f}, \sigma} \mathcal{O}_F \quad (8.211)$$

We have the action of complex conjugation, i.e. of  $\pi_0(G(\mathbb{R}))$ , on the cohomology  $H_!^1(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})_{\text{int}}(\mathcal{I}(\Pi_f))$  we get the decomposition (up to an isogeny of degree  $2^m$ )

$$H_!^1(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})_{\text{int}}(\mathcal{I}(\Pi_f)) \supset H_{!,+}^1(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})_{\text{int}}(\mathcal{I}(\Pi_f)) \oplus H_{!,-}^1(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})_{\text{int}}(\mathcal{I}(\Pi_f)) \quad (8.212)$$

and after taking the tensor product by  $\mathbb{Q}$  both summands become one dimensional vector spaces over  $\mathbb{Q}(\mathcal{I}(\Pi_f))$ . But it is by no means clear that the integral modules are isomorphic.

This becomes a little bit better if tensor by  $\mathcal{O}_F$  then then we have again

$$H_1^1(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathcal{O}_F})_{\text{int}}(\pi_f) \supset H_{1,+}^1(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathcal{O}_F})_{\text{int}}(\pi_f) \oplus H_{1,-}^1(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\mathcal{O}_F})_{\text{int}}(\pi_f) \tag{8.213}$$

and now the two summands are  $\mathcal{O}_F$  modules of rank one and get their structure as Hecke-modules from the homomorphism  $\sigma$ . ( In a sense  $\pi_f = (\Pi_f, \sigma)$ ) But still they are not necessarily isomorphic. If we want to define the periods we need class number one. But instead of defining a period we define a local periods. If we tensor the semilocal ring  $\mathcal{O}_{F,p} = \mathcal{O}_F \otimes \mathbb{Z}_{(p)}$  then the class number problem disappears we can choose a period such that we get an isomorphism

$$\Omega_{\pm}^{(p)}(\pi_f)^{-1} \cdot \mathcal{F}_1^{(1)}(\omega_{\pm}) : \bigotimes_p \mathcal{W}_{\mathcal{O}_{F,p}(\pi_f)}(\pi_f, \tau) \xrightarrow{\sim} H_{1,\pm}^1(\mathcal{S}_{K_f}^G, \mathcal{M}_{\mathcal{O}_{F,p}})(\pi_f) \tag{8.214}$$

Recall that we viewed  $\pi_f$  as a modular form  $f$  of weight  $k$  we change the notation for the periods slightly and denote them by  $\Omega_{\pm}^{(p)}(f)$ . Our character  $\chi$  will now be unramified which implies that it is uniquely determined by its type  $\mu$ . We put  $\nu = \mu + 1$  then we get for  $\nu = 1, 2, \dots, k - 1$  the following integrality statement

$$\Delta(f) \frac{L(f, \nu)}{\Omega_{\pm}(f)} \in \mathcal{O}_{F,p} \tag{8.215}$$

But we can still do a little bit better. Recall that we have to evaluate our integral cohomology class on a modular symbol  $c_{\mu}$ . This modular symbol is a relative cycle from 0 to  $i\infty$  (just along the imaginary axis) loaded by an element  $e_{\mu} = X^{\mu}Y^{n-\mu}$ , we denote it by  $[0, i\infty] \times e_{\nu}$ . The index  $\mu$  runs from zero to  $n$ . This is a relative cycle and defines a class in  $H_1(\mathcal{S}_{K_f}^G, \partial(\mathcal{S}_{K_f}^G), \tilde{\mathcal{M}})$ . We have the boundary operator

$$\partial : H_1(\mathcal{S}_{K_f}^G, \partial(\mathcal{S}_{K_f}^G), \tilde{\mathcal{M}}) \rightarrow H_0(\partial(\mathcal{S}_{K_f}^G), \tilde{\mathcal{M}}). \tag{8.216}$$

We represent the boundary by the circle at  $i\infty$  then it is clear that

$$\partial(e_{\mu}) = e_{\mu} - we_{\mu} \tag{8.217}$$

and we see that  $\partial(e_{\mu})$  is a torsion class if  $\mu \neq 0, n$ . Not only that it is a torsion class it is annihilated by a power of the Hecke-operator  $T_p^n$ . This implies that  $T_p^n([0, i\infty] \times e_{\mu})$  can be lifted to a homology class in  $\tilde{E}_{\mu} \in H_1(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}})$ . But then it is clear that the evaluation of our generator  $\xi_{\pm}$  in  $H_{1,\pm}^1(\mathcal{S}_{K_f}^G, \mathcal{M}_{\mathcal{O}_{F,p}})(\pi_f)$  on

this lifted cycle gives an integral value. Since  $\xi_{\pm}$  is an eigenvalue for the Hecke operator we get for  $\mu = 1, \dots, n-1$  and  $\nu = \mu + 1$

$$\langle \xi_{\pm}, \tilde{E}_{\mu} \rangle = \pi_f(T_p)^n \langle \xi, e_{\mu} \rangle = \pi_f(T_p)^n \frac{L(f, \nu)}{\Omega_{\epsilon(\nu)}(f)} \in \mathcal{O}_{F,p} \quad (8.218)$$

This means that we do not need the factor  $\Delta(f)$  in front.

We choose a prime  $\mathfrak{p}$  in  $\mathcal{O}_F$  lying above  $p$ . Let us now assume that  $\pi_f(T_p)$  is a unit, i.e.  $f$  is ordinary at  $\mathfrak{p}$  then we can conclude that

$$\frac{L(f, \nu)}{\Omega_{\pm}^{(p)}(f)} \in \mathcal{O}_{F,\mathfrak{p}}$$

and consequently

$$\text{ord}_{\mathfrak{p}}\left(\frac{L(f, \nu)}{\Omega_{\pm}^{(p)}(f)}\right) \geq 0 \text{ for all } 2 \leq \nu \leq k-2 \quad (8.219)$$

We also know what we should expect at the argument  $\nu = k-1$ . In this case  $\partial(e_n)$  is not a torsion element, but we know that for all primes  $\ell$  the element  $(\ell^{k-1} + 1 - \pi_f(T_{\ell}))\partial(e_n)$  is annihilated by a power of  $T_p$ . If  $b_{\mathfrak{p}}(f)$  is the minimum of the numbers  $\text{ord}_{\mathfrak{p}}(\ell^{k-1} + 1 - \pi_f(T_{\ell}))$  then we can conclude that

$$\text{ord}_{\mathfrak{p}}\left(\frac{L(f, \mu)}{\Omega_{\pm}^{(p)}(f)}\right) + b_{\mathfrak{p}}(f) \geq 0 \text{ for } \mu = 1, k-1 \quad (8.220)$$

Hence we can say (still a little bit conjecturally and using Poincaré-duality and the fact that the modular symbols  $c_{\mu}$  generate the relative homology. (H. Gebertz, Diploma Thesis Bonn .)

*If  $\mathfrak{p}$  is ordinary then the numbers  $\Omega_{\pm}^{(p)}(f)$  are the right periods at  $\mathfrak{p}$  if and only if one of the non negative numbers in the + or - part of the lists (8.219), (8.220)*

$$\mathcal{L}_{f,\mathfrak{p}} = \left\{ \text{ord}_{\mathfrak{p}}\left(\frac{L(f, k-1)}{\Omega_{-}^{(p)}(f)}\right) + b_{\mathfrak{p}}(f), \text{ord}_{\mathfrak{p}}\left(\frac{L(f, k-2)}{\Omega_{+}^{(p)}(f)}\right), \dots, \text{ord}_{\mathfrak{p}}\left(\frac{L(f, \nu)}{\Omega_{\pm}^{(p)}(f)}\right), \dots \right\}$$

*is zero.*

This discussion is interesting in view of the conjectures on congruences in [31]. In this note we make conjectures about some congruences between Siegel and elliptic modular forms, these congruences are congruences modulo a "large" prime and I do not really say what a large prime should be. Already in [31] I address the issue that we have to choose the right period, but there the choice is rather ad hoc.

Now we have a better recipe. The heuristic argument for the existence of the congruences only works if the prime is ordinary for the modular form  $f$ . But in this case we have now a much more precise rule to compute the period. For an ordinary prime  $\mathfrak{p}$  we should expect a congruence if for one of the members in the above lists we find a strictly positive value. Here we should still be a little bit more careful, my heuristic argument predicts congruences if  $\mathfrak{p}$  occurs in the denominator of a ratio

$$\text{ord}_p\left(\frac{\mathcal{L}_{f,p}(\nu)}{\mathcal{L}_{f,p}(\nu+1)}\right) < 0, \nu = k-2, k-3, \dots, k/2+1$$

so we should pay attention to possible cancellations.

Checking the list of the list of the modular forms of weight 12,16,18,20,22,26 we find that the only cases of ordinary primes for which we expect congruences are indeed the cases  $k = 22, \ell = 41$  and  $k = 26, \ell = 29, 43, 97$  and they are already in [31]. Here is no cancellation.

It will be very interesting to check the case of the two dimensional space of cusp forms of weight 24. In this case the field  $F = \mathbb{Q}(\sqrt{144169})$ . Again we find very few instances of ordinary candidates, these are the primes dividing 73, 179 and the congruences have been checked.

But apart from these two cases we have the two divisors of 13, they occur rather frequently in our list  $\mathcal{L}_{f,p}$  and it seems to be interesting to see what happens.

The modular form  $f$  of weight 24 has an expansion with coefficients in  $\mathbb{Q}(\omega)$  where  $\omega^2 = 144169$ , we write the first few terms

$$f(q) = q + 12(45 - \omega)q^2 + 36(4715 + 16 \cdot \omega)q^3 + 32(395729 - 405 \cdot \omega)q^4 + 1410(25911 + 128 \cdot \omega)q^5 \dots + 658(3325311035 - 23131008 \cdot \omega)q^{13} \dots \tag{8.221}$$

and this provides the two modular forms  $f^{(+)}$  (resp.  $f^{(-)}$ ) with real coefficients which we get if we send  $\omega$  to the positive root  $\sqrt{144169}$  (resp. negative root).

We have the periods  $\Omega_{\pm}(f^{(+)})$ ,  $\Omega_{\pm}(f^{(-)})$  and we know that

$$\frac{L(f^{(+)}, \nu)}{\Omega_{\epsilon(\nu)}(f^{(+)})}, \frac{L(f^{(-)}, \nu)}{\Omega_{\epsilon(\nu)}(f^{(-)})} \in \mathbb{Q}(\sqrt{144169}) \tag{8.222}$$

Looking at the norms of these numbers we find some factors of 13. The prime 13 decomposes in  $\mathbb{Z}[\omega]$  and we see that the two prime factors above thirteen are given by the homomorphism  $\phi_5 : \omega \mapsto 5 \pmod{13}$ . and  $\phi_8 : \omega \mapsto 8 \pmod{13}$ . We check that  $f^{(+)}$  is ordinary at  $\phi_8$  but not at  $\phi_5$ . But if we look at the prime factor decomposition of  $\frac{L(f^{(+)}, \nu)}{\Omega_{\epsilon(\nu)}(f^{(+)})}$  then we see that  $\phi_5$  occurs non trivially but  $\phi_8$  does not. Hence we do not expect the existence of a Siegel modular form and a congruence modulo  $\phi_5$  because  $\phi_5$  is not ordinary for  $f^{(+)}$ . The prime  $\phi_8$  is ordinary for  $f^{(+)}$  but this prime does not occur in the  $L$ -values.

**Anton’s Congruence**

The issue to fix the period becomes even more delicate once we allow ramification. Let us consider the case of the congruence subgroup  $\Gamma_0(p)$ , this means that our open compact subgroup will be  $K_{0,f}(p) = \prod_{q:p \nmid q} \text{Gl}_2(\mathbb{Z}_q) \times K_0(p)$ . Again we can determine the periods locally at a prime  $\ell$  by evaluating period integrals against certain modular symbols. The point is that we have more modular symbols, because we allow ramification. To get control over these modular symbols we consider the representation  $\text{Ind}_{K_{0,f}(p)}^{K_f} \mathbf{1}$ , i.e. the induced from the trivial representation of  $K_{0,f}(p)$  to the maximal compact subgroup  $K_f$ . This

representation can be viewed as a representation of  $\mathrm{Gl}_2(\mathbb{F}_p)$ , it is of dimension  $p+1$  and it has the Steinberg-module  $\mathrm{St}_p$  of dimension  $p$ . Then we can consider the cohomology  $H^1(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_n \otimes \mathrm{St}_p)$ , and new forms  $f$  for  $\Gamma_0(p)$  correspond to eigenclasses in  $H_1^1(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_n \otimes \mathrm{St}_p)$ .

We can construct modular symbols with coefficients in  $\tilde{\mathcal{M}}_n \otimes \mathrm{St}_p$ . The standard torus  $T(\mathbb{F}_p)$  acts on  $\mathrm{St}_p$  and under this action we get a decomposition into eigenspaces (we invert the divisors of  $p(p-1)$  let  $R = \mathbb{Z}[\frac{1}{p(p-1)}]$ )

$$\mathrm{St}_p \otimes R = \bigoplus_{\chi: \mathbb{F}_p^\times \rightarrow \mu_{p-1}} Re_\chi \tag{8.223}$$

(The trivial character occurs two times)

Hence we can define modular symbols  $e_\mu \otimes e_\chi$  where  $e_\mu$  is as above. Then we get integrality for the values

$$\frac{L(f \otimes \chi, \mu)}{\Omega_{\epsilon(\mu, \chi)}(f)} G(\chi, \tau) \tag{8.224}$$

Since we inverted  $p$  the Gaussian sum does not play any role. We assume that the modular symbols  $e_\mu \otimes e_\chi$  generate the relative homology  $H_1(\mathcal{S}_{K_f}^G, \partial(\mathcal{S}_{K_f}^G), \tilde{\mathcal{M}}_n \otimes \mathrm{St}_p \otimes R)$ . Hence we can fix the periods locally at a prime  $\ell$  which does not divide  $p(p-1)$  and which is ordinary for  $f$ . We compute the  $L$ -values and then we must have

$$\mathrm{ord}_\ell\left(\frac{L(f \otimes \chi, \mu)}{\Omega_{\epsilon(\mu, \chi)}(f)}\right) \geq 0 \tag{8.225}$$

and for both signs  $\epsilon(\mu, \chi)$  at least one of these numbers has to be zero. Here  $\ell$  runs over the divisors of  $\ell$  in  $\mathcal{O}_F[\zeta_{p-1}]$ .

We want to consider the special case of modular forms of weight 4 for  $\Gamma_0(p)$ . In this case we have only three critical values  $L(f \otimes \chi, \mu)$  for  $\mu = 1, 2, 3$ .

We are interested in this case because we want to understand the conjectures in [31] also in the case of a non regular coefficient system, especially we want to look at the case of the trivial coefficient system, i.e. the case where the representation is one dimensional. Then we find modular forms of weight four in the boundary cohomology and this forces us to allow ramification. But we want to keep the ramification as small as possible.

We start from the group  $G = \mathrm{GSp}_2/\mathbb{Z}$ , we choose as level subgroup the group  $K_f = K_{f,p}^G = \prod_{q:q \neq p} G(\mathbb{Z}_q) \times K_0(p)$ , where  $K_0(p)$  is the group of  $\mathbb{Z}_p$  valued points of the unique non special maximal parahoric subgroup scheme  $\mathcal{P}_{\gamma_1}$ . (Here  $\gamma_1$  is the fundamental weight attached to the short root viewed as a cocharacter, we have  $\langle \gamma_1, \alpha_1 \rangle = 1, \langle \gamma_1, \alpha_2 \rangle = 0$  ). This choice  $K_{f,p}^G$  defines an arithmetic subgroup  $\Gamma_p \subset \mathrm{GSp}_2(\mathbb{Q})$  which is called the paramodular group.

We consider the homomorphism

$$H^3(\mathcal{S}_{K_f}^G, R) \xrightarrow{r} H^3(\partial(\mathcal{S}_{K_f}^G), R) \tag{8.226}$$

The right hand side contains a contribution coming from the cuspidal cohomology of the stratum of the Siegel parabolic subgroup, this is the contribution  $H_1^1(\mathcal{S}_{K_f}^M, H^2(\mathfrak{u}_P, R))$ . The point is that now that  $K_f^M = K_{0,f}(p) =$

$\prod_{q:q \neq p} \text{Gl}_2(\mathbb{Z}_q) \times K_0(p)$ , which we introduced above. The  $M$ -module  $H^2(\mathfrak{u}_P, R)$  is the standard three dimensional representation. Hence this cohomology is described by the space of modular forms of weight 4 for the group  $\Gamma_0(p)$ .

Any modular (new) form  $f$  of weight 4 for  $\Gamma_0(p)$ , yields a contribution

$$H^1_!(\mathcal{S}^M_{K_f}, H^2(\mathfrak{u}_P, R))[f]$$

of rank one over  $R \otimes \mathcal{O}_F$ . Let us consider the inverse image  $H^3(\mathcal{S}^G_{K_f}, R)[f] = r^{-1}(H^1(\mathcal{S}^M_{K_f}, H^2(\mathfrak{u}_P, R)[f]))$ . We consider the restriction

$$H^3(\mathcal{S}^G_{K_f}, R)[f] \xrightarrow{r_f} H^1(\mathcal{S}^M_{K_f}, H^2(\mathfrak{u}_P, R)[f]) \tag{8.227}$$

We invoke results from Eisenstein cohomology. Schwermer has shown: This restriction map is surjective if and only if we have  $L(f, 2) = 0$  otherwise we encounter a pole of an Eisenstein class.

I also discuss an analogous situation in the appendix of [Ha-Eis]. There I assume that we have no ramification, but I discuss non trivial non regular coefficient systems. A rather speculative computation using the comparison between the Lefschetz and the topological trace formula suggests that in this case

*$r_f$  has a non trivial kernel  $H^3_!(\mathcal{S}^G_{K_f}, R)[f]$  if and only if the sign of the functional equation for  $L(f, s)$  is minus one.*

Let us believe that the same is true in this case (and if we do not believe in the trace formula we could also try to explain this kernel as a Gritsenko lift) and we get the exact sequence

$$0 \rightarrow H^3_!(\mathcal{S}^G_{K_f}, R)[f] \rightarrow H^3(\mathcal{S}^G_{K_f}, R)[f] \xrightarrow{r_f} H^1(\mathcal{S}^M_{K_f}, H^2(\mathfrak{u}_P, R)[f]), \tag{8.228}$$

where  $H^3_!(\mathcal{S}^G_{K_f}, R)[f]$  is the Scholl motive attached to  $f$ . This yields an extension class of motives

$$\mathcal{X}(f) \in \text{Ext}^1(R(-2), H^3_!(\mathcal{S}^G_{K_f}, R)[f]). \tag{8.229}$$

Tony Scholl suggests to attach a number to such an extension. More precisely he suggests to construct a suitable biextension, this can be done by the Anderson construction introducing an auxiliary prime  $p_0$ .) and then this number should be essentially

$$\frac{\frac{L'(f, 2)}{\Omega_+(f)}}{\frac{L(f, 3)}{\Omega_-(f)}} \tag{8.230}$$

Under this assumption the denominator  $\frac{L(f, 3)}{\Omega_-(f)}$  becomes interesting. Since we fixed the period, we can ask whether ordinary primes  $l$  dividing this number yield denominators of Eisenstein classes and hence congruences. Such a congruence has been detected by Anton Mellit in the case  $p = 61$  and  $\ell = 43$ . Checking the tables of W. Stein we find that for  $p = 61$  the cohomology  $H^1_!(\mathcal{S}^M_{K_f}, H^2(\mathfrak{u}_P, R))$  is of rank  $2 \times 15$  and decomposes into a 12-dimensional and a 18 dimensional

piece (over  $\mathbb{Q}$ ). The 6 dimensional piece corresponds to a modular cusp form  $f$  of weight 4 for  $\Gamma_0(61)$  its coefficients lie in a field of degree 6 over  $\mathbb{Q}$ . The sign in the functional equation is  $-1$  and we should look for the prime decomposition of the number

$$\frac{L(f, 3)}{\Omega_-(f)} \tag{8.231}$$

over  $\ell = 43$ . We know that there is a Siegel modular form for  $\Gamma_{61}$  which is not a Gritsenko lift and satisfies the congruence (Poor-Yuen). The question is whether a divisor  $l|\ell$  occurs in the value above. But then it becomes clear that we have to obey strict rules to fix the period.

We may also check some other primes  $p$  and compute the ratios in (??) and look whether they are divisible by interesting primes and whether these primes yield congruences for non Gritsenko lifts.

### 8.2.6 The $L$ -functions

Again I have to say a few words concerning  $L$ -functions.

To get the automorphic  $L$ -functions at the unramified places we have to introduce the dual group  $G^\vee(\mathbb{C})$  ( this is  $\mathrm{Gl}_2(\mathbb{C})$  in this case ) and a finite dimensional representation  $r$  of this group. The definition of the dual group is designed in such a way that the Satake parameter  $\omega_p$  of an unramified representation at  $p$  can be interpreted as a semi simple conjugacy class in  $G^\vee(\mathbb{C})$  (see [La]). Therefore we can form the expression

$$L(\pi_p, r, s) = \det(\mathrm{Id} - r(\omega_p)p^{-s})^{-1}$$

and then the global  $L$  function  $L(\pi, r, s)$  is defined as the product over all these unramified  $L$ -factors times a product over suitable  $L$ -factors at the finite primes. If we do this for our automorphic forms on  $\mathrm{Gl}_2$  and if  $r = r_1$  is the tautological representation of  $\mathrm{Gl}_2(\mathbb{C})$  then we get the local  $L$ -factors

$$L(\pi_p, r_1, s) = \frac{1}{(1 - \lambda_{p,2}(p)p^{-s})(1 - \lambda_{p,1}(p)p^{-s})}$$

and we see that it differs by a shift by  $1/2$  from our previous definition. Our earlier  $L$ -function was the motivic  $L$ -function, its definition does not require the additional datum  $r$ . Our automorphic form  $\pi$  defines a motive  $\mathbb{M}(\pi)$ . This motive has the disadvantage that it does not occur in the cohomology of a variety, it occurs only after we apply a Tate twist to it. The central character  $\omega(\pi)$  has type  $x \mapsto x^n$  and defines a Tate motive. The automorphic form  $\pi \otimes \omega(\pi)^{-1} = \pi^\vee$  occurs in the cohomology

$$H^1(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}[-n]) \supset H^1(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}[-n])(\pi \otimes \omega(\pi)^{-1}) = H^1(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}[-n])(\pi^\vee)$$

where  $\mathcal{M}_n[-n]$  is obtained by twisting the original module by the  $-n$ -th power of the determinant. (See [Ha-Eis], III). This motive occurs in the cohomology of a quasiprojective scheme ( See also [Scholl] ) Now we adopt the point of view that  $\pi_f$  is a pair  $(\Pi_f, \iota)$  (See 1.2.6) and then  $\mathbb{M}(\pi)$  defines a system of  $\ell$ -adic

representations  $\rho(\pi)_\ell$  which are also labelled by the  $\iota : \mathbb{Q}(\pi_f) \rightarrow \bar{\mathbb{Q}}$ . Then it is Delignes theorem that for unramified primes

$$L(\pi_p, r_1, s - \frac{1}{2}) = L_p((\mathbb{M}(\pi^\vee), s) = \det(\text{Id} - \rho(F_p)_p^{-1} | \mathbb{M}(\pi^\vee)_\ell p^{-s})$$

for a suitable choice of  $\ell \neq p$ .

**Weights and Hodge numbers**

We may of course look at the motives  $\mathbb{M}(\pi)$  which are attached to an eigenspace in  $H_1^1(S_{K_f}^G, \tilde{\mathcal{M}}[-k])(\pi)$  in other words we twisted the natural module  $\mathcal{M}_n$  by the  $-k$ -th power of the determinant. Again we get an  $\ell$ -adic representation  $\rho_\ell$  and the Weil conjectures imply that the eigenvalues of the inverse Frobenius  $\rho_\ell(F_p^{-1})$  all have the same absolute value  $p^{\frac{2k-n+1}{2}}$ . The number  $2k - n + 1$  is usually called the weight  $w(\rho_\ell)$  of the Galois representation or also the weight  $w(\mathbb{M}(\pi))$  of the motive  $\mathbb{M}(\pi)$ .

The central character  $\omega(\pi)$  of  $\pi$  has a type and if we make the natural identification of  $G_m$  with the centre then the type of  $\omega(\pi)$  is an integer  $\text{type}(\omega(\pi)) \in \mathbb{Z}$  and the formula for the weight is

$$w(\mathbb{M}(\pi)) = -\text{type}(\omega(\pi)) + 1.$$

This weight plays a role if we want to get a first understanding of the analytic properties of the motivic  $L$ -functions. Its abcizza of convergence is the line  $\Re(s) = w(\mathbb{M}(\pi)) + 1$ .

The special case  $k = n$  is special, because in this case our motive occurs in the cohomology of a variety. The eigenvalues of the Frobenius are algebraic integers and the non zero Hodge numbers are  $h^{n+1,0}$  and  $h^{0,n+1}$ . If  $k$  is arbitrary then the centre acts on  $\mathcal{M}_n[-k]$  by the character  $t(k) = n - 2k$  and the non zero Hodge numbers will be  $h^{1+\frac{n-t(k)}{2}, -\frac{n+t(k)}{2}}$ . We notice that for an isotypic component  $H_1^1(S_{K_f}^G, \tilde{\mathcal{M}}[-k])(\pi)$  the number  $t(k)$  is the type of the central character  $\omega(\pi)$ .

**8.2.7 The special values of  $L$ -functions**

We now observe that the local  $L$  factors  $L(\mathbb{M}(\pi^\vee \otimes (\chi^{(1)})^{-1}), s)$  which we introduced in 2.2.6 are actually the local  $L$ -factors of the motivic  $L$ -function, i.e.

$$L(\mathbb{M}(\pi^\vee \otimes (\chi^{(1)})^{-1}), s) = L(\mathbb{M}(\pi^\vee \otimes (\chi^{(1)})^{-1}), s)$$

**Theorem 8.2.2.** *With these notations we can give a formula for the composition*

$$J_{c_{\chi, \iota}} \circ \Omega_\varepsilon(\pi_f)^{-1} \cdot \mathcal{F}_1^{(1)}(\omega_\varepsilon) = \frac{L(\mathbb{M}(\pi^\vee \otimes (\chi^{(1)})^{-1}), 1)}{\Omega_\varepsilon(\pi_f)} \cdot I^{loc}(\pi_f, \chi_f^{-1})$$

**Applications**

We evaluate this formula at elements  $\psi_f \in \mathcal{W}(\pi_f, \tau)_{\mathcal{O}(\pi_f, \chi)}$  and an element  $\underline{g}_f \in G(\mathbb{A}_f)$ . We get  $\Omega_\varepsilon(\pi_f)^{-1} \cdot \mathcal{F}_1^{(1)}(\omega_\varepsilon)(\psi_f) = \tilde{\psi}_f \in H_{1, \varepsilon}^1(S_{K_f}^G, \tilde{\mathcal{M}})_{\mathcal{O}(\pi_f, \chi)}$  and

$$J_{c_{\chi, \iota}}(\psi_f)(\underline{g}_f) = \frac{L(\mathbb{M}(\pi^\vee \otimes (\chi^{(1)})^{-1}), 1)}{\Omega_\varepsilon(\pi_f)} \cdot I^{loc}(\pi_f, \chi_f^{-1})(\psi_f)(\underline{g}_f)$$

We have seen that  $J_{c_{x,!}}(\psi_f)(\underline{g}_f)d(\underline{g}_f)$  (Lemma 2.2) is an integer and it is obvious that  $d(\underline{g}_f) = \prod_p d(g_p)$ . If we choose for  $\psi_f$  an element which is also a product  $\psi_f(\underline{g}_f) = \prod_p \psi_p(g_p)$  then we get

$$J_{c_{x,!}}(\psi_f)(\underline{g}_f) \prod_p d(g_p) = \frac{L(\mathbb{M}(\pi^\vee \otimes (\chi^{(1)})^{-1}), 1)}{\Omega_\varepsilon(\pi_f)} \cdot \prod_p I_p^{loc}(\pi_p, \chi_p^{-1})(\psi_p)(g_p)d(g_p)$$

The factors in the products over all primes are equal to one at almost all places. Then we have to optimize the choices of  $\psi_p$  and  $g_p$ . First of all we can choose these data such that all local factors are different from zero. Then we conclude that we have an invariance under Galois for the  $L$ -values

$$\left( \frac{L(\mathbb{M}(\pi^\vee \otimes (\chi^{(1)})^{-1}), 1)}{\Omega_\varepsilon(\pi_f)} \right)^\sigma = \chi^{(1)}(\underline{t}_\sigma) \frac{L(\mathbb{M}((\pi^\vee \otimes (\chi^{(1)})^{-1})^\sigma), 1)}{\Omega_\varepsilon(\pi_f^\sigma)}$$

We may observe that the characters  $\chi^{(1)}$  can be written as product of a Dirichlet character and a power of the Tate character, i.e.  $\chi^{(1)} = \phi \cdot \alpha^{-\nu}$  where  $\nu = 0, \dots, n$ . Now we can write

$$\mathbb{M}(\pi^\vee \otimes (\chi^{(1)})^{-1}) = \mathbb{M}(\pi^\vee \otimes \phi^{-1}) \otimes \mathbb{Z}(\nu)$$

and

$$L(\mathbb{M}(\pi^\vee \otimes (\chi^{(1)})^{-1}), 1) = L(\mathbb{M}(\pi^\vee \otimes \phi^{-1}), 1 + \nu)$$

and the arguments  $1 + \nu$  are exactly the critical arguments for the motive  $\mathbb{M}(\pi^\vee \otimes \phi^{-1})$  in the sense of Deligne.

Of course we are now able to prove also some integrality results, it is clear that the left hand side is integral, more precisely it is an element in  $\mathcal{O}(\pi_f, \chi_f)$ . Now we have to work with local representations to find out under which conditions we can force the product of local factors to be a unit or at least to bound the primes dividing it. Hence we have a tool to show that

$$\frac{L(\mathbb{M}(\pi^\vee \otimes (\chi^{(1)})^{-1}), 1)}{\Omega_\varepsilon(\pi_f)} \in \mathcal{O}(\pi_f, \chi_f)$$

at least if we invert a few more primes.

### The arithmetic interpretation

It is clear that we have some control of the primes that have to be inverted. I call them *small* primes. The main reason why I am interested in the integrality statement for these special values is, that I want to understand what it means if a *large* prime divides these values.

I strongly believe that the large primes dividing these  $L$ -values are related to the denominators of Eisenstein classes for the cohomology of the symplectic group, what this means will be explained in 5.6 and we also refer to the notes [kolloquium.pdf]. In the following section I want to give some idea how such a relationship between the arithmetic properties of the  $L$ -values and the integral structure of the cohomology as a Hecke-module should look like.

# Chapter 9

## Eisenstein cohomology

Our starting point is a smooth group scheme  $\mathcal{G}/\mathrm{Spec}(\mathbb{Z})$  whose generic fiber  $G = \mathcal{G} \times_{\mathbb{Z}} \mathbb{Q}$  is reductive and quasisplit. We assume the group scheme is reductive over the largest possible open subset of  $\mathrm{Spec}(\mathbb{Z})$  and at the remaining places it is given by a maximal parahoric group scheme structure. If  $G$  is split, then we assume that  $\mathcal{G}$  is split. We define  $K_f = \mathcal{G}(\hat{\mathbb{Z}}) = \prod_p \mathcal{G}(\mathbb{Z}_p) \subset G(\mathbb{A}_f)$

We choose a Borel subgroup  $B/\mathbb{Q}$  and a torus  $T/\mathbb{Q} \subset B/\mathbb{Q}$ . We assume that  $T(\mathbb{A}_f) \cap K_f = T(\hat{\mathbb{Z}})$  is maximal compact in  $T(\mathbb{A}_f)$ . Let  $\lambda \in X^*(T)$  be a highest weight, let  $\mathcal{M}_\lambda$  be a highest weight module attached to this weight. It is a  $\mathbb{Z}$ -module, the module  $\mathcal{M}_\lambda \otimes \mathbb{Q}$  is a highest weight module for the group  $G/\mathbb{Q}$ . We consider

### 9.1 The Borel-Serre compactification

We consider our space

$$\mathcal{S}_{K_f}^G = G(\mathbb{Q}) \backslash G(\mathbb{A}) / K_\infty K_f$$

and its Borel-Serre compactification

$$i : \mathcal{S}_{K_f}^G \rightarrow \bar{\mathcal{S}}_{K_f}^G.$$

Our highest weight module  $\mathcal{M}_\lambda$  provides a sheaf  $\tilde{\mathcal{M}}_\lambda$  on these spaces.

We have an isomorphism

$$H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \xrightarrow{\sim} H^\bullet(\bar{\mathcal{S}}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$$

for any coefficient system  $\tilde{\mathcal{M}}_\lambda$  coming from a rational representation  $\mathcal{M}$  of  $G(\mathbb{Q})$ . The boundary  $\partial \bar{\mathcal{S}}_K$  is a manifold with corners. It is stratified by submanifolds

$$\partial \bar{\mathcal{S}}_K = \bigcup_P \partial_P \mathcal{S}_{K_f}^G,$$

where  $P$  runs over the  $G(\mathbb{Q})$  conjugacy classes of proper parabolic subgroups defined over  $\mathbb{Q}$ . We identify the set of conjugacy classes of parabolic subgroups

with the set of representatives given by the parabolic subgroups that contain our standard Borel subgroup  $B/\mathbb{Q}$ . Then we have

$$H^\bullet(\partial_P \mathcal{S}_{K_f^G}^G, \tilde{\mathcal{M}}_\lambda) = H^\bullet(P(\mathbb{Q}) \backslash G(\mathbb{A}) / K_\infty K_f, \tilde{\mathcal{M}}_\lambda)$$

We have a finite coset decomposition

$$G(\mathbb{A}_f) = \bigcup_{\xi_f} P(\mathbb{A}_f) \xi_f K_f,$$

for any  $\xi_f$  put  $K_f^P(\xi_f) = P(\mathbb{A})_f \cap \xi_f K_f \xi_f^{-1}$ . Then we have

$$P(\mathbb{Q}) \backslash X \times G(\mathbb{A}_f) / K_f = \bigcup_{\xi_f} P(\mathbb{Q}) \backslash X \times P(\mathbb{A}_f) / K_f^P(\xi_f) \xi_f,$$

If  $R_u(P) \subset P$  is the unipotent radical, then

$$M = P/R_u(P)$$

is a reductive group. For any open compact subgroup  $K_f \subset G(\mathbb{A}_f)$  (resp. for  $K_\infty \subset G_\infty$ ) we define  $K_f^M(\xi_f) \subset M(\mathbb{A}_f)$  (resp.  $K_\infty^M \subset M_\infty$ ) to be the image of  $K_f^P(\xi_f)$  in  $M(\mathbb{A}_f)$  (resp.  $M_\infty$ ). We put

$$\mathcal{S}_{K_f(\xi_f)}^M = M(\mathbb{Q}) \backslash M(\mathbb{A}) / K_\infty^M K_f^M(\xi_f)$$

and get a fibration

$$\pi_P : P(\mathbb{Q}) \backslash X \times P(\mathbb{A}_f) / K_f^P(\xi_f) \rightarrow M(\mathbb{Q}) \backslash M(\mathbb{A}) / M(\mathbb{Q}) \backslash K_\infty^M \times K_f^M(\xi_f)$$

where the fibers are of the form  $\Gamma_U \backslash R_u(P)(\mathbb{R})$  and where  $\Gamma_U \subset U(\mathbb{Z})$  is of finite index and defined by some congruence condition dictated by  $K_f^P(\xi_f)$ . The Lie-algebra  $\mathfrak{u}$  of  $R_u(P)$  is a free  $\mathbb{Z}$ -module and it is clear that we have an integral version of the van Est theorem which says:

*If  $R = \mathbb{Z}[\frac{1}{N}]$  where a suitable set of primes has been inverted then*

$$H^\bullet(\Gamma_U \backslash R_u(P)(\mathbb{R}), \tilde{\mathcal{M}}_R) \xrightarrow{\sim} H^\bullet(\mathfrak{u}, \tilde{\mathcal{M}}_R).$$

*More precisely we know that the local coefficient system  $R^\bullet \pi_{P*}(\tilde{\mathcal{M}})$  is obtained from the rational representation of  $M$  on  $H^\bullet(\mathfrak{u}, \mathcal{M})$ .*

Hence we get

$$H^\bullet(\partial_P \mathcal{S}, \tilde{\mathcal{M}}_R) = \bigcup_{\xi_f} H^\bullet(\mathcal{S}_{K_f^M(\xi_f)}^M, \widetilde{H^\bullet(\mathfrak{u}, \mathcal{M})}_R),$$

and

$$H^\bullet(\mathfrak{u}, \mathcal{M}_R) = \bigoplus_{w \in W^P} H^{l(w)}(\mathfrak{u}, \mathcal{M}_R)(w \cdot \lambda),$$

where  $W^P$  is the set of Kostant representatives of  $W/W^M$  and where  $w \cdot \lambda = (\lambda + \rho)^w - \rho$  and  $\rho$  is the half sum of positive roots.

The primes which we have to be inverted should be those which are smaller than the coefficients of the dominant weights in the highest weight of  $\mathcal{M}$ . But at this point we may have to enlarge the set of small primes.

We conclude

*The cohomology of the boundary strata  $\partial_P \mathcal{S}_{K_f}^G$  with coefficients in  $\mathcal{M}$  can be computed in terms of the cohomology of the reductive quotient, where we have coefficients in the cohomology of the Lie algebra of the unipotent radical with coefficients in  $\mathcal{M}$*

In the following considerations we sometimes suppress the subscripts  $K_f, K_{K_f}^M$  and so on. Then we mean that the considerations are valid for a fixed level or that we have taken the limit over the  $K_f$ . (See the remarks below concerning induction)

### 9.1.1 The two spectral sequences

The covering of the boundary by the strata  $\partial_P \mathcal{S}$  provides a spectral sequence, which converges to the cohomology of the boundary. We can introduce the simplex  $\Delta$  of types of parabolic subgroups, the vertices correspond to the maximal ones and the full simplex corresponds to the minimal parabolic. To any type of a parabolic  $P$  let  $d(P)$  its rank, we make the convention that  $d(P) - 1$  is equal to the dimension of the corresponding face in the simplex. Let  $M = M_P = P/R_u(P)$  be the reductive quotient (the Levi quotient). If  $Z_M/\mathbb{Q}$  is the connected component of the identity of the center of  $M/\mathbb{Q}$  then  $d(P)$  is also the dimension of the maximal split subtorus of  $Z_M/\mathbb{Q}$  minus the dimension of the maximal split subtorus of  $Z_G/\mathbb{Q}$ . The covering yields a spectral sequence whose  $E_1^{\bullet, \bullet}$  term together with the differentials of our spectral sequence is given by

$$0 \rightarrow E_1^{0,q} = \bigoplus_{P,d(P)=1} H^q(\partial_P \mathcal{S}, \mathcal{M}) \xrightarrow{d_1^{0,q}} \dots \rightarrow \bigoplus_{P,d(P)=p+1} H^q(\partial_P \mathcal{S}, \mathcal{M}) \xrightarrow{d_1^{p,q}} \dots \tag{9.1}$$

where the boundary map  $d_1^{p,q}$  is obtained from the restriction maps (See [Gln]). There is also a homological spectral sequence which converges to the cohomology of the boundary. It can be written as a spectral sequence for the cohomology with compact supports. Let  $d$  be the dimension of  $\mathcal{S}$  then we have a complex

$$\rightarrow \bigoplus_{P,d(P)=p+1} H_c^{d-1-p-q-1}(\partial_P \mathcal{S}, \mathcal{M}) \xrightarrow{\delta_1} \bigoplus_{P,d(P)=p} H_c^{d-1-p-q}(\partial_P \mathcal{S}, \mathcal{M}) \rightarrow \dots \tag{9.2}$$

and therefore the  $E_{\bullet, \bullet}^1$  term is

$$E_{p,q}^1 = \bigoplus_{P,d(P)=p} H_c^{d-1-p-q}(\partial_P \mathcal{S}, \mathcal{M})$$

the (higher) differential go from  $(p, q)$  to  $(p - r, q + 1 - r)$ .

### 9.1.2 Induction

The description of the cohomology of a boundary stratum is a little bit clumsy, since we are working with the coset decomposition. The reason is that we are working on a fixed level, if we consider cohomology with integral coefficients. If we have rational coefficients then we can pass to the limit. Then

$$H^\bullet(\partial_P \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}) = \lim_{K_f} H^\bullet(P(\mathbb{Q}) \backslash G(\mathbb{A}) / K_\infty K_f, \tilde{\mathcal{M}}) = \\ \text{Ind}_{\pi_0(M(\mathbb{R})) \times P(\mathbb{A}_f)}^{\pi_0(G(\mathbb{R}) \times G(\mathbb{A}_f))} \lim_{K_f^M} H^\bullet(\mathcal{S}_{K_f^M}^M, \widetilde{H^\bullet(\mathfrak{u}, \mathcal{M})}) = \text{Ind}_{\pi_0(M(\mathbb{R})) \times P(\mathbb{A}_f)}^{\tilde{G}(\mathbb{A})} H^\bullet(\mathcal{S}^M, \widetilde{H^\bullet(\mathfrak{u}, \mathcal{M})}),$$

where the induction is ordinary group theoretic induction. We should keep in our mind that the  $\pi_0(M(\mathbb{R})) \times P(\mathbb{A}_f)$ -modules are in fact  $\pi_0(M(\mathbb{R})) \times M(\mathbb{A}_f)$ -modules. We need some simplification in the notation and we will write for any such  $\pi_0(M(\mathbb{R})) \times M(\mathbb{A}_f)$  module  $H$

$$\text{Ind}_{\pi_0(M(\mathbb{R})) \times P(\mathbb{A}_f)}^{\tilde{G}(\mathbb{A})} H = I_M^G H$$

We will use the same notation for an induction from the torus  $T$  to  $M$ .

Under certain conditions we also have the notion of induction for Hecke - modules and we can work with integral coefficient systems. This will be discussed at another occasion.

But I want to mention that in the case that  $K_f$  is a hyperspecial maximal compact subgroup ( in the cases where we are dealing with a split semi-simple group scheme over  $\text{Spec}(\mathbb{Z})$  we can take  $K_f = \prod \mathbb{G}(\mathbb{Z}_p)$  (see 1.1)) then  $G(\mathbb{Q}_p) = P(\mathbb{Z}_p)K_p = B(\mathbb{Z}_p)K_p$  the group theoretic induction followed by taking  $K_f$  invariants gives back the original module. In this case we do not have to induce!

Of course we have to understand the coefficient systems  $H^\bullet(\mathfrak{u}, \mathcal{M})$ , for this we need the theorem of Kostant which will be discussed in the next section.

### 9.1.3 A review of Kostants theorem

At this point we can make the assumption that our group  $G/\mathbb{Q}$  is quasisplit, we also assume that  $G^{(1)}/\mathbb{Q}$  is simply connected. Then we may assume that  $\mathcal{M}_{\mathbb{Z}}$  is irreducible and of highest weight  $\lambda$ . Let  $B/\mathbb{Q}$  be a Borel subgroup, we choose a torus  $T/\mathbb{Q} \subset B/\mathbb{Q}$ . Let  $X^*(T) = \text{Hom}(T \times_{\mathbb{Q}} \bar{\mathbb{Q}}, \mathbb{G}_m \times_{\mathbb{Q}} \bar{\mathbb{Q}})$  be the character module, it comes with an action of a finite Galois group  $\text{Gal}(F/\mathbb{Q})$ , here  $F$  is the smallest sub field of  $\bar{\mathbb{Q}}$  over which  $G/\mathbb{Q}$  splits. Let  $T^{(1)}/\mathbb{Q} \subset T/\mathbb{Q}$  the maximal torus in  $G^{(1)}/\mathbb{Q}$ , then  $X^*(T^{(1)})$  contains the set  $\Delta$  of roots, the subset  $\Delta^+$  of positive roots (with respect to  $B$ .) The set of simple roots is identified to a finite index set  $I = \{1, 2, \dots, r\}$ , i.e we write the set of simple roots as  $\pi = \{\alpha_1, \dots, \alpha_i, \dots, \alpha_r\} \subset \Delta^+$ . We assume that the numeration is somehow adapted the Dynkin diagram. The finite Galois group  $\text{Gal}(F/\mathbb{Q})$  acts on  $I$  and  $\pi$  by permutations. Attached to the simple roots we have the dominant fundamental weights  $\{\dots, \gamma_i, \dots, \gamma_j, \dots\}$  they are related to the simple roots by the rule

$$2 \frac{\langle \gamma_i, \beta_j \rangle}{\langle \beta_j, \beta_j \rangle} = \delta_{i,j}.$$

The dominant fundamental weights form a basis of  $X^*(T^{(1)})$ .

Our maximal torus  $T/\mathbb{Q}$  is up to isogeny the product of  $T^{(1)}$  and the central torus  $C/\mathbb{Q}$ , i.e.  $T = T^{(1)} \cdot C$  and the restriction of characters yields an injection

$$j : X^*(T) \rightarrow X^*(T^{(1)}) \oplus X^*(C),$$

this becomes an isomorphism if we tensorize by the rationals

$$X_{\mathbb{Q}}^*(T) = X^*(T) \otimes \mathbb{Q} \xrightarrow{\sim} X_{\mathbb{Q}}^*(T^{(1)}) \oplus X_{\mathbb{Q}}^*(C).$$

This isomorphism gives us canonical lifts of elements in  $X^*(T^{(1)})$  or  $X^*(C)$  to elements in  $X_{\mathbb{Q}}^*(T)$  which will be denoted by the same letter. Especially the fundamental weights  $\gamma_1, \dots, \gamma_i, \dots$  are elements in  $X_{\mathbb{Q}}^*(T)$ .

Let  $\lambda \in X^*(T)$  be a dominant weight, our decomposition allows us to write it as

$$\lambda = \sum_{i \in I} a_i \gamma_i + \delta = \lambda^{(1)} + \delta$$

we have  $a_i \in \mathbb{Z}, a_i \geq 0$  and  $\delta \in X^*(C)$ . To such a dominant weight  $\lambda$  we have an absolutely irreducible  $G \times F$ -module  $\mathcal{M}_{\lambda}$ .

We consider maximal parabolic subgroups  $P/\mathbb{Q} \supset B/\mathbb{Q}$ . These parabolic subgroups are given by the choice of a  $\text{Gal}(F/\mathbb{Q})$  orbit  $\tilde{i} = J \subset I$ . Such an orbit yields a character  $\gamma_J = \sum_{i \in J} \gamma_i$ . The parabolic subgroup  $P/\mathbb{Q}$  provided by this datum is determined by its root system  $\Delta^P = \{\beta \in \Delta \mid \langle \beta, \gamma_J \rangle \geq 0\}$ . The choice of the maximal torus  $T \subset P$  also provides a Levi subgroup  $M \subset P$  but actually it is better to consider  $M$  as the quotient  $P/U_P$ .

The set of simple roots of  $M^{(1)}$  is the subset  $\pi_M = \{\dots, \alpha_i, \dots\}_{i \in I_M}$ , where of course  $I_M = I \setminus J$ . We also consider the group  $G^{(1)} \cap M = M_1$ . It is a reductive group, it has  $T^{(1)}$  as its maximal torus. We apply our previous considerations to this group  $M_1$ . It has a non trivial central torus  $C_1/\mathbb{Q}$ . This torus has a simple description, we pick a root  $\alpha_i, i \in J$ , we know that  $J$  is an orbit under  $\text{Gal}(F/\mathbb{Q})$ . We have the subfield  $F_{\alpha_i} \subset F$  such that  $\text{Gal}(F/F_{\alpha_i})$  is the stabilizer of  $\alpha_i$ . Then it is clear that

$$C_1 \xrightarrow{\sim} R_{F_{\alpha_i}/\mathbb{Q}}(\mathbb{G}_m/F_{\alpha_i}),$$

up to isogeny it is a product of an anisotropic torus  $C_1^{(1)}/\mathbb{Q}$  and a copy of  $\mathbb{G}_m$ . The character module  $X_{\mathbb{Q}}^*(C_1)$  is a direct sum

$$X_{\mathbb{Q}}^*(C_1) = X_{\mathbb{Q}}^*(C_1^{(1)}) \oplus \mathbb{Q}\gamma_J. \tag{9.3}$$

Here  $X_{\mathbb{Q}}^*(C_1^{(1)}) = \{\gamma \in X_{\mathbb{Q}}^*(C_1) \mid \langle \gamma, \sum_{i \in J} \alpha_i \rangle = 0\}$ . The half sum of positive roots in the unipotent radical is

$$\rho_U = f_P \gamma_J \tag{9.4}$$

where  $2f_P > 0$  is an integer.

We also have the semi simple part  $T^{(1,M)} \subset M^{(1)}$  and again we get the orthogonal decomposition

$$X_{\mathbb{Q}}^*(T^{(1)}) = X_{\mathbb{Q}}^*(T^{(1,M)}) \oplus X_{\mathbb{Q}}^*(C_1) = \bigoplus_{i \in I_M} \mathbb{Q}\alpha_i \oplus \bigoplus_{i \in J} \mathbb{Q}\gamma_i = \bigoplus_{i \in I_M} \mathbb{Q}\gamma_i^M \oplus \bigoplus_{i \in J} \mathbb{Q}\gamma_i.$$

Here we have to observe that the  $\gamma_i^M, i \in I_M$  are the dominant fundamental weights for the group  $M^{(1)}$ , they are the orthogonal projections of the  $\gamma_i$  to the first summand in the above decomposition. We have a relation

$$\gamma_j = \gamma_j^M + \sum_{i \in \tilde{i}} c(j, i)\gamma_i, \text{ for } j \in I_M$$

and we have  $c(j, i) \geq 0$  for all  $i \in J$ .

Let  $W$  be absolute Weylgroup and subgroup  $W_M \subset W$  the Weyl group of  $M$ . For the quotient  $W_M \backslash W$  we have a canonical system of representatives

$$W^P = \{w \in W \mid w^{-1}(\pi_M) \subset \Delta^+\}.$$

To any  $w \in W$  we define  $w \cdot \lambda = w(\lambda + \rho) - \rho$  where  $\rho$  us the half sum of positive roots. If we do this with an element  $w \in W^P$  then  $\mu = w \cdot \lambda$  is a highest weight for  $M^{(1)}$  and  $w \cdot \lambda$  defines us a module for  $M$ . Then Kostants theorem says

$$H^\bullet(\mathfrak{u}_P, \mathcal{M}_\lambda) = \bigoplus_{w \in W^P} H^{\ell(w)}(\mathfrak{u}_P, \mathcal{M})(w \cdot \lambda),$$

the summands on the right hand side are the irreducible modules attached to  $w \cdot \lambda$ , they sit in degree

$$l(w) = \#\{\alpha \in \Delta^+ \mid w^{-1}\alpha \in \Delta^-\} \tag{9.5}$$

Each isomorphism class occurs only once.

We write

$$\begin{aligned} w \cdot \lambda &= \underbrace{\mu^{(1,M)} + \delta_1}_{\in X_{\mathbb{Q}}^*(T^{(1,M)}) \oplus X_{\mathbb{Q}}^*(C_1)} + \delta \tag{9.6} \\ &\in X_{\mathbb{Q}}^*(T^{(1,M)}) \oplus X_{\mathbb{Q}}^*(C_1) \oplus X^*(C) \end{aligned}$$

We decompose  $\delta_1$  and define the numbers  $a(w, \lambda)$  (see (9.3))

$$\delta_1 = \delta'_1 + a(w, \lambda)\gamma_J.$$

Then we get

$$w(\lambda + \rho) - \rho = \mu^{(1,M)} + a(w, \lambda)\gamma_J \tag{9.7}$$

We also consider the extended Weyl group  $\tilde{W}$ , this is the group of automorphisms of the root system. Let  $w_0 \in \tilde{W}$  be the element sending all positive roots into negative ones. We have an automorphism  $\Theta_- \in \tilde{W}$  inducing  $t \mapsto t^{-1}$  on the torus. Let  $\Theta = w_0 \circ \Theta_-$ . This element induces a permutation on the set  $\pi$  of positive roots, which may be the identity and induces  $-1$  on the determinant. Then

$$\Theta\lambda = \sum_{i \in I} a_{\Theta i} \gamma_i - \delta$$

is a dominant weight and the resulting highest weight module is dual module to  $\mathcal{M}_\lambda$ . Therefore we get a non degenerate pairing

$$H^\bullet(\mathfrak{u}_P, \mathcal{M}_\lambda) \times H^\bullet(\mathfrak{u}_P, \mathcal{M}_{\Theta\lambda}) \rightarrow H^{d_{U_P}}(\mathfrak{u}_P, F) = F(-2\rho_U),$$

which respects the decomposition, i.e. we get a bijection  $w \mapsto w'$  such that  $l(w) + l(w') = d_{U_P}$  and such

$$H^{l(w)}(\mathfrak{u}_P, \mathcal{M}_\lambda)(w \cdot \lambda) \times H^{l(w')}(\mathfrak{u}_P, \mathcal{M}_{\Theta\lambda})(w' \cdot \Theta\lambda) \rightarrow H^{d_{U_P}}(\mathfrak{u}_P, F) \quad (9.8)$$

is non degenerate. We conclude

$$a(w, \lambda) + a(w', \Theta\lambda) = -2f_P. \quad (9.9)$$

We say that  $w \cdot \lambda$  is in the positive chamber if

$$a(w, \lambda) \leq -f_P \quad (9.10)$$

The element  $\Theta$  conjugates the parabolic subgroup  $P$  into the parabolic subgroup  $Q$ , which may be equal to  $P$  or not. If  $P = Q$  resp.  $P \neq Q$  then we say that  $P$  is (resp. not) conjugate to its opposite parabolic. If  $\Theta_-$  is in the Weyl group then all parabolic subgroups are conjugate to their opposite. In this case we have  $\Theta = 1$ .

Conjugating by the element  $\Theta$  provides an identification  $\theta_{P,Q} : W^P \xrightarrow{\sim} W^Q$ . We have two specific Kostant representatives, namely the identity  $e \in W^P$  and the element  $w_P \in W^P$ , this is the element which sends all the roots in  $U_P$  to negative roots (the longest element). Its length  $l(w_P)$  is equal to the dimension  $d_P = \dim(U_P)$ .

Any element in  $w \in W^P$  can be written as product of reflections

$$w = s_{\alpha_{i_1}} \dots s_{\alpha_{i_\nu}} \quad (9.11)$$

where  $\nu = l(w)$  and the first factor  $\alpha_{i_1} \in J$ . We always can complement this product to a product giving the longest element

$$s_{\alpha_{i_1}} \dots s_{\alpha_{i_\nu}} s_{\alpha_{i_{\nu+1}}} \dots s_{\alpha_{i_{d_P}}} = w s_{\alpha_{i_{\nu+1}}} \dots s_{\alpha_{i_{d_P}}} = w_P, \quad (9.12)$$

The inverse of the element  $s_{\alpha_{i_{\nu+1}}} \dots s_{\alpha_{i_{d_P}}}$  is

$$w' = s_{\alpha_{i_{d_P}}} \dots s_{\alpha_{i_{\nu+1}}} \in W^Q$$

This defines a second bijection  $i_{P,Q} : W^P \xrightarrow{\sim} W^Q$  which is defined by the relation

$$w = w_P \cdot i_{P,Q}(w) = w_P \cdot w', \quad l(w) + l(w') = d_P \quad (9.13)$$

The composition  $\theta_{P,Q}^{-1} \circ i_{P,Q} : W^P \rightarrow W^P$  is the bijection provided by duality.

The element  $w_P$  conjugates the Levi subgroup  $M$  of  $P$  into the Levi subgroup of  $Q = w_P P w_P^{-1}$ . The element  $\tilde{w}_P = \Theta w_P$  conjugates the parabolic subgroup  $P$  into its opposite (which is conjugate to  $Q$ ) and induces an automorphism on the subgroup  $M$  which is a common Levi-subgroup of  $P$  and its opposite.

If we choose  $w = e$  then

$$\sum_{i \in I} a_i \gamma_i + \delta = \sum_{i \in I_M} a_i \gamma_i^M + \sum_{j \in J} \left( \sum_{i \in I_M} a_i c(i, j) + n_j \right) \gamma_j + \delta.$$

Since  $J$  is the orbit of an element  $i \in I$  we see that  $\langle \gamma_j, \alpha_j \rangle$  is independent of  $j$  and hence we get easily

$$\sum_{j \in J} \left( \sum_{i \in I_M} a_i c(i, j) + n_j \right) \gamma_j = \frac{1}{\#J} \left( \sum_{j \in J} \left( \sum_{i \in I_M} a_i c(i, j) + n_j \right) \right) \gamma_J + \delta'$$

and hence

$$a(e, \lambda) = \frac{1}{\#J} \left( \sum_{j \in J} \left( \sum_{i \in I_M} a_i c(i, j) + a_j \right) \right)$$

If we choose  $\Theta_P$  then as an  $M$ -module  $\mathcal{M}_{\Theta_P \cdot \lambda}$  is dual to  $\mathcal{M}_{\Theta \lambda}(-2f_J \gamma_J)$ . We write  $\Theta \lambda + \rho = \sum_{i \in I} a_{\Theta i} \gamma_i - \delta$  and then

$$w_P \left( \sum_{i \in I} a_i \gamma_i + \delta \right) = \sum_{i \in I_M} n_{\Theta i} \gamma_i^M - \sum_{j \in J} \left( \sum_{\Theta i \in I_M} a_{\Theta i} c(\Theta i, \Theta j) + a_{\Theta j} \right) \gamma_j - 2f_J \gamma_J - \delta.$$

and especially we find

$$a(w_P, \lambda) = - \left( \frac{1}{\#J} \left( \sum_{j \in J} \left( \sum_{i \in I_M} a_{\Theta i} c(\Theta i, \Theta j) + a_{\Theta j} \right) \right) + 2f_J \right) \gamma_J$$

In general we have the inequalities

$$a(\Theta_P, \lambda) \leq a(w, \lambda) \leq a(e, \lambda).$$

We can write our relation (9.7) slightly differently. We can move the half sum of positive roots to the right and split into  $\rho = \rho^M + f_P \gamma_J$ . We put  $\tilde{\mu}^{(1)} = \mu^{(1, M)} + \rho^M$  and then we write

$$w(\lambda + \rho) = \tilde{\mu}^{(1)} + (a(w, \lambda) + f_P) \gamma_J = \tilde{\mu}^{(1)} + b(w, \lambda) \gamma_J \quad (9.14)$$

and of course now we have

$$b(w, \lambda) + b(w', \Theta \lambda) = 0. \quad (9.15)$$

#### 9.1.4 The inverse problem

Later we will encounter the following problem. Our data are as above and we start from a highest weight for  $M$ , we write

$$\mu = \mu^{(1)} + \delta_1 + a \gamma_J + \delta = \sum_{i \in I_M} n_{\Theta i} \gamma_i^M + \delta_1 + a \gamma_J + \delta.$$

We ask whether we can find a  $\lambda$  such that we can solve the equation (*Kost*). More precisely: We give ourselves only the semi simple component  $\mu^{(1)}$  of  $\mu$  and we ask for the solutions

$$w(\lambda + \rho) = \tilde{\mu}^{(1)} + \dots$$

where  $w \in W^P$  and  $\lambda$  dominant, i.e. we only care for the semi simple component.

Let us consider the case where  $J = \{i_0\}$ , i.e. it is just one simple root. Then the term  $\delta_1$  disappears and our equation becomes

$$w(\lambda + \rho) = \tilde{\mu}^{(1)} + b\gamma_{i_0} + \delta,$$

of course the  $\delta$  is irrelevant, but we want to know the range of the values  $b = b(\lambda, w)$  when  $\tilde{\mu}^{(1)}$  is fixed, but  $\lambda, w$  vary. Of course it may be empty. Let us fix a  $w$  and let us assume we have solved  $w(\lambda + \rho) = \tilde{\mu}^{(1)} + \dots$ . Then it is clear that the other solutions are of the form  $\lambda + \rho + \nu$  where  $w\nu \in \mathbb{Q}\gamma_{i_0}$ . These  $\nu$  are of the form  $\nu = c\nu_0$  with  $c \in \mathbb{Z}$ . We write  $\nu_0 = \sum_{i \in I} b_i \gamma_i$  and it is easy to see that there must be some  $b_i > 0$  and some  $b_j < 0$ . This implies that  $\lambda + c\nu_0$  is dominant if and only if  $c \in [M, N]$ , an interval with integers as boundary point. This of course implies that -still for a given  $w$  - the values  $b = b(\lambda, w)$  also have to lie in a fixed finite interval

$$b = b(w, \lambda) \in [b_{\min}(w, \tilde{\mu}^{(1)}), a_{\max}(w, \tilde{\mu}^{(1)})] = I(w, \tilde{\mu}^{(1)}).$$

This will be of importance because these intervals will be related to intervals of critical values of  $L$ -functions.

## 9.2 The goal of Eisenstein cohomology

The goal of the Eisenstein cohomology is to provide an understanding of the restriction map  $r$  in theorem ( 6.2.1). More precisely we assume that we understand (can describe) the cohomology  $H^\bullet(\partial\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$  then we want to understand the image  $H_{\text{Eis}}^\bullet(\partial\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$  in terms of this description. Under certain conditions we will construct a section  $\text{Eis} : H_{\text{Eis}}^i(\partial\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda, \mathbb{C}}) \rightarrow H^i(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda, \mathbb{C}})$ . It is clear from the previous considerations that understanding of  $H^\bullet(\partial\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$  requires understanding cohomology of  $H^\bullet(\mathcal{S}_{K_f^M}^M, \widetilde{H^\bullet(\mathfrak{u}, \mathcal{M})})$  and we have to compute the differentials in the spectral sequence. These differentials will depend on the Eisenstein cohomology of  $H^\bullet(\mathcal{S}_{K_f^M}^M, \widetilde{H^\bullet(\mathfrak{u}, \mathcal{M})})$ . Under certain conditions the spectral sequence degenerates at  $E_2$  level and I do not know whether this is true in general. In a certain sense it would be much more interesting if this is not the case.

We consider certain submodules in the cohomology of the Borel-Serre compactification for which we can construct a section as above. We start from a maximal parabolic subgroup  $P/\mathbb{Q}$ , let  $M/\mathbb{Q}$  be its reductive quotient. We define

$$H_!^\bullet(\partial_P \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) = \bigoplus_{w \in W^P} H_!^{\bullet-l(w)}(\mathcal{S}_{K_f^M}^M, H^{l(w)}(\mathfrak{u}_P, \tilde{\mathcal{M}})(w \cdot \lambda)) \subset H^\bullet(\partial_P \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \tag{9.16}$$

We will abbreviate  $H^{l(w)}(\mathfrak{u}_P, \tilde{\mathcal{M}})(w \cdot \lambda) = \tilde{\mathcal{M}}(w \cdot \lambda)$  where always keep in mind that the element  $w \in W^P$  knows what the actual parabolic subgroup is and that  $\tilde{\mathcal{M}}(w \cdot \lambda)$  sits in degree  $l(w)$ .

By definition the inner cohomology is the image of the cohomology with compact supports. This implies that the submodule

$$\bigoplus_{P:d(P)=1} H_!^q(\partial_P \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \subset \bigoplus_{P:d(P)=1} H^q(\partial_P \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) = E_1^{0,q}$$

is annihilated by all differentials  $d_{\nu'}^{0,q}$  and hence we get an inclusion

$$i_P : \bigoplus_{w \in W^P} I_P^G H_!^{\bullet-l(w)}(\mathcal{S}_{K_f}^M, \mathcal{M}(w \cdot \lambda)) \rightarrow H^\bullet(\partial \mathcal{S}_{K_f}^G, \mathcal{M}_\lambda) \quad (9.17)$$

Taking the direct sum over the maximal parabolic subgroups yields a submodule

$$H_!^\bullet(\partial \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \hookrightarrow H^\bullet(\partial \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \quad (9.18)$$

The Hecke algebra acts on these two modules. Let us assume that this submodule when tensorized by  $\mathbb{Q}$  is isotypical in  $H_!^\bullet(\partial \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{Q})$ . Then we get a decomposition

$$H_!^\bullet(\partial \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{Q}) \oplus H_{\text{non!}}^\bullet(\partial \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{Q}) = H^\bullet(\partial \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{Q}). \quad (9.19)$$

We formulated the goal of the Eisenstein cohomology, we described an isotypical subspace and we know can ask: What is the intersection of  $H_{\text{Eis}}^\bullet(\partial \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{Q})$  with this subspace, or what amounts to the same, what is  $H_{!, \text{Eis}}^\bullet(\partial \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{Q})$ .

The element  $\Theta$  induces an involution on the set of parabolic subgroups containing  $B$  ( $=$  set of  $G(\mathbb{Q})$  conjugacy classes of parabolic subgroups) two parabolic subgroups  $P, Q \supset B$  are called associate if  $\Theta P = Q$ . We can decompose the cohomology  $H_!^\bullet(\partial \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{Q})$  into summands attached to the classes of associated parabolic subgroups

$$H_!^\bullet(\partial \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{Q}) = \bigoplus_{P:P=\Theta P} H_!^\bullet(\partial_P \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \oplus \bigoplus_{[P,Q]} H_!^\bullet(\partial_P \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \oplus H_!^\bullet(\partial_Q \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \quad (9.20)$$

where in the second sum  $Q = \Theta P$ . Each summand is a sum over the elements of  $W^P$  and then we can decompose under the action of the Hecke algebra. We choose a sufficiently large extension  $F/\mathbb{Q}$  and in the case  $P = \Theta P$  we get

$$H_!^\bullet(\partial_P \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F) = \bigoplus_{w \in W^P} \bigoplus_{\sigma_f} H_!^{\bullet-l(w)}(\mathcal{S}_{K_f}^M, \tilde{\mathcal{M}}(w \cdot \lambda) \otimes F)(\sigma_f) \quad (9.21)$$

In the case  $P \neq \Theta P = Q$  we group the contributions from the two parabolic subgroups together. To any  $w \in W^P$  we have the element  $i_{P,Q}(w) = w' \in W^Q$ . We also group the terms corresponding to  $w$  and  $w'$  together. To any  $\sigma_f$  which occurs in  $H_!^{\bullet-l(w)}(\mathcal{S}_{K_f}^M, H^{l(w)}(\mathfrak{u}_P, \tilde{\mathcal{M}})(w \cdot \lambda) \otimes F)$  we find a  $\sigma'_f = \sigma_f^{w_P} | \gamma_{\Theta j} |_f^{2f_Q}$ , which occurs in the second summand.

The decomposition into isotypical pieces becomes

$$\bigoplus_{\sigma_f} (H_!^{\bullet-l(w)}(\mathcal{S}_{K_f}^M, \tilde{\mathcal{M}}(w \cdot \lambda) \otimes F)(\sigma_f) \oplus H_!^{\bullet-l(w')}(\mathcal{S}_{K_f}^{M'}, \tilde{\mathcal{M}}(w' \cdot \lambda) \otimes F)(\sigma'_f)) \quad (9.22)$$

We can define the second step in the filtration ( 6.20) as the inverse image of  $H_!^\bullet(\partial \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$  under the restriction  $r$ .

### 9.2.1 Induction and the local intertwining operator at finite places

Our modules  $\sigma_f$  are modules for the Hecke algebras  $\mathcal{H}_{K_f^M}^M = \otimes_p \mathcal{H}_{K_p^M}^M$ . Therefore we can write them as tensor product  $\sigma_f = \otimes_p \sigma_p$ . We consider a prime  $p$  where  $\sigma_f$  is unramified then we get can give a standard model for this isomorphism class. The module  $H_{\sigma_p}$  is the rank one  $\mathcal{O}_F$ -module  $\mathcal{O}_F$ , i.e. it comes with a distinguished generator 1. The Hecke algebra acts by a homomorphism (See 6.3.2)

$$h(\sigma_p) : \mathcal{H}_{K_p^M, \mathbb{Z}}^{(M, w \cdot \lambda)} \rightarrow \mathcal{O}_F \tag{9.23}$$

and gives us the Hecke-module structure on  $H_{\sigma_p}$ . We can induce  $H_{\sigma_p}$  to a  $\mathcal{H}_{K_p^G}^G$  module. This is actually the same  $\mathcal{O}_F$  module but now with an action of the algebra  $\mathcal{H}_{K_p^G, \mathbb{Z}}^{(G, \lambda)}$ . We simply observe that we have an inclusion  $\mathcal{H}_{K_p^G, \mathbb{Z}}^{(G, \lambda)} \hookrightarrow \mathcal{H}_{K_p^M, \mathbb{Z}}^{(M, w' \cdot \lambda)}$  and induction simply means restriction.

It follows easily from the description of the description of the spherical (unramified) Hecke modules via their Satake-parameters that the induced modules  $H_{\sigma_p}$  and  $H_{\sigma'_p}$  are isomorphic as  $\mathcal{H}_{K_p^G, \mathbb{Z}}^{(G, \lambda)}$ -modules and hence we get that after induction the two summands in (9.22) become isomorphic. We choose a local intertwining operator

$$T_p^{\text{loc}} : H_{\sigma_p} \rightarrow H_{\sigma'_p} \tag{9.24}$$

simply the identity.

We postpone the discussion of a local intertwining operator at ramified places.

## 9.3 The Eisenstein intertwining operator

We start from an irreducible unitary module  $H_{\sigma_\infty} \times H_{\sigma_f} = H_\sigma$  and assume that we have an inclusion  $\Phi : H_\sigma \hookrightarrow L_{\text{cusp}}^2(M(\mathbb{Q}) \backslash M(\mathbb{A}))$ . We assume that  $\sigma_f$  occurs in the cohomology  $H_{\text{cusp}}^\bullet(\mathcal{S}_{K_f^M}^M, \tilde{\mathcal{M}}(w \cdot \lambda)_{\mathbb{C}})$  and we assume that  $w \cdot \lambda$  is in the positive chamber. We consider  $\Phi$  as an element of  $W(\sigma)$  and for the moment we identify  $H_\sigma$  to its image under  $\Phi$ . We stick to our assumption that  $\sigma$  occurs with multiplicity one in the cuspidal spectrum.

Then we we can consider the induced module, recall that this is the space of functions

$$\{f : G(\mathbb{A}) \rightarrow H_\sigma \mid f(\underline{p}g) = \bar{p}f(g)\} \tag{Ind}$$

where  $\bar{p}$  is the image of  $\underline{p}$  in  $M(\mathbb{A})$ . We can define the subspace  $H_\sigma^{(\infty)}$  consisting of those  $f$  which satisfy some suitable smoothness conditions and then we can define a submodule  $\text{Ind}_{P(\mathbb{A})}^{G(\mathbb{A})} H_\sigma^{(\infty)}$  where the  $f(g) \in H_\sigma^{(\infty)}$  and the  $f$  themselves also satisfy some smoothness conditions.

We embed this space into the space  $\mathcal{A}(P(\mathbb{Q}) \backslash G(\mathbb{A}))$  by sending

$$f \mapsto \{g \mapsto f(g)(e_M)\},$$

here  $\mathcal{A}$  denotes some space of automorphic forms. This an embedding of  $G(\mathbb{A})$ -modules or an embedding of Hecke modules if we fix a level.

We have the character  $\gamma_P : M \rightarrow G_m$ , for any complex number  $z$  this yields a homomorphism  $|\gamma_P|^z : M(\mathbb{A}) \rightarrow \mathbb{R}^\times$  which is given by  $|\gamma_P| : \underline{m} \mapsto |\gamma_P(\underline{m})|^z$ . As usual we denote by  $\mathbb{C}(|\gamma_P|^z)$  the one dimensional  $\mathbb{C}$  vector space on which  $M(\mathbb{A})$  acts by the character  $|\gamma_P|^z$ . Then we may twist the representation  $H_\sigma$  by this character and put  $H_\sigma \otimes |\gamma_P|^z = H \otimes \mathbb{C}(|\gamma_P|^z)$ . An element  $\underline{g} \in G(\mathbb{A})$  can be written as  $\underline{g} = \underline{p}\underline{k}, \underline{p} \in P(\mathbb{A}), \underline{k} \in K_f^0$  where  $K_f^0 \supset K_f$  is a suitable maximal compact subgroup and now we define  $h(\underline{g}) = |\gamma_P|(\underline{p})$ .

Eisenstein summation yields embeddings

$$\text{Eis} : \text{Ind}_{P(\mathbb{A})}^{G(\mathbb{A})} H_\sigma^{(\infty)} \otimes |\gamma_P|^z \rightarrow \mathcal{A}(G(\mathbb{Q}) \backslash G(\mathbb{A})), \tag{9.25}$$

where

$$\text{Eis}(f)(\underline{g}) = \sum_{\gamma \in P(\mathbb{Q}) \backslash G(\mathbb{Q})} f(\gamma \underline{g})(e_M) h(\gamma \underline{g})^z,$$

it is well known that this is locally uniformly convergent provided  $\Re(z) \gg 0$  and it has meromorphic continuation into the entire  $z$  plane (See [Ha-Ch]).

We assumed that  $H_\sigma$  is in the cuspidal spectrum. We get important information concerning these Eisenstein series, if we compute their constant Fourier coefficient with respect to parabolic subgroups: For any parabolic subgroup  $P_1/\mathbb{Q} \subset G/\mathbb{Q}$  with unipotent radical  $U_1 \subset P_1$  we define (See [Ha-Ch], 4)

$$\mathcal{F}^{P_1}(\text{Eis}(f))(\underline{g}) = \int_{U_1(\mathbb{Q}) \backslash U_1(\mathbb{A})} \text{Eis}(f)(\underline{u}\underline{g})(e_M) d\underline{u}.$$

This essentially only depends on the  $G(\mathbb{Q})$ -conjugacy class of  $P_1/\mathbb{Q}$ . It is also in [Ha-Ch], 4 that this constant term is zero unless  $P_1$  is maximal and the conjugacy class of  $P_1$  is equal to the conjugacy class of  $P/\mathbb{Q}$  or the conjugacy class of  $Q/\mathbb{Q}$ . (which may or may not be equal to the conjugacy class of  $P/\mathbb{Q}$ .)

These constant Fourier coefficients have been computed by Langlands, we have to distinguish the two cases:

a) The parabolic subgroup  $P/\mathbb{Q}$  is conjugate to an opposite parabolic  $Q/\mathbb{Q}$ .

In this case we have a Kostant representative  $w^P \in W^P$  which conjugates  $Q/\mathbb{Q}$  into  $P/\mathbb{Q}$  and it induces an automorphism of  $M/\mathbb{Q}$ . We get a twisted representation  $w^P(\sigma)$  of  $M(\mathbb{A})$ . In the computation of the the constant term we have to exploit that  $\sigma$  is cuspidal and we get two terms:

$$\begin{aligned} \mathcal{F}^P \circ \text{Eis} : \text{Ind}_{P(\mathbb{A})}^{G(\mathbb{A})} H_\sigma \otimes |\gamma_P|^z &\rightarrow \\ \text{Ind}_{P(\mathbb{A})}^{G(\mathbb{A})} H_\sigma \otimes |\gamma_P|^z \oplus \text{Ind}_{P(\mathbb{A})}^{G(\mathbb{A})} H_{w^P(\sigma)} \otimes |\gamma_Q|^{2f_P-z} &\subset \mathcal{A}(P(\mathbb{Q}) \backslash G(\mathbb{A})). \end{aligned} \tag{9.26}$$

We can describe the image. It is well known, that we can define a holomorphic family

$$T^{\text{loc}}(z) : \text{Ind}_{P(\mathbb{A})}^{G(\mathbb{A})} H_\sigma \otimes |\gamma_P|^z \rightarrow \text{Ind}_{P(\mathbb{A})}^{G(\mathbb{A})} H_{\sigma w^P} \otimes |\gamma_Q|^{2f_P-z}$$

which is defined in a neighborhood of  $z = 0$  and which is nowhere zero. This local intertwining operator is unique up to a nowhere vanishing holomorphic function  $h(z)$ . It is the tensor product over all places  $T^{\text{loc}}(z) = \otimes_v T_v^{\text{loc}}(z)$ .

For the unramified finite places the local operator is constant, i.e. does not depend on  $z$  and is equal to  $T_p^{\text{loc}}$  in section (9.2.1) and  $T^{\text{loc}}(0) = \otimes_p T_p^{\text{loc}}$ . At the remaining factors there is a certain arbitrariness for the choice of the local operator and some fine tuning is appropriate.

We also assume that we have chosen nice model spaces  $H_{\sigma_\infty}, H_{\sigma'_\infty}$ , and an intertwining operator

$$T_\infty^{\text{loc}} : H_{\sigma_\infty} \rightarrow H_{\sigma'_\infty} \tag{9.27}$$

which is normalized by the requirement that it induces the "identity" on a certain fixed  $K_\infty^M$  type.

Then we get the classical formula of Langlands for the constant term: For  $f \in \text{Ind}_{P(\mathbb{A})}^{G(\mathbb{A})} H_\sigma \otimes |\gamma_P|^z$  we get

$$\mathcal{F}^P \circ \text{Eis}(f) = f + C(\sigma, z) T^{\text{loc}}(z)(f), \tag{9.28}$$

where  $C(\sigma, \lambda, z)$  is a product of local factors  $C(\sigma_v, z)$  and where  $C(\sigma_v, z)$  is a function in  $z$  which is holomorphic for  $\Re(z) \geq 0$  (here we need that  $w \cdot \lambda$  is in the positive chamber.) This function compares our local intertwining operator to an intertwining operator which is defined by the integral.

The computation of this factor is carried out in H. Kims paper in [C-K-M], chap. 6. He expresses the factor in terms of the automorphic  $L$  function attached to  $\sigma_f$ . To formulate the result of this computation we have to recall the notion of the dual group (7.0.1). Inside the dual group  ${}^L G$  we have the dual group  ${}^L M$  which acts by conjugation on the Lie algebra  $\mathfrak{u}_P^\vee$ . The set of roots  $\Delta_{U_P}^+$  is a set of cocharacters of  $T/\mathbb{Q}$ , a coroot  $\alpha^\vee \in \Delta_{U_P}^+$  defines a one-dimensional root subgroup  $\mathfrak{u}_{P, \alpha^\vee}^\vee$ . The  ${}^L M$ -module  $\mathfrak{u}_P^\vee$  decomposes into submodules. We recall that the maximal parabolic subgroup  $P/\mathbb{Q}$  was obtained from the choice of a Galois-orbit  $\tilde{i} \subset I$  (9.1.3) and any

$$\alpha^\vee \in \Delta_{U_P}^+, \chi = a(\alpha^\vee, \tilde{i}) \chi_{\tilde{i}} + \sum_{j \notin \tilde{i}} m_{\tilde{i}, j} \chi_j. \tag{9.29}$$

Here the coefficients are integers  $\geq 0$  and  $a(\alpha^\vee, \tilde{i}) > 0$ . For a given integer  $a > 0$  we define

$$\mathfrak{u}_P^\vee[a] = \bigoplus_{\alpha^\vee : a(\alpha^\vee, \tilde{i}) = a} \mathfrak{u}_{P, \alpha^\vee}^\vee \tag{9.30}$$

it is rather obvious that  $\mathfrak{u}_P^\vee[a]$  is an invariant submodule under the action of  $M$  and actually it is even irreducible. Let us denote the representation of  $M/\mathbb{Q}$  on  $\mathfrak{u}_P^\vee[a]$  by  $r_a^{\mathfrak{u}_P^\vee}$ . In the following  $\eta_a$  will be the highest weight of  $r_a^{\mathfrak{u}_P^\vee}$ .

With these notations we get the following formula for the local factor at  $p$  (See[Kim])

$$C_p(\sigma, z) = \prod_{a=1}^r \frac{L^{\text{aut}}(\sigma_p, r_a^{\mathfrak{u}_P^\vee}, a(z - f_P))}{L^{\text{aut}}(\sigma_p, r_a^{\mathfrak{u}_P^\vee}, a(z - f_P) + 1)} T_p^{\text{loc}}(z)(f) \tag{9.31}$$

We do not discuss the ramified finite places, from now on we assume that  $\sigma_f$  is unramified. Then we get

$$C(\sigma, z) = C(\sigma_\infty, z) \prod_p C_p(\sigma_p, z) = C(\sigma_\infty, z) \prod_{a=1}^r \frac{L^{\text{aut}}(\sigma_f, r_a^{\text{u}_P^\vee}, a(z - f_P))}{L^{\text{aut}}(\sigma_f, r_a^{\text{u}_P^\vee}, a(z - f_P) + 1)}$$

The local factor at infinity depends on the choice of  $T_\infty^{\text{loc}}$ , in 1.2.4 we gave some rules how to fix it, if it is not zero on cohomology.

b) The opposite group  $Q/\mathbb{Q}$  is not conjugate to  $P/\mathbb{Q}$ , then we have to compute two Fourier coefficients namely  $\mathcal{F}^P$  and  $\mathcal{F}^Q$  in this case we get

$$\mathcal{F} : \text{Ind}_{P(\mathbb{A})}^{G(\mathbb{A})} H_\sigma \otimes |\gamma_P|^z \xrightarrow{\mathcal{F}^P \oplus \mathcal{F}^Q}$$

$$\text{Ind}_{P(\mathbb{A})}^{G(\mathbb{A})} H_\sigma \otimes |\gamma_P|^z \oplus \text{Ind}_{Q(\mathbb{A})}^{G(\mathbb{A})} H_\sigma \otimes |\gamma_Q|^{2f_P - z} \subset \mathcal{A}(P(\mathbb{Q}) \backslash G(\mathbb{A})) \oplus \mathcal{A}(Q(\mathbb{Q}) \backslash G(\mathbb{A})).$$

and again we get

$$\mathcal{F} \circ \text{Eis}(f) = f + C(\sigma_\infty, z) \prod_a \frac{L^{\text{aut}}(\sigma_f, r_a^{\text{u}_P^\vee}, a(z - f_P))}{L^{\text{aut}}(\sigma_f, r_a^{\text{u}_P^\vee}, a(z - f_P) + 1)} T^{\text{loc}}(z)(f), \quad (9.32)$$

where now  $T^{\text{loc}}(z)$  is a product of local intertwining operators

$$T_v^{\text{loc}} : \text{Ind}_{P(\mathbb{Q}_v)}^{G(\mathbb{Q}_v)} H_{\sigma_v} \otimes |\gamma_P|^z \rightarrow \text{Ind}_{Q(\mathbb{Q}_v)}^{G(\mathbb{Q}_v)} H_{\sigma_{w^P}} \otimes (2f_P - z).$$

It is also due to Langlands that the Eisenstein intertwining operator is holomorphic at  $z = 0$  if the factor in front of the second term is holomorphic at  $z = 0$ . Up to here  $\sigma$  can be any representation occurring in the cuspidal spectrum of  $M$ .

Now we assume that we have a coefficient system  $\mathcal{M} = \mathcal{M}_\lambda$  and a  $w \in W^P$  such that our  $\sigma_f$  occurs in  $H_1^{\bullet - l(w)}(\mathcal{S}_{K_f^M}^M, \tilde{\mathcal{M}}(w \cdot \lambda) \otimes F)$ . Then we find a  $(\mathfrak{m}, K_\infty^M)$ -module  $H_{\sigma_\infty}$  such that  $H^\bullet(\mathfrak{m}, K_\infty^M, H_{\sigma_\infty} \otimes \mathcal{M}(w \cdot \lambda)) \neq 0$ . We also find an embedding

$$\Phi_\iota : H_{\sigma_\infty} \otimes H_{\sigma_f} \otimes_{F, \iota} \mathbb{C} \hookrightarrow L_{\text{cusp}}^2(M(\mathbb{Q}) \backslash M(\mathbb{A})) \quad (9.33)$$

Let us assume that  $w \cdot \lambda$  or equivalently  $\sigma_f$  are in the positive chamber. In case a) we have holomorphicity at  $z = 0$  if the weight  $\lambda$  is regular (See [Schw]) and in case b) the Eisenstein series is always holomorphic at  $z = 0$ . In this section that we assume that the Eisenstein series is holomorphic at  $z = 0$  and hence we can evaluate at  $z = 0$  in (9.179) and get an intertwining operator

$$\text{Eis} \circ \Phi_\iota : \text{Ind}_{P(\mathbb{A})}^{G(\mathbb{A})} H_\sigma \rightarrow \mathcal{A}(G(\mathbb{Q}) \backslash G(\mathbb{A})). \quad (9.34)$$

We get a homomorphism on the de-Rham complexes

$$\text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \text{Ind}_{P(\mathbb{A})}^{G(\mathbb{A})} H_\sigma \otimes_{F, \iota} \mathbb{C} \otimes \mathcal{M}_\lambda) \rightarrow \text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \mathcal{A}(G(\mathbb{Q}) \backslash G(\mathbb{A})) \otimes \tilde{\mathcal{M}}_\lambda) \quad (9.35)$$

We introduce the abbreviation  $H_{\iota\circ\sigma_f} = H_{\sigma_f} \otimes_{F,\iota} \mathbb{C}$  and decompose  $H_{\iota\circ\sigma} = H_{\sigma_\infty} \otimes H_{\iota\circ\sigma_f}$ . We compose (9.35) with the constant term and get

$$\begin{aligned} \mathcal{F} \circ \text{Eis}^\bullet : \text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \text{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})} H_{\sigma_\infty} \otimes \mathcal{M}_\lambda) \otimes H_{\iota\circ\sigma_f} \rightarrow \\ \text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \text{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})} H_{\sigma_\infty} \otimes \mathcal{M}_\lambda) \otimes H_{\iota\circ\sigma_f} \oplus \text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \text{Ind}_{Q(\mathbb{R})}^{G(\mathbb{R})} H_{\sigma'_\infty} \otimes \mathcal{M}_\lambda) \otimes H_{\iota\circ\sigma'_f} \end{aligned} \tag{9.36}$$

where  $P = Q$  in case a).

We choose an  $\omega \in \text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \text{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})} H_{\sigma_\infty} \otimes \mathcal{M}_\lambda)$  and consider classes  $\omega \otimes \psi_f$  and map them by the Eisenstein intertwining operator to the cohomology (or the de-Rham complex) on  $\mathcal{S}_{K_f}^G$ . Then the restriction of of the Eisenstein cohomology to the boundary is given by the classes

$$\Phi_\iota(\omega \otimes \psi_f + \frac{1}{\Omega(\sigma_f)} C(\sigma_\infty, \lambda) C(\sigma_f, \lambda) T_\infty^{\text{loc}}(\omega) \otimes T_f^{\text{loc}}(\psi_f)) \tag{9.37}$$

Here the factor  $C(\sigma_f, \lambda)$  can be expressed in terms of the cohomological  $L$ -function. Translating the formula (9.31) yields (see 9.14)

$$C(\sigma_f, \lambda) = \prod_a \frac{L^{\text{coh}}(\sigma_f, r_a^{\text{u}_P^\vee}, \langle \eta_a, \tilde{\mu}^{(1)} \rangle - b(w, \lambda) \langle \eta_a, \gamma_P \rangle)}{L^{\text{coh}}(\sigma_f, r_a^{\text{u}_P^\vee}, \langle \eta_a, \tilde{\mu}^{(1)} \rangle - b(w, \lambda) \langle \eta_a, \gamma_P \rangle + 1)} \tag{9.38}$$

We may complete the cohomological  $L$ -function by the correct factor at infinity and replace the ratio of  $L$ -values by the corresponding ratio of values for the completed  $L$ -function. By definition we have  $\langle \eta_a, \gamma_P \rangle = a$  and then our formula for the second term in (9.37) becomes

$$\frac{1}{\Omega(\sigma_f)} \prod_a \frac{\Lambda^{\text{coh}}(\sigma_f, r_a^{\text{u}_P^\vee}, \langle \eta_a, \tilde{\mu}^{(1)} \rangle - ab(w, \lambda))}{\Lambda^{\text{coh}}(\sigma_f, r_a^{\text{u}_P^\vee}, \langle \eta_a, \tilde{\mu}^{(1)} \rangle - ab(w, \lambda) + 1)} C^*(\sigma_\infty, \lambda) T_\infty^{\text{loc}}(\omega) \otimes T_f^{\text{loc}}(\psi_f) \tag{9.39}$$

This formula needs some comments. The factor  $C^*(\sigma_\infty) T_\infty^{\text{loc}}$  is a representation theoretic contribution it is not easy to understand. Experience shows that becomes very simple at the end. In SecOps.pdf we discuss the special case of the symplectic group.

The number  $\Omega(\sigma_f)$  is a period, it will be discussed later.

We see that the constant term is the sum of two terms. The first term reproduces the original class from which we started. We assumed that  $w$  or  $w \cdot \lambda$  it is in the positive chamber (see(9.10)). The second term is some kind of scattering term which is the image of the first term under an intertwining operator. In case a) the restriction of the second term gives a class in the same stratum, in case b) the restriction of the second term gives a class in a second stratum.

At this point I formulate a general principle

**Under certain circumstances the second term is of fundamental arithmetic interest, it contains relevant arithmetic information.**

To exploit this information we have to understand several aspects of the behavior of this second term in the constant term. We have to recall that is obtained as the evaluation of a meromorphic function  $C(\sigma_f, \lambda, z)$  at  $z = 0$ , i.e. we have to know whether it has pole at  $z = 0$  or not. We also have to understand the contribution  $C(\sigma_\infty, \lambda)T_\infty^{\text{loc}}$ , and we have to understand the arithmetic nature of this term, it is a product and some of the factors yield an algebraic number and the rest will have a motivic interpretation. This is explained further down and in [Mix-Mot-2013.pdf].

We give some more detailed indications how such arithmetic applications may look like. We assume that  $w \cdot \lambda$  is in the positive chamber and  $l(w) \geq l(w')$ . Let us also assume that the Eisenstein intertwining operator is holomorphic at  $z = 0$ . Then we have to look at

$$T_\infty^{\text{loc}, \bullet} : \text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \text{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})} H_{\sigma_\infty} \otimes \mathcal{M}_\lambda) \rightarrow \text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \text{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})} H_{\sigma'_\infty} \otimes \mathcal{M}_\lambda) \tag{9.40}$$

The two complexes can be described by the Delorme isomorphism

$$\text{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \text{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})} H_{\sigma_\infty} \otimes \mathcal{M}_\lambda) \xrightarrow{\sim} \bigoplus_{w \in W^P} \text{Hom}_{K_\infty^M}(\Lambda^{\bullet-l(w)}(\mathfrak{m}_\mathbb{C}^{(1)}/\mathfrak{k}^M), H_{\sigma_\infty} \otimes \mathcal{M}(w \cdot \lambda)) \tag{9.41}$$

Our intertwining operator respects this decomposition and we get

$$T_\infty^{\text{loc}, \bullet}(w) : \text{Hom}_{K_\infty^M}(\Lambda^{\bullet-l(w)}(\mathfrak{m}_\mathbb{C}^{(1)}/\mathfrak{k}^M), H_{\sigma_\infty} \otimes \mathcal{M}(w \cdot \lambda)) \rightarrow \text{Hom}_{K_\infty^M}(\Lambda^{\bullet-l(w')}(\mathfrak{m}_\mathbb{C}^{(1)}/\mathfrak{k}^M), H_{\sigma'_\infty} \otimes \mathcal{M}(w' \cdot \lambda))$$

Now we know that for regular representations  $\mathcal{M}_\lambda$  the cohomology  $H^\nu(\mathfrak{m}, K_\infty^M, H_{\sigma_\infty} \otimes \mathcal{M}(w \cdot \lambda))$  is non zero only for  $\nu$  in a very narrow interval around the middle degree (See [Vo-Zu], Thm. 5.5). If the difference  $|l(w) - l(w')|$  is greater than the length of this interval, then the following condition is fulfilled

*In any degree  $T_\infty^{\text{loc}, \bullet}(w)$  induces zero on the cohomology. (Tzero)*

In this cases (and under the assumption that the Eisenstein series is holomorphic at  $z = 0$ ) the Eisenstein intertwining operator gives us a section for the Hecke-modules

$$\text{Eis}_\mathbb{C} : H^{q-l(w)}(\mathcal{S}_{K_f^M}^M, \mathcal{M}(w \cdot \lambda) \otimes \mathbb{C})(\sigma_f) \rightarrow H^q(\mathcal{S}_{K_f^G}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{C}) \tag{9.42}$$

### 9.4 The special case $\text{Gl}_n$

Our group is  $\text{Gl}_n/Q$  and we choose a parabolic subgroup  $P$  containing the standard Borel subgroup and with reductive quotient  $M = \text{Gl}_{n_1} \times \text{Gl}_{n_2} \times \dots \times \text{Gl}_{n_r}$ . We want to construct Eisenstein cohomology classes in  $H^\bullet(\mathcal{S}_{K_f^G}^G, \tilde{\mathcal{M}}_{\lambda, \mathbb{C}})$

starting from cuspidal classes in  $H^\bullet(\partial_P \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda, \mathbb{C}})$ . For an element  $w \in W^P$  we write

$$w(\lambda + \rho) = \underline{\mu}^{(1)} - b_1(w, \lambda)\gamma_{n_1} - b_2(w, \lambda)\gamma_{n_1+n_2} + \dots - b_r(w, \lambda)\gamma_{n_1+\dots+n_{r-1}} + d\delta. \tag{9.43}$$

It is the sum of the semi simple part (with respect to  $M$ )

$$\underline{\mu}^{(1)} = (b_1\gamma_1^M + \dots + b_{n_1-1}\gamma_{n_1-1}^M) + (b_{n_1+1}\gamma_{n_1+1}^M + \dots + b_{n_1+n_2-1}\gamma_{n_1+n_2-1}^M) + \dots \tag{9.44}$$

$$= \mu_1^{(1)} + \dots + \mu_r^{(1)} \tag{9.45}$$

and the abelian part  $\underline{\mu}^{\text{ab}}$ .

We assume that  $b_i(w, \lambda) \geq 0$  i.e.  $w(\lambda + \rho)$  is in the negative chamber and we also assume that the  $\mu_i^{(1)}$  are self dual, this is a condition on  $\lambda, w$ . We decompose the strongly inner cohomology

$$H_{\text{cusp}}^\bullet(\partial_P \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) = \bigoplus_{w \in W^P} \bigoplus_{\underline{\sigma}_f} \text{Ind}_P^G H_{\text{cusp}}^{\bullet-l(w)}(\mathcal{S}_{K_f^M}^M, \tilde{\mathcal{M}}_{w \cdot \lambda})(\underline{\sigma}_f) \tag{9.46}$$

The Künneth-theorem implies that  $\underline{\sigma}_f = \sigma_{1,f} \otimes \sigma_{2,f} \otimes \dots \otimes \sigma_{r,f}$ . At an unramified place  $p$  then this module has a Satake parameter

$$\omega_p(\underline{\sigma}_f) = \{\omega_{1,p}, \dots, \omega_{n_1,p}, \omega_{n_1+1,p}, \dots, \omega_{n_1+n_2,p}, \dots\}$$

where the first  $n_1$  entries are the Satake parameters of  $\sigma_{1,f}$  and so on.

We choose an  $\iota : E \rightarrow \mathbb{C}$ . We take an irreducible submodule  $H_{\underline{\sigma}_f}$  then we find an irreducible  $(\mathfrak{g}, K_\infty^M)$ -module  $H_{\underline{\sigma}_\infty}$  and an embedding

$$\Phi : H_{\underline{\sigma}_\infty} \otimes H_{\underline{\sigma}_f} \otimes_{E, \iota} \mathbb{C} = H_{\underline{\sigma}} \hookrightarrow \mathcal{C}_{\text{cusp}}(M(\mathbb{Q}) \backslash M(\mathbb{A})) \tag{9.47}$$

For  $\underline{z} = (z_1, z_2, \dots, z_{r-1}), z_i \in \mathbb{C}$  we define the character

$$|\gamma_P|^{\underline{z}} = |\gamma_{n_1}|^{z_1} |\gamma_{n_1+n_2}|^{z_2} \dots |\gamma_{n_1+n_2+\dots+n_{r-1}}|^{z_{r-1}} : M(\mathbb{A}) \rightarrow \mathbb{C}^\times$$

By the usual summation process we get an Eisenstein intertwining operator

$$\text{Eis}(\underline{\sigma}, \underline{z}) : I_P^G H_{\underline{\sigma}} \otimes |\gamma_P|^{\underline{z}} \rightarrow \mathcal{A}(G(\mathbb{Q}) \backslash G(\mathbb{A})) \tag{9.48}$$

the series is locally uniformly converging in a region where all  $\Re(z_i) \gg 0$  and hence the Eisenstein intertwining operator is holomorphic in this region. We know that it admits a meromorphic extension into the entire  $\mathbb{C}^{r-1}$ .

We want to evaluate at  $\underline{z} = 0$  this is possible if  $\text{Eis}(\underline{\sigma}, \underline{z})$  is holomorphic at  $\underline{z} = 0$ , we have to find out what happens at  $\underline{z} = 0$  we have to consider the constant term (constant Fourier coefficient) of  $\text{Eis}(\underline{\sigma}, \underline{z})$  along parabolic subgroups  $P_1$ . (See [H-C] ) These constant Fourier coefficients are given by integrals

$$\mathcal{F}^{P_1} : f(\underline{g}) \mapsto \int_{U_{P_1}(\mathbb{Q}) \backslash U_{P_1}(\mathbb{A})} f(\underline{u}\underline{g}) d\underline{u}. \tag{9.49}$$

It suffices to compute these constant terms only for parabolic subgroups containing our given maximal torus. It is shown in [H-C] that the constant term evaluated at  $\text{Eis}(\underline{\sigma}, \underline{z})(f)$  is zero unless  $P$  and  $P_1$  are associate, this means that the Levi subgroups  $M$  and  $M_1$  are isomorphic. (For this we need the cuspidality condition (See [H-C], )) ( But then we can find an element in the Weyl group which conjugates  $M$  into  $M_1$  and hence we may assume that  $P$  and  $P_1$  both contain our given Levi subgroup  $M$ . Of course now  $P_1$  may not contain the standard Borel subgroup.)

We may also assume that  $n_1 = n_2 = \dots = n_{j_1} < n_{j_1+1} = \dots = n_{j_1+j_2} < \dots < n_{j_1+\dots+j_{s-1}+1} = \dots = n_{j_1+\dots+j_s} = n_r$ , Then it is easy to see that the number of conjugacy classes of parabolic subgroups which contain  $M$  is equal to  $r!/j_1!j_2!\dots j_s!$ .

We compute  $\mathcal{F}^{P_1} \circ \text{Eis}(\underline{\sigma}, \underline{z})(f)$  following [H-C], . By definition (adelic variables in  $U(\mathbb{A}), P(\mathbb{A}), \dots$  are underlined)

$$\mathcal{F}^{P_1} \circ \text{Eis}(\underline{\sigma}, \underline{z})(f)(\underline{g}) = \int_{U_{P_1}(\mathbb{Q}) \backslash U_{P_1}(\mathbb{A})} \sum_{a \in P(\mathbb{Q}) \backslash G(\mathbb{Q})} f_{\underline{z}}(\underline{a} \underline{u} \underline{g}) d\underline{u} \quad (9.50)$$

Let  $W_M$  be the Weyl group of  $M$ , the Bruhat decomposition yields  $G(\mathbb{Q}) = \bigcup_{w \in W} P(\mathbb{Q}) \backslash w P_1(\mathbb{Q})$ , put  $P_1^{(w)}(\mathbb{Q}) = w^{-1} P(\mathbb{Q}) w \cap P_1(\mathbb{Q})$  then our expression becomes (we pull the summation over  $W$  to the front)

$$\mathcal{F}^{P_1} \circ \text{Eis}(\underline{\sigma}, \underline{z})(f)(\underline{g}) = \sum_{W_{M_1} \backslash W^{M, M_1} / W_M} \int_{U_{P_1}(\mathbb{Q}) \backslash U_{P_1}(\mathbb{A})} \sum_{b \in P_1^{(w)}(\mathbb{Q}) \backslash P_1(\mathbb{Q})} f_{\underline{z}}(w b \underline{u} \underline{g}) d\underline{u} \quad (9.51)$$

where  $W_M$  is the Weyl group of  $M$ . If now for a given  $w$  the intersection of algebraic groups  $w^{-1} U_1 w \cap M = V$  has dimension  $> 0$ , then this intersection is the unipotent radical of a proper parabolic subgroup of  $M$ . Since  $\sigma$  is cuspidal the integral over  $V(\mathbb{Q}) \backslash V(\mathbb{A})$  is zero, therefore this  $w$  contributes by zero. Hence we can restrict our summation over those  $w \in W$  which satisfy  $w M w^{-1} = M_1$ . let us call this set  $W^{M, M_1}$ . But then

$$P_1^{(w)}(\mathbb{Q}) \backslash P_1(\mathbb{Q}) = w^{-1} U_P(\mathbb{Q}) w \cap U_{P_1}(\mathbb{Q}) \backslash U_{P_1}(\mathbb{Q})$$

and the above expression becomes

$$\begin{aligned} \mathcal{F}^{P_1} \circ \text{Eis}(\underline{\sigma}, \underline{z})(f)(\underline{g}) &= \sum_{W_M \backslash W^{M, M_1} / W_M} \int_{U_{P_1}(\mathbb{Q}) \backslash U_{P_1}(\mathbb{A})} \sum_{v \in U_{P_1}^{(w)}(\mathbb{Q}) \backslash U_{P_1}(\mathbb{Q})} f_{\underline{z}}(w v \underline{u} \underline{g}) d\underline{u} = \\ & \sum_{W_M \backslash W^M / W^{M, M_1}} \int_{(w^{-1} U_P w \cap U_{P_1} \backslash U_{P_1})(\mathbb{A})} f_{\underline{z}}(\underline{u} \underline{g}) d\underline{u} \end{aligned} \quad (9.52)$$

Our parabolic subgroup  $P$  contains the standard Borel subgroup, let  $U_P^-$  be the unipotent radical of the opposite group. In the argument of  $f_{\underline{z}}$  we conjugate by  $w$ , then  $U_P \cap w U_{P_1} w^{-1} \backslash w U_{P_1} w^{-1} = w U_{P_1} w^{-1} \cap U_P^- = U_{P, P_1}^{-, w}$ .

$$\mathcal{F}^{P_1} \circ \text{Eis}(\underline{\sigma}, \underline{z})(f)(\underline{g}) = \sum_{W_{M_1} \backslash W^{M, M_1} / W_M} \int_{U_{P, P_1}^{-, w}(\mathbb{A})} f_{\underline{z}}(\underline{u} \underline{g}) d\underline{u} \quad (9.53)$$

We pick a  $w$ , the group  $M$  acts by the adjoint action on  $w^{-1}U_{P,P_1}^{-,w}w$  and hence by a character  $\delta_{P,P_1}^{(w)}$  on the highest exterior power of the Lie-algebra of this group. Then this operator sends

$$\mathcal{F}^{P_1,w} \circ \text{Eis}(\underline{\sigma}, \underline{z}) : I_P^G H_{\underline{\sigma}} \otimes |\gamma_P|^{\underline{z}} \rightarrow I_{P_1}^G H_{\underline{\sigma}^{w^{-1}}} \otimes (|\gamma_P|^{\underline{z}})^{w^{-1}} |\delta_{P,P_1}^{(w)}| \quad (9.54)$$

The integral is a product of local integrals over all places, we may assume that  $f_{\underline{z}} = f_{\infty,\underline{z}} \prod_{p:\text{prime}} f_{p,\underline{z}}$ , and then

$$\int_{U_{P,P_1}^{-,w}(\mathbb{A})} f_{\underline{z}}(\underline{u}w\underline{g})d\underline{u} = \int_{U_{P,P_1}^{-,w}(\mathbb{R})} f_{\infty,\underline{z}}(u_{\infty}wg_{\infty}) \prod_p \int_{U_{P,P_1}^{-,w}(\mathbb{Q}_p)} f_{p,\underline{z}}(u_pwg_p) \quad (9.55)$$

and here the local integrals yield intertwining operators

$$T_v^{P,P_1,w}(\sigma_v, \underline{z}) : I_P^G H_{\sigma_v} \otimes |\gamma_P|_v^{\underline{z}} \rightarrow I_{P_1}^G H_{\sigma_v^{w^{-1}}} \otimes |\gamma_P|_v^{w^{-1}\underline{z}} \otimes |\delta_{P,P_1}^{(w)}|_v \quad (9.56)$$

**Proposition 9.4.1.** *We can find local intertwining operators*

$$T_v^{P,P_1,w,\text{loc}}(\sigma_v, \underline{z}) : I_P^G H_{\sigma_v} \otimes |\gamma_P|_v^{\underline{z}} \rightarrow I_{P_1}^G H_{\sigma_v^{w^{-1}}} \otimes |\gamma_P|_v^{w^{-1}\underline{z}} \otimes |\delta_{P,P_1}^{(w)}|_v \quad (9.57)$$

which have the following properties

- a) They are holomorphic and nowhere zero in  $\Re z_i \geq 0$  (we are still assuming that  $\underline{\mu}$  is in the negative chamber.)
- b) They have a certain rationality property ([35], for the case of finite places 7.3.2.1, for the infinite places 8.4.5 and Chapter 9 (Weselmann).)
- c) At the unramified primes  $v = p$  they map the spherical vector to the spherical vector.

and finally we have

$$\mathcal{F}^{P_1,w} \circ \text{Eis}(\underline{\sigma}, \underline{z}) = C(w, P, P_1, \underline{\sigma}, \underline{z}) T_{\infty}^{P,P_1,w,\text{loc}}(\sigma_{\infty}, \underline{z}) \otimes \bigotimes_{p:\text{primes}} T_p^{P,P_1,w,\text{loc}}(\sigma_p, \underline{z}) \quad (9.58)$$

where  $C(w, P, P_1, \underline{\sigma}, \underline{z})$  is a meromorphic function in the variable  $\underline{z}$ . Therefore these functions  $C(w, P, P_1, \underline{\sigma}, \underline{z})$  decide whether  $\text{Eis}(\underline{\sigma}, \underline{z})$  is holomorphic at  $\underline{z} = 0$ , the poles of  $\text{Eis}(\underline{\sigma}, \underline{z})$  at  $\underline{z}$  are the poles of the  $C(w, P, P_1, \underline{\sigma}, \underline{z})$ .

We compute these factors  $C(w, P, P_1, \underline{\sigma}, \underline{z})$ . By definition the group  $U_{P,P_1}^{-,w}$  is a subgroup of  $U_P^-$  and as such it is easy to describe. Recall that our group  $M$  is  $GL_{n_1} \times \dots \times GL_{n_r}$  and this corresponds to a decomposition of  $\mathbb{Q}^n = X_1 \oplus X_2 \oplus \dots \oplus X_r$  into subspaces and for any two indices  $1 \leq i < j \leq r$  we define  $G_{i,j}$  to be the subgroup  $GL(X_i \oplus X_j)$  acting trivially on all other summands. For all pairs  $i, j$  we define the cocharacters  $\chi_{i,j} : \mathbb{G}_m \rightarrow T$  where  $\chi_{i,j}(t)$  is the diagonal matrix having  $t$  as entry at place  $i$ , and  $t^{-1}$  at place  $j$  and 1 everywhere else. We define  $\mathbf{w}_{i,j} := \langle \chi_{i,j}, \underline{\mu}^{(1)} \rangle$ .

The intersection  $G_{i,j} \cap U_{P,P_1}^{-,w}$  is either trivial or it is the full left lower block unipotent group  $U_{i,i+1}^-$

This tells us that the above integral can be written as iterated integral over subgroups of the form  $U_{\nu,\mu}(\mathbb{A})$ . To be more precise: If  $U_{P,P_1}^{-,w} \neq 1$  then we find an index  $i$  such that  $U_{i,i+1}$  is not trivial. In a first step we compute the local integral  $\int_{U_{i,i+1}(\mathbb{Q}_p)} f_{p,\underline{z}}^{(0)}(u_p w g_p) du_p$  at finite places where our representation  $\underline{\sigma}_p$  is unramified. We are basically in the situation, that our parabolic subgroup is maximal. The group  $P' = P \cap G_{i,i+1}$  contains the standard Borel subgroup,  $P'_1 = P_1 \cap G_{i,i+1}$  is the opposite and  $w = e$ . Then

$$C_p(e, P', P'_1, \underline{\sigma}, \underline{z}) = \frac{L^{\text{coh}}(\sigma_{i,p} \times \sigma_{i+1,p}^{\vee}, \frac{\mathbf{w}_{i,i+1}}{2} + b_i(w, \lambda) + \langle \chi_{i,i+1}, \underline{z} \rangle - 1)}{L^{\text{coh}}(\sigma_{i,p} \times \sigma_{i+1,p}^{\vee}, \frac{\mathbf{w}_{i,i+1}}{2} + b_i(w, \lambda) + \langle \chi_{i,i+1}, \underline{z} \rangle)} \quad (9.59)$$

A standard argument (See Langlands, Kim, Shahidi ) tells us that we can reduce the computation of the iterated integral to situations like the one above and then we get at unramified places

$$C_p(w, P, P_1, \underline{\sigma}, \underline{z}) = \prod_{i,j} \frac{L^{\text{coh}}(\sigma_{i,p} \times \sigma_{j,p}^{\vee}, \frac{\mathbf{w}_{i,j}}{2} + b_{i,j}(w, \lambda) + \langle \chi_{i,j}, \underline{z} \rangle - 1)}{L^{\text{coh}}(\sigma_{i,p} \times \sigma_{j,p}^{\vee}, \frac{\mathbf{w}_{i,j}}{2} + b_{i,j}(w, \lambda) + \langle \chi_{i,j}, \underline{z} \rangle)} \quad (9.60)$$

Here the indices  $i, j$  run over those indices for which  $U_{i,j} \subset U_{P,P_1}^{-,w}$ , and  $b_{i,j}(w, \lambda) = \langle \chi_{i,j}, \underline{\mu}^{\text{ab}} \rangle$ .

Now we define  $C_v(w, P, P_1, \underline{\sigma}, \underline{z})$  for all places  $v$  by the above expression, where we express the the cohomological  $L$  factor by the automorphic Rankin-Selberg  $L$  factor with the shift in the variable  $s$ . We go back to equation (9.58) and define

$$C(w, P, P_1, \underline{\sigma}, \underline{z}) = \prod_v C_v(w, P, P_1, \underline{\sigma}, \underline{z}). \quad (9.61)$$

We from the above proposition (9.4.1) that the factors  $C(w, P, P_1, \underline{\sigma}, \underline{z})$  determine the analytic behavior of  $Eis(\underline{\sigma}, \underline{z})$  at  $\underline{z} = 0$ . We have to exploit the analytic properties of the Rankin-Selberg  $L$ -functions. Here we have to use Shahidi's theorem which yields -(always remember that  $\underline{\mu}$  is in the negative chamber-)

$$L^{\text{coh}}(\sigma_{i,p} \times \sigma_{j,p}^{\vee}, \frac{\mathbf{w}_{i,j}}{2} + b_{i,j}(w, \lambda) + \langle \chi_{i,j}, \underline{z} \rangle - 1) \quad (9.62)$$

is holomorphic at  $\underline{z} = 0$  unless we are in the following special case:

a) In the product in formula ( 9.60) we have factors  $(i, i + 1)$  where  $n_i = n_{i+1}$ ,  $\mu_i^{(1)} = \mu_{i+1}^{(1)}$  and  $b_i(w, \lambda) = 1$ .

b) The pair  $\sigma_i \times \sigma_{i+1}$  is a segment, this means that  $\sigma_i \otimes \det_i = \sigma_{i+1}$

If these two conditions are fulfilled then  $C(w, P, P_1, \underline{\sigma}, \underline{z})$  has first order pole along  $z_i = 0$ .

The denominator is always holomorphic and never zero at  $\underline{z} = 0$ . (This is a deep theorem: it is the prime number theorem for Rankin-Selberg  $L$ -functions.)

### 9.4.1 Rationality of $L$ -values and some questions

We see that we get an abundant supply of cohomology classes: Starting from any parabolic  $P$  and an isotypical subspace  $\text{Ind}_P^G H_{\text{cusp}}^{\bullet-l(w)}(\mathcal{S}_{K_f^M}, \tilde{\mathcal{M}}_{w \cdot \lambda})(\underline{\sigma}_f)$  we get the Eisenstein intertwining operator (See equation (9.48)). We analyze what happens at  $\underline{z} = 0$ . If it is holomorphic we get a Hecke invariant homomorphism

$$\text{Eis}^\bullet(0) : H^\bullet(\mathfrak{g}, K_\infty, \text{Ind}_P^G \sigma_\infty \otimes \tilde{\mathcal{M}}) \otimes \text{Ind}_P^G H_{\underline{\sigma}_f} \rightarrow H^\bullet(\mathcal{S}_{K_f^G}, \tilde{\mathcal{M}}_{\mathbb{C}}) \quad (9.63)$$

We can restrict these cohomology classes to the boundary and even to boundary strata  $\partial_Q(\mathcal{S}_{K_f^G}, \tilde{\mathcal{M}})$  where  $Q$  runs over the parabolic subgroups associate to  $P$ , or more generally those parabolic subgroups which contain an associate to  $P$ . This means that the class "spreads out" over different boundary strata. These restrictions to these other strata are given by certain linear maps which are product of "local intertwining operators" times certain special values of  $L$  functions.

In certain cases this "spreading out" is highly non trivial. We have to clarify some local issues. First of all we have to find out whether the local intertwining operators are non zero and have certain rationality properties. Especially we have to show that these local operators at the infinite places induce non zero maps between the cohomology groups of certain induced Harish-Chandra modules. And we have to show that these maps on the level of cohomology have rationality properties. ([35], 7.3, )

If these local issues are settled then we can argue: The image of the cohomology  $H^\bullet(\mathcal{S}_{K_f^G}, \tilde{\mathcal{M}})$  in the cohomology of the boundary is defined over  $\mathbb{Q}$  (or some number field depending on our data). Since the  $L$ - values enter in the description of this image we get rationality statements for special values of  $L$ -functions.

This has been exploited in some cases ([?], [?], [32]-Mum) and a very general result in this direction is in [35](See previous section).

But in case we have a pole we may also produce cohomology classes by taking residues, again starting from one boundary stratum. The restriction of these classes to the boundary will spread out over other strata in the boundary and we may play the same game. In this case the non vanishing issue of intertwining operators on cohomological level comes up again and will be discussed in the following section. (See Thm. 9.6.1)

We also will encounter situation where a pole along a plane  $z_i = 0$  (or may be even several such planes ) "fights" with a zero along some other planes containing zero. Then this influences the structure of the cohomology. But how? This question has been discussed in [Ha-Gln]. Is the order of vanishing along this zero visible in the structure of the cohomology? Or is it visible in the structure of the cohomology of the boundary, or in the spectral sequence?

## 9.5 Residual classes

We have seen that our Eisenstein classes may be singular at  $\underline{z} = 0$ . In this section we look at the extremal case that  $\text{Eis}(\sigma, \underline{z})$  has simple poles along the lines  $z_i = \langle \chi_{n_i, n_i+1}, \underline{z} \rangle = 0$ . In this case we call these Eisenstein classes residual.

It follows from the work of Mœglin-Waldspurger [M-W] that this can only happen under some very special conditions.

We start from a factorization  $n = uv$  we look the parabolic subgroup  $P_{u,v}$  which contains the standard Borel subgroup and has reductive quotient  $\mathrm{Gl}_u \times \mathrm{Gl}_u \times \cdots \times \mathrm{Gl}_u$ . The standard maximal torus is a product  $T = \prod_{i=1}^{i=v} T_i$  and accordingly we have  $X^*(T) = \bigoplus_{i=1}^{i=v} X^*(T_i)$ . We have an obvious identification  $T_i = \mathbb{G}_m^u$ .

We choose a highest weight  $\lambda = \sum a_i \gamma_i + d\delta$ , we assume that it is self dual, i.e.  $a_i = a_{n-i}$ . We have a restriction on the character  $\mu = w \cdot \lambda = w(\lambda + \rho_N) - \rho_N$ , we must have

$$\begin{aligned} w(\lambda + \rho_N) - \rho_N &= b_1 \gamma_1^M + b_2 \gamma_2^M + \cdots + b_{u-1} \gamma_{u-1}^M - d_0 \det^{(1)} \\ &+ b_1 \gamma_{1+u}^M + b_2 \gamma_{2+u}^M + \cdots + b_{u-1} \gamma_{2u-1}^M - (d_0 + 1) \det^{(2)} + \cdots \\ &\cdots \\ &b_1 \gamma_{(v-1)u+1}^M + b_2 \gamma_2^M + \cdots + b_{u-1} \gamma_{vu-1}^M - (d_0 + v - 1) \det^{(v)} \end{aligned} \quad (9.64)$$

where  $\det^{(\nu)}$  is the determinant on the  $\nu$ -th block. Our highest weight is a sum  $\mu = \sum \mu_i$  where

$$\mu_i = \mu^{(1)} - (d_0 + i) \det^{(i)} \quad (9.65)$$

where the semi simple component  $\mu^{(1)} = b_1 \gamma_1^M + b_2 \gamma_2^M + \cdots + b_{u-1} \gamma_{u-1}^M = b_1 \gamma_{1+u}^M + b_2 \gamma_{2+u}^M + \cdots + b_{u-1} \gamma_{2u-1}^M \dots$  is "always the same". We notice that of course we have the self duality condition  $b_i = b_{u-i}$ . Furthermore we have  $\sum d_i = -d$ .

We define

$$\mathbb{D}_\mu = \bigotimes_{i=1}^{i=v} \mathbb{D}_{\mu_i} \quad (9.66)$$

and start from our isotypical  $H_{\mathrm{cusp}}^\bullet(\mathcal{S}_{K_f^M}^M, \mathcal{M}_{w \cdot \lambda})(\sigma_f)$ . The Künneth formula yields that we can write  $\sigma_f = \sigma_{1,f} \times \sigma_{2,f} \times \cdots \times \sigma_{v,f}$  where all the  $\sigma_{i,f}$  occur in the cuspidal cohomology of  $\mathrm{Gl}_u$ , hence they may be compared. The relation (9.65) allows us to require that  $\sigma_{i+1,f} = \sigma_{i,f} \otimes |\delta|$ . If this is satisfied we say that  $\sigma_f$  is a segment. We assume  $v \neq 1$  and hence  $P \neq G$ .

We know that under the assumption that  $\sigma_f$  is a segment (and only under this assumption) the factor  $C(\sigma, w_P, \underline{z})$  has a simple poles along the lines  $z_i = 0$ , and this is the only term in (??) having these poles. The operator  $T^{\mathrm{loc}}(\sigma, \underline{s})$  is a product of local operators at all places

$$T^{\mathrm{loc}}(\sigma, \underline{z}) = T_\infty^{\mathrm{loc}}(\sigma_\infty, \underline{s}) \times \prod_p T_p^{\mathrm{loc}}(\sigma_p, \underline{z}),$$

and the local factors are holomorphic as long as  $\Re(z_i) \geq 0$ . We take the residue at  $\underline{z} = 0$  i.e. we evaluate

$$\left( \prod z_i \right) \mathcal{F}^P \circ \mathrm{Eis}(\sigma \otimes \underline{s})|_{\underline{z}=0} = \left( \prod z_i \right) C(\sigma, w_P, \underline{z})|_{\underline{z}=0} T^{\mathrm{loc}}(\sigma, w_P, \underline{0})(f) \quad (9.67)$$

This tells us that the residue of the Eisenstein class gives us an intertwining operator

$$\text{Res}_{z=0} \text{Eis}(\sigma \otimes z) : {}^a\text{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})} \mathbb{D}_\mu \otimes V_{\sigma_f} \rightarrow L_{\text{disc}}^2(G(\mathbb{Q}) \backslash G(\mathbb{A}) / K_f, \omega_{\mathcal{M}_\lambda}^{-1} |_{S(\mathbb{R})^0}) \quad (9.68)$$

The image  $J_{\sigma_\infty} \otimes J_{\sigma_f}$  is an irreducible module ( this is a Langlands quotient) and via the constant Fourier coefficient it injects into  ${}^a\text{Ind}_{P(\mathbb{A})}^{G(\mathbb{A})} \mathbb{D}_{\mu'} \otimes V_{\sigma_f}$ . At the infinite place we get a diagram

$$\begin{array}{ccc} \text{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})} \mathbb{D}_\mu & \xrightarrow{T^{(\text{loc})}(D_\mu)} & J_{\sigma_\infty} \\ & & \downarrow \\ & & \text{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})} \mathbb{D}_{\mu'} \end{array} \quad (9.69)$$

It is a - not completely trivial - exercise to write down the solutions for the system of equations (9.64). This means starting from our highest weight  $\mu$  we have to find  $w, \lambda$ . The answer is that  $\lambda$  must be of the special form

$$\lambda = a_1 \gamma_v + a_2 \gamma_{2v} + \dots + a_{u-1} \gamma_{(u-1)v} + d\delta \quad (9.70)$$

which in addition is essentially self dual, i.e.  $a_i = a_{v-i}$  the number  $d$  is uninteresting and only serves to satisfy the parity condition.

We choose a specific Kostant representative  $w'_{u,v} \in W^P$ , it is the permutation in the letters  $1, 2, \dots, n$  given by the following rule: write  $\nu = i + (j - 1)v$  with  $1 \leq i \leq v$  then  $w'_{u,v}(\nu) = j + (i - 1)v$ . Then we compute  $w'_{u,v}(\lambda + \rho_N) - \rho_N \in X^*(T \times E)$  and we get

$$\begin{aligned} & (w'_{u,v}(\lambda + \rho_N) - \rho_N) = \\ & (a_1 + v - 1)\gamma_1^M + (a_2 + v - 1)\gamma_2^M + (a_{u-1} + v - 1)\gamma_{u-1}^M \\ & (a_1 + v - 1)\gamma_{1+u}^M + (a_2 + v - 1)\gamma_{2+u}^M + (a_{u-1} + v - 1)\gamma_{u-1+u}^M \\ & \vdots \\ & (a_1 + v - 1)\gamma_{1+(v-1)u}^M + (a_2 + v - 1)\gamma_{2+(v-1)u}^M + \dots + (a_{u-1} + v - 1)\gamma_{u-1+(v-1)u}^M + \\ & -(u - 1)(\gamma_u + \gamma_{2u} + \dots + \gamma_{(v-1)u}) + d\delta \end{aligned} \quad (9.71)$$

The length of this Kostant representative is

$$l(w'_{u,v}) = n(u - 1)(v - 1)/4.$$

Let  $w_P$  be the longest Kostant representative which sends all the roots in  $U_P$  to negative roots. Then we define the (reflected) Kostant representative  $w_{u,v} = w_P w'_{u,v}$ . We get

$$\begin{aligned} w_{u,v}(\lambda + \rho) - \rho = \mu = & (a_1 + v - 1)(\gamma_1^M + \gamma_{1+u}^M + \dots + \gamma_{1+(v-1)u}^M) + \\ & (a_2 + v - 1)(\gamma_2^M + \gamma_{2+u}^M + \dots + \gamma_{2+(v-1)u}^M) + \\ & \vdots \\ & (a_{u-1} + v - 1)(\gamma_{u-1}^M + \gamma_{u-1+u}^M + \dots + \gamma_{u-1+(v-1)u}^M) + \\ & -(u + 1)(\gamma_u + \gamma_{2u} + \dots + \gamma_{(v-1)u}) + d\delta \end{aligned} \quad (9.72)$$

Hence we see that the semi simple component stays the same and the abelian parts differ by  $2(\gamma_u + \gamma_{2u} + \dots + \gamma_{(v-1)u})$ . We see that we can solve (9.64) provided  $b_i \geq v - 1$ .

**The identification**  $J_{\sigma_\infty} \xrightarrow{\sim} A_q(\lambda)$

Of course we expect

$$H^\bullet(\mathfrak{g}, K_\infty, J_{\sigma_\infty} \otimes \mathcal{M}_\lambda) \neq 0. \quad (9.73)$$

In the paper [Vo-Zu] the authors give a list of irreducible  $(\mathfrak{g}, K_\infty)$  modules  $A_q(\lambda)$  which have non trivial cohomology  $H^\bullet(\mathfrak{g}, K_\infty, A_q(\lambda) \otimes \mathcal{M}_\lambda) \neq 0$ . This list contains all unitary modules having this property. On the other hand we know that any such unitary  $A_q(\lambda)$  can be written as a Langlands quotient. In the paper of Vogan and Zuckerman it is explained how we can get a given unitary  $A_q(\lambda)$  as Langlands quotient, basically this means we construct a diagram of the form (9.69) but where now we have  $A_q(\lambda)$  in the upper right corner instead of  $J_{\sigma_\infty}$ . In the following section we describe a specific  $A_q(\lambda)$  and write it as a Langlands quotient (i.e. we find its Langlands parameters) this means we determine the upper left and lower right entries and then check that these entries are the ones in diagram (9.69). From this we will derive the following

*The map*

$$H^\bullet(\mathfrak{g}, K_\infty, J_{\sigma_\infty} \otimes \mathcal{M}_\lambda) \otimes J_{\sigma_f} \rightarrow H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \quad (9.74)$$

is non zero in degree  $l(w'_{u,v}) = n(u-1)(v-1)/4$ .

See Theorem (9.6.1)

## 9.6 Detour: $(\mathfrak{g}, K_\infty)$ - modules with cohomology for $G = \mathrm{Gl}_n$

I want to fix some notations and conventions.

Let  $T/\mathbb{Q}$  be the maximal torus in  $\mathrm{Gl}_n/\mathbb{Q}$ , let  $T^{(1)} = \mathrm{Sl}_n \cap T$ . We put  $r = n-1$ . We have the standard basis for the character-module  $X^*(T)$ :

$$e_i : T \rightarrow G_m, t \mapsto t_i.$$

The positive (resp. simple roots) roots are  $\alpha_{i,j} = e_i - e_j, i < j$ , (resp.  $\alpha_i = e_i - e_{i+1}$ .) We have the determinant  $\delta = \sum_{i=1}^n e_i$ .

The fundamental weights are elements in  $X^*(T) \otimes \mathbb{Q}$ , they are defined by

$$\gamma_i = \sum_{\nu=1}^i e_\nu - \frac{i}{n} \delta,$$

these  $\gamma_i$  are the fundamental weights if we restrict to  $\mathrm{Sl}_n$ , the image of  $\gamma_i$  under the restriction map lies in  $X^*(T^{(1)})$ .

From now on my natural basis for  $X^*(T) \otimes \mathbb{Q}$  will be

$$\{\gamma_1, \dots, \gamma_i, \dots, \gamma_r, \delta\}.$$

This basis respects the decomposition of  $T$  into  $T^{(1)} \cdot G_m$ , the first factor is its component in  $Sl_n$  and the second one is the central torus.

We also have the cocharacters  $\chi_i \in X_*(T^{(1)})$  which are given by

$$\chi_i : t \mapsto \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & \ddots & \dots & \dots & \dots \\ 0 & 0 & t & 0 & \dots & 0 \\ 0 & \dots & 0 & t^{-1} & \dots & 0 \\ \dots & \dots & \dots & \dots & \ddots & \dots \\ 0 & 0 & 0 & \dots & 0 & 1 \end{pmatrix}$$

and the central cocharacter

$$\zeta : t \mapsto \begin{pmatrix} t & 0 & 0 \dots & 0 \\ 0 & t & \dots & \dots \\ 0 & 0 & \ddots & \dots \\ 0 & \dots & 0 & \dots \end{pmatrix}$$

We have the standard pairing  $(\chi, \gamma) \mapsto \langle \chi, \gamma \rangle$  between cocharacters and characters which is defined by  $\gamma \circ \chi = \{t \mapsto t^{\langle \chi, \gamma \rangle}\}$ . We have the relations

$$\langle \chi_j, \gamma_i \rangle = \delta_{ij}, \quad \langle \chi_i, \alpha_i \rangle = 2$$

the character  $\delta$  is trivial on the  $\chi_i$  and  $\delta \circ \zeta = \{t \mapsto t^n\}$ . It is clear that an element  $\gamma = \sum_i a_i \gamma_i + d\delta \in X^*(T)$  if and only if the  $a_i, nd \in \mathbb{Z}$  and we have the congruence

$$\sum i a_i \equiv nd \pmod n.$$

We identify the center of  $Gl_n$  with  $G_m$  via the cocharacter  $\zeta$ , the character module of  $G_m$  is  $\mathbb{Z}$ . Hence the central character  $\omega_\lambda$  is an integer and we find

$$\omega_\lambda = nd.$$

Actually this central character should be considered as an element in  $\mathbb{Z} \pmod n$  because we can replace  $d$  by  $r + d$  and then the central character changes by a multiply of  $n$ . If  $\lambda \in X^+(T^{(1)})$  is a dominant weight then we write it as

$$\lambda = \sum a_i \gamma_i$$

then we have  $a_i \geq 0$ .

### 9.6.1 The tempered representation at infinity

We consider the group  $Gl_n/\mathbb{R}$ , we choose a essentially selfdual highest weight  $\lambda = \sum_1^{n-1} a_i \gamma_i + d\delta$  (i.e.  $a_i = a_{n-i}$ ). The  $a_i$  are integers and  $d$  is a half integer which satisfies the parity condition

$$d \in \mathbb{Z} \text{ if } n \text{ is odd, } \frac{n}{2} a_{\frac{n}{2}} \equiv nd \pmod n \text{ if } n \text{ is even}$$

We want to recall the construction of a specific  $(\mathfrak{g}, K_\infty)$ -module  $\mathbb{D}_\lambda$  such that

$$H^\bullet(\mathfrak{g}, K_\infty, \mathbb{D}_\lambda \otimes \mathcal{M}_\lambda) \neq 0$$

and we will also determine the structure of this cohomology. This module is the only tempered Harish-Chandra module which has non trivial cohomology with coefficients in  $\mathcal{M}_\lambda$ . The center  $\mathbb{G}_m$  of  $\mathrm{Gl}_n$  acts on the module  $\mathcal{M}_\lambda$  by the character  $\omega_\lambda : x \mapsto x^{nd}$ . Since we want no zero cohomology the center  $S(\mathbb{R})$  of  $\mathrm{Gl}_n(\mathbb{R})$  acts by the central character  $(\omega_\lambda)_\mathbb{R}^{-1}$  on  $\mathbb{D}_\lambda$ . The module  $\mathbb{D}_\lambda$  will be essentially unitary with respect to that character.

We construct our representation  $\mathbb{D}_\lambda$  by inducing from discrete series representations. We consider the parabolic subgroup  ${}^\circ P$  whose simple root system is described by the diagram

$$\circ - \times - \circ - \times - \dots - \circ(-\times) \tag{9.75}$$

i.e. the set of simple roots  $I_{\circ M}$  of the semi simple part of the Levi quotient  ${}^\circ M$  consists of those which have an odd index. Let  $m$  be the largest odd integer less or equal to  $n - 1$  then  $\alpha_m$  is the last root in the system of simple roots in  $I_{\circ M}$ . Of course  $m = n - 1$  if  $n$  is even and  $m = n - 2$  else.

The reductive quotient is equal to  $\mathrm{Gl}_2 \times \mathrm{Gl}_2 \times \dots \times \mathrm{Gl}_2(\times \mathbb{G}_m)$ , where the last factor occurs if  $n$  is odd. This product decomposition of  ${}^\circ M$  induces a product decomposition of the standard maximal torus  $T = \prod_{i:i\text{ odd}} T_i(\times \mathbb{G}_m)$  and for the character module we get

$$X^*(T) = \bigoplus_{i:i\text{ odd}} X^*(T_i) \oplus X^*(\mathbb{G}_m) \tag{9.76}$$

The semi simple reductive quotient  ${}^\circ M^{(1)}(\mathbb{R})$  is  $A_1 \times A_1 \times \dots \times A_1$ , the number of factors is

$${}^\circ r = (m + 1)/2 = \begin{cases} \frac{n}{2} & \text{if } n \text{ is even} \\ \frac{n-1}{2} & \text{if } n \text{ is odd} \end{cases}$$

We also introduce the number

$$\epsilon(n) = \begin{cases} 0 & \text{if } n \text{ even} \\ 1 & \text{if } n \text{ odd} \end{cases} \tag{9.77}$$

We have a very specific Kostant representative  $w_{\mathrm{un}} \in W^{\circ P}$ . The inverse of this permutation it is given by

$$w_{\mathrm{un}}^{-1} = \{1 \mapsto 1, 2 \mapsto n, 3 \mapsto 2, 4 \mapsto n - 1, \dots\}.$$

The length of this element is equal to  $1/2$  the number of roots in the unipotent radical of  ${}^\circ P$ , i.e.

$$l(w_{\mathrm{un}}) = \begin{cases} \frac{1}{4}n(n - 2) & \text{if } n \text{ is even} \\ \frac{1}{4}(n - 1)^2 & \text{if } n \text{ is odd} \end{cases} \tag{9.78}$$

9.6. DETOUR:  $(\mathfrak{G}, K_\infty)$ -MODULES WITH COHOMOLOGY FOR  $G = \mathrm{GL}_N$  377

We compute

$$w_{\mathrm{un}}(\lambda + \rho) - \rho = \sum_{i:i \text{ odd}} b_i \gamma_i^{\circ M^{(1)}} + d\delta = \sum_{i:i \text{ odd}} b_i \frac{\alpha_i}{2} + d\delta = \tilde{\mu}^{(1)} + d\delta. \quad (9.79)$$

(The subscript  $\mathrm{un}$  refers to unitary, it refers also to the length being half the dimension of the unipotent radical. Here we have to observe that  $w \cdot \lambda$  is an element in  $X^*(T)$  but the individual summands may only lie in  $X^*(T) \otimes \mathbb{Q} = X_{\mathbb{Q}}^*(T)$ . Any element  $\gamma \in X^*(T)$  also defines a quasicharacter  $\gamma_{\mathbb{R}} : T(\mathbb{R}) \rightarrow \mathbb{R}^\times$  (by definition). But an element  $\gamma \in X_{\mathbb{Q}}^*(T)$  only defines a quasicharacter  $|\gamma|_{\mathbb{R}} : T(\mathbb{R}) \rightarrow \mathbb{R}_{>0}^\times$  which is defined by  $|\gamma|_{\mathbb{R}}(x) = |m\gamma(x)|^{1/m}$ .)

To compute the coefficients  $b_j$  we use the pairing (See 7.1) and observe that  $\langle \chi_i, \gamma_j \rangle = \delta_{i,j}$ . Then

$$b_j = \langle \chi_j, w_{\mathrm{un}}(\lambda + \rho) - \rho \rangle = \langle w_{\mathrm{un}}^{-1} \chi_j, \lambda + \rho \rangle - \langle \chi_j, \rho \rangle. \quad (9.80)$$

Now the choice of  $w_{\mathrm{un}}$  becomes clear. It is designed in such a way that

$$w_{\mathrm{un}}^{-1} \chi_1(t) = \begin{pmatrix} t & 0 & 0 & \dots & 0 \\ 0 & \ddots & \dots & \dots & \dots \\ 0 & 0 & 1 & 0 & \dots & 0 \\ 0 & \dots & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \ddots & 0 \\ 0 & 0 & 0 & \dots & 0 & t^{-1} \end{pmatrix}, w_{\mathrm{un}}^{-1} \chi_3(t) = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & t & \dots & \dots & \dots \\ 0 & 0 & 1 & 0 & \dots & 0 \\ 0 & \dots & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & t^{-1} & 0 \\ 0 & 0 & 0 & \dots & 0 & 1 \end{pmatrix}$$

and for the general odd index  $j$  we have  $w_{\mathrm{un}}^{-1} \chi_j(t) = h_{(j+1)/2}$  where for all  $1 \leq \nu \leq n/2$  we denote by  $h_\nu(t)$  the diagonal matrix which has a 1 at all entries different from  $\nu, n+1-\nu$  and which has entry  $t$  at  $\nu$  and  $t^{-1}$  at  $n+1-\nu$ . Then  $h_\nu = \{t \mapsto h_\nu(t)\}$  is a cocharacter. It is clear that

$$\gamma_i(h_\nu(t)) = \begin{cases} t & \text{if } \nu \leq i \leq n-\nu \\ 1 & \text{else} \end{cases}$$

This yields for  $j = 1, \dots, r$

$$b_{2j-1} = \sum_{\nu} (a_\nu + 1) \langle h_j, \gamma_\nu \rangle - \langle \chi_j, \rho \rangle = \left( \sum_{j \leq \nu \leq n-j} (a_\nu + 1) \right) - 1.$$

We should keep in mind that we assume  $a_\nu = a_{n-\nu}$ . Then we can rewrite the expressions for the  $b_\nu$  :

$$b_{2j-1} = \begin{cases} 2a_j + 2a_{j+1} + \dots + 2a_{\frac{n}{2}-1} + a_{\frac{n}{2}} + n - 2j & \text{if } n \text{ is even} \\ 2a_j + 2a_{j+1} + \dots + 2a_{\frac{n-1}{2}} + n - 2j & \text{if } n \text{ is odd} \end{cases} \quad (9.81)$$

The  $b_{2j+1}$  will be called the *cuspidal parameters* and we summarize

*The  $b_{2j-1}$  have the same parity, this parity is odd if  $n$  is odd. If  $n$  is even then  $b_{2j-1}$  has parity of  $a_{\frac{n}{2}}$ . We have  $b_1 > b_3 > \dots > b_m > 0$ . They only depend on the semi simple part  $\lambda^{(1)}$ .*

By Kostants theorem

$$w_{\text{un}} \cdot \lambda = w_{\text{un}}(\lambda + \rho) - \rho$$

is the highest weight of an irreducible representation of  ${}^\circ M$ . This irreducible representation occurs with multiplicity one in  $H^{l(w_{\text{un}})}(\mathfrak{u} \circ P, \mathcal{M}_\lambda)$ .

The highest weight of this representation is

$$w_{\text{un}} \cdot \lambda = w_{\text{un}}(\lambda + \rho) - \rho = \sum_{i:i \text{ odd}} b_i \gamma_i^{\circ M^{(1)}} + d\delta - (2\gamma_2 + 2\gamma_4 + \cdots + 2\gamma_{m-1} + \frac{3}{2}\gamma_{m+1}) \quad (9.82)$$

Digression: *Discrete series representations of  $\text{Gl}_2(\mathbb{R})$ , some conventions*

We consider the group  $\text{Gl}_2/\text{Spec}(\mathbb{Z})$ , the standard torus  $T$  and the standard Borel subgroup  $B$ . We have  $X^*(T) = \{\gamma = a\gamma_1 + d\delta \mid a \in \mathbb{Z}, d \in \frac{1}{2}\mathbb{Z}; a + 2d \equiv 0 \pmod{2}\}$  where

$$\gamma\left(\begin{pmatrix} t_1 & 0 \\ 0 & t_2 \end{pmatrix}\right) = t_1^{\frac{a}{2}+d} t_2^{-\frac{a}{2}+d} = \left(\frac{t_1}{t_2}\right)^{\frac{a}{2}} (t_1 t_2)^d$$

(Note that the exponents in the expression in the middle term are integers)

A dominant weight  $\lambda = a\gamma_1 + d\delta$  is a character where  $a \geq 0$ . These dominant weights parameterize the finite dimensional representations of  $\text{Gl}_2/\mathbb{Q}$ . The dual representation is given by  $\lambda^\vee = a\gamma_1 - d\delta$ . But these highest weights also parameterize the discrete series representations of  $\text{Gl}_2(\mathbb{R})$ , (or better the discrete series Harish-Chandra modules). The highest weight  $\lambda$  defines a line bundle  $\mathcal{L}_{-a\gamma+d\delta}$  on  $B \backslash G$  and

$$\mathcal{M}_\lambda = H^0(B \backslash G, \mathcal{L}_{-a\gamma+d\delta})$$

Then we get an embedding and a resulting exact sequence

$$0 \rightarrow \mathcal{M}_\lambda \rightarrow I_B^G((-a\gamma_1 + d\delta)_{\mathbb{R}}) \rightarrow \mathcal{D}_{\lambda^\vee} \rightarrow 0$$

and  $\mathcal{D}_{\lambda^\vee}$  is the discrete series representation attached to  $\lambda^\vee$ . (Note the subscript  $\mathbb{R}$  can not be pulled inside the bracket!).

A basic argument in representation theory yields a pairing

$$I_B^G((-a\gamma_1 - d\delta)_{\mathbb{R}}) \times I_B^G(((a+2)\gamma_1 + d\delta)_{\mathbb{R}}) \rightarrow \mathbb{R}$$

(here observe that  $2\gamma_1 = 2\rho \in X^*(T)$ ).

From this we get another exact sequence which gives the more familiar definition of the discrete series representation

$$0 \rightarrow \mathcal{D}_\lambda \rightarrow I_B^G(((a+2)\gamma_1 + d\delta)_{\mathbb{R}}) \rightarrow \mathcal{M}_\lambda \rightarrow 0. \quad (9.83)$$

The module  $\mathbb{D}_\lambda$  is also a module for the group  $K_\infty = \text{SO}(2)$  and it is well known that it decomposes into  $K_\infty$  types

$$D_\lambda = \cdots \oplus \mathbb{C}\psi_\nu \cdots \mathbb{C}\psi_{-a-4} \oplus \mathbb{C}\psi_{-a-2} \oplus \mathbb{C}\psi_{+a+2} \oplus \mathbb{C}\psi_{+a+4} \cdots \quad (9.84)$$

(End of digression)

9.6. DETOUR:  $(\mathfrak{G}, K_\infty)$ -MODULES WITH COHOMOLOGY FOR  $G = \mathrm{GL}_N$  379

We return to our formula (9.82). The group

$${}^\circ M = \prod_{i:\text{odd}} M_i \times (\mathbb{G}_m)$$

where  $M_i = \mathrm{Gl}_2$ . If  $T_i$  is the maximal torus in the  $i$ -th factor, then the highest weight is  $\gamma_i^{\circ M^{(1)}}$  and let  $\delta_i$  be the determinant on that factor. The indices  $i$  run over the odd numbers  $1, 3, \dots, m$ . If  $n$  is odd then let  $\delta_n : T \rightarrow \mathbb{G}_m$  be the character given by the last entry. Then we have for the determinant

$$\delta = \delta_1 + \delta_3 + \dots + \delta_m + \begin{cases} 0 \\ \delta_n \end{cases} \quad (9.85)$$

We want to write the character  $2\gamma_2 + 2\gamma_4 + \dots + 2\gamma_{m-1} + \frac{3}{2}\gamma_{m+1}$  in terms of the  $\delta_i$ . We recall that

$$\begin{aligned} \gamma_2 &= \delta_1 - \frac{2}{n}\delta \\ \gamma_4 &= \delta_1 + \delta_3 - \frac{4}{n}\delta \\ &\vdots \\ \gamma_{m-1} &= \delta_1 + \delta_3 \dots + \delta_{m-2} - \frac{m-1}{n}\delta \\ &\text{and if } n \text{ is odd} \\ \gamma_{m+1} &= \delta_1 + \delta_3 \dots + \delta_m - \frac{m+1}{n}\delta \end{aligned} \quad (9.86)$$

Then the summation over the  $\delta$ -terms on the right hand side yields

$$-\frac{1}{n}(4 + 8 + \dots + 2(m-1) - \begin{cases} 0 \\ \frac{3}{2}(m+1) \end{cases}) = -[\frac{n-1}{2}] \quad (9.87)$$

and if we take our formula (9.85) into account and also count the number of times a  $\delta_i$  occurs in the summation we get

$$2\gamma_2 + 2\gamma_4 + \dots + 2\gamma_{m-1} + \frac{3}{2}\gamma_{m+1} = \begin{cases} (\frac{n}{2}-1)\delta_1 + (\frac{n}{2}-3)\delta_3 + \dots + (-\frac{n}{2}+1)\delta_{m-2} & n \equiv 0 \pmod{2} \\ \frac{n-2}{2}\delta_1 + \dots + \frac{-n+4}{2}\delta_m - \frac{n-1}{2}\delta_n & \text{else} \end{cases} \quad (9.88)$$

Let us denote the coefficient of  $\delta_i$  in the expressions on the right hand side by  $c(i, n)$ . We recall that we still have the summand  $d\delta$  in our formula (??). Then

$$\underline{mu} = w_{\text{un}} \cdot \lambda = \sum_{i:\text{odd}} b_i \gamma_i^{\circ M^{(1)}} + (c(i, n) + d)\delta_i + \begin{cases} d\delta \\ (-\frac{n-1}{2} + d)\delta_n \end{cases} \quad (9.89)$$

We claim that the individual summands are in the character modules  $X^*(T_i)$  (resp.  $X^*(\mathbb{G}_m)$ ). This means that

$$b_i \gamma_i^{\circ M^{(1)}} + (c(i, n) + d)\delta_i \in X^*(T_i), \quad -\frac{n-1}{2} + d \in \mathbb{Z}. \quad (9.90)$$

We have to verify the parity conditions. If  $n$  is odd the the parity condition for  $\lambda$  says that  $d \in \mathbb{Z}$ . On the other hand we know that in this case the  $b_i$  are

odd and since the  $c(i, n)$  are also odd the parity condition is satisfied for the individual summands.

If  $n$  is even then the parity condition for  $\lambda$  says that  $\frac{n}{2}a_{\frac{n}{2}} \equiv nd \pmod n$ . We know that the  $b_i$  all have the same parity:  $b_i \equiv a_{\frac{n}{2}} \pmod 2$ . Hence need that  $a_{\frac{n}{2}} \equiv 2d \pmod 2$ , but this is the parity condition for  $\lambda$ .

For any of the characters  $\mu_i$  we have the induced representations  $I_{B_i}^{\circ M_i}(\mu_i + 2\rho_i)$  the discrete series representation  $\mathcal{D}_{\mu_i}$  and the exact sequence

$$0 \rightarrow \mathcal{D}_{\mu_i} \rightarrow I_{B_i}^{\circ M_i}(\mu_i + 2\rho_i) \rightarrow \mathcal{M}_{\mu_i} \rightarrow 0. \tag{9.91}$$

The tensor product

$$\mathcal{D}_{\mu} = \bigotimes_{i:\text{odd}} \mathcal{D}_{\mu_i} \otimes \mathbb{C}(-\frac{n-1}{2} + d) \tag{9.92}$$

is a module for  ${}^{\circ}M$ .

Here we have to work with  $K_{\infty}^{\circ M} = K_{\infty} \cap {}^{\circ}M$ . This compact group is not necessarily connected, its connected component of the identity is

$$K_{\infty}^{\circ M} \cap {}^{\circ}M^{(1)}(\mathbb{R}) = \text{SO}(2) \times \text{SO}(2) \times \cdots \times \text{SO}(2) = K_{\infty}^{\circ M, (1)}.$$

An easy computation shows

$$K_{\infty}^{\circ M} = \begin{cases} S(\text{O}(2) \times \text{O}(2) \times \cdots \times \text{O}(2)) & \text{if } n \text{ is even} \\ \text{O}(2) \times \text{O}(2) \times \cdots \times \text{O}(2) & \text{if } n \text{ is odd} \end{cases} \hat{E}, \tag{9.93}$$

since  $K_{\infty} \subset \text{Sl}_n(\mathbb{R})$  we have the determinant condition in the even case, in the odd case we have the  $\{\pm 1\}$  in the last factor and this relaxes the determinant condition.

Under the action of  $K_{\infty}^{\circ M, (1)}$  we get a decomposition

$$\mathcal{D}_{\underline{mu}} = \bigoplus_{\underline{\varepsilon}} \bigotimes_{i=1}^{\circ_r} \left( \bigoplus_{\nu_i=0}^{\infty} \mathbb{C}\psi_{\varepsilon_i(b_i+2+2\nu_i)} \right) \tag{9.94}$$

occur with multiplicity one. Here  $\underline{\varepsilon} = (\dots, \varepsilon_i, \dots)$  is an array of signs  $\pm 1$ . The induced representation (algebraic induction)

$$\text{Ind}_{\circ P(\mathbb{R})}^{G(\mathbb{R})} \mathcal{D}_{\underline{mu}} = \mathbb{D}_{\lambda} \tag{9.95}$$

is an irreducible essentially unitary  $(\mathfrak{g}, K_{\infty})$ -module, this is the module we wanted to construct. (To be more precise: We first construct the induced representation of  $G(\mathbb{R})$  where  $G(\mathbb{R})$  is acting on vectors space  $V_{\infty}$  consisting of a suitable class of functions from  $G(\mathbb{R})$  with values in  $\mathcal{D}_{\underline{mu}}$  and then we take the  $K_{\infty}$  finite vectors in  $V_{\infty}$ .) The restriction of this module to  $K_{\infty}^{(1)}$  is given by

$$\text{Ind}_{K_{\infty}^{\circ M(1)}}^{K_{\infty}^{(1)}} \mathcal{D}_{\underline{mu}} = \bigoplus_{\underline{\varepsilon}} \bigotimes_{i=1}^{\circ_r} \left( \bigoplus_{\nu_i=0}^{\infty} \text{Ind}_{K_{\infty}^{\circ M(1)}}^{K_{\infty}^{(1)}} \mathbb{C}\psi_{\varepsilon_i(b_i+2+2\nu_i)} \right) \tag{9.96}$$

(The last induced module is defined in terms of the theory of algebraic groups. We consider  $K_{\infty}^{(1)}$  as the group of real points of an algebraic group, namely the

9.6. DETOUR:  $(\mathfrak{G}, K_\infty)$ -MODULES WITH COHOMOLOGY FOR  $G = \mathrm{GL}_N$  381

connected group of the identity of the fixed points under the Cartan involution  $\Theta$ . Then  $K_\infty^{\circ M^{(1)}}$  is the group of real points of a maximal torus. Then

$$\begin{aligned} & \mathrm{Ind}_{K_\infty^{\circ M^{(1)}}}^{K_\infty^{(1)}} \mathbb{C} \psi_{\varepsilon_i(b_i+2+2\nu_i)} = \\ & \{f \mid f \text{ regular function } f(tk) = \prod_j e_i(t)^{\varepsilon_i(b_i+2+2\nu_i)} f(k), \text{ for all } t \in K_\infty^{\circ M^{(1)}}, k \in K_\infty\} \end{aligned} \quad (9.97)$$

)

We compute the cohomology of this module

$$\mathrm{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), D_\lambda \otimes \mathcal{M}_\lambda) = H^\bullet(\mathfrak{g}, K_\infty, D_\lambda \otimes \mathcal{M}_\lambda),$$

i.e. the differentials in the complex on the left hand side are all zero. (Reference to 4.1.4)

We apply Delorme to compute this cohomology. We can decompose  ${}^\circ \mathfrak{m} = {}^\circ \mathfrak{m}^{(1)} \oplus \mathfrak{a}$  then  ${}^\circ \mathfrak{k} \subset {}^\circ \mathfrak{m}^{(1)}$  and

$$\begin{aligned} \mathrm{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), D_\lambda \otimes \mathcal{M}_\lambda) &= \mathrm{Hom}_{K_\infty^{\circ M}}(\Lambda^\bullet({}^\circ \mathfrak{m}/{}^\circ \mathfrak{k}), \mathcal{D}_{\bar{\mu}} \otimes \mathcal{M}_{w_{\mathrm{un}} \cdot \lambda}) = \\ & \mathrm{Hom}_{K_\infty^{\circ M}}(\Lambda^\bullet({}^\circ \mathfrak{m}^{(1)}/{}^\circ \mathfrak{k}), \mathcal{D}_{\bar{\mu}} \otimes \mathcal{M}_{w_{\mathrm{un}} \cdot \lambda}) \otimes \Lambda^\bullet(\mathfrak{a}). \end{aligned} \quad (9.98)$$

If we replace  $K_\infty^{\circ M}$  on the right hand side by its connected component of the identity then we have an obvious decomposition

$$\mathrm{Hom}_{K_\infty^{\circ M, (1)}}(\Lambda^\bullet({}^\circ \mathfrak{m}^{(1)}/{}^\circ \mathfrak{k}), \mathcal{D}_\mu \otimes \mathcal{M}_{w_{\mathrm{un}} \cdot \lambda}) = \bigotimes_{i:i \text{ odd}} \mathrm{Hom}_{K_\infty^{i, \circ M, (1)}}(\Lambda^\bullet({}^\circ \mathfrak{m}^{(i,1)}/{}^\circ \mathfrak{k}^i), \mathcal{D}_{b_i} \otimes \mathcal{M}_{b_i}) \quad (9.99)$$

the factors on the right hand side are of rank two: We have  $K_\infty^{i, \circ M, (1)} = \mathrm{SO}(2)$  and under the adjoint action of  $K_\infty^{i, \circ M, (1)}$  the module  $\mathfrak{m}^{(i,1)}/{}^\circ \mathfrak{k}^i \otimes \mathbb{C}$  decomposes

$$\mathfrak{m}^{(i,1)}/{}^\circ \mathfrak{k}^i \otimes \mathbb{C} = \mathbb{C} P_{i,+}^\vee \oplus \mathbb{C} P_{i,-}^\vee$$

(See [Sltwo.pdf]) Then the two summands are generated by the tensors

$$\omega_{i,+} = P_{i,+}^\vee \otimes \psi_{b_i+2} \otimes m_{-b_i}, \bar{\omega}_{i,-} = P_{i,-}^\vee \otimes \psi_{-b-2} \otimes m_{b_i} \quad (9.100)$$

where  $m_{\pm(b_i)}$  is a highest (resp.) lowest weight vector for  $K_\infty^{i, \circ M}$  acting on  $\mathcal{M}_{w_{\mathrm{un}} \cdot \lambda}$ . On the tensor product on the right we have an action of the maximal compact subgroup  $\mathrm{O}(2) \times \mathrm{O}(2) \times \cdots \times \mathrm{O}(2)$  and under this action it decomposes into eigenspaces of dimension one. These eigenspaces are given by the product of sign characters  $\underline{\epsilon} = (\epsilon_1, \epsilon_2, \dots)$ .

Then it becomes clear that  $\mathrm{Hom}_{K_\infty^{\circ M}}(\Lambda^\bullet({}^\circ \mathfrak{m}^{(1)}/{}^\circ \mathfrak{k}), \mathcal{D}_{\underline{mu}} \otimes \mathcal{M}_{w_{\mathrm{un}} \cdot \lambda})$  is of rank one if  $n$  is odd and for  $n$  even it decomposes into two eigenspaces for the action of the group  $\mathrm{O}(2) \times \mathrm{O}(2) \times \cdots \times \mathrm{O}(2)/S(\mathrm{O}(2) \times \mathrm{O}(2) \times \cdots \times \mathrm{O}(2)) = \{\pm 1\}$

$$\begin{aligned} & \mathrm{Hom}_{K_\infty^{\circ M}}(\Lambda^\bullet({}^\circ \mathfrak{m}^{(1)}/{}^\circ \mathfrak{k}), \mathcal{D}_{\underline{mu}} \otimes \mathcal{M}_{w_{\mathrm{un}} \cdot \lambda}) = \\ & \mathrm{Hom}_{K_\infty^{\circ M}}(\Lambda^\bullet({}^\circ \mathfrak{m}^{(1)}/{}^\circ \mathfrak{k}), \mathcal{D}_{\underline{mu}} \otimes \mathcal{M}_{w_{\mathrm{un}} \cdot \lambda})_+ \oplus \mathrm{Hom}_{K_\infty^{\circ M}}(\Lambda^\bullet({}^\circ \mathfrak{m}^{(1)}/{}^\circ \mathfrak{k}), \mathcal{D}_{\underline{mu}} \otimes \mathcal{M}_{w_{\mathrm{un}} \cdot \lambda})_- \end{aligned}$$

We have to recall that  $\mathcal{M}_{\lambda_{\circ M}^{\text{un}}} = H^{l(w_{\text{un}})}(\mathfrak{u}_{\circ P}, \mathcal{M}_{\lambda})$  is a cohomology group in degree  $l(w_{\text{un}})$ . The classes in the factors of the last tensor product lie in degree 1, hence the multiply up to classes in degree  ${}^{\circ}r$ . This means that

$$H^q(\mathfrak{g}, K_{\infty}, \mathbb{D}_{\lambda} \otimes \mathcal{M}_{\lambda}) \neq 0 \text{ exactly for } q \in [l(w_{\text{un}}) + {}^{\circ}r, l(w_{\text{un}}) + n] \quad (9.101)$$

in the minimal degree  ${}^{\circ}r$  it is of rank 2 or 1 depending on the parity of  $n$ .

### 9.6.2 The lowest $K_{\infty}$ type in $\mathbb{D}_{\lambda}$

The maximal compact subgroup  $K_{\infty}$  is the fixed group of the standard Cartan-involution  $\Theta : g \mapsto {}^t g^{-1}$ . The subgroup  ${}^{\circ}M$  is fixed under  $\Theta$  and the subgroup  $\text{SO}(2) \times \text{SO}(2) \times \cdots \times \text{SO}(2) = K_{\infty}^{\circ M, (1)} = T_1^c$  is a maximal torus in  $K_{\infty}$ . It is the stabilizer of a direct sum decompositions of  $\mathbb{R}^n$  into two dimensional oriented planes  $V_i$  plus a line  $\mathbb{R}z$  if  $n$  is odd, we write

$$\mathbb{R}^n = \bigoplus V_i \oplus (\mathbb{R}z) \quad (9.102)$$

The Cartan involution is the identity on our torus  $T_1^c/\mathbb{R}$ . This torus can be supplemented to a  $\Theta$ -stable maximal torus by multiplying it by the torus  $T_{1, \text{split}}$  which is the product of the diagonal tori acting on the  $V_i$  in (9.102) times another copy of  $\mathbb{G}_m$  acting on  $\mathbb{R}z$  (if necessary). So we get a maximal torus  $T_1 = T_1^c \cdot T_{1, \text{split}}$ . Obviously  $T_1$  is the centralizer of  $T_1^c$  and the centralizer of  $T_{1, \text{split}}$  is the group  ${}^{\circ}M$ .

If we base change to  $\mathbb{C}$  then  $T_1^c$  splits. We identify

$$\text{SO}(2) \xrightarrow{\sim} \begin{pmatrix} a & b \\ -b & a \end{pmatrix} \quad (9.103)$$

and then the character group  $X^*(T_1^c \times \mathbb{C}) = \bigoplus \mathbb{Z}e_{\nu}$  where on the  $\nu$ -th component  $e_{\nu} : \begin{pmatrix} a & b \\ -b & a \end{pmatrix} \mapsto a + bi = a + b\sqrt{-1}$ . Then this choice provides a Borel subgroup  $B_c \supset T_1^c \times \mathbb{C}$ , for which the simple roots  $\alpha_1^c, \alpha_2^c, \dots, \alpha_r^c$  are

$$\begin{cases} e_1 - e_2, e_2 - e_3, \dots, e_{\circ r-1} - e_{\circ r}, e_{\circ r-1} + e_{\circ r} & \text{for } n \text{ even} \\ e_1 - e_2, e_2 - e_3, \dots, e_{\circ r} & \text{if } n \text{ is odd} \end{cases}$$

(See [Bou]). For  $n$  even we get the fundamental dominant weights

$$\gamma_{\nu}^c = \begin{cases} e_1 + e_2 + \cdots + e_{\nu}, & \text{if } \nu < {}^{\circ}r - 1 \\ \frac{1}{2}(e_1 + e_2 + \cdots + e_{\circ r-1} - e_{\circ r}) & \text{if } \nu = {}^{\circ}r - 1 \\ \frac{1}{2}(e_1 + e_2 + \cdots + e_{\circ r-1} + e_{\circ r}) & \text{if } \nu = {}^{\circ}r \end{cases} \quad (9.104)$$

and for  $n$  odd we get

$$\gamma_{\nu}^c = \begin{cases} e_1 + e_2 + \cdots + e_{\nu}, & \text{if } \nu < {}^{\circ}r \\ \frac{1}{2}(e_1 + e_2 + \cdots + e_{\circ r}) & \text{last weight} \end{cases} \quad (9.105)$$

An easy calculation shows

$$\sum_{i=1}^{\circ r} g_i e_i = \begin{cases} (g_1 - g_2)\gamma_1^c + (g_2 - g_3)\gamma_2^c + \cdots + (g_{\circ r-1} - g_{\circ r})\gamma_{\circ r-1}^c + (g_{\circ r-1} + g_{\circ r})\gamma_{\circ r}^c & n \text{ even} \\ (g_1 - g_2)\gamma_1^c + (g_2 - g_3)\gamma_2^c + \cdots + (g_{\circ r-1} - g_{\circ r})\gamma_{\circ r-1}^c + 2g_{\circ r}\gamma_{\circ r}^c & n \text{ odd} \end{cases} \quad (9.106)$$

The character  $\sum_{i=1}^{\circ r} g_i e_i$  is dominant (with respect to  $B_c$ ) if

$$\begin{cases} g_1 \geq g_2 \geq \cdots \geq g_{\circ r-1} \geq \pm g_{\circ r} & \text{if } n \text{ is even} \\ g_1 \geq g_2 \geq \cdots \geq g_{\circ r-1} \geq g_{\circ r} \geq 0 & \end{cases} \quad (9.107)$$

Under the action of  $K_\infty^{(1)}$  the  $(\mathfrak{g}, K_\infty^{(1)})$ -module  $\mathbb{D}_\lambda$  decomposes into a direct sum

$$\mathbb{D}_\lambda = \bigoplus_{\mu^c} \mathbb{D}_\lambda(\Theta_{\mu^c}) \quad (9.108)$$

where  $\mu^c \in X^*(T^c \times \mathbb{C})$  is a highest weight,  $\Theta_{\mu^c}$  is the resulting irreducible  $K_\infty$ -module and  $\mathbb{D}_\lambda(\Theta_{\mu^c})$  is the isotypical component.

We introduce the highest weight (see (9.81))

$$\mu_0^c(\lambda) = (b_1 + 2)e_1 + (b_3 + 2)e_2 + \cdots + (b_{2\circ r-1} + 2)e_{\circ r} \quad (9.109)$$

and in terms of our dominant weight  $\lambda$  this is

$$\mu_0^c(\lambda) = \begin{cases} 2(a_1 + 1)\gamma_1^c + \cdots + 2(a_{\circ r-1} + 1)\gamma_{\circ r-1}^c + 2(a_{\circ r-1} + a_{\circ r} + 3)\gamma_{\circ r}^c & \text{if } n \text{ is even} \\ 2(a_1 + 1)\gamma_1^c + \cdots + 2(a_{\circ r} + 3)\gamma_{\circ r}^c & \text{if } n \text{ is odd} \end{cases} \quad (9.110)$$

For  $\lambda = 0$  we get an expression (not depending on the parity of  $n$ )

$$\mu_0^c(0) = 2\gamma_1^c + \cdots + 2\gamma_{\circ r-1}^c + 6\gamma_{\circ r}^c \quad (9.111)$$

In the case that  $n$  is even the group  $K_\infty$  contains the element  $\theta$  which maps  $e_i \rightarrow e_i$  for  $i \leq \circ r - 1$  and  $e_{\circ r} \rightarrow -e_{\circ r}$  or what amounts to the same exchanges  $\gamma_{\circ r-1}^c$  and  $\gamma_{\circ r}^c$  and fixes the other fundamental dominant weights. Then

$$\bar{\mu}_0^c(\lambda) := \vartheta(\mu_0^c(\lambda)) = 2\gamma_1^c + \cdots + 6\gamma_{\circ r-1}^c + 2\gamma_{\circ r}^c + \vartheta(\lambda^c) \quad (9.112)$$

**Proposition 9.6.1.** *If  $n$  is odd then the  $K_\infty^{(1)}$ -type  $\Theta_{\mu_0^c(\lambda)}$  occurs in  $\mathbb{D}_\lambda$  with multiplicity one. All other occurring  $K_\infty^{(1)}$ -types are "larger", i.e. their highest weight satisfies  $\mu^c = \mu_0^c(\lambda) + \sum n_i \alpha_i^c$  with  $n_i \geq 0$ . We have*

$$H^\bullet(\mathfrak{g}, K_\infty, \mathbb{D}_\lambda \otimes \mathcal{M}_\lambda) = \mathrm{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \Theta_{\mu_0^c(\lambda)} \otimes \mathcal{M}_\lambda)$$

*If  $n$  is even then the  $(\mathfrak{g}, K_\infty^{(1)})$  module  $\mathbb{D}_\lambda$  decomposes into two irreducible sub modules*

$$\mathbb{D}_\lambda = \mathbb{D}_\lambda^+ \oplus \mathbb{D}_\lambda^-.$$

The  $K_\infty^{(1)}$  types  $\Theta_{\mu_0^c(\lambda)}$  resp.  $\Theta_{\bar{\mu}_0^c(\lambda)}$  occur with multiplicity one (resp. zero) in  $\mathbb{D}_\lambda^+$  (resp.  $\mathbb{D}_\lambda^-$ ). They are the lowest  $K_\infty^{(1)}$  types respectively. We have

$$H^\bullet(\mathfrak{g}, K_\infty^{(1)}, \mathbb{D}_\lambda \otimes \mathcal{M}_\lambda) = H^\bullet(\mathfrak{g}, K_\infty^{(1)}, \mathbb{D}_\lambda^+ \otimes \mathcal{M}_\lambda) \oplus H^\bullet(\mathfrak{g}, K_\infty^{(1)}, \mathbb{D}_\lambda^- \otimes \mathcal{M}_\lambda) = \\ \text{Hom}_{K_\infty^{(1)}}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \Theta_{\mu_0^c(\lambda)} \otimes \mathcal{M}_\lambda) \oplus \text{Hom}_{K_\infty^{(1)}}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \Theta_{\bar{\mu}_0^c(\lambda)} \otimes \mathcal{M}_\lambda)$$

*Proof.* For two fundamental weights we write  $\mu^c \geq \mu_1^c$  if  $\mu^c$  is "larger" than  $\mu_1^c$  in the above sense. We start from (9.96) and consider a single summand  $\text{Ind}_{K_\infty^{(1)} M^{(1)}}^{K_\infty^{(1)}} \mathbb{C}\psi_{\varepsilon_i(b_i+2+2\nu_i)}$ . This induced module decomposes into isotypical modules

$$\text{Ind}_{K_\infty^{(1)} M^{(1)}}^{K_\infty^{(1)}} \mathbb{C}\psi_{\varepsilon_i(b_i+2+2\nu_i)} = \bigoplus_{\mu^c} \text{Ind}_{K_\infty^{(1)} M^{(1)}}^{K_\infty^{(1)}} \mathbb{C}\psi_{\varepsilon_i(b_i+2+2\nu_i)}(\Theta_{\mu^c}) \quad (9.113)$$

where  $\mu^c$  runs over the set of dominant weights, where  $\Theta_{\mu^c}$  is the irreducible module of highest weight  $\mu^c$  and where  $\text{Ind}_{K_\infty^{(1)} M^{(1)}}^{K_\infty^{(1)}} \mathbb{C}\psi_{\varepsilon_i(b_i+2+2\nu_i)}(\Theta_{\mu^c})$  is the isotypical component. If we pick any dominant weight  $\mu^c$  then Frobenius reciprocity yields that

$$\Theta_{\mu^c} \text{ occurs in } \text{Ind}_{K_\infty^{(1)} M^{(1)}}^{K_\infty^{(1)}} \mathbb{C}\psi_{\varepsilon_i(b_i+2+2\nu_i)} \text{ with multiplicity } k \iff \\ t \mapsto \prod_j e_i(t)^{\varepsilon_i(b_i+2+2\nu_i)} \text{ occurs in } \Theta_{\mu^c} \text{ with multiplicity } k \quad (9.114)$$

and if  $k > 0$  this implies  $\mu^c \geq t \mapsto \prod_j e_i(t)^{\varepsilon_i(b_i+2+2\nu_i)}(t)$ . It is easy to see that we get minimal  $K_\infty^{(1)}$  types only if all  $\nu_i = 0$ . But

$$t \mapsto \prod_j e_i(t)^{\varepsilon_i(b_i+2)} \text{ is dominant} \iff \begin{cases} \varepsilon = (1, 1, \dots, 1, \pm 1) \text{ if } n \text{ even} \\ \varepsilon = (1, 1, \dots, 1, 1) \text{ if } n \text{ odd} \end{cases} \quad (9.115)$$

and in the  $n$  even case these two characters are exactly  $\mu_0^c(\lambda)$  and  $\bar{\mu}_0^c(\lambda)$  and in the  $n$  odd case this character is  $\mu_0^c(\lambda)$ .  $\square$

### 9.6.3 The unitary modules with cohomology, cohomological induction.

We start from an essentially self dual highest weight  $\lambda$  and the attached highest weight module  $\mathcal{M}_\lambda$ . In their paper [Vo-Zu] Vogan and Zuckerman construct a finite family of  $(\mathfrak{g}, K_\infty)$  modules denoted by  $A_q(\lambda)$  which have non trivial cohomology with coefficients in  $\mathcal{M}_\lambda$ , i.e.

$$H^\bullet(\mathfrak{g}, K_\infty, A_q(\lambda) \otimes \mathcal{M}_\lambda) \neq 0$$

They also show that all unitary irreducible  $(\mathfrak{g}, K_\infty)$ -modules with non trivial cohomology in with coefficients in  $\mathcal{M}_\lambda$  are of this form. We briefly recall their construction and translate it into our language and our way of thinking about these issues.

9.6. DETOUR:  $(\mathfrak{g}, K_\infty)$ -MODULES WITH COHOMOLOGY FOR  $G = GL_N$  385

We introduce the torus  $\mathbb{S}^1/\mathbb{R}$  whose group of real points is the unit circle in  $\mathbb{C}^\times$  and we chose once for all an isomorphism

$$i_0 : \mathbb{S}^1 \times_{\mathbb{R}} \mathbb{C} \xrightarrow{\sim} \mathbb{G}_m/\mathbb{C} \tag{9.116}$$

We consider the free  $\mathbb{Z}$  module

$$\text{Hom}_{\mathbb{R}}(\mathbb{S}^1, T_1^c) = \text{Hom}_{\mathbb{R}}(\mathbb{S}^1, T_1) = X_*(T_1^c \times_{\mathbb{R}} \mathbb{C})$$

where of course the last identification depends on the choice of  $i_0$ . We have the standard pairing  $\langle \cdot, \cdot \rangle : X_*(T_1 \times_{\mathbb{R}} \mathbb{C}) \times X^*(T_1 \times_{\mathbb{R}} \mathbb{C}) \rightarrow \mathbb{Z}$ .

The first ingredient in the construction of an  $A_{\mathfrak{q}}(\lambda)$  is the choice of a cocharacter  $\chi : \mathbb{S}^1 \rightarrow T_c$  (defined over  $\mathbb{R}$ ). From this cocharacter we get the centralizer  $Z_\chi$ , this is a reductive subgroup whose set of roots is

$$\Delta_\chi = \{\alpha \in \Delta \subset X^*(T_1 \times_{\mathbb{R}} \mathbb{C}) \mid \langle \chi, \alpha \rangle = 0\}.$$

We can also define

$$\Delta_\chi^+ = \{\alpha \mid \langle \chi, \alpha \rangle > 0\},$$

this set depends on the choice of  $i_0$  (see (3.23)). This provides a parabolic subgroup  $P_\chi \subset G \times_{\mathbb{R}} \mathbb{C}$  whose system of roots is  $\Delta_\chi \cup \Delta_\chi^+$ . Clearly  $\Theta(P_\chi) = P_\chi$  hence  $P_\chi$  is the  $\Theta$ -stable parabolic subgroup attached to the datum  $\chi$ . This parabolic subgroup is only defined over  $\mathbb{C}$ , if we intersect it with its conjugate  $\bar{P}_\chi$  then we get the centralizer  $Z_\chi$  of  $\chi$ . We relate this to the notations in [Vo-Zu]: the  $\mathfrak{q}$  in  $A_{\mathfrak{q}}(\lambda)$  is the Lie-algebra of  $P_\chi$ , the group  $Z_\chi$  is the  $L$ . Let  $\mathfrak{u}_\chi$  be the Lie algebra of  $U_\chi$ . The datum  $\chi$  determines the  $\mathfrak{q}$  in  $A_{\mathfrak{q}}(\lambda)$ . We could introduce the notation  $A_{\mathfrak{q}}(\lambda) = A_\chi(\lambda)$ . Since  $T_1$  is the centralizer of  $T_c$  we can find a generic cocharacter  $\chi_{\text{gen}}$  such that  $P_{\chi_{\text{gen}}} = B_c$  our chosen Borel subgroup in  ${}^\circ M$ .

To a highest weight  $\lambda$  which is trivial on the semi-simple part  $Z_\chi^{(1)}$  Vogan-Zuckerman attach an irreducible unitary  $(\mathfrak{g}, K_\infty)$  module  $A_{\mathfrak{q}}(\lambda)$  such that

$$H^\bullet(\mathfrak{g}, K_\infty, A_{\mathfrak{q}}(\lambda) \otimes \mathcal{M}_\lambda) \neq 0.$$

Vogan and Zuckerman show (based on results of many others) that all the unitary irreducible  $(\mathfrak{g}, K_\infty)$  modules with non trivial cohomology in  $\mathcal{M}_\lambda$  are isomorphic to an  $A_{\mathfrak{q}}(\lambda)$ .

Furthermore they give a description of the  $K_\infty$  types occurring in  $A_{\mathfrak{q}}(\lambda)$  especially they show that  $A_{\mathfrak{q}}(\lambda)$  contains a lowest  $K_\infty$  type. This lowest  $K_\infty$ -type is given by a dominant weight which obtained by the following rule:

We consider the action of the group  $K_\infty$  on the unipotent radical  $U_\chi$  and on the Lie algebra  $\mathfrak{u}_\chi$  and the restriction of this action to  $T_1^c$ . The torus  $T_1$  also acts on  $\mathfrak{u}_\chi$  and under this action we get a decomposition into one dimensional eigenspaces

$$\mathfrak{u}_\chi = \bigoplus_{\alpha \in \Delta_\chi^+} \mathfrak{u}_\alpha$$

let us choose generators  $X_\alpha$  in these eigenspaces. We observe that the roots  $\alpha, \Theta\alpha \in \Delta^+$  induce the same root  $\alpha_c$  on  $T_1^c$ . The vector  $V_{\alpha_c} = X_\alpha - \Theta X_\alpha \in \mathfrak{u}_\chi$  is a non zero eigenvector for  $T_1^c$  and

$$\mathfrak{u}_\chi \cap (\mathfrak{p} \otimes \mathbb{C}) = \bigoplus_{(\alpha, \Theta\alpha) \in \Delta_\chi^+} \mathbb{C}V_{\alpha_c}$$

the sum runs over the unordered pairs. Then

$$\mu_c(\chi, \lambda) = \sum_{(\alpha, \Theta\alpha) \in \Delta_\chi^+} \alpha_c + \lambda_c \tag{9.117}$$

is a highest weight of a representation  $\Theta_{\mu_c(\chi, \lambda)}$  of  $K_\infty^{(1)}$  and this is the lowest  $K_\infty^{(1)}$  type in  $A_q(\lambda)$ . We get

$$H^\bullet(\mathfrak{g}, K_\infty^{(1)}, A_q(\lambda) \otimes \mathcal{M}_\lambda) = \text{Hom}_{K_\infty^{(1)}}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), A_q(\lambda) \otimes \mathcal{M}_\lambda) = \text{Hom}_{K_\infty^{(1)}}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \Theta_{\mu_c(\chi, \lambda)} \otimes \mathcal{M}_\lambda) \tag{9.118}$$

The module is determined by these properties:

- 1) It has non trivial cohomology with coefficients in  $\mathcal{M}_\lambda$
- 2) It has  $\mu_c(\chi, \lambda)$  as highest weight of a minimal  $K_\infty$  type. (See Thm. 5. 3 in [Vo-Zu].)

Recall that our aim at this moment is to identify the module  $J_{\sigma_\infty}$  to an  $A_q(\lambda)$ , and to achieve this goal we exhibit a list of very specific  $A_q(\lambda)$ 's.

**Comparison of two tori**

We need to compute  $\mu_c(\chi, \lambda)$  and to achieve this goal the author of this book modifies the Cartan involution in order to do the computation in a split group. Our standard torus  $T$  is contained in the standard Borel subgroup  $B$  of upper triangular matrices. Let  $w_0$  be an element in the normalizer of  $T$  which conjugates  $B$  into its opposite Borel subgroup. If we replace our Cartan involution by  $\Theta_1 = w_0\Theta$  then  $\Theta_1$  fixes  $T$  and the Borel subgroup  $B$ . This is not a Cartan involution, but it is easily seen that it is conjugate to  $\Theta$  over  $\text{Gl}_n(\mathbb{C})$ . and

$$\Theta_1 : \begin{pmatrix} t_1 & 0 & 0 & \dots & \\ 0 & t_2 & \dots & \dots & \\ 0 & 0 & \ddots & 0 & \dots \\ 0 & \dots & t_{n-1} & & \\ 0 & & & & t_n \end{pmatrix} \mapsto \begin{pmatrix} t_n^{-1} & 0 & 0 & \dots & \\ 0 & t_{n-1}^{-1} & \dots & \dots & \\ 0 & 0 & \ddots & 0 & \dots \\ 0 & \dots & t_2^{-1} & & \\ 0 & & & & t_1^{-1} \end{pmatrix} \tag{9.119}$$

We can decompose  $T$  up to isogeny into a torus  $T_c$  on which  $\Theta_1$  acts by the identity and a torus  $T_{\text{split}}$  where it acts by  $x \mapsto x^{-1}$  :

$$T_c = \left\{ \begin{pmatrix} t_1 & 0 & 0 & \dots & \\ 0 & t_2 & \dots & \dots & \\ 0 & 0 & \ddots & 0 & \dots \\ 0 & \dots & t_2^{-1} & & \\ 0 & & & & t_1^{-1} \end{pmatrix} \right\} \text{ resp. } T_{\text{split}} = \left\{ \begin{pmatrix} t_1 & 0 & 0 & \dots & \\ 0 & t_2 & \dots & \dots & \\ 0 & 0 & \ddots & 0 & \dots \\ 0 & \dots & t_2 & & \\ 0 & & & & t_1 \end{pmatrix} \right\}$$

It is clear that a suitable permutation matrix conjugates  $T_{1,\text{split}}$  into  $T_{\text{split}}$ . This permutation matrix maps the centralizer of  $T_{1,\text{split}}$  (which is  ${}^\circ M$ ) to the

centralizer  ${}^\circ M'$  of  $T_{\text{split}}$  and the anisotropic torus  $T_1^c$  to an anisotropic torus  $T_1^{c'}$  in  ${}^\circ M'$ . Then we can find an element  $m \in {}^\circ M'(\mathbb{C})$  which conjugates  $T_1^{c'} \times \mathbb{C}$  into  $T_c$ .

The composition of these conjugations provides an identification of the character modules  $X^*(T_1 \times \mathbb{C}) = X^*(T)$  which respects the product decompositions and hence we get

$$X^*(T_1^c \times \mathbb{C}) = X^*(T_c). \tag{9.120}$$

We choose our conjugating element  $m$  such that the  $e_i \in X^*(T_1^c \times \mathbb{C})$  are mapped to the element  $t \mapsto t_i$  (for  $i = 1$  to  ${}^\circ r$ ).

Inside  $X^*(T)$  we have the dominant fundamental weights  $\gamma_1, \dots, \gamma_{n-1}$ , let  $\bar{\gamma}_i$  be the restriction of  $\gamma_i$  to  $T_1^c$  then we have  $\bar{\gamma}_i = \bar{\gamma}_{n-i}$ . We can interpret the  $\bar{\gamma}_i$  also as elements in  $X^*(T_1 \times \mathbb{C}) \otimes \mathbb{Q}$  we require that the restriction of  $\bar{\gamma}_i$  to  $T_{1,\text{split}}$  is trivial. Then we can write

$$\bar{\gamma}_i = \begin{cases} \frac{1}{2}(\gamma_i + \gamma_{n-i}) & \text{if } i \neq \frac{n}{2} \\ \gamma_i & \text{else} \end{cases} \tag{9.121}$$

We can relate the dominant weights  $\gamma_i^c$  and the  $\bar{\gamma}_i$ : If  $n$  is even then

$$\gamma_\nu^c = \bar{\gamma}_\nu \text{ for } 1 \leq \nu < {}^\circ r - 1, \gamma_{{}^\circ r - 1}^c = \bar{\gamma}_{{}^\circ r - 1} - \frac{1}{2}\bar{\gamma}_{{}^\circ r}, \gamma_{{}^\circ r}^c = \frac{1}{2}\bar{\gamma}_{{}^\circ r} \tag{9.122}$$

For  $n$  odd we get

$$\gamma_\nu^c = \bar{\gamma}_\nu \text{ for } 1 \leq \nu < {}^\circ r, \gamma_{{}^\circ r}^c = \frac{1}{2}\bar{\gamma}_{{}^\circ r}$$

The Borel subgroup  $B$  is invariant under  $\Theta_1$ , the root subgroup  $U_{i,j}; 1 \leq i < l \leq n$  is conjugated into  $U_{n+1-j, n+1-i}$ . Inside the unipotent radical we have the half diagonal of spots  $({}^\circ r, {}^\circ r + 1 + 2\epsilon(n)), \dots, (2, n-1), (1, n)$  The involution is a reflection along this half diagonal and the spots on the left of the half diagonal form a system of representatives for  $\sim \Theta_1$ . Of course we have a corresponding Borel subgroup  $B_1 \supset T_1 \times \mathbb{C}$  of  $G \times \mathbb{C}$ .

**Proposition 9.6.2.** *Under the above identification the restrictions of the  $\gamma_{2i-1}^M$  to  $T_c$  are equal to the  $e_i$  in  $X^*(T_1^c \times \mathbb{C})$ .*

We want to compute  $\mu_c(\chi, \lambda)$ . By definition this is an element in  $X^*(T_c \times \mathbb{C})$  using the identification in (9.6.3) we carry out this computation in  $X^*(T_c)$ . A cocharacter  $\chi : \mathbb{G}_m \rightarrow T_c$  is of the form

$$\chi : t \mapsto \begin{pmatrix} t^{m_1} & 0 & 0 & \dots & \\ 0 & t^{m_2} & \dots & \dots & \\ 0 & 0 & \ddots & 0 & \dots \\ 0 & & \dots & t^{-m_2} & \\ 0 & & & & t^{-m_1} \end{pmatrix}$$

since we want  $P_\chi \supset B_1$  we require  $m_1 \geq m_2 \geq m_{{}^\circ r} \geq 0$ . (If  $n$  is odd then there is an  $m_{{}^\circ r + 1} = 0$ ). Let us start with the regular case, this means that all  $\geq$  signs are actually strict, i.e.  $>$  signs. Then it an easy computation that

$$\mu_c(\chi_{\text{reg}}, \lambda) = \begin{cases} ne_1 + (n-2)e_2 + \dots + 2e_{{}^\circ r} + \lambda_c & \text{if } n \text{ is even} \\ ne_1 + (n-2)e_2 + \dots + 3e_{{}^\circ r} + \lambda_c & \text{if } n \text{ is odd} \end{cases} \tag{9.123}$$

The set  $\Delta_{\chi_{\text{reg}}}^+$  is the set of roots of  $B$  modulo the conjugation  $\Theta_1$ . Hence we see that

$$\mu_c(\chi_{\text{reg}}, \lambda) = \mu_0^c(\lambda).$$

The interesting contribution is in fact  $\mu_c(\chi_{\text{reg}}, 0)$  and this is the number  $\mu_0^c$  in (9.111) We can express  $\mu_c(\chi_{\text{reg}}, 0)$  in terms of the fundamental weights  $\gamma_i$  (or the  $\bar{\gamma}_i$ ) we use the formulas (9.122). We get

$$\mu_c(\chi_{\text{reg}}, 0) = 2\bar{\gamma}_1 + 2\bar{\gamma}_2 + \cdots + 2\bar{\gamma}_{\circ r-1} + \begin{cases} 2\bar{\gamma}_{\circ r} & n \equiv 0 \pmod{2} \\ 6\bar{\gamma}_{\circ r} & n \equiv 1 \pmod{2} \end{cases} \quad (9.124)$$

If  $\chi$  is not regular then the relevant information extracted from  $\chi$  is the list

$$t_\chi = (t_1, t_2, \dots, t_s; t_0)$$

(the type of  $\chi$ ) where the  $t_i$  are the length of the intervals where the  $m_i > 0$  are constant, i.e.  $m_1 = m_2 = \cdots = m_{t_1} > m_{t_1+1} = \cdots = m_{t_1+t_2} > \dots$ . The number  $t_0$  is the length of the interval where  $m_i = 0$ . The  $\Theta$  stable parabolic  $P_\chi$  subgroup only depends on  $t_\chi$ . The types  $t_\chi$  have to satisfy the (only) constraint

$$2 \sum t_\nu + t_0 = n \quad (9.125)$$

The regular case corresponds to the list  $(1, 1, \dots, 1; 0$  or  $1)$ . In the general case we get a decorated Dynkin diagram where the crossed out roots are those where the  $m_i$  jump.

$$- \times - \circ - \circ - \circ - \circ - \times - \times - \circ - \dots - \circ - \times \cdots \times - \circ - \dots - \circ$$

This decorated diagram is symmetric under the reflection  $i \mapsto n - i$ . We look at the connected component of  $\circ$ -s. These components come in pairs unless the component is invariant under the reflection, i.e. it is central. The non central pairs

$$\pi_{\chi, \nu} = \begin{array}{ccccccc} \times - & \circ - & \dots - & \circ & - \times - \dots \times - & \circ & - \dots - & \circ \\ & \alpha_{i_\nu} & & \alpha_{j_\nu} & & \alpha_{n-j_\nu} & & \alpha_{n-i_\nu} \end{array} \quad (9.126)$$

are labelled by the indices  $\nu$  for which  $t_\nu > 1$ , and are of length  $t_\nu - 1 = j_\nu - i_\nu + 1$ . (The meaning of the indices  $i_\nu, j_\nu$  is explained in the diagram). The central connected component is of length  $t_0 - 1$ , of course it may be empty. We write it as

$$\pi_{\chi, 0} = \begin{array}{cccc} \times - & \circ & - \dots - & \circ & - \times \\ & \alpha_{i_0} & & \alpha_{j_0} & \end{array} \quad (9.127)$$

where of course  $i_0 = n - j_0$ . Let  $\pi_\chi$  be the union of these connected components. Let  $\Delta_\nu^+$  be the set of positive roots which are sums of roots in  $\pi_\nu$ .

To compute  $\mu_c(\chi, 0)$  we have to subtract from  $\mu_c(\chi_{\text{reg}}, 0)$  the sum of roots in  $\Delta_\nu^+$  with  $j_\nu <^\circ r$  and the sum of roots in  $\Delta_0^+ / \{\Theta_1\}$ .

A simple calculation shows that for  $\nu > 0$

$$2\rho^{(\nu)} = \sum_{i=i_\nu}^{i=j_\nu} \gamma_i + \gamma_{n-i} - (t_\nu - 1)(\gamma_{i_\nu-1} + \gamma_{i_\mu+1}) \quad (9.128)$$

where we put  $\gamma_{-1} = \gamma_n = 0$ . This means that subtracting  $2\rho^{(\nu)}$  from the sum which yields  $\mu_c(\chi_{\mathrm{reg}}, 0)$  has the effect that the sum  $\sum_{i=i_\nu}^{i=j_\nu} \gamma_i + \gamma_{n-i} = 2 \sum \bar{\gamma}_i$  cancels out and we have to add  $(t_\nu - 1)(\gamma_{i_\nu-1} + \gamma_{i_\mu+1})$ . Observe that  $i_{\nu-1}, j_{\mu+1} \notin \pi_\chi$ . We still have to subtract the contribution from the central component  $\Delta_0^+$ . We have to sum the roots in  $\Delta_0^+ / \{\Theta_1\}$  this means that we take half the sum of all roots and add half the sum of the symmetric roots. This yields

$$2\rho^{(0)} = \frac{1}{2}((j_0 - i_0 + 1)\alpha_{i_0} + \cdots + (j_0 - i_0 + 1)\alpha_{j_0}) + \frac{1}{2}(\alpha_{i_0} + \cdots + \cdots + \alpha_{j_0}) = ((j_0 - i_0 + 2)\bar{\alpha}_{i_0} + \cdots + (\dots)\bar{\alpha}_{o_r})$$

we see again that the sum  $\sum_{i=i_0}^{n-i_0} \bar{\gamma}_i$  drops out and we have to add a term  $t_0(\gamma_{i_0-1} + \gamma_{i_0+1})$ .

Hence we get: Let  $\pi_\chi^c$  be the union of the  $\pi_\nu^c$  and  $\pi_0^c$ . Then

$$\mu_c(\chi, 0) = \sum_{i \notin \pi_\chi^c} (2 + c_i(\chi, 0))\gamma_i^c$$

where

$$c_i(\chi, 0) = \begin{cases} (t_{\nu^-} - 1) + (t_{\nu^+} - 1) & \text{if } \nu \neq 0 \\ (t_{\nu^-} - 1) + t_{\nu^+} & \text{if } \nu = 0 \end{cases} \quad (9.129)$$

and where  $t_{\nu^-} - 1$  is the length of connected component directly to the left of  $i_\nu - 1$  and  $t_{\nu^+} - 1$  is the length of the component directly to the right of  $i_\nu - 1$ .

If we have chosen a highest weight  $\lambda = \sum a_i \gamma_i$  then we require  $a_i = a_{n+1-i} \geq 0$  and we must have  $a_i = 0$  for all  $i \in \pi_\chi$ . Then

$$\mu_c(\chi, \lambda) = \sum_{i \notin \pi_\chi} (2 + c_i(\chi, 0) + 2a_i)\gamma_i^c.$$

For us a special case is of interest. We decompose  $n = uv$  and take  $\chi_{u,v} = \chi$  of type  $t_\chi = (v, v, \dots, v)$ . Hence the reductive quotient of the  $\Theta$  stable parabolic subgroup is  $M^\vee = \mathrm{Gl}_v \times \mathrm{Gl}_v \times \cdots \times \mathrm{Gl}_v$ , the number of factors is  $u$ . In this case we get

$$\begin{array}{cccccccccccc} \circ & - & \circ & \cdots - & \circ - & \times & - \circ - & \circ - \cdots - & \circ - \times & - \circ \cdots \\ \alpha_1 & & \alpha_2 & & \alpha_{v-1} & \alpha_v & \alpha_{v+1} & & \alpha_{2v-1} & \alpha_{2v} & \end{array} \quad (9.130)$$

so the indices outside  $\pi_\chi$  are the multiples of  $v$ . Let us denote by  $\mathfrak{q}$  the Lie-algebra of  $P_{\chi_{u,v}}$ .

$$\mu_c(\chi_{u,v}, \lambda) = \sum_{\nu: \nu v \leq \frac{u}{2}} (2 + 2(v-1) + e(\nu))\gamma_{\nu v}^c + \lambda_c \quad (9.131)$$

where  $e(\nu) = 0$  except in the case that  ${}^\circ r \in [\nu v, (\nu+1)v]$  and then it is equal to 1.

### 9.6.4 The $A_{\mathfrak{q}_{u,v}}(\lambda)$ as Langlands quotients

Let  $n = uv$  and  $\mathfrak{q} = \mathfrak{q}_{u,v}$  as above. The parabolic is  $P_{\chi_{u,v}}$ . To realize  $A_{\mathfrak{q}_{u,v}}(\lambda)$  as Langlands quotient we apply the procedure described in [Vo-Zu], p.82-83. We have to find a parabolic subgroup  $P \subset \mathrm{Gl}_n/\mathbb{R}$  and a tempered representation  $\sigma_\infty$  of  $M = P/U$  such that

- a) our  $\lambda$  is a character on  $P$ ,
- b) the module  ${}^a\mathrm{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})}\sigma_\infty$  has the right infinitesimal character,
- c) the module  $\mathrm{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})}\sigma_\infty$  restricted to  $K_\infty$  contains  $\mu_c(\chi_{u,v}, \lambda_c)$  as minimal  $K_\infty$  type.

To get our parabolic subgroup we choose a cocharacter  $\eta_{u,v} : \mathbb{G}_m \rightarrow T$ , this cocharacter is defined as

$$t \mapsto \eta_{u,v}(t) = \begin{pmatrix} t^v & 0 & 0 & & \dots \\ 0 & t^{v-1} & \dots & \dots & \\ 0 & 0 & \ddots & 0 & \dots \\ 0 & & \dots & t^1 & \\ 0 & & & & t^v \\ 0 & & & & & t^{v-1} \\ & & & & & & \ddots \\ 0 & & & & & & & \ddots \end{pmatrix} \quad (9.132)$$

i.e. we have  $u$  copies of the diagonal matrix  $\mathrm{diag}(t^v, t^{v-1}, \dots, t)$  on the diagonal.

This cocharacter  $\eta = \eta_{u,v}(t)$  yields a parabolic subgroup  $P_\eta$  which contains the torus and has as roots  $\Delta_\eta = \{\alpha < \eta, \alpha \geq 0\}$ . Its reductive quotient is  $\mathrm{Gl}_u \times \mathrm{Gl}_u \times \dots \times \mathrm{Gl}_u$  where the number of factors is  $v$ . The embedding into  $\mathrm{Gl}_n$  is not the obvious one and  $P_\eta$  does not contain the standard Borel subgroup of upper triangular matrices.

To describe the relation between these two groups we denote by  $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$  the standard orthonormal basis of our underlying vector space  $\mathbb{R}^n$ . Then we group these basis elements

$$\{\{\mathbf{e}_1, \dots, \mathbf{e}_v\}, \{\mathbf{e}_{v+1}, \dots, \mathbf{e}_{2v}\}, \dots, \{\mathbf{e}_{(u-1)v+1}, \dots, \mathbf{e}_{uv}\}\}$$

and this grouping yields a direct sum decomposition

$$\begin{aligned} \mathbb{R}^n &= (\mathbb{R}\mathbf{e}_1 \oplus \dots \oplus \mathbb{R}\mathbf{e}_v) \oplus (\mathbb{R}\mathbf{e}_{v+1} \oplus \dots \oplus \mathbb{R}\mathbf{e}_{2v}) \oplus \dots \oplus (\mathbb{R}\mathbf{e}_{(u-1)v+1}, \dots, \mathbb{R}\mathbf{e}_{uv}) = \\ &V_1 \oplus V_2 \oplus \dots \oplus V_u \end{aligned} \quad (9.133)$$

and then  $M^\vee = \mathrm{Gl}(V_1) \times \dots \times \mathrm{Gl}(V_u)$ .

We get a second grouping of the basis elements

$$\{\{\mathbf{e}_1, \mathbf{e}_{v+1}, \dots, \mathbf{e}_{(u-1)v+1}\}, \{\mathbf{e}_2, \mathbf{e}_{v+2}, \dots, \mathbf{e}_{(u-1)v+2}\}, \dots\}, \{\dots, \mathbf{e}_{uv}\} \quad (9.134)$$

which yields direct sum decomposition

$$\begin{aligned} \mathbb{R}^n &= (\mathbb{R}\mathbf{e}_1 \oplus \mathbb{R}\mathbf{e}_{v+1} \oplus \dots \oplus \mathbb{R}\mathbf{e}_{(u-1)v+1}) \oplus (\mathbb{R}\mathbf{e}_2 \oplus \mathbb{R}\mathbf{e}_{v+2} \oplus \dots \oplus \mathbb{R}\mathbf{e}_{(u-1)v+2}) \oplus \dots \\ &W_1 \oplus W_2 \oplus \dots \oplus W_v \end{aligned} \quad (9.135)$$

and then  $M = \mathrm{Gl}(W_1) \times \mathrm{Gl}(W_2) \times \cdots \times \mathrm{Gl}(W_v) = \mathrm{Gl}_u \times \mathrm{Gl}_u \times \cdots \times \mathrm{Gl}_u$ . The groups  $M^\vee$  and  $M$  are mutual centralizers of each other.

The two groupings define two different Borel subgroups, the first one defines the standard Borel  $B$  of upper triangular matrices and the second Borel  $B^*$  fixes the flag  $\{\mathbf{e}_1\}, \{\mathbf{e}_1, \mathbf{e}_{v+1}\}, \dots$ . Let us denote by  $\lambda^*, \rho^*, w_{u,v}^*, \dots$  the dominant weight with respect to  $B^*$ , the half sum of positive roots and so on. Our highest weight  $\lambda$  is trivial on the semi simple part of  $M^\vee$  it must be of the form (9.70) Now we consider the highest weight for the group  $M$

$$\begin{aligned} w_{u,v}^*(\lambda^* + \rho^*) - \rho^* = \underline{\mu}^* &= (a_1 + v - 1)(\gamma_1^{*,M} + \gamma_{1+u}^{*,M} + \cdots + \gamma_{1+(v-1)u}^{*,M}) + \\ &\quad (a_2 + v - 1)(\gamma_2^{*,M} + \gamma_{2+u}^{*,M} + \cdots + \gamma_{2+(v-1)u}^{*,M}) + \\ &\quad \vdots \\ &\quad (a_{u-1} + v - 1)(\gamma_{u-1}^{*,M} + \gamma_{u-1+u}^{*,M} + \cdots + \gamma_{u-1+(v-1)u}^{*,M}) + \\ &\quad -(u+1)(\gamma_u^* + \gamma_{2u}^* + \cdots + \gamma_{(v-1)u}^*) + d\delta \quad . \end{aligned} \quad (9.136)$$

We choose  $\sigma_\infty = \mathbb{D}_{\underline{\mu}^*}$ . (See (9.66))

We check the lowest  $K_\infty$  type in  $\mathrm{Ind}_{P^*}^G \mathbb{D}_{\underline{\mu}^*}$ . To compute this lowest  $K_\infty$  type we write  $M = \prod M_\nu$  where of course each  $M_\nu = \mathrm{Gl}_u$ . Accordingly we write  $T = \prod T_\nu$ . The weight  $\mu^* = \sum \mu_\nu^*$  where the semi simple part is "always the same". We apply the considerations in section 9.6.1 to the factors  $M_\nu$ . We take  $\nu = 1$  then

$$\mu_1^* = (a_1 + v - 1)\gamma_1^* + (a_2 + v - 1)\gamma_1^* + \cdots + (a_{u-1} + v - 1)\gamma_{u-1}^* + d^* \det_u$$

Inside  $M_1$  we have the subgroup  ${}^\circ M_1$  which is the reductive Levi factor of  ${}^\circ P_1$  as in section 9.6.1 and we have the Kostant element  $w_{1,\mathrm{un}}$ . Then we consider the character

$$\tilde{\mu}_1^* = w_{1,\mathrm{un}}(\mu_1^* + \rho_1^*) - \rho_1^* = \sum_{i:i \text{ odd}} b_i^* \gamma_i^{{}^\circ M_1^{(1)}} + \tilde{\mu}_1^{*,\mathrm{ab}} \quad (9.137)$$

where again the  $b_i^*$  are the cuspidal parameters and they are given by

$$b_{2j-1}^* = v(u+1-2j) - 1 + \begin{cases} 2a_j + 2a_{j+1} + \cdots + 2a_{\frac{u}{2}-1} + a_{\frac{u}{2}} & \text{if } u \text{ is even} \\ 2a_j + 2a_{j+1} + \cdots + 2a_{\frac{u-1}{2}} & \text{if } u \text{ is odd} \end{cases} \quad (9.138)$$

The abelian part  $\tilde{\mu}_1^{*,\mathrm{ab}}$  does not play any role in the following (The  $\lambda$  in section (9.6.1) is now  $\mu_1^*$  and the  $\underline{mu}$  in formula (9.89) is now  $\tilde{\mu}_1^*$ ) We renumber our basis (9.134)

$$\{\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_{u-1}, \mathbf{f}_u, \dots\} = \{\mathbf{e}_1, \mathbf{e}_{v+1}, \dots, \mathbf{e}_{(u-1)v+1}, \mathbf{e}_2, \dots\} \quad (9.139)$$

and decompose the space  $\mathbb{R}^n$  into a direct sum of euclidian planes (plus a line if  $n$  is odd)

$$\mathbb{R}^n = (\mathbb{R}\mathbf{f}_1 \oplus \mathbb{R}\mathbf{f}_2) \oplus (\mathbb{R}\mathbf{f}_3 \oplus \mathbb{R}\mathbf{f}_4) \oplus \cdots \oplus (\mathbb{R}\mathbf{f}_n).$$

and this provides a maximal anisotropic torus

$$T_c^* = \mathrm{SO}(2) \times \mathrm{SO}(2) \times \cdots \times \mathrm{SO}(2)$$

In analogy with section 9.6.2 we write

$$X^*(T_c^* \otimes \mathbb{C}) = \bigoplus \mathbb{Z} f_j \quad (9.140)$$

where  $f_j$  is defined in analogy with the  $e_\nu$  in section 9.6.2.

We have

$$M = \mathrm{Gl}(\mathbb{R}f_1 \oplus \mathbb{R}f_2 \oplus \cdots \oplus \mathbb{R}f_u) \times \cdots \times \mathrm{Gl}(\mathbb{R}f_{(v-1)u+1} \oplus \cdots \oplus \mathbb{R}f_{uv})$$

and the intersection  $T_c^{*,M} = T_c^* \cap M$  is a maximal anisotropic torus in  $M$ . It is equal to  $T_c^*$  if  $u$  is even. If  $u$  is odd (and  $v > 1$ ) then it is a proper sub torus, if  $\circ r_u = \frac{u-1}{2}$  then

$$T_c^{*,M} = \underbrace{\mathrm{SO}(2) \times \cdots \times \mathrm{SO}(2)}_{\circ r_u \text{ factors}} \times \{\pm 1\} \times \underbrace{\mathrm{SO}(2) \times \cdots \times \mathrm{SO}(2)}_{\circ r_u \text{ factors}} \times \{\pm 1\} \times \text{spot } u \text{ and } u+1 \quad (9.141)$$

where the product of signs is one. To get the torus  $T_c^*$  we have to put another  $\mathrm{SO}(2)$  at the spots  $(u, u+1), (2u, 2u+1), \dots$ . We apply the reasoning of section (9.6.2) to the factors  $M_\nu$ .

The representation  $\mathbb{D}_{\mu_1^*} = \mathrm{Ind}_{P_\nu}^{M_1} \mathcal{D}_{\bar{\mu}_1^*}$  contains as lowest  $K_\infty^{M_\nu}$  type the representation with highest weight

$$(b_1^* + 2)f_1 + (b_3^* + 2)f_2 + \cdots + (b_{2 \circ r_u - 1}^* + 2)f_{\circ r_u}$$

where the  $b_{2j-1}^*$  are taken from (9.138). This weight occurs in  $\mathcal{D}_{\bar{\mu}_1^*}$ . Hence we see that as a  $T_c^*$  module the representation  $\otimes \mathcal{D}_{\bar{m}u^*}$  contains the weight (depending on  $u$  even or odd)

$$\left\{ \begin{aligned} &((b_1^* + 2)f_1 + (b_3^* + 2)f_2 + \cdots + (b_{2 \circ r_u - 1}^* + 2)f_{\circ r_u}) + ((b_1^* + 2)f_{\circ r_u + 1} + \cdots) + \cdots \\ &((b_1^* + 2)f_1 + (b_3^* + 2)f_2 + \cdots + (b_{2 \circ r_u - 1}^* + 2)f_{\circ r_u - 1}) + ((b_1^* + 2)f_{\circ r_u + 1} + \cdots) + \cdots \end{aligned} \right. \quad (9.142)$$

This weight is not dominant, to get a dominant weight we have to reorder the  $f_\nu$  according to the size of the coefficient in front. Then we get a dominant weight

$$(b_1^* + 2)(f_1^\dagger + f_2^\dagger + \cdots + f_v^\dagger) + (b_3^* + 2)(f_{v+1}^\dagger + f_{v+2}^\dagger + \cdots + f_{2v}^\dagger) + \cdots \quad (9.143)$$

and then formula (9.123) and the formula for the  $b_j^*$  give us the following dominant weight expressed in terms of the fundamental dominant weights

$$\sum_{\nu: \nu v \leq \frac{u}{2}} (2v + e(\nu) + 2a_\nu) \gamma_{\nu v}^c \quad (9.144)$$

This is now the weight  $\mu_c(\chi_{u,v}, \lambda)$  in (9.123). Hence we see that  $\Theta_{\mu_c(\chi_{u,v}, \lambda)}$  occurs with multiplicity one in  $\mathbb{D}_{\underline{mu}} : \mathrm{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})} \mathbb{D}_\mu$  and we get

**Theorem 9.6.1.** *We have a nonzero intertwining operator  $T^{(\mathrm{loc})}(\mathbb{D}_{\underline{mu}}) : \mathrm{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})}\mathbb{D}_{\underline{\mu}} \rightarrow \mathrm{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})}\mathbb{D}_{\underline{\mu}'}$  and get a diagram*

$$\begin{array}{ccc} \mathrm{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})}\mathbb{D}_{\underline{mu}} & \xrightarrow{T^{(\mathrm{loc})}(D_\mu)} & A_{\mathfrak{q}}(\lambda) \\ & & \downarrow \\ & & \mathrm{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})}\mathbb{D}_{\underline{\mu}'} \end{array} \quad (9.145)$$

The horizontal arrow is surjective, and the vertical arrow is injective. The map induced by the vertical arrow in cohomology

$$H^q(\mathfrak{g}, K_\infty; A_{\mathfrak{q}}(\lambda) \otimes \mathcal{M}_\lambda) \longrightarrow H^q(\mathfrak{g}, K_\infty; {}^a\mathrm{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})}\mathbb{D}_{\underline{\mu}'} \otimes \mathcal{M}_\lambda)$$

is a bijection in the lowest degree of nonzero cohomology; this lowest degree is

$$q = v \left[ \frac{u^2}{4} \right] + \frac{n(u-1)(v-1)}{4}.$$

*Proof.* We have an inclusion between the two complexes

$$\mathrm{Hom}_{K_\infty^0}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), A_{\mathfrak{q}}(\lambda) \otimes \mathcal{M}_\lambda) \rightarrow \mathrm{Hom}_{K_\infty^0}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \mathrm{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})}\mathbb{D}_{\underline{\mu}'} \otimes \mathcal{M}_\lambda).$$

In the complex on the left all differentials are zero. It follows from the work of Kostant that we have a splitting

$$\mathrm{Hom}(\Lambda^\bullet(\mathfrak{u}_P), \mathcal{M}_\lambda) = \mathbb{H}^\bullet(\mathfrak{u}_P, \mathcal{M}_\lambda) \oplus AC^\bullet$$

where  $\mathbb{H}^\bullet(\mathfrak{u}_P, \mathcal{M}_\lambda)$  is the space of harmonic forms (and this space is isomorphic to the cohomology  $H^\bullet(\mathfrak{u}_P, \mathcal{M}_\lambda)$ .) and where  $AC^\bullet$  is an acyclic complex.

We have Delorme's formula

$$\begin{aligned} \mathrm{Hom}_{K_\infty^0}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \mathrm{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})}\mathbb{D}_{\underline{\mu}'} \otimes \mathcal{M}_\lambda) &= \mathrm{Hom}_{K_\infty^M}(\Lambda^\bullet(\mathfrak{m}/\mathfrak{k}^M), \mathbb{D}_{\underline{\mu}'} \otimes \mathrm{Hom}(\Lambda^\bullet(\mathfrak{u}_P), \mathcal{M}_\lambda)) = \\ &= \mathrm{Hom}_{K_\infty^M}(\Lambda^\bullet(\mathfrak{m}/\mathfrak{k}^M), \mathbb{D}_{\underline{\mu}'} \otimes \mathbb{H}^\bullet(\mathfrak{u}_P, \mathcal{M}_\lambda)) \oplus \mathrm{Hom}_{K_\infty^M}(\Lambda^\bullet(\mathfrak{m}/\mathfrak{k}^M), \mathbb{D}_{\underline{\mu}'} \otimes AC^\bullet) \end{aligned} \quad (9.146)$$

The  $(\mathfrak{m}/K_\infty^M)$  has a lowest  $K_\infty^M$  type  $\vartheta(\mu')$ , which can be computed easily from 3.1.4 and we have

$$\mathrm{Hom}_{K_\infty^M}(\Lambda^\bullet(\mathfrak{m}/\mathfrak{k}^M), \mathbb{D}_{\underline{\mu}'} \otimes \mathbb{H}^\bullet(\mathfrak{u}_P, \mathcal{M}_\lambda)) = \mathrm{Hom}_{K_\infty^M}(\Lambda^\bullet(\mathfrak{m}/\mathfrak{k}^M), \mathbb{D}_{\underline{\mu}'}(\vartheta(\mu')) \otimes \mathbb{H}^\bullet(\mathfrak{u}_P, \mathcal{M}_\lambda)).$$

Using the formula in [Vo-Zu] for the highest weight of the lowest  $K_\infty$ -type  $\Theta(\mathfrak{q}, \lambda)$  in  $A_{\mathfrak{q}}(\lambda)$  we see that  $\Theta(\mathfrak{q}, \lambda)$  is the lowest  $K_\infty$  type in  $\mathrm{Ind}_{K_\infty^M}^{K_\infty}$ . This implies that the map

$$\mathrm{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), A_{\mathfrak{q}}(\lambda)(\Theta(\mathfrak{q}, \lambda) \otimes \mathcal{M}_\lambda) \rightarrow \mathrm{Hom}_{K_\infty^M}(\Lambda^\bullet(\mathfrak{m}/\mathfrak{k}^M), \mathbb{D}_{\underline{\mu}'} \otimes \mathbb{H}^\bullet(\mathfrak{u}_P, \mathcal{M}_\lambda)) \quad (9.147)$$

is an isomorphism of vector spaces (but not of complexes). But since the complex on the right is zero in degrees  $\bullet < q$  it follows that the classes in the image of  $\mathrm{Hom}_{K_\infty}(\Lambda^q(\mathfrak{g}/\mathfrak{k}), A_{\mathfrak{q}}(\lambda)(\Theta(\mathfrak{q}, \lambda) \otimes \mathcal{M}_\lambda)$  survive in cohomology.  $\square$

We got to the global context, we have a diagram

$$\begin{array}{ccc} J_{\sigma_\infty} \otimes J_{\sigma_f}^{K_f} & \hookrightarrow & L_{\text{disc}}^2(G(\mathbb{Q}) \backslash G(\mathbb{A}) / K_f, \omega_{\mathcal{M}_\lambda}^{-1} |_{S(\mathbb{R})^0}) \\ \downarrow & & \downarrow \mathcal{F}^P \\ {}^a\text{Ind}_{P(\mathbb{A})}^{G(\mathbb{A})} \mathbb{D}_{\mu'} \otimes V_{\sigma_f}^{K_f} & \hookrightarrow & \mathcal{A}(P(\mathbb{Q})U(\mathbb{A}) \backslash G(\mathbb{A}) / K_f) \end{array} \quad (9.148)$$

This induces maps in cohomology

$$\begin{array}{ccc} H^\bullet(\mathfrak{g}, K_\infty, J_{\sigma_\infty} \otimes \mathcal{M}_\lambda) \otimes J_{\sigma_f}^{K_f} & \rightarrow & H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \\ \downarrow & & \downarrow \mathcal{F}^P \\ H^\bullet(\mathfrak{g}, K_\infty, {}^a\text{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})} \mathbb{D}_{\mu'} \otimes \mathcal{M}_\lambda) \otimes V_{\sigma_f}^{K_f} & \hookrightarrow & H^\bullet(\partial_P \mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda) \end{array} \quad (9.149)$$

The left vertical arrow is non zero for  $\bullet = q$ , the horizontal arrow in the bottom line is injective for all values of  $\bullet$  (Borel see ) hence the horizontal arrow in the top line is non zero in degree  $\bullet = q$ .

Of course we also should investigate the horizontal arrow in the top line in all degrees, this question becomes delicate. To answer it we should invoke the results in Franke's paper [ ] or we could work with proposition (8.1.4) or its corollary (8.1.1).

In the extremal case  $u = n, v = 1$  the parabolic subgroup  $P$  is all of  $G$  and  $A_q(\lambda) = \mathbb{D}_\lambda$ . In this case, and only this case, the representation  $A_q(\lambda)$  is tempered.

In the other extremal case that  $u = 1, v = n$  the representation  $J_{\sigma_\infty}$  is one dimensional - (basically it is the space of constant functions twisted by a character on the group of connected components) - in this case the map in the top row is understood in terms of the topological model (Franke + Diploma students).

### 9.6.5 Congruences

We formulate a condition ( $NUQuot$ ) (No unitarizable quotient) for the induced module:

*The induced module  $I_P^G(\sigma_f)$  as module under the Hecke-algebra does not have a non trivial quotient which admits a unitary scalar product (here it may be necessary to pass to the corresponding representation of  $G(\mathbb{A}_f)$ ).*

The negation of this condition ( $UQuot$ ) says that for all primes  $p$  the induced module  $I_P^G \sigma_p$  has a unitarizable quotient.

This condition has been discussed in [Ha-Eis] Kap. II, 2.3.

If we have ( $NUQuot$ ) then

$$\text{Hom}_{\mathcal{H}_{K_f}^G}(I_P^G(\sigma_f), H_!^\bullet(\mathcal{S}_{K_f}^G, \mathcal{M} \otimes \mathbb{C})) = 0 \quad (9.150)$$

this implies that the Manin-Drinfeld is valid and this implies that our above section is defined over  $F$ , i.e. we get a unique section of Hecke-modules

$$\text{Eis} : H^{q-l(w)}(\mathcal{S}_{K_f^M}^M, \mathcal{M}(w \cdot \lambda) \otimes F)(\sigma_f) \rightarrow H^q(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F). \quad (9.151)$$

Then it looks as if the second term is completely uninteresting, but in fact it is not. In the lecture notes volume [Ha-Eis] we raise the question whether it influences the structure of the integral cohomology  $H_{\mathrm{int}}^q(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F)$ . In some cases we have convincing experimental evidence that "arithmetic" of the ratio of special values

$$\frac{1}{\Omega(\sigma_f)} \prod_a \frac{\Lambda^{\mathrm{coh}}(\sigma_f, r_a^{\check{u}}, \langle \eta_a, \tilde{\mu}^{(1)} \rangle - ab(w, \lambda))}{\Lambda^{\mathrm{coh}}(\sigma_f, r_a^{\check{u}}, \langle \eta_a, \tilde{\mu}^{(1)} \rangle - ab(w, \lambda) + 1)} \quad (9.152)$$

has influence on the structure integral of the cohomology. Under certain conditions the above expression is a product of an algebraic part and the value of a motivic extension class. Primes dividing the denominator of the algebraic part may occur in the denominator of the Eisenstein class and we will have congruences (See (8.2.3), (??)). This will be explained in the next section in the special case of the group  $\mathrm{GSp}_2/\mathbb{Z}$ .

### Attaching motives to $\sigma_f$ ???

The condition  $(NUQuot)$  will be true if  $\lambda$  is sufficiently regular but for non regular weights it fails. If the weight is not regular then we may have a pole of the Eisenstein series at  $z = 0$ . This possibility has to be discussed, it can only happen if we have  $(UQuot)$ . But even if we have  $(UQuot)$  we may not have a pole.

Let us assume that we have  $(UQuot)$  and the Eisenstein operator is holomorphic at  $z = 0$ . Then we may have several copies of  $J(\sigma_f)$  in  $H_!^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{C})$ . This defines again an isotypical submodule  $H_!^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F)(\bar{\sigma}_f)$ . We get an exact sequence

$$0 \rightarrow H_!^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F)(\bar{\sigma}_f) \rightarrow \mathcal{X}(\sigma_f) \rightarrow J(\sigma_f) \rightarrow 0 \quad (9.153)$$

This is a sequence of Hecke-modules over  $F$ , the section (9.42) provides a section over  $\mathbb{C}$ .

If our locally symmetric space  $\mathcal{S}_{K_f}^G$  the set of complex points of a Shimura variety then we can interpret this sequence as a mixed motive. This motive has an extension class in the category of mixed Hodge-structures

$$[\mathcal{X}(\sigma_f)]_{B-dRh} \in \mathrm{Ext}_{B-dRh}^1(J(\sigma_f), H_!^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F)(\bar{\sigma}_f)) \quad (9.154)$$

and in some cases we can compute this class (we have to look at a suitable bi-extension) and express it in terms of the second term in the constant term (See [MixMot-2013.pdf].)

We have seen that in many situations the space  $\mathcal{S}_{K_f^M}^M$  is not the set of complex points of a Shimura variety and therefore we do not know how to attach a motive or an  $\ell$  adic Galois representation to it. (Sometimes we know how to attach a motive to it but it is simply a Tate motive). But if it happens that the module  $J(\sigma_f)$  produces a non trivial submodule  $H_!^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F)(\bar{\sigma}_f)$  then the situation changes and we can attach a Galois-module  $H_!^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F_\lambda)(\bar{\sigma}_f)$  to it which contains a lot of information about  $\sigma_f$ . Again we refer to ([MixMot-2013.pdf].) We have seen in [Ha-Eis] (3.1.4.) that this can happen.

**The motivic interpretation of Shahidis theorem**

We go back to a general submodule  $\sigma_f = \sigma_f^{(1)} \times \sigma_f^{(2)} = \sigma_f \in \text{Coh}(H_{\text{cusp}}^\bullet(\mathcal{S}_{K_f^M}^M, \tilde{\mathcal{M}}_{w,\lambda}))$ ,

we drop the assumptions above. We assume that we can attach motives  $\mathbb{M}(\sigma_f^{(1)}, r_1), \mathbb{M}(\sigma_f^{(2)}, r_1)$  where  $r_1$  is the tautological representation. (Actually we do not need the motives it suffices to have the compatible systems of  $l$ -adic representations) Then we can attach the Rankin-Selberg motive to this pair

$$\mathbb{M}_{\text{RS}}(\sigma_f, \text{Ad}) = \mathbb{M}(\sigma_f^{(1)}, r_1) \times \mathbb{M}(\sigma_f^{(2)}, r_1)^\vee = \text{Hom}(\mathbb{M}(\sigma_f^{(2)}, r_1), \mathbb{M}(\sigma_f^{(1)}, r_1)) \otimes \mathbb{Z}(-\mathbf{w}(\mu^{(2)}, r_2)) \tag{9.155}$$

Under the assumption of the theorem the we have  $\mathbb{M}(\sigma_f^{(1)}, r_1) \xrightarrow{\sim} \mathbb{M}(\sigma_f^{(2)}, r_1)$  and we see that the Galois module  $\text{Hom}(\mathbb{M}(\sigma_f^{(2)}, r_1), \mathbb{M}(\sigma_f^{(1)}, r_1))$  contains a copy of  $\mathbb{Z}_l(0)$  and therefore we get an exact sequence of Galois modules

$$0 \rightarrow \mathbb{Z}(-\mathbf{w}(\mu^{(2)}, r_2)) \rightarrow \mathbb{M}_{\text{RS}}(\sigma_f, \text{Ad})_{\text{ét,Ad}} \rightarrow \mathbb{M}_{\text{RS}}^{(0)}(\sigma_f, \text{Ad})_{\text{ét,Ad}} \rightarrow 0$$

Hence the motivic  $L$  function is a product

$$L(\mathbb{M}_{\text{RS}}(\sigma_f, \text{Ad})_{\text{ét,Ad}}, s) = L(\mathbb{Z}(-\mathbf{w}(\mu^{(2)}, r_2)), s) L(\mathbb{M}_{\text{RS}}^{(0)}(\sigma_f, \text{Ad})_{\text{ét,Ad}}, s)$$

If we substitute for  $s$  the expression

$$\frac{\mathbf{w}(r_1, \mu_1^{(1)}) + \mathbf{w}(r_2, \mu_2^{(1)})}{2} - b(w, \lambda) + s = \mathbf{w}(r_2, \mu_2^{(1)}) - b(w, \lambda) + s$$

then we find

$$L(\mathbb{M}_{\text{RS}}(\sigma_f, \text{Ad})_{\text{ét,Ad}}, s) = \zeta(-b(w, \lambda) + s) L(\mathbb{M}_{\text{RS}}^{(0)}(\sigma_f, \text{Ad})_{\text{ét,Ad}}, s)$$

Then the motivic interpretation of Shahidis theorem is, that  $L(\mathbb{M}_{\text{RS}}^{(0)}(\sigma_f, \text{Ad})_{\text{ét,Ad}}, \mathbf{w}(r_2, \mu_2^{(1)}) - b(w, \lambda) + s)$  is holomorphic at  $s = 0$  and non zero (this is in a sense the prime number theorem for this  $L$  function) and therefore - if we have  $b(w, \lambda) = -1$  - the pole comes from the first order pole of the Riemann  $-\zeta$  function. If now  $\sigma_f^{(1)} \times \sigma_f^{(2)} = \sigma_f$  occurs in the cuspidal cohomology then we have an inclusion

$$\mathbb{D}_\mu \times H_{\sigma_f} \hookrightarrow \mathcal{A}(M(\mathbb{Q}) \backslash M(\mathbb{A}) / K_f^M)$$

We form the Eisenstein intertwining operator and compose it with constant Fourier coefficient, then we get

$$\mathcal{F}^P \circ \text{Eis}(s) : f \mapsto f + C(\sigma, s) T^{\text{loc}}(s)(f) \tag{9.156}$$

The operator  $T^{\text{loc}}(s) = T_\infty^{\text{loc}}(s) \otimes \bigotimes_p T_p^{\text{loc}}(s)$  is holomorphic at  $s = 0$ . Under our assumptions the function  $C(\sigma, s)$  has a first order pole at  $s = 0$  and we get a residual intertwining operator

$$\text{Res}_{s=0} : \text{Ind}_P^G \mathbb{D}_\mu \times H_{\sigma_f} \otimes (0) \rightarrow \mathcal{A}^{(2)}(G(\mathbb{Q}) \backslash G(\mathbb{A}) / K_f) \tag{9.157}$$

**Rationality results**

Finally we want to discuss the case that  $P \neq \Theta(P) = Q$ . If this happens then  $\mathcal{S}_{K_f}^G$  is never a Shimura variety. We have isotypical pieces (see (9.22) )

$$H_1^{\bullet-l(w)}(\mathcal{S}_{K_f^M}^M, \tilde{\mathcal{M}}(w \cdot \lambda) \otimes F)(\sigma_f) \oplus H_1^{\bullet-l(w')}( \mathcal{S}_{K_f^{M'}}^{M'}, \tilde{\mathcal{M}}(w' \cdot \lambda) \otimes F)(\sigma'_f) \tag{9.158}$$

and we know that component of the Eisenstein cohomology consists of the classes

$$\{ \psi_f \oplus \mathcal{L}(\sigma_f) T_f^{\mathrm{loc}}(\psi_f) \} \tag{9.159}$$

where  $\mathcal{L}(\sigma_f)$  is an element of  $F$  and for all  $\iota : F \rightarrow \mathbb{C}$  we have

$$\iota(\mathcal{L}(\sigma_f)) = \frac{1}{\Omega(\iota \circ \sigma_f)} C(\sigma_\infty, \lambda) C(\iota \circ \sigma_f, \lambda) \tag{9.160}$$

If the factor at infinity  $C(\sigma_\infty, \lambda) \neq 0$  then we get from this rationality results for the ratios of  $L$ -values. (See [32],[35]) These rationality results will be important when we discuss the arithmetic nature of the numbers in??

Combining the results of Borel–Garland [5] and Mœglin–Waldspurger [58] we get that the homomorphism

$$\bigoplus_{u|n} \bigoplus_{\sigma_f:\mathrm{segment}} H^\bullet(\mathfrak{g}, K_\infty; A_q(\lambda) \otimes \mathcal{M}_\lambda) \otimes J_{\sigma_f} \rightarrow H_{(2)}^\bullet(\mathcal{S}_{K_f}^G, \mathcal{M}_\lambda) \tag{9.161}$$

is surjective. This gives us the decomposition into isotypical spaces of  $H_{(2)}^\bullet(\mathcal{S}_{K_f}^G, \mathcal{M}_\lambda)$ . We separate the cuspidal part ( $v = 1$ ) from the residual part and get

$$H_{(2)}^\bullet(\mathcal{S}_{K_f}^G, \mathcal{M}_\lambda) = \bigoplus_{\pi_f:\mathrm{cuspidal}} H_{\mathrm{cusp}}^\bullet(\mathcal{S}_{K_f}^G, \mathcal{M}_\lambda)(\pi_f) \oplus \bigoplus_{\substack{u|n \\ u < n}} \bigoplus_{\sigma_f:\mathrm{segment}} \overline{H^\bullet(\mathfrak{g}, K_\infty; A_q(\lambda) \otimes \mathcal{M}_\lambda)} \otimes J_{\sigma_f},$$

where the bar on top means we have gone to its image via the map in (9.161). It follows from the theorem of Jacquet–Shalika [47] that there are no intertwining operators between the summands.


In the extremal case  $u = n, v = 1$  the parabolic subgroup  $P$  is all of  $G$  and  $A_q(\lambda) = \mathbb{D}_\lambda$ . In this case and only this case the representation  $A_q(\lambda)$  is tempered, and the lowest degree of nonvanishing cohomology is the number  $b_n^F$ . An easy computation shows that in the case  $v > 1$  the number  $q < b_n^F$ . Then our theorem above implies that in degree  $q$

$$H^q(\gamma, K_\infty; A_q(\lambda) \otimes \mathcal{M}_\lambda) \otimes J_{\sigma_f} \rightarrow H^q(\mathcal{S}_{K_f}^G, \mathcal{M}_\lambda)$$

is injective. This has also been proved by Grobner [20]. The above result, which we announced earlier (??), can be sharpened as in the following theorem. During the induction argument we use Thm. ?? for the reductive quotients  $M$  of the parabolic subgroups.

### 9.7 The example $G = \mathrm{Sp}_2/\mathbb{Z}$

#### 9.7.1 Some notations and structural data



The maximal torus is

$$T_0/\mathbb{Z} = t = \left\{ \begin{pmatrix} t_1 & 0 & 0 & 0 \\ 0 & t_2 & 0 & 0 \\ 0 & 0 & t_2^{-1} & 0 \\ 0 & 0 & 0 & t_1^{-1} \end{pmatrix} \right\}$$

the simple roots are

$$\alpha_1(t) = t_1/t_2, \alpha_2(t) = t_2^2$$

and the fundamental dominant weights are

$$\gamma_1(t) = t_1, \gamma_2(t) = t_1 t_2$$

and finally we have

$$2\gamma_1^M = t_1/t_2$$

We choose a highest weight  $\lambda = n_1\gamma_1 + n_2\gamma_2$  let  $\mathcal{M}_\lambda$  be a resulting module for  $G/\mathrm{Spec}(\mathbb{Z})$ . We get the following list of Kostant representatives for the Siegel parabolic subgroup  $P$  and they provide the following list of weights.

$$\begin{aligned} 1 \cdot \lambda &= \lambda = \frac{1}{2}(2n_2 + n_1)\gamma_2 + n_1\gamma_1^{M_1} \\ s_2 \cdot \lambda &= \frac{1}{2}(-2 + n_1)\gamma_2 + (2n_2 + n_1 + 2)\gamma_1^{M_1} \\ s_2 s_1 \cdot \lambda &= \frac{1}{2}(-4 - n_1)\gamma_2 + (2 + 2n_2 + n_1)\gamma_1^{M_1} \\ s_2 s_1 s_2 \cdot \lambda &= \frac{1}{2}(-6 - 2n_2 - n_1)\gamma_2 + n_1\gamma_1^{M_1}, \end{aligned}$$

We choose for  $K_\infty \subset \mathrm{Sp}_2(\mathbb{R})$  the standard maximal compact subgroup  $U(2)$ , it is the centralizer of the matrix

$$\begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}$$

which defines a complex structure. With this choice we can define  $\mathcal{S}_{K_f}^G = G(\mathbb{Q}) \backslash G(\mathbb{R}) / K_\infty \times G(\mathbb{A}_f) / K_f$ .

### 9.7.2 The cuspidal cohomology of the Siegel-stratum

We consider the cohomology groups  $H^\bullet(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda)$  and the resulting fundamental exact sequence. We have the boundary stratum  $\partial_P(\mathcal{S}_{K_f}^G)$  with respect to the Siegel parabolic. Let us assume that we are in the unramified case, then we get

$$H^\bullet(\partial_P(\mathcal{S}_{K_f}^G), \tilde{\mathcal{M}}_\lambda) = \bigoplus_{w \in W^P} H^{\bullet-l(w)}(\mathcal{S}_{K_f}^M, H^{l(w)}(\mathfrak{u}_P, \mathcal{M}_\lambda)) \quad (9.162)$$

We look at the case  $w = s_2 s_1$  in this case we know how to describe the corresponding summand in terms of automorphic forms on  $\mathrm{Gl}_2$ . We introduce the usual abbreviation  $H^{l(w)}(\mathfrak{u}_P, \mathcal{M}_\lambda) = \mathcal{M}_\lambda(w \cdot \lambda)$ .

Our coefficient modules are the modules attached to the highest weight

$$w \cdot \lambda = \mu = (2 + 2n_2 + n_1)\gamma_1^{M_1} + \frac{1}{2}(-4 - n_1)\gamma_2$$

Let us put  $k = 4 + 2n_2 + n_1$  and  $m = \frac{1}{2}n_1$ . We give the usual concrete realization for these modules as  $\mathcal{M}_{2+2n_2+n_1}[n_2 - 3 - k] = \mathcal{M}_{k-2}[n_2 - 3 - k]$

Let us look at the space  $\mathcal{S}_{K_f}^M$ . The group  $M/\mathrm{Spec}(\mathbb{Z})$  is isomorphic to  $\mathrm{Gl}_2$ , it is the Levi-quotient of the Siegel parabolic. The group  $K_\infty^M$  is the image of  $P(\mathbb{R}) \cap K_\infty$  under the projection  $P(\mathbb{R}) \rightarrow M(\mathbb{R})$ . This is the group  $\mathbb{O}(2)$  it contains the standard choice  $K_\infty^M(1) = \mathrm{SO}(2)$  as a subgroup of index 2. Hence we get a covering of degree 2

$$\mathcal{S}_{K_f}^{\tilde{M}} = M(\mathbb{Q}) \backslash M(\mathbb{R})/K_\infty^M(1) \times M(\mathbb{A}_f)/K_f^M \rightarrow \mathcal{S}_{K_f}^M \quad (9.163)$$

We get an inclusion

$$i : H^1(\mathcal{S}_{K_f}^M, \mathcal{M}_\lambda(w \cdot \lambda)) \hookrightarrow H^1(\mathcal{S}_{K_f}^{\tilde{M}}, \mathcal{M}_\lambda(w \cdot \lambda)). \quad (9.164)$$

On the cohomology on the right we have the action of  $\mathbb{O}(2)/\mathrm{SO}(2) = \mathbb{Z}/2\mathbb{Z}$  and the cohomology decomposes into a + and a - eigenspace. The inclusion  $i$  provides an isomorphism of the left hand side and the + eigenspace.

This inclusion is of course compatible with the action of the Hecke algebra. If we pass to a suitable extension  $F/\mathbb{Q}$  we get the decompositions into isotypic subspaces if we tensor our coefficient system by  $F$ . An isomorphism type  $\sigma_f$  occurs with multiplicity one on the left hand side and with multiplicity two on the right hand side. Over the ring  $\mathcal{O}_F$  the modules  $H_{\pm, \mathrm{int}}^1(\mathcal{S}_{K_f}^M, \mathcal{M}_\lambda(w \cdot \lambda)_F)(\sigma_f)$  are of rank one, hence we can find locally in the base  $\mathrm{Spec}(\mathcal{O}_F)$  an isomorphism

$$T^{\mathrm{arith}}(\sigma_f) : H_+^1(\mathcal{S}_{K_f}^{\tilde{M}}, \mathcal{M}_\lambda(w \cdot \lambda)_F)(\sigma_f) \xrightarrow{\sim} H_-^1(\mathcal{S}_{K_f}^{\tilde{M}}, \mathcal{M}_\lambda(w \cdot \lambda)_F)(\sigma_f) \quad (9.165)$$

The isomorphism given by the fundamental class (see(7.25) interchanges the + and the - eigenspace, hence we can arrange our arithmetic intertwining operator such that it satisfies

$$T^{\mathrm{arith}}(\sigma_f \otimes |\delta_f|) = T^{\mathrm{arith}}(\sigma_f \otimes |\delta_f|)^{-1} \quad (9.166)$$

We consider the transcendental description of the cohomology groups

$$H^1(\mathcal{S}_{K_f^M}^{\tilde{M}}, \mathcal{M}_\lambda(w \cdot \lambda)_{\mathbb{C}}) = \bigoplus_{\sigma_f} H_+^1(\mathcal{S}_{K_f^M}^{\tilde{M}}, \mathcal{M}_\lambda(w \cdot \lambda)_{\mathbb{C}})(\sigma_f) \oplus H_-^1(\mathcal{S}_{K_f^M}^{\tilde{M}}, \mathcal{M}_\lambda(w \cdot \lambda)_{\mathbb{C}})(\sigma_f) \quad (9.167)$$

We consider the standard Borel subgroup  $B \subset M$  the standard split torus  $T_0 \subset B$  it contains our torus  $Z_0$ . We define the character

$$\chi_\mu = (k, m+2) : B(\mathbb{R}) \rightarrow \mathbb{C}^\times, \chi(t) = \gamma_1^M(t)^k |\gamma_2|^{m+2}.$$

It yields the induced Harish-Chandra module  $I_{B(\mathbb{R})}^{M(\mathbb{R})} \chi_\mu$  : We consider the functions

$$f : M(\mathbb{R}) \rightarrow \mathbb{C}; f(bg) = \chi(b)f(g); f|T_1 \text{ is of finite type .}$$

This is in fact a  $(\mathfrak{m}, K_\infty^{M,0})$ -module, it contains the discrete representation  $\mathcal{D}_{\chi_\mu}$ . We have the decomposition

$$\mathcal{D}_{\chi_\mu} = \bigoplus_{\nu \equiv 0(2), |\nu| \geq k} F\phi_{\chi, \nu}$$

where

$$\phi_{\chi, \nu}(g) = \phi_{\chi, \nu}(b \begin{pmatrix} \cos(\phi) & \sin(\phi) \\ -\sin(\phi) & \cos(\phi) \end{pmatrix}) = \chi(b)e^{2\pi i \nu \phi}.$$

Of course  $K_\infty^{M,0} = T_1(\mathbb{R}) = \{e(\phi) = \begin{pmatrix} \cos(\phi) & \sin(\phi) \\ -\sin(\phi) & \cos(\phi) \end{pmatrix}\}$  and we can write  $e(\phi)^\nu = e^{2\pi i \nu \phi}$ .

We have the well known formula for the  $((\mathfrak{m}, K_\infty^{M,0})$  cohomology

$$H^1((\mathfrak{m}, K_\infty^{M,0}), \mathcal{D}_{\chi_\mu} \otimes \mathcal{M}_\lambda(w \cdot \lambda)) = \text{Hom}_{K_\infty^{M,0}}(\Lambda^1(\mathfrak{m}/\mathfrak{k}^M), \mathcal{D}_{\chi_\mu} \otimes \mathcal{M}_\lambda(w \cdot \lambda)) = \quad (9.168)$$

$$\mathbb{C}P_+^\vee \otimes \phi_{\chi, -k} \otimes v_{k-2} + \mathbb{C}P_-^\vee \otimes \phi_{\chi, k} \otimes v_{-k+2} = \mathbb{C}\omega_{k,m} + \mathbb{C}\bar{\omega}_{k,m} \quad (9.169)$$

Here  $v_{k-2} = (X + iY)^{k-2}$ , resp.  $v_{-k} = (X - iY)^{k-2}$  are two carefully chosen highest (resp. lowest) weight vectors with respect to the action of  $K_\infty^{M,0}$ . The elements  $P_\pm$  are the usual elements in  $\mathfrak{m}/\mathfrak{k}$ . We choose a model space  $H_{\sigma_f}$  for  $\sigma_f$  i.e. a free rank one  $\mathcal{O}_F$ -module on which the Hecke algebra acts by the homomorphism  $\sigma_f : \mathcal{H}_{K_f^M}^M \rightarrow \mathcal{O}_F$ . We also choose an embedding  $\iota : F \hookrightarrow \mathbb{C}$  and an  $(\mathfrak{m}, K_\infty^{M,0}) \times K_\infty^M \times \mathcal{H}_{K_f^M}^M$ -invariant embedding

$$\Phi_\iota : \mathcal{D}_{\chi_\mu} \otimes H_{\sigma_f} \otimes_{F, \iota} \mathbb{C} \rightarrow L_0^2(M(\mathbb{Q}) \backslash M(\mathbb{A})) \quad (9.170)$$

this is unique up to a scalar in  $\mathbb{C}^\times$  because the representation is irreducible and occurs with multiplicity one in the right hand side. This yields an isomorphism

$$\Phi_\iota^1 : H^1((\mathfrak{m}, K_\infty^{M,0}), \mathcal{D}_{\chi_\mu} \otimes \mathcal{M}_\lambda(w \cdot \lambda)) \otimes_{H_{\sigma_f} \otimes_{F, \iota} \mathbb{C}} \xrightarrow{\sim} H^1(\mathcal{S}_{K_f^M}^{\tilde{M}}, \mathcal{M}_\lambda(w \cdot \lambda)_{\mathbb{C}})(\iota \circ \sigma_f)$$

We observe that the element  $\epsilon = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \in K_\infty^M$  has the following effect

$$\mathrm{Ad}(\epsilon)(P_+) = P_-, \epsilon(\phi_{\chi,k}) = \phi_{\chi,-k} \quad \text{and} \quad \epsilon(v_{k-2}) = (-1)^m v_{2-k} \quad (9.171)$$

Hence we see that

$$\omega_{k,m}^{(+)} = \omega_{k,m} + (-1)^m \bar{\omega}_{k,m} \quad \text{resp.} \quad \omega_{k,m}^{(-)} = \omega_{k,m} - (-1)^m \bar{\omega}_{k,m} \quad (9.172)$$

are generators of the  $+$  and the  $-$  eigenspace in  $H^1(\mathfrak{m}, K_\infty^{M,0}, \mathcal{D}_{\chi_\mu} \otimes \mathcal{M}_\lambda(w \cdot \lambda))$ . Therefore our map  $\Phi$  and the choice of these generators provide isomorphisms

$$\Phi_\iota^{(+)} : H_{\sigma_f} \otimes_{F,\iota} \mathbb{C} \xrightarrow{\sim} H_+^1(\mathcal{S}_{K_f^M}^{\tilde{M}}, \mathcal{M}_\lambda(w \cdot \lambda)_\mathbb{C})(\iota \circ \sigma_f), \quad (9.173)$$

$$\Phi_\iota^{(-)} : H_{\sigma_f} \otimes_{F,\iota} \mathbb{C} \xrightarrow{\sim} H_-^1(\mathcal{S}_{K_f^M}^{\tilde{M}}, \mathcal{M}_\lambda(w \cdot \lambda)_\mathbb{C})(\iota \circ \sigma_f) \quad (9.174)$$

The choice of  $P_+, P_-$  and  $\phi_{\chi,-\nu}$  is canonic, hence we see that the identifications depend only on  $\Phi_\iota$ , which is unique up to a scalar. This means that the composition

$$\begin{aligned} T^{\mathrm{trans}}(\iota \circ \sigma_f) &= \Phi_\iota^{(-)} \circ (\Phi_\iota^{(+)})^{-1} \\ &: H_+^1(\mathcal{S}_{K_f^M}^{\tilde{M}}, \mathcal{M}_\lambda(w \cdot \lambda)_\mathbb{C})(\iota \circ \sigma_f) \xrightarrow{\sim} H_-^1(\mathcal{S}_{K_f^M}^{\tilde{M}}, \mathcal{M}_\lambda(w \cdot \lambda)_\mathbb{C})(\iota \circ \sigma_f) \end{aligned}$$

yields a second (canonical) identification between the  $\pm$  eigenspaces in the cohomology. Our arithmetic intertwining operator (See (9.165)) yields an array of intertwining operators

$$T^{\mathrm{arith}}(\sigma_f) \otimes_{F,\iota} \mathbb{C} : H_+^1(\mathcal{S}_{K_f^M}^{\tilde{M}}, \mathcal{M}_\lambda(w \cdot \lambda)_F)(\sigma_f) \otimes_{F,\iota} \mathbb{C} \xrightarrow{\sim} H_-^1(\mathcal{S}_{K_f^M}^{\tilde{M}}, \mathcal{M}_\lambda(w \cdot \lambda)_F)(\sigma_f) \otimes_{F,\iota} \mathbb{C} \quad (9.175)$$

Hence get an array of periods which compare these two arrays of intertwining operators

$$\Omega(\sigma_f, \iota) T^{\mathrm{trans}}(\iota \circ \sigma_f) = T^{\mathrm{arith}}(\sigma_f) \otimes_{F,\iota} \mathbb{C} \quad (9.176)$$

Our formula (9.166) tells us that we can arrange the intertwining operators such that

$$\Omega(\sigma_f \otimes |\delta_f|, \iota) = \Omega(\sigma_f, \iota)^{-1} \quad (9.177)$$

These periods are uniquely defined up to a unit in  $\mathcal{O}_F^\times$ .

### The Eisenstein intertwining

We pick a  $\sigma_f$  which occurs in  $H_+^1(\mathcal{S}_{K_f^M}^{\tilde{M}}, \mathcal{M}_\lambda(w \cdot \lambda)_F)$ , we choose a  $\iota : F \hookrightarrow \mathbb{C}$  and we choose an embedding

$$\Phi_\iota : \mathcal{D}_{\chi_\mu} \otimes H_{\sigma_f} \otimes_{F,\iota} \mathbb{C} \hookrightarrow L_{\mathrm{cusp}}^2(M(\mathbb{Q}) \backslash M(\mathbb{A})) \quad (9.178)$$

and from this we get the Eisenstein intertwining

$$\mathrm{Eis} \circ \Phi_\iota : \mathrm{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})}(\mathcal{D}_{\chi_\mu}) \otimes H_{\sigma_f} \otimes_{F,\iota} \mathbb{C} \rightarrow \mathcal{A}(G(\mathbb{Q}) \backslash G(\mathbb{A})) \quad (9.179)$$

(Here we use that  $K_f = \mathrm{GSp}_2(\hat{\mathbb{Z}})$ .) Hence we get an intertwining operator

$$\mathrm{Eis}^\bullet : \mathrm{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), I_P^G(\mathcal{D}_{\chi_\mu}) \otimes \mathcal{M}_\lambda) \otimes H_{\sigma_f} \otimes_{F,\iota} \mathbb{C} \rightarrow \mathrm{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \mathcal{A}(G(\mathbb{Q}) \backslash G(\mathbb{A})) \otimes \mathcal{M}_\lambda) \quad (9.180)$$

and this induces a homomorphism in cohomology

$$H^3(\mathfrak{g}, K_\infty, I_P^G(\mathcal{D}_{\chi_\mu}) \otimes \mathcal{M}_\lambda) \otimes H_{\sigma_f} \otimes_{F,\iota} \mathbb{C} \rightarrow H^3(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_{\lambda,\mathbb{C}}) \quad (9.181)$$

and we want to compose it with the restriction to the cohomology of the boundary. We have to compose it with the constant Fourier coefficient  $\mathcal{F}^P : \mathcal{A}(G(\mathbb{Q}) \backslash G(\mathbb{A})) \rightarrow \mathcal{A}(P(\mathbb{Q})U(\mathbb{A}) \backslash G(\mathbb{A}))$ . We know that  $\mathcal{F}^P$  maps into the subspace

$$I_P^G \mathcal{D}_{\chi_\mu} \otimes H_{\sigma_f} \otimes_{F,\iota} \mathbb{C} \xrightarrow{\mathcal{F}^P} I_P^G \mathcal{D}_{\chi_\mu} \otimes H_{\sigma_f} \otimes_{F,\iota} \mathbb{C} \bigoplus I_P^G \mathcal{D}_{\chi_{\mu'}} \otimes H_{\sigma_f^{w_P} |\gamma_{P,f}|^{2f_P}} \otimes_{F,\iota} \mathbb{C} \quad (9.182)$$

where  $\mu' = w_P w \cdot \lambda = s_2 \cdot \lambda = (2 + 2n_2 + n_1)\gamma_1^{M_1} + \frac{1}{2}(-2 + n_1)\gamma_2$ . More precisely we know that for  $h \in I_P^G \mathcal{D}_{\chi_\mu} \otimes H_{\sigma_f} \otimes_{F,\iota} \mathbb{C}$

$$\mathcal{F}^P(h) = h + C(\sigma, 0)T^{\mathrm{loc}}(0)(h) \quad (9.183)$$

where  $T^{\mathrm{loc}}(0) = T_\infty^{\mathrm{loc}} \otimes \otimes_p T_p^{\mathrm{loc}}$ . The local intertwining operator at the finite primes is normalized, it maps the standard spherical function into the standard spherical function. The operator  $T_\infty^{\mathrm{loc}}$  will be discussed below.

Our general formula for the constant term yields for an  $h = h_\infty \times h_f$

**Explain in more detail**

$$\mathcal{F}^P(h) = h + C(\sigma_\infty, \lambda)T_\infty^{\mathrm{loc}}(h_\infty) \frac{L^{\mathrm{coh}}(f, n_1 + n_2 + 2) \zeta(n_1 + 1)}{L^{\mathrm{coh}}(f, n_1 + n_2 + 3) \zeta(n_1 + 2)} \times T_f^{\mathrm{loc}}(0)(h_f) \quad (9.184)$$

(For the following compare SecOps.pdf) We analyze the factor  $C(\sigma_\infty, \lambda)T_\infty^{\mathrm{loc}}$  more precisely we study the effect of this operator on the cohomology. Let us look at the map between complexes

$$T_\infty^{\mathrm{loc}, \bullet} : \mathrm{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), I_P^G \mathcal{D}_{\chi_\mu} \otimes \mathcal{M}_\lambda) \rightarrow \mathrm{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), I_P^G \mathcal{D}_{\chi_{\mu'}} \otimes \mathcal{M}_\lambda) \quad (9.185)$$

The intertwining operator  $T_\infty^{\mathrm{loc}} : I_P^G \mathcal{D}_{\chi_\mu} \rightarrow I_P^G \mathcal{D}_{\chi_{\mu'}}$  has a kernel  $\mathbb{D}_{\chi_\mu}$ , this is a discrete series representation. We know that

$$\mathrm{Hom}_{K_\infty}(\Lambda^\bullet(\mathfrak{g}/\mathfrak{k}), \mathbb{D}_{\chi_\mu} \otimes \mathcal{M}_\lambda) = \mathrm{Hom}_{K_\infty}(\Lambda^3(\mathfrak{g}/\mathfrak{k}), \mathbb{D}_{\chi_\mu} \otimes \mathcal{M}_\lambda) = \quad (9.186)$$

$$H^3(\mathfrak{g}, K_\infty, \mathbb{D}_{\chi_\mu} \otimes \mathcal{M}_\lambda) = \mathbb{C}\Omega_{2,1} \oplus \mathbb{C}\Omega_{1,2} \quad (9.187)$$

We have the surjective homomorphism

$$H^3(\mathfrak{g}, K_\infty, \mathbb{D}_{\chi_\mu} \otimes \mathcal{M}_\lambda) \rightarrow H^3(\Lambda^3(\mathfrak{g}/\mathfrak{k}), I_P^G \mathcal{D}_{\chi_\mu} \otimes \mathcal{M}_\lambda) = H^1(\mathfrak{m}, K_\infty^M, \mathcal{D}_{\chi_\mu} \otimes H^2(\mathfrak{u}_P, \mathcal{M}_\lambda)) = \mathbb{C}\omega^{(3)} \quad (9.188)$$

the differential form  $\Omega_{2,1} + \epsilon(\lambda)\Omega_{1,2}$  maps to a non zero multiple  $A(\lambda)\omega^{(3)}$ . (The factor  $\epsilon(\lambda)$  is a sign depending on  $\lambda$ ). We can write  $\Omega_{2,1} - \epsilon(\lambda)\Omega_{1,2} = d\psi$  where

$$\psi \in \mathrm{Hom}_{K_\infty}(\Lambda^2(\mathfrak{g}/\mathfrak{k}), \mathbb{D}_{\chi_\mu} \otimes \mathcal{M}_\lambda) \quad (9.189)$$

and  $\omega = T_\infty^{\mathrm{loc},2}(\psi) \in \mathrm{Hom}_{K_\infty}(\Lambda^2(\mathfrak{g}/\mathfrak{k}), \mathcal{D}_{\chi_{\mu'}} \otimes \mathcal{M}_\lambda)$  is a closed form, hence it provides a cohomology class. Let us denote this cohomology class by  $\kappa(\omega^{(3)})$ .

Choosing  $\omega^{(3)}$  as a basis element and applying the Eisenstein intertwining operator (9.180) yields a homomorphism

$$\mathrm{Eis}^{(3)} \circ \Phi_\iota : H_!^1(\mathcal{S}_{K_f^M}^{\tilde{M}}, \mathcal{M}_\lambda(w \cdot \lambda)_F)(\sigma_f \circ \iota) \rightarrow H^3(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes \mathbb{C}) \quad (9.190)$$

The local intertwining operator  $T_\infty^{\mathrm{loc}}$  maps  $\omega^{(3)}$  to zero and hence it follows that the composition  $r \circ \mathrm{Eis}^{(3)}$  is the identity, the Eisenstein intertwining operator yields a section on  $H_!^1(\mathcal{S}_{K_f^M}^{\tilde{M}}, \mathcal{M}_\lambda(w \cdot \lambda)_F)(\sigma_f)$ . (Remember  $w = s_2s_1$ ). If we define

$$H^3(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F)(\sigma_f) = r^{-1}(H_!^1(\mathcal{S}_{K_f^M}^{\tilde{M}}, \mathcal{M}_\lambda(w \cdot \lambda)_F)(\sigma_f)) \quad (9.191)$$

(Induction does not play a role since the level is one) then we get the decomposition

$$H_!^3(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F) \oplus H_{\mathrm{Eis}}^3(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F)(\sigma_f) = H_!^3(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F)(\sigma_f) \quad (9.192)$$

### The denominator of the Eisenstein class

We restrict this decomposition to the integral cohomology (better the image of the integral cohomology in the cohomology with rational coefficients)

$$H_{\mathrm{int}}^3(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F)(\sigma_f) \supset H_{\mathrm{int}}^3(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F)(\sigma_f) \oplus H_{\mathrm{int}, \mathrm{Eis}}^3(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F)(\sigma_f) \quad (9.193)$$

The image of  $H_{\mathrm{int}, \mathrm{Eis}}^3(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F)(\sigma_f)$  under  $r$  is a submodule of finite index in  $H_{\mathrm{int}}^1(\mathcal{S}_{K_f^M}^{\tilde{M}}, \mathcal{M}_\lambda(w \cdot \lambda)_F)(\sigma_f)$  and the quotient is

$$H_{\mathrm{int}}^3(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F)(\sigma_f) / (H_{\mathrm{int}}^3(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F)(\sigma_f) \oplus H_{\mathrm{int}, \mathrm{Eis}}^3(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F)(\sigma_f)) = H_{\mathrm{int}}^1(\mathcal{S}_{K_f^M}^{\tilde{M}}, \mathcal{M}_\lambda(w \cdot \lambda)_F)(\sigma_f) / \mathrm{image}(r). \quad (9.194)$$

The quotient on the right hand side is  $\mathcal{O}_F/\Delta(\sigma_f)$  where  $\Delta(\sigma_f)$  is the denominator ideal. Tensoring the exact sequence

$$0 \rightarrow H_{\mathrm{int}}^3(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F)(\sigma_f) \oplus H_{\mathrm{int}, \mathrm{Eis}}^3(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F)(\sigma_f) \rightarrow H_{\mathrm{int}}^1(\mathcal{S}_{K_f^M}^{\tilde{M}}, \mathcal{M}_\lambda(w \cdot \lambda)_F)(\sigma_f) \rightarrow \mathcal{O}_F/\Delta(\sigma_f) \rightarrow 0 \quad (9.195)$$

by  $\mathcal{O}_F/\Delta(\sigma_f)$  yields an inclusion

$$\mathrm{Tor}_{\mathcal{O}_F}^1(\mathcal{O}_F/\Delta(\sigma_f), \mathcal{O}_F/\Delta(\sigma_f) = \mathcal{O}_F/\Delta(\sigma_f)) \hookrightarrow H_{\mathrm{int}}^3(\mathcal{S}_{K_f}^G, \tilde{\mathcal{M}}_\lambda \otimes F)(\sigma_f) \otimes \mathcal{O}_F/\Delta(\sigma_f) \quad (9.196)$$

and this explains the congruences.

**The secondary class**

We choose generators  $\omega^{(3)}(\sigma_f)$  ( resp.  $\omega^{(2)}(\sigma_f^{w_P}|\gamma_{P,f}|^{2f_P})$ ) for  $H_{\text{int}}^1(\mathcal{S}_{K_f}^{\tilde{M}}, \mathcal{M}_\lambda(w \cdot \lambda)_F)$  ( resp.  $H_{\text{int}}^1(\mathcal{S}_{K_f}^{\tilde{M}}, \mathcal{M}_\lambda(s_2 \cdot \lambda))(\sigma_f)$ ) (Perhaps we can do this only locally on  $\text{Spec}(\mathcal{O}_F)$ .) We may arrange these generators such that  $T^{\text{arith}}(\sigma_f)(\omega^{(3)}(\sigma_f)) = \omega^{(2)}(\sigma_f^{w_P}|\gamma_{P,f}|^{2f_P})$ . The isomorphism

$$\Phi_\iota^{(3)} : H^3(\mathfrak{g}, K_\infty, \mathcal{D}_{\chi_\mu} \otimes \mathcal{M}_\lambda) \otimes H_{\sigma_f} \otimes_{F,\iota} \mathbb{C} \xrightarrow{\sim} H_{\text{int}}^1(\mathcal{S}_{K_f}^{\tilde{M}}, \mathcal{M}_\lambda(w \cdot \lambda)_F)(\iota \circ \sigma_f) \quad (9.197)$$

maps

$$(\Omega_{2,1} + \epsilon(\lambda)\Omega_{1,2}) \otimes \omega^{(3)}(\iota \circ \sigma_f) \mapsto \Omega_+(\sigma_f, \iota)\omega(\sigma_f)$$

where  $\Omega_+(\sigma_f, \iota)$  is a period depending on the choice of  $\Phi_\iota$ . The element

$$(\Omega_{2,1} - \epsilon(\lambda)\Omega_{1,2}) \otimes \omega^{(3)}(\iota \circ \sigma_f) = d\psi \otimes \omega^{(3)}(\iota \circ \sigma_f).$$

where  $\psi \in \text{Hom}_{K_\infty}(\Lambda^2(\mathfrak{g}/\mathfrak{k}), I_P^G \mathcal{D}_{\chi_\mu} \otimes \mathcal{M}_\lambda)$ . The operator  $T^{\text{loc}}(0)$  in (9.183) provides a homomorphism (9.185)

$$T^{\text{loc},2} \otimes T_f^{\text{loc}} : \text{Hom}_{K_\infty}(\Lambda^2(\mathfrak{g}/\mathfrak{k}), I_P^G \mathcal{D}_{\chi_\mu} \otimes \mathcal{M}_\lambda) \otimes H_{\iota \circ \sigma_f} \rightarrow \text{Hom}_{K_\infty}(\Lambda^2(\mathfrak{g}/\mathfrak{k}), I_P^G \mathcal{D}_{\chi_{\mu'}} \otimes \mathcal{M}_\lambda) \otimes H_{\iota \circ \sigma_f^{w_P}|\gamma_P}$$

Under this homomorphism the class  $\psi$  is mapped to a multiple of  $\omega^{(2)}(\sigma_f^{w_P}|\gamma_{P,f}|^{2f_P})$ .

We can calculate this multiple, during this calculation we see a second period  $\Omega_-(\sigma_f, \iota)$  depending on  $\Phi_\iota$  and the ratio of these periods will be our period  $\Omega(\iota \circ \sigma_f)$  in formula (9.176).

This period is independent of  $\Phi_\iota$ . To state the final result we denote by  $f$  the modular cusp form attached to  $\sigma_f$ , this is a modular form with coefficients in  $F$ , then  $\iota \circ f$  is a modular form with coefficients in  $\mathbb{C}$ . By  $\Lambda(f, s)$  we denote the usual completed  $L$ -function. We get

$$C(\sigma, 0)T^{\text{loc}}(\kappa(\omega^{(3)}(\iota \circ \sigma_f))) = \left( \frac{1}{\Omega(\sigma_f, \iota)^{\epsilon(k,m)}} \frac{\Lambda^{\text{coh}}(\iota \circ f, n_1 + n_2 + 2)}{\Lambda^{\text{coh}}(\iota \circ f, n_1 + n_2 + 3)} \frac{1}{\zeta(-1 - n_1)} \right) \frac{\zeta'(-n_1)}{\pi} \omega^{(2)}(\sigma_f^{w_P}|\gamma_{P,f}|^{2f_P})$$

The factor inside the large brackets is essentially rational ( it is in  $F$  and behaves invariantly under the action of the Galois group) the factor  $\frac{\zeta'(-n_1)}{\pi}$  should be viewed as a generator of a group of extension classes of mixed motives.

For me the most difficult part in the calculation is the treatment of the intertwining operator at  $\infty$ , this is carried out in SecOps.pdf. At the end of SecOps.pdf. I discuss the arithmetic applications and the conjectural relationship between the primes dividing the denominator of the expression in the large brackets and the denominators of the Eisenstein classes in (??)

# Bibliography

- [1] BATEMAN MANUSCRIPT PROJECT
- [2] J. Bewersdorff *Eine Lefschetzsche Fixpunktformel für Hecke-Operatoren. (German) [A Lefschetz fixed point formula for Hecke operators]* Dissertation, University of Bonn, Bonn, 1985. Bonner Mathematische Schriften 164. Universität Bonn, Mathematisches Institut, Bonn, 1985
- [3] A. Borel, *Regularization theorems in Lie algebra cohomology. Applications.* Duke Math. J. 50 (1983), no. 3, 605–623.
- [4] A. Borel and W. Casselman, *L<sup>2</sup>-cohomology of locally symmetric manifolds of finite volume.* Duke Math. J. 50 (1983), no. 3, 625–647.
- [5] A. Borel and H. Garland, *Laplacian and the discrete spectrum of an arithmetic group.* Amer. J. Math. 105 (1983), no. 2, 309–335.
- [6] A. Borel and N. Wallach, *Continuous cohomology, discrete subgroups, and representations of reductive groups.* Second edition. Mathematical Surveys and Monographs, 67. American Mathematical Society, Providence, RI, 2000.
- [7] A. Borel and J.-P. Serre, *Corners and arithmetic groups,* *Commen. Math. Helvetici* 48 (1973), 436–491.
- [8] N. Bourbaki, *Groupes et algèbres de Lie* Chapitres 4,5, et 6 *Elements de Mathematiques*
- [9] W. Casselman *On Some Results of Atkin and Lehner* *Math. Ann.* 201,301–3t4 (1973)
- [10] G. Chenevier, J. Lannes *Automorphic Forms and Even Unimodular Lattices* *Ergebnisse der Mathematik und ihrer Grenzgebiete* Springer 2019
- [11] L. Clozel, *Motifs et formes automorphes: applications du principe de fonctorialité.* *Automorphic forms, Shimura varieties, and L-functions, Vol. I* (Ann Arbor, MI, 1988), 77–159, *Perspect. Math.*, 10, Academic Press, Boston, MA, 1990.
- [12] P. Deligne, *Formes modulaires et représentations l-adiques.* (French) [Modular forms and  $\ell$ -adic representations] *Séminaire Bourbaki. Vol. 1968/69: Exposés 347-363, Exp. No. 355, 139-172, Lecture Notes in Math.*, 175, Springer, Berlin, 1971

- [13] P. Deligne, *Valeurs de fonctions  $L$  et périodes d'intégrales (French)*, With an appendix by N. Koblitz and A. Ogus. Proc. Sympos. Pure Math., XXXIII, Automorphic forms, representations and  $L$ -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Oregon, 1977), Part 2, pp. 313–346, Amer. Math. Soc., Providence, R.I., 1979.
- [14] T. Deng *Torsions in Cohomology of  $Sl_2(\mathbb{Z})$  and Congruence of Modular Forms* arXiv:1905.05260
- [15] Elstrodt, J., Grunewald, F., Mennicke, J. *Groups Acting on Hyperbolic Space* Springer Monographs in Mathematics
- [16] Faber, Carel; van der Geer, Gerard: *Sur la cohomologie des systèmes locaux sur les espaces de modules des courbes de genre 2 et des surfaces abéliennes. I*, C. R. Math. Acad. Sci. Paris 338 (2004), no. 5, 381–384. II, no. 6, 467–470
- [17] Franke, Jens *Harmonic analysis in weighted  $L^2$ -spaces*. Ann. Sci. École Norm. Sup. (4) 31 (1998), no. 2, 181–279.
- [18] Godement, R. *Lecture Notes on Jacquet-Langlands* IAS Princeton
- [19] M. Goresky, ; R. MacPherson, *The topological trace formula*. J. Reine Angew. Math. 560 (2003), 77-150
- [20] H. Grobner, *A cohomological injectivity result for the residual automorphic spectrum of  $GL(n)$* . Pacific J. Math., 268 (2014), no. 1, 33–46.
- [21] K. H. H. Haberland, *Perioden von Modulformen einer Variabler and Gruppencohomologie. I, II, III* Math. Nachr. 112 (1983), 245-282, 283-295, 297-315.
- [22] G. Harder, *A Gauss-Bonnet formula for discrete arithmetically defined groups* Annales scientifiques de l'É.N.S. 4e série, tome 4, no 3 (1971), p. 409-455
- [23] Harder, G., *General aspects in the theory of modular symbols*. Seminarberichte des Séminaire DPP, Birkhäuser, 73 - 88, 1983
- [24] G. Harder, *Eisenstein cohomology of arithmetic groups. The case  $GL_2$* . Invent. Math. 89, no. 1, 37–118 (1987).
- [25] Harder, G. *Eisensteinkohomologie und die Konstruktion gemischter Motive*, Springer Lecture Notes 1562
- [26] G. Harder, *Some results on the Eisenstein cohomology of arithmetic subgroups of  $GL_n$* , in Cohomology of arithmetic groups and automorphic forms (ed. J.-P. Labesse and J. Schwermer), Springer Lecture Notes in Mathematics, vol. 1447, pp. 85–153 (1990).
- [27] G. Harder- N. Schappacher *Special values of Hecke  $L$ -functions and abelian integrals*. Workshop Bonn 1984 (Bonn, 1984), 17–49, Lecture Notes in Math., 1111, Springer, Berlin, 1985.

- [28] G. Harder, *Lectures on Algebraic Geometry I* Aspects of Mathematics, E 35 viewegteubner
- [29] G. Harder, *Lectures on Algebraic Geometry II* Aspects of Mathematics, E 35 viewegteubner
- [30] G. Harder, *Interpolating coefficient systems and  $p$ -ordinary cohomology of arithmetic groups*. Groups Geom. Dyn., 5 (2011), no. 2, 393–444.
- [31] G. Harder *A congruence between a Siegel and an Elliptic Modular Form* The 1-2-3 of Modular Forms, Springer Unitext
- [32] G. Harder *Arithmetic Aspects of Rank One Eisenstein Cohomology* Proceedings of the International Colloquium on Cycles, Motives and Shimura Varieties, Mumbai 2008
- [33] G. Harder, *Arithmetic Harish-Chandra modules*. Preprint available at <http://arxiv.org/abs/1407.0574>
- [34] G. Harder *The topological trace formula* [www.math.uni-bonn.de/harder/Manuscripts/Eisenstein/chap5.pdf](http://www.math.uni-bonn.de/harder/Manuscripts/Eisenstein/chap5.pdf)
- [35] G. Harder and A. Raghuram, *Eisenstein cohomology and ratios of critical values of Rankin–Selberg  $L$ -functions*. C. R. Math. Acad. Sci. Paris 349 (2011), no. 13-14, 719–724.
- [36] G. Harder *Eisenstein cohomology for  $Sl_2(\mathbb{Z}[i])$  and special values of  $L$ -functions*. Cohomology of arithmetic groups, 51-82, Springer Proc. Math. Stat., 245, Springer, Cham, 2018.
- [37] Harish-Chandra, *Automorphic Forms on Semisimple Lie Groups* Springer Lecture Notes in Mathematics 62, 1968
- [38] E. Hecke . *Über Modulfunktionen und die Dirichletschen Reihen mit Eulerscher Produktentwicklung. I.* (German) Math. Ann. 114 (1937), no. 1, 1?28
- [39] S. Helgason, *Differential geometry and symmetric spaces*. Pure and Applied Mathematics, Vol. XII. Academic Press, New York-London 1962
- [40] Hida, Haruzo *Congruence of cusp forms and special values of their zeta functions*. Invent. Math. 63 (1981), no. 2, 225-261.
- [41] H. Jacquet, *Principal  $L$ -functions of the linear group*. Automorphic forms, representations and  $L$ -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2, pp. 63-86, Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979.
- [42] H. Jacquet, *On the residual spectrum of  $GL(n)$* . Lie group representations, II (College Park, Md., 1982/1983), 185-208, Lecture Notes in Math., 1041, Springer, Berlin, 1984.

- [43] Jacquet- Langlands, *Automorphic forms on  $Gl_2$* , Springer Lecture Notes 114
- [44] H. Jacquet and J.A. Shalika, *On Euler products and the classification of automorphic representations. I.* Amer. J. Math. 103 (1981), no. 3, 499–558.
- [45] H. Kasten and C.-G. Schmidt *The critical values of Rankin-Selberg convolutions.* Int. J. Number Theory 9 (2013), no. 1, 205–256
- [46] D. Kazhdan; B. Mazur and C.-G. Schmidt *Relative modular symbols and Rankin-Selberg convolutions.* J. Reine Angew. Math. 519 (2000), 97–141, p. 97-141
- [47] H. Jacquet and J.A. Shalika, *On Euler products and the classification of automorphic forms. II.* Amer. J. Math. 103 (1981), no. 4, 777–815.
- [48] H. Kim, *Automorphic  $L$ -functions.* Lectures on automorphic  $L$ -functions, 97–201, Fields Inst. Monogr., 20, Amer. Math. Soc., Providence, RI, 2004.
- [49] B. Kostant, *Lie algebra cohomology and the generalized Borel-Weil theorem* Annals of Math., Vol 74, No. 2 (1961) 329–387.
- [50] A. Knapp, *Local Langlands correspondence: the Archimedean case*, Motives (Seattle, WA, 1991), 393–410, Proc. Sympos. Pure Math., 55, Part 2, Amer. Math. Soc., Providence, RI, 1994.
- [51] Kneser, Martin *Strong approximation in Algebraic Groups and Discontinuous Subgroups* (Proc. Sympos. Pure Math., Boulder, Colo., 1965) pp. 187-196 Amer. Math. Soc., Providence, R.I.
- [52] J.-P. Labesse, R. Langlands,  *$L$ -indistinguishability for  $SL(2)$* , Canadian J. Math. 31 (1979), pp. 726–785
- [53] R. Langlands, *Euler products*, Yale University Press (1971).
- [54] R. Langlands, *On the functional equations satisfied by Eisenstein series.* Lecture Notes in Mathematics, Vol. 544., Springer-Verlag, Berlin-New York, 1976.
- [55] J.-S. Li- J. Schwermer, *On the Eisenstein cohomology of arithmetic groups.* Duke Math. J. 123 (2004), no. 1, 141-169.
- [56] R. Langlands *Automorphic representations, Shimura varieties, and motives. Ein Märchen.* Automorphic forms, representations and  $L$ -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2, pp. 205-246, Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979
- [57] J.-S. Li-, J. Schwermer, *On the Eisenstein cohomology of arithmetic groups.* Duke Math. J. 123 (2004), no. 1, 141-169.
- [58] C. Mœglin and J.-L. Waldspurger, *Le spectre résiduel de  $GL(n)$ .* (French) [The residual spectrum of  $GL(n)$ ] Ann. Sci. École Norm. Sup. (4) 22 (1989), no. 4, 605–674.

- [59] G. Nebe, B. Venkov, *On Siegel modular forms of weight 12*, J. reine angew. Math. 351 (2001), pp. 49-60.
- [60] J. Neukirch *Algebraic Number Theory* Grundlehren der mathematischen Wissenschaften, Springer 1999
- [61] H.-V. Niemeier, *Definite quadratische Formen der Dimension 24 und Diskriminante 1*. J. Number Theory 5 (1973), 142-178
- [62] Pink, R. *On  $l$ -adic sheaves on Shimura varieties and their higher direct images in the Baily-Borel compactification*. Math. Ann. 292 (1992), no. 2, 197-240.
- [63] Platonov, V. P. (1969), *The problem of strong approximation and the Kneser-Tits hypothesis for algebraic groups* Izvestiya Akademii Nauk SSSR. Seriya Matematicheskaya, 33: 1211-1219
- [64] De-Rham, G, *Differentiable manifolds. Forms, currents, harmonic forms* Springer Grundlehren, 266, 1984
- [65] Ribet, Kenneth A. *A modular construction of unramified  $p$ -extensions of  $\mathbb{Q}(\mu_p)$* . Inv. Math. 34 (3): 151-162
- [66]  *$\mathcal{L}$ -modules and the conjecture of Rapoport and Goresky-MacPherson*. Automorphic forms. I. Astérisque No. 298 (2005)
- [67] Satake, I. *Spherical functions and Ramanujan conjecture*. Algebraic Groups and Discontinuous Subgroups (Proc. Sympos. Pure Math., Boulder, Colo., 1965) pp. 258-264 Amer. Math. Soc., Providence, R.I.
- [68] Satake, I. *The Gauss-Bonnet theorem for  $V$ -manifolds*. J. Math. Soc. Japan 9 (1957), 464-492.
- [69] Scholl, A., *Motives for modular forms*. Invent. Math. 100 (1990), no. 2, 419-430.
- [70] J. Schwermer, *Kohomologie arithmetisch definierter Gruppen und Eisensteinreihen*. (German) [Cohomology of arithmetically defined groups and Eisenstein series] Lecture Notes in Mathematics, 988. Springer-Verlag, Berlin, 1983. iv+170 pp.
- [71] J. Schwermer, *Cohomology of arithmetic groups, automorphic forms and  $L$ -functions*. Cohomology of arithmetic groups and automorphic forms (Luminy-Marseille, 1989), 1-29, Lecture Notes in Math., 1447, Springer, Berlin, 1990.
- [72] J. Schwermer, *On Euler products and residual Eisenstein cohomology classes for Siegel modular varieties*. (English. English summary) Forum Math. 7 (1995), no. 1, 1-28.
- [73] Serre, Jean-Pierre, *A Course in Arithmetic*, Graduate Texts in Mathematics, 7, Springer-Verlag

- [74] J.-P. Serre, *Facteurs locaux des fonctions zeta des variétés algébriques (définitions et conjectures)*, in Séminaire Delange-Pisot-Poitou, 1969/70, Collected Papers II, vol. 19, p. 581–592.
- [75] F. Shahidi, *Whittaker models for real groups*. Duke Math. J. 47 (1980), no. 1, 99–125.
- [76] F. Shahidi, *Local coefficients as Artin factors for real groups*. Duke Math. J. 52 (1985), no. 4, 973–1007.
- [77] F. Shahidi, *Eisenstein series and automorphic L-functions*. American Mathematical Society Colloquium Publications, 58. American Mathematical Society, Providence, RI, (2010).
- [78] G. Shimura, *The special values of the zeta functions associated with Hilbert modular forms*. Duke Math. J. 45 (1978), no. 3, 637–679.
- [79] C.L. Siegel *Gesammelte Werke*
- [80] T.A. Springer *Linear Algebraic Groups* Modern Birkhäuser Classics
- [81] R. Steinberg *Lectures on Chevalley Groups* Yale University 1967
- [82] H.P.F. Swinnerton-Dyer *On  $\ell$ -adic representations and congruences for coefficients of modular forms* In Modular Functions of One Variable III, SLN 350
- [83] Tate, J. T. *Fourier analysis in number fields, and Hecke's zeta-functions*. Algebraic Number Theory (Proc. Instructional Conf., Brighton, 1965), 305-347, Thompson, Washington, D.C., 1967.
- [84] D. Vogan and G. Zuckerman, *Unitary representations with nonzero cohomology*. Comp. Math. 53 (1984), no. 1, 51–90.
- [85] R. Weissauer, *Endoscopy for  $\mathrm{GSp}(4)$  and the Cohomology of Siegel Modular Threefolds* Springer Lecture Notes 1968 (2009)
- [86] U. Weselmann, *Siegel Modular Varieties and the Eisenstein Cohomology of  $\mathrm{PGL}_{2g+1}$* , Manuscripta Math. 145 (2014), no. 1-2, 175–220.
- [87] A. Wiles *Modular elliptic curves and Fermat's last theorem*. Ann. of Math. (2) 141 (1995), no. 3, 443–551.
- [88] E. Witt *Über eine Identität zwischen Modulformen zweiten Grades* Gesammelte Abhandlungen, Springer Verlag

## References

[G-J] Gelbart, S.-Jacquet, H, A relation between automorphic representations of  $\mathrm{GL}(2)$  and  $\mathrm{GL}(3)$ . Ann. Sci. cole Norm. Sup. (4) 11 (1978), no. 4, 471–542

The following items can be obtained from my home page [www.math.uni-bonn.de/harder/Manuscripts/Eisenstein](http://www.math.uni-bonn.de/harder/Manuscripts/Eisenstein)

[MixMot-2015.pdf] Modular Construction of Mixed Motives

[SecOps.pdf] Secondary Operations on the Cohomology of Harish-Chandra Modules

[