

ON INTEGRAL LOCAL SHIMURA VARIETIES

GEORGIOS PAPPAS AND MICHAEL RAPOPORT

ABSTRACT. We give a construction of *integral local Shimura varieties* which are formal schemes that generalize the well-known integral models of the Drinfeld p -adic upper half spaces. The construction applies to all classical groups, at least for odd p . These formal schemes also generalize the formal schemes defined by Rapoport-Zink via moduli of p -divisible groups, and are characterized purely in group-theoretic terms.

More precisely, for a local p -adic Shimura datum (G, b, μ) and a quasi-parahoric group scheme \mathcal{G} for G , Scholze has defined a functor on perfectoid spaces which parametrizes p -adic shtukas. He conjectured that this functor is representable by a normal formal scheme which is locally formally of finite type and flat over $O_{\mathbb{E}}$. Scholze-Weinstein proved this conjecture when (G, b, μ) is of (P)EL type by using Rapoport-Zink formal schemes. We prove this conjecture for any (G, μ) of abelian type when $p \neq 2$, and when $p = 2$ and G is of type A or C . We also relate the generic fiber of this formal scheme to the local Shimura variety, a rigid-analytic space attached by Scholze to (G, b, μ, \mathcal{G}) .

CONTENTS

1. Introduction	1
2. Statements of the main results	5
3. Preliminaries	10
4. Parahoric vs Quasi-parahoric group schemes	17
5. Ad-isomorphisms and integral moduli spaces of local shtukas	28
6. The case of trivial μ_{ad}	42
7. The crucial Hodge type case	42
8. Existence of Hodge embeddings	48
9. Proofs of Theorems 2.5.5 and 2.5.4	51
References	54

1. INTRODUCTION

Recall the mechanism of Shimura varieties: to a Shimura datum (G, X) (here the first entry is a reductive group over \mathbb{Q}), there is associated a tower of algebraic varieties $(\text{Sh}_K(G, X) \mid K \subset G(\mathbb{A}_f))$ over the reflex field $E = E(G, X)$. Here, K runs through the open compact subgroups of $G(\mathbb{A}_f)$. Furthermore, at least if (G, X) is of PEL-type, there exist integral p -adic models if K is of the form $K = K^p K_p$, where K_p satisfies some additional conditions. More precisely, fix a prime number p and a p -adic place v of E . Then if $K_p \subset G(\mathbb{Q}_p)$ is the stabilizer of a point in the extended Bruhat-Tits building of $G_{\mathbb{Q}_p}$, there exists in the PEL-type case

an integral model $\mathcal{S}_K(G, X)$ over $O_{E,v}$. The study of these integral models has been the focus of interest for many years, with spectacular applications in arithmetic. Furthermore, in recent years such models have been constructed even for Shimura varieties of abelian type, comp. [KP18], [KZ21], see [Pa18]. In the present paper, we are concerned with a local p -adic analogue of this mechanism.

We fix a prime number p . There is the notion of a local Shimura datum (G, b, μ) [RV14], a p -adic analogue of the notion of a (global) Shimura datum. Here G is a reductive group over \mathbb{Q}_p , and $b \in G(\check{\mathbb{Q}}_p)$ and μ is a conjugacy class of minuscule cocharacters of G . Let $E = E(G, \mu)$ be the local reflex field. In [RV14], it is postulated that there should be an associated tower of rigid-analytic varieties $(\text{Sht}_K(G, b, \mu) \mid K \subset G(\mathbb{Q}_p))$ over the completion \check{E} of the maximal unramified extension of E . This idea was put into reality by P. Scholze, see [Sch18]. He defined functors on the category Perfd_k of perfectoid spaces over the residue field k of \check{E} and showed that they are representable by rigid-analytic spaces. Furthermore, if \mathcal{G} is a smooth model of G over \mathbb{Z}_p , Scholze defines a functor $\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}$ on Perfd_k which he shows to be a v -sheaf. Let \mathcal{G} be a quasi-parahoric group scheme, in the sense of [SW20, §21.5]. This is a natural class of smooth group scheme models over \mathbb{Z}_p of G . In the analogous function field case, T. Richarz [Ri16] has proved that quasi-parahoric group schemes can be characterized as those smooth group scheme models of G for which the corresponding affine Grassmannian is ind-proper. Scholze has conjectured that $\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}$ is representable by a formal scheme $\mathcal{M}_{\mathcal{G}, b, \mu}$ which is normal and flat and locally formally of finite type over $O_{\check{E}}$. Assuming this conjecture, Scholze-Weinstein [SW20] show that when \mathcal{G} is a parahoric group scheme for G and $K = \mathcal{G}(\mathbb{Z}_p)$, then $\mathcal{M}_{\mathcal{G}, b, \mu}$ is an integral model of $\text{Sht}_K(G, b, \mu)$ where $K = \mathcal{G}(\mathbb{Z}_p)$, i.e., the rigid-analytic generic fiber $\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{rig}}$ of $\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}$ can be identified with $\text{Sht}_K(G, b, \mu)$. In [SW20] the representability conjecture is proved when (G, b, μ) is of EL-type, and in many cases when (G, b, μ) is of PEL-type, by relating the functor $\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}$ to Rapoport-Zink formal schemes. The rigid-analytic variety $\text{Sht}_K(G, b, \mu)$ is called the *local Shimura variety* (associated to the local Shimura datum (G, b, μ) and the open compact subgroup K of $G(\mathbb{Q}_p)$) and the functor $\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}$, resp. the formal scheme $\mathcal{M}_{\mathcal{G}, b, \mu}$ the *integral local Shimura variety* (for the quasi-parahoric group scheme \mathcal{G} for G).

It is interesting to observe that the “classical” approach of [RZ96] proceeds in the reverse way compared to the approach in [SW20]. Namely, when (G, b, μ) arises from (P)EL-data, then for certain quasi-parahorics \mathcal{G} for G one constructs a formal scheme $\mathcal{M}_{\mathcal{G}, b, \mu}$ by posing a certain moduli problem of p -divisible groups with additional structure on the category $\text{Nilp}_{O_{\check{E}}}$. Then $\text{Sht}_{\mathcal{G}(\mathbb{Z}_p)}(G, b, \mu)$ is *defined* to be the generic fiber of $\mathcal{M}_{\mathcal{G}, b, \mu}$ and the rest of the tower $\text{Sht}_K(G, b, \mu)$ is constructed by imposing level K structures on the p -adic Tate module of the generic fiber of the universal p -divisible group over $\mathcal{M}_{\mathcal{G}, b, \mu}$. This definition depends a priori on the choice of (P)EL data and is not obviously functorial in the triple (\mathcal{G}, b, μ) . On the other hand, this approach has the dividend that the structure of the formal schemes $\mathcal{M}_{\mathcal{G}, b, \mu}$ can be studied by making use of the theory of p -divisible groups. The model for such a structure result is Drinfeld’s description of his integral model of the Drinfeld p -adic halfspace in [Dr77]. This approach has been used in a number of problems of arithmetic, e.g. in applications to the Zhang Arithmetic Fundamental Lemma conjecture [Zha12] and the Arithmetic Transfer conjecture [RSZ18] and to the Kudla-Rapoport Divisor Intersection conjecture [KR11]. The Rapoport-Zink approach has been generalized to certain Hodge type

cases by W. Kim [Kim18], Howard-Pappas [HP17], Hamacher-Kim [HK19], and to some abelian type cases by Shen [Sh20]. Also, there is a purely group-theoretical definition by Bültel-Pappas [BP20] of the moduli problem on $\mathrm{Nilp}_{O_{\bar{E}}}$ for $\mathcal{M}_{\mathcal{G},b,\mu}$ for hyperspecial parahoric group schemes \mathcal{G} . In the Hodge type case, it is shown in [BP20] that this moduli problem, restricted to Noetherian test rings, is representable by a formal scheme. It coincides with the formal schemes in [Kim18], [HP17] and [HK19].

The Scholze-Weinstein approach has the advantage that the input data are purely group-theoretical and that the result is functorial. In this way, they are able in certain cases to identify two RZ formal schemes for different (P)EL data which define closely related group-theoretical data. For instance, using this approach they prove the conjectures of Rapoport-Zink [RZ17] and of Kudla-Rapoport-Zink [KRZ20] which postulated such hidden identifications, cf. [SW20, §25.4-25.5]. The downside of this approach is that the global structure of the formal schemes $\mathcal{M}_{\mathcal{G},b,\mu}$ is harder to study.

We note that one expects a precise relation between integral local Shimura varieties and formal completions of global Shimura varieties along isogeny loci of their reduction modulo p . This is provided by the theory of non-archimedean uniformization in the Rapoport-Zink framework; something analogous is conjectured to hold in the general Scholze-Weinstein context, and is known in a number of cases, cf. [PR21, Thm. 1.3.3].

In this paper, we are concerned with passing from the PEL case to the more general case when the local Shimura datum (G, b, μ) is of abelian type. Here the definition of this last term is modeled on the case of (global) Shimura varieties, cf. [HPR20], [HLR18]. Namely, (G, b, μ) is of abelian type if the associated adjoint local Shimura datum $(G_{\mathrm{ad}}, b_{\mathrm{ad}}, \mu_{\mathrm{ad}})$ comes from a local Shimura datum (G_1, b_1, μ_1) of Hodge type (i.e., (G_1, μ_1) admits an embedding into (GL_n, μ_d) , where μ_d is a minuscule coweight of GL_n). In particular, G_{ad} is a classical group. We prove that Scholze's conjecture holds true when (G, b, μ) is of abelian type and either $p \neq 2$ or $p = 2$ and G_{ad} is of type A or C . We allow here general quasi-parahoric group schemes, even those outside the class singled out in [SW20, §25.3] (those for which the group $\Pi_{\mathcal{G}}$ below is trivial). Allowing general quasi-parahorics is important for two related reasons: moduli problems leading to Rapoport-Zink spaces often correspond to quasi-parahorics, and allowing quasi-parahorics is important for devissage in the proofs.

We also show that $\mathcal{M}_{\mathcal{G},b,\mu}$ is an integral model of $\mathrm{Sht}_K(G, b, \mu)$, in the following sense. Let

$$\Pi_{\mathcal{G}} = \ker(\mathrm{H}_{\mathrm{\acute{e}t}}^1(\mathbb{Z}_p, \mathcal{G}) \longrightarrow \mathrm{H}_{\mathrm{\acute{e}t}}^1(\mathbb{Q}_p, G)), \quad (1.0.1)$$

a finite abelian group, cf. [SW20, 25.3]. To every $\bar{\beta} \in \Pi_{\mathcal{G}}$, we associate a quasi-parahoric group scheme $\mathcal{G}_{\bar{\beta}}$ for G over \mathbb{Z}_p such that $\check{K}_{\bar{\beta}} = \mathcal{G}_{\bar{\beta}}(\check{\mathbb{Z}}_p)$ is conjugate to $\check{K} = \mathcal{G}(\check{\mathbb{Z}}_p)$ in $G(\check{\mathbb{Q}}_p)$, and we prove that

$$\mathcal{M}_{\mathcal{G},b,\mu}^{\mathrm{rig}} \simeq \bigsqcup_{\bar{\beta} \in \Pi_{\mathcal{G}}} \mathrm{Sht}_{K_{\bar{\beta}}}(G, b, \mu). \quad (1.0.2)$$

This formula is reminiscent of the formula of Kottwitz [Ko92], according to which the generic fiber of a PEL-moduli scheme in [Ko92] is a disjoint sum of copies of Shimura varieties enumerated by $\ker(\mathrm{H}^1(\mathbb{Q}, G) \rightarrow \prod_v \mathrm{H}^1(\mathbb{Q}_v, G))$ (these copies are mutually isomorphic in types A and C , but not necessarily in type D).

In fact, we prove that the decomposition (1.0.2) comes by passing to the generic fiber of a decomposition of functors

$$\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}} \simeq \bigsqcup_{\bar{\beta} \in \Pi_{\mathcal{G}}} \mathcal{M}_{\mathcal{G}_{\bar{\beta}}^o,b,\mu}^{\text{int}} / \pi_0(\mathcal{G}_{\bar{\beta}})^{\phi}. \quad (1.0.3)$$

Here $\mathcal{G}_{\bar{\beta}}^o$ denotes the parahoric group scheme associated to the quasi-parahoric group scheme $\mathcal{G}_{\bar{\beta}}$ (the neutral connected component). This formula also gives a reduction of Scholze's conjecture from quasi-parahoric group schemes to parahoric group schemes.

We know quite a bit about the local structure of the formal scheme $\mathcal{M}_{\mathcal{G},b,\mu}$. Indeed, let $\mathbb{M}_{\mathcal{G}^o,\mu}^{\text{loc}}$ be the local model associated to the parahoric group scheme \mathcal{G}^o associated to \mathcal{G} . Here the associated v -sheaf on Perfd_k is defined in [SW20, §21] and the representability by a weakly normal scheme $\mathbb{M}_{\mathcal{G}^o,\mu}^{\text{loc}}$ flat over $O_{\bar{E}}$ is established in the case of a general local Shimura datum by J. Anschütz, I. Gleason, J. Lourenço, T. Richarz in [AGLR22]. In fact, $\mathbb{M}_{\mathcal{G}^o,\mu}^{\text{loc}}$ is always normal with reduced special fiber (by [AGLR22] and [GLo22] which settled some remaining cases for $p = 2$ and $p = 3$). When $p \neq 2$, this local model also coincides for local Shimura data of abelian type with the local model in the style of Pappas and Zhu [PZ13] (modified in [HPR20]), as extended to groups which arise by restriction of scalars from wild extensions by B. Levin [Le01]. When $p = 2$ and G_{ad} is of type A or C , this local model also coincides with the local model obtained by taking the closure of the generic fiber in the naive local model of [RZ96]. We also know that, if $p \neq 2$, $\mathbb{M}_{\mathcal{G},\mu}^{\text{loc}}$ is Cohen-Macaulay [HR19]. Then we prove under our assumptions above that for every $x \in \mathcal{M}_{\mathcal{G},b,\mu}(k)$, there exists $y \in \mathbb{M}_{\mathcal{G},\mu}^{\text{loc}}(k)$ and an isomorphism of formal completions

$$\widehat{\mathcal{M}_{\mathcal{G},b,\mu}/x} \simeq \widehat{\mathbb{M}_{\mathcal{G},\mu}^{\text{loc}}/y}.$$

When \mathcal{G} is a parahoric group scheme, Gleason has defined the formal completion $\widehat{\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}/x}$ as a v -sheaf. In our approach, we first show that his definition extends to the case of an arbitrary quasi-parahoric group scheme and then show that $\widehat{\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}/x}$ is representable (by the formal spectrum of a complete Noetherian local ring which is the completion of a corresponding local model, as above). The representability of $\widehat{\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}/x}$ for all x is closely related to the representability of $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}$. In fact, in [PR21], we show in the Hodge type case under certain hypotheses that, conversely, if all formal completions are representable, then so is $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}$. Using this statement, we prove in [PR21] the representability of $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}$ in many cases of Hodge type by using global methods.

By contrast, the approach in the present paper is purely local and direct, with more general results. The key case for $p > 2$ occurs for (G, b, μ, \mathcal{G}) of Hodge type when the closed immersion $G \hookrightarrow \text{GL}_n$ extends to a closed immersion $\mathcal{G} \hookrightarrow \text{GL}(\Lambda)$ for a \mathbb{Z}_p -lattice $\Lambda \subset \mathbb{Q}_p^n$. In this case, the representability of $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}$ is established by imitating the construction in [KP18], as extended in [KZ21], of integral models of global Shimura varieties of Hodge type. The crucial step here is the construction of a suitable versal deformation of a p -divisible group equipped with crystalline tensors. This is done by using Zink's displays and requires the assumption $p > 2$. The general Hodge type case is reduced to this case by devissage. In this devissage, there are two steps. In a first step, one shows that the assertion is independent of the quasi-parahoric \mathcal{G} within the class of all quasi-parahorics sharing a fixed parahoric as their neutral component. In a second step, one shows that the assertion is independent of the group G within the class

of all groups sharing the same adjoint group. In these devissage steps one has to deal with affine Deligne-Lusztig varieties for quasi-parahorics, as already considered by U. Görtz, X. He and S. Nie in [GHN20]. Finally, the general abelian type is reduced to the Hodge type case by following Deligne's analogous reduction [De79] in the case of global Shimura varieties.

What remains to be done for the construction of integral local Shimura varieties and Scholze's representability conjecture in all generality? For local Shimura data of abelian type, there are still cases open for $p = 2$. Some of these appear accessible but there are several technical complications. Outside the abelian type cases, we have only cases that involve exceptional groups of type E_6 or E_7 , even orthogonal groups with cocharacters which are "mixed", of type $\mathbf{D}^{\mathbb{R}}$ with $\mathbf{D}^{\mathbb{H}}$, and trialitarian forms. These are more mysterious. Especially the cases of exceptional groups are wide open.

We thank P. Scholze for helpful comments, and R. Zhou for an e-mail message which clarified a point in [KZ21]. The first author also acknowledges support by NSF grant #DMS-2100743.

2. STATEMENTS OF THE MAIN RESULTS

2.1. Hodge and Abelian type. Let $(G, b, \{\mu\})$ be a local Shimura datum over \mathbb{Q}_p , cf. [RV14]. Recall that this means that G is a reductive group, that $b \in G(\check{\mathbb{Q}}_p)$ and that $\{\mu\}$ is a conjugacy class of a minuscule cocharacter $\mu : \mathbb{G}_{m/\check{\mathbb{Q}}_p} \rightarrow G_{\check{\mathbb{Q}}_p}$. It is assumed that b is neutral acceptable, i.e., the σ -conjugacy class $[b]$ lies in $B(G, \mu^{-1})$. We will often omit the bracket and simply write (G, b, μ) .

We will be concerned with local Shimura data of a particular type.

Definition 2.1.1. The local Shimura datum (G, b, μ) is called of *Hodge type* if there is an embedding

$$i : (G, \mu) \hookrightarrow (\mathrm{GL}_h, \mu_d^{-1}),$$

for some $0 \leq d \leq h$, where $\mu_d(a) = \mathrm{diag}(a^{(d)}, 1^{(h-d)})$ is the standard minuscule cocharacter of GL_h .

Remark 2.1.2. This is the local analogue of the corresponding terminology in the classical theory of Shimura varieties, where one requires an embedding into a similitude symplectic group.

Definition 2.1.3. ([HPR20, §2.7], [HLR18, Def. 9.6]) The local Shimura datum (G, b, μ) is called of *abelian type* if there is a local Shimura datum (G_1, b_1, μ_1) of Hodge type and an isomorphism $(G_{1,\mathrm{ad}}, b_{1,\mathrm{ad}}, \mu_{1,\mathrm{ad}}) \simeq (G_{\mathrm{ad}}, b_{\mathrm{ad}}, \mu_{\mathrm{ad}})$. In this case, (G_1, b_1, μ_1) is called a *central lift of Hodge type* for (G, b, μ) , cf. [HLR18, §9].

Note that the existence of b_1 that lifts b_{ad} is automatic since $B(G, \mu) \simeq B(G_{\mathrm{ad}}, \mu_{\mathrm{ad}})$ by [Ko97, (6.5.1)], so this is really a property of the pair (G, μ) .

Remark 2.1.4. Note that this is weaker than the analogous notion in the global case, where one asks that there is a morphism $(G_1, \mu_1) \rightarrow (G_{\mathrm{ad}}, \mu_{\mathrm{ad}})$ such that the induced morphism $G_{1,\mathrm{der}} \rightarrow G_{\mathrm{ad}}$ admits a factorization through G_{der} and induces an isomorphism $(G_{1,\mathrm{ad}}, \mu_{1,\mathrm{ad}}) \simeq (G_{\mathrm{ad}}, \mu_{\mathrm{ad}})$, comp. [KP18]. In [HPR20], it is this stronger version that is imposed also in the local case; however, in the local case, this stronger notion seems unnecessary, as pointed out in [HLR18].

2.2. Quasi-parahoric subgroups. Let \check{G} be a reductive group over $\check{\mathbb{Q}}_p$. By definition, a quasi-parahoric subgroup of $\check{G}(\check{\mathbb{Q}}_p)$ is a subgroup \check{K} which is squeezed as

$$\check{G}(\check{\mathbb{Q}}_p)^0 \cap \text{Stab}_{\mathfrak{F}} \subset \check{K} \subset \check{G}(\check{\mathbb{Q}}_p)^1 \cap \text{Stab}_{\mathfrak{F}}.$$

Here $\text{Stab}_{\mathfrak{F}}$ is the stabilizer of a facet \mathfrak{F} in the building $\mathcal{B}(\check{G}_{\text{ad}}, \check{\mathbb{Q}}_p)$, and

$$\check{G}(\check{\mathbb{Q}}_p)^0 = \ker(\kappa : \check{G}(\check{\mathbb{Q}}_p) \longrightarrow \pi_1(\check{G})_I), \quad \check{G}(\check{\mathbb{Q}}_p)^1 = \ker(\kappa : \check{G}(\check{\mathbb{Q}}_p) \longrightarrow \pi_1(\check{G})_I \otimes \mathbb{Q}).$$

Here, κ is the Kottwitz homomorphism. Equivalently, it is a subgroup of finite index in $\check{G}(\check{\mathbb{Q}}_p)^1 \cap \text{Stab}_{\mathfrak{F}}$. Still another way of characterizing quasi-parahoric subgroups is to say that they are of finite index in the stabilizer of a point in the extended building $\mathcal{B}^e(\check{G}, \check{\mathbb{Q}}_p)$. In the case that the quasi-parahoric subgroup coincides with $\check{G}(\check{\mathbb{Q}}_p)^1 \cap \text{Stab}_{\mathbf{x}}$, with $\mathbf{x} \in \mathcal{B}(\check{G}_{\text{ad}}, \check{\mathbb{Q}}_p)$ (equivalently, if it coincides with the stabilizer subgroup in $\check{G}(\check{\mathbb{Q}}_p)$ of a point in the extended building $\mathcal{B}^e(\check{G}, \check{\mathbb{Q}}_p)$), it is called a *stabilizer quasi-parahoric*. The parahoric subgroup $\check{K}^\circ = \check{G}(\check{\mathbb{Q}}_p)^0 \cap \text{Stab}_{\mathfrak{F}}$ is called the parahoric subgroup *associated* to the quasi-parahoric subgroup \check{K} . Note that if $\pi_1(\check{G})_I$ is torsion-free, then any quasi-parahoric is a parahoric.

By Bruhat-Tits theory [BTII], there is a unique smooth group scheme $\check{\mathcal{G}}$ over $\check{\mathbb{Z}}_p$ with generic fiber \check{G} such that $\check{\mathcal{G}}(\check{\mathbb{Z}}_p) = \check{K}$.

Now let G be a reductive group over \mathbb{Q}_p . Then, in analogy with the Bruhat-Tits definition of a \mathbb{Q}_p -parahoric subgroup [BTII, right after Def. 5.2.6], there is the notion of a \mathbb{Q}_p -quasi-parahoric subgroup of G . It can be equivalently defined as a quasi-parahoric subgroup \check{K} of $G(\check{\mathbb{Q}}_p)$ which is invariant under Frobenius, or as a subgroup of finite index in $G(\check{\mathbb{Q}}_p)^1 \cap \text{Stab}_{\mathfrak{F}}$, where $\text{Stab}_{\mathfrak{F}}$ is the stabilizer of a facet \mathfrak{F} in the building $\mathcal{B}(G_{\text{ad}}, \mathbb{Q}_p)$. By descent from $\check{\mathbb{Z}}_p$ to \mathbb{Z}_p , we have a corresponding smooth group scheme \mathcal{G} over $\text{Spec}(\mathbb{Z}_p)$. Then $\mathcal{G}(\mathbb{Z}_p) = \check{K} \cap G(\mathbb{Q}_p)$.

In the sequel, we simply call \mathcal{G} a *quasi-parahoric group scheme* for G . Note that a parahoric group scheme for G is uniquely defined by its associated subgroup $\mathcal{G}(\mathbb{Z}_p)$ of $G(\mathbb{Q}_p)$, cf. [BTII, Prop. 5.2.8]. However, for general quasi-parahoric group schemes for G , the association $\mathcal{G} \mapsto \mathcal{G}(\mathbb{Z}_p)$ ceases to be injective. Still, we write $K = \mathcal{G}(\mathbb{Z}_p) = \check{K} \cap G(\mathbb{Q}_p)$ and sometimes use the notation K to refer to our choice of \mathcal{G} . This does not lead to confusions.

2.3. The local model. Recall the Scholze-Weinstein v -sheaf local model $\mathbb{M}_{\mathcal{G}, \mu}^v$ over $\text{Spd}(O_E)$, cf. [SW20, §§20, 21], see also [AGLR22] (in [SW20] it is denoted by $\text{Gr}_{\mathcal{G}, \text{Spd}(O_E), \leq \mu}$).

Assume μ is minuscule. In this case, $\text{Gr}_{\mathcal{G}, \text{Spd}(O_E), \leq \mu} = \text{Gr}_{\mathcal{G}, \text{Spd}(O_E), \mu}$ and the v -sheaf local model $\mathbb{M}_{\mathcal{G}, \mu}^v = \text{Gr}_{\mathcal{G}, \text{Spd}(O_E), \mu}$ is given as the closure inside the Beilinson-Drinfeld style affine Grassmannian $\text{Gr}_{\mathcal{G}, \text{Spd}(O_E)}$ of the v -sheaf X_μ^\diamond associated to the symmetric space $X_\mu = \text{Gr}_{G, \mu}$ of parabolics of type μ over $\text{Spec}(E)$. Then $\mathbb{M}_{\mathcal{G}, \mu}^v$ is representable by a normal flat projective O_E -scheme $\mathbb{M}_{\mathcal{G}, \mu}^{\text{loc}}$ with reduced special fiber. This was conjectured by Scholze-Weinstein [SW20, Conj. 21.4.1] and was shown in [AGLR22] and [GLo22] (which settled the normality in some remaining cases in characteristics 2 and 3). Here $\mathbb{M}_{\mathcal{G}, \mu}^{\text{loc}}$ is uniquely determined. Assume in addition that (G, μ) is of abelian type and satisfies Condition (A) or (B), cf. Definition 2.5.2 below. Then the normality of $\mathbb{M}_{\mathcal{G}, \mu}^{\text{loc}}$ is given by [AGLR22, Thm. 7.23]. Furthermore, in the case (A) $\mathbb{M}_{\mathcal{G}, \mu}^{\text{loc}}$ can be obtained by the procedure of [PZ13], extended to restrictions of scalars from wild extensions in [Le01] and as modified in [HPR20] when $p \mid |\pi_1(G_{\text{der}})|$. This statement uses Remark 2.5.3. In the case (B) of Definition 2.5.2, $\mathbb{M}_{\mathcal{G}, \mu}^{\text{loc}}$ can also be obtained by taking the closure of the generic fiber in the naive local model of [RZ96]. There is no

difference when discussing quasi-parahorics because the natural map is an isomorphism,

$$\mathbb{M}_{\mathcal{G}^o, \mu}^v \xrightarrow{\sim} \mathbb{M}_{\mathcal{G}, \mu}^v, \quad (2.3.1)$$

cf. [SW20, Prop. 21.4.3]. (This proposition assumes that μ is minuscule, but the argument applies to general μ , see also [AGLR22, §4]). If μ is minuscule, this also yields an isomorphism for the corresponding scheme local models,

$$\mathbb{M}_{\mathcal{G}^o, \mu}^{\text{loc}} \xrightarrow{\sim} \mathbb{M}_{\mathcal{G}, \mu}^{\text{loc}}.$$

2.4. The integral local Shimura variety $\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}$. Let \mathcal{G} be a smooth affine group scheme over \mathbb{Z}_p with generic fiber a reductive group G , let $b \in G(\check{\mathbb{Q}}_p)$, and let μ be a conjugacy class of cocharacters of G . It is assumed that the Frobenius-conjugacy class of b lies in $B(G, \mu^{-1})$. We call the triple (\mathcal{G}, b, μ) an *integral local shtuka datum* and (G, b, μ) a *rational local shtuka datum*. As usual, we denote by E the field of definition of μ and by \check{E} the completion of the maximal unramified extension.

Then Scholze-Weinstein associate to (\mathcal{G}, b, μ) an “integral” moduli space of shtukas $\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}$ over $O_{\check{E}}$, [SW20, Def. 25.1.1]. It is given as a “ v -sheaf moduli space” of certain \mathcal{G} -shtukas with one leg bounded by μ with a fixed associated Frobenius element, cf. [SW20, §§23.1, 23.2, 23.3], as follows.

Definition 2.4.1. The *integral moduli space of local shtuka* $\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}$ is the functor that sends $S \in \text{Perfd}_k$ to the set of isomorphism classes of tuples

$$(S^\sharp, \mathcal{P}, \phi_{\mathcal{P}}, i_r),$$

where

- 1) S^\sharp is an untilt of S over $\text{Spa}(O_{\check{E}})$,
- 2) $(\mathcal{P}, \phi_{\mathcal{P}})$ is a \mathcal{G} -shtuka over S with one leg along S^\sharp bounded by μ ,
- 3) i_r is a “framing”, i.e. an isomorphism of G -torsors

$$i_r : G_{\mathcal{Y}_{[r, \infty)}(S)} \xrightarrow{\sim} \mathcal{P}_{|\mathcal{Y}_{[r, \infty)}(S)} \quad (2.4.1)$$

for large enough r (for an implicit choice of pseudouniformizer ϖ), under which $\phi_{\mathcal{P}}$ is identified with $\phi_b = b \times \text{Frob}_S$.

Here, $\mathcal{Y}_{[r, \infty)}(S)$ is as defined in [SW20], [FS21, II]. We have denoted by $G_{\mathcal{Y}_{[r, \infty)}(S)}$ the trivial G -torsor over $\mathcal{Y}_{[r, \infty)}(S)$ (denoted $G \times \mathcal{Y}_{[r, \infty)}(S)$ in [SW20, App. to §19]), and by

$$\phi_b = \phi_{G, b} : \phi^*(G_{\mathcal{Y}_{[r, \infty)}(S)}) = G_{\mathcal{Y}_{[\frac{1}{p}r, \infty)}(S)} \xrightarrow{b\phi} G_{\mathcal{Y}_{[r, \infty)}(S)} \quad (2.4.2)$$

the isomorphism given by the ϕ -linear isomorphism induced by right multiplication by b (denoted by $b \times \text{Frob}$ in [SW20, Def. 23.1.1]). In 3) we mean more precisely an equivalence class, where i_r and $i_{r'}$ are called equivalent if there exists $r'' \geq r, r'$ such that $i_r|_{\mathcal{Y}_{[r'', \infty)}(S)} = i_{r'}|_{\mathcal{Y}_{[r'', \infty)}(S)}$. Also, in 2) the precise definition of “bounded by μ ” is given via the local model $\mathbb{M}_{\mathcal{G}, \mu}^v$ (see [SW20, Def. 25.1.1], [PR21].)

Note that the formation of $\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}$ is functorial, i.e., for a morphism $(\mathcal{G}, b, \mu) \rightarrow (\mathcal{G}', b', \mu')$ we have $\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}} \rightarrow \mathcal{M}_{\mathcal{G}', b', \mu'}^{\text{int}} \times_{\text{Spd}(O_{\check{E}})} \text{Spd}(O_{\check{E}'})$. It is also compatible with products, i.e., $\mathcal{M}_{\mathcal{G}_1 \times \mathcal{G}_2, b_1 \times b_2, \mu_1 \times \mu_2}^{\text{int}}$ is isomorphic to the product $\mathcal{M}_{\mathcal{G}_1, b_1, \mu_1}^{\text{int}} \times \mathcal{M}_{\mathcal{G}_2, b_2, \mu_2}^{\text{int}}$ after base changing

to the compositum of the reflex fields (in this, we omit the base changes from the notation, for simplicity).

By loc. cit., $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}$ is a v -sheaf over $\text{Spd}(O_{\tilde{E}})$. In fact, by [Gl20, Prop. 2.3.5], $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}$ is a small v -sheaf. In addition, $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}$ supports a Weil descent datum from $O_{\tilde{E}}$ down to O_E , as explained in [PR21, 3.1, 3.2]. In this paper, even though this is not always pointed out explicitly, the smooth group scheme \mathcal{G} will always be a quasi-parahoric group scheme for G .

In fact, most of the time we are interested in the case that (G, b, μ) is a local Shimura datum, i.e. we also assume that μ is minuscule. In addition, we take \mathcal{G} to be a quasi-parahoric group scheme for G . In this case we call $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}$ the *integral local Shimura variety* associated to the local Shimura datum (G, b, μ) and the quasi-parahoric group scheme \mathcal{G} .

2.5. Statement of the main results. Our concern is with the following conjecture.

Conjecture 2.5.1. (Scholze) *Let (G, b, μ) be a local Shimura datum and \mathcal{G} a quasi-parahoric group scheme for G . There exists a formal scheme $\mathcal{M}_{\mathcal{G},b,\mu}$ which is normal and flat locally formally of finite type over $O_{\tilde{E}}$ with*

$$\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}} = \mathcal{M}_{\mathcal{G},b,\mu}^{\diamond},$$

as v -sheaves over $\text{Spd}(O_{\tilde{E}})$. Then, $\mathcal{M}_{\mathcal{G},b,\mu}$ is unique ([SW20, Prop. 18.4.1]).

Our main result is a proof of this conjecture when (G, b, μ) is of abelian type under certain mild hypotheses.

Definition 2.5.2. Let (G, b, μ) be of abelian type. We introduce two kinds of conditions on p , G and μ .

(A) $p \neq 2$.

(B) $p = 2$ and $G_{\text{ad}} = \prod_i \text{Res}_{F_i/\mathbb{Q}_p} H_i$, where for each i , $H_i = B_i^\times / F_i^\times$ with B_i a simple algebra with center F_i , or $H_i = \text{PGSp}_{2n_i}$, or the corresponding component $\mu_{\text{ad},i}$ of μ_{ad} is trivial.

Note that if $G = T$ is a torus, T trivially satisfies (A) or (B).

Remark 2.5.3. Recall from [PR21, §5] that G is called acceptable if $G_{\text{ad}} \simeq \prod_i \text{Res}_{F_i/\mathbb{Q}_p} H_i$, where H_i is absolutely simple and splits over a tamely ramified extension of F_i . Note that G is automatically acceptable if $p \geq 5$, cf. [PR21, §5]. Moreover, if $p = 3$, the acceptability condition only excludes groups whose adjoint groups contain a ramified triality group (type $D_4^{(3)}$ or $D_4^{(6)}$) as a factor. We note that by Serre [Se79, §3, Cor. 2] this last possibility does not occur when (G, b, μ) is of abelian type, unless the corresponding component of μ_{ad} is trivial. In other words, Condition (A) implies that the simple factors of G_{ad} on which the projections of μ_{ad} are non-trivial, are acceptable. Indeed, let (G_1, μ_1) be a central lift of Hodge type for (G, μ) . Let G'_1 be the minimal normal subgroup of G_1 containing the conjugacy class μ_1 . Then by [Se79, §3, Cor. 2], $G'_{1,\text{ad}}$ contains no ramified triality group as a factor. On the other hand, $G'_{1,\text{ad}} = \prod_{\{i | \mu_{\text{ad},i} \neq 1\}} \text{Res}_{F_i/\mathbb{Q}_p} H_i$, which proves the claim.

Theorem 2.5.4. *Let (G, b, μ) be a local Shimura datum of abelian type satisfying Condition (A) or (B), and let \mathcal{G} be a quasi-parahoric group scheme for G . Then $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}$ satisfies Conjecture 2.5.1.*

Theorem 2.5.4 is closely related to the following result. In fact, it was shown in [PR21] that the following result implies Theorem 2.5.4 in the Hodge type case under some mild additional hypotheses, cf. [PR21, proof of Thm. 3.7.1].

Theorem 2.5.5. *Let (G, b, μ) be a local Shimura datum of abelian type satisfying Condition (A) or (B), and let \mathcal{G} be a quasi-parahoric group scheme for G . For any $\mathrm{Spd}(k)$ -valued point x of $\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}}$, there is a k -valued point y of $\mathbb{M}_{\mathcal{G}, \mu}^{\mathrm{loc}}$ such that*

$$\widehat{\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}}}_{/x} \simeq (\widehat{\mathbb{M}_{\mathcal{G}, \mu}^{\mathrm{loc}}}_{/y})^{\diamond}, \quad (2.5.1)$$

as v -sheaves over $\mathrm{Spd}(O_{\check{E}})$.

In this, y is the k -valued point of $\mathbb{M}_{\mathcal{G}, \mu}^{\mathrm{loc}}$ obtained by fixing a trivialization of the \mathcal{G} -torsor underlying the \mathcal{G} -shtuka at x . The notation $\widehat{\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}}}_{/x}$ stands for the formal completion of the v -sheaf $\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}}$ at x , as defined in [Gl20, Def. 1.4.24], cf. [PR21, §3.3.1], see also §3.4 (Gleason calls these “tubular neighborhoods”). On the RHS, $(\widehat{\mathbb{M}_{\mathcal{G}, \mu}^{\mathrm{loc}}}_{/y})^{\diamond}$ denotes the v -sheaf attached to the adic space $\mathrm{Spa}(A, A)$, where A is the completion of the local ring of the scheme $\mathbb{M}_{\mathcal{G}, \mu}^{\mathrm{loc}}$ at y , taken with the topology defined by the maximal ideal.

The combination of these two results also gives

Corollary 2.5.6. *Let (G, b, μ) be a local Shimura datum of abelian type satisfying Condition (A) or (B), and let \mathcal{G} be a quasi-parahoric group scheme for G . Let $\mathcal{M}_{\mathcal{G}, b, \mu}$ be the formal scheme that represents $\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}}$ by Theorem 2.5.4. For any k -valued point x of $\mathcal{M}_{\mathcal{G}, b, \mu}$, there is a k -valued point y of $\mathbb{M}_{\mathcal{G}, \mu}^{\mathrm{loc}}$ such that $\widehat{\mathcal{M}_{\mathcal{G}, b, \mu}}_{/x} \simeq \widehat{\mathbb{M}_{\mathcal{G}, \mu}^{\mathrm{loc}}}_{/y}$.*

Note here that, by the full-faithfulness result of [SW20, Prop. 18.4.1], the Weil descent datum for the v -sheaf $\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}}$ gives a corresponding Weil descent datum for the formal scheme $\mathcal{M}_{\mathcal{G}, b, \mu}$, from $\mathcal{O}_{\check{E}}$ down to O_E .

The following result shows that $\mathcal{M}_{\mathcal{G}, b, \mu}$ is an integral model of the local Shimura variety in a certain sense. Recall from the introduction that to \mathcal{G} is associated the finite abelian group $\Pi_{\mathcal{G}}$, and that to every $\bar{\beta} \in \Pi_{\mathcal{G}}$, there is associated a quasi-parahoric group scheme $\mathcal{G}_{\bar{\beta}}$ over \mathbb{Z}_p , as well as its associated parahoric group scheme $\mathcal{G}_{\bar{\beta}}^{\circ}$, and the finite group of its connected components $\pi_0(\mathcal{G}_{\bar{\beta}})$ with its action by the Frobenius ϕ . Also we let $K_{\bar{\beta}} = \mathcal{G}_{\bar{\beta}}(\mathbb{Z}_p)$.

Theorem 2.5.7. *Let (G, b, μ) be a local Shimura datum of abelian type satisfying Condition (A) or (B), and let \mathcal{G} be a quasi-parahoric group scheme for G . There is an isomorphism of rigid-analytic varieties over \check{E} ,*

$$\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{rig}} \simeq \bigsqcup_{\bar{\beta} \in \Pi_{\mathcal{G}}} \mathrm{Sht}_{K_{\bar{\beta}}}(G, b, \mu).$$

This isomorphism is induced by an isomorphism of formal schemes

$$\mathcal{M}_{\mathcal{G}, b, \mu} \simeq \bigsqcup_{\bar{\beta} \in \Pi_{\mathcal{G}}} \mathcal{M}_{\mathcal{G}_{\bar{\beta}}, b, \mu} / \pi_0(\mathcal{G}_{\bar{\beta}})^{\phi}.$$

2.6. The plan of the proof. The proof proceeds in the following steps.

- *Step 1.* We develop a devissage procedure that allows us to pass
 - a) between \mathcal{G} and \mathcal{G}° .

b) between groups linked via an ad-isomorphism.

In most of the statements in this step, we deal with integral local shtuka data (\mathcal{G}, b, μ) , i.e. we do not assume that μ is minuscule.

- *Step 2.* We show the results in “good” Hodge type cases: In such cases, the proof follows from the case of GL_n (already treated in [SW20] by relating to RZ formal schemes) and from the constructions in [KP18], [KZ21] (which use Zink displays and Breuil-Kisin modules).
- *Step 3.* Using Step 1 we reduce the general case to:
 - a) the Hodge type cases handled in Step 2, (in case (A)),
 - b) EL/PEL cases for which we give directly $\mathcal{M}_{\mathcal{G}, b, \mu}$ as a Rapoport-Zink formal scheme, (in case (B)).

2.7. The lay-out of the paper. In §3, we generalize the Anschütz purity theorem from parahoric group schemes to quasi-parahoric group schemes, and use this to transfer to this more general context the formalism of specialization, formal completion and v -sheaf local model diagram that Gleason had established for hyperspecial parahoric \mathcal{G} . We also make the link between the reduced locus of $\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}}$ and affine Deligne-Lusztig varieties for quasi-parahorics. In §4, we study the devissage step of varying the quasi-parahorics with a given associated parahoric. We also prove at this point Theorem 2.5.7. In §5, we study the devissage step of varying the groups with a given adjoint group. In §6 we treat the case when μ_{ad} is trivial. In §7 we consider the crucial Hodge type case mentioned in the introduction. In §8, we develop methods to relate Hodge type cases and abelian type cases to the crucial Hodge type case. Everything comes together in §9, where we give the proofs of Theorems 2.5.4 and 2.5.5; this is done separately for G satisfying Condition (A) and (B).

3. PRELIMINARIES

3.1. Torsors under quasi-parahoric group schemes. Let $\Omega = \Omega_G = \pi_1(G)_I$. We recall the Kottwitz homomorphism

$$\kappa: G(\check{\mathbb{Q}}_p) \longrightarrow \Omega_G.$$

A parahoric subgroup lies in the kernel of κ and in fact, the kernel of κ is generated by all parahoric subgroups, cf. [HR08, Lem. 17]. Also, for any quasi-parahoric subgroup \check{K} of $G(\check{\mathbb{Q}}_p)$ with associated parahoric subgroup \check{K}^o we have $\check{K}^o = \ker(\kappa|_{\check{K}}: \check{K} \rightarrow \Omega)$. Let \mathcal{G} be the corresponding quasi-parahoric group scheme and consider the injective map

$$\pi_0(\mathcal{G}) = \check{K}/\check{K}^o \hookrightarrow \Omega_G. \quad (3.1.1)$$

In [SW20, 25.3.1], Scholze-Weinstein point out the importance of the finite abelian group

$$\Pi_{\mathcal{G}} := \ker(\pi_0(\mathcal{G})_{\phi} \longrightarrow \Omega_{\phi}) \quad (3.1.2)$$

Lemma 3.1.1. *There is a natural isomorphism*

$$\Pi_{\mathcal{G}} \cong \ker(\mathrm{H}_{\mathrm{\acute{e}t}}^1(\mathbb{Z}_p, \mathcal{G}) \longrightarrow \mathrm{H}_{\mathrm{\acute{e}t}}^1(\mathbb{Q}_p, G)).$$

Proof. This is sketched in [SW20, 25.3]. The exact sequence $0 \rightarrow \check{K}^o \rightarrow \check{K} \rightarrow \pi_0(\mathcal{G}) \rightarrow 0$ induces an exact sequence

$$\mathrm{H}_{\mathrm{\acute{e}t}}^1(\mathbb{Z}_p, \mathcal{G}^o) \longrightarrow \mathrm{H}_{\mathrm{\acute{e}t}}^1(\mathbb{Z}_p, \mathcal{G}) \longrightarrow \mathrm{H}_{\mathrm{\acute{e}t}}^1(\mathbb{Z}_p, \pi_0(\mathcal{G})).$$

Here the left term vanishes. The second map is surjective since $H_{\text{ét}}^1(\mathbb{Z}_p, \mathcal{G}) = \check{K}/_{\phi} \check{K}$. We therefore obtain

$$H_{\text{ét}}^1(\mathbb{Z}_p, \mathcal{G}) \simeq H_{\text{ét}}^1(\mathbb{F}_p, \pi_0(\mathcal{G})) \simeq H^1(\langle \phi \rangle, \pi_0(\mathcal{G})) \cong \pi_0(\mathcal{G})_{\phi}.$$

On the other hand, we have an injection

$$H_{\text{ét}}^1(\mathbb{Q}_p, G) \hookrightarrow H^1(\langle \phi \rangle, G(\check{\mathbb{Q}}_p)) \cong B(G) \quad (3.1.3)$$

with the image landing in the basic elements $B(G)_{\text{basic}} \subset B(G)$, cf. [Ko97]. Now

$$B(G)_{\text{basic}} \simeq \pi_1(G)_{\Gamma} = (\pi_1(G)_I)_{\phi} = \Omega_{\phi},$$

and the map (3.1.3) induces an identification

$$H_{\text{ét}}^1(\mathbb{Q}_p, G) \simeq \Omega_{\phi, \text{tors}},$$

where $\Omega_{\phi, \text{tors}}$ is the torsion subgroup of Ω_{ϕ} . It remains to observe that the following diagram is commutative,

$$\begin{array}{ccccc} H_{\text{ét}}^1(\mathbb{Z}_p, \mathcal{G}) & \longrightarrow & H_{\text{ét}}^1(\mathbb{Q}_p, G) & \hookrightarrow & B(G) \\ \simeq \downarrow & & \downarrow \simeq & & \downarrow \simeq \\ \pi_0(\mathcal{G})_{\phi} & \longrightarrow & \Omega_{\phi, \text{tors}} & \hookrightarrow & \Omega_{\phi}. \end{array} \quad (3.1.4)$$

This commutativity follows from the fact that the vertical homomorphisms on the outer left and the outer right are both induced by the Kottwitz homomorphism $G(\check{\mathbb{Q}}_p) \rightarrow \Omega$: this is obvious from the construction for the outer left map and follows from [Ko97, §7.5] for the outer right map. \square

3.2. Purity of torsors. The following purity/extension result is crucial. It is a quick generalization of the corresponding result for parahoric subgroups due to Anschütz ([An18]).

Proposition 3.2.1. *Suppose \mathcal{G} is a quasi-parahoric group scheme over \mathbb{Z}_p , and consider a pair (C, C^+) with C an algebraically closed non-archimedean field of characteristic p and C^+ an open and bounded valuation ring of C . Then every \mathcal{G} -torsor over $U = \text{Spec}(W(C^+)) \setminus V(p, [\varpi])$ extends to a \mathcal{G} -torsor over $\text{Spec}(W(C^+))$ and hence is trivial.*

Proof. We have an exact sequence

$$1 \longrightarrow \mathcal{G}^o \longrightarrow \mathcal{G} \longrightarrow i_*(\pi_0(\mathcal{G})) \longrightarrow 1$$

of étale sheaves, where $i : \text{Spec}(k) \hookrightarrow \text{Spec}(\check{\mathbb{Z}}_p)$. Note $U \times_{\text{Spec}(\check{\mathbb{Z}}_p)} \text{Spec}(k) = \text{Spec}(C^+[1/\varpi]) = \text{Spec}(C)$. This gives the exact sequence

$$H^1(U, \mathcal{G}^o) \longrightarrow H^1(U, \mathcal{G}) \longrightarrow H^1(U, i_*(\pi_0(\mathcal{G}))) = H^1(\text{Spec}(C), \pi_0(\mathcal{G})) = (0).$$

By Anschütz's theorem [An18], $H^1(U, \mathcal{G}^o) = (0)$ and the result follows. \square

This extends as follows to strictly totally disconnected affinoid perfectoids which are given as “products of points”.

Proposition 3.2.2. *Let (R, R^+) be a “product of the points (C_i, C_i^+) , $i \in I$ ”, with all C_i algebraically closed of characteristic p , i.e., $R^+ = \prod_{i \in I} C_i^+$, $R = R^+[1/\varpi]$, with the ϖ -topology, where $\varpi = (\varpi_i)_i$, with ϖ_i pseudouniformizers of C_i^+ . Then every \mathcal{G} -torsor over $\text{Spec}(W(R^+)) \setminus V(p, [\varpi])$ extends to a \mathcal{G} -torsor over $\text{Spec}(W(R^+))$ and is trivial.*

Proof. In the parahoric case $\mathcal{G} = \mathcal{G}^o$, the extension is shown in [Gl20, Prop. 2.1.19]. Observe that all étale covers of

$$\mathrm{Spec}(R^+) = \mathrm{Spec}\left(\prod_{i \in I} C_i^+\right)$$

split. Indeed, as in the proof of [BS17, Lem. 6.2], we can consider

$$\mathrm{Spec}\left(\prod_{i \in I} C_i^+\right) \longrightarrow \pi_0(\mathrm{Spec}\left(\prod_i C_i^+\right)) = \pi_0(\mathrm{Spec}\left(\prod_i k_i\right)) = \beta I$$

where $k_i = C_i^+/\mathfrak{m}_i$ is the (algebraically closed) residue field. Here, βI is the Stone-Čech compactification of the discrete set I . Each connected component of $\mathrm{Spec}(R^+)$ (i.e. fiber of this map) is the spectrum of a valuation ring V with algebraically closed fraction field K ; this valuation ring is an ultraproduct of C_i^+ . It follows that all étale covers of $\mathrm{Spec}(V)$ and then also of $\mathrm{Spec}(R^+)$ split. (We see that R^+ is “strictly w -local” in the terminology of [BS15, 2.2], and, in fact, $\mathrm{Spec}(R^+)$ is “ w -contractible”, [BS15, Lem. 2.4.8].)

A similar picture holds for the localization $R = R^+[1/\varpi]$: in this case, we have $\pi_0(\mathrm{Spec}(R)) \simeq \pi_0(\mathrm{Spec}(R^+)) \simeq \beta I$ (by considering idempotents) and

$$\mathrm{Spec}(R) \longrightarrow \pi_0(\mathrm{Spec}(R)) \simeq \beta I,$$

has every fiber isomorphic to $\mathrm{Spec}(V[1/\varpi])$, with $V[1/\varpi]$ a valuation ring with (algebraically closed) fraction field K . Étale covers over each such $\mathrm{Spec}(V[1/\varpi])$ split, and then étale covers over $\mathrm{Spec}(R)$ also split.

Now it also follows that every \mathcal{G} -torsor over $\mathrm{Spec}(W(R^+))$ is trivial. Indeed, since \mathcal{G} is smooth and $W(R^+)$ is p -adically complete, it is enough to show that all \mathcal{G} -torsors over $R^+ = \prod_{i \in I} C_i^+$ are trivial. As above, we see that all étale covers of $\mathrm{Spec}(R^+)$ split and so this follows using the smoothness of \mathcal{G} ; the same applies of course to \mathcal{G}^o -torsors. The argument in the proof of Proposition 3.2.1 above now extends to $U = \mathrm{Spec}(W(R^+)) \setminus V(p, [\varpi])$, to complete the proof. \square

3.3. The affine Witt Grassmannian and affine Deligne-Lusztig varieties. Let

$$X_{\mathcal{G}} := \mathrm{Gr}_{\mathcal{G}}^W = LG/L^+\mathcal{G} \tag{3.3.1}$$

be the Witt vector affine Grassmannian for \mathcal{G} , cf. [BS17, Zhu17]. Here $LG(R) = G(W(R)[\frac{1}{p}])$ and $L\mathcal{G}(R) = \mathcal{G}(W(R))$ for any perfect k -algebra R . Note that here \mathcal{G} is only assumed to be a quasi-parahoric group scheme, whereas in loc. cit. it is assumed that \mathcal{G} is a parahoric group scheme. Note that k -points of $X_{\mathcal{G}}$ are given by isomorphism classes of pairs (\mathcal{P}, α) of a \mathcal{G} -torsor \mathcal{P} over $W(k)$ with a trivialization α of its restriction to $W(k)[1/p]$.

Let (\mathcal{G}, b, μ) be an integral local shtuka datum such that \mathcal{G} is a quasi-parahoric group scheme for G . Inside the Witt vector affine Grassmannian we consider the affine Deligne-Lusztig variety $X_{\mathcal{G}}(b, \mu^{-1})$ (the μ^{-1} -admissible locus). This is a perfect scheme over k with

$$X_{\mathcal{G}}(b, \mu^{-1})(k) = \{g\check{K} \in G(\check{\mathbb{Q}}_p)/\check{K} \mid g^{-1}b\phi(g) \in \mathrm{Adm}^K(\mu^{-1})\} \subset X_{\mathcal{G}}(k). \tag{3.3.2}$$

Here $\mathrm{Adm}^K(\mu^{-1}) = \check{K}\mathrm{Adm}(\mu^{-1})\check{K}$, where $\mathrm{Adm}(\mu^{-1}) \subset \widetilde{W}$ is the μ^{-1} -admissible subset of the Iwahori Weyl group of G .

Proposition 3.3.1. *Let (\mathcal{G}, b, μ) be an integral local shtuka datum such that \mathcal{G} is a quasi-parahoric group scheme for G . Then the reduced locus $(\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}})_{\text{red}}$ is represented by the perfect k -scheme $X_{\mathcal{G}}(b, \mu^{-1})$, and hence*

$$\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}(\text{Spd}(k)) = X_{\mathcal{G}}(b, \mu^{-1})(k).$$

Proof. Here, the reduced locus \mathcal{F}_{red} of a small v -sheaf \mathcal{F} is a “scheme-theoretic” v -sheaf which is defined as in [Gl20, 1.4]. For hyperspecial parahoric \mathcal{G} the proposition is shown in [Gl20], cf. [Gl20, Prop. 2.3.10]. One main ingredient in the proof of this is the Anschütz purity theorem for \mathcal{G} and its extension to products of points. By Propositions 3.2.1 and 3.2.2, these purity statements remain true for quasi-parahoric group schemes. Also, by [AGLR22, Thm. 6.16] combined with (2.3.1), the reduced locus $(\mathbb{M}_{\mathcal{G}, \mu}^v)_{\text{red}}$ is represented by the perfect k -scheme which is the μ -admissible locus in $X_{\mathcal{G}}$. With these ingredients, the proof in [Gl20] now extends to this case. \square

Remark 3.3.2. In the identification of Prop. 3.3.1 above, the inverse μ^{-1} appears on the RHS because of the convention in the definition of “bounded by μ ” for \mathcal{G} -shtuka, comp. [PR21].

3.4. The specialization map and formal completions. Gleason explains certain conditions on a small v -sheaf \mathcal{F} to construct a continuous specialization map on the underlying spaces,

$$\text{sp}_{\mathcal{F}}: |\mathcal{F}| \longrightarrow |\mathcal{F}_{\text{red}}|,$$

cf. [Gl20, Prop. 1.4.15], see also [AGLR22, §2.3].

Futhermore, he proves that these conditions are satisfied when $\mathcal{F} = \mathbb{M}_{\mathcal{G}, \mu}^v$ with \mathcal{G} reductive (hyperspecial parahoric). This result is extended to parahoric \mathcal{G} by [AGLR22, Prop. 4.14]. In view of (2.3.1), it also holds for quasi-parahoric \mathcal{G} . The scheme-theoretic v -sheaf $(\mathbb{M}_{\mathcal{G}, \mu}^v)_{\text{red}}$ is represented by a perfect k -scheme (by [AGLR22] this is the μ^{-1} -admissible locus) which is a closed subscheme of the Witt affine Grassmannian $X_{\mathcal{G}} = \text{Gr}_{\mathcal{G}}^W$.

By [Gl20, Prop. 2.3.5], these conditions are also satisfied in the case when $\mathcal{F} = \mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}$ where \mathcal{G} is a hyperspecial parahoric subgroup. Again, one main ingredient is the Anschütz purity theorem for \mathcal{G} and its extension to product of points. By Propositions 3.2.1 and 3.2.2, these purity statements remain true for quasi-parahoric group schemes, and the proof extends.

In fact, by Proposition 3.3.1, $(\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}})_{\text{red}}$ is represented by the affine Deligne-Lusztig variety (ADLV) $X_{\mathcal{G}}(b, \mu^{-1})$ in the Witt affine Grassmannian $X_{\mathcal{G}}$. Hence we get a continuous map

$$\text{sp}_{\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}}: |\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}| \longrightarrow |(\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}})_{\text{red}}| = |X_{\mathcal{G}}(b, \mu^{-1})|. \quad (3.4.1)$$

Using this, we define the formal completion of $\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}$ along a point $x \in \mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}(\text{Spd}(k)) = X_{\mathcal{G}}(b, \mu^{-1})(k)$ as the sub- v -sheaf of $\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}$, with

$$\widehat{\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}}_{/x}(S) = \{y: S \longrightarrow \mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}} \mid \text{sp}_{\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}} \circ y(|S|) \subset \{x\}\}.$$

Similarly, we can define the formal completion $\widehat{\mathbb{M}_{\mathcal{G}, \mu}^v}_{/y}$ of $\mathbb{M}_{\mathcal{G}, \mu}^v$ along a point

$$y \in \mathbb{M}_{\mathcal{G}, \mu}^v(\text{Spd}(k)) \subset X_{\mathcal{G}}(k) = G(W(k)[1/p])/\mathcal{G}(W(k)).$$

If μ is minuscule, then $\mathbb{M}_{\mathcal{G}, \mu}^v$ is representable by the O_E -scheme $\mathbb{M}_{\mathcal{G}, \mu}^{\text{loc}}$ ([AGLR22]), and the formal completion $\widehat{\mathbb{M}_{\mathcal{G}, \mu}^v}_{/y}$ is given as $\text{Spd}(A, A)$, where A is the completion of the local ring

of $\mathbb{M}_{\mathcal{G},\mu}^{\text{loc}} \otimes_{O_E} \check{O}_E$ at the corresponding point, taken with the topology given by the maximal ideal.

3.5. Change of base point. Note that, when $b \in \text{Adm}^K(\mu^{-1})$, then $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}$ has a canonical $\text{Spd}(k)$ -valued “base point” x_0 . Under the identification $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}(\text{Spd}(k)) = X_{\mathcal{G}}(b, \mu^{-1})(k) \subset X_{\mathcal{G}}(k)$ it corresponds to the “base point” of the ADLV given by the trivial coset.

In general, let $x \in \mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}(\text{Spd}(k)) = X_{\mathcal{G}}(b, \mu^{-1})(k) \subset X_{\mathcal{G}}(k)$ correspond to the isomorphism class of a pair (\mathcal{P}, α) , where \mathcal{P} is a \mathcal{G} -torsor over $W(k)$ and

$$\alpha : \mathcal{P}[1/p] \xrightarrow{\sim} \mathcal{G} \times \text{Spec}(W(k)[1/p])$$

is a trivialization of the restriction $\mathcal{P}[1/p]$ of \mathcal{P} to $\text{Spec}(W(k)[1/p])$ such that

$$\phi_{\mathcal{P}} = \alpha^{-1} \cdot \phi_b \cdot \phi^*(\alpha) : \phi^*(\mathcal{P})[1/p] \xrightarrow{\sim} \mathcal{P}[1/p]$$

has pole at $p = 0$ bounded by μ . Choose a trivialization of the \mathcal{G} -torsor \mathcal{P} over $W(k)$. Then α is given by $g \in \mathcal{G}(W(k)[1/p])$ and $\phi_{\mathcal{P}}$ by $b_x \times \phi : \mathcal{G}[1/p] \rightarrow \mathcal{G}[1/p]$ such that $b_x = g^{-1}b\phi(g)$. Hence we have $b_x = g^{-1}b\phi(g) \in \text{Adm}^K(\mu^{-1})$. We obtain an isomorphism

$$\tau_g : \mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}} \longrightarrow \mathcal{M}_{\mathcal{G},b_x,\mu}^{\text{int}}, \quad (\mathcal{P}, \phi_{\mathcal{P}}, i_r) \longmapsto (\mathcal{P}, \phi_{\mathcal{P}}, i_r \circ g^{-1})$$

which sends x to the base point x_0 of $\mathcal{M}_{\mathcal{G},b_x,\mu}^{\text{int}}$, cf. [PR21, proof of Prop. 3.4.1, p. 56]. This gives an isomorphism (depending on our choices)

$$\widehat{\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}}_{/x} \simeq \widehat{\mathcal{M}_{\mathcal{G},b_x,\mu}^{\text{int}}}_{/x_0}. \quad (3.5.1)$$

In particular, $\widehat{\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}}_{/x}$ is representable (by a normal complete Noetherian local ring) for given \mathcal{G} and any b in a given ϕ -conjugacy class in $B(G)$ and arbitrary $x \in \mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}(\text{Spd}(k))$ if and only if this holds for the base point x_0 of $\widehat{\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}}$, for given \mathcal{G} and any $b \in \text{Adm}^K(\mu^{-1})$ in the given ϕ -conjugacy class. Note that, by He’s theorem [He16], any ϕ -conjugacy class $[b] \in B(G, \mu^{-1})$ contains elements $b \in \text{Adm}^K(\mu^{-1})$.

3.6. A v -sheaf “local model diagram”. Let \mathcal{H} be an affine group scheme over $\check{\mathbb{Z}}_p$. For an affinoid perfectoid (R, R^+) over k , consider

$$\mathbb{W}^+\mathcal{H}(\text{Spa}(R, R^+)) = \mathcal{H}(W(R^+)),$$

and

$$\widehat{\mathbb{W}^+\mathcal{H}}(R, R^+) = \{h \in \mathcal{H}(W(R^+)) \mid h \equiv 1 \pmod{[\varpi_h]}\},$$

(here ϖ_h stands for some pseudo-uniformizer of R^+ that depends on h).

These definitions extend to perfectoid spaces over k and define v -sheaves of groups $\mathbb{W}^+\mathcal{H}$ and $\widehat{\mathbb{W}^+\mathcal{H}}$ on Perfd_k . We set

$$\widehat{L_W^+}\mathcal{H} = \widehat{\mathbb{W}^+\mathcal{H}} \times \text{Spd}(\mathbb{Z}_p)$$

which is a v -sheaf of groups over $\text{Spd}(\mathbb{Z}_p)$. (This definition appears in [Gl20, 2.3.15].) We have

$$\widehat{L_W^+}\mathcal{H}(S) = \{((S^\sharp, y), h) \mid h \in \mathcal{H}(W(R^+)), h \equiv 1 \pmod{[\varpi_h]}\}, \quad (3.6.1)$$

where $S = \text{Spa}(R, R^+)$ and (S^\sharp, y) is an S -valued point of $\text{Spd}(\mathbb{Z}_p)$, and where ϖ_h is a pseudouniformizer of R^+ (that depends on h). We can think of $\widehat{L_W^+}\mathcal{H}$ as the formal completion at the identity of the v -sheaf of groups $\mathbb{W}^+\mathcal{H} \times \text{Spd}(\mathbb{Z}_p)$.

We will later need the following lemma.

Lemma 3.6.1. *For each $h \in \widehat{\mathbb{W}}^+ \mathcal{H}(R, R^+)$, there is a unique $\lambda \in \widehat{\mathbb{W}}^+ \mathcal{H}(R, R^+)$ such that $h = \lambda^{-1} \cdot \phi(\lambda)$.*

Proof. The ring $W(R^+)$ is complete and separated for the $[\varpi]$ -topology, where ϖ is a pseudo-uniformizer of R^+ . Set inductively $\lambda_0 = 1$, $\eta_0 = h^{-1}$, and

$$\eta_n = \lambda_n^{-1} \phi(\lambda_n) \cdot h^{-1}, \quad \lambda_{n+1} = \lambda_n \cdot \eta_n,$$

Then we have

$$\eta_{n+1} = h \cdot \phi(\eta_n) \cdot h^{-1}.$$

Since $\eta_0 \equiv 1 \pmod{[\varpi_h]}$, this gives $\eta_n \equiv 1 \pmod{[\varpi_h^{p^n}]}$. Hence, η_n converges to 1 and λ_n converges to an element λ with

$$h = \lambda^{-1} \phi(\lambda).$$

(cf. [Gl20, Lem. 2.1.27].) If $\lambda = \phi(\lambda)$ with $\lambda \equiv 1 \pmod{[\varpi]}$, then we easily see inductively $\lambda \equiv 1 \pmod{[\varpi^{p^n}]}$ for all n , so $\lambda = 1$, so uniqueness follows. \square

Let (\mathcal{G}, b, μ) be an integral local shtuka datum such that \mathcal{G} is quasi-parahoric. We define as follows a functor $\widehat{L\mathcal{G}}_{b, \mu}$ on Perfd_k over $\text{Spd}(O_{\check{E}})$, cf. [Gl20]. It assigns to a affinoid perfectoid $S = \text{Spa}(R, R^+)$ over k the set

$$\widehat{L\mathcal{G}}_{b, \mu}(S) = \{\text{isomorphism classes of } ((S^\sharp, y), \mathcal{P}, \psi, \sigma)\}$$

where

- (S^\sharp, y) is an untile of S over $O_{\check{E}}$,
- \mathcal{P} is a \mathcal{G} -torsor over $\text{Spec}(W(R^+))$,
- $\psi : \mathcal{P}[1/\xi_{R^\sharp}] \xrightarrow{\sim} \mathcal{G} \times \text{Spec}(W(R^+)[1/\xi_{R^\sharp}])$, and $\sigma : \mathcal{P} \xrightarrow{\sim} \phi^*(\mathcal{G} \times \text{Spec}(W(R^+)))$ are both \mathcal{G} -isomorphisms such that:
 - 1) (\mathcal{P}, ψ) is bounded by μ along $\xi_{R^\sharp} = 0$,
 - 2) there is a pseudo-uniformizer $\varpi \in R^+$ such that $\phi_b \circ \sigma \equiv \psi \pmod{[\varpi]}$.

Here, “bounded by μ ” means, by definition, that the point of the \mathbb{B}_{dR} -affine Grassmannian $\text{Gr}_{\mathcal{G}, \text{Spd}(\check{O}_E)}$ given by $((S^\sharp, y), \mathcal{P}, \psi)$ factors through the v -sheaf local model $\mathbb{M}_{\mathcal{G}, \mu}^v \subset \text{Gr}_{\mathcal{G}, \text{Spd}(\check{O}_E)}$. More precisely, choose (locally) a section $\tau : \mathcal{G} \xrightarrow{\sim} \mathcal{P}$ and consider $g = \psi \circ \tau(1)$; we ask that $g^{-1} \mathcal{G}(\mathbb{B}_{\text{dR}}^+(S^\sharp))$ lies in $\mathbb{M}_{\mathcal{G}, \mu}^v(R^\sharp)$.

As in [Gl20], we can see that $\widehat{L\mathcal{G}}_{b, \mu}$ is a v -sheaf over $\text{Spd}(O_{\check{E}})$ (this v -sheaf is denoted by $\widehat{\text{LSht}}_{\mathcal{D}}^{\leq \mu}$ in [Gl20, Def. 2.3.15 (3)], and by $\widehat{L\mathcal{M}}_{\mathcal{G}, b, \mu/x_0}^{\text{int}}$ in [PR21]).

It is useful to observe that by using the mapping

$$(\mathcal{P}, \psi, \sigma) \longmapsto h = (\psi \circ \sigma^{-1})(1) \in \mathcal{G}(W(R^+)[1/\xi_{R^\sharp}]),$$

we obtain the following simpler description:

$$\widehat{L\mathcal{G}}_{b, \mu}(S) = \{((S^\sharp, y), h) \mid h \in \mathcal{G}(W(R^+)[1/\xi_{R^\sharp}]), h \equiv b \pmod{[\varpi_h]}, [h^{-1}] \in \mathbb{M}_{\mathcal{G}, \mu}^v(S)\}, \quad (3.6.2)$$

where (S^\sharp, y) is an untile of S over $O_{\check{E}}$, ϖ_h is a pseudo-uniformizer of R^+ , and $[h^{-1}]$ is the S -point of the \mathbb{B}_{dR} -affine Grassmannian $\text{Gr}_{\mathcal{G}, \text{Spd}(\check{O}_E)}$ defined by the coset $h^{-1} \mathcal{G}(\mathbb{B}_{\text{dR}}^+(R^\sharp))$.

Theorem 3.6.2. *There is a diagram of v -sheaves over $\mathrm{Spd}(\check{O}_E)$*

$$\begin{array}{ccc} & \widehat{L\mathcal{G}}_{b,\mu} & \\ \pi_\bullet \swarrow & & \searrow \pi_\star \\ \widehat{\mathcal{M}}_{\mathcal{G},b,\mu/x_0}^{\mathrm{int}} & & \widehat{\mathbb{M}}_{\mathcal{G},\mu/x_0}^v \end{array} \quad (3.6.3)$$

where both π_\bullet, π_\star are $\widehat{L_W^+ \mathcal{G}}$ -torsors (for the v -topology) for two corresponding actions (see (d) below).

a) Here $x_0 \in \widehat{\mathcal{M}}_{\mathcal{G},b,\mu}^{\mathrm{int}}(\mathrm{Spd}(k))$ denotes the base point as in §3.5 above. Similarly $x_0 \in \widehat{\mathbb{M}}_{\mathcal{G},\mu}^v(\mathrm{Spd}(k))$ denotes the corresponding point of $\widehat{\mathbb{M}}_{\mathcal{G},\mu}^v(\mathrm{Spd}(k)) \subset X_{\mathcal{G}}(k)$ given by the pair (\mathcal{G}, b^{-1}) , i.e. by the coset $b^{-1} \mathcal{G}(W(k))$. Also, $\widehat{\mathcal{M}}_{\mathcal{G},b,\mu/x_0}^{\mathrm{int}}$, resp. $\widehat{\mathbb{M}}_{\mathcal{G},\mu/x_0}^v$, denotes the v -sheaf given by the formal completion of $\mathcal{M}_{\mathcal{G},b,\mu}^{\mathrm{int}}$, resp. $\mathbb{M}_{\mathcal{G},\mu}^v$, at these points. (See §3.4 above.)

b) The map π_\star is given by

$$\pi_\star(((S^\sharp, y), \mathcal{P}, \psi, \sigma)) = ((S^\sharp, y), \mathcal{P}, \psi).$$

c) The map π_\bullet is given by

$$\pi_\bullet(((S^\sharp, y), \mathcal{P}, \psi, \sigma)) = ((S^\sharp, y), (\mathcal{P}, \phi_{\mathcal{P}}), i_r).$$

Here, $(\mathcal{P}, \phi_{\mathcal{P}})$ is the \mathcal{G} -shtuka over S with leg at y given as follows: The \mathcal{G} -torsor \mathcal{P} is $(\phi^{-1})^*(\mathcal{P})$ restricted to $\mathcal{Y}_{[0,\infty)}(S)$, i.e.

$$\mathcal{P} = (\phi^{-1})^*(\mathcal{P})|_{\mathcal{Y}_{[0,\infty)}(S)}.$$

The Frobenius $\phi_{\mathcal{P}}$ is given by a similar restriction of the composition

$$\Phi : \mathcal{P}[1/\xi_{R^\sharp}] \xrightarrow{\psi} \mathcal{G} \times \mathrm{Spec}(W(R^+)[1/\xi_{R^\sharp}]) \xrightarrow{((\phi^{-1})^* \sigma)^{-1}} (\phi^{-1})^*(\mathcal{P})[1/\xi_{R^\sharp}].$$

(Note that $\phi^*(\mathcal{P}) = \mathcal{P}|_{\mathcal{Y}_{[0,\infty)}(S)}$. The framing i_r is constructed in [Gl20].

d) The v -sheaf of groups $\widehat{L_W^+ \mathcal{G}}$ acts on $\widehat{L\mathcal{G}}_{b,\mu}$ on the right by

$$(\mathcal{P}, \psi, \sigma) \star g = (\mathcal{P}, \psi, \phi^*(r_{\phi^{-1}(g)}) \circ \sigma), \quad (\mathcal{P}, \psi, \sigma) \bullet g = (\mathcal{P}, r_g \circ \psi, \phi^*(r_g) \circ \sigma),$$

where, r_a is the map given by right multiplication by a , and, for simplicity, we omit the untill (S^\sharp, y) from the notation.

Note that the composition $\Phi : \mathcal{P}[1/\xi_{R^\sharp}] \rightarrow (\phi^{-1})^*(\mathcal{P})[1/\xi_{R^\sharp}]$ in c), is fixed under the \bullet -action. This follows from the identity

$$((\phi^{-1})^*(\phi^*(r_g) \circ \sigma))^{-1} \circ (r_g \circ \psi) = (r_g \circ (\phi^{-1})^* \sigma)^{-1} \circ r_g \circ \psi = ((\phi^{-1})^* \sigma)^{-1} \circ \psi.$$

Proof. The statement of the theorem is shown by Gleason for \mathcal{G} reductive (see [Gl20, Thm. 2.3.14, Lem. 2.3.16, Lem. 2.3.17]). The proof generalizes to \mathcal{G} quasi-parahoric by using Proposition 3.2.1 and its extension Proposition 3.2.2 to products of points. \square

It is convenient to express the torsors π_\star, π_\bullet , using the description (3.6.2) of $\widehat{L\mathcal{G}}_{b,\mu}$. In the description (3.6.2), π_\star is given by projection to the coset $[h^{-1}] = h^{-1} \mathcal{G}(\mathbb{B}_{\mathrm{dR}}^+(R^\sharp))$, and π_\bullet is given by sending h to the shtuka $\mathcal{P}(h) = \mathcal{G} \times \mathcal{Y}_{[0,\infty)}(S) = \mathcal{G}_{\mathcal{Y}_{[0,\infty)}(S)}$ (the trivial torsor) with Frobenius defined by $\phi_{\mathcal{P}(h)} = \phi_h = r_h \phi : (\phi^* \mathcal{G}_{\mathcal{Y}_{[0,\infty)}(S)})[1/\xi_{R^\sharp}] \xrightarrow{\sim} \mathcal{G}_{\mathcal{Y}_{[0,\infty)}(S)}[1/\xi_{R^\sharp}]$. To check

this description of π_\bullet , note that, since $\phi_h = r_h \phi = \psi \circ \sigma^{-1}$, and $\Phi = ((\phi^{-1})^*(\sigma))^{-1} \circ \psi$, the following diagram commutes

$$\begin{array}{ccc} \mathcal{P}[1/\xi_{R^\sharp}] & \xrightarrow{\Phi} & (\phi^{-1})^*(\mathcal{P})[1/\xi_{R^\sharp}] \\ \sigma \downarrow & & \downarrow (\phi^{-1})^* \sigma \\ \phi^*(\mathcal{G}_{W(R^+)})[1/\xi_{R^\sharp}] & \xrightarrow{\phi_h} & \mathcal{G}_{W(R^+)}[1/\xi_{R^\sharp}]. \end{array}$$

Hence, the \mathcal{G} -shtukas $(\mathcal{P}(h), \phi_{\mathcal{P}(h)})$ and $(\mathcal{P}, \phi_{\mathcal{P}})$ given above, are isomorphic.

Under the above isomorphism, the framing i_r of $(\mathcal{P}, \phi_{\mathcal{P}})$ corresponds to a framing $i_r(h)$ of $(\mathcal{P}(h), \phi_{\mathcal{P}(h)})$; this is given by the unique lift of the identity trivialization modulo $[\varpi_h]$, cf. Lemma 3.6.1. More precisely, for $((S^\sharp, y), h) \in \widehat{L\mathcal{G}}_{b,\mu}(S)$, the framing of $(\mathcal{P}(h), \phi_{\mathcal{P}(h)})$ is given by the unique element $i_r(h) \in G(B_{(R,R^+)}^{[r,\infty)})$ with $i(h) \equiv 1 \pmod{[\varpi_h]}$ and the property

$$h = i_r(h)^{-1} \cdot b \cdot \phi(i_r(h)).$$

(See [Gl20, Lem. 2.1.28].) Finally, the actions correspond to $h \star g = g^{-1}h$ and $h \bullet g = \phi(g)^{-1}hg$. Note that $[(h \star g)^{-1}] = [(g^{-1}h)^{-1}] = [h^{-1}g] = [h^{-1}]$.

Remark 3.6.3. a) Gleason calls the diagram (3.6.3) a (v -sheaf) “local model diagram”. However, we warn the reader that (3.6.3) does not compare directly to the local model diagrams in the theory of Shimura varieties and of Rapoport-Zink spaces; these are of a different nature. In particular, the group acting there is \mathcal{G} which is, in a sense, “finite dimensional”, while here we have torsors for $\widehat{L_W^+ \mathcal{G}}$.

b) The existence of the diagram (3.6.3) with the properties listed in the above theorem, does not imply the isomorphism (2.5.1) of Theorem 2.5.5: for example, we cannot deduce from general principles that the torsors π_\star and π_\bullet split, since the formal completions are not “sufficiently local” for the v -topology.

4. PARAHORIC VS QUASI-PARAHORIC GROUP SCHEMES

4.1. The aim of this section. The aim of this section is to prove the following devissage result.

Theorem 4.1.1. *Let (G, b, μ) be a local Shimura datum. The following are equivalent:*

- 1) $\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}$ satisfies the representability conjecture 2.5.1, for all parahoric \mathcal{G} .
- 2) $\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}$ satisfies the representability conjecture 2.5.1, for all stabilizer group schemes \mathcal{G} ,
- 3) $\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}$ satisfies the representability conjecture 2.5.1, for all quasi-parahoric \mathcal{G} .

Here by a stabilizer group scheme we understand the BT-group scheme associated to the stabilizer of a point in the extended building $\mathcal{B}^e(G, \check{\mathbb{Q}}_p)$ (a particular type of quasi-parahoric group schemes, cf. section 2.2). The proof proceeds by the comparing formal completions of $\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}$ and of $\mathcal{M}_{\mathcal{G}^\circ, b, \mu}^{\text{int}}$ (cf. Section 4.2) and of their underlying reduced sets (cf. Section 4.3).

4.2. Formal completions. Let (\mathcal{G}, b, μ) be an integral local shtuka datum such that \mathcal{G} is quasi-parahoric. For simplicity of notation, we set $O = O_{\bar{E}}$ with residue field $k = \bar{k}_E$. We consider the natural v -sheaf morphism

$$\pi : \mathcal{M}_{\mathcal{G}^o, b, \mu}^{\text{int}} \longrightarrow \mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}} \quad (4.2.1)$$

over $\text{Spd}(O)$.

Proposition 4.2.1. *a) The map π is qcqs (quasi-compact quasi-separated [Sch17]).*

b) The map π induces an isomorphism

$$\widehat{\mathcal{M}_{\mathcal{G}^o, b, \mu}^{\text{int}}}_{/x} \xrightarrow{\sim} \widehat{\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}}_{/\pi(x)}$$

for each $x \in \mathcal{M}_{\mathcal{G}^o, b, \mu}^{\text{int}}(\text{Spd}(k))$.

Proof. (a) Observe that $\mathcal{G}^o \rightarrow \mathcal{G}$ is an isomorphism on the generic fibers and can be identified with the dilation \mathcal{G}^{dil} of \mathcal{G} along its closed subscheme given by the neutral component $(\mathcal{G} \otimes_{\mathbb{Z}_p} \mathbb{F}_p)^o$ of the special fiber $\mathcal{G} \otimes_{\mathbb{Z}_p} \mathbb{F}_p$. (Indeed, both \mathcal{G}^o and the dilation \mathcal{G}^{dil} are smooth, have the same generic fiber and the same $\check{\mathbb{Z}}_p$ -points.) In particular, we have $\mathcal{O}_{\mathcal{G}} \hookrightarrow \mathcal{O}_{\mathcal{G}^o} = \mathcal{O}_{\mathcal{G}}[f_1, \dots, f_m]$, and there is $N \geq 1$, such that $p^N f_i \in \mathcal{O}_{\mathcal{G}}$, for all i . Then the argument in [PR21, proof of Prop. 3.6.2] applies to prove the claim.

(b) As in section 3.5 we have an isomorphism

$$\widehat{\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}}_{/x} \simeq \widehat{\mathcal{M}_{\mathcal{G}, b_x, \mu}^{\text{int}}}_{/x_0} \quad (4.2.2)$$

and similarly for \mathcal{G}^o . Hence it is enough to show the isomorphism for the base point $x = x_0 \in \mathcal{M}_{\mathcal{G}^o, b, \mu}^{\text{int}}(\text{Spd}(k))$ (at the cost of changing $b = b_{x_0}$ to b_x). By Theorem 3.6.2

$$\widehat{L\mathcal{G}_{b, \mu}^o} \longrightarrow \widehat{\mathcal{M}_{\mathcal{G}^o, b, \mu}^{\text{int}}}_{/x_0}$$

is a $\widehat{\mathbb{W}}^+ \mathcal{G}^o \times \text{Spd}(O_{\bar{E}})$ -torsor. Also, $\widehat{L\mathcal{G}_{b, \mu}^o}$ is a $\widehat{\mathbb{W}}^+ \mathcal{G}^o \times \text{Spd}(O_{\bar{E}})$ -torsor over $\widehat{\mathbb{M}_{\mathcal{G}^o, \mu}^v}_{/x_0}$, and there are corresponding statements for \mathcal{G} . By (2.3.1), the natural map $\mathcal{G}^o \rightarrow \mathcal{G}$ induces an isomorphism

$$\mathbb{M}_{\mathcal{G}^o, \mu}^v \xrightarrow{\sim} \mathbb{M}_{\mathcal{G}, \mu}^v.$$

On the other hand, it is easy to see that $\mathcal{G}^o \rightarrow \mathcal{G}$ induces $\widehat{\mathbb{W}}^+ \mathcal{G}^o \xrightarrow{\sim} \widehat{\mathbb{W}}^+ \mathcal{G}$ and, hence, $\widehat{L\mathcal{G}_{b, \mu}^o} \xrightarrow{\sim} \widehat{L\mathcal{G}_{b, \mu}}$. The result follows. \square

Remark 4.2.2. Suppose κ is an algebraically closed field over k and set $O(\kappa) = O_{\bar{E}} \otimes_{W(k)} W(\kappa)$. We can consider the base change

$$(\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}})_{O(\kappa)} := \mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}} \times_{\text{Spd}(O)} \text{Spd}(O(\kappa)),$$

and similarly for \mathcal{G}^o . Given $x \in \mathcal{M}_{\mathcal{G}^o, b, \mu}^{\text{int}}(\text{Spd}(\kappa))$, we obtain a corresponding point $x \in (\mathcal{M}_{\mathcal{G}^o, b, \mu}^{\text{int}})_{O(\kappa)}(\text{Spd}(\kappa))$. The formal completions $(\widehat{\mathcal{M}_{\mathcal{G}^o, b, \mu}^{\text{int}}})_{O(\kappa)/x}$ and $(\widehat{\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}})_{O(\kappa)/\pi(x)}$ now make sense and the argument in the proof of (b) above also applies to give

$$(\widehat{\mathcal{M}_{\mathcal{G}^o, b, \mu}^{\text{int}}})_{O(\kappa)/x} \xrightarrow{\sim} (\widehat{\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}})_{O(\kappa)/\pi(x)}. \quad (4.2.3)$$

4.3. ADLV for quasi-parahorics. In this subsection, we express the ADLV $X_{\mathcal{G}}(b, \mu)$ for a quasi-parahoric \mathcal{G} in terms of ADLV attached to parahoric subgroups. Note that the results will eventually be applied for μ replaced by μ^{-1} , to relate to integral local Shimura varieties via Prop. 3.3.1.

Recall the Kottwitz map $\kappa_G : G(\check{\mathbb{Q}}_p) \rightarrow \Omega_G$. This map can be enhanced to a morphism $LG \rightarrow \Omega_G$ which factors through $X_{\mathcal{G}^\circ} = LG/L^+\mathcal{G}^\circ$, and induces a map with connected fibers,

$$\kappa_G : X_{\mathcal{G}^\circ} \longrightarrow \Omega_G, \quad (4.3.1)$$

cf. [Zhu17, Prop. 1.21], comp. also [PR08, §5].

Recall $\pi_0(\mathcal{G}) = \check{K}/\check{K}^\circ \subset \Omega$, cf. (3.1.1). Then $\pi_0(\mathcal{G})$ acts on $X_{\mathcal{G}^\circ}$ by $g\check{K}^\circ \mapsto g\dot{\gamma}\check{K}^\circ$, where $\dot{\gamma} \in \check{K}$ is a representative of a given element in $\pi_0(\mathcal{G})$. This action is permuting connected components and the quotient is $X_{\mathcal{G}}$,

$$X_{\mathcal{G}^\circ} \longrightarrow X_{\mathcal{G}} = X_{\mathcal{G}^\circ}/\pi_0(\mathcal{G}).$$

Each connected component of $X_{\mathcal{G}^\circ}$ maps isomorphically to a connected component of $X_{\mathcal{G}}$.

Now $X_{\mathcal{G}^\circ}(b, \mu)$ lies in the union of those connected components of $X_{\mathcal{G}^\circ}$, where κ_G falls in the class of $\kappa_G(b) - \mu^\sharp$ in $\pi_1(G)_\Gamma = \Omega_\phi$. Here $\mu^\sharp \in \Omega_\phi$ is the element associated to the conjugacy class μ , i.e., the residue class of $\kappa(\mu) \in \Omega$. Use the exact sequence

$$1 \longrightarrow \Omega^\phi \longrightarrow \Omega \xrightarrow{\phi-1} \Omega \longrightarrow \Omega_\phi \longrightarrow 1$$

to deduce that this is a coset of Ω^ϕ of the form $c_{b,\mu} + \Omega^\phi \subset \Omega$, where $c_{b,\mu} \in \Omega$ is an element with $\phi(c_{b,\mu}) - c_{b,\mu} = \kappa_G(b) - \mu^\sharp$, comp. [HZ20, §6]. In fact, by [HZ20, Lem. 6.1], the inclusion is an equality,

$$\kappa_G(X_{\mathcal{G}^\circ}(b, \mu)) = c_{b,\mu} + \Omega^\phi. \quad (4.3.2)$$

We now extend these considerations to quasi-parahorics. Let $C = C_{\mathcal{G}} = \Omega/\pi_0(\mathcal{G})$. Then the Kottwitz homomorphism induces a map

$$\kappa_G : X_{\mathcal{G}} \longrightarrow C_{\mathcal{G}}. \quad (4.3.3)$$

The image of $X_{\mathcal{G}}(b, \mu)$ in C_ϕ is equal to the residue class of $\kappa_G(b) - \mu^\sharp$, so

$$\kappa_G(X_{\mathcal{G}}(b, \mu)) \subset \bar{c}_{b,\mu} + C^\phi, \quad (4.3.4)$$

where $\bar{c}_{b,\mu} \in C_{\mathcal{G}}$ is the image of $c_{b,\mu}$. This inclusion is an equality, cf. Corollary 4.3.6 below.

Recall $\Pi_{\mathcal{G}} = \ker(\pi_0(\mathcal{G})_\phi \rightarrow \Omega_\phi)$, cf. (3.1.2). From the exact sequence $0 \rightarrow \pi_0(\mathcal{G}) \rightarrow \Omega \rightarrow C_{\mathcal{G}} \rightarrow 0$ we have an exact sequence

$$0 \longrightarrow \Omega^\phi/\pi_0(\mathcal{G})^\phi \longrightarrow C_{\mathcal{G}}^\phi \longrightarrow \Pi_{\mathcal{G}} \longrightarrow 0. \quad (4.3.5)$$

Let $\bar{\beta} \in \Pi_{\mathcal{G}}$. Take $\beta \in \pi_0(\mathcal{G}) \subset \Omega$ lifting $\bar{\beta}$. Then

$$\beta = (1 - \phi)\gamma$$

for some $\gamma \in \Omega$. Note that, since $\phi(\gamma) - \gamma$ is in $\pi_0(\mathcal{G})$, the image of γ in C is ϕ -invariant and maps to $\bar{\beta}$ under the connecting homomorphism $C^\phi \rightarrow \Pi_{\mathcal{G}}$.

Recall that the quasi-parahoric subgroup \check{K} is defined via a point \mathbf{x} in the building of G_{ad} . Write the Iwahori-Weyl group as $\widetilde{W} = W_a \rtimes \Omega$ and

$$1 \longrightarrow T(\check{\mathbb{Q}}_p)_1 \longrightarrow N(\check{\mathbb{Q}}_p) \longrightarrow \widetilde{W} = W_a \rtimes \Omega \longrightarrow 1$$

with $T = Z_G(S_{\check{\mathbb{Q}}_p})$ as in [HR08]. Here the choices are made such that \mathbf{x} is in the base alcove \mathfrak{a}_0 of the apartment of S . Then $T(\check{\mathbb{Q}}_p)_1 \subset \check{K}^o$. When we consider Ω as a subset of \widetilde{W} , we write the group law in a multiplicative way.

Lemma 4.3.1. *There is a lift $\dot{\gamma} \in N(\check{\mathbb{Q}}_p)$ of $\gamma \in \Omega \subset \widetilde{W}$ such that*

$$\phi(\dot{\gamma})^{-1}\dot{\gamma} \in \check{K}.$$

Proof. Lift $\gamma \in \Omega$ to $\dot{\gamma} \in N(\check{\mathbb{Q}}_p)$. Then $\phi(\dot{\gamma})^{-1}\dot{\gamma}$ lifts $\beta \in \pi_0(\mathcal{G}) \subset \Omega$. Now we use the exact sequence

$$1 \longrightarrow T(\check{\mathbb{Q}}_p)_1 \longrightarrow N(\check{\mathbb{Q}}_p) \cap \check{K} \longrightarrow \pi_0(\mathcal{G}) \longrightarrow 1.$$

This holds, since, denoting by $U = U_{K^o}$ the pro-unipotent radical of \check{K}^o , we have $\check{K} \subset U \cdot N(\check{\mathbb{Q}}_p)$, cf. [HR08, proof of Prop. 8]. We can lift β to $\dot{\beta} \in N(\check{\mathbb{Q}}_p) \cap \check{K}$. Then $\phi(\dot{\gamma})^{-1}\dot{\gamma}\dot{\beta}^{-1} \in T(\check{\mathbb{Q}}_p)_1$. Since $T(\check{\mathbb{Q}}_p)_1 \subset \check{K}^o$, the result follows. \square

It follows that

$$\check{K}\dot{\gamma}^{-1} = \check{K}\phi(\dot{\gamma})^{-1}.$$

Consider now the conjugates of \check{K} , resp. \check{K}^o ,

$$\check{K}_\gamma := \dot{\gamma}\check{K}\dot{\gamma}^{-1}, \quad \check{K}_\gamma^o := \dot{\gamma}\check{K}^o\dot{\gamma}^{-1}. \quad (4.3.6)$$

We have

$$\phi(\dot{\gamma}\check{K}\dot{\gamma}^{-1}) = \dot{\gamma}\check{K}\dot{\gamma}^{-1}, \quad \phi(\dot{\gamma}\check{K}^o\dot{\gamma}^{-1}) = \dot{\gamma}\check{K}^o\dot{\gamma}^{-1}.$$

Hence, \check{K}_γ and \check{K}_γ^o are rational, i.e., correspond to subgroups of $G(\mathbb{Q}_p)$. For varying β , the parahorics K_γ^o are conjugate to K^o in $G(\check{\mathbb{Q}}_p)$ but not necessarily in $G(\mathbb{Q}_p)$.

Proposition 4.3.2. *The subgroups \check{K}_γ and \check{K}_γ^o of $G(\check{\mathbb{Q}}_p)$ are independent of the lift $\dot{\gamma}$ of γ . Furthermore, their $G(\mathbb{Q}_p)$ -conjugacy class only depends on the class β in $\Pi_{\mathcal{G}}$.*

Proof. We go through all choices made in the construction.

- *independence of $\dot{\gamma}$:* If $\dot{\gamma}$ is replaced by $\dot{\gamma}\dot{\delta}$, with $\dot{\delta} \in T(\check{\mathbb{Q}}_p)_1$, then $\dot{\gamma}\check{K}^o\dot{\gamma}^{-1}$ is replaced by $\dot{\gamma}\dot{\delta}\check{K}^o\dot{\delta}^{-1}\dot{\gamma}^{-1} = \dot{\gamma}\check{K}^o\dot{\gamma}^{-1}$.
- *independence of γ :* If γ is replaced by $\gamma + \delta$, with $\delta \in \Omega^\phi$, then as choice for $\dot{\gamma}$, we can take $\dot{\delta}\dot{\gamma}$, where $\dot{\delta} \in N(\mathbb{Q}_p) \cap \check{K}$ (i.e., \mathbb{Q}_p -rational). This follows from [HR08, Rem. 9]. Hence $\dot{\gamma}\check{K}^o\dot{\gamma}^{-1}$ is replaced by $\delta\dot{\gamma}\check{K}^o\dot{\gamma}^{-1}\delta^{-1}$, hence is conjugate under $G(\mathbb{Q}_p)$ to \check{K}_β^o .
- *independence of β :* If β is replaced by $\beta + (\phi - 1)\delta$ with $\delta \in \pi_0(\mathcal{G})$, then γ is replaced by $\gamma + \delta$. Then we may replace $\dot{\gamma}$ by $\dot{\gamma}\dot{\delta}$, where $\dot{\delta} \in N(\check{\mathbb{Q}}_p) \cap \check{K}$. Since $\dot{\delta}$ normalizes \check{K}^o , the parahoric \check{K}_β^o is unchanged.

\square

Remark 4.3.3. The above proof seems to use that we multiply $\dot{\gamma}$ by $\dot{\delta}$ from the right instead of from the left. However, Ω is an abelian group and what is being used here is that in the first instance $\dot{\delta} \in T(\check{\mathbb{Q}}_p)_1$ acts trivially on \mathfrak{a}_0 , and in the third instance that $\dot{\delta} \in N(\check{\mathbb{Q}}_p) \cap \check{K}$ preserves \mathfrak{a}_0 and fixes x and hence also $\dot{\gamma}x$, by the commutativity of Ω .

In the sequel, we make a fixed choice of $\dot{\gamma}$, for each element $\bar{\beta} \in \Pi_{\mathcal{G}}$. We denote the corresponding groups by \check{K}_{β} , resp. $\check{\mathcal{G}}_{\beta}$, resp. $\check{K}_{\beta}^{\circ}$, resp. $\check{\mathcal{G}}_{\beta}^{\circ}$, by slightly abusing notation. We consider the map

$$\begin{aligned} X_{\mathcal{G}_{\beta}} &= LG/L^{+}\check{K}_{\beta} = LG/L^{+}(\dot{\gamma}\check{K}\dot{\gamma}^{-1}) \longrightarrow LG/L^{+}\check{K} = X_{\mathcal{G}}, \\ g(\dot{\gamma}\check{K}\dot{\gamma}^{-1}) &\longmapsto g\dot{\gamma}\check{K}. \end{aligned} \quad (4.3.7)$$

Proposition 4.3.4. *The above map defines by restriction a map*

$$\pi_{\beta} : X_{\mathcal{G}_{\beta}}(b, \mu) \longrightarrow X_{\mathcal{G}}(b, \mu).$$

Proof. Suppose that $g\check{K}_{\beta} \in X_{\mathcal{G}_{\beta}}(b, \mu)$, i.e.,

$$g^{-1}b\phi(g) \in \check{K}_{\beta}\text{Adm}(\mu)\check{K}_{\beta} = \dot{\gamma}\check{K}\dot{\gamma}^{-1}\text{Adm}(\mu)\dot{\gamma}\check{K}\dot{\gamma}^{-1}.$$

Since $\text{Adm}(\mu)$ is stable under the conjugation action of Ω , we see that

$$\check{K}\dot{\gamma}^{-1}\text{Adm}(\mu)\dot{\gamma}\check{K} = \check{K}\text{Adm}(\mu)\check{K}.$$

Hence

$$g^{-1}b\phi(g) \in \dot{\gamma}\check{K}\text{Adm}(\mu)\check{K}\dot{\gamma}^{-1},$$

so

$$\dot{\gamma}^{-1}g^{-1}b\phi(g)\phi(\dot{\gamma}) \in \check{K}\text{Adm}(\mu)\check{K}\dot{\gamma}^{-1}\phi(\dot{\gamma}) \subset \check{K}\text{Adm}(\mu)\check{K},$$

since $\check{K}\dot{\gamma}^{-1}\phi(\dot{\gamma}) = \check{K}$. Hence

$$(g\dot{\gamma})^{-1}b\phi(g\dot{\gamma}) \in \check{K}\text{Adm}(\mu)\check{K},$$

i.e., $g\dot{\gamma}\check{K} \in X_{\mathcal{G}}(b, \mu)$, as had to be shown. \square

Composing the above map with $X_{\mathcal{G}_{\beta}}(b, \mu) \rightarrow X_{\mathcal{G}_{\beta}}(b, \mu)$ and letting $\bar{\beta}$ vary, we obtain the map

$$\pi : \bigsqcup_{\bar{\beta} \in \Pi_{\mathcal{G}}} X_{\mathcal{G}_{\beta}}(b, \mu) \longrightarrow X_{\mathcal{G}}(b, \mu). \quad (4.3.8)$$

In the sequel, we call a *component* of $X_{\mathcal{G}_{\beta}}(b, \mu)$, resp. of $X_{\mathcal{G}}(b, \mu)$, the non-empty fibers of the Kottwitz map to Ω , resp. to $C_{\mathcal{G}} = \Omega/\pi_0(\mathcal{G})$.

Proposition 4.3.5. (i) *For $\bar{\beta}_1 \neq \bar{\beta}_2$, the images of $X_{\mathcal{G}_{\beta_1}}(b, \mu)$ and $X_{\mathcal{G}_{\beta_2}}(b, \mu)$ under π fall into different components of $X_{\mathcal{G}}(b, \mu)$.*

(ii) *The map π is surjective.*

Proof. For (i), let $x_1 \in X_{\mathcal{G}_{\beta_1}}(b, \mu)$ and $x_2 \in X_{\mathcal{G}_{\beta_2}}(b, \mu)$ be such that their images in $C_{\mathcal{G}}$ are the same. We deduce that

$$\kappa_G(x_1) + \gamma_1 = \kappa_G(x_2) + \gamma_2 \pmod{\pi_0(\mathcal{G})}.$$

On the other hand, $\kappa_G(x_i) = c_{b, \mu} - \lambda_i$ with $\lambda_i \in \Omega^{\phi}$, for $i = 1, 2$. We therefore get

$$\gamma_1 - \gamma_2 = \lambda_1 - \lambda_2 + \mu, \quad \mu \in \pi_0(\mathcal{G}).$$

Applying $1 - \phi$ to this identity, we obtain

$$\beta_1 - \beta_2 = (1 - \phi)\mu,$$

i.e., $\bar{\beta}_1 = \bar{\beta}_2$, as desired.

For (ii), let $g\check{K} \in X_{\mathcal{G}}(b, \mu)$. By [GHN20, Lem. 5.10, (iii)],

$$\check{K}^o \text{Adm}(\mu) \check{K} = \check{K} \text{Adm}(\mu) \check{K}. \quad (4.3.9)$$

Hence $g^{-1}b\phi(g) \in \check{K} \text{Adm}(\mu) \check{K} = \check{K}^o \text{Adm}(\mu) \check{K}$. Write accordingly

$$g^{-1}b\phi(g) = k_1^o \omega k.$$

Apply $\kappa : G(\check{\mathbb{Q}}_p) \rightarrow \Omega$ to get

$$(\phi - 1)\kappa(g) + \kappa(b) = \kappa(k) + \kappa(\mu).$$

Hence we get

$$(\phi - 1)\kappa(g) = \kappa(k) - (\phi - 1)c_{b, \mu}. \quad (4.3.10)$$

Let $\beta = -\kappa(k) \in \pi_0(\mathcal{G})$ and denote by $\bar{\beta} \in \pi_0(\mathcal{G})_\phi$ its image. Then the last equation shows that $\bar{\beta} \in \Pi_{\mathcal{G}}$. We again write $\beta = (1 - \phi)\gamma$ and choose a lift $\dot{\gamma} \in N(\check{\mathbb{Q}}_p) \cap \check{K}$. Then $g\dot{\gamma}^{-1}(\dot{\gamma}K^o\dot{\gamma}^{-1})$ is in $X_{\dot{\gamma}K^o\dot{\gamma}^{-1}}(b, \mu)$ and maps to $gK \in X_K(b, \mu)$. Indeed, $(g\dot{\gamma}^{-1})^{-1}b\phi(g\dot{\gamma}^{-1})$ lies in $(\dot{\gamma}K^o\dot{\gamma}^{-1})\text{Adm}(\mu)(\dot{\gamma}K^o\dot{\gamma}^{-1})$ because

$$\begin{aligned} (g\dot{\gamma}^{-1})^{-1}b\phi(g\dot{\gamma}^{-1}) &= \dot{\gamma}g^{-1}b\phi(g)\phi(\dot{\gamma}^{-1}) = \dot{\gamma}k_1^o\omega k\phi(\dot{\gamma}^{-1}) = \\ &= (\dot{\gamma}k_1^o\dot{\gamma}^{-1})(\dot{\gamma}\omega\dot{\gamma}^{-1})(\dot{\gamma}k\phi(\dot{\gamma}^{-1})). \end{aligned}$$

The last factor lies in $\dot{\gamma}\check{K}\dot{\gamma}^{-1}$. The result follows because the last factor lies even in $\dot{\gamma}\check{K}^o\dot{\gamma}^{-1}$ because $\kappa(\dot{\gamma}k\phi(\dot{\gamma}^{-1})) = (1 - \phi)\gamma - \beta = 0$. \square

Corollary 4.3.6. *The map κ induces a surjective map $X_{\mathcal{G}}(b, \mu) \rightarrow \bar{c}_{b, \mu} + C^\phi$.*

Proof. Let $C^\phi \rightarrow \Pi_{\mathcal{G}}$ be the natural surjective map from (4.3.5), and let $(C^\phi)_{\bar{\beta}}$ be the fiber over $\bar{\beta} \in \Pi_{\mathcal{G}}$. Via our fixed choice of γ for $\bar{\beta}$, this can be identified with $\Omega^\phi/\pi_0(\mathcal{G}_\beta)^\phi$. By (4.3.2), the set of components of the inverse image of $\bar{c}_{b, \mu} + (C^\phi)_{\bar{\beta}}$ in $X_{\mathcal{G}_\beta}(b, \mu)$ can then be identified with $c_{b, \mu} + \Omega^\phi$. The assertion follows. \square

We introduce the intermediate group $K^o \subset K' \subset K$, corresponding to the subgroup $\pi_0(\mathcal{G})^\phi$ of $\pi_0(\mathcal{G})$. We denote the corresponding group scheme by \mathcal{G}' . Then we obtain a factorization

$$X_{\mathcal{G}^o} \longrightarrow X_{\mathcal{G}'} \longrightarrow X_{\mathcal{G}}.$$

Each of the maps induces an isomorphism of a component onto its image. In fact, the first map is the quotient by the finite abelian group $\pi_0(\mathcal{G})^\phi$, the second map is the quotient by the finite abelian group $\pi_0(\mathcal{G})/\pi_0(\mathcal{G})^\phi$, and the composed map is the quotient by the finite abelian group $\pi_0(\mathcal{G})$. Here the actions of these covering groups is trivial, in the sense that they only permute connected components.

Similarly, for arbitrary $\bar{\beta} \in \Pi_{\mathcal{G}}$, we introduce the intermediate subgroup $K_\beta^o \subset K'_\beta \subset K_\beta$, corresponding to the subgroup $\pi_0(\mathcal{G}_\beta)^\phi$ of $\pi_0(\mathcal{G}_\beta)$. We obtain a sequence of maps,

$$X_{\mathcal{G}_\beta^o} \longrightarrow X_{\mathcal{G}'_\beta} \longrightarrow X_{\mathcal{G}_\beta}, \quad (4.3.11)$$

where the first map is the quotient by the finite group $\pi_0(\mathcal{G}_\beta)^\phi$, the second map is the quotient by the finite abelian group $\pi_0(\mathcal{G}_\beta)/\pi_0(\mathcal{G}_\beta)^\phi$, and the composed map is the quotient by the finite abelian group $\pi_0(\mathcal{G}_\beta)$.

Proposition 4.3.7. *Let (\mathcal{G}, b, μ) be an integral local shtuka datum such that \mathcal{G} is a quasi-parahoric group scheme for G .*

- (i) *The ADLV $X_{\mathcal{G}_\beta^\circ}(b, \mu)$ is invariant under the action of $\pi_0(\mathcal{G}_\beta)^\phi$.*
- (ii) *The map (4.3.8) induces an isomorphism*

$$\bar{\pi} : \bigsqcup_{\bar{\beta} \in \Pi_{\mathcal{G}}} X_{\mathcal{G}_\beta^\circ}(b, \mu) / \pi_0(\mathcal{G}_\beta)^\phi \xrightarrow{\sim} X_{\mathcal{G}}(b, \mu).$$

Proof. (i) Let $\gamma \in \check{K}'_\beta$. Then $\phi(\gamma) = \gamma k$, with $k \in \check{K}_\beta^\circ$. The invariance follows from

$$(g\gamma)^{-1}b\phi(g\gamma) = \gamma^{-1}g^{-1}b\phi(g)\phi(\gamma) = \gamma^{-1}g^{-1}b\phi(g)\gamma k,$$

because the last element lies in $\check{K}_\beta^\circ \gamma^{-1} \text{Adm}(\mu) \gamma \check{K}_\beta^\circ = \check{K}_\beta^\circ \text{Adm}(\mu) \check{K}_\beta^\circ$.

(ii) By Proposition 4.3.5, (i), this becomes a question one $\bar{\beta}$ at the time. But if the components of $X_{\mathcal{G}_\beta^\circ}(b, \mu)$ corresponding to λ_1 and λ_2 in Ω go to the same component of $X_{\mathcal{G}}(b, \mu)$, it follows that $\lambda_1 - \lambda_2 \in \pi_0(\mathcal{G})$. But by (4.3.2), this element lies in Ω^ϕ , hence also in $\pi_0(\mathcal{G})^\phi$. \square

Remark 4.3.8. Proposition 4.3.5 and Corollary 4.3.6 imply that the map $X_{\mathcal{G}^\circ}(b, \mu) \rightarrow X_{\mathcal{G}}(b, \mu)$ is surjective if and only if $\Pi_{\mathcal{G}}$ is trivial. The question of the surjectivity of this map is also analyzed in [GHN20, §5]. By [GHN20, Prop. 5.11] it holds if there is a product decomposition $\Omega = \pi_0(\mathcal{G}) \times C$ into subgroups stable under ϕ (then, of course, $\Pi_{\mathcal{G}} = (0)$). In [GHN20, §5] it is also analyzed when $X_{\mathcal{G}^\circ}(b, \mu)$ is the full inverse image of $X_{\mathcal{G}}(b, \mu)$ under $X_{\mathcal{G}^\circ} \rightarrow X_{\mathcal{G}}$. By [GHN20, Prop. 5.13] this holds when $\pi_0(\mathcal{G})^\phi = \pi_0(\mathcal{G})$ and $\Omega^\phi \rightarrow C_{\mathcal{G}}^\phi$ is surjective (again, these conditions imply $\Pi_{\mathcal{G}} = (0)$).

4.4. Integral moduli spaces of local shtukas. The map $\mathcal{G}_\beta^\circ \rightarrow \mathcal{G}$, given by $g \mapsto \dot{\gamma}^{-1}g\dot{\gamma}$ induces a morphism on integral LSV

$$\mathcal{M}_{\mathcal{G}_\beta^\circ, b, \mu}^{\text{int}} \longrightarrow \mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}},$$

and hence a morphism

$$\pi : \bigsqcup_{\bar{\beta} \in \Pi_{\mathcal{G}}} \mathcal{M}_{\mathcal{G}_\beta^\circ, b, \mu}^{\text{int}} \longrightarrow \mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}, \quad (4.4.1)$$

compatible, via the specialization map from the integral LSV to the ADLV, with the morphism (4.3.8) for μ^{-1} .

Theorem 4.4.1. *Let (\mathcal{G}, b, μ) be an integral local shtuka datum such that \mathcal{G} is a quasi-parahoric group scheme for G . The morphism (4.4.1) induces an isomorphism*

$$\bar{\pi} : \bigsqcup_{\bar{\beta} \in \Pi_{\mathcal{G}}} \mathcal{M}_{\mathcal{G}_\beta^\circ, b, \mu}^{\text{int}} / \pi_0(\mathcal{G}_\beta)^\phi \xrightarrow{\sim} \mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}},$$

where the quotient is for an action of $\pi_0(\mathcal{G}_\beta)^\phi$ which permutes connected components.

Let us first define the action of $\pi_0(\mathcal{G}_\beta)^\phi$ on $\mathcal{M}_{\mathcal{G}_\beta^\circ, b, \mu}^{\text{int}}$. We use the exactness of the sequence

$$0 \longrightarrow \mathcal{G}_\beta^\circ(\mathbb{Z}_p) \longrightarrow \mathcal{G}_\beta(\mathbb{Z}_p) \longrightarrow \pi_0(\mathcal{G}_\beta)^\phi \longrightarrow 0.$$

Let $\gamma \in G(\mathbb{Q}_p)$ be in the normalizer of \mathcal{G}_β° . Recall that

$$\mathcal{M}_{\mathcal{G}^\circ, b, \mu}^{\text{int}}(S) = \{\text{isomorphism classes of } (S^\sharp, \mathcal{P}, \phi_{\mathcal{P}}, i_r)\}.$$

We define a new \mathcal{G}_β^o -torsor $\mathcal{P}' = \mathcal{P}_\gamma$ by twisting the \mathcal{G}_β^o -action by γ ,

$$g \star x = (\gamma^{-1}g\gamma) \cdot x.$$

The new Frobenius $\phi_{\mathcal{P}'}$ is taken to be identical to $\phi_{\mathcal{P}}$. This is indeed a morphism of \mathcal{G}_β^o -torsors since

$$\phi_{\mathcal{G}'}(g \star x) = \phi_{\mathcal{G}}((\gamma^{-1}g\gamma) \cdot x) = (\gamma^{-1}\phi(g)\gamma) \cdot \phi_{\mathcal{G}}(x) = \phi(g) \star \phi_{\mathcal{G}'}(x),$$

where we used that $\gamma \in G(\mathbb{Q}_p)$. Finally, the framing i'_r is given by $i'_r(g) = i_r(\gamma^{-1}g\gamma)$. This is indeed an isomorphism of \mathcal{G}_β^o -bundles, since

$$i'_r(hg) = i_r(\gamma^{-1}hg\gamma) = i_r((\gamma^{-1}h\gamma)(\gamma^{-1}g\gamma)) = (\gamma^{-1}h\gamma) \cdot i_r(\gamma^{-1}g\gamma) = h \star i'_r(g).$$

This expression makes sense since $\gamma^{-1}h\gamma \in \mathcal{G}_\beta^o$.

Note that if $\gamma \in \mathcal{G}_\beta^o(\mathbb{Z}_p)$, then the tuple $(S^\sharp, \mathcal{P}', \phi_{\mathcal{P}'}, i'_r)$ is isomorphic to $(S^\sharp, \mathcal{P}, \phi_{\mathcal{P}}, i_r)$. Indeed, the map $x \mapsto \gamma \cdot x$ defines an isomorphism $\alpha : \mathcal{P} \rightarrow \mathcal{P}'$ compatible with $\phi_{\mathcal{G}}$ and $\phi_{\mathcal{G}'}$, and with i_r and i'_r . For instance

$$\phi_{\mathcal{G}'}(\alpha(x)) = \phi_{\mathcal{G}}(\gamma \cdot x) = \phi(\gamma) \cdot \phi_{\mathcal{G}}(x) = \gamma \cdot \phi_{\mathcal{G}}(x) = \alpha(\phi_{\mathcal{G}}(x)).$$

Therefore we obtain an action of the factor group $N_{G(\mathbb{Q}_p)}(\mathcal{G}_\beta^o)/\mathcal{G}_\beta^o(\mathbb{Z}_p)$ on $\mathcal{M}_{\mathcal{G}_\beta^o, b, \mu}^{\text{int}}$. Restricting this action to $\mathcal{G}_\beta^o(\mathbb{Z}_p)/\mathcal{G}_\beta^o(\mathbb{Z}_p)$, we obtain the desired action of $\pi_0(\mathcal{G})^\phi$ on $\mathcal{M}_{\mathcal{G}_\beta^o, b, \mu}^{\text{int}}$.

The same argument as the one above shows that $\gamma \in \mathcal{G}_\beta(\mathbb{Z}_p)$ induces an isomorphism between the images of $(S^\sharp, \mathcal{P}, \phi_{\mathcal{P}}, i_r)$ and $(S^\sharp, \mathcal{P}', \phi_{\mathcal{P}'}, i'_r)$ in $\mathcal{M}_{\mathcal{G}_\beta, b, \mu}^{\text{int}}$, hence we obtain a morphism $\mathcal{M}_{\mathcal{G}_\beta^o, b, \mu}^{\text{int}}/\pi_0(\mathcal{G}_\beta)^\phi \rightarrow \mathcal{M}_{\mathcal{G}_\beta, b, \mu}^{\text{int}}$. Letting now $\bar{\beta}$ vary, we obtain a morphism from the LHS to the RHS in Theorem 4.4.1.

Proof of Theorem 4.4.1: We deduce from Proposition 3.3.1 and Proposition 4.3.7 that the morphism induces a bijection on the set of $\text{Spd}(\kappa)$ -points, for all algebraically closed extensions κ/k . Also it is qcqs by Proposition 4.2.1 (a). Using relative properness and [SW20, Cor. 17.4.10] we see it is enough to check that it gives a bijection on $\text{Spa}(C, O_C)$ -points. Note that any $\text{Spa}(C, O_C)$ -point \tilde{x} of $\mathcal{M}_{\mathcal{G}_\beta, b, \mu}^{\text{int}}$ factors through the formal completion $(\widehat{\mathcal{M}_{\mathcal{G}_\beta, b, \mu}^{\text{int}}})_{O(k_C)/\bar{x}}$ of the base change of $\mathcal{M}_{\mathcal{G}_\beta, b, \mu}^{\text{int}}$ by $O(k_C) = O_{\tilde{E}} \otimes_{W(k)} W(k_C)$, at its specialization at $\bar{x} = \text{sp}(\tilde{x})$. Here, $k_C = O_C/\mathfrak{m}_C$ is the residue field. The corresponding fact is also true for $\text{Spa}(C, O_C)$ -points of $\mathcal{M}_{\mathcal{G}_\beta^o, b, \mu}^{\text{int}}$.

We can now complete the proof. First we show injectivity: Suppose \tilde{x}_i , $i = 1, 2$, are two points in $(\sqcup_{\bar{\beta} \in \Pi_{\mathcal{G}}} \mathcal{M}_{\mathcal{G}_\beta^o, b, \mu}^{\text{int}}/\pi_0(\mathcal{G}_\beta)^\phi)(C, O_C)$, mapping by $\bar{\pi}$ to the same point in $\mathcal{M}_{\mathcal{G}_\beta, b, \mu}^{\text{int}}(C, O_C)$. By Proposition 4.3.7, these two points have the same specialization $\bar{x} = \bar{x}_1 = \bar{x}_2$. By the above, they factor through the formal completion of (the base change) of $\mathcal{M}_{\mathcal{G}_\beta, b, \mu}^{\text{int}}$ at a corresponding \bar{x} , for some common $\bar{\beta} \in \Pi_{\mathcal{G}}$. By (4.2.3) for $\kappa = k_C$, we see that these two points agree in the formal completion of $\mathcal{M}_{\mathcal{G}_\beta, b, \mu}^{\text{int}}$ at \bar{x} , and therefore $\tilde{x}_1 = \tilde{x}_2$. The proof of surjectivity onto $\mathcal{M}_{\mathcal{G}_\beta, b, \mu}^{\text{int}}(C, O_C)$ is by a similar argument. \square

We define a map of v -sheaves

$$\kappa_{\mathcal{G}} \circ \text{sp} : \mathcal{M}_{\mathcal{G}_\beta, b, \mu}^{\text{int}} \longrightarrow \underline{C_{\mathcal{G}}},$$

where $\text{sp}: |\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}| \rightarrow |X_{\mathcal{G}}(b, \mu^{-1})|$ denotes the (continuous) specialization map under the identification $(\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}})_{\text{red}} = X_{\mathcal{G}}(b, \mu^{-1})$, cf. Proposition 3.3.1. For $\tau \in C_{\mathcal{G}}$, we denote by

$$\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau} = \mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}} \times_{C_{\mathcal{G}}} \{\tau\} \subset \mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}} \quad (4.4.2)$$

the fiber over τ . This is an open and closed v -subsheaf of $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}$. We call $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau}$ the *component* of $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}$ corresponding to τ .

Corollary 4.4.2. *Let (\mathcal{G}, b, μ) be an integral local shtuka datum such that \mathcal{G} is a quasi-parahoric group scheme for G and let $\mathcal{G} \rightarrow \mathcal{G}'$ be a morphism extending the identity morphism of G in the generic fibers of quasi-parahoric group schemes for G with the same associated parahoric group scheme. Then for every $\tau \in \Omega_G/\pi_0(\mathcal{G})$, with image $\tau' \in \Omega_G/\pi_0(\mathcal{G}')$, the natural morphism $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}} \rightarrow \mathcal{M}_{\mathcal{G}',b,\mu}^{\text{int}}$ induces an isomorphism*

$$\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau} \xrightarrow{\sim} \mathcal{M}_{\mathcal{G}',b,\mu}^{\text{int},\tau'}$$

of v -sheaves. □

Proof of Theorem 4.1.1: For the representability conjecture, we assume that (G, b, μ) is a local Shimura datum, i.e., μ is minuscule. Now $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}$ is representable if and only if $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau}$ is representable for every $\tau \in \Omega_G/\pi_0(\mathcal{G})$. Hence the assertion follows from Corollary 4.4.2 and the fact that every component $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau}$ is isomorphic to a component $\mathcal{M}_{\mathcal{G}^o,b,\mu}^{\text{int},\tilde{\tau}}$ of $\mathcal{M}_{\mathcal{G}^o,b,\mu}^{\text{int}}$ for $\tilde{\tau} \in \Omega_G$ mapping to τ and, conversely, every component $\mathcal{M}_{\mathcal{G}^o,b,\mu}^{\text{int},\tilde{\tau}}$ is isomorphic to the component $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\bar{\tau}}$ of $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}$, where $\bar{\tau} \in \Omega_G/\pi_0(\mathcal{G})$ is the image of $\tilde{\tau}$. □

4.5. Generic fibers. In this subsection, we interpret the induced decomposition in Theorem 4.4.1 in the generic fibers, in the case when μ is minuscule. Let (G, b, μ) be a local Shimura datum and let \mathcal{G} be a quasi-parahoric for G . We consider the generic fiber

$$(\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}})_{\eta} = \mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}} \times_{\text{Spd}(O_{\check{E}})} \text{Spd}(\check{E})$$

of $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}$. Theorem 4.4.1 implies

$$(\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}})_{\eta} \simeq \bigsqcup_{\beta \in \Pi_{\mathcal{G}}} (\mathcal{M}_{\mathcal{G}_{\beta},b,\mu}^{\text{int}})_{\eta} / \pi_0(\mathcal{G}_{\beta})^{\phi} \quad (4.5.1)$$

Since \mathcal{G}_{β}^o is parahoric, it follows that

$$(\mathcal{M}_{\mathcal{G}_{\beta}^o,b,\mu}^{\text{int}})_{\eta} = \text{Sht}_{\mathcal{G}_{\beta}^o(\mathbb{Z}_p)}^{\diamond}(G, b, \mu),$$

cf. [SW20, §23.3, §25.3]. Here, $\text{Sht}_K^{\diamond}(G, b, \mu)$ denotes, for any compact open subgroup $K \subset G(\mathbb{Q}_p)$, the diamond local Shimura variety, represented by a smooth rigid analytic variety over $\text{Sp}(\check{E})$, cf. [SW20, §24.1]. We use the notation $\text{Sht}_K(G, b, \mu)$ for the representing rigid-analytic variety. The morphism

$$\text{Sht}_{\mathcal{G}_{\beta}^o(\mathbb{Z}_p)}(G, b, \mu) \longrightarrow \text{Sht}_{\mathcal{G}_{\beta}(\mathbb{Z}_p)}(G, b, \mu)$$

is a Galois étale cover for the finite abelian group

$$\mathcal{G}_{\beta}(\mathbb{Z}_p) / \mathcal{G}_{\beta}^o(\mathbb{Z}_p) = (\mathcal{G}_{\beta}(\check{\mathbb{Z}}_p) / \mathcal{G}_{\beta}^o(\check{\mathbb{Z}}_p))^{\phi} = \pi_0(\mathcal{G}_{\beta})^{\phi},$$

cf. [SW20, 23.3]. We deduce the following identification which is the desired interpretation of the decomposition in Theorem 4.4.1 in the generic fiber.

Theorem 4.5.1. *Let (G, b, μ) be a local Shimura datum and let \mathcal{G} be a quasi-parahoric for G . Then*

$$(\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}})_\eta \simeq \bigsqcup_{\tilde{\beta} \in \Pi_{\mathcal{G}}} \text{Sht}_{\mathcal{G}_{\tilde{\beta}}(\mathbb{Z}_p)}^\diamond(G, b, \mu),$$

In particular, if $\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}$ is representable by the formal scheme $\mathcal{M}_{\mathcal{G}, b, \mu}$ over $O_{\check{E}}$, then

$$\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{rig}} \simeq \bigsqcup_{\tilde{\beta} \in \Pi_{\mathcal{G}}} \text{Sht}_{\mathcal{G}_{\tilde{\beta}}(\mathbb{Z}_p)}(G, b, \mu).$$

□

4.6. Interpretation in terms of local systems. Consider a perfectoid $S \rightarrow \text{Spd}(\check{E})$, i.e. with untilt S^\sharp in char. 0. Suppose $r > 0$ is sufficiently small, so that $\mathcal{Y}_{[0, r]}(S)$ avoids the divisor corresponding to the untilt. A \mathcal{G} -shtuka over S with leg at S^\sharp , restricts to a vector bundle over $\mathcal{Y}_{[0, r]}(S)$ with an isomorphism covering the action of ϕ^{-1} on $\mathcal{Y}_{[0, r]}(S)$; as in [SW20, §22.1], we call this a ϕ^{-1} module over $\mathcal{Y}_{[0, r]}(S)$. We see that an S -point of $\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}}$ over $\text{Spd}(\check{E})$ gives an exact tensor functor

$$\text{Rep}_{\mathbb{Z}_p}(\mathcal{G}) \longrightarrow \phi^{-1}\text{-mod} / \mathcal{Y}_{[0, r]}(S)$$

On the other hand, by [SW20, Prop. 22.3.2], we have an exact equivalence of tensor categories,

$$\phi^{-1}\text{-mod} / \mathcal{Y}_{[0, r]}(S) \xrightarrow{\sim} \mathbb{Z}_p\text{-Loc}(S).$$

Here, $\mathbb{Z}_p\text{-Loc}(S)$ stands for the category of pro-étale local systems for \mathbb{Z}_p over S , i.e. \mathbb{Z}_p -torsors for the pro-étale topology on S .

Composing these two functors, we obtain an exact tensor functor

$$\text{Rep}_{\mathbb{Z}_p}(\mathcal{G}) \longrightarrow \mathbb{Z}_p\text{-Loc}(S). \quad (4.6.1)$$

The composition of this functor with $\mathbb{Z}_p\text{-Loc}(S) \rightarrow \mathbb{Q}_p\text{-Loc}(S)$ is pro-étale locally on S isomorphic to the forgetful functor (so, pro-étale locally, it gives the trivial torsor), cf. [SW20, Thm. 22.5.2].

This leads to an alternative description of the generic fiber analogous to [SW20, Prop. 23.3.1]. In the statement below, $X_{FF, S} := \mathcal{Y}_{(0, \infty)}(S) / \phi^{\mathbb{Z}}$ denotes the Fargues-Fontaine curve over S , [FS21]. We will denote by $\mathcal{E}^b \times_{\text{Spd}(k)} S$, or simply \mathcal{E}^b , the descent of $(G_{\mathcal{Y}_{(r, \infty)}(S)}, \phi_b)$ to $X_{FF, S}$.

Proposition 4.6.1. *The S -points of $(\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}})_\eta$ over $\text{Spd}(\check{E})$ are in bijection with isomorphism classes of 5-tuples $(S^\sharp, \mathcal{E}, \alpha, \mathbb{P}, \iota)$ where*

- S^\sharp is an untilt of S over \check{E} ;
- \mathcal{E} is a G -torsor over $X_{FF, S}$, trivial at every geometric point of S ;
- α is a G -torsor isomorphism

$$\mathcal{E}|_{X_{FF, S} \setminus S^\sharp} \xrightarrow{\sim} \mathcal{E}^b|_{X_{FF, S} \setminus S^\sharp},$$

which is meromorphic along S^\sharp and bounded by μ ;

- \mathbb{P} is a fully faithful exact tensor functor

$$\text{Rep}_{\mathbb{Z}_p}(\mathcal{G}) \longrightarrow \mathbb{Z}_p\text{-Loc}(S);$$

- ι is an isomorphism of the base change of \mathbb{P} to \mathbb{Q}_p with the functor given by the $\underline{G(\mathbb{Q}_p)}$ -torsor associated to \mathcal{E} by [SW20, Thm. 22.5.2].

Proof. This follows by the argument in the proof of [SW20, Prop. 23.3.1]. Observe that, since we do not assume that \mathcal{G} has connected fibers, \mathcal{G} -torsors over $\mathrm{Spec}(\mathbb{Z}_p)$ are not necessarily trivial. So instead of $\underline{\mathcal{G}(\mathbb{Z}_p)}$ -torsors as in loc. cit., we just obtain functors \mathbb{P} as above, compare also to [SW20, proof of Prop. 22.6.1]. \square

We can use Proposition 4.6.1 to reinterpret the isomorphism in Theorem 4.5.1. For S perfectoid over k , we can consider the fiber functor at $s : \mathrm{Spa}(C, O_C) \rightarrow S$,

$$F_s : \mathbb{Z}_p\text{-Loc}(S) \longrightarrow \mathbb{Z}_p\text{-mod.}$$

Given an S -point of $\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}}$ over $\mathrm{Spd}(\check{E})$ as above, the composition of (4.6.1) and F_s defines an exact tensor functor

$$\mathrm{Rep}_{\mathbb{Z}_p}(\mathcal{G}) \longrightarrow \mathbb{Z}_p\text{-mod.} \quad (4.6.2)$$

It defines a \mathcal{G} -torsor \mathcal{T}_s and we can consider the contracted product

$$\mathcal{G}_s := \mathrm{Aut}(\mathcal{T}_s) = \mathcal{G} \wedge^{\mathcal{G}} \mathcal{T}_s,$$

where \mathcal{G} acts on itself by conjugation. Since \mathcal{G} is smooth, we have

$$\mathcal{G}_s \otimes_{\mathbb{Z}_p} \check{\mathbb{Z}}_p \simeq \mathcal{G} \otimes_{\mathbb{Z}_p} \check{\mathbb{Z}}_p.$$

The class $[\mathcal{T}_s]$ of the torsor \mathcal{T}_s lies in $\ker(\mathrm{H}^1(\mathbb{Z}_p, \mathcal{G}) \rightarrow \mathrm{H}^1(\mathbb{Q}_p, G)) = \Pi_{\mathcal{G}}$, comp. Lemma 3.1.1.

A 4-tuple $(S^{\sharp}, \mathcal{E}, \alpha, \mathbb{P}) \in (\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}})_{\eta}(S)$ defines now the function

$$S \longrightarrow \Pi_{\mathcal{G}}, \quad s \longmapsto [\mathcal{T}_s]. \quad (4.6.3)$$

Assuming a theory of pro-étale systems in this situation that resembles the classical theory of (étale) \mathbb{Z}_p -local systems, it would follow that the isomorphism class of the torsor \mathcal{T}_s is constant on connected components of S . More precisely, if we choose another point s' of S , then $F_s \simeq F_{s'}$ (by a “path connecting s to s' ”) and this should give $\mathcal{T}_s \simeq \mathcal{T}_{s'}$ and $\mathcal{G}_s(\mathbb{Z}_p) = \mathcal{G}_{s'}(\mathbb{Z}_p)$ as subgroups of $G(\mathbb{Q}_p)$. This would explain how to associate to a point $s \in (\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}})_{\eta}(C, O_C)$ the index $\bar{\beta}(s) \in \Pi_{\mathcal{G}}$ on the RHS of (4.5.1). It would also explain how to associate to s the point of $\mathrm{Sht}_{\mathcal{G}_{\beta}(Z_p)}^{\diamond}(G, b, \mu)$ on the RHS of (4.5.1) (note that $\mathcal{G}_s = \mathcal{G}_{\beta(s)}$).

4.7. Comparison with Wintenberger’s theorem. Let us discuss “classical” points $x : \mathrm{Spd}(F) \rightarrow (\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}})_{\eta}$, where F/\check{E} is a finite field extension. Consider the crystalline period map

$$\pi_{\mathrm{GM}} : (\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}})_{\eta} \longrightarrow (X_{\mu} \times_{\mathrm{Sp}(E)} \mathrm{Sp}(\check{E}))^{\diamond}.$$

Under the identification of Theorem 4.5.1, it is the disjoint sum of the period maps, one for each $\bar{\beta} \in \Pi_{\mathcal{G}}$,

$$\pi_{\mathrm{GM}, \bar{\beta}} : \mathrm{Sht}_{\mathcal{G}_{\beta}(Z_p)}(G, b, \mu) \longrightarrow X_{\mu} \times_{\mathrm{Sp}(E)} \mathrm{Sp}(\check{E}). \quad (4.7.1)$$

On the image of $\pi_{\mathrm{GM}, \bar{\beta}}$ (the admissible locus), there is the corresponding local $G(\mathbb{Q}_p)$ -system. Let $x \in \mathrm{Sht}_{\mathcal{G}_{\beta}(Z_p)}(G, b, \mu)(F)$. Specializing this local system at $\pi_{\mathrm{GM}, \bar{\beta}}(x)$ defines a Galois representation

$$\rho_x : \mathrm{Gal}(\bar{F}/F) \longrightarrow K_{\beta} = \mathcal{G}_{\beta}(\mathbb{Z}_p) \subset G(\mathbb{Q}_p). \quad (4.7.2)$$

Recall the theorem of Wintenberger (comp. [KP18, Prop. 3.3.4]), according to which the image of the Galois representation ρ_x lies in the neutral component $\mathcal{G}_\beta^o(\mathbb{Z}_p)$, i.e.,

$$\rho_x : \text{Gal}(\bar{F}/F) \longrightarrow \mathcal{G}_\beta^o(\mathbb{Z}_p) \subset G(\mathbb{Q}_p). \quad (4.7.3)$$

This may be surprising at first glance, given the construction of (4.7.2). However, consider the Galois étale cover for the group $\pi_0(\mathcal{G})^\phi$,

$$\pi : \text{Sht}_{\mathcal{G}_\beta^o(\mathbb{Z}_p)}(G, b, \mu) \longrightarrow \text{Sht}_{\mathcal{G}_\beta(\mathbb{Z}_p)}(G, b, \mu).$$

By our earlier discussion on connected components, this cover is totally split. This fits with the above result of Wintenberger which directly implies that, for every F -valued point of $\text{Sht}_{\mathcal{G}_\beta(\mathbb{Z}_p)}(G, b, \mu)$, the inverse image $\pi^{-1}(x)$ is split over F .

5. AD-ISOMORPHISMS AND INTEGRAL MODULI SPACES OF LOCAL SHTUKAS

5.1. Ad-isomorphisms. Recall from [Ko97] that a homomorphism $f : G \rightarrow G'$ is an *ad-isomorphism* if the center Z_G of G is mapped to the center $Z_{G'}$ of G' and the induced homomorphism $G_{\text{ad}} \rightarrow G'_{\text{ad}}$ between the adjoint groups is an isomorphism. Equivalently, $\ker f$ is a central subgroup and $\text{Im } f$ is a normal subgroup of G' with torus cokernel.

Lemma 5.1.1. *An ad-isomorphism is the composition of ad-isomorphisms of one of the following types.*

- (i) $f : G \rightarrow G'$ is the inclusion of a normal subgroup with torus cokernel.
- (ii) $f : G \rightarrow G'$ is a surjection with kernel a central torus which is a product of induced tori (a “quasi-trivial” torus).

Proof. Let $f : G \rightarrow G'$ be an ad-isomorphism, and let $Z = \ker f$. Find an embedding $Z \hookrightarrow T$ where T is a quasi-trivial torus. Consider the push out of

$$1 \longrightarrow Z \longrightarrow G \longrightarrow G/Z \longrightarrow 1$$

by $Z \hookrightarrow T$. This gives

$$1 \longrightarrow T \longrightarrow G'' \longrightarrow G/Z \longrightarrow 1$$

and we can view $G \rightarrow G'$ as a composition of ad-isomorphisms

$$G \hookrightarrow G'' \longrightarrow G/Z \hookrightarrow G'$$

where the first map is of type (i), the second map of type (ii) and the third map of type (i). \square

In this section, we consider an ad-isomorphism $f : G \rightarrow G'$ and an extension of f to a morphism $f : \mathcal{G} \rightarrow \mathcal{G}'$ between quasi-parahoric group schemes. It is assumed that the corresponding parahoric group schemes (the neutral connected components) \mathcal{G}^o and \mathcal{G}'^o correspond to the same point in the common building of G_{ad} and G'_{ad} . Let (\mathcal{G}, b, μ) and (\mathcal{G}', b', μ') be two integral local shtuka data such that $f(b) = b'$ and $\{\mu'\} = f(\{\mu\})$. We then say that f is an *ad-isomorphism of integral local shtuka data*. We obtain a morphism

$$f : \mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}} \longrightarrow \mathcal{M}_{\mathcal{G}', b', \mu'}^{\text{int}} \times_{\text{Spd}(O')} \text{Spd}(O) \quad (5.1.1)$$

where $O = O_{\check{E}}$ and $O' = O_{\check{E}'}$. Consider the commutative diagram

$$\begin{array}{ccc} |\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}| & \longrightarrow & |\mathcal{M}_{\mathcal{G}',b',\mu'}^{\text{int}} \times_{\text{Spd}(O')} \text{Spd}(O)| \\ \text{sp} \downarrow & & \downarrow \text{sp} \\ |X_{\mathcal{G}}(b, \mu^{-1})| & \longrightarrow & |X_{\mathcal{G}'}(b', \mu'^{-1})|. \end{array} \quad (5.1.2)$$

Composing with the Kottwitz homomorphisms (4.3.3) gives a commutative diagram

$$\begin{array}{ccc} \mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}} & \longrightarrow & \mathcal{M}_{\mathcal{G}',b',\mu'}^{\text{int}} \times_{\text{Spd}(O)} \text{Spd}(O') \\ \kappa \circ \text{sp} \downarrow & & \downarrow \kappa \circ \text{sp} \\ \underline{C}_{\mathcal{G}} & \longrightarrow & \underline{C}_{\mathcal{G}'}. \end{array} \quad (5.1.3)$$

Recall the components $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau}$ for $\tau \in C_{\mathcal{G}}$, resp. $\mathcal{M}_{\mathcal{G}',b',\mu'}^{\text{int},\tau'}$ for $\tau' \in C_{\mathcal{G}'}$, cf. (4.4.2). The aim of this section is to prove the following theorem.

Theorem 5.1.2. *Let $f: (\mathcal{G}, b, \mu) \rightarrow (\mathcal{G}', b', \mu')$ be an ad-isomorphism of integral local shtuka data. The morphism (5.1.1) induces an isomorphism,*

$$f: \mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau} \xrightarrow{\sim} \mathcal{M}_{\mathcal{G}',b',\mu'}^{\text{int},\tau'} \times_{\text{Spd}(O')} \text{Spd}(O),$$

where $\tau' = f(\tau) \in C_{\mathcal{G}'}$.

The proof proceeds in three steps. In subsection 5.3 we show that f induces an isomorphism of formal completions; in subsection 5.4 we analyze the map on ADLV induced by f ; in subsection 5.5 we complete the proof by showing that the map in question is qcqs.

We note first that for ad-isomorphisms of type (i) and for the claim of qcqs, there is no need to consider components separately.

Proposition 5.1.3. *Assume that f is of type (i). Then the morphism*

$$f: \mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}} \rightarrow \mathcal{M}_{\mathcal{G}',b',\mu'}^{\text{int}} \times_{\text{Spd}(O')} \text{Spd}(O)$$

is qcqs. If in addition $\mathcal{G}(\check{\mathbb{Z}}_p) = G(\check{\mathbb{Q}}_p) \cap f^{-1}(\mathcal{G}'(\check{\mathbb{Z}}_p))$, it is a closed immersion of v -sheaves.

Proof. We first show that the morphism is qcqs. Quasi-separateness follows from Lemma 5.5.3 below and the fact that $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}} \rightarrow \text{Spd}(O_{\check{E}})$ is qs ([Gl20, Prop. 2.3.7]). It remains to show quasi-compactness and the additional statement about closed immersions. Under our assumption of type (i), we have a short exact sequence

$$1 \longrightarrow G \longrightarrow G' \longrightarrow T \longrightarrow 1$$

with T a torus. This gives an exact sequence

$$1 \longrightarrow \pi_1(G) \longrightarrow \pi_1(G') \longrightarrow X_*(T) \longrightarrow 0.$$

By taking I -coinvariants we obtain an exact sequence

$$\Omega_G = \pi_1(G)_I \longrightarrow \Omega_{G'} = \pi_1(G')_I \longrightarrow X_*(T)_I \longrightarrow 0$$

where the kernel of the first map is a finite abelian group. Hence, the inverse image Y of $\pi_0(\mathcal{G}') \subset \Omega_{G'}$ in Ω_G is a finite group which contains the subgroup $\pi_0(\mathcal{G})$

$$\pi_0(\mathcal{G}) \subset Y \subset \Omega_G.$$

Using this and Bruhat-Tits theory, cf. the construction of [BTII, Prop. (4.6.18)], we obtain a quasi-parahoric \mathcal{G}_1 for G with neutral component $\mathcal{G}_1^o = \mathcal{G}^o$ and $\pi_0(\mathcal{G}_1) = Y$. Then, $\mathcal{G}_1(\check{\mathbb{Z}}_p) = G(\check{\mathbb{Q}}_p) \cap f^{-1}(\mathcal{G}'(\check{\mathbb{Z}}_p))$. We have

$$\mathcal{G}_1^o = \mathcal{G}^o \subset \mathcal{G} \subset \mathcal{G}_1$$

and $\mathcal{G} \rightarrow \mathcal{G}'$ factors as $\mathcal{G} \subset \mathcal{G}_1 \rightarrow \mathcal{G}'$. By applying Proposition 4.2.1 to \mathcal{G} and \mathcal{G}_1 , we see that it is enough to show quasi-compactness for the morphism induced by $\mathcal{G}_1 \rightarrow \mathcal{G}'$. Hence we may assume from the start that $\mathcal{G}(\check{\mathbb{Z}}_p) = G(\check{\mathbb{Q}}_p) \cap f^{-1}(\mathcal{G}'(\check{\mathbb{Z}}_p))$. Then, as in [PR21, Lem. 3.6.1], $f : \mathcal{G} \rightarrow \mathcal{G}'$ is a dilated immersion of smooth group schemes, i.e. f identifies \mathcal{G} with the Neron smoothening of the Zariski closure $\bar{\mathcal{G}}$ of $f(G)$ in \mathcal{G}' . Now the proof of [PR21, Prop. 3.6.2] applies to show that under this assumption the morphism is a closed immersion. \square

5.2. Ad-isomorphisms and generic fibers. If (G, b, μ) is a rational local shtuka datum and \mathcal{G} is a smooth group scheme model over \mathbb{Z}_p with connected special fiber, we set, for $K = \mathcal{G}(\mathbb{Z}_p)$,

$$\mathrm{Sht}_K(G, b, \mu) = (\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}})_{\eta}.$$

This is a locally spatial diamond over $\mathrm{Spd}(\check{E})$, cf. [SW20, §23.3]. Note that here, in contrast to section 4.5, $\mathrm{Sht}_K(G, b, \mu)$ denotes a diamond which is not representable by a rigid-analytic space, unless μ is minuscule. We also set

$$\mathrm{Sht}_{\infty}(G, b, \mu) = \varprojlim_K \mathrm{Sht}_K(G, b, \mu),$$

as in [SW20, §23.3].

Proposition 5.2.1. *Let $f : (G, b, \mu) \rightarrow (G', b', \mu')$ be an ad-isomorphism of rational local shtuka data. The natural morphism*

$$\mathrm{Sht}_{\infty}(G, b, \mu) \longrightarrow \mathrm{Sht}_{\infty}(G', b', \mu') \times_{\mathrm{Spd}(\check{E}')} \mathrm{Spd}(\check{E})$$

induces an isomorphism of v -sheaves over $\mathrm{Spd}(\check{E})$,

$$\mathrm{Sht}_{\infty}(G, b, \mu) \xrightarrow{\frac{G(\mathbb{Q}_p)}{\times}} \underline{G'(\mathbb{Q}_p)} \xrightarrow{\sim} \mathrm{Sht}_{\infty}(G', b', \mu') \times_{\mathrm{Spd}(\check{E}')} \mathrm{Spd}(\check{E}).$$

Proof. In the case of minuscule μ this is given by [PR21, Prop. 3.1.2]. The argument extends provided we explain that the corresponding admissible loci in the “Schubert varieties” $\mathrm{Gr}_{G, \mathrm{Spd}(\check{E}), \leq \mu}$, resp. $\mathrm{Gr}_{G', \mathrm{Spd}(\check{E}'), \leq \mu'} \times_{\mathrm{Spd}(\check{E}')} \mathrm{Spd}(\check{E})$ agree. Recall that the latter objects are spatial diamonds [SW20, Thm. 19.2.4] which are proper over $\mathrm{Spd}(\check{E})$, cf. [SW20, Prop. 19.2.3]. The admissible loci are open subobjects, cf. [SW20, §23.3].

Since $G \rightarrow G'$ is an ad-isomorphism, the morphism

$$\mathrm{Gr}_{G, \mathrm{Spd}(\check{E})} \longrightarrow \mathrm{Gr}_{G', \mathrm{Spd}(\check{E}')} \times_{\mathrm{Spd}(\check{E}')} \mathrm{Spd}(\check{E})$$

induces an isomorphism between corresponding connected components. In particular, the morphism

$$\mathrm{Gr}_{G, \mathrm{Spd}(\check{E}), \leq \mu} \xrightarrow{\sim} \mathrm{Gr}_{G', \mathrm{Spd}(\check{E}'), \leq \mu'} \times_{\mathrm{Spd}(\check{E}')} \mathrm{Spd}(\check{E})$$

is an isomorphism. Therefore, by [Sch17, Prop. 11.15, or Prop. 12.9], it remains to show that the induced map on admissible loci is bijective on points with values in $\mathrm{Spa}(C, C^+)$, for any algebraically closed non-archimedean field C and open bounded valuation ring C^+ . Consider

the following commutative diagram of (pro-systems of) locally spatial diamonds over \check{E} , in which the vertical arrows are the period maps ([SW20, §23.3]),

$$\begin{array}{ccc} \mathrm{Sht}_\infty(G, b, \mu) & \longrightarrow & \mathrm{Sht}_\infty(G', b', \mu') \times_{\mathrm{Spa}(\check{E}')} \mathrm{Spa}(\check{E}) \\ \downarrow & & \downarrow \\ \mathrm{Gr}_{G, \mathrm{Spd}(\check{E}), \leq \mu} & \xrightarrow{\sim} & \mathrm{Gr}_{G', \mathrm{Spd}(\check{E}'), \leq \mu'} \times_{\mathrm{Spa}(\check{E}')} \mathrm{Spa}(\check{E}). \end{array}$$

The images of the vertical maps are the *admissible sets*. We claim that under the lower horizontal map the admissible sets correspond to each other. We have $\mathrm{Sht}_K(G, b, \mu)(C, C^+) = \mathrm{Sht}_K(G, b, \mu)(C, O_C)$ and $\mathrm{Sht}_{K'}(G', b', \mu')(C, C^+) = \mathrm{Sht}_{K'}(G', b', \mu')(C, O_C)$, by their definitions in terms of shtuka. Hence, it is enough to check that the $\mathrm{Spa}(C, O_C)$ -points of the admissible sets coincide. Now a point x of $\mathrm{Gr}_{G, \mathrm{Spd}(\check{E}), \leq \mu}$ with values in $\mathrm{Spa}(C, O_C)$ lies in the admissible locus if and only if the corresponding modification \mathcal{E}_x^b at ∞ of the G -bundle \mathcal{E}^b over the FF curve is trivial, cf. [SW20, Thm. 22.6.2]. This in turn is equivalent to \mathcal{E}_x^b being a semi-stable G -bundle on the FF curve (use $[b] \in B(G, \mu^{-1})$, cf. [PR21, proof of Prop. 3.1.1]). The image of x lies in the admissible set in $\mathrm{Gr}_{G', \mathrm{Spd}(\check{E}'), \leq \mu'}$ if and only if the G' -bundle $\mathcal{E}_{f_*(x)}^{b'} = f_*(\mathcal{E}_x^b)$ is a semi-stable G' -bundle on the FF curve. But since f induces a bijection between parabolic subgroups of G , resp. G' , the semi-stability of $f_*(\mathcal{E}_x^b)$ is equivalent to the semi-stability of \mathcal{E}_x^b (apply this bijection to the Harder-Narasimhan parabolics and the fact that semi-stability is equivalent to the fact that the HN-parabolic is the whole group). Our claim follows. Now consider the following diagram, where in the lower line appears the isomorphism between admissible sets,

$$\begin{array}{ccc} \mathrm{Sht}_\infty(G, b, \mu) \xrightarrow{G(\mathbb{Q}_p) \times} \underline{G'}(\mathbb{Q}_p) & \longrightarrow & \mathrm{Sht}_\infty(G', b', \mu') \times_{\mathrm{Spd}(\check{E}')} \mathrm{Spd}(\check{E}) \\ \downarrow & & \downarrow \\ (\mathrm{Gr}_{G, \mathrm{Spd}(\check{E}), \leq \mu})^{\mathrm{adm}} & \xrightarrow{\sim} & (\mathrm{Gr}_{G', \mathrm{Spd}(\check{E}'), \leq \mu'})^{\mathrm{adm}} \times_{\mathrm{Spd}(\check{E}')} \mathrm{Spd}(\check{E}). \end{array}$$

Now the fibers of the left vertical arrow are identified with $G(\mathbb{Q}_p) \times^{G(\mathbb{Q}_p)} G'(\mathbb{Q}_p) = G'(\mathbb{Q}_p)$, and hence map bijectively to the fibers of the right vertical arrow. \square

Remark 5.2.2. In [PR21, proof of Prop. 3.1.1], there is a different argument for the proof of the isomorphism between the admissible loci of the lower horizontal map in the last diagram.

5.3. Ad-isomorphisms and formal completions.

Proposition 5.3.1. *Let $f: (\mathcal{G}, b, \mu) \rightarrow (\mathcal{G}', b', \mu')$ be an ad-isomorphism of integral local shtuka data. For each $x \in \mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}}(\mathrm{Spd}(k))$, f induces an isomorphism*

$$\widehat{f}: \widehat{\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}}}_{/x} \xrightarrow{\sim} \widehat{\mathcal{M}_{\mathcal{G}', b', \mu'}^{\mathrm{int}}}_{/x'} \times_{\mathrm{Spd}(O')} \mathrm{Spd}(O),$$

where $x' = f(x)$.

Proof. As in [PR21, proof of Prop. 3.4.1, p. 56] we have

$$\widehat{\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}}}_{/x} \simeq \widehat{\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}}}_{/x_0}$$

and it is enough to show the isomorphism for the base point $x = x_0$. By Lemma 5.1.1, it is enough to show this when f is of type (i) or of type (ii).

Assume first that f is of type (ii). Then we have an exact sequence

$$1 \longrightarrow Z \longrightarrow G \longrightarrow G' \longrightarrow 1,$$

with $Z = T$ a quasi-trivial central torus. In this case, we have an exact sequence, where \mathcal{Z} is the flat closure of Z ,

$$1 \longrightarrow \mathcal{Z} \longrightarrow \mathcal{G} \longrightarrow \mathcal{G}'.$$

Note that since $Z = T$ is quasi-trivial, i.e. $Z \simeq \prod_i \text{Res}_{K_i/\mathbb{Q}_p} \mathbb{G}_m$, we have that $\Omega_Z = \pi_1(Z)_I$ is torsion free and Z is an R -smooth torus in the sense of [KZ21, Prop. 2.4.3]. By [KZ21, Prop. 2.4.12], cf. [PR08, Lem. 6.7], we conclude that \mathcal{Z} is smooth and connected and, in fact, is the connected Neron model of the torus T and that we have a short fppf exact sequence of smooth connected group schemes between the parahoric neutral components,

$$1 \longrightarrow \mathcal{Z} \longrightarrow \mathcal{G}^o \longrightarrow \mathcal{G}'^o \longrightarrow 1.$$

Note that in this situation $\mathcal{Z} \simeq \prod_i \text{Res}_{\mathcal{O}_{K_i}/\mathbb{Z}_p} \mathbb{G}_m$. The image of $\pi_0(\mathcal{G}) \subset \Omega_G$ under $\Omega_G \rightarrow \Omega_{G'}$ is a finite subgroup Y' of $\Omega_{G'}$. We can find a short fppf exact sequence of (smooth) group schemes

$$1 \longrightarrow \mathcal{Z} \longrightarrow \mathcal{G} \longrightarrow \mathcal{G}'_1 \longrightarrow 1,$$

where $\mathcal{G}'_1 \subset \mathcal{G}'$ is a quasi-parahoric group scheme with the same neutral component \mathcal{G}'^o and $\pi_0(\mathcal{G}'_1) = Y'$. By Proposition 4.2.1, there is an isomorphism $\widehat{\mathcal{M}}_{\mathcal{G}'_1, b'_x, \mu' / x'_0}^{\text{int}} \simeq \widehat{\mathcal{M}}_{\mathcal{G}', b'_x, \mu' / x'_0}^{\text{int}}$. Hence we may replace \mathcal{G}' by \mathcal{G}'_1 , i.e., we may assume from the beginning that $\mathcal{G} \rightarrow \mathcal{G}'$ is surjective.

Note that

$$\widehat{\mathbb{W}}^+ \mathcal{G}' = \widehat{\mathbb{W}}^+ \mathcal{G}'^o.$$

We obtain a short exact sequence of group v -sheaves

$$1 \longrightarrow \widehat{\mathbb{W}}^+ \mathcal{Z} \longrightarrow \widehat{\mathbb{W}}^+ \mathcal{G} \longrightarrow \widehat{\mathbb{W}}^+ \mathcal{G}' \longrightarrow 1. \quad (5.3.1)$$

The local models for (\mathcal{G}, μ) and (\mathcal{G}', μ') “agree”, i.e.

$$f : \mathbb{M}_{\mathcal{G}, \mu}^v \xrightarrow{\sim} \mathbb{M}_{\mathcal{G}', \mu'}^v \times_{\text{Spd } \mathcal{O}'} \text{Spd } \mathcal{O},$$

([SW20, 21.5.1]), so we have

$$\widehat{f}_0 : \widehat{\mathbb{M}}_{\mathcal{G}, \mu / y_0}^v \xrightarrow{\sim} \widehat{\mathbb{M}}_{\mathcal{G}', \mu' / y_0}^v \times_{\text{Spd } \mathcal{O}'} \text{Spd } \mathcal{O}.$$

We now use Theorem 3.6.2 for \mathcal{G} and \mathcal{G}' : The morphism

$$\widehat{L\mathcal{G}}_{b, \mu} \longrightarrow \widehat{\mathbb{M}}_{\mathcal{G}, \mu / y_0}^v \xrightarrow{\sim} \widehat{\mathbb{M}}_{\mathcal{G}', \mu' / y_0}^v \times_{\text{Spd } \mathcal{O}'} \text{Spd } \mathcal{O}$$

is a $\widehat{\mathbb{W}}^+ \mathcal{G} \times \text{Spd } (\mathcal{O})$ -torsor and

$$\widehat{L\mathcal{G}}'_{b', \mu'} \times_{\text{Spd } (\mathcal{O}')} \text{Spd } (\mathcal{O}) \longrightarrow \widehat{\mathbb{M}}_{\mathcal{G}, \mu / y_0}^v \xrightarrow{\sim} \widehat{\mathbb{M}}_{\mathcal{G}', \mu' / y_0}^v \times_{\text{Spd } \mathcal{O}'} \text{Spd } \mathcal{O},$$

a $\widehat{\mathbb{W}}^+ \mathcal{G}' \times_{\text{Spd } (\mathcal{O}')} \text{Spd } (\mathcal{O})$ -torsor. Hence, we have an isomorphism of $\widehat{\mathbb{W}}^+ \mathcal{G}'$ -torsors,

$$\widehat{L\mathcal{G}}'_{b', \mu'} \times_{\text{Spd } (\mathcal{O}')} \text{Spd } (\mathcal{O}) \simeq \widehat{L\mathcal{G}}_{b, \mu} \times^{\widehat{\mathbb{W}}^+ \mathcal{G}} \widehat{\mathbb{W}}^+ \mathcal{G}'. \quad (5.3.2)$$

It follows that, v -locally,

$$\widehat{L\mathcal{G}}_{b,\mu} = g_0 \cdot \widehat{\mathbb{W}}^+ \mathcal{G}, \quad \widehat{L\mathcal{G}}'_{b',\mu'} = g'_0 \cdot \widehat{\mathbb{W}}^+ \mathcal{G}',$$

(after we choose a section g_0 of the first torsor). By Theorem 3.6.2 also, we can recover $\widehat{\mathcal{M}}_{\mathcal{G},b,\mu/x_0}^{\text{int}}$ as the quotient of $\widehat{L\mathcal{G}}_{b,\mu}$ by the ϕ -conjugation action of $\widehat{\mathbb{W}}^+ \mathcal{G} \times \text{Spd}(O)$. More precisely,

$$\widehat{L\mathcal{G}}_{b,\mu} \longrightarrow \widehat{\mathcal{M}}_{\mathcal{G},b,\mu/x_0}^{\text{int}}$$

is a $\widehat{\mathbb{W}}^+ \mathcal{G} \times \text{Spd}(O)$ -torsor for the ϕ -conjugation action. The corresponding statement for (\mathcal{G}', b', μ') also holds. The exact sequence (5.3.1) and the above now implies that the map \widehat{f}_0 is v -locally surjective.

To show that \widehat{f}_0 is injective, it is enough to check injectivity for ϕ -conjugacy classes as follows (for simplicity, we omit the base change to O from the notation). Suppose that $g_0 \cdot g_1$ and $g_0 \cdot g_2 \in \widehat{L\mathcal{G}}_{b,\mu}(R, R^+)$ map, after replacing (R, R^+) by a v -cover, to the same ϕ -conjugacy class (by $\widehat{\mathbb{W}}^+ \mathcal{G}'(R, R^+)$) in $\widehat{L\mathcal{G}}'_{b',\mu'}(R, R^+)$. Then

$$g_0 \cdot g_1 = \phi(g'^+)^{-1} \cdot (g_0 \cdot g_2) \cdot g'^+ \in \widehat{L\mathcal{G}}'_{b',\mu'}(R, R^+) \subset \mathcal{G}'(W(R^+)[1/\xi]), \quad (5.3.3)$$

for some $g'^+ \in \widehat{\mathbb{W}}^+ \mathcal{G}'(R, R^+)$. Using (5.3.1), after further replacing (R, R^+) by a v -cover, we can lift g'^+ to $g^+ \in \widehat{\mathbb{W}}^+ \mathcal{G}(R, R^+)$. The identity (5.3.3) gives

$$z = [g_1 \cdot (g^+)^{-1} \cdot g_2^{-1}] \cdot g_0^{-1} \cdot \phi(g^+) \cdot g_0 \quad (5.3.4)$$

in $\mathcal{G}(W(R^+)[1/\xi])$ with $z \in \mathcal{Z}(W(R^+)[1/\xi])$ (central in $\mathcal{G}(W(R^+)[1/\xi])$). In this, the term $k^+ := g_1 \cdot (g^+)^{-1} \cdot g_2^{-1}$ in the bracket belongs to $\widehat{\mathbb{W}}^+ \mathcal{G}(R, R^+)$. Write the above

$$z = k^+ \cdot (g_0^{-1} \cdot \phi(g^+) \cdot g_0).$$

We claim that this implies that $z = z^+ \in \widehat{\mathbb{W}}^+ \mathcal{Z}(R, R^+)$. Indeed, for all $m \geq 1$, we obtain

$$z^m = (k^+)^m \cdot (g_0^{-1} \cdot \phi(g^+)^m \cdot g_0).$$

Hence, there is $N \geq 1$ depending on g_0 only, such that $z^m \in \mathcal{Z}(W(R^+)[1/\xi])$ has matrix entries (after applying some faithful representation of \mathcal{G}) with denominators which are powers of ξ bounded by N , for all m . This implies that $z \in \mathcal{Z}(W(R^+))$. Indeed, write $\mathcal{Z} = \prod_i \text{Res}_{O_{K_i}/\mathbb{Z}_p} \mathbb{G}_m$. Then, if $z^m \in (O_{K_i} \otimes_{\mathbb{Z}_p} W(R^+)[1/\xi])^\times$ has bounded denominators for all $m \geq 1$, then $z \in (O_{K_i} \otimes_{\mathbb{Z}_p} W(R^+))^\times$. We can see using (5.3.4) that we also have $z \equiv 1 \pmod{[\varpi]}$, for some pseudo-uniformizer ϖ of R^+ . Hence, we conclude $z = z^+ \in \widehat{\mathbb{W}}^+ \mathcal{Z}(R, R^+)$. By applying Lemma 3.6.1 for \mathcal{Z} to z^+ , we can write $z^+ = z_1^+ \phi(z_1^+)^{-1}$. Set $h^+ = g^+ \cdot z_1^+$. Using that Z is central, we can rewrite (5.3.4) as

$$g_0 g_1 = \phi(h^+)^{-1} \cdot g_0 g_2 \cdot h^+.$$

This gives that $g_0 g_1$ and $g_0 g_2$ are ϕ -conjugate by $h^+ \in \widehat{\mathbb{W}}^+ \mathcal{G}(R, R^+)$. This shows injectivity and concludes the proof.

Assume now that f is of type (i). Using Proposition 4.2.1, we can replace $\mathcal{G}, \mathcal{G}'$ by their parahoric neutral components $\mathcal{G}^o, \mathcal{G}'^o$. For these we have $\mathcal{G}^o(\check{\mathbb{Z}}_p) \subset G(\check{\mathbb{Q}}_p) \cap f^{-1}(\mathcal{G}'^o(\check{\mathbb{Z}}_p))$ with the intersection the $\check{\mathbb{Z}}_p$ -points of a quasi-parahoric $\tilde{\mathcal{G}}$ with $\tilde{\mathcal{G}}^o = \mathcal{G}^o$. The morphism f factors

$$\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}} \longrightarrow \mathcal{M}_{\tilde{\mathcal{G}},b,\mu}^{\text{int}} \xrightarrow{\tilde{f}} \mathcal{M}_{\mathcal{G}',b',\mu'}^{\text{int}} \times_{\text{Spd}(O')} \text{Spd}(O).$$

By Proposition 5.1.3 applied to $\tilde{\mathcal{G}} \rightarrow \mathcal{G}'$, the morphism \tilde{f} is a closed immersion. Using [Gl20, Prop. 1.4.26] we see that the base change of \tilde{f} along the completion $\widehat{\mathcal{M}_{\mathcal{G}', b', \mu'}^{\text{int}}}_{/x'_0} \rightarrow \mathcal{M}_{\mathcal{G}', b', \mu'}^{\text{int}}$ is the completion of \tilde{f} at the base points

$$\hat{f}_0 : \widehat{\mathcal{M}_{\tilde{\mathcal{G}}, b, \mu}^{\text{int}}}_{/\tilde{x}_0} \longrightarrow \widehat{\mathcal{M}_{\mathcal{G}', b', \mu'}^{\text{int}}}_{/x'_0} \times_{\text{Spd}(O')} \text{Spd}(O).$$

Hence, this is also a closed immersion. By Proposition 4.2.1 applied to the quasi-parahoric $\tilde{\mathcal{G}}$ we see that the source of this morphism can be identified with $\widehat{\mathcal{M}_{\tilde{\mathcal{G}}, b, \mu}^{\text{int}}}_{/x_0}$ and the morphism with \hat{f}_0 . Hence, \hat{f}_0 is also a closed immersion.

Lemma 5.3.2. *Suppose $\tilde{x}_1, \tilde{x}_2 \in (\widehat{\mathcal{M}_{\tilde{\mathcal{G}}, b, \mu}^{\text{int}}})_{\eta}(C, O_C)$ have the same image $\pi(\tilde{x}_1) = \pi(\tilde{x}_2)$ under the period map. Then there is $g \in G(\mathbb{Q}_p)^0$ such that $\tilde{x}_2 = g \cdot \tilde{x}_1$.*

Proof. Since the period map $\pi : (\mathcal{M}_{\mathcal{G}^o, b, \mu}^{\text{int}})_{\eta} \rightarrow \text{Gr}_{G, \text{Spd}(\check{E}), \leq \mu}^{\text{adm}}$ is étale with geometric fibers $G(\mathbb{Q}_p)/K^o$, there is $g \in G(\mathbb{Q}_p)$ with $x_2 = g \cdot x_1$. We have $\text{sp}(\tilde{x}_1) = \text{sp}(\tilde{x}_2) = x$ and it remains to show that $\kappa_G(g) = 1$. We first observe that the statement is true when $G = T$ is a torus. Indeed, in this case the period map is

$$(\mathcal{M}_{\mathcal{T}^o, b, \mu}^{\text{int}})_{\eta} \longrightarrow \text{Spd}(\check{E})$$

with geometric fibers given by $\kappa_T : T(\mathbb{Q}_p)/\mathcal{T}^o(\mathbb{Z}_p) \xrightarrow{\sim} \Omega_T$ while $(\widehat{\mathcal{M}_{\mathcal{T}^o, b, \mu}^{\text{int}}})_{\eta} \simeq \text{Spd}(\check{E})$, see [SW20, 25.2]. Then, the result follows when G_{der} is simply connected by applying functoriality to $G \rightarrow G_{\text{ab}} = D$ and observing that, in this case, $\Omega_G \simeq \Omega_D$. Finally, we reduce the general case to the case when the derived group is simply-connected by employing a z -extension

$$1 \longrightarrow Z \longrightarrow \tilde{G} \longrightarrow G \longrightarrow 1$$

and lifting (\mathcal{G}^o, b, μ) to a corresponding triple $(\tilde{\mathcal{G}}^o, \tilde{b}, \tilde{\mu})$. Here, \tilde{G}_{der} is simply connected and Z is a quasi-trivial torus. By case (ii) applied to $\tilde{G} \rightarrow G$ we obtain

$$(\widehat{\mathcal{M}_{\tilde{\mathcal{G}}^o, \tilde{b}, \tilde{\mu}}^{\text{int}}})_{\eta} \xrightarrow{\sim} (\widehat{\mathcal{M}_{\mathcal{G}^o, b, \mu}^{\text{int}}})_{\eta},$$

and the admissible sets for \tilde{G} and G coincide by the proof of Proposition 5.2.1. Since we are assuming that the result holds for \tilde{G} , we deduce it for G . \square

Lemma 5.3.3. *The map f induces a surjection $G(\mathbb{Q}_p)^0/K^o \rightarrow G'(\mathbb{Q}_p)^0/K'^o$.*

Proof. For any reductive group G over \mathbb{Q}_p we have

$$G(\mathbb{Q}_p)^0 = K \cdot \varphi(G_{\text{sc}}(\mathbb{Q}_p)),$$

where K is an arbitrary parahoric subgroup of $G(\mathbb{Q}_p)$ and $\phi : G_{\text{sc}} \rightarrow G_{\text{der}} \rightarrow G$ is the natural map. Note that both factors on the RHS are contained in the LHS, and the second factor is a normal subgroup, so the inclusion of the RHS into the LHS is clear. For the other inclusion, we choose an apartment A^{\natural} of the building over \mathbb{Q}_p as in [BTII] such that K fixes a facet in A^{\natural} . Associated to A^{\natural} is the smooth connected group scheme \mathcal{Z}^o over \mathbb{Z}_p . Now Bruhat-Tits [BTII] prove the following two facts:

- $G(\mathbb{Q}_p)^0 = \mathcal{Z}^o(\mathbb{Z}_p) \cdot \varphi(G_{\text{sc}}(\mathbb{Q}_p))$, cf. last line of [BTII, (5.2.11)].
- $\mathcal{Z}^o(\mathbb{Z}_p) \subset K$, cf. first display in [BTII, (5.2.4)].

Hence, we also obtain the other inclusion. By applying the above to G' and K'^o and G , noting that $\varphi' : G_{\text{sc}} = G'_{\text{sc}} \rightarrow G'$ factors through $f : G \rightarrow G'$, we obtain

$$G'(\mathbb{Q}_p)^0 = f(G(\mathbb{Q}_p)^0) \cdot K'^o,$$

which gives the result. \square

The above two lemmas, together with Proposition 5.2.1, imply that the map

$$(\hat{f}_0)_\eta : (\widehat{\mathcal{M}_{\mathcal{G}, b, \mu/x_0}^{\text{int}}})_\eta \longrightarrow (\widehat{\mathcal{M}_{\mathcal{G}', b', \mu'/x'_0}^{\text{int}}})_\eta$$

induced by f , gives a surjection on (C, O_C) -points. Since it is also a closed immersion between partially proper v -sheaves, it then follows that it is an isomorphism.

Both the source $X := \widehat{\mathcal{M}_{\mathcal{G}, b, \mu/x_0}^{\text{int}}}$ and the target $X' := \widehat{\mathcal{M}_{\mathcal{G}', b', \mu'/x'_0}^{\text{int}}}$ of the closed immersion

$$\hat{f}_0 : X \longrightarrow X'$$

are topologically flat by [PR21, Prop. 3.4.9], as extended in [PR21, Rem. 3.4.12] also for the non-minuscule case (this extension uses the results of [AGLR22]). By the definition of topological flatness, this implies that $|X_\eta|$, resp. $|X'_\eta|$, is dense in $|X|$, resp. $|X'|$. Since $(\hat{f}_0)_\eta : X_\eta \simeq X'_\eta$ is an isomorphism, we obtain $|X| \simeq |X'|$. The result that $\hat{f}_0 : X \rightarrow X'$ is an isomorphism now follows from [SW20, Lem. 17.4.1]. \square

5.4. Ad-isomorphisms and ADLV. Let $f : (\mathcal{G}, b, \mu) \rightarrow (\mathcal{G}', b', \mu')$ be an ad-isomorphism of integral local shtuka data. We have a commutative diagram with surjective vertical arrows,

$$\begin{array}{ccc} X_{\mathcal{G}^o} & \longrightarrow & X_{\mathcal{G}'^o} \\ \kappa_G \downarrow & & \downarrow \kappa_{G'} \\ \Omega_G & \longrightarrow & \Omega_{G'} \end{array} \quad (5.4.1)$$

where the lower row are discrete schemes (which we treat like sets).

Lemma 5.4.1. *The upper arrow induces an isomorphism between corresponding connected components. The diagram is cartesian.*

Proof. An ad-isomorphism induces an isomorphism $f_{\text{sc}} : G_{\text{sc}} \xrightarrow{\sim} G'_{\text{sc}}$. This easily implies the first assertion. The second assertion follows since $\pi_0(X_{\mathcal{G}^o}) = \Omega_G$ and $\pi_0(X_{\mathcal{G}'^o}) = \Omega_{G'}$. \square

Let us consider the effect of an ad-isomorphism on Iwahori Weyl groups. An ad-isomorphism induces maps $N(\check{\mathbb{Q}}_p) \rightarrow N'(\check{\mathbb{Q}}_p)$, and $T(\check{\mathbb{Q}}_p)_1 \rightarrow T'(\check{\mathbb{Q}}_p)_1$, and $\widetilde{W} \rightarrow \widetilde{W}'$ and an identification of affine Weyl groups $W_a = W'_a$, compatible with the semi-direct product decompositions,

$$\widetilde{W} = W_a \rtimes \Omega_G \longrightarrow W_a \rtimes \Omega_{G'} = \widetilde{W}'.$$

Denoting by K^o , resp. K'^o the parahoric subgroups of $G(\mathbb{Q}_p)$, resp. $G'(\mathbb{Q}_p)$, we have subgroups $W^{K^o} \subset W_a$ and $W^{K'^o} \subset W_a$. These subgroups of \widetilde{W} , resp. \widetilde{W}' , can be identified.

The admissible sets are contained in a single W_a -coset and hence the map $\widetilde{W} \rightarrow \widetilde{W}'$ induces a bijection

$$\text{Adm}(\mu) \xrightarrow{\sim} \text{Adm}(\mu').$$

Similarly denoting by $\text{Adm}_{K^o}(\mu)$, resp. $\text{Adm}_{K'^o}(\mu')$, the image of $\text{Adm}(\mu)$ in $W^{K^o} \backslash \widetilde{W} / W^{K^o}$, resp. of $\text{Adm}(\mu')$ in $W^{K'^o} \backslash \widetilde{W}' / W^{K'^o}$, we have a bijection

$$\text{Adm}_{K^o}(\mu) \xrightarrow{\sim} \text{Adm}_{K'^o}(\mu'). \quad (5.4.2)$$

The commutative diagram (5.4.1) induces a commutative diagram, cf. (4.3.2),

$$\begin{array}{ccc} X_{\mathcal{G}^o}(b, \mu) & \longrightarrow & X_{\mathcal{G}'^o}(b', \mu') \\ \kappa_G \downarrow & & \downarrow \kappa_{G'} \\ c_{b, \mu} + \Omega_G^\phi & \longrightarrow & c_{b', \mu'} + \Omega_{G'}^\phi. \end{array} \quad (5.4.3)$$

Lemma 5.4.2. *The upper arrow induces an isomorphism between corresponding components. The diagram is cartesian, with surjective vertical arrows.*

Proof. Here recall that by a component of $X_{\mathcal{G}}(b, \mu)$, we mean an intersection of $X_{\mathcal{G}}(b, \mu)$ with a connected component of $X_{\mathcal{G}}$. The surjectivity of the vertical arrows follows from (4.3.2).

Let $\tau \in c_{b, \mu} + \Omega_G^\phi$, with image $\tau' = f(\tau) \in c_{b', \mu'} + \Omega_{G'}^\phi$. By Lemma 5.4.1, for every $g' \check{K}'^o \in X_{\mathcal{G}'^o}(b', \mu') \cap X_{\mathcal{G}'^o}^{\tau'}$, there exists a unique point $g \check{K}^o \in X_{\mathcal{G}^o}^\tau$ lying over it. We have to compare the double cosets of $h := g^{-1} b \phi(g)$ in $\check{K}^o \backslash G(\mathbb{Q}_p) / \check{K}^o = W^{K^o} \backslash \widetilde{W} / W^{K^o}$ and of $f(h) = g'^{-1} b' \phi(g')$ in $\check{K}'^o \backslash G(\mathbb{Q}_p) / \check{K}'^o = W^{K'^o} \backslash \widetilde{W}' / W^{K'^o}$. But by (5.4.2) it follows that the class of h lies in $\text{Adm}_{K^o}(\mu)$ if and only if the class of $f(h)$ lies in $\text{Adm}_{K'^o}(\mu')$. Hence $g \check{K}^o \in X_{\mathcal{G}^o}(b, \mu)^\tau$, as desired. \square

Now we pass from the parahorics \mathcal{G}^o and \mathcal{G}'^o to the quasi-parahorics \mathcal{G} and \mathcal{G}' . The homomorphism $f : G \rightarrow G'$ induces homomorphisms

$$\Omega_G \longrightarrow \Omega_{G'}, \quad \pi_0(\mathcal{G}) \longrightarrow \pi_0(\mathcal{G}'), \quad C_{\mathcal{G}} \longrightarrow C_{\mathcal{G}'}, \quad \Pi_{\mathcal{G}} \longrightarrow \Pi_{\mathcal{G}'}. \quad (5.4.4)$$

Here recall that $C_{\mathcal{G}} = \Omega_G / \pi_0(\mathcal{G})$ and $C_{\mathcal{G}'} = \Omega_{G'} / \pi_0(\mathcal{G}')$, and $\Pi_{\mathcal{G}} = \ker(\pi_0(\mathcal{G})_\phi \rightarrow \Omega_{G, \phi})$ and $\Pi_{\mathcal{G}'} = \ker(\pi_0(\mathcal{G}')_\phi \rightarrow \Omega_{G', \phi})$. We obtain from Corollary 4.3.6 a diagram with surjective vertical arrows,

$$\begin{array}{ccc} X_{\mathcal{G}}(b, \mu) & \longrightarrow & X_{\mathcal{G}'}(b', \mu') \\ \kappa_G \downarrow & & \downarrow \kappa_{G'} \\ \bar{c}_{b, \mu} + C_{\mathcal{G}}^\phi & \longrightarrow & \bar{c}_{b', \mu'} + C_{\mathcal{G}'}^\phi. \end{array} \quad (5.4.4)$$

Proposition 5.4.3. *The upper arrow in (5.4.4) induces an isomorphism between corresponding components. The diagram is cartesian, with surjective vertical arrows.*

Proof. By Theorem 4.4.1 and the proof of Corollary 4.3.6, the diagram can be rewritten as

$$\begin{array}{ccc} \bigsqcup_{\bar{\beta} \in \Pi_{\mathcal{G}}} X_{\mathcal{G}_\beta^o}(b, \mu) / \pi_0(\mathcal{G}_\beta)^\phi & \longrightarrow & \bigsqcup_{\bar{\beta}' \in \Pi_{\mathcal{G}'}} X_{\mathcal{G}'_{\beta'}^o}(b', \mu') / \pi_0(\mathcal{G}'_{\beta'})^\phi \\ \kappa_G \downarrow & & \downarrow \kappa_{G'} \\ \bigsqcup_{\bar{\beta} \in \Pi_{\mathcal{G}}} (c_{b, \mu} + \Omega_G^\phi) / \pi_0(\mathcal{G}_\beta)^\phi & \longrightarrow & \bigsqcup_{\bar{\beta}' \in \Pi_{\mathcal{G}'}} (c_{b', \mu'} + \Omega_{G'}^\phi) / \pi_0(\mathcal{G}'_{\beta'})^\phi. \end{array}$$

The maps in this diagram respect the disjoint sum decompositions. Hence the assertion follows from Lemma 5.4.2. \square

5.5. Proof of Theorem 5.1.2. The plan of the proof is as follows. We will show in Proposition 5.5.2 below that

$$f^\tau : \mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau} \longrightarrow \mathcal{M}_{\mathcal{G}',b',\mu'}^{\text{int}} \times_{\text{Spd}(O')} \text{Spd}(O)$$

is qcqs. Since $\mathcal{M}_{\mathcal{G}',b',\mu'}^{\text{int},\tau'} \rightarrow \mathcal{M}_{\mathcal{G}',b',\mu'}^{\text{int}}$ is an open and closed immersion,

$$f^{\tau,\tau'} : \mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau} \longrightarrow \mathcal{M}_{\mathcal{G}',b',\mu'}^{\text{int},\tau'} \times_{\text{Spd}(O')} \text{Spd}(O)$$

is also qcqs (cf. the proof of Lemma 5.5.3 below). Assuming this, by [SW20, Cor. 17.4.10] and the partial properness of $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau}$ and $\mathcal{M}_{\mathcal{G}',b',\mu'}^{\text{int},\tau'}$ over $\text{Spd}(O)$, resp. $\text{Spd}(O')$, it is enough to show the following statement.

Proposition 5.5.1. *Under the above assumptions, $f^{\tau,\tau'}$ induces a bijection on $\text{Spa}(C, O_C)$ -points.*

Proof. By Proposition 3.3.1, if κ is a discrete, algebraically closed field of characteristic p , we have bijections

$$\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau}(\text{Spd}(\kappa)) = X_{\mathcal{G}}^\tau(b, \mu^{-1})(\kappa), \quad \mathcal{M}_{\mathcal{G}',b',\mu'}^{\text{int},\tau'}(\text{Spd}(\kappa)) = X_{\mathcal{G}'}^{\tau'}(b', \mu'^{-1})(\kappa)$$

and $f^{\tau,\tau'}(\text{Spd}(\kappa))$ is identified with the corresponding map

$$X_{\mathcal{G}}^\tau(b, \mu^{-1})(\kappa) \rightarrow X_{\mathcal{G}'}^{\tau'}(b', \mu'^{-1})(\kappa). \quad (5.5.1)$$

But this map is a bijection by Proposition 5.4.3. Hence, $f^{\tau,\tau'}$ induces a bijection on $\text{Spd}(\kappa)$ -points. We still have to treat $\text{Spa}(C, O_C)$ -points.

Set $O(\kappa) = O \otimes_{W(k)} W(\kappa)$, and similarly for $O'(\kappa)$. Note that the reduced locus of the base change

$$(\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau})_{O(\kappa)} := \mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau} \times_{\text{Spd}(O)} \text{Spd}(O(\kappa))$$

is $X_{\mathcal{G}}^\tau(b, \mu^{-1})_\kappa := X_{\mathcal{G}}^\tau(b, \mu^{-1}) \times_{\text{Spec}(k)} \text{Spec}(\kappa)$, and similarly for $(\mathcal{M}_{\mathcal{G}',b',\mu'}^{\text{int},\tau'})_{O'(\kappa)}$.

Now note that any $\text{Spa}(C, O_C)$ -point $\tilde{x} : \text{Spa}(C, O_C) \rightarrow \mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau}$ factors through the formal completion $(\widehat{\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau}})_{O(\kappa)/x}$ of $(\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau})_{O(\kappa)}$ at $x := \text{sp}(\tilde{x})$, where κ is the residue field $k(C) = O_C/\mathfrak{m}_C$. (Observe that x now gives a closed point of the reduced locus of $(\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau})_{O(\kappa)}$ which is $X_{\mathcal{G}}^\tau(b, \mu^{-1})_\kappa$.) Similarly, the corresponding fact is true for $\text{Spa}(C, O_C)$ -points of $\mathcal{M}_{\mathcal{G}',b',\mu'}^{\text{int},\tau'}$. If x , resp. x' , is a $\text{Spd}(\kappa)$ -point of $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau}$, resp. $\mathcal{M}_{\mathcal{G}',b',\mu'}^{\text{int},\tau'}$, we have

$$(\widehat{\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau}})_{O(\kappa)/x} \simeq (\widehat{\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}})_{O(\kappa)/x}, \quad \text{resp.} \quad (\widehat{\mathcal{M}_{\mathcal{G}',b',\mu'}^{\text{int},\tau'}})_{O'(\kappa)/x'} \simeq (\widehat{\mathcal{M}_{\mathcal{G}',b',\mu'}^{\text{int}}})_{O'(\kappa)/x'}.$$

The result now follows from a simple extension of Proposition 5.3.1 to the base changes by $W(\kappa)$. Indeed, this gives

$$f^{\tau,\tau'} : (\widehat{\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau}})_{O(\kappa)/x} \xrightarrow{\sim} (\widehat{\mathcal{M}_{\mathcal{G}',b',\mu'}^{\text{int},\tau'}})_{O'(\kappa)/f(x)} \times_{\text{Spd}(O')} \text{Spd}(O).$$

This, combined with the bijection (5.5.1) above for $\kappa = k(C)$, implies the result. \square

It remains to show the following statement.

Proposition 5.5.2. *The map of v -sheaves*

$$f^\tau : \mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}, \tau} \longrightarrow \mathcal{M}_{\mathcal{G}', b', \mu'}^{\text{int}} \times_{\text{Spd}(O')} \text{Spd}(O)$$

is qcqs.

Proof. In the following, we occasionally just write $\mathcal{M}_{\mathcal{G}}^{\text{int}}$, $\mathcal{M}_{\mathcal{G}'}^{\text{int}}$, etc., for notational simplicity. First we show that $f : \mathcal{M}_{\mathcal{G}}^{\text{int}} \rightarrow \mathcal{M}_{\mathcal{G}'}^{\text{int}} \times_{\text{Spd}(O')} \text{Spd}(O)$ is quasi-separated (qs). This quickly implies that the same is true for f^τ . For this we use that $f : \mathcal{M}_{\mathcal{G}}^{\text{int}} \rightarrow \text{Spd}(O)$ is qs ([Gl20, Prop. 2.3.7]) and the following lemma.

Lemma 5.5.3. *Consider morphisms of small v -sheaves*

$$X \xrightarrow{f} Y \xrightarrow{g} Z.$$

- 1) *If $g \circ f$ is qc and g is qs, then f is qc.*
- 2) *If $g \circ f$ is qs, then so is f .*

Proof. Note that, by definition, $f : X \rightarrow Y$ is called qs when the diagonal $X \rightarrow X \times_Y X$ is qc, cf. [Sch17, §8]. Also note that compositions and base changes of qc, resp. qs, maps are also qc, resp. qs.

Part 1): Write f as the composition $X \rightarrow X \times_Z Y \rightarrow Y$, where the first map is $\text{id} \times f$ and the second the projection. The projection is qc as the base change of the qc map $X \rightarrow Z$, while the first map is a section of the projection $X \times_Z Y \rightarrow X$; this projection is qs as the base change of $Y \rightarrow Z$. It remains to observe that a section $s : S \rightarrow T$ of a qs map $T \rightarrow S$ is qc since it can be viewed as the base change of the qc diagonal $T \rightarrow T \times_S T$ by $s \times \text{id} : T = S \times_S T \rightarrow T \times_S T$.

Part 2): Since $g \circ f : X \rightarrow Z$ is qs, the map $X \rightarrow X \times_Z X$ is qc. We can write this as a composition

$$X \longrightarrow X \times_Y X \longrightarrow X \times_Z X.$$

Here $X \times_Y X \rightarrow X \times_Z X$ is an injection and hence it is qs, since then the diagonal map is an isomorphism. We now apply (a) to this composition to deduce that $X \rightarrow X \times_Y X$ is qc, hence $X \rightarrow Y$ is qs. \square

It remains to show that f^τ is qc. For this we write again $G \rightarrow G'$ as a composition of ad-isomorphisms which are either of type (i), i.e. closed embeddings (with cokernel a torus), or type (ii), i.e. fppf surjections with kernel a central quasi-trivial torus $Z = T$. Using functoriality and the fact that compositions of qcqs morphisms are again qcqs, we see that it is enough to treat these two cases separately. In the type (i) case the result follows from Proposition 5.1.3. It remains to deal with cases of type (ii). Then we have an exact

$$1 \longrightarrow Z \longrightarrow G \longrightarrow G' \longrightarrow 1$$

with $Z = T$ a quasi-trivial torus. For a morphism $\mathcal{G}'_1 \rightarrow \mathcal{G}'$ of quasi-parahoric group schemes corresponding to a fixed parahoric group scheme for G' , the morphism $\mathcal{M}_{\mathcal{G}'_1}^{\text{int}} \rightarrow \mathcal{M}_{\mathcal{G}'}^{\text{int}}$ is qc. Hence the argument in the proof of Proposition 5.3.1 (for type (ii)) shows that we may assume that the sequence

$$1 \longrightarrow \mathcal{Z} \longrightarrow \mathcal{G} \longrightarrow \mathcal{G}' \longrightarrow 1$$

is exact.

To establish quasi-compactness we use a “sequence of points” argument (compare to [SW20, proof of Thm. 21.2.1]). Let $Y = \mathrm{Spa}(R, R^+)$ be affinoid perfectoid over k and let

$$Y \longrightarrow \mathcal{M}_{\mathcal{G}', b', \mu'}^{\mathrm{int}} \times_{\mathrm{Spd}(O')} \mathrm{Spd}(O)$$

be a morphism given by an untilt of Y over O and a \mathcal{G}' -shtuka over $\mathcal{Y}_{[0, \infty)}(R, R^+)$ with a framing. Consider also the small v -sheaf

$$X = Y \times_{(\mathcal{M}_{\mathcal{G}', b', \mu'}^{\mathrm{int}} \times_{\mathrm{Spd}(O')} \mathrm{Spd}(O))} \mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}, \tau} \longrightarrow Y.$$

Take $I = |X|$ and, for each $i \in I$, choose a point of X given by $x_i : \mathrm{Spa}(C_i, C_i^+) \rightarrow \mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}, \tau}$ and $y_i : \mathrm{Spa}(C_i, C_i^+) \rightarrow Y = \mathrm{Spa}(R, R^+)$, with matching compositions to $\mathcal{M}_{\mathcal{G}', b', \mu'}^{\mathrm{int}} \times_{\mathrm{Spd}(O')} \mathrm{Spd}(O)$, which corresponds to $i \in I = |X|$. Consider the product of points

$$(D, D^+) = ((\prod_{i \in I} C_i^+)[1/(\varpi_i)], \prod_{i \in I} C_i^+).$$

Here the pseudo-uniformizers $\varpi_i \in C_i^+$ are given by a pseudouniformizer of R^+ . The collection of y_i extends to $y : \mathrm{Spa}(D, D^+) \rightarrow \mathrm{Spa}(R, R^+)$. The compositions

$$\mathrm{Spa}(C_i, C_i^+) \longrightarrow \mathrm{Spa}(D, D^+) \longrightarrow \mathrm{Spa}(R, R^+) \longrightarrow \mathrm{Spd}(O)$$

specify untits of (C_i, C_i^+) given by $\xi_i \in W(C_i^+)$. So, we have a \mathcal{G}' -shtuka \mathcal{P}' with framing over $\mathrm{Spa}(D, D^+)$ obtained from

$$\mathrm{Spa}(D, D^+) \longrightarrow \mathrm{Spa}(R, R^+) \longrightarrow \mathcal{M}_{\mathcal{G}', b', \mu'}^{\mathrm{int}} \times_{\mathrm{Spd}(O')} \mathrm{Spd}(O),$$

and a collection of \mathcal{G} -shtukas \mathcal{P}_i with framings over each (C_i, C_i^+) , and with legs at ξ_i , given by x_i . These are compatible via the pushout of torsors by $\mathcal{G} \rightarrow \mathcal{G}'$. We want to show that (x_i) extend to

$$x : \mathrm{Spa}(D, D^+) \longrightarrow \mathcal{M}_{\mathcal{G}}^{\mathrm{int}, \tau}.$$

Since the resulting map

$$\mathrm{Spa}(D, D^+) \longrightarrow X = Y \times_{(\mathcal{M}_{\mathcal{G}', b', \mu'}^{\mathrm{int}} \times_{\mathrm{Spd}(O')} \mathrm{Spd}(O))} \mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}, \tau}$$

is a v -cover, this fact will imply the quasi-compactness of X and hence of f^τ .

In the above, using the partial properness of $\mathcal{M}_{\mathcal{G}}^{\mathrm{int}}$ and $\mathcal{M}_{\mathcal{G}'}^{\mathrm{int}}$, we can replace C_i^+ by O_{C_i} . Using Proposition 3.2.1, we see that the pairs (Φ_i, i_{r_i}) of \mathcal{G} -shtuka with framings given by x_i , are described by pairs (Φ_i, g_i) , where $\Phi_i \in \mathcal{G}(W(O_{C_i})[1/\xi_i])$ and $g_i \in G(\mathcal{Y}_{[r_i, \infty)}(C_i, O_{C_i}))$. We denote by (Φ'_i, g'_i) the images of these pairs under the map induced by $\mathcal{G} \rightarrow \mathcal{G}'$. We have

$$\Phi_i = g_i^{-1} \cdot b \cdot \phi(g_i). \quad (5.5.2)$$

Set

$$\Phi = (\Phi_i) \in \prod_i \mathcal{G}(W(O_{C_i})[1/\xi_i]) = \mathcal{G}(\prod_i (W(O_{C_i})[1/\xi_i])).$$

Since the ξ_i -denominators of Φ_i are uniformly bounded in terms of the coweight μ , we see that

$$\Phi \in \mathcal{G}((\prod_i O_{C_i})[1/\xi_i]) \subset \mathcal{G}(\prod_i (W(O_{C_i})[1/\xi_i])).$$

This says

$$\Phi \in \mathcal{G}(W(D^+)[1/\xi]), \quad \text{with} \quad \xi = (\xi_i)_i.$$

The element Φ defines a \mathcal{G} -shtuka $\mathcal{P}(\Phi)$ over $\mathrm{Spa}(D, D^+)$. We claim that the corresponding \mathcal{G}' -shtuka $\mathcal{P}(\Phi')$ given by Φ' is isomorphic to the \mathcal{G}' -shtuka \mathcal{P}' , given by $\mathrm{Spa}(D, D^+) \rightarrow \mathcal{M}_{\mathcal{G}'}^{\mathrm{int}}$ above. Indeed, let \mathcal{P}' be given by $\Psi' \in \mathcal{G}'(W(D^+)[1/\xi])$ (Note that all \mathcal{G} -torsors, resp. \mathcal{G}' -torsors, over $W(D^+)$ are trivial by Prop. 3.2.2.) By construction, the images Φ'_i , Ψ'_i , or Φ' , Ψ' , under the projections $W(D^+)[1/\xi] \rightarrow W(O_{C_i})[1/\xi_i]$ satisfy

$$\Phi'_i = h'_i \cdot \Psi'_i \cdot \phi(h'_i)^{-1}$$

with $h'_i \in \mathcal{G}'(W(O_{C_i}))$. Since

$$W(D^+)[1/\xi] \hookrightarrow \prod_i (W(O_{C_i})[1/\xi_i]).$$

this gives $\Phi' = h' \cdot \Psi' \cdot \phi(h')^{-1}$, for $h' = (h'_i)_i$ in $\mathcal{G}'(W(D^+)) = \prod_i \mathcal{G}'(W(O_{C_i}))$, and the claim follows.

It remains to show that the framings $(i_{r_i})_i$ extend to a framing of $\mathcal{P}(\Phi)$ which lifts the framing i'_r of $\mathcal{P}(\Phi')$ given by our $\mathrm{Spa}(D, D^+)$ -point of $\mathcal{M}_{\mathcal{G}'}^{\mathrm{int}}$. We can understand framings as isomorphisms of corresponding G -bundles \mathcal{E} , resp. G' -bundles \mathcal{E}' over the Fargues-Fontaine curve. Denote by \mathcal{E}^b , $\mathcal{E}^{b'}$ the G -bundle, resp. G' -bundle, over $X_{\mathrm{Spd}(k)}$ given by b , resp. b' . Then we have isomorphisms of G -bundles on $X_{\mathrm{Spd}(D, D^+)}$, resp. $X_{\mathrm{Spd}(C_i, O_{C_i})}$,

$$g' : \mathcal{E}' \xrightarrow{\sim} \mathcal{E}^{b'} \times_{\mathrm{Spd}(k)} \mathrm{Spa}(D, D^+), \quad g_i : \mathcal{E}_i \xrightarrow{\sim} \mathcal{E}^b \times_{\mathrm{Spd}(k)} \mathrm{Spa}(C_i, O_{C_i}),$$

and we would like to find

$$g : \mathcal{E} \xrightarrow{\sim} \mathcal{E}^b \times_{\mathrm{Spd}(k)} \mathrm{Spa}(D, D^+)$$

which lifts g' and projects to g_i by $\mathrm{Spa}(C_i, O_{C_i}) \rightarrow \mathrm{Spa}(D, D^+)$, for all $i \in I$. (Here, we denote these framings again by g' , g_i , hopefully this does not introduce confusion.) For a perfectoid space S over k and a G -torsor \mathcal{E} over the FF curve X_S which has all geometric fibers isomorphic to \mathcal{E}^b , we consider the torsors over S under $\tilde{G}_b = \underline{\mathrm{Aut}}(\mathcal{E})$, resp. $\tilde{G}'_{b'} = \underline{\mathrm{Aut}}(\mathcal{E}')$,

$$Q_S := \underline{\mathrm{Isom}}_{\mathrm{fil}}(\mathcal{E}, \mathcal{E}^b \times_{\mathrm{Spd}(k)} S), \quad Q'_S := \underline{\mathrm{Isom}}_{\mathrm{fil}}(\mathcal{E}', \mathcal{E}^{b'} \times_{\mathrm{Spd}(k)} S).$$

Note here that there is a HN filtration on \mathcal{E} and \mathcal{E}' , see [FS21, proof of Prop. III. 5.3], and we ask the isomorphisms to respect the filtrations. This preservation is automatic over a point $S = \mathrm{Spa}(C, O_C)$, and also for \mathcal{E}' over $S = \mathrm{Spa}(D, D^+)$ since all the fibers have the same HN polygon given by b' , cf. [FS21, Thm. II.2.19]. We have a map of torsors $Q_S \rightarrow Q'_S$ covering the homomorphism $\tilde{G}_b \rightarrow \tilde{G}'_{b'}$. By [FS21, proof of Prop. III.5.3], these are \tilde{G}_b -, resp. $\tilde{G}'_{b'}$ -torsors which are trivial pro-étale locally on S .

In the description of \tilde{G}_b and $\tilde{G}'_{b'}$ given by [FS21, Prop. III.5.1], we see that $\tilde{G}_b \rightarrow \tilde{G}'_{b'}$ induces an isomorphism $\tilde{G}_b^{>0} \xrightarrow{\sim} \tilde{G}'_{b'}^{>0}$. Indeed, the central torus Z acts trivially on $\mathrm{Lie}(G)$ with the adjoint action and so for $\lambda > 0$, we have

$$(\mathrm{Lie}(G) \otimes_{\mathbb{Q}_p} \check{\mathbb{Q}}_p, \mathrm{Ad}(b)\sigma)^{-\lambda} = (\mathrm{Lie}(G') \otimes_{\mathbb{Q}_p} \check{\mathbb{Q}}_p, \mathrm{Ad}(b')\sigma)^{-\lambda}$$

with the notations as in loc. cit.. It follows by loc. cit. that the kernel of $\tilde{G}_b \rightarrow \tilde{G}'_{b'}$ is $\underline{Z}(\mathbb{Q}_p)$. This is also the kernel of $\underline{G}_b(\mathbb{Q}_p) \rightarrow \underline{G}'_{b'}(\mathbb{Q}_p)$.

Let $T = \mathrm{Spa}(D, D^+)$. For $i \in I$, the projection $(D, D^+) \rightarrow (C_i, O_{C_i})$ gives $\mathrm{Spa}(C_i, O_{C_i}) \rightarrow T$. These combine to give an injection

$$I \hookrightarrow |T|.$$

This identifies the discrete set I with a dense subspace of $|T|$ and of $\pi_0(|T|)$. As in the proof of [Gl20, Prop. 1.1.9], we see that

$$\pi_0(|T|) \simeq \beta I$$

is the Stone-Ćech compactification of the discrete set I .

By pulling back the map of torsors $f : Q_T \rightarrow Q'_T$ over T along

$$q : T \longrightarrow Q'_T$$

given by g , we obtain a $Z(\mathbb{Q}_p)$ -torsor $\mathcal{Q} = T \times_{Q'_T} Q_T \rightarrow T$. Since the perfectoid space T is strictly totally disconnected, the $Z(\mathbb{Q}_p)$ -torsor \mathcal{Q} is trivial, cf. [FS21, Lem. III.2.6] (see also the argument on top of loc. cit. p. 90.). The framings corresponding to g_i , $i \in I$, give $\mathrm{Spa}(C_i, O_{C_i})$ -points q_i of \mathcal{Q} . We can think of $(q_i)_{i \in I}$ as giving a section of \mathcal{Q} over I .

On the other hand, we have a morphism $\tilde{y} : \mathcal{Q} \rightarrow \mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}}$ which lifts the T -point y of $\mathcal{M}_{\mathcal{G}', b', \mu'}^{\mathrm{int}}$, i.e. it fits in a commutative diagram

$$\begin{array}{ccc} \mathcal{Q} & \xrightarrow{\tilde{y}} & \mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}} \\ \downarrow & & \downarrow f \\ T & \xrightarrow{y} & \mathcal{M}_{\mathcal{G}', b', \mu'}^{\mathrm{int}}. \end{array} \quad (5.5.3)$$

The morphism \tilde{y} is obtained by combining the \mathcal{G} -shtuka $\mathcal{P}(\Phi)$ with the framing provided by the universal point of \mathcal{Q} . We claim that there is an extension of the given section of \mathcal{Q} over I to a section of \mathcal{Q} over T . This would give the desired lift g of the framing g' and finish the proof.

Since \mathcal{Q} is the trivial torsor, such a section is given by a continuous function $|T| \rightarrow Z(\mathbb{Q}_p)$ extending the given function $I \rightarrow Z(\mathbb{Q}_p)$. To construct it, we use a compactness argument. Consider the composition

$$\omega : |\mathcal{Q}| \xrightarrow{|\tilde{y}|} |\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}}| \xrightarrow{\kappa_G \circ \mathrm{osp}} C_{\mathcal{G}} = \Omega_G / \pi_0(\mathcal{G}).$$

Note that

$$\omega(z \cdot q) = \kappa_G(z) + \omega(q)$$

for $z \in Z(\mathbb{Q}_p)$.

Now observe that, since the points x_i in our construction lie in $\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}, \tau}(\mathrm{Spa}(C_i, O_{C_i}))$, the corresponding points q_i of \mathcal{Q} satisfy $\omega(q_i) = \tau$, for all $i \in I$. Set $\mathfrak{T} := \omega^{-1}(\{\tau\}) \subset |\mathcal{Q}|$. This is a quasi-compact subset of $|\mathcal{Q}|$ since $|\mathcal{Q}| \simeq |T| \times Z(\mathbb{Q}_p)$, where $|T|$ is quasi-compact, and where the restriction $Z(\mathbb{Q}_p) \rightarrow C_{\mathcal{G}}$ of κ_G has compact fibers. The last fact follows because $\Omega_Z = Z(\mathbb{Q}_p)/Z(\mathbb{Z}_p)$ and $\Omega_Z \rightarrow \Omega_G \rightarrow C_{\mathcal{G}}$ has finite kernel, so each fiber is given by a finite union of cosets of $Z(\mathbb{Z}_p)$ in $Z(\mathbb{Q}_p)$.

Since $\pi_0(\mathfrak{T})$ is compact, by the universal property of the Stone-Ćech compactification we see that the composed map $I \rightarrow \mathfrak{T} \rightarrow \pi_0(\mathfrak{T})$, given by $i \mapsto q_i$, uniquely extends to a continuous map $\beta I \simeq \pi_0(|T|) \rightarrow \pi_0(\mathfrak{T}) \subset \pi_0(|T|) \times Z(\mathbb{Q}_p)$. This corresponds to the desired continuous extension $|T| \rightarrow \pi_0(|T|) \rightarrow Z(\mathbb{Q}_p)$. As above, this produces a section $T \rightarrow \mathcal{Q}$ whose composition with $\tilde{y} : \mathcal{Q} \rightarrow \mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}}$ gives $T \rightarrow \mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}}$. This factors through the open and closed $\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}, \tau} \hookrightarrow \mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}}$ and provides the desired $x : T \rightarrow \mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}, \tau}$. \square

6. THE CASE OF TRIVIAL μ_{ad}

In this section we prove Theorems 2.5.4 and 2.5.5 in the case when $\mu_{\text{ad}} = 1$ is trivial. Note that when $\mu_{\text{ad}} = 1$ is trivial, $B(G, \mu^{-1})$ consists of the unique basic element contained in it. In this section, b denotes a representative of this unique element.

6.1. The case of trivial μ . Let first \mathcal{G} be a parahoric for G . It follows from the definition that $\mathbb{M}_{\mathcal{G},1}^{\text{loc}} = \text{Spec}(\mathbb{Z}_p)$. Furthermore, we have

$$\mathcal{M}_{\mathcal{G},1,1}^{\text{int}} \simeq \underline{G(\mathbb{Q}_p)/\mathcal{G}(\mathbb{Z}_p)} \quad (6.1.1)$$

over $\text{Spd}(\check{\mathbb{Z}}_p)$, cf. [SW20, Prop. 25.2.1] (in loc. cit. only the case of a torus group is considered but the proof is valid in the general case). It follows that $\mathcal{M}_{\mathcal{G},1,1}^{\text{int}}$ is representable by the formal scheme

$$\mathcal{M}_{\mathcal{G},1,1} = \coprod_{G(\mathbb{Q}_p)/\mathcal{G}(\mathbb{Z}_p)} \text{Spf}(\check{\mathbb{Z}}_p),$$

and that the formal completion at a k -point is isomorphic to $\check{\mathbb{Z}}_p$. This proves Theorems 2.5.4 and 2.5.5 in the case when \mathcal{G} is a parahoric group scheme. The case of a general quasi-parahoric follows from Theorem 4.1.1, Proposition 4.2.1 and (2.3.1).

6.2. The general case. The case when $\mu_{\text{ad}} = 1$ is reduced to the case of the adjoint group, as follows. Choose an extension $\mathcal{G} \rightarrow \mathcal{G}'$ to quasi-parahorics of the natural morphism $G \rightarrow G' := G_{\text{ad}}$. Consider the corresponding morphism of v -sheaves

$$\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}} \longrightarrow \mathcal{M}_{\mathcal{G}',1,1}^{\text{int}}.$$

We now use the following functorialities:

- there is an isomorphism $f : \mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau} \xrightarrow{\sim} \mathcal{M}_{\mathcal{G}',1,1}^{\text{int},\tau'} \times_{\text{Spd}(\check{\mathbb{Z}}_p)} \text{Spd}(O_{\check{E}})$, for each $\tau \in C_{\mathcal{G}}$, cf. Theorem 5.1.2. Hence the representability of $\mathcal{M}_{\mathcal{G}',1,1}^{\text{int}}$ implies the representability of $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}$.
- there is an isomorphism $\hat{f} : \widehat{\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}} \xrightarrow{\sim} \widehat{\mathcal{M}_{\mathcal{G}',1,1}^{\text{int}}} \times_{\text{Spd}(\check{\mathbb{Z}}_p)} \text{Spd}(O_{\check{E}})$, for each $x \in \mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}(\text{Spd } k)$, cf. Proposition 5.3.1.
- there is an isomorphism $\mathbb{M}_{\mathcal{G},\mu}^v \xrightarrow{\sim} \mathbb{M}_{\mathcal{G}',1}^v \times_{\text{Spd}(\mathbb{Z}_p)} \text{Spd}(O_E)$, cf. [SW20, Prop. 21.5.1] (attributed in loc. cit. to J. Lourenço).

Hence the case $(\mathcal{G}', 1, 1)$ implies the case (\mathcal{G}, b, μ) .

7. THE CRUCIAL HODGE TYPE CASE

We now give the first non-banal cases when we can show the isomorphism of Theorem 2.5.5 and show representability of the formal completions. Let (G, b, μ) be a local Shimura datum. Also, throughout this section, we let $\mathcal{G} = \mathcal{G}_{\mathbf{x}}$ be a stabilizer Bruhat-Tits group scheme for a corresponding point \mathbf{x} in the extended building. We denote by \mathcal{G}^o the corresponding parahoric. We assume we are in case (A), so in particular $p > 2$.

7.1. The crucial Hodge type case. Let $\iota : (G, \mu) \hookrightarrow (\mathrm{GL}_h, \mu_d)$ be a Hodge embedding. We assume that there is a \mathbb{Z}_p -lattice $\Lambda \subset \mathbb{Q}_p^h$ such that the homomorphism ι extends to a closed immersion $\iota : \mathcal{G} \hookrightarrow \mathrm{GL}(\Lambda)$. Then we also have

$$\mathcal{G}(\check{\mathbb{Z}}_p) = G(\check{\mathbb{Q}}_p) \cap \iota^{-1}(\mathrm{GL}(\Lambda \otimes_{\mathbb{Z}_p} \check{\mathbb{Z}}_p)).$$

By [PR21, Prop. 3.6.2], this gives a closed immersion

$$\iota : \mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}} \hookrightarrow (\mathcal{M}_{\mathrm{GL}(\Lambda), \iota(b), \mu_d^{-1}}^{\mathrm{int}})_O := \mathcal{M}_{\mathrm{GL}(\Lambda), \iota(b), \mu_d^{-1}}^{\mathrm{int}} \times_{\mathrm{Spd}(W(k))} \mathrm{Spd}(O).$$

The formal completion $\widehat{\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}}}_{/x_0}$ at the base point x_0 fits in a cartesian diagram

$$\begin{array}{ccc} \widehat{\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}}}_{/x_0} & \xrightarrow{\hat{\iota}} & (\mathcal{M}_{\mathrm{GL}(\Lambda), \iota(b), \mu_d^{-1}}^{\mathrm{int}})_{/ \iota(x_0)}^{\mathrm{int}}_O \\ \downarrow & & \downarrow \\ \mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}} & \xrightarrow{\iota} & (\mathcal{M}_{\mathrm{GL}(\Lambda), \iota(b), \mu_d^{-1}}^{\mathrm{int}})_O. \end{array} \quad (7.1.1)$$

Hence, we also have a closed immersion

$$\hat{\iota} : \widehat{\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}}}_{/x_0} \hookrightarrow \mathcal{M}_{\mathrm{GL}(\Lambda), \iota(b), \mu_d^{-1}}^{\mathrm{int}} \times_{\mathrm{Spd}(W(k))} \mathrm{Spd}(O).$$

Here, by [SW20, Thm. 25.1.2] and the definition of formal completion, we have

$$\mathcal{M}_{\mathrm{GL}(\Lambda), \iota(b), \mu_d^{-1}}^{\mathrm{int}} \times_{\mathrm{Spd}(W(k))} \mathrm{Spd}(O) \simeq \mathrm{Spd}(R \otimes_{W(k)} O), \quad (7.1.2)$$

where $R \simeq W(k)[[x_1, \dots, x_m]]$, for $m = d(h-d)$. The point $\iota(x_0)$ gives a p -divisible group \mathcal{G}_0 of height h over k and R is naturally identified with the universal formal deformation ring of \mathcal{G}_0 .

Proposition 7.1.1. *There is a commutative diagram*

$$\begin{array}{ccc} (\widehat{\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}}}_{/x_0})_\eta & \xrightarrow{\hat{\iota}_\eta} & \mathrm{Spf}(R)_{\check{E}}^{\mathrm{rig}} \\ \pi_{\mathcal{G}} \downarrow & & \downarrow \pi_{\mathrm{GL}_h} \\ X_\mu^{\mathrm{an}} & \xrightarrow{\iota^{\mathrm{an}}} & \mathrm{Gr}(d, h)_{\check{E}}^{\mathrm{an}} \end{array} \quad (7.1.3)$$

of morphisms of smooth rigid analytic spaces over $\mathrm{Sp}(\check{E})$. The vertical maps are the restrictions of the étale period maps to the tubular neighborhoods. The bottom horizontal map is the analytification of the Zariski closed immersion

$$X_\mu \hookrightarrow \mathrm{Gr}(d, h)_E$$

which is obtained from $\iota : G \hookrightarrow \mathrm{GL}_h$. The top horizontal map is a closed immersion.

Here $\mathrm{Spf}(R)_{\check{E}}^{\mathrm{rig}}$ is the rigid analytic generic fiber of $\mathrm{Spf}(R)$ (in Berthelot's sense). For simplicity, we use the same symbol for the v -sheaf $(\widehat{\mathcal{M}_{\mathcal{G}, b, \mu}^{\mathrm{int}}})_\eta$, and for the smooth rigid analytic space over $\mathrm{Sp}(\check{E})$ that represents it.

Proof. Consider the commutative diagram of v -sheaves

$$\begin{array}{ccc} (\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}})_\eta & \xrightarrow{\iota} & \mathcal{M}_{\text{GL}_h,\iota(b),\mu_d^{-1}}^{\text{int}} \times_{\text{Spd}(\mathbb{Z}_p)} \text{Spd}(\check{E}) \\ \pi_{\mathcal{G}} \downarrow & & \downarrow \pi_{\text{GL}_h} \\ X_\mu^\diamond & \xrightarrow{\iota} & \text{Gr}(d,h)^\diamond \times_{\text{Spd}(\mathbb{Q}_p)} \text{Spd}(\check{E}). \end{array} \quad (7.1.4)$$

In this, the two horizontal maps are closed immersions. In particular, the bottom horizontal map is represented by a Zariski closed immersion. The two vertical period maps are given by the étale period maps (on the left side we combine the period maps on the components, as in Theorem 4.5.1). Their images are the corresponding open admissible sets and we have

$$X_\mu^{\text{adm}} = (\text{Gr}(d,h) \times_{\text{Spd}(\mathbb{Q}_p)} \text{Spd}(\check{E}))^{\text{adm}} \cap X_\mu,$$

cf. [PR21, proof of Prop. 3.1.1]. Note that $(\widehat{\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}})_\eta \hookrightarrow (\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}})_\eta$ is an open immersion by the argument of [Gl20, Prop. 1.4.29]. Hence, all the v -sheaves in (7.1.3) are representable by smooth rigid analytic spaces. By full-faithfulness [SW20, Prop. 10.2.3] the top horizontal map is also representable by a morphism of rigid analytic spaces and the result follows. \square

Theorem 7.1.2. *Let (G, b, μ) be a local Shimura datum and let \mathcal{G} be a quasi-parahoric stabilizer group scheme for G . Suppose $p > 2$ and assume that there exists a Hodge embedding $\iota : (G, \mu) \hookrightarrow (\text{GL}_h, \mu_d^{-1})$ and a \mathbb{Z}_p -lattice $\Lambda \subset \mathbb{Q}_p^h$ such that:*

- a) *The homomorphism ι extends to a closed immersion $\iota : \mathcal{G} \hookrightarrow \text{GL}(\Lambda)$.*
- b) *The Zariski closure \overline{X}_μ of $X_\mu \subset \text{Gr}(d, h)_E$ in $\text{Gr}(d, \Lambda)_{O_E}$ is normal.*
- c) *The image $\iota(G)$ contains the scalars.*

Then $\mathbb{M}_{\mathcal{G},\mu}^{\text{loc}} \simeq \overline{X}_\mu$ and for any $x \in \mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}(\text{Spd}(k))$ there is $y \in \mathbb{M}_{\mathcal{G},\mu}^{\text{loc}}(k)$ such that

$$\widehat{\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}}_{/x} \simeq (\widehat{\mathbb{M}_{\mathcal{G},\mu}^{\text{loc}}})_y^\diamond. \quad (7.1.5)$$

Proof. By section 3.5, we may assume that $x = x_0$ is the base point. For simplicity, we write

$$\widehat{\mathcal{M}_{/x_0}^{\text{int}}} = \widehat{\mathcal{M}_{\mathcal{G},b,\mu/x_0}^{\text{int}}}, \quad \mathbb{M}^{\text{loc}} = \mathbb{M}_{\mathcal{G},\mu}^{\text{loc}}.$$

We first show $\mathbb{M}^{\text{loc}} \simeq \overline{X}_\mu$. By [SW20, Thm. 21.2.1], ι induces a closed immersion of v -sheaves over $\text{Spd}(\mathbb{Z}_p)$,

$$\iota : \text{Gr}_{\mathcal{G},\text{Spd}(\mathbb{Z}_p)} \longrightarrow \text{Gr}_{\text{GL}_h,\text{Spd}(\mathbb{Z}_p)}.$$

This gives a closed immersion of v -sheaves over $\text{Spd}(O_E)$,

$$\mathbb{M}^v \longrightarrow \text{Gr}(d, h)_{O_E}^\diamond.$$

Indeed, $\mathbb{M}^v = \mathbb{M}_{\mathcal{G},\mu}^v$ is, by definition, the v -sheaf closure of X_μ^\diamond in

$$\text{Gr}(d, h)_{O_E}^\diamond \hookrightarrow \text{Gr}_{\text{GL}_h} \times_{\text{Spd}(\mathbb{Z}_p)} \text{Spd}(O_E).$$

(See [AGLR22, 2.1] for a discussion of v -sheaf closures.) Since $\mathbb{M}^v = (\mathbb{M}^{\text{loc}})^\diamond$ and \mathbb{M}^{loc} is normal, we deduce from full-faithfulness (using [SW20, Prop. 18.4.1] and formal GAGA) that the closed immersion $\mathbb{M}^v \rightarrow \text{Gr}(d, h)_{O_E}^\diamond$ is represented by a morphism of schemes,

$$\mathbb{M}^{\text{loc}} \longrightarrow \text{Gr}(d, h)_{O_E}.$$

This morphism extends $X_\mu \hookrightarrow \mathrm{Gr}(d, h)_E$ on the generic fibers. Now $(\overline{X}_\mu)^\diamond$ is topologically flat ([PR21, Lem. 3.4.2]) and so it is the v -sheaf closure of the generic fiber X_μ^\diamond in $\mathrm{Gr}(d, h)_{O_E}^\diamond$. We obtain an isomorphism

$$\mathbb{M}^v = (\mathbb{M}^{\mathrm{loc}})^\diamond \xrightarrow{\sim} (\overline{X}_\mu)^\diamond \hookrightarrow \mathrm{Gr}(d, h)_{O_E}^\diamond.$$

Since $\mathbb{M}^{\mathrm{loc}}$ is normal and we are also assuming that \overline{X}_μ is normal, the isomorphism $(\mathbb{M}^{\mathrm{loc}})^\diamond \xrightarrow{\sim} (\overline{X}_\mu)^\diamond$ is obtained from an isomorphism $\mathbb{M}^{\mathrm{loc}} \xrightarrow{\sim} \overline{X}_\mu$ of O_E -schemes, as claimed.

We now proceed to show the isomorphism (7.1.5). By [Ki10, Prop. 1.3.2], there is a finite set of tensors $\{s_a\}_a \subset \Lambda^\otimes$, such that \mathcal{G} is the scheme theoretic pointwise stabilizer of s_a ,

$$\mathcal{G} = \{g \in \mathrm{GL}(\Lambda) \mid g \cdot s_a = s_a, \forall a\}.$$

By the Tannakian formalism applied to $\mathcal{G} \rightarrow \mathrm{GL}(\Lambda)$, each \mathcal{G} -torsor over $\mathrm{Spec}(R)$ gives a finite projective R -module M of rank equal to $\mathrm{rank}_{\mathbb{Z}_p}(\Lambda)$ with tensors $\tilde{s}_a \in M^\otimes$.

Consider the universal \mathcal{G} -shtuka over $\widehat{\mathcal{M}}_{/x_0}^{\mathrm{int}}$; its push-out by $\mathcal{G} \rightarrow \mathrm{GL}(\Lambda)$ is the shtuka which is obtained from the Breuil-Kisin-Fargues module of the universal p -divisible group, as in the proof of [SW20, Theorem 25.1.2]. By specializing to the base point x_0 and using the Tannakian formalism we see that the \mathcal{G} -shtuka over $\mathrm{Spd}(k)$ that corresponds to x_0 equips the Dieudonné module $\mathbb{D} := \mathbb{D}(\mathcal{G}_0)(\check{\mathbb{Z}}_p)$ of \mathcal{G}_0 with Frobenius invariant tensors $s_{a,0} \in \mathbb{D}^\otimes$.

Write $R_{\mathcal{G}}$ for the completion of $\mathbb{M}^{\mathrm{loc}} \otimes_{O_E} O$ at y_0 . The O -algebra $R_{\mathcal{G}}$ is normal and is a quotient of $R \otimes_W O$. Note that, since $\tilde{X}_\mu \simeq \mathbb{M}^{\mathrm{loc}}$, we see that $\mathbb{M}^{\mathrm{loc}} \otimes_{O_E} O$ is identified with the reduced Zariski closure of a G -orbit $G \cdot y$ in $\mathrm{Gr}(d, h)_O$, where y is an F -point that corresponds to a filtration induced by a G -valued cocharacter μ_y conjugate to μ^{-1} . Here F is a finite extension of $\check{\mathbb{Q}}_p$.

The following statement encapsulates a construction of [KP18], as extended in [KZ21].

Proposition 7.1.3. ([KP18], [KZ21]) *There exists a p -divisible group $\mathcal{G}^{\mathrm{univ}}$ over $R \otimes_W O$ which is a versal formal deformation of \mathcal{G}_0 and which satisfies the following property:*

Let K/\check{E} be a finite field extension and $\tilde{x}^ : R \otimes_W O \rightarrow O_K$ a local O -algebra homomorphism satisfying:*

- a) The filtration on $\mathbb{D} \otimes_{\check{\mathbb{Z}}_p} K$ which corresponds to the deformation $\mathcal{G}_{\tilde{x}} := \tilde{x}^*(\mathcal{G}^{\mathrm{univ}})$ of the p -divisible group \mathcal{G}_0 to O_K given by base change of $\mathcal{G}^{\mathrm{univ}}$ via \tilde{x}^* , is induced by a G -valued cocharacter which is G -conjugate to μ^{-1} ;*
- b) The tensors $s_{\alpha,0} \in \mathbb{D}^\otimes$ correspond to tensors $s_{\alpha,\mathrm{ét}} \in T_p \mathcal{G}_{\tilde{x}}^{\vee \otimes}$ under the p -adic (étale-crystalline) comparison isomorphism.*

Then the homomorphism $\tilde{x}^ : R \otimes_W O \rightarrow O_K$ factors through the quotient $R \otimes_W O \rightarrow R_{\mathcal{G}}$.*

Proof. Let us give a very quick overview to orient the reader. The construction of $\mathcal{G}^{\mathrm{univ}}$ uses Zink's theory of displays. By the argument of [KP18, §3.2.12] we obtain a “Dieudonné display triple” $(M_{R_{\mathcal{G}}}, M_{R_{\mathcal{G}},1}, \Psi_{R_{\mathcal{G}}})$, as in loc. cit. (see Remark 7.1.5 below and the proof of [KZ21, Prop. 4.1.7] for the removal of Condition 3.2.2 in [KP18]). This triple gives, by [KP18, Lem. 3.1.5], a Dieudonné display over $R_{\mathcal{G}}$ and then one constructs also a versal display $(M_R, M_{R,1}, \Psi_R)$ over $R \otimes_W O$ which lifts $(M_{R_{\mathcal{G}}}, M_{R_{\mathcal{G}},1}, \Psi_{R_{\mathcal{G}}})$. The display $(M_R, M_{R,1}, \Psi_R)$ gives the desired versal formal deformation $\mathcal{G}^{\mathrm{univ}}$ of \mathcal{G}_0 over $\mathrm{Spf}(R \otimes_W O)$.

We can then see that the p -divisible group $\mathcal{G}^{\mathrm{univ}}$, as constructed as above, has the property stated in the proposition. For this first note that, by [KZ21, Prop. 4.2.4], a deformation $\mathcal{G}_{\tilde{x}}$

of \mathcal{G}_0 which satisfies (a) and (b) is a “ (\mathcal{G}, μ^{-1}) -adapted lifting” in the sense of [KZ21, Def. 4.1.4]. Then the claim follows from [KZ21, Prop. 4.1.7] (the same statement under additional tameness hypotheses on G also appears in [KP18], see [KP18, Prop. 3.3.13]). \square

The versal formal deformation $\mathcal{G}^{\text{univ}}$ over $R \otimes_W O$ of the p -divisible group \mathcal{G}_0 induces an identification of $\text{Spd}(R \otimes_W O)$ with $\widehat{\mathcal{M}_{\text{GL}(\Lambda), \iota(b), \mu_d / \iota(x_0)}^{\text{int}}} \times_{\text{Spd}(W)} \text{Spd}(O)$. We will now use Proposition 7.1.3 to check that the rigid analytic closed subspace

$$\text{Spf}(R_{\mathcal{G}})^{\text{rig}} \hookrightarrow \text{Spf}(R \otimes_W O)^{\text{rig}} = \text{Spf}(R)_{\check{E}}^{\text{rig}}$$

agrees with the rigid analytic closed subspace $(\widehat{\mathcal{M}_{/x_0}^{\text{int}}})_{\eta}$, under the identification induced by diagram (7.1.3).

Let K/\check{E} be a finite extension. We will compare $(\widehat{\mathcal{M}_{/x_0}^{\text{int}}})_{\eta}(K)$ and $\text{Spf}(R_{\mathcal{G}})^{\text{rig}}(K)$ as subsets of $\text{Spf}(R)_{\check{E}}^{\text{rig}}(K)$.

Proposition 7.1.4. *For all finite field extensions K/\check{E} , there is an inclusion*

$$(\widehat{\mathcal{M}_{/x_0}^{\text{int}}})_{\eta}(K) \subset \text{Spf}(R_{\mathcal{G}})^{\text{rig}}(K).$$

Proof. A K -point x of $(\widehat{\mathcal{M}_{/x_0}^{\text{int}}})_{\eta}$ gives, after composing with $(\widehat{\mathcal{M}_{/x_0}^{\text{int}}})_{\eta} \hookrightarrow \text{Spf}(R \otimes_W O)^{\text{rig}}$ a formal scheme morphism $\tilde{x} : \text{Spf}(O_K) \rightarrow \text{Spf}(R \otimes_W O)$. The point x also gives a crystalline representation

$$\rho_x : \text{Gal}(\bar{K}/K) \longrightarrow \text{GL}(T_p(\mathcal{G}_{\tilde{x}})^{\vee}) \quad (7.1.6)$$

on the linear dual of the Tate module of the p -divisible group $\mathcal{G}_{\tilde{x}}$ obtained from $\mathcal{G}^{\text{univ}}$ as above, by pulling back by \tilde{x} . Take $C = \hat{K}$ which supports a $\text{Gal}(\bar{K}/K)$ -action and consider the corresponding point $\bar{x} : \text{Spa}(C, O_C) \rightarrow \text{Spd}(K) \rightarrow (\widehat{\mathcal{M}_{/x_0}^{\text{int}}})_{\eta}$ which gives a (\mathcal{G}, μ) -shtuka $(\mathcal{P}_{\bar{x}}, \phi_{\mathcal{P}_{\bar{x}}})$ with framing. By Proposition 4.6.1 we obtain a functor

$$\mathbb{P}_{\bar{x}} : \text{Rep}_{\mathbb{Z}_p}(\mathcal{G}) \longrightarrow \mathbb{Z}_p\text{-Loc}(\text{Spa}(C, O_C)) \cong \mathbb{Z}_p\text{-mod}$$

such that $\mathbb{P}_{\bar{x}}(\Lambda) = T_p(\mathcal{G}_{\tilde{x}})^{\vee}$. For $\gamma \in \text{Gal}(\bar{K}/K)$, there is an isomorphism $\mathbb{R}_x(\gamma) : \mathbb{P}_{\bar{x}} \simeq \mathbb{P}_{\bar{x} \cdot \gamma} = \mathbb{P}_{\bar{x}}$ giving the Galois action (7.1.6), i.e., $\mathbb{R}_x(\gamma)(\Lambda) = \rho_x(\gamma)$.

The \mathcal{G} -invariant tensors $s_a \in \Lambda^{\otimes}$ give, by applying the functor \mathbb{P} , corresponding tensors $s_{a, \text{ét}} \in T_p(\mathcal{G}_{\tilde{x}})^{\vee \otimes}$; these are invariant under the action of $\text{Gal}(\bar{K}/K)$ through ρ_x . As in §4.7, we see that the Galois representation ρ_x factors

$$\rho_x : \text{Gal}(\bar{K}/K) \longrightarrow \mathcal{G}_{\beta}^{\circ}(\mathbb{Z}_p) \subset \mathcal{G}_{\beta}(\mathbb{Z}_p) \subset G(\mathbb{Q}_p),$$

where $\bar{\beta} \in \Pi_{\mathcal{G}}$ is such that x lies in the component $\text{Sh}_{\mathcal{G}_{\beta}(\mathbb{Z}_p)}(G, b, \mu)$ of $(\mathcal{M}_{\mathcal{G}, b, \mu}^{\text{int}})_{\eta}$, cf. Theorem 4.5.1. Note that

$$\mathcal{G}_{\beta} \otimes_{\mathbb{Z}_p} \check{\mathbb{Z}}_p \simeq \mathcal{G} \otimes_{\mathbb{Z}_p} \check{\mathbb{Z}}_p.$$

Choose a uniformizer π_K of K and a collection of roots π_K^{1/p^n} in $\bar{K} \subset C$. We can now see, as in [PR21, §3.5], [KP18, (3.3.3)], that the Breuil-Kisin module $\mathfrak{M}_{\tilde{x}}$ of $\mathcal{G}_{\tilde{x}}$ obtained using these choices can be refined to a $\mathcal{G}_{\beta}^{\circ}$ -Breuil-Kisin module. By definition, this is a $\mathcal{G}_{\beta}^{\circ}$ -torsor over $\text{Spec}(W(k)[[u]])$ with meromorphic Frobenius structure, see loc. cit.. Refined here is meant in the sense that the push-out by $\mathcal{G}_{\beta}^{\circ} \rightarrow \mathcal{G} \rightarrow \text{GL}(\Lambda)$ recovers the $\text{GL}(\Lambda)$ -torsor over $\text{Spec}(W(k)[[u]])$ which corresponds to $\mathfrak{M}_{\tilde{x}}$. By pushing out the torsor by $\mathcal{G}_{\beta}^{\circ} \rightarrow \mathcal{G}$, we obtain a \mathcal{G} -Breuil-Kisin module \mathcal{P}_{BK} over $\text{Spec}(W(k)[[u]])$. The tensors s_a induce tensors $\tilde{s}_a \in \mathfrak{M}_{\tilde{x}}^{\otimes}$

over $W(k)[[u]]$, comp. [KP18, (3.3.3)]. (Note that after base changing by $W(k)[[u]] \rightarrow W(O_C)$ given by $(\pi_K^{1/p^n})_n$, \mathcal{P}_{BK} gives a \mathcal{G} -Breuil-Kisin-Fargues module. By the proof of [PR21, Prop. 3.5.1], the restriction of this \mathcal{G} -Breuil-Kisin-Fargues module from $\text{Spec}(W(O_C))$ to $\mathcal{Y}_{[0,\infty)}(C, O_C)$ is isomorphic to the initial (\mathcal{G}, μ) -shtuka $(\mathcal{P}_{\tilde{x}}, \phi_{\mathcal{P}_{\tilde{x}}})$ which corresponds to the point x .) Recall that the tensors s_a also induce $s_{a,0} \in \mathbb{D}^\otimes$, where, as above, $\mathbb{D} = \mathbb{D}(\mathcal{G}_0)(\check{\mathbb{Z}}_p)$ is the Dieudonné module of the special fiber $\mathcal{G}_0 = \mathcal{G}_{\tilde{x}} \otimes_{O_K} k$. The existence of \tilde{s}_a above, together with the compatibility properties of the Breuil-Kisin functor (see for example, [KP18, Thm. 3.3.2, Prop. 3.3.8]), implies that $s_{a,\text{ét}}$ and $s_{a,0}$ correspond under the comparison isomorphism between p -adic-étale and crystalline cohomology. It also implies that the Hodge filtration on $\mathbb{D} \otimes_{\check{\mathbb{Z}}_p} K$ which corresponds to the deformation $\mathcal{G}_{\tilde{x}}$ is induced by a G -cocharacter. The above compatibility of the \mathcal{G} -Breuil-Kisin module \mathcal{P}_{BK} with the initial (\mathcal{G}, μ) -shtuka $(\mathcal{P}_{\tilde{x}}, \phi_{\mathcal{P}_{\tilde{x}}})$ implies that this cocharacter is conjugate to μ^{-1} .

The conditions a) and b) of Proposition 7.1.3 are now satisfied for \tilde{x}^* (the p -divisible group $\mathcal{G}_{\tilde{x}}$ is “ (\mathcal{G}, μ^{-1}) -adapted”, in the terminology of [KZ21]).¹ It follows that the morphism $\tilde{x}^* : R \otimes_W O \rightarrow O_K$ inducing $\mathcal{G}_{\tilde{x}}$ factors through $R_{\mathcal{G}} \rightarrow O_K$. This gives a K -valued point of $\text{Spf}(R_{\mathcal{G}})^{\text{rig}}$ and hence x belongs to $\text{Spf}(R_{\mathcal{G}})^{\text{rig}}(K)$. \square

Now $(\widehat{\mathcal{M}_{/x_0}^{\text{int}}})_\eta$ and $\text{Spf}(R_{\mathcal{G}})^{\text{rig}}$ are both smooth rigid analytic spaces over $\text{Sp}(\check{E})$ of the same dimension, both closed in $\text{Spf}(R_{\check{E}})^{\text{rig}}$. Since $R_{\mathcal{G}}$ is normal, $\text{Spf}(R_{\mathcal{G}})^{\text{rig}}$ is connected, cf. [deJ95, Lem. 7.3.5]. From Proposition 7.1.4 it now follows that, under our identifications,

$$(\widehat{\mathcal{M}_{/x_0}^{\text{int}}})_\eta = \text{Spf}(R_{\mathcal{G}})^{\text{rig}}. \quad (7.1.7)$$

Let us now complete the proof. Both $\text{Spd}(R_{\mathcal{G}}) \hookrightarrow \text{Spd}(R \otimes_W O)$ and $\widehat{\mathcal{M}_{/x_0}^{\text{int}}} \hookrightarrow \text{Spd}(R \otimes_W O)$ are closed immersions of v -sheaves. By Proposition 4.2.1 (b) and [PR21, Prop. 3.4.9], $(\widehat{\mathcal{M}_{/x_0}^{\text{int}}})_\eta$ is “topologically flat”, i.e. $|(\widehat{\mathcal{M}_{/x_0}^{\text{int}}})_\eta|$ is dense in $|\widehat{\mathcal{M}_{/x_0}^{\text{int}}}|$. Similarly, $\text{Spd}(R_{\mathcal{G}})$ is topologically flat by [PR21, Lem. 3.4.2]. Since by (7.1.7), $|(\widehat{\mathcal{M}_{/x_0}^{\text{int}}})_\eta| = |\text{Spd}(R_{\mathcal{G}})_\eta|$, we deduce that $|\widehat{\mathcal{M}_{/x_0}^{\text{int}}}| = |\text{Spd}(R_{\mathcal{G}})|$. Hence, by [SW20, Lem. 17.4.1], comp. also [Sch17, Prop. 12.15 (iii)], we have

$$\widehat{\mathcal{M}_{/x_0}^{\text{int}}} = \text{Spd}(R_{\mathcal{G}}),$$

under the identification of $\text{Spd}(R \otimes_W O)$ with $\widehat{\mathcal{M}_{\text{GL}(\Lambda), \iota(b), \mu_d / \iota(x_0)}^{\text{int}}} \times_{\text{Spd}(W)} \text{Spd}(O)$ induced by $\mathcal{G}^{\text{univ}}$. (Compare to the argument in the proof of Prop. 5.3.1, case (i).) \square

Remark 7.1.5. In [KP18, §§3.2, 3.3], there is always the blanket assumption that G splits over a tamely ramified extension of \mathbb{Q}_p . This is because essential use is made of “purity” of \mathcal{G}^o -torsors over $\text{Spec}(W(k)[[u]]) \setminus \{(u, p)\}$ which was shown in [KP18, Prop. 1.4.3] under this tameness hypothesis on G . This purity is now proven in all cases by Anschütz [An18]. With this additional ingredient, the proofs of essentially all the statements in [KP18, §§3.1, 3.2, 3.3] go through without the tameness assumption, with no change in the proofs. This observation is the main point in the proofs of [KZ21, Prop. 4.1.7, Prop. 4.2.4] that we use

¹Note that in [KZ21], the stabilizer is denoted by $\tilde{\mathcal{G}}$ and the corresponding parahoric by \mathcal{G} . In [KZ21] the authors often assume that the stabilizers are of points which are generic in their facet but their proofs of the statements we use go through without this assumption.

in the argument above (the conditions (a), (b), (c), appear in [KZ21, (4.1.5)]). Note that when G is acceptable, then in [PR21, §5] there is a simpler proof of the purity statement in question.

Remark 7.1.6. In the PEL-case, the assertion of Theorem 7.1.2 follows from the identification [SW20, §25] of $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}$ with the v -sheaf associated to an RZ-space and the “classical” local model diagram. However, the proof of Theorem 7.1.2 given above does not seem to simplify substantially in the PEL case.

8. EXISTENCE OF HODGE EMBEDDINGS

8.1. The good Hodge type case. The next statement guarantees in many cases that the hypotheses of Theorem 7.1.2 are satisfied.

Proposition 8.1.1. *Let (G, b, μ) be a local Shimura datum. Suppose $p > 2$ and that (G, b, μ) is of Hodge type. Assume that there is a Hodge embedding $\iota_0 : G \hookrightarrow \text{GL}_n$ such that $\iota_0(G)$ contains the scalars. Suppose, in addition, that all the following conditions are satisfied:*

- 1) p does not divide $|\pi_1(G_{\text{der}})|$.
- 2) *There exists a closed embedding $G \hookrightarrow G' = \prod_{i=1}^r \text{Res}_{F_i/\mathbb{Q}_p} H_i$ which induces an isomorphism $G_{\text{der}} \xrightarrow{\sim} G'_{\text{der}}$, where, for each i , H_i is a reductive group over F_i which splits over a tamely ramified extension of F_i , and the Hodge embedding $\iota_0 : G \hookrightarrow \text{GL}_n$ extends to a closed immersion $\iota'_0 : G' \hookrightarrow \text{GL}_n$.*
- 3) *The centralizer of a maximal split torus of $G \otimes_{\mathbb{Q}} \check{\mathbb{Q}}_p$ is R -smooth over $\check{\mathbb{Q}}_p$ (in the sense of [KZ21, Def. 2.4.3]).*

Then for any stabilizer group scheme \mathcal{G} of G , there exists a Hodge embedding ι which satisfies conditions (a), (b) and (c) of Theorem 7.1.2.

In particular, for arbitrary $x \in \mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}(\text{Spd}(k))$, the formal completion $\widehat{\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}}_{/x}$ is representable and there is $y \in \mathbb{M}_{\mathcal{G},\mu}^{\text{loc}}(k)$ such that

$$\widehat{\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}}_{/x} \simeq (\widehat{\mathbb{M}_{\mathcal{G},\mu}^{\text{loc}}}_{/y})^{\diamond}.$$

Remark 8.1.2. When G splits over a tamely ramified extension of \mathbb{Q}_p , conditions (2) and (3) are satisfied: (2) is obvious (one can take $r = 1$, $F_1 = \mathbb{Q}_p$ and $H_1 = G$) and (3) holds because any torus that splits over a tamely ramified extension is R -smooth, cf. [KZ21, Prop. 2.4.6]. Under this additional tame condition, Proposition 8.1.1 follows from the results in [KP18, §1, §2].

The conjunction of conditions 2) and 3) corresponds to the notion of a *regular pair* (G, μ) in [KZ21, Def. 3.4.2]. The conjunction of conditions 1)–3) corresponds to the condition (5.1.2.1) on (G, μ) in [KZ21, §5.1]. We say “corresponds” because in [KZ21], “local Hodge embedding” requires an immersion into a symplectic similitude group. More precisely, in [KZ21] and [KP18], a “Hodge embedding” for $(G, \{\mu\})$ is, by definition, given by $\iota : G \hookrightarrow \text{GSp}(V)$ that maps $\{\mu\}$ to the unique conjugacy class of minuscule cocharacters of $\text{GSp}(V)$ and with $\iota(G)$ containing the scalars. However, the alternating form plays no role here and the arguments in the proof of [KZ21, Cor. 3.4.9] go through in our situation.

Proof. This follows from [KZ21, §3.4]; see in particular the proof of [KZ21, Cor. 3.4.9]. Here is a brief overview of the argument.

Assumptions (1), (2) and (3) imply that there is a (possibly different) Hodge embedding $\iota' : G \hookrightarrow \mathrm{GL}_{n'}$ which extends to a closed embedding $\mathcal{G} \hookrightarrow \mathcal{GL}$ over \mathbb{Z}_p , where \mathcal{GL} is some parahoric group scheme for $\mathrm{GL}_{n'}$, and which also induces an embedding of corresponding local models. This uses, among other ingredients, Landvogt's results [La00] on embeddings of Bruhat-Tits buildings and [KP18, Prop. 1.3.3]. Note here that in [KP18], [KZ21], the local models for $(G, \{\mu\}, \mathcal{G})$ are given using the local models constructed in [PZ13], or, more generally, in [Le01]. See [KZ21, (3.4.4)] for the definition of the local models employed in this construction; this involves passing to the group G' as above. Under our assumptions, these are normal with reduced special fiber, cf. [Le01, Thm. 4.2.7] and [KZ21, Thm. 3.1.6]. Hence, we can see that they agree with the local models $\mathbb{M}_{\mathcal{G}, \mu}^{\mathrm{loc}}$ that represent $\mathbb{M}_{\mathcal{G}, \mu}^v$ as considered above. In the final step of the argument, the Hodge embedding ι' is further adjusted by composing with a suitable homomorphism $\mathrm{GL}_{n'} \hookrightarrow \prod_{j=1}^f \mathrm{GL}_{n'} \subset \mathrm{GL}_{fn'}$ such that the parahoric \mathcal{GL} can be taken to be hyperspecial, i.e. $\mathcal{GL} = \mathrm{GL}(\Lambda)$, for some lattice Λ . Then the corresponding local model for GL_h with $h = fn'$ is a (smooth) Grassmannian $\mathrm{Gr}(d, \Lambda)$. This gives (a) and (b) and one can see that (c) remains true. \square

8.2. Central lifts of good Hodge type. The next proposition relates in case (A) a local Shimura datum of abelian type to the good Hodge type cases treated in Proposition 8.1.1.

Proposition 8.2.1. *Let (G, b, μ) be a local Shimura datum of abelian type and $p > 2$. Write*

$$(G_{\mathrm{ad}}, \mu_{\mathrm{ad}}) \simeq \prod_{i=1}^m (\mathrm{Res}_{F_i/\mathbb{Q}_p} H_i, \mu_i), \quad (8.2.1)$$

with all H_i absolutely simple. Assume that all μ_i are non-trivial. Then there exists a central lift (G_1, b_1, μ_1) of Hodge type for (G, b, μ) as in Definition 2.1.3 with the following properties:

- 1) *The hypotheses of Proposition 8.1.1 are satisfied for (G_1, μ_1) .*
- 2) *$E_1 = E_{\mathrm{ad}}$.*

Proof. This closely follows the blueprint of the proof of Proposition 2.3.10 in [De79] (see also [KZ21, Prop. 5.2.7] and [KP18, Lem. 4.6.22]). In this p -adic case, things are actually simpler for two reasons: we do not care about giving a symplectic form and we only keep track of μ and not of a Hodge structure.

We first observe that we may assume that G_{ad} is \mathbb{Q}_p -simple. Indeed, we can construct (G_1, μ_1) by taking the product over the factors in (8.2.1). From now on let $G_{\mathrm{ad}} = \mathrm{Res}_{F/\mathbb{Q}_p}(H)$, with H absolutely simple over F .

We index by I the set $\{\phi_v\}_{v \in I}$ of embeddings $\phi_v : F \hookrightarrow \bar{\mathbb{Q}}_p$. For each $v \in I$, we set \mathcal{D}_v for the Dynkin diagram of $H \otimes_{F, \phi_v} \bar{\mathbb{Q}}_p$. Also write

$$\mu : \mathbb{G}_m/\bar{\mathbb{Q}}_p \longrightarrow (\mathrm{Res}_{F/\mathbb{Q}_p} H) \otimes_{\mathbb{Q}_p} \bar{\mathbb{Q}}_p = \prod_{v \in I} H \otimes_{F, \phi_v} \bar{\mathbb{Q}}_p, \quad \mu = (\mu_v)_{v \in I}.$$

By the arguments in [De79, 1.3.6, 1.3.8], the condition that $(G_{\mathrm{ad}}, \mu_{\mathrm{ad}})$ is of abelian type implies that for each $v \in I$, μ_v is either trivial, or it corresponds to a node s_v in the Dynkin diagram \mathcal{D}_v which is among the “encircled nodes” of the diagrams displayed in the table [De79, 1.3.9]. In fact, $(G_{\mathrm{ad}}, \mu_{\mathrm{ad}})$ is one of types **A**, **B**, **C**, **D^R**, **D^H** as described in [De79,

2.3.8], see also [Se79, §3, Annexe]. For example, if $n > 4$, $\mathbf{D}^{\mathbb{H}}$ means that each factor is (D_n, ω_n^\vee) or (D_n, ω_{n-1}^\vee) and $\mathbf{D}^{\mathbb{R}}$ means that each factor is (D_n, ω_1^\vee) , cf. [HLR18, §5]. Note that, by [Se79, §3, Cor. 2], since μ is not trivial, H cannot be a trialitarian form $D_4^{(3)}$ or $D_4^{(6)}$. Recall that we assume $p > 2$. Hence, it follows that H splits over a tamely ramified extension of F .

Denote by I_c the set of $v \in I$ for which μ_v is trivial and set $I_{nc} = I \setminus I_c$. Set

$$\mathcal{D} = \sqcup_{v \in I} \mathcal{D}_v.$$

The Galois group $\Gamma = \text{Gal}(\bar{\mathbb{Q}}_p/\mathbb{Q}_p)$ acts on \mathcal{D} compatibly with its action on the set of embeddings I .

By our assumption μ is not trivial, so $I_{nc} \neq \emptyset$. Let $S \subset \mathcal{D}$ be a Galois stable subset such that, for each $v \in I_{nc}$, $S \cap \mathcal{D}_v$ is a non-empty set of nodes in \mathcal{D}_v which are among the underlined nodes in the table [De79, 1.3.9] for (\mathcal{D}_v, s_v) .² A node $s \in S \cap \mathcal{D}_v$ corresponds to an irreducible representation

$$\rho_s : G_{\text{sc}, \bar{\mathbb{Q}}_p} \longrightarrow H_{\text{sc}} \otimes_{F, \phi_v} \bar{\mathbb{Q}}_p \longrightarrow \text{GL}(V(s))$$

over $\bar{\mathbb{Q}}_p$. Here, G_{sc} and H_{sc} are the simply connected covers. Consider

$$\bigoplus_{s \in S} V(s)^{\oplus n}$$

which, for sufficiently divisible n , gives a representation V of G_{sc} defined over \mathbb{Q}_p ; denote by V_s the unique irreducible factor of $V_{\bar{\mathbb{Q}}_p}$ isomorphic to $V(s)^{\oplus n}$; the Galois group Γ permutes the factors V_s by an action compatible with its action on $S \subset \mathcal{D}$. There is a finite étale \mathbb{Q}_p -algebra K_S such that $\text{Hom}_{\mathbb{Q}_p}(K_S, \bar{\mathbb{Q}}_p) \simeq S$ as Γ -sets and so the decomposition

$$V \otimes_{\mathbb{Q}_p} \bar{\mathbb{Q}}_p \simeq \bigoplus_{s \in S} V_s$$

is induced by a corresponding K_S -module structure on V which is such that on V_s , K_S acts via the map $K_S \rightarrow \bar{\mathbb{Q}}_p$ that corresponds to s . Note that since $S \rightarrow I$ is Γ -equivariant, K_S is naturally an F -algebra. It is a product $K_S = \prod_j K_j$ of field extensions K_j of F which are all at most tamely ramified over F . The units $K_S^\times = \prod_j K_j^\times$ are the points of a torus $T' = \text{Res}_{F/\mathbb{Q}_p} \prod_j T_j$ with T_j/F splitting over the tame extension K_j/F . We have $K_S^\times \subset \text{GL}(V)$, i.e. $T' \hookrightarrow \text{GL}(V)$, and this centralizes the map $G_{\text{sc}} \rightarrow \text{GL}(V)$. Denote by G'_{sc} the quotient of G_{sc} which acts faithfully on V ; then

$$G_{\text{sc}} \longrightarrow G'_{\text{sc}} \hookrightarrow \text{GL}(V).$$

The group G'_{sc} is the restriction of scalars $G'_{\text{sc}} = \text{Res}_{F/\mathbb{Q}_p} H'_{\text{sc}}$, with H'_{sc} a quotient of H_{sc} . In fact, we see that $H'_{\text{sc}} = H_{\text{sc}}$ in all cases, except for groups of type D_n . In the case of type D_n , the kernel of $H_{\text{sc}} \rightarrow H'_{\text{sc}}$ is either trivial, or of order 2. Set

$$G_1 := G'_{\text{sc}} \cdot T' \hookrightarrow \text{GL}(V).$$

As in [De79], we see that $\mu : \mathbb{G}_{m/\bar{\mathbb{Q}}_p} \rightarrow (G_{\text{ad}})_{\bar{\mathbb{Q}}_p}$ lifts to a fractional cocharacter μ' of $G'_{\text{sc}, \bar{\mathbb{Q}}_p}$. The composition of this fractional character with $G'_{\text{sc}, \bar{\mathbb{Q}}_p} \hookrightarrow \text{GL}(V_{\bar{\mathbb{Q}}_p}) \rightarrow \text{GL}(V_s)$ is trivial when s maps to $v \in I_c$, in which case we set $V_s = V_s(0)$. When s maps to $v \in I_{nc}$, then this

²The table in [De79, 1.3.9] requires a small correction which is explained in the translation of the paper on Milne's webpage: on the first line (case **A**) all nodes should be underlined if $p = 1$.

composition has exactly two distinct weights r_v and $r_v - 1$, with r_v as indicated in the table [De79, 1.3.9], so that

$$V_s = V_s(r_v - 1) \oplus V_s(r_v).$$

We can now consider the direct sum decomposition of $V_{\mathbb{Q}_p}$ into two summands: one, $V_{\mathbb{Q}_p}(0)$, given as the sum of all $V_s(0)$ and all $V_s(r_v - 1)$, and the other, $V_{\mathbb{Q}_p}(1)$, given as the sum of all $V_s(r_v)$. This decomposition defines a cocharacter μ_1 of $\mathrm{GL}(V)_{\mathbb{Q}_p}$ acting by weights 0 and 1 on these two summands and μ_1 factors through $(G_1)_{\mathbb{Q}_p} = (G'_{\mathrm{sc}} \cdot T')_{\mathbb{Q}_p}$.

We now verify that the pair (G_1, μ_1) satisfies the desired assumptions: First of all, we see that $E_1 = E_{\mathrm{ad}}$ and that $\pi_1(G_{1,\mathrm{der}})$ is a 2-group, hence G_1 satisfies condition (1) in Proposition 8.1.1. It is clear that the image $G_1 \hookrightarrow \mathrm{GL}(V)$ contains the scalars (since it contains K_S^\times). We have $G_1 = \mathrm{Res}_{F/\mathbb{Q}_p} H_1$, with $H_1 = H'_{\mathrm{sc}} \cdot (\prod_j T_j)$, so we can take $G_1 = G'_1$ in Proposition 8.1.1 (2). It remains to show that the condition of (3) of Proposition 8.1.1 is satisfied, i.e. that the centralizer of a maximal split torus of $G_1 \otimes_{\mathbb{Q}_p} \check{\mathbb{Q}}_p$ is R -smooth over $\check{\mathbb{Q}}_p$. By the above, this centralizer is the Weil restriction of scalars of a torus which splits over a tamely ramified extension. Such tori are R -smooth over $\check{\mathbb{Q}}_p$ by [KZ21, (2.4.5), Prop. 2.4.6]. \square

9. PROOFS OF THEOREMS 2.5.5 AND 2.5.4

In this section we prove our main theorems. We treat the cases (A) and (B) separately.

9.1. Proof of Theorem 2.5.5 in case (A). Let (G, b, μ) be of abelian type and of type (A), and let \mathcal{G} be quasi-parahoric. Write $(G_{\mathrm{ad}}, b_{\mathrm{ad}}, \mu_{\mathrm{ad}}) = \prod_i (\mathrm{Res}_{F_i/\mathbb{Q}_p} H_i, b_i, \mu_i)$, where H_i is absolutely simple. We split $(G_{\mathrm{ad}}, b_{\mathrm{ad}}, \mu_{\mathrm{ad}})$ into the product of two factors: in the first factor we lump together all components $(\mathrm{Res}_{F_i/\mathbb{Q}_p} H_i, b_i, \mu_i)$ where μ_i is trivial, and in the second factor we lump together all components $(\mathrm{Res}_{F_i/\mathbb{Q}_p} H_i, b_i, \mu_i)$ where μ_i is non-trivial. Let us first assume that the first factor is trivial.

Write $\mathcal{G}_{\mathbf{x}}^0 \subset \mathcal{G} \subset \mathcal{G}_{\mathbf{x}}$, with $\mathcal{G}_{\mathbf{x}}(\check{\mathbb{Z}}_p)$ the stabilizer in $G(\check{\mathbb{Q}}_p)$ of a point \mathbf{x} in the extended building $\mathcal{B}^e(G, \mathbb{Q}_p)$ of $G(\mathbb{Q}_p)$. Let \mathbf{x}_{ad} be the corresponding point in the building $\mathcal{B}(G_{\mathrm{ad}}, \mathbb{Q}_p) = \mathcal{B}^e(G_{\mathrm{ad}}, \mathbb{Q}_p)$. Using Proposition 8.2.1, we produce (G_1, b_1, μ_1) of Hodge type which satisfies the assumptions (1)-(2). We can lift \mathbf{x}_{ad} to \mathbf{x}_1 under the canonical map $\mathcal{B}^e(G_1, \mathbb{Q}_p) \rightarrow \mathcal{B}(G_{\mathrm{ad}}, \mathbb{Q}_p)$. Consider the stabilizer $\mathcal{G}_1(\check{\mathbb{Z}}_p)$ of \mathbf{x}_1 ; this is a quasi-parahoric subgroup of $G_1(\check{\mathbb{Q}}_p)$ which comes with a map $\mathcal{G}_1 \rightarrow \mathcal{G}_{\mathrm{ad}}$ extending $G_1 \rightarrow G_{\mathrm{ad}}$. We similarly have $\mathcal{G} \rightarrow \mathcal{G}_{\mathrm{ad}}$, giving $G \rightarrow G_{\mathrm{ad}}$.

We can now apply Proposition 8.1.1 to $(\mathcal{G}_1, b_1, \mu_1)$ and obtain an isomorphism

$$\widehat{\mathcal{M}_{\mathcal{G}_1, b_1, \mu_1/x}^{\mathrm{int}}} \simeq (\widehat{\mathbb{M}_{\mathcal{G}_1, \mu_1/y}^{\mathrm{loc}}})^\diamond.$$

On the other hand, Proposition 5.3.1 applied to the maps $(\mathcal{G}, b, \mu) \rightarrow (\mathcal{G}_{\mathrm{ad}}, b_{\mathrm{ad}}, \mu_{\mathrm{ad}})$ and $(\mathcal{G}_1, b_1, \mu_1) \rightarrow (\mathcal{G}_{\mathrm{ad}}, b_{\mathrm{ad}}, \mu_{\mathrm{ad}})$, gives

$$\begin{aligned} \widehat{\mathcal{M}_{\mathcal{G}, b, \mu/x}^{\mathrm{int}}} &\xrightarrow{\sim} \widehat{\mathcal{M}_{\mathcal{G}_{\mathrm{ad}}, b_{\mathrm{ad}}, \mu_{\mathrm{ad}}/x}^{\mathrm{int}}} \times_{\mathrm{Spd} \check{O}_{\mathrm{ad}}} \mathrm{Spd} \check{O}, \\ \widehat{\mathcal{M}_{\mathcal{G}_1, b_1, \mu_1/x}^{\mathrm{int}}} &\xrightarrow{\sim} \widehat{\mathcal{M}_{\mathcal{G}_{\mathrm{ad}}, b_{\mathrm{ad}}, \mu_{\mathrm{ad}}/x}^{\mathrm{int}}}. \end{aligned} \tag{9.1.1}$$

(Recall here that $E_1 = E_{\mathrm{ad}}$, so $O_1 = O_{\mathrm{ad}}$.) Combining these with the isomorphisms

$$\widehat{\mathbb{M}_{\mathcal{G}, \mu/y}^{\mathrm{loc}}} \simeq \widehat{\mathbb{M}_{\mathcal{G}_{\mathrm{ad}}, \mu_{\mathrm{ad}}/y}^{\mathrm{loc}}} \times_{\mathrm{Spd}(\check{O}_{\mathrm{ad}})} \mathrm{Spd}(\check{O}) \simeq \widehat{\mathbb{M}_{\mathcal{G}_1, \mu_1/y}^{\mathrm{loc}}} \times_{\mathrm{Spd}(\check{O}_1)} \mathrm{Spd}(\check{O}) \tag{9.1.2}$$

obtained from [SW20, Prop. 21.5.1] gives the isomorphism

$$\widehat{\mathcal{M}_{\mathcal{G},b,\mu/x}^{\text{int}}} \xrightarrow{\sim} (\widehat{\mathbb{M}_{\mathcal{G},\mu/y}^{\text{loc}}})^\diamond.$$

This concludes the proof of Theorem 2.5.5 when the first factor of $(G_{\text{ad}}, b_{\text{ad}}, \mu_{\text{ad}})$ is trivial.

Now let us consider the general adjoint case. Using the compatibility of $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}$ and $\mathbb{M}_{\mathcal{G},\mu}^v$ with products (cf. §2.4), it suffices to consider each factor separately. The second factor has been treated above. For the first factor the assertion follows from section 6. Now the passage from the adjoint case to the general case follows by the same argument as above, making use of the isomorphism in the first line in (9.1.1) and the first isomorphism in (9.1.2). \square

9.2. Proof of Theorem 2.5.4 in case (A). Via the ad-isomorphism $G \rightarrow G_{\text{ad}}$ it follows using Theorem 5.1.2 that $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}}$ is representable by a normal formal scheme locally formally of finite type over \check{O} , provided this representability holds for the adjoint group. As in the above proof, we may consider separately the case where μ_{ad} is trivial and the case where all components of μ_{ad} are non-trivial. The first case follows from section 6. Let us consider the second case.

As in the above proof, using Proposition 8.2.1 we construct (G_1, b_1, μ_1) of Hodge type with \mathcal{G}_1 a stabilizer group scheme such that there is a closed “Hodge” embedding

$$\mathcal{G}_1 \hookrightarrow \text{GL}(\Lambda).$$

Then, using Theorem 2.5.5 we deduce that $\widehat{\mathcal{M}_{\mathcal{G}_1,b_1,\mu_1/x}^{\text{int}}}$ is representable by the formal spectrum of a noetherian normal complete local ring, for any $x \in \mathcal{M}_{\mathcal{G}_1,b_1,\mu_1}^{\text{int}}(\text{Spd}(k))$. The proof of [PR21, Thm. 3.7.1] now applies and we obtain that $\mathcal{M}_{\mathcal{G}_1,b_1,\mu_1}^{\text{int}}$ is representable by a formal scheme which is normal and flat locally formally of finite type over \check{O} . (Note that [PR21, Thm. 3.7.1] is stated for parahoric \mathcal{G} . However, given the construction of the specialization map for quasi-parahorics as in §3.4 above, the argument extends in a straightforward fashion to our situation, in which \mathcal{G}_1 is quasi-parahoric. In fact, we do not need to use that \mathcal{G}_1 is a stabilizer group scheme. The argument in loc. cit. only requires $\mathcal{G}_1(\check{\mathbb{Z}}_p) = \text{GL}(\Lambda \otimes_{\mathbb{Z}_p} \check{\mathbb{Z}}_p) \cap G_1(\check{\mathbb{Q}}_p)$ and this holds here since $\mathcal{G}_1 \hookrightarrow \text{GL}(\Lambda)$ is a closed immersion.) Since the maps $G \rightarrow G_{\text{ad}}$ and $G_1 \rightarrow G_{\text{ad}}$ are both ad-isomorphisms, it now follows using Theorem 5.1.2 twice that $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau}$, for each $\tau \in \Omega_G$, is representable by such a formal scheme. It follows that $\mathcal{M}_{\mathcal{G},b,\mu}^{\text{int}} = \sqcup_{\tau} \mathcal{M}_{\mathcal{G},b,\mu}^{\text{int},\tau}$ is representable by a formal scheme which is normal and flat locally formally of finite type over \check{O} . \square

9.3. Proof of Theorems 2.5.5, 2.5.4 in case (B). Recall that in case (B) we have $p = 2$ and $G_{\text{ad}} = \prod_{i=1}^m \text{Res}_{F_i/\mathbb{Q}_2} H_i$, with $H_i = B_i^\times / F_i^\times$, or $H_i = \text{PGSp}_{2n_i}$, or μ_i trivial. As in the proofs of Theorems 2.5.4 and 2.5.5 in the case (A), we can easily reduce to the case that $m = 1$ and assume $G_{\text{ad}} = \text{Res}_{F/\mathbb{Q}_2} H$ with $H = B^\times / F^\times$, or $H = \text{PGSp}_{2n}$. In the first subcase, we take $G_1 = \text{Res}_{F/\mathbb{Q}_2} B^\times$. In the second subcase, we first set

$$\langle v, w \rangle = \text{Tr}_{F/\mathbb{Q}_2}(v, w)$$

where (v, w) is the standard perfect alternating F -bilinear form on F^{2n} . Then we take $G_1 = J$, with the group J defined by

$$J(R) = \{g \in \text{GL}_{2n}(F \otimes_{\mathbb{Q}_p} R) \mid \langle gv, gw \rangle = c(g)\langle v, w \rangle, \ c(g) \in R^\times\}.$$

In each subcase, we lift μ_{ad} to a corresponding minuscule μ_1 .

The quasi-parahoric \mathcal{G} gives a corresponding point \mathbf{x} in the extended Bruhat-Tits building $\mathcal{B}^e(G, \mathbb{Q}_2)$, so that $\mathcal{G}^o = \mathcal{G}_{\mathbf{x}}^o(\check{\mathbb{Z}}_2) \subset \mathcal{G}(\check{\mathbb{Z}}_2) \subset \mathcal{G}_{\mathbf{x}}(\check{\mathbb{Z}}_2)$. Consider the stabilizer group scheme $\mathcal{G}_1 := \mathcal{G}_{1, \mathbf{x}_1}$ of G_1 which corresponds to a point \mathbf{x}_1 in $\mathcal{B}^e(G_1, \mathbb{Q}_2)$ that lifts the point \mathbf{x}_{ad} in the building $\mathcal{B}(G_{\text{ad}}, \mathbb{Q}_2)$, as in the argument in the proof for the case (A) above. The devissage results of §4 (Theorem 4.1.1 and its proof), allows us to replace $\mathcal{G}_{1, \mathbf{x}_1}$ by the stabilizer group scheme of a point which is generic in the smallest facet that contains it, and still has the same parahoric neutral component. Therefore, we can assume that \mathbf{x}_1 already has this “genericity” property. We can now see, using [RZ96, App. to Chapt. 3] and the standard explicit description of the buildings for these groups ([BT84], [BT87]), that there is a lattice chain (\mathcal{L}) , resp. a self-dual lattice chain (\mathcal{L}) , such that the stabilizer group scheme above is given as a scheme theoretic stabilizer of (\mathcal{L}) in G_1 . The data (G_1, b_1, μ_1) together with the lattice chain (\mathcal{L}) determine integral EL-, resp. PEL-data \mathcal{D} as in [RZ96], see also [SW20, Def. 24.3.3].

Consider the corresponding RZ formal scheme $\mathcal{M}_{\mathcal{D}}^{\text{naive}}$ (as defined in [RZ96]; the hypothesis $p \neq 2$ is not needed in this case). Then $\mathcal{M}_{\mathcal{D}}^{\text{naive}}$ is a formal scheme locally formally of finite type over $O_{\check{E}}$. Under our assumptions, $\mathcal{M}_{\mathcal{D}}^{\text{naive}}$ has formal completions at closed points which agree with those of the naive local model $\mathbb{M}_{\mathcal{D}}^{\text{naive}}$ ([RZ96, Prop. 3.33]); recall that this follows by Grothendieck-Messing deformation theory which applies for $p = 2$. Note that in general, $\mathbb{M}_{\mathcal{D}}^{\text{naive}}$ and $\mathcal{M}_{\mathcal{D}}^{\text{naive}}$ are not flat over $O_{\check{E}}$ (this accounts for the terminology “naive”).

Set $\mathcal{M}_{\mathcal{D}} := \mathcal{M}_{\mathcal{D}}^{\text{flat}}$ to be the closed formal subscheme of $\mathcal{M}_{\mathcal{D}}^{\text{naive}}$ given by chains of 2-divisible groups which are “ $\mathbb{M}_{\mathcal{G}_1, \mu_1}^{\text{loc}}$ -admissible” in the sense of [SW20, Lect. 24, 25]. Note that under our assumptions, $\mathbb{M}_{\mathcal{G}_1, \mu_1}^{\text{loc}}$ is a closed subscheme of the “naive” local model $\mathbb{M}_{\mathcal{D}}^{\text{naive}}$ which is identified with the flat closure of its generic fiber; in particular, this flat closure has reduced special fiber. Indeed, in the case where $G_1 = \text{Res}_{F/\mathbb{Q}_2} B^\times$, this follows from [PR05, Thm. 7.3] (based on Görtz [G01]). In the case where $G_1 = J$, this follows from [PR05, Thm. 12.4] (based on [G03, G05]). (These two results also follow from the proof of the coherence conjecture by Zhu [Zhu14].)

By [SW20, Cor. 25.1.3], we have

$$(\mathcal{M}_{\mathcal{D}})^\diamond \simeq \mathcal{M}_{\mathcal{G}_1, b_1, \mu_1}^{\text{int}}$$

as v -sheaves over $\text{Spd}(O_{\check{E}})$. It follows that $\mathcal{M}_{\mathcal{G}_1, b_1, \mu_1}^{\text{int}}$ is represented by the formal scheme $\mathcal{M}_{\mathcal{D}}$. By its construction and the above discussion, we obtain that the formal scheme $\mathcal{M}_{\mathcal{D}}$ has formal completions at closed points which agree with those of $\mathbb{M}_{\mathcal{G}_1, \mu_1}^{\text{loc}}$; in particular, $\mathcal{M}_{\mathcal{D}}$ is normal and flat over $O_{\check{E}}$ since these properties hold for $\mathbb{M}_{\mathcal{G}_1, \mu_1}^{\text{loc}}$. Hence the representability conjecture 2.5.1 holds and $\mathcal{M}_{\mathcal{G}_1, b_1, \mu_1} = \mathcal{M}_{\mathcal{D}}$. We can now pass from (G_1, b_1, μ_1) and \mathcal{G}_1 , to (G, b, μ) and \mathcal{G} , by the same devissage as in the proofs in case (A). This completes the proofs of Theorems 2.5.5 and 2.5.4 in case (B). \square

Remark 9.3.1. Our strategy for the proofs of Theorems 2.5.5 and 2.5.4 in case (B) could be applied to more EL/PEL cases than the ones currently given, provided that in the corresponding cases the results of [RZ96, App. to Chapt. 3] can be suitably modified for $p = 2$. An example is given by the unramified unitary group, cf. [RSZ21, App. A]. On the other hand, it is also reasonable to expect that the strategy of the proof in case (A) could be extended to $p = 2$ (at least assuming that G is acceptable), if the constructions of [KP18], [KZ21] can be

extended to cover $p = 2$. Such an extension was given in [KMP16] in the hyperspecial case, i.e. when \mathcal{G} is reductive over \mathbb{Z}_2 .

REFERENCES

- [An18] J. Anschütz, *Extending torsors on the punctured $\mathrm{Spec}(A_{\mathrm{inf}})$* . J. Reine Angew. Math. 783 (2022), 227–268.
- [AGLR22] J. Anschütz, I. Gleason, J. N. P. Lourenço, T. Richarz, *On the p -adic theory of local models*. Preprint, arXiv:2201.01234
- [BS17] B. Bhatt, P. Scholze, *Projectivity of the Witt vector affine Grassmannian*. Invent. Math. 209 (2017), 329–423.
- [BS15] B. Bhatt, P. Scholze, *The pro-étale topology for schemes*. Astérisque No. 369 (2015), 99–201.
- [BTII] F. Bruhat and J. Tits, *Groupes réductifs sur un corps local. II. Schémas en groupes. Existence d’une donnée radicielle valuée*. Publ. Math. IHES, 60 (1984), 197–376.
- [BT84] F. Bruhat and J. Tits, *Schémas en groupes et immeubles des groupes classiques sur un corps local*. Bull. Soc. Math. France 112 (1984), no. 2, 259–301.
- [BT87] F. Bruhat and J. Tits, *Schémas en groupes et immeubles des groupes classiques sur un corps local. II. Groupes unitaires*. Bull. Soc. Math. France 115 (1987), no. 2, 141–195.
- [BP20] O. Bültel, G. Pappas, *(G, μ) -displays and Rapoport-Zink spaces*. J. Inst. Math. Jussieu 19 (2020), no. 4, 1211–1257.
- [deJ95] A. J. de Jong, *Crystalline Dieudonné module theory via formal and rigid geometry*. Inst. Hautes Études Sci. Publ. Math. No. 82 (1995), 5–96.
- [De79] P. Deligne, *Variétés de Shimura: interprétation modulaire, et techniques de construction de modèles canoniques*. Automorphic forms, representations and L -functions, Part 2, Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979, pp. 247–289.
- [Dr77] V. Drinfeld, *Coverings of p -adic symmetric regions*. Funct. Anal. Appl. 10 (1977), 29–40.
- [FS21] L. Fargues, P. Scholze, *Geometrization of the local Langlands correspondence*. Preprint, arXiv:2102.13459.
- [Gl20] I. Gleason, *Specialization maps for Scholze’s category of diamonds*. Preprint, arXiv:2012.05483.
- [GLo22] I. Gleason, J. N. P. Lourenço, *Tubular neighborhoods of local models*. Preprint, arXiv:2204.05526
- [G01] U. Görtz, *On the flatness of models of certain Shimura varieties of PEL-type*. Math. Ann. 321 (2001), no. 3, 689–727.
- [G03] U. Görtz, *On the flatness of local models for the symplectic group*. Adv. Math. 176 (2003), no. 1, 89–115.
- [G05] U. Görtz, *Topological flatness of local models in the ramified case*. Math. Z. 250 (2005), no. 4, 775–790.
- [GHN20] U. Görtz, X. He, S. Nie *Basic loci of Coxeter type with arbitrary parahoric level*. Preprint, arXiv:2006.08838
- [HR08] T. Haines, M. Rapoport, *On parahoric subgroups*, Appendix to [PR08].
- [HLR18] T. Haines, J. Lourenço, T. Richarz, *On the normality of Schubert varieties: Remaining cases in positive characteristics*. Preprint, arXiv:1806.11001
- [HR19] T. Haines, T. Richarz, *Normality and Cohen-Macaulayness of parahoric local models*. Preprint, arXiv:1903.10585
- [HK19] P. Hamacher, W. Kim, *l -adic étale cohomology of Shimura varieties of Hodge type with non-trivial coefficients*. Math. Ann. 375 (2019), no. 3–4, 973–1044.
- [He16] X. He, *Kottwitz-Rapoport conjecture on unions of affine Deligne-Lusztig varieties*. Ann. Sci. ENS 49 (2016), 1125–1141.
- [HPR20] X. He, G. Pappas, M. Rapoport, *Good and semi-stable reductions of Shimura varieties*. J. de l’Ecole Polytechnique Math. 7 (2020), 497–571.
- [HZ20] X. He, R. Zhou, *On the connected components of affine Deligne-Lusztig varieties*. Duke Math. J. 169 (2020), 2697–2765.
- [HP17] B. Howard, G. Pappas, *Rapoport-Zink spaces for spinor groups*. Compos. Math. 153 (2017), no. 5, 1050–1118.

- [Kim18] W. Kim, *Rapoport-Zink spaces of Hodge type*. Forum of Math., Sigma 6 (2018), paper No. e8, 110 pp.
- [KMP16] W. Kim, K. Madapusi Pera, *2-adic integral canonical models*. Forum Math. Sigma 4 (2016), Paper No. e28, 34 pp.
- [Ki10] M. Kisin, *Integral models for Shimura varieties of abelian type*. J. Amer. Math. Soc. 23 (2010), no. 4, 967–1012.
- [KP18] M. Kisin, G. Pappas, *Integral models of Shimura varieties with parahoric level structure*. Publ. Math. IHES 128 (2018), 121–218.
- [KZ21] M. Kisin, R. Zhou, *Independence of ℓ for Frobenius conjugacy classes attached to abelian varieties*. Preprint, arXiv:2103.09945
- [Ko92] R. Kottwitz, *Points on some Shimura varieties over finite fields*. J. Amer. Math. Soc. 5 (1992), no. 2, 373–444.
- [Ko97] R. Kottwitz, *Isocrystals with additional structure. II*. Compositio Math. 109 (1997), no. 3, 255–339.
- [KR11] S. Kudla, M. Rapoport, *Special cycles on unitary Shimura varieties, I. Unramified local theory*. Invent. Math. 184 (2011), no. 3, 629–682.
- [KRZ20] S. Kudla, M. Rapoport, Th. Zink, *On the p -adic uniformization of unitary Shimura curves*. Preprint, arXiv:2007.05211
- [La00] E. Landvogt, *Some functorial properties of the Bruhat-Tits building*. J. Reine Angew. Math. 518 (2000), 213–241.
- [Le01] B. Levin, *Local models for Weil-restricted groups*. Compositio Math. 152 (2016), no. 12, 2563–2601.
- [Pa18] G. Pappas, *Arithmetic models for Shimura varieties*. Proceedings of the International Congress of Mathematicians (Rio de Janeiro, 2018). Vol. II (Hackensack, NJ), World Sci. Publ., 2018, p. 377–398.
- [PR05] G. Pappas, M. Rapoport, *Local models in the ramified case. II. Splitting models*. Duke Math. J. 127 (2005), no. 2, 193–250.
- [PR08] G. Pappas, M. Rapoport, *Twisted loop groups and their affine flag varieties*. Adv. Math. 219 (2008), no. 1, 118–198. With an appendix by Th. Haines and Rapoport.
- [PR21] G. Pappas, M. Rapoport, *p -adic shtukas and the theory of global and local Shimura varieties*. Preprint, arXiv:2106.08270
- [PZ13] G. Pappas, X. Zhu, *Local models of Shimura varieties and a conjecture of Kottwitz*. Invent. math. 194 (2013), 147–254.
- [Ra18] M. Rapoport, *Accessible and weakly accessible period domains*. Appendix to: P. Scholze, *On the p -adic cohomology of the Lubin-Tate tower*, Ann. Sci. ENS 51 (2018), 856–863.
- [RSZ21] M. Rapoport, B. Smithling, W. Zhang, *On Shimura varieties for unitary groups*. Pure Applied Math. Quarterly 17 (2021), 773–837.
- [RSZ18] M. Rapoport, B. Smithling, W. Zhang, *Regular Formal Moduli Spaces and Arithmetic Transfer Conjectures*. Math. Ann. 370 (2018), 1079–1175.
- [RV14] M. Rapoport, E. Viehmann, *Towards a theory of local Shimura varieties*. Münster J. of Math. 7 (2014), 273–326.
- [RZ17] M. Rapoport, Th. Zink, *On the Drinfeld moduli problem of p -divisible groups*. Cambridge J. Math. 5 (2017), 229–279.
- [RZ96] M. Rapoport, T. Zink, *Period spaces for p -divisible groups*. Ann. of Math. Studies 141, Princeton University Press, Princeton, NJ, 1996.
- [Ri16] T. Richarz, *Affine Grassmannians and geometric Satake equivalences*. Int. Math. Res. Not. 12 (2016), 3717–3767.
- [Sch18] P. Scholze, *p -adic geometry*. Proceedings of the International Congress of Mathematicians–Rio de Janeiro 2018. Vol. I. Plenary lectures, 899–933, World Sci. Publ., Hackensack, NJ, 2018.
- [SW20] P. Scholze, J. Weinstein, *Berkeley lectures on p -adic geometry*. Ann. of Math. Studies, 207, Princeton University Press, Princeton, 2020.
- [Sch17] P. Scholze, *Étale cohomology of diamonds*. Preprint, arXiv:1709.07343
- [Se79] J.-P. Serre, *Groupes algébriques associés aux modules de Hodge-Tate*. Journées de Géométrie Algébrique de Rennes. (Rennes, 1978), Vol. III, pp. 155–188, Astérisque, 65, Soc. Math. France, Paris, 1979.

- [Sh20] X. Shen, *On some generalized Rapoport-Zink spaces*. Canad. J. Math. 72 (2020), no. 5, 1111–1187.
- [T79] J. Tits, *Reductive groups over local fields*. Automorphic forms, representations and L-functions, Proc. Sympos. Pure Math., XXXIII, part 1, 29–69, Amer. Math. Soc., Providence, R.I., 1979.
- [Zha12] W. Zhang, *On arithmetic fundamental lemmas*. Invent. Math. 188 (2012), 197–252.
- [Zhu14] X. Zhu, *On the coherence conjecture of Pappas and Rapoport*. Ann. of Math. 180 (2014), 1–85.
- [Zhu17] X. Zhu, *Affine Grassmannians and the geometric Satake in mixed characteristic*. Ann. of Math. (2) 185 (2017), no. 2, 403–492.

DEPT. OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, E. LANSING, MI 48824, USA

E-mail address: `pappasg@msu.edu`

MATHEMATISCHES INSTITUT DER UNIVERSITÄT BONN, ENDENICHER ALLEE 60, 53115 BONN, GERMANY,
AND UNIVERSITY OF MARYLAND, DEPARTMENT OF MATHEMATICS, COLLEGE PARK, MD 20742, USA

E-mail address: `rapoport@math.uni-bonn.de`