

Lectures on Analytic Geometry

Peter Scholze

Contents

Analytic Geometry	5
Preface	5
1. Lecture I: Introduction	6
2. Lecture II: Solid modules	11
3. Lecture III: Condensed \mathbb{R} -vector spaces	16
4. Lecture IV: \mathcal{M} -complete condensed \mathbb{R} -vector spaces	20
Appendix to Lecture IV: Quasiseparated condensed sets	26
5. Lecture V: Entropy and a real B_{dR}^+	28
6. Lecture VI: Statement of main result	33
Appendix to Lecture VI: Recollections on analytic rings	39
7. Lecture VII: $\mathbb{Z}((T))_{>r}$ is a principal ideal domain	42
8. Lecture VIII: Reduction to “Banach spaces”	47
Appendix to Lecture VIII: Completions of normed abelian groups	54
Appendix to Lecture VIII: Derived inverse limits	56
9. Lecture IX: End of proof	57
Appendix to Lecture IX: Some normed homological algebra	65
10. Lecture X: Some computations with liquid modules	68
11. Lecture XI: Towards localization	72
12. Lecture XII: Localizations	78
Appendix to Lecture XII: Topological invariance of analytic ring structures	85
Appendix to Lecture XII: Frobenius	88
Appendix to Lecture XII: Normalizations of analytic animated rings	92
13. Lecture XIII: Analytic spaces	94
14. Lecture XIV: Varia	102
Bibliography	109

Analytic Geometry

Preface

These are lectures notes (to be updated regularly) for a course on analytic geometry taught in the winter term 2019/20 at the University of Bonn. The material presented is part of joint work with Dustin Clausen.

The goal of this course is to use the formalism of analytic rings as defined in the course on condensed mathematics to define a category of analytic spaces that contains (for example) adic spaces and complex-analytic spaces, and to adapt the basics of algebraic geometry to this context; in particular, the theory of quasicohherent sheaves.

October 2019, Peter Scholze

1. Lecture I: Introduction

Mumford writes in *Curves and their Jacobians*: “[Algebraic geometry] seems to have acquired the reputation of being esoteric, exclusive, and very abstract, with adherents who are secretly plotting to take over all the rest of mathematics. In one respect this last point is accurate.”

For some reason, this secret plot has so far stopped short of taking over analysis. The goal of this course is to launch a new attack, turning functional analysis into a branch of commutative algebra, and various types of analytic geometry (like manifolds) into algebraic geometry. Whether this will make these subjects equally esoteric will be left to the reader’s judgement.

What do we mean by analytic geometry? We mean a version of algebraic geometry that

- (1) instead of merely allowing polynomial rings as its basic building blocks, allows rings of convergent power series as basic building blocks;
- (2) instead of being able to define open subsets only by the nonvanishing of functions, one can define open subsets by asking that a function is small, say less than 1;
- (3) strictly generalizes algebraic geometry in the sense that the category of schemes, the theory of quasicoherent sheaves over them, etc., all embed fully faithfully into the corresponding analytic category.

The author always had the impression that the highly categorical techniques of algebraic geometry could not possibly be applied in analytic situations; and certainly not over the real numbers.¹ The goal of this course is to correct this impression.

How can one build algebraic geometry? One perspective is that one starts with the abelian category Ab of abelian groups, with its symmetric monoidal tensor product. Then one can consider rings R in this category (which are just usual rings), and over any ring R one can consider modules M in that category (which are just usual R -modules). Now to any R , one associates the space $\text{Spec } R$, essentially by declaring that (basic) open subsets of $\text{Spec } R$ correspond to localizations $R[f^{-1}]$ of R . One can then glue these $\text{Spec } R$ ’s together along open subsets to form schemes, and accordingly the category of R -modules glues to form the category of quasicoherent sheaves.

Now how to build analytic geometry? We are basically stuck with the first step: We want an abelian category of some kind of topological abelian groups (together with a symmetric monoidal tensor product that behaves reasonably). Unfortunately, topological abelian groups do not form an abelian category: A map of topological abelian groups that is an isomorphism of underlying abelian groups but merely changes the topology, say

$$(\mathbb{R}, \text{discrete topology}) \rightarrow (\mathbb{R}, \text{natural topology}),$$

has trivial kernel and cokernel, but is not an isomorphism.

This problem was solved in the course on condensed mathematics last semester, by replacing the category of topological spaces with the much more algebraic category of condensed sets, and accordingly topological abelian groups with abelian group objects in condensed sets, i.e. condensed abelian groups. Let us recall the definition.

¹In the nonarchimedean case, the theory of adic spaces goes a long way towards fulfilling these goals, but it has its shortcomings: Notably, it lacks a theory of quasicoherent sheaves, and in the nonnoetherian case the general formalism does not work well (e.g. the structure presheaf fails to be sheaf). Moreover, the language of adic spaces cannot easily be modified to cover complex manifolds. This is possible by using Berkovich spaces, but again there is no theory of quasicoherent sheaves etc.

DEFINITION 1.1. Consider the pro-étale site $*_{\text{proét}}$ of the point $*$, i.e. the category $\text{ProFin} \cong \text{Pro}(\text{Fin})$ of profinite sets S with covers given by finite families of jointly surjective maps. A condensed set is a sheaf on $*_{\text{proét}}$; similarly, a condensed abelian group, ring, etc. is a sheaf of abelian groups, rings, etc. on $*_{\text{proét}}$.²

As for sheaves on any site, it is then true that a condensed abelian group, ring, etc. is the same as an abelian group object/ring object/etc. in the category of condensed sets: for example, a condensed abelian group is a condensed set M together with a map $M \times M \rightarrow M$ (the addition map) of condensed sets making certain diagrams commute that codify commutativity and associativity, and admits an inverse map.

More concretely, a condensed set is a functor

$$X : \text{ProFin}^{\text{op}} \rightarrow \text{Sets}$$

such that

- (1) one has $X(\emptyset) = *$, and for all profinite sets S_1, S_2 , the map $X(S_1 \sqcup S_2) \rightarrow X(S_1) \times X(S_2)$ is bijective;
- (2) for any surjection $S' \rightarrow S$ of profinite sets, the map

$$X(S) \rightarrow \{x \in X(S') \mid p_1^*(x) = p_2^*(x) \in X(S' \times_S S')\}$$

is bijective, where $p_1, p_2 : S' \times_S S' \rightarrow S'$ are the two projections.

How to think about a condensed set? The value $X(*)$ should be thought of as the underlying set, and intuitively $X(S)$ is the space of continuous maps from S into X . For example, if T is a topological space, one can define a condensed $X = \underline{T}$ via

$$\underline{T}(S) = \text{Cont}(S, T),$$

the set of continuous maps from S into T . One can verify that this defines a condensed set.³ Part (1) is clear, and for part (2) the key point is that any surjective map of profinite sets is a quotient map.

For example, in analysis a central notion is that of a sequence x_0, x_1, \dots converging to x_∞ . This is codified by maps from the profinite set $S = \{0, 1, \dots, \infty\}$ (the one-point compactification $\mathbb{N} \cup \{\infty\}$ of \mathbb{N}) into X . Allowing more general profinite sets makes it possible to capture more subtle convergence behaviour. For example, if T is a compact Hausdorff space and you have any sequence of points $x_0, x_1, \dots \in T$, then it is not necessarily the case that it converges in T ; but one can always find convergent subsequences. More precisely, for each ultrafilter \mathcal{U} on \mathbb{N} , one can take the limit along the ultrafilter. In fact, this gives a map $\beta\mathbb{N} \rightarrow T$ from the space $\beta\mathbb{N}$ of ultrafilters on \mathbb{N} . The space $\beta\mathbb{N}$ is a profinite set, known as the Stone-Ćech compactification of \mathbb{N} ; it is the initial compact Hausdorff space to which \mathbb{N} maps (by an argument along these lines).

²As discussed last semester, this definition has minor set-theoretic issues. We explained then how to resolve them; we will mostly ignore the issues in these lectures.

³There are some set-theoretic subtleties with this assertion if T does not satisfy the separation axiom $T1$, i.e. its points are not closed; we will only apply this functor under this assumption.

We have the following result about the relation between topological spaces and condensed sets, proved last semester except for the last assertion in part (4), which can be found for example in [BS15, Lemma 4.3.7].⁴

PROPOSITION 1.2. *Consider the functor $T \mapsto \underline{T}$ from topological spaces to condensed sets.*

- (1) *The functor has a left adjoint $X \mapsto X(*)_{\text{top}}$ sending any condensed set X to its underlying set $X(*)$ equipped with the quotient topology arising from the map*

$$\bigsqcup_{S, a \in X(S)} S \rightarrow X(*).$$

- (2) *Restricted to compactly generated (e.g., first-countable, e.g., metrizable) topological spaces, the functor is fully faithful.*
 (3) *The functor induces an equivalence between the category of compact Hausdorff spaces, and quasicompact quasiseparated condensed sets.*
 (4) *The functor induces a fully faithful functor from the category of compactly generated weak Hausdorff spaces (the standard “convenient category of topological spaces” in algebraic topology), to quasiseparated condensed sets. The category of quasiseparated condensed sets is equivalent to the category of ind-compact Hausdorff spaces “ $\varinjlim_i T_i$ ” where all transition maps $T_i \rightarrow T_j$ are closed immersions. If $X_0 \hookrightarrow X_1 \hookrightarrow \dots$ is a sequence of compact Hausdorff spaces with closed immersions and $X = \varinjlim_n X_n$ as a topological space, the map*

$$\varinjlim_n X_n \rightarrow \underline{X}$$

is an isomorphism of condensed sets. In particular, $\varinjlim_n X_n$ comes from a topological space.

Here the notions of quasicompactness/quasiseparatedness are general notions applying to sheaves on any (coherent) site. In our case, a condensed set X is quasicompact if there is some profinite S with a surjective map $S \rightarrow X$. A condensed set X is quasiseparated if for any two profinite sets S_1, S_2 with maps to X , the fibre product $S_1 \times_X S_2$ is quasicompact.

The functor in (4) is close to an equivalence, and in any case (4) asserts that quasiseparated condensed sets can be described in very classical terms. Let us also mention the following related result.

LEMMA 1.3. *Let $X_0 \hookrightarrow X_1 \hookrightarrow \dots$ and $Y_0 \hookrightarrow Y_1 \hookrightarrow \dots$ be two sequences of compact Hausdorff spaces with closed immersions. Then, inside the category of topological spaces, the natural map*

$$\bigcup_n X_n \times Y_n \rightarrow \left(\bigcup_n X_n \right) \times \left(\bigcup_n Y_n \right)$$

is a homeomorphism; i.e. the product on the right is equipped with its compactly generated topology.

PROOF. The map is clearly a continuous bijection. In general, for a union like $\bigcup_n X_n$, open subsets U are the subsets of the form $\bigcup_n U_n$ where each $U_n \subset X_n$ is open. Thus, let $U \subset \bigcup_n X_n \times Y_n$ be any open subset, written as a union of open subset $U_n \subset X_n \times Y_n$, and pick any point $(x, y) \in U$. Then for any large enough n (so that $(x, y) \in X_n \times Y_n$), we can find open neighborhoods $V_n \subset X_n$

⁴Again, some of these assertions run into minor set-theoretic problems; we refer to the notes from last semester. Importantly, everything is valid on the nose when restricted to topological spaces with closed points, and quasiseparated condensed sets; in particular, points (3) and (4) are true as stated.

of x in X_n and $W_n \subset Y_n$ of y in Y_n , such that $V_n \times W_n \subset U_n$. In fact, we can ensure that even $\overline{V_n} \times \overline{W_n} \subset U_n$ by shrinking V_n and W_n . Constructing the V_n and W_n inductively, we may then moreover ensure $\overline{V_n} \subset V_{n+1}$ and $\overline{W_n} \subset W_{n+1}$. Then $V = \bigcup_n V_n \subset \bigcup_n X_n$ and $W = \bigcup_n W_n \subset \bigcup_n Y_n$ are open, and $V \times W = \bigcup_n V_n \times W_n \subset U$ contains (x, y) , showing that U is open in the product topology. \square

We asserted above that one should think of $X(*)$ as the underlying set of X , and about $X(S)$ as the continuous maps from S into X . Note however that in general the map

$$X(S) \rightarrow \prod_{s \in S} X(\{s\}) = \prod_{s \in S} X(*) = \text{Map}(S, X(*))$$

may not be injective, and in fact it may and does happen that $X(*) = *$ but $X(S) \neq *$ for large S . However, for quasiseparated condensed sets, this map is always injective (as follows from (4) above), and in particular a quasiseparated condensed set X is trivial as soon as $X(*)$ is trivial.

The critical property of condensed sets is however exactly that they can also handle non-Hausdorff situations (i.e., non-quasiseparated situations in the present technical jargon) well. For example, condensed abelian groups, like sheaves of abelian groups on any site, form an abelian category. Considering the example from above and passing to condensed abelian groups, we get a short exact sequence

$$0 \rightarrow (\mathbb{R}, \text{discrete topology}) \rightarrow (\mathbb{R}, \text{natural topology}) \rightarrow Q \rightarrow 0$$

of condensed abelian groups, for a condensed abelian group Q satisfying

$$Q(S) = \text{Cont}(S, \mathbb{R}) / \{\text{locally constant maps } S \rightarrow \mathbb{R}\}$$

for any profinite set S .⁵ In particular $Q(*) = 0$ while $Q(S) \neq 0$ for general S : There are plenty of non-locally constant maps $S \rightarrow \mathbb{R}$ for profinite sets S , say any convergent sequence that is not eventually constant. In particular, Q is not quasiseparated. We see that enlarging topological abelian groups into an abelian category precisely forces us to include non-quasiseparated objects, in such a way that a quotient M_1/M_2 still essentially remembers the topology on both M_1 and M_2 .

At this point, we have our desired abelian category, the category $\text{Cond}(\text{Ab})$ of condensed abelian groups. Again, like for sheaves of abelian groups on any site, it has a symmetric monoidal tensor product (representing bilinear maps in the obvious way). We could then follow the same steps as for schemes. However, the resulting theory does not yet achieve our stated goals:

- (1) The basic building blocks in algebraic geometry, the polynomial rings, are exactly the free rings on some set I . As sets are generated by finite sets, really the case of a finite set is relevant, giving rise to the polynomial algebras $\mathbb{Z}[X_1, \dots, X_n]$. Similarly, the basic building blocks are now the free rings on a condensed set; as these are generated by profinite sets, one could also just take the free rings on a profinite set S . But for a profinite set S , the corresponding free condensed ring generated by S is not anything like a ring of (convergent) power series. In fact, the underlying ring is simply the free ring on the underlying set of S , i.e. an infinite polynomial algebra.

To further illustrate this point, let us instead consider the free condensed ring A equipped with an element $T \in A$ with the sequence $T, T^2, \dots, T^n, \dots$ converging 0: This

⁵It is nontrivial that this formula for Q is correct: A priori, this describes the quotient on the level of presheaves, and one might have to sheafify. However, using $H^1(S, M) = 0$ for any profinite set S and discrete abelian group M , as proved last semester, one gets the result by using the long exact cohomology sequence.

is given by $A = \mathbb{Z}[S]/([\infty] = 0)$ for the profinite set $S = \mathbb{N} \cup \{\infty\}$. The underlying ring of A is then still the polynomial algebra $\mathbb{Z}[T]$; it is merely equipped with a nonstandard condensed ring structure.

- (2) One cannot define open subsets by “putting bounds on functions”. If say over \mathbb{Q}_p we want to make the locus $\{|T| \leq 1\}$ into an open subset of the affine line, and agree that this closed unit disc should correspond to the algebra of convergent power series

$$\mathbb{Q}_p\langle T \rangle = \left\{ \sum_{n \geq 0} a_n T^n \mid a_n \rightarrow 0 \right\}$$

while the affine line corresponds to $\mathbb{Q}_p[T]$, then being a localization should mean that the multiplication map

$$\mathbb{Q}_p\langle T \rangle \otimes_{\mathbb{Q}_p[T]} \mathbb{Q}_p\langle T \rangle \rightarrow \mathbb{Q}_p\langle T \rangle$$

should be an isomorphism. However, this is not an isomorphism of condensed abelian groups: On underlying abelian groups, the tensor product is just the usual algebraic tensor product.

These failures are not unexpected: We did not yet put in any nontrivial analysis. Somewhere we have to specify which kinds of convergent power series we want to use. Concretely, in a condensed ring R with a sequence $T, T^2, \dots, T^n, \dots$ converging to 0, we want to allow certain power series $\sum_{n \geq 0} r_n T^n$ where not almost all of the coefficients r_n are zero. Hopefully, this specification maintains a nice abelian category that acquires its own tensor product, in such a way that now the tensor product computation in point (2) above works out.

Last semester, we developed such a formalism that works very well in nonarchimedean geometry: This is the formalism of solid abelian groups that we will recall in the next lecture. In particular, the solidification of the ring A considered in (1) is exactly the power series ring $\mathbb{Z}[[T]]$ (with its condensed ring structure coming from the usual topology on the power series ring), and after solidification the tensor product equation in (2) becomes true. Unfortunately, the real numbers \mathbb{R} are not solid: Concretely, $T = \frac{1}{2} \in \mathbb{R}$ has the property that its powers go to 0, but not any sum $\sum r_n (\frac{1}{2})^n$ with coefficients $r_n \in \mathbb{Z}$ converges in \mathbb{R} . Thus, the solid formalism breaks down completely over \mathbb{R} . Now maybe that was expected? After all, the whole formalism is based on the paradigm of resolving nice compact Hausdorff spaces like the interval $[0, 1]$ by profinite sets, and this does not seem like a clever thing to do over the reals; over totally disconnected rings like \mathbb{Q}_p it of course seems perfectly sensible.

However, last semester we stated a conjecture on how the formalism might be adapted to cover the reals. The first main goal of this course will be to prove this conjecture.

2. Lecture II: Solid modules

In this lecture, we recall the theory of solid modules that was developed last semester.

Recall from the last lecture that we want to put some “completeness” condition on modules in such a way that the free objects behave like some kind of power series. To understand the situation, let us first analyze the structure of free condensed abelian groups.

PROPOSITION 2.1. *Let $S = \varprojlim_i S_i$ be a profinite set, written as an inverse limit of finite sets S_i . For any n , let $\mathbb{Z}[S_i]_{\leq n} \subset \mathbb{Z}[S_i]$ be the subset of formal sums $\sum_{s \in S_i} n_s[s]$ such that $\sum |n_s| \leq n$; this is a finite set, and the natural transition maps $\mathbb{Z}[S_j] \rightarrow \mathbb{Z}[S_i]$ preserve these subsets. There is a natural isomorphism of condensed abelian groups*

$$\mathbb{Z}[S] \cong \bigcup_n \varprojlim_i \mathbb{Z}[S_i]_{\leq n} \subset \varprojlim_i \mathbb{Z}[S_i].$$

In particular, $\mathbb{Z}[S]$ is a countable union of the profinite sets $\mathbb{Z}[S]_{\leq n} := \varprojlim_i \mathbb{Z}[S_i]_{\leq n}$.

Note that the right-hand side indeed defines a subgroup: The addition on $\varprojlim_i \mathbb{Z}[S_i]$ takes

$$\mathbb{Z}[S_i]_{\leq n} \times \mathbb{Z}[S_i]_{\leq n'}$$

into $\mathbb{Z}[S_i]_{\leq n+n'}$. We also remark that the bound imposed is as an ℓ^1 -bound, but in fact it is equivalent to an ℓ^0 -bound, as only finitely many n_s (in fact, n of them) can be nonzero.

PROOF. By definition, $\mathbb{Z}[S]$ is the free condensed abelian group on S , and this is formally given by the sheafification of the functor $T \mapsto \mathbb{Z}[\text{Cont}(T, S)]$. First, we check that the map

$$\mathbb{Z}[S] \rightarrow \varprojlim_i \mathbb{Z}[S_i]$$

is an injection. First, we observe that the map of underlying abelian groups is injective. This means that given any finite formal sum $\sum_{j=1}^k n_j[s_j]$ where the $s_j \in S$ are distinct elements and $n_j \neq 0$ are integers, one can find some projection $S \rightarrow S_i$ such that the image in $\mathbb{Z}[S_i]$ is nonzero. But we can arrange that the images of the s_j in S_i are all distinct, giving the result.

Now, assume that $f \in \mathbb{Z}[\text{Cont}(T, S)]$ maps to 0 in $\varprojlim_i \mathbb{Z}[S_i](T)$. In particular, for all $t \in T$, the specialization $f(t) \in \mathbb{Z}[S]$ is zero by the injectivity on underlying abelian groups. We have to see that there is some finite cover of T by profinite sets $T_m \rightarrow T$ such that the preimage of f in each $\mathbb{Z}[\text{Cont}(T_m, S)]$ is zero. Write $f = \sum_{j=1}^k n_j[g_j]$ where $g_j : T \rightarrow S$ are distinct continuous functions and $n_j \neq 0$. We argue by induction on k . For each pair $1 \leq j < j' \leq k$, let $T_{jj'} \subset T$ be the closed subset where $g_j = g_{j'}$. Then the $T_{jj'}$ cover T : Indeed, if $t \in T$ does not lie in any $T_{jj'}$, then all $g_j(t) \in S$ are pairwise distinct, and then $\sum_{j=1}^k n_j[g_j(t)] \in \mathbb{Z}[S]$ is nontrivial. Thus, we may pass to the cover by the $T_{jj'}$ and thereby assume that $g_j = g_{j'}$ for some $j \neq j'$. But this reduces k , so we win by induction.

As observed before, $\bigcup_n \varprojlim_i \mathbb{Z}[S_i]_{\leq n}$ defines a condensed abelian group, and it admits a map from $S = \varprojlim_i S_i$; in particular, as $\mathbb{Z}[S]$ is the free condensed abelian group on S , the map $\mathbb{Z}[S] \rightarrow \varprojlim_i \mathbb{Z}[S_i]$ factors over $\bigcup_n \varprojlim_i \mathbb{Z}[S_i]_{\leq n}$. It remains to see that the induced map

$$\mathbb{Z}[S] \rightarrow \bigcup_n \varprojlim_i \mathbb{Z}[S_i]_{\leq n}$$

is surjective. For this, consider the map

$$S^n \times \{-1, 0, 1\}^n = \varprojlim_i S_i^n \times \{-1, 0, 1\}^n \rightarrow \varprojlim_i \mathbb{Z}[S_i]_{\leq n}$$

given by $(x_1, \dots, x_n, a_1, \dots, a_n) \mapsto a_1[x_1] + \dots + a_n[x_n]$. This is a cofiltered limit of surjections of finite sets, and so a surjection of profinite sets. This map sits in a commutative diagram

$$\begin{array}{ccc} S^n \times \{-1, 0, 1\}^n & \longrightarrow & \varprojlim_i \mathbb{Z}[S_i]_{\leq n} \\ \downarrow & & \downarrow \\ \mathbb{Z}[S] & \longrightarrow & \bigcup_m \varprojlim_i \mathbb{Z}[S_i]_{\leq m} \end{array}$$

and so implies that the lower map contains $\mathbb{Z}[S_i]_{\leq n}$ in its image. As this works for any n , we get the desired result. \square

REMARK 2.2. In particular, we see that the condensed set $\mathbb{Z}[S]$ is quasiseparated, and in fact comes from a compactly generated weak Hausdorff topological space $\mathbb{Z}[S]_{\text{top}}$, by Proposition 1.2 (4). Moreover, by Lemma 1.3, the addition

$$\mathbb{Z}[S]_{\text{top}} \times \mathbb{Z}[S]_{\text{top}} \rightarrow \mathbb{Z}[S]_{\text{top}}$$

is continuous (not only when the source is equipped with its compactly generated topology), so $\mathbb{Z}[S]$ really comes from a topological abelian group $\mathbb{Z}[S]_{\text{top}}$.

EXERCISE 2.3. Prove that for any compact Hausdorff space S , the condensed abelian group $\mathbb{Z}[S]$ can naturally be written as a countable union $\bigcup_n \mathbb{Z}[S]_{\leq n}$ of compact Hausdorff spaces $\mathbb{Z}[S]_{\leq n}$, and comes from a topological abelian group $\mathbb{Z}[S]_{\text{top}}$.

The idea of solid modules is that we would want to enlarge $\mathbb{Z}[S]$, allowing more sums deemed “convergent”, and an obvious possibility presents itself:

$$\mathbb{Z}[S]^{\blacksquare} := \varprojlim_i \mathbb{Z}[S_i].$$

In particular, this ensures that

$$\mathbb{Z}[\mathbb{N} \cup \{\infty\}]^{\blacksquare} / ([\infty] = 0) = \varprojlim_n \mathbb{Z}\{0, 1, \dots, n-1, \infty\} / ([\infty] = 0) = \varprojlim_n \mathbb{Z}[T]/T^n = \mathbb{Z}[[T]]$$

is the power series algebra.⁶

In other words, we want to pass to a subcategory $\text{Solid} \subset \text{Cond}(\text{Ab})$ with the property that the free solid abelian group on a profinite set S is $\mathbb{Z}[S]^{\blacksquare}$. This is codified in the following definition.

DEFINITION 2.4. A condensed abelian group M is solid if for any profinite set S with a map $f : S \rightarrow M$, there is a unique map of condensed abelian groups $\tilde{f} : \mathbb{Z}[S]^{\blacksquare} \rightarrow M$ such that the composite $S \rightarrow \mathbb{Z}[S]^{\blacksquare} \rightarrow M$ is the given map f .

Let us immediately state the main theorem on solid abelian groups.

⁶Here, $\mathbb{Z}[T]$ denotes the polynomial algebra in a variable T , not the free condensed module on a profinite set T .

THEOREM 2.5 ([Sch19, Theorem 5.8 (i), Theorem 6.2 (i)]). *The category Solid of solid abelian groups is, as a subcategory of the category $\text{Cond}(\text{Ab})$ of condensed abelian groups, closed under all limits, colimits, and extensions. The inclusion functor admits a left adjoint $M \mapsto M^\blacksquare$ (called solidification), which, as an endofunctor of $\text{Cond}(\text{Ab})$, commutes with all colimits and takes $\mathbb{Z}[S]$ to $\mathbb{Z}[S]^\blacksquare$. The category of solid abelian groups admits compact projective generators, which are exactly the condensed abelian groups of the form $\prod_I \mathbb{Z}$ for some set I . There is a unique symmetric monoidal tensor product \otimes^\blacksquare on Solid making $\text{Cond}(\text{Ab}) \rightarrow \text{Solid} : M \mapsto M^\blacksquare$ symmetric monoidal.*

The theorem is rather nontrivial. Indeed, a small fraction of it asserts that \mathbb{Z} is solid. What does this mean? Given a profinite set $S = \varprojlim_i S_i$ and a continuous map $f : S \rightarrow \mathbb{Z}$, there should be a unique map $\tilde{f} : \mathbb{Z}[S]^\blacksquare \rightarrow \mathbb{Z}$ with given restriction to S . Note that

$$\text{Cont}(S, \mathbb{Z}) = \varinjlim_i \text{Cont}(S_i, \mathbb{Z}) = \varinjlim_i \text{Hom}(\mathbb{Z}[S_i], \mathbb{Z}) \rightarrow \text{Hom}(\varprojlim_i \mathbb{Z}[S_i], \mathbb{Z}) = \text{Hom}(\mathbb{Z}[S]^\blacksquare, \mathbb{Z}).$$

In particular, the existence of f is clear, but the uniqueness is not. We need to see that any map $\mathbb{Z}[S]^\blacksquare \rightarrow \mathbb{Z}$ of condensed abelian groups factors over $\mathbb{Z}[S_i]$ for some i . This expresses some ‘‘compactness’’ of $\mathbb{Z}[S]^\blacksquare$ that seems to be hard to prove by a direct attack (if S is a countable limit of finite sets, it is however possible). We note in particular that we do not know whether the similar result holds with \mathbb{Z} replaced by any ring A (and in fact the naive translation of the theorem above fails for general rings).

Let us analyze the structure of $\mathbb{Z}[S]^\blacksquare$. It turns out that it can be regarded as a space of measures. More precisely,

$$\mathbb{Z}[S]^\blacksquare = \varprojlim_i \mathbb{Z}[S_i] = \varprojlim_i \underline{\text{Hom}}(C(S_i, \mathbb{Z}), \mathbb{Z}) = \underline{\text{Hom}}(\varinjlim_i C(S_i, \mathbb{Z}), \mathbb{Z}) = \underline{\text{Hom}}(C(S, \mathbb{Z}), \mathbb{Z})$$

is the space dual to the (discrete) abelian group of continuous functions $S \rightarrow \mathbb{Z}$. Accordingly, we will often denote elements of $\mathbb{Z}[S]^\blacksquare$ as $\mu \in \mathbb{Z}[S]^\blacksquare$ and refer to them as measures.

Thus, in a solid abelian group, it holds true that whenever $f : S \rightarrow M$ is a continuous map, and $\mu \in \mathbb{Z}[S]^\blacksquare$ is a measure, one can form the integral $\int_S f \mu := \tilde{f}(\mu) \in M$. Again, an important special case is when $S = \mathbb{N} \cup \{\infty\}$. In that case, f can be thought of as a convergent sequence $m_0, m_1, \dots, m_n, \dots, m_\infty$ in M , and a measure μ on S can be characterized by the masses $a_0, a_1, \dots, a_n, \dots$ it gives to the finite points (which are isolated in S), as well as the mass a it gives to all of whose S . Then formally we have

$$\int_S f \mu = a_0(m_0 - m_\infty) + a_1(m_1 - m_\infty) + \dots + a_n(m_n - m_\infty) + \dots + a m_\infty ;$$

in other words, the infinite sums on the right are defined in M . In particular, if $m_\infty = 0$, then any sum $\sum a_i m_i$ with coefficients $a_i \in \mathbb{Z}$ is defined in M . Note that the sense in which it is defined is a tricky one: It is not directly as any kind of limit of the finite sums; rather, it is characterized by the uniqueness of the map $\mathbb{Z}[S]^\blacksquare \rightarrow M$ with given restriction to S . Roughly, this says that there is only one way to consistently define such sums for all measures μ on S simultaneously; and then one evaluates for any given μ .

Now we want to explain the proof of Theorem 2.5 *when restricted to \mathbb{F}_p -modules* for some prime p . (The arguments below could be adapted to $\mathbb{Z}/n\mathbb{Z}$ or \mathbb{Z}_p with minor modifications.) This leads to some important simplifications. Most critically, the underlying condensed set of $\mathbb{F}_p[S]^\blacksquare$ is profinite.

In other words, we set

$$\mathbb{F}_p[S]^\blacksquare := \varprojlim_i \mathbb{F}_p[S_i] = \mathbb{Z}[S]^\blacksquare/p$$

and define the category of solid \mathbb{F}_p -modules $\text{Solid}(\mathbb{F}_p) \subset \text{Cond}(\mathbb{F}_p)$ as the full subcategory of all condensed \mathbb{F}_p -modules such that for all profinite sets S with a map $f : S \rightarrow M$, there is a unique extension to a map $\mathbb{F}_p[S]^\blacksquare \rightarrow M$ of condensed \mathbb{F}_p -modules. We note that equivalently we are simply considering the p -torsion full subcategories inside Solid resp. $\text{Cond}(\text{Ab})$.

In the following, we use two simple facts about condensed sets:

- (1) If M is a discrete set, considered as a condensed set, then for any profinite set $S = \varprojlim_i S_i$,

$$\underline{M}(S) = C(S, M) = \varinjlim_i C(S_i, M) = \varinjlim_i \underline{M}(S_i).$$

- (2) If $X_i, i \in I$, is any filtered system of condensed sets, then for any profinite set S ,

$$(\varinjlim_i X_i)(S) = \varinjlim_i X_i(S).$$

For the latter, one has to see that the filtered colimit is still a sheaf on $*_{\text{proét}}$, which follows from the commutation of equalizers with filtered colimits.

PROPOSITION 2.6. *The discrete \mathbb{F}_p -module \mathbb{F}_p is solid.*

PROOF. We have to prove that

$$\text{Hom}(\varprojlim_i \mathbb{F}_p[S_i], \mathbb{F}_p) = \varinjlim_i C(S_i, \mathbb{F}_p).$$

But any map $\varprojlim_i \mathbb{F}_p[S_i] \rightarrow \mathbb{F}_p$ of condensed abelian groups can be regarded as a map of condensed sets. As the source is profinite and the target is discrete, it follows that the map factors over $\mathbb{F}_p[S_i]$ for some i . A priori this factorization is only as condensed sets, but if we assume that the transition maps are surjective, it is automatically a factorization as condensed abelian groups. This gives the desired result. \square

COROLLARY 2.7. *For any set I , the profinite \mathbb{F}_p -vector space $\prod_I \mathbb{F}_p$ is solid.*

PROOF. It follows directly from the definition that the class of solid \mathbb{F}_p -modules is stable under all limits. \square

Let us say that a condensed \mathbb{F}_p -vector space V is profinite if the underlying condensed set is profinite. One checks easily that any such V can be written as a cofiltered limit of finite-dimensional \mathbb{F}_p -vector spaces, and then that $V \mapsto V^* = \text{Hom}(V, \mathbb{F}_p)$ defines an anti-equivalence between profinite \mathbb{F}_p -vector spaces and discrete \mathbb{F}_p -vector spaces. In particular, any profinite \mathbb{F}_p -vector space is isomorphic to $\prod_I \mathbb{F}_p$ for some set I . From here, it is not hard to verify the following proposition:

PROPOSITION 2.8. *The class of profinite \mathbb{F}_p -vector spaces forms an abelian subcategory of $\text{Cond}(\mathbb{F}_p)$ stable under all limits, cokernels, and extensions.*

PROOF. Let $f : V \rightarrow W$ be a map of profinite \mathbb{F}_p -vector spaces. This can be written as a cofiltered limit of maps $f_i : V_i \rightarrow W_i$ of finite-dimensional \mathbb{F}_p -vector spaces. One can take kernels and cokernels of each f_i , and then pass to the limit to get the kernel and cokernel of f , which will then again be profinite \mathbb{F}_p -vector spaces. For stability under extensions, we only have to check that the middle term is profinite as a condensed set, which is clear. \square

THEOREM 2.9. *A condensed \mathbb{F}_p -module M is solid if and only if it is a filtered colimit of profinite \mathbb{F}_p -vector spaces. These form an abelian category stable under all limits and colimits.*

One could also prove the rest of Theorem 2.5 in this setting.

PROOF. We know that profinite \mathbb{F}_p -vector spaces are solid. As $\text{Hom}(\mathbb{F}_p[S]^\blacksquare, -)$ (as a functor from $\text{Cond}(\mathbb{F}_p)$ to sets) commutes with filtered colimits as $\mathbb{F}_p[S]^\blacksquare$ is profinite, the class of solid \mathbb{F}_p -vector spaces is stable under all filtered colimits, and as observed before it is stable under all limits. Next, we observe that the class of filtered colimits of profinite \mathbb{F}_p -vector spaces forms an abelian subcategory of $\text{Cond}(\mathbb{F}_p)$. For this, note that any map $f : V \rightarrow W$ of such is a filtered colimit of maps $f_i : V_i \rightarrow W_i$ of profinite \mathbb{F}_p -vector spaces, and one can form the kernel and cokernel of each f_i and then pass to the filtered colimit again. In particular, they are stable under all colimits (which are generated by cokernels, finite direct sums, and filtered colimits).

It remains to see that any solid M is a filtered colimit of profinite \mathbb{F}_p -vector spaces. Any M admits a surjection $\bigoplus_j \mathbb{F}_p[S_j] \rightarrow M$ for certain profinite sets $S_j \rightarrow M$. As M is solid, this gives a surjection $V \rightarrow M$ where $V = \bigoplus_j \mathbb{F}_p[S_j]^\blacksquare$ is a filtered colimit of profinite \mathbb{F}_p -vector spaces. As solid modules are stable under limits, in particular kernels, the kernel of this map is again solid, and so by repeating the argument we find a presentation $W \rightarrow V \rightarrow M \rightarrow 0$ where W and V are filtered colimits of profinite \mathbb{F}_p -vector spaces. As this class is stable under quotients, we see that also M is such a filtered colimit, as desired. \square

3. Lecture III: Condensed \mathbb{R} -vector spaces

The first goal of this course is to define an analogue of solid modules over the real numbers. Roughly, we want to define a notion of liquid \mathbb{R} -vector spaces that is close to the notion of complete locally convex \mathbb{R} -vector spaces, but is a nice abelian category (which topological vector spaces of any kind are not).

To get started, we will translate some of the classical theory of topological vector spaces into the condensed setup. The most familiar kind of topological \mathbb{R} -vector space are the Banach spaces.

DEFINITION 3.1. A Banach space is a topological \mathbb{R} -vector space V that admits a norm $\|\cdot\|$, i.e. a continuous function

$$\|\cdot\| : V \rightarrow \mathbb{R}_{\geq 0}$$

with the following properties:

- (1) For any $v \in V$, the norm $\|v\| = 0$ if and only if $v = 0$;
- (2) For all $v \in V$ and $a \in \mathbb{R}$, one has $\|av\| = |a|\|v\|$;
- (3) For all $v, w \in V$, one has $\|v + w\| \leq \|v\| + \|w\|$;
- (4) The sets $\{v \in V \mid \|v\| < \epsilon\}$ for varying $\epsilon \in \mathbb{R}_{>0}$ define a basis of open neighborhoods of 0;
- (5) For any sequence $v_0, v_1, \dots \in V$ with $\|v_i - v_j\| \rightarrow 0$ as $i, j \rightarrow \infty$, there exists a (necessarily unique) $v \in V$ with $\|v - v_i\| \rightarrow 0$.

We note that this notion is very natural from the topological point of view in the sense that it is easy to say what the open subsets of V are – they are the unions of open balls with respect to the given norm. On the other hand, V is clearly a metrizable topological space (with distance $d(x, y) = \|y - x\|$), so in particular first-countable, so in particular compactly generated, so the passage to the condensed vector space \underline{V} does not lose information. Let us try to understand \underline{V} ; in other words, we must understand how profinite sets map into V .

PROPOSITION 3.2. *Let V be a Banach space, or more generally a complete locally convex topological vector space. Let S be a profinite set, or more generally a compact Hausdorff space, and let $f : S \rightarrow V$ be a continuous map. Then f factors over a compact absolutely convex subset $K \subset V$, i.e. a compact Hausdorff subspace of V such that for all $x, y \in K$ also $ax + by \in K$ whenever $|a| + |b| \leq 1$.*

In other words, as a condensed set \underline{V} is the union of its compact absolutely convex subsets.

Recall that a topological vector space V is locally convex if it has a basis of neighborhoods U of 0 such that for all $x, y \in U$ also $ax + by \in U$ whenever $|a| + |b| \leq 1$. It is complete if every Cauchy net converges, i.e. for any directed index set I and any map $I \rightarrow V : i \mapsto x_i$, if $x_i - x_j$ converges to 0, then there is a unique $x \in V$ so that $x - x_i$ converges to 0. (Note that V may fail to be metrizable, so one needs to pass to nets.)

To prove the proposition, we need to find K . It should definitely contain all convex combinations of images of points in S ; and all limit points of such. This quickly leads to the idea of integrating a (suitably bounded) measure on S against f . The intuition is that if a profinite set S maps into V , then one can also integrate any measure on S against this map, to produce a map from the space of measures on S towards V .

More precisely, in case $S = \varprojlim_i S_i$ is profinite, consider the space of (“signed Radon”) measures of norm ≤ 1 ,

$$\mathcal{M}(S)_{\leq 1} := \varprojlim_i \mathcal{M}(S_i)_{\leq 1},$$

where $\mathcal{M}(S_i)_{\leq 1} \subset \mathbb{R}[S_i]$ is the subspace of ℓ^1 -norm ≤ 1 . More generally, increasing the norm, we define

$$\mathcal{M}(S) = \bigcup_{c>0} \mathcal{M}(S)_{\leq c}, \quad \mathcal{M}(S)_{\leq c} = \varprojlim_i \mathcal{M}(S_i)_{\leq c}.$$

This can be defined either in the topological or condensed setting, and is known as the space of “signed Radon” measures. We note that we do not regard $\mathcal{M}(S)$ as a Banach space with the norm given by c . We also stress the strong similarity with the description of $\mathbb{Z}[S]$.

EXERCISE 3.3. For any compact Hausdorff space S , there is a notion of signed Radon measure on S – look up the official definition. Show that the corresponding topological vector space $\mathcal{M}(S)$, with the weak topology, is a union of compact Hausdorff subspaces $\mathcal{M}(S)_{\leq c}$ (but beware that the weak topology is not the induced colimit topology). Moreover, show that if S is a profinite set, then a signed Radon measure on S is equivalent to a map μ assigning to each open and closed subset $U \subset S$ of S a real number $\mu(U) \in \mathbb{R}$, so that $\mu(U \sqcup V) = \mu(U) + \mu(V)$ for two disjoint U, V , and there is some constant $C = C(\mu)$ such that for all disjoint decompositions $S = U_1 \sqcup \dots \sqcup U_n$,

$$\sum_{i=1}^n |\mu(U_i)| \leq C.$$

Moreover, for a general compact Hausdorff space S , choose a surjection $\tilde{S} \rightarrow S$ from a profinite set, so that $S = \tilde{S}/R$ for the equivalence relation $R = \tilde{S} \times_S \tilde{S} \subset \tilde{S} \times \tilde{S}$. Show that $\mathcal{M}(S)$ is the coequalizer of $\mathcal{M}(R) \rightrightarrows \mathcal{M}(\tilde{S})$. Thus, the rather complicated notion of signed Radon measures on general compact Hausdorff spaces is simply a consequence of descent from profinite sets.

PROPOSITION 3.4. *Let V be a complete locally convex \mathbb{R} -vector space. Then any continuous map $f : S \rightarrow V$ from a profinite set S extends uniquely to a map of topological \mathbb{R} -vector spaces $\mathcal{M}(S) \rightarrow V : \mu \mapsto \int_S f \mu$. The image of $\mathcal{M}(S)_{\leq 1}$ is a compact absolutely convex subset of V containing S .*

REMARK 3.5. The exercise implies that the same holds true for any compact Hausdorff S . Also note that maps $\mathcal{M}(S) \rightarrow V$ of topological vector spaces are equivalent to maps between the corresponding condensed vector spaces, as the source is compactly generated.

PROOF. The final sentence is clear as V is Hausdorff and so the image of any compact set is compact; and clearly $\mathcal{M}(S)_{\leq 1}$ is absolutely convex (hence so is its image), and the image contains S (as $S \subset \mathcal{M}(S)_{\leq 1}$ as Dirac measures).

We have to construct the map $\mathcal{M}(S) \rightarrow V$, so take $\mu \in \mathcal{M}(S)$, and by rescaling assume that $\mu \in \mathcal{M}(S)_{\leq 1}$. Write $S = \varprojlim_i S_i$ as a limit of finite sets S_i with surjective transition maps, and pick any lift $t_i : S_i \rightarrow S$ of the projection $\pi : S \rightarrow S_i$ (we ask for no compatibilities between different t_i). We define a net in V as follows: For each i , let

$$v_i = \sum_{s \in S_i} f(t_i(s)) \mu(\pi_i^{-1}(s)) \in V.$$

We have to see that this is a Cauchy net, so pick some absolutely convex neighborhood U of 0. Choosing i large enough, we can (by continuity of f) ensure that for any two choices $t_i, t'_i : S_i \rightarrow S$, one has $f(t_i(s)) - f(t'_i(s)) \in U$ (in other words, f varies at most within a translate of U on the preimages of $S \rightarrow S_i$). As $\sum_{s \in S_i} |\mu(\pi_i^{-1}(s))| \leq 1$, we get a convex combination of such differences,

which still lies in U . This verifies that each v_i does not depend on the choice of the t_i up to translation by an element in U , and a slight refinement then proves that one gets a Cauchy net. By completeness of V , we get a unique limit $v \in V$ of the v_i . The proof essentially also gives continuity: Note that the choice of i depending on U only depended on f , not on μ . \square

Thus, from the condensed point of view, the following concepts seem fundamental: Compact absolutely convex sets, and the spaces of measures $\mathcal{M}(S)$ with their compactly generated topology. Note that the latter are ill-behaved from the topological point of view: It is hard to say what a general open subset of $\mathcal{M}(S)$ looks like; one cannot do better than naively taking the increasing union $\bigcup_{c>0} \mathcal{M}(S)_{\leq c}$ of compact Hausdorff spaces.

These concepts lead one to the definition of a Smith space:

DEFINITION 3.6. A Smith space is a complete locally convex topological \mathbb{R} -vector space V that admits a compact absolutely convex subset $K \subset V$ such that $V = \bigcup_{c>0} cK$ with the induced compactly generated topology on V .

One can verify that $\mathcal{M}(S)$ is a Smith space: One needs to check that it is complete and locally convex. We will redefine Smith spaces in the next lecture, and check that it defines a Smith space in the latter sense.

COROLLARY 3.7. *Let V be a complete locally convex topological \mathbb{R} -vector space, and consider the category of Smith spaces $W \subset V$. Then this category is filtered, and as condensed \mathbb{R} -vector spaces*

$$\underline{V} = \varinjlim_{W \subset V} \underline{W}.$$

PROOF. To see that the category is filtered, let $W_1, W_2 \subset V$ be two Smith subspaces, with compact generating subsets K_1, K_2 . Then $K_1 \sqcup K_2 \rightarrow V$ factors over a compact absolutely convex subset $K \subset V$ by the proposition above, and $W = \bigcup_{c>0} cK \subset V$ (with the inductive limit topology) is a Smith space containing both W_1 and W_2 . By the same token, if $f : S \rightarrow V$ is any map from a profinite set, then f factors over a compact absolutely convex subset $K \subset V$, and hence over the Smith space $W = \bigcup_{c>0} cK \subset V$, proving that

$$\underline{V} = \varinjlim_{W \subset V} \underline{W}.$$

\square

This realizes the idea that from the condensed point of view, Smith spaces are the basic building blocks. On the other hand, it turns out that Smith spaces are closely related to Banach spaces:

THEOREM 3.8 (Smith, [Smi52]). *The categories of Smith spaces and Banach spaces are anti-equivalent. More precisely, if V is a Banach space, then $\text{Hom}(V, \mathbb{R})$ is a Smith space; and if W is a Smith space, then $\text{Hom}(W, \mathbb{R})$ is a Banach space, where in both cases we endow the dual space with the compact-open topology. The corresponding biduality maps are isomorphisms.*

This will be proved in the next lecture.

REMARK 3.9. This gives one sense in which Banach spaces are always reflexive, i.e. isomorphic to their bidual. Note, however, that this does not coincide with the usual notion of reflexivity: It is customary to make the dual of a Banach space itself into a Banach space (by using the norm

$\|f\| = \sup_{v \in V, \|v\| \leq 1} |f(v)|$), and then reflexivity asks whether a Banach space is isomorphic to its corresponding bidual. Even if that happens, the two notions of dual are still different: If a topological vector space is both Banach and Smith, it is finite-dimensional.

4. Lecture IV: \mathcal{M} -complete condensed \mathbb{R} -vector spaces

The discussion of the last lecture motivates the following definition.

DEFINITION 4.1. Let V be a condensed \mathbb{R} -vector space. Then V is \mathcal{M} -complete if (the underlying condensed set of) V is quasiseparated, and for all maps $f : S \rightarrow V$ from a profinite set, there is an extension to a map

$$\tilde{f} : \mathcal{M}(S) \rightarrow V$$

of condensed \mathbb{R} -vector spaces.

DEFINITION 4.2 (slight return). A Smith space is an \mathcal{M} -complete condensed \mathbb{R} -vector space V such that there exists some compact Hausdorff $K \subset V$ with $V = \bigcup_{c>0} cK$.

EXERCISE 4.3. Prove that this definition of Smith spaces is equivalent to the previous definition.

We note that it is clear that Smith spaces defined in this way form a full subcategory of topological \mathbb{R} -vector spaces (using Proposition 1.2 and Lemma 1.3). Moreover, Proposition 3.4 implies that any Smith space in the sense of the last lecture is a Smith space in the current sense. It remains to see that for any Smith space in the current sense, the corresponding topological vector space is complete and locally convex. Note that it follows easily that in the current definition one can take K to be absolutely convex.

Let us check first that for any profinite set T , the condensed \mathbb{R} -vector space $\mathcal{M}(T)$ is a Smith space. We only need to see that it is \mathcal{M} -complete. Any map $f : S \rightarrow \mathcal{M}(T)$ factors over $\mathcal{M}(T)_{\leq c}$ for some c . Then writing $T = \varprojlim_i T_i$ as an inverse limit of finite sets T_i , f is an inverse limit of maps $f_i : S \rightarrow \mathcal{M}(T_i)_{\leq c}$. This reduces us to the case that T is finite (provided we can bound the norm of the extension); but then we can apply Proposition 3.4 (and observing that the extension maps $\mathcal{M}(S)_{\leq 1}$ into $\mathcal{M}(T_i)_{\leq c}$).

Our initial hope was that \mathcal{M} -complete condensed \mathbb{R} -vector spaces, without the quasiseparation condition, would behave as well as solid \mathbb{Z} -modules. We will see in the next lecture that this is not so. In this lecture, we however want to show that with the quasiseparation condition, the category behaves as well as it can be hoped for: It is not an abelian category (because cokernels are problematic under quasiseparatedness), but otherwise nice.

First, we check that the extension \tilde{f} is necessarily unique.

PROPOSITION 4.4. *Let V be a quasiseparated condensed \mathbb{R} -vector space and let $g : \mathcal{M}(S) \rightarrow V$ be a map of condensed \mathbb{R} -vector space for some profinite set S . If the restriction of g to S vanishes, then $g = 0$.*

PROOF. The preimage $g^{-1}(0) \subset \mathcal{M}(S)$ is a quasicompact injection of condensed sets. This means that it is a closed subset in the topological sense, see the appendix to this lecture.

On the other hand, $g^{-1}(0)$ contains the \mathbb{R} -vector space spanned by S . This contains a dense subset of $\mathcal{M}(S)_{\leq c}$ for all c , and thus its closure by the preceding. \square

EXERCISE 4.5. Show that in the definition of \mathcal{M} -completeness, one can restrict to extremally disconnected S .

PROPOSITION 4.6. *Any \mathcal{M} -complete condensed \mathbb{R} -vector space V is the filtered colimit of the Smith spaces $W \subset V$; conversely, any filtered colimit of Smith spaces along injections is quasiseparated and \mathcal{M} -complete. For any map $f : V \rightarrow V'$ between \mathcal{M} -complete condensed \mathbb{R} -vector spaces,*

the kernel and image of f (taken in condensed \mathbb{R} -vector spaces) are \mathcal{M} -complete condensed \mathbb{R} -vector spaces. Moreover, f admits a cokernel in the category of \mathcal{M} -complete condensed \mathbb{R} -vector spaces.

PROOF. Let V be an \mathcal{M} -complete condensed \mathbb{R} -vector space and let $f : S \rightarrow V$ be a map from a profinite set. Then f extends uniquely to $\tilde{f} : \mathcal{M}(S) \rightarrow V$. The subset $\mathcal{M}(S)_{\leq 1} \subset \mathcal{M}(S)$ is compact Hausdorff (i.e., a quasicompact quasiseparated condensed set); thus, its image in the quasiseparated V is still quasicompact and quasiseparated, i.e. a compact Hausdorff $K \subset V$. Let $W = \bigcup_{c>0} cK \subset V$. This is a condensed \mathbb{R} -vector space, and it is itself \mathcal{M} -complete. Indeed, assume that T is some profinite set with a map $T \rightarrow W$; by rescaling, we can assume that T maps into K . We get a unique extension to a map $\mathcal{M}(T) \rightarrow V$, and we need to see that it factors over W . We may assume that T is extremally disconnected (as any surjection $T' \rightarrow T$ from an extremally disconnected T' induces a surjection $\mathcal{M}(T') \rightarrow \mathcal{M}(T)$). Then the map $T \rightarrow K$ lifts to a map $T \rightarrow \mathcal{M}(S)_{\leq 1}$. Thus, we get a map $\mathcal{M}(T) \rightarrow \mathcal{M}(S)$ whose composite with $\mathcal{M}(S) \rightarrow V$ must agree with the original map $\mathcal{M}(T) \rightarrow V$ by uniqueness. This shows that the image of $\mathcal{M}(T) \rightarrow V$ is contained in W , as desired. It follows that W is a Smith space.

It is now easy to see that the such of W is filtered, and then V is the filtered colimit, as we have seen that any map from a profinite set S factors over one such W . It is clear that any filtered colimit $\varinjlim_i W_i$ of Smith spaces W_i along injections is quasiseparated (as quasiseparatedness is preserved under filtered colimits of injections). It is also \mathcal{M} -complete, as any map $f : S \rightarrow \varinjlim_i W_i$ factors over one W_i , and then extends to a map $\tilde{f} : \mathcal{M}(S) \rightarrow W_i \rightarrow \varinjlim_i W_i$.

For the assertions about a map f , the claim about the kernel is clear. For the image, we may assume that V is a Smith space, by writing it as a filtered colimit of such; and then that $V = \mathcal{M}(S)$ by picking a surjection $S \rightarrow K \subset V$ where K is a generating compact Hausdorff subset of V . But then the image of f is exactly the Smith space $W \subset V$ constructed in the first paragraph.

For the cokernel of f , we may first take the cokernel Q in the category of condensed \mathbb{R} -vector spaces. Now we may pass to the maximal quasiseparated quotient Q^{qs} of Q , which is still a condensed \mathbb{R} -vector space, by the appendix to this lecture. It is then formal that this satisfies the condition of being \mathcal{M} -complete when tested against extremally disconnected S , as any map $S \rightarrow Q^{\text{qs}}$ lifts to $S \rightarrow W$, and so extends to $\mathcal{M}(S) \rightarrow W \rightarrow Q^{\text{qs}}$. The automatic uniqueness of this extension then implies the result for general profinite S (by covering S by an extremally disconnected $\tilde{S} \rightarrow S$). \square

We will now prove the anti-equivalence between Smith spaces in the current sense, with Banach spaces. (In particular, using the original form of Smith's theorem, this also solves the exercise above.)

THEOREM 4.7. *For any Smith space W , the internal dual $\underline{\text{Hom}}_{\mathbb{R}}(W, \mathbb{R})$ is isomorphic to \underline{V} for a Banach space V . Conversely, for any Banach space V , the internal dual $\underline{\text{Hom}}_{\mathbb{R}}(\underline{V}, \mathbb{R})$ is a Smith space. The corresponding biduality maps are isomorphisms.*

PROOF. Assume first that $W = \mathcal{M}(S, \mathbb{R})$ for some profinite set S . Then $\text{Hom}_{\mathbb{R}}(\mathcal{M}(S, \mathbb{R}), \mathbb{R}) = C(S, \mathbb{R})$ as we have seen that any continuous map $S \rightarrow \mathbb{R}$ extends uniquely to a map of condensed \mathbb{R} -vector spaces $\mathcal{M}(S, \mathbb{R}) \rightarrow \mathbb{R}$. More generally, for any profinite set T ,

$$\underline{\text{Hom}}_{\mathbb{R}}(\mathcal{M}(S, \mathbb{R}), \mathbb{R})(T) = \text{Hom}_{\mathbb{R}}(\mathcal{M}(S, \mathbb{R}), \underline{\text{Hom}}_{\mathbb{R}}(\mathbb{R}[T], \mathbb{R})),$$

and $\underline{\text{Hom}}_{\mathbb{R}}(\mathbb{R}[T], \mathbb{R}) = \underline{C}(T, \mathbb{R})$, for the Banach space $C(T, \mathbb{R})$ with the sup-norm. Indeed, this Banach space has the property that continuous maps $T' \rightarrow C(T, \mathbb{R})$ are equivalent to continuous

maps $T' \times T \rightarrow \mathbb{R}$. Thus, using that Banach spaces are complete locally convex, we see

$$\mathrm{Hom}_{\mathbb{R}}(\mathcal{M}(S, \mathbb{R}), \underline{\mathrm{Hom}}_{\mathbb{R}}(\mathbb{R}[T], \mathbb{R})) = \mathrm{Hom}_{\mathbb{R}}(\mathcal{M}(S, \mathbb{R}), \underline{C}(T, \mathbb{R})) = C(T, \mathbb{R})(S) = C(T \times S, \mathbb{R}),$$

showing that indeed $\underline{\mathrm{Hom}}_{\mathbb{R}}(\mathcal{M}(S, \mathbb{R}), \mathbb{R}) = \underline{C}(S, \mathbb{R})$.

Now assume that $V = \overline{C(S, \mathbb{R})}$. We need to see that for any profinite set T ,

$$\mathrm{Hom}_{\mathbb{R}}(\underline{C}(S, \mathbb{R}), \underline{C}(T, \mathbb{R})) = \mathcal{M}(S)(T).$$

But the left-hand side can be computed in topological vector spaces, and any map of Banach spaces has bounded norm. It suffices to see that maps $C(S, \mathbb{R}) \rightarrow C(T, \mathbb{R})$ of Banach spaces of norm ≤ 1 are in bijection with maps $T \rightarrow \mathcal{M}(S)_{\leq 1}$. But any such map of Banach spaces is uniquely determined by its restriction to $\varinjlim_i C(S_i, \mathbb{R})$, where $S = \varprojlim_i S_i$ is written as a limit of finite sets, where again we assume that the operator norm is bounded by 1. Similarly, maps $T \rightarrow \mathcal{M}(S)_{\leq 1}$ are limits of maps $T \rightarrow \mathcal{M}(S_i)_{\leq 1}$. We can thus reduce to the case S is finite, where the claim is clear. We see also that the biduality maps are isomorphisms in this case.

Now let W be a general Smith space, with a generating subset $K \subset W$. Pick a profinite set S with a surjection $S \rightarrow K$. We get a surjection $\mathcal{M}(S) \rightarrow W$. Its kernel $W' \subset \mathcal{M}(S)$ is automatically quasiseparated, and in fact a Smith space again: It is generated by $W' \cap \mathcal{M}(S)_{\leq 1}$, which is compact Hausdorff. Picking a further surjection $\mathcal{M}(S') \rightarrow W'$, we get a resolution

$$\mathcal{M}(S') \rightarrow \mathcal{M}(S) \rightarrow W \rightarrow 0.$$

Taking $\underline{\mathrm{Hom}}_{\mathbb{R}}(-, \mathbb{R})$, we get

$$0 \rightarrow \underline{\mathrm{Hom}}_{\mathbb{R}}(W, \mathbb{R}) \rightarrow \underline{C}(S, \mathbb{R}) \rightarrow \underline{C}(S', \mathbb{R}).$$

The latter map corresponds to a map of Banach space $C(S, \mathbb{R}) \rightarrow C(S', \mathbb{R})$, whose kernel is a closed subspace that is itself a Banach space V , and we find $\underline{\mathrm{Hom}}_{\mathbb{R}}(W, \mathbb{R}) = \underline{V}$.

Conversely, if V is any Banach space, equipped with a norm, the unit ball $B = \{f \in \mathrm{Hom}(V, \mathbb{R}) \mid \|f\| \leq 1\}$ is compact Hausdorff when equipped with the weak topology.⁷ Picking a surjection $S \rightarrow B$ from a profinite set S , we get a closed immersion $V \rightarrow C(S, \mathbb{R})$ (to check that it is a closed immersion, use that it is an isometric embedding by the Hahn-Banach theorem), whose quotient will then be another Banach space; thus, any Banach space admits a resolution

$$0 \rightarrow V \rightarrow C(S, \mathbb{R}) \rightarrow C(S', \mathbb{R})$$

We claim that taking $\underline{\mathrm{Hom}}_{\mathbb{R}}(-, \mathbb{R})$ gives an exact sequence

$$\mathcal{M}(S') \rightarrow \mathcal{M}(S) \rightarrow \underline{\mathrm{Hom}}_{\mathbb{R}}(\underline{V}, \mathbb{R}) \rightarrow 0.$$

For this, using that we already know about spaces of measures, we have to see that for any extremally disconnected T , the sequence

$$\mathrm{Hom}_{\mathbb{R}}(C(S', \mathbb{R}), C(T, \mathbb{R})) \rightarrow \mathrm{Hom}_{\mathbb{R}}(C(S, \mathbb{R}), C(T, \mathbb{R})) \rightarrow \mathrm{Hom}_{\mathbb{R}}(V, C(T, \mathbb{R})) \rightarrow 0$$

⁷This is known as the Banach-Alaoglu theorem, and follows from Tychonoff by using the closed embedding $B \hookrightarrow \prod_{v \in V} [-1, 1]$.

is exact. This follows from the fact that the Banach space $C(T, \mathbb{R})$ is injective in the category of Banach spaces, cf. [ASC⁺16, Section 1.3]: If $V \subset V'$ is a closed embedding of Banach spaces, then any map $V \rightarrow C(T, \mathbb{R})$ of Banach spaces extends to a map $V' \rightarrow C(T, \mathbb{R})$.⁸

Using these resolutions, it also follows that the biduality maps are isomorphisms in general. \square

We remark that if S is extremally disconnected, then $\mathcal{M}(S)$ is a projective object in the category of Smith spaces (in fact, in the category of \mathcal{M} -complete condensed \mathbb{R} -vector spaces), as S is a projective object in the category of condensed sets. Dually, this means that $C(S, \mathbb{R})$ is an injective object in the category of Banach spaces. For solid abelian groups, it turned out that $\mathbb{Z}[S]^\blacksquare$ is projective in the category of solid abelian groups, for all profinite sets S . Could a similar thing happen here? It turns out, that no:

PROPOSITION 4.8. *For “most” profinite sets S , for example $S = \mathbb{N} \cup \{\infty\}$, or a product $S = S_1 \times S_2$ of two infinite profinite sets S_1, S_2 , the Banach space $C(S, \mathbb{R})$ is not injective; equivalently, the Smith space $\mathcal{M}(S)$ is not projective.*

PROOF. Note that $C(\mathbb{N} \cup \{\infty\}, \mathbb{R})$ is the product of the Banach space c_0 of null-sequences, with \mathbb{R} . One can show that $C(S_1 \times S_2, \mathbb{R})$ contains c_0 as a direct factor, cf. [Cem84], so it is enough to show that c_0 is not injective, which is [ASC⁺16, Theorem 1.25]. \square

In fact, cf. [ASC⁺16, Section 1.6.1], there is no known example of an injective Banach space that is not isomorphic to $C(S, \mathbb{R})$ where S is extremally disconnected!

Next, let us discuss tensor products. Before making the connection with the existing notions for Banach spaces, let us follow our nose and define a tensor product on the category of \mathcal{M} -complete condensed \mathbb{R} -vector spaces.

PROPOSITION 4.9. *Let V and W be \mathcal{M} -complete condensed \mathbb{R} -vector spaces. Then there is an \mathcal{M} -complete condensed \mathbb{R} -vector space $V \otimes_\pi W$ equipped with a bilinear map*

$$V \times W \rightarrow V \otimes_\pi W$$

of condensed \mathbb{R} -vector spaces, which is universal for bilinear maps; i.e. any bilinear map $V \times W \rightarrow L$ to a \mathcal{M} -complete condensed \mathbb{R} -vector space extends uniquely to a map $V \otimes_\pi W \rightarrow L$ of condensed \mathbb{R} -vector spaces.

The functor $(V, W) \mapsto V \otimes_\pi W$ from pairs of \mathcal{M} -complete condensed \mathbb{R} -vector spaces to \mathcal{M} -complete condensed \mathbb{R} -vector spaces commutes with colimits in each variable, and satisfies

$$\mathcal{M}(S) \otimes_\pi \mathcal{M}(S') \cong \mathcal{M}(S \times S')$$

for any profinite sets S, S' .

Note that the case of $S_1 \times S_2$ in Proposition 4.8 means that a tensor product $\mathcal{M}(S_1 \times S_2)$ of two projectives $\mathcal{M}(S_1), \mathcal{M}(S_2)$ is not projective anymore. This is a subtlety that will persist, and that we have to live with.

PROOF. It is enough to show that $\mathcal{M}(S \otimes S')$ represents bilinear maps $\mathcal{M}(S) \times \mathcal{M}(S') \rightarrow L$; indeed, the other assertions will then follow via extending everything by colimits to the general case.

⁸The case $T = *$ is the Hahn-Banach theorem. Part of the Hahn-Banach theorem is that the extension can be bounded by the norm of the original map. This implies the injectivity of $l^\infty(S_0, \mathbb{R}) = C(\beta S_0, \mathbb{R})$ for any set S_0 , and then the result for general T follows by writing T as a retract of βS_0 for some S_0 .

In other words, we have to show that bilinear maps $\mathcal{M}(S) \times \mathcal{M}(S') \rightarrow L$ are in bijection with maps $S \times S' \rightarrow L$. Any map $S \times S' \rightarrow L$ extends to $\mathcal{M}(S \times S') \rightarrow L$, and thus gives a bilinear map $\mathcal{M}(S) \times \mathcal{M}(S') \rightarrow L$, as one can directly define a natural bilinear map $\mathcal{M}(S) \times \mathcal{M}(S') \rightarrow \mathcal{M}(S \times S')$. Thus, it is enough to show that any bilinear map $\mathcal{M}(S) \times \mathcal{M}(S') \rightarrow L$ that vanishes on $S \times S'$ is actually zero. For this, we again observe that it vanishes on a dense subset of $\mathcal{M}(S)_{\leq c} \times \mathcal{M}(S')_{\leq c}$ for any c , and thus on the whole of it, as the kernel is a closed subspace when L is quasiseparated. \square

Tensor products of Banach spaces were defined by Grothendieck, [Gro55], and interestingly he has defined several tensor products. There are two main examples. One is the projective tensor product $V_1 \otimes_{\pi} V_2$; this is a Banach space that represents bilinear maps $V_1 \times V_2 \rightarrow W$.

PROPOSITION 4.10. *Let V_1 and V_2 be two Banach spaces. Then $\underline{V_1} \otimes_{\pi} \underline{V_2} \cong \underline{V_1 \otimes_{\pi} V_2}$.*

REMARK 4.11. The proposition implies that any compact absolutely convex subset $K \subset V_1 \otimes_{\pi} V_2$ is contained in the closed convex hull of $K_1 \times K_2$ for compact absolutely convex subsets $K_i \subset V_i$.

PROOF. Any Banach space admits a projective resolution by spaces of the form $\ell^1(I)$ for some set I ; this reduces us formally to that case. One can even reduce to the case that I is countable, by writing $\ell^1(I)$ as the ω_1 -filtered colimit of $\ell^1(J)$ over all countable $J \subset I$. Thus, we may assume that $V_1 = V_2 = \ell^1(\mathbb{N})$ is the space of ℓ^1 -sequences. In that case $V_1 \otimes_{\pi} V_2 = \ell^1(\mathbb{N} \times \mathbb{N})$.

In that case,

$$\underline{\ell^1(\mathbb{N})} = \varinjlim_{(\lambda_n)_n} \mathcal{M}(\mathbb{N} \cup \{\infty\}) / (\mathbb{R} \cdot [\infty])$$

where the filtered colimit is over all null-sequences of positive real numbers $0 < \lambda_n \leq 1$. Now the result comes down to the observation that any null-sequence $\lambda_{n,m}$ of positive real numbers in $(0, 1]$ parametrized by pairs $n, m \in \mathbb{N}$ can be bounded from above by a sequence of the form $\lambda_n \lambda'_m$ where λ_n and λ'_m are null-sequences of real numbers in $(0, 1]$. Indeed, one can take

$$\lambda_n = \lambda'_n = \sqrt{\max_{n', m' \leq n} \lambda_{n', m'}}. \quad \square$$

On the other hand, there is the injective tensor product $V_1 \otimes_{\epsilon} V_2$; this satisfies $C(S_1, \mathbb{R}) \otimes_{\epsilon} C(S_2, \mathbb{R}) \cong C(S_1 \times S_2, \mathbb{R})$. There is a map $V_1 \otimes_{\pi} V_2 \rightarrow V_1 \otimes_{\epsilon} V_2$, that is however far from an isomorphism.

PROPOSITION 4.12. *Let V_i , $i = 1, 2$, be Banach spaces with dual Smith spaces W_i . Let $W = W_1 \otimes_{\pi} W_2$, which is itself a Smith space, and let V be the Banach space dual to W . Then there is a natural map*

$$V_1 \otimes_{\epsilon} V_2 \rightarrow V$$

that is a closed immersion of Banach spaces. It is an isomorphism if V_1 or V_2 satisfies the approximation property.

We recall that most natural Banach spaces have the approximation property; in fact, it had been a long-standing open problem whether all Banach spaces have the approximation property, until a counterexample was found by Enflo, [Enf73] (for which he was awarded a live goose by Mazur).

PROOF. The injective tensor product has the property that if $V_1 \hookrightarrow V_1'$ is a closed immersion of Banach spaces, then $V_1 \otimes_\epsilon V_2 \hookrightarrow V_1' \otimes_\epsilon V_2$ is a closed immersion. However, it is not in general true that if

$$0 \rightarrow V_1 \rightarrow V_1' \rightarrow V_1''$$

is a resolution, then

$$0 \rightarrow V_1 \otimes_\epsilon V_2 \rightarrow V_1' \otimes_\epsilon V_2 \rightarrow V_1'' \otimes_\epsilon V_2$$

is a resolution: Exactness in the middle may fail. The statement is true when $V_2 = C(S_2, \mathbb{R})$ is the space of continuous functions on some profinite set S_2 , as in that case $W \otimes_\epsilon V_2 = C(S_2, W)$ for any Banach space W , and $C(S_2, -)$ preserves exact sequences of Banach spaces for profinite sets S_2 .

In any case, such resolutions prove the desired statement: Note that if $V_i = C(S_i, \mathbb{R})$, one has $W_i = \mathcal{M}(S_i)$, and then $W = \mathcal{M}(S_1 \times S_2)$ and so $V = C(S_1 \times S_2, \mathbb{R})$, which is indeed $V_1 \otimes_\epsilon V_2$; and these identifications are functorial. Now the resolution above gives the same statement as long as $V_2 = C(S_2, \mathbb{R})$; and in general one sees that V is the kernel of the map $V_1' \otimes_\epsilon V_2 \rightarrow V_1'' \otimes_\epsilon V_2$.

It remains to see that the map $V_1 \otimes_\epsilon V_2 \rightarrow V$ is an isomorphism if one of V_1 and V_2 has the approximation property. Assume that V_2 has the approximation property. Note that

$$\underline{V} = \underline{\text{Hom}}_{\mathbb{R}}(W_1 \otimes_{\mathbb{R}} W_2, \mathbb{R}) = \underline{\text{Hom}}_{\mathbb{R}}(W_1, \underline{\text{Hom}}_{\mathbb{R}}(W_2, \mathbb{R})) = \underline{\text{Hom}}_{\mathbb{R}}(W_1, \underline{V}_2).$$

In the classical literature, this is known as the weak- $*$ -to-weak compact operators from V_1^* to V_2 . On the other hand, $V_1 \otimes_\epsilon V_2 \subset V$ is the closed subspace generated by algebraic tensors $V_1 \otimes V_2$; equivalently, this is the closure of the space of maps $W_1 \rightarrow \underline{V}_2$ of finite rank. The condition that V_2 has the approximation property precisely ensures that this is all of $\underline{\text{Hom}}_{\mathbb{R}}(W_1, \underline{V}_2)$, cf. [DFS08, Theorem 1.3.11]. \square

In other words, when Banach spaces are covariantly embedded into \mathcal{M} -complete condensed \mathbb{R} -spaces, one gets the projective tensor product; while under the duality with Smith spaces, one (essentially) gets the injective tensor product. Thus, in the condensed setting, there is only one tensor product, but it recovers both tensor products on Banach spaces.

Appendix to Lecture IV: Quasiseparated condensed sets

In this appendix, we make some remarks about the category of quasiseparated condensed sets. We start with the following critical observation, giving an interpretation of the topological space $X(*)_{\text{top}}$ purely in condensed terms.

PROPOSITION 4.13. *Let X be a quasiseparated condensed set. Then quasicompact injections $i : Z \hookrightarrow X$ are equivalent to closed subspaces $W \subset X(*)_{\text{top}}$ via $Z \mapsto Z(*)_{\text{top}}$, resp. sending a closed subspace $W \subset X(*)_{\text{top}}$ to the subspace $Z \subset X$ with*

$$Z(S) = X(S) \times_{\text{Map}(S, X(*)_{\text{top}})} \text{Map}(S, W).$$

In the following, we will sometimes refer to quasicompact injections $i : Z \hookrightarrow X$ as closed subspaces of X , noting that by the proposition this agrees with the topological notion.

PROOF. The statement reduces formally to the case that X is quasicompact by writing X as the filtered union of its quasicompact subspaces. In that case, X is equivalent to a compact Hausdorff space. If $i : Z \hookrightarrow X$ is a quasicompact injection, then Z is again quasicompact and quasiseparated, so again (the condensed set associated to) a compact Hausdorff space. But injections of compact Hausdorff spaces are closed immersions. In other words, the statement reduces to the assertion that under the equivalence of quasicompact quasiseparated condensed sets with compact Hausdorff spaces, injections correspond to closed subspaces, which is clear. \square

LEMMA 4.14. *The inclusion of the category of quasiseparated condensed sets into all condensed sets admits a left adjoint $X \mapsto X^{\text{qs}}$, with the unit $X \rightarrow X^{\text{qs}}$ being a surjection of condensed sets. The functor $X \mapsto X^{\text{qs}}$ preserves finite products. In particular, it defines a similar left adjoint for the inclusion of quasiseparated condensed A -modules into all condensed A -modules, for any quasiseparated condensed ring A .*

PROOF. Choose a surjection $X' = \bigsqcup_i S_i \rightarrow X$ from a disjoint union of profinite sets S_i , giving an equivalence relation $R = X' \times_X X' \subset X' \times X'$. For any map $X \rightarrow Y$ with Y quasiseparated, the induced map $X' \rightarrow Y$ has the property that $X' \times_Y X' \subset X' \times X'$ is a quasicompact injection (i.e., is a closed subspace) and is an equivalence relation that contains R ; thus, it contains the minimal closed equivalence relation $\overline{R} \subset X' \times X'$ generated by R (which exists, as any intersection of closed equivalence relations is again a closed equivalence relation). This shows that $X^{\text{qs}} = X'/\overline{R}$ defines the desired adjoint.

To check that it preserves finite products, we need to check that if $R \subset X \times X$ and $R' \subset X' \times X'$ are two equivalence relations on quasiseparated condensed sets X, X' , then the minimal closed equivalence relation $\overline{R \times R'}$ on $X \times X'$ containing $R \times R'$ is given by $\overline{R} \times \overline{R'}$. To see this, note first that for fixed $x' \in X'$, it must contain $\overline{R} \times (x', x') \subset X \times X \times X' \times X'$. Similarly, for fixed $x \in X$, it must contain $(x, x) \times \overline{R'}$. But now if (x_1, x'_1) and (x_2, x'_2) are two elements of $X \times X'$ such that x_1 is \overline{R} -equivalent to x_2 and x'_1 is $\overline{R'}$ -equivalent to x'_2 , then (x_1, x'_1) is $\overline{R \times R'}$ -equivalent to (x_2, x'_1) , which is $\overline{R \times R'}$ -equivalent to (x_2, x'_2) . Thus, $\overline{R \times R'} \subset \overline{R} \times \overline{R'}$, and the reverse inclusion is clear. \square

COROLLARY 4.15. *The inclusion of the category of \mathcal{M} -complete condensed \mathbb{R} -vector spaces into all condensed \mathbb{R} -vector spaces admits a left adjoint, the “ \mathcal{M} -completion”.*

PROOF. Let V be any condensed \mathbb{R} -vector space, and pick a resolution

$$\bigoplus_j \mathbb{R}[S'_j] \rightarrow \bigoplus_i \mathbb{R}[S_i] \rightarrow V \rightarrow 0.$$

The left adjoint exists for $\mathbb{R}[S]$, with S profinite, and takes the value $\mathcal{M}(S)$, essentially by definition of \mathcal{M} -completeness. It follows that the left adjoint for V is given by the quasiseparation of the cokernel of

$$\bigoplus_j \mathcal{M}(S'_j) \rightarrow \bigoplus_i \mathcal{M}(S_i). \quad \square$$

5. Lecture V: Entropy and a real B_{dR}^+

in memoriam Jean-Marc Fontaine

The discussion of the last lecture along with the definition of solid abelian groups begs the following question:

QUESTION 5.1. Does the category of condensed \mathbb{R} -vector spaces V such that for all profinite sets S with a map $f : S \rightarrow V$, there is a unique extension to a map $\tilde{f} : \mathcal{M}(S) \rightarrow V$ of condensed \mathbb{R} -vector spaces, form an abelian category stable under all kernels, cokernels, and extensions?

We note that stability under kernels is easy to see. We will see that stability under cokernels and stability under extensions fails, so the answer is no.

Note that the condition imposed is some form of local convexity. It is a known result in Banach space theory that there are extensions of Banach spaces that are not themselves Banach spaces, and in fact not locally convex. Let us recall the construction, due to Ribe, [Rib79].

Let $V = \ell^1(\mathbb{N})$ be the Banach space of ℓ^1 -sequences of real numbers x_0, x_1, \dots . We will construct a non-split extension

$$0 \rightarrow \mathbb{R} \rightarrow V' \rightarrow V \rightarrow 0.$$

The construction is based on the following two lemmas.

LEMMA 5.2. *Let V be a Banach space, let $V_0 \subset V$ be a dense subvector space and let $\phi : V_0 \rightarrow \mathbb{R}$ be a function that is almost linear in the sense that for some constant C , we have for all $v, w \in V_0$*

$$|\phi(v+w) - \phi(v) - \phi(w)| \leq C(\|v\| + \|w\|);$$

moreover, $\phi(av) = a\phi(v)$ for all $a \in \mathbb{R}$ and $v \in V_0$. Then one can turn the abstract \mathbb{R} -vector space $V'_0 = V_0 \times \mathbb{R}$ into a topological vector space by declaring a system of open neighborhoods of 0 to be

$$\{(v, r) \mid \|v\| + |r - \phi(v)| < \epsilon\}.$$

The completion V' of V'_0 defines an extension

$$0 \rightarrow \mathbb{R} \rightarrow V' \rightarrow V \rightarrow 0$$

of topological vector spaces (which stays exact as a sequence of condensed vector spaces).

This extension of topological vector spaces is split if and only if there is a linear function $f : V_0 \rightarrow \mathbb{R}$ such that $|f(v) - \phi(v)| \leq C'\|v\|$ for all $v \in V_0$ and some constant C' .

One can in fact show that all extensions arise in this way.

PROOF. All statements are immediately verified. Note that the extension $V' \rightarrow V$ splits as topological spaces: Before completion we have the nonlinear splitting $V_0 \rightarrow V'_0 : v \mapsto (v, \phi(v))$, and this extends to completions. \square

We see that non-split extensions are related to functions that are almost linear locally, but not almost linear globally. An example is given by entropy. (We note that a relation between entropy and real analogues of p -adic Hodge-theoretic rings has been first proposed by Connes and Consani, cf. e.g. [CC15], [Con11]. There is some relation between the constructions of this and the next lecture, and their work.) Recall that if p_1, \dots, p_n are real numbers in $[0, 1]$ with sum 1 (considered as a probability distribution on the finite set $\{1, \dots, n\}$), its entropy is

$$H = - \sum_{i=1}^n p_i \log p_i.$$

The required local almost linearity comes from the following lemma.

LEMMA 5.3. *For all real numbers s and t , one has*

$$|s \log |s| + t \log |t| - (s + t) \log |s + t|| \leq 2 \log 2 (|s| + |t|).$$

We note that $0 \log 0 := 0$ extends the function $s \mapsto s \log |s|$ continuously to 0.

PROOF. Rescaling both s and t by a positive scalar λ multiplies the left-hand side by λ (Check!). We may thus assume that $s, t \in [-1, 1]$ and at least one of them has absolute value equal to 1. Changing sign and permuting, we assume that $t = 1$. It then suffices to see that the left-hand side is bounded by $2 \log 2$ for all $s \in [-1, 1]$. Note that some bound is now clear, as the left-hand side is continuous; to get $2 \log 2$, note that $s \log |s|$ and $(s + 1) \log |s + 1|$ take opposite signs for $s \in [-1, 1]$, so it suffices to bound both individually by $2 \log 2$, which is easy. \square

COROLLARY 5.4. *Let $V_0 \subset V = \ell^1(\mathbb{N})$ be the subspace spanned by sequences with finitely many nonzero terms. The function*

$$H : (x_0, x_1, \dots) \in V_0 \mapsto s \log |s| - \sum_{i \geq 0} x_i \log |x_i|, \text{ where } s = \sum_{i \geq 0} x_i,$$

is locally almost linear but not globally almost linear, and so defines a nonsplit extension

$$0 \rightarrow \mathbb{R} \rightarrow V' \rightarrow V \rightarrow 0.$$

PROOF. Local almost linearity follows from the lemma (the scaling invariance $H(ax) = aH(x)$ uses the addition of $s \log |s|$). For global non-almost linearity, assume that H was close to $\sum \lambda_i x_i$ for certain $\lambda_i \in \mathbb{R}$. Looking at the points $(0, \dots, 0, 1, 0, \dots)$, one sees that the λ_i are bounded (as $H = 0$ on such points). On the other hand, looking at $(\frac{1}{n}, \dots, \frac{1}{n}, 0, \dots)$ with n occurrences of $\frac{1}{n}$, global almost linearity requires

$$|H(n) - \frac{1}{n} \sum_{i=0}^{n-1} \lambda_i| \leq C$$

for some constant C . This would require $H(n)$ to be bounded (as the λ_i are), but one computes $H(n) = \log n$. \square

Now we translate this extension into the condensed picture. Ideally, we would like to show that there is an extension

$$0 \rightarrow \mathcal{M}(S) \rightarrow \widetilde{\mathcal{M}}(S) \rightarrow \mathcal{M}(S) \rightarrow 0$$

of Smith spaces, functorial in the profinite set S . Taking $S = \mathbb{N} \cup \{\infty\}$ and writing $\mathcal{M}(S) = W_1 \oplus \mathbb{R} \cdot [\infty]$ by splitting off ∞ , the space W_1 is a Smith space containing $\ell^1(\mathbb{N})$ as a subspace with the same underlying \mathbb{R} -vector space (a compact convex generating set of W is the space of sequences $x_0, x_1, \dots \in [-1, 1]$ with $\sum |x_i| \leq 1$). Then we get an extension

$$0 \rightarrow W \rightarrow \widetilde{W} \rightarrow W \rightarrow 0$$

and we can take the pullback along $\ell^1 \rightarrow W$ and the pushout along $W \rightarrow \mathbb{R}$ (summing all x_i)⁹ to get an extension

$$0 \rightarrow \mathbb{R} \rightarrow ? \rightarrow \ell^1 \rightarrow 0.$$

⁹Warning: This map is not well-defined, but after pullback to ℓ^1 the extension can be reduced to a self-extension of $\ell^1 \subset W$ by itself, and there is a well-defined map $\ell^1 \rightarrow \mathbb{R}$ along which one can the pushout.

This will be the extension constructed above.

We would like to make the following definition.¹⁰

DEFINITION 5.5 (does not work). For a finite set S , let

$$\widetilde{\mathbb{R}[S]}_{\leq c} = \{(x_s, y_s) \in \mathbb{R}[S] \times \mathbb{R}[S] \mid \sum_{s \in S} (|x_s| + |y_s - x_s \log |x_s||) \leq c\}.$$

For a profinite set $S = \varprojlim_i S_i$, let

$$\widetilde{\mathcal{M}(S)} = \bigcup_{c>0} \varprojlim_i \widetilde{\mathbb{R}[S]}_{\leq c}.$$

The problem with this definition is that the transition maps in the limit over i do not preserve the subspaces $\leq c$. It would be enough if there was some universal constant C such that for any map of finite set $S \rightarrow T$, the $\widetilde{\mathbb{R}[S]}_{\leq c}$ maps into $\widetilde{\mathbb{R}[T]}_{\leq Cc}$. Unfortunately, even this is not true. We will see in the next lecture that replacing the ℓ^1 -norm implicit in the definition of $\widetilde{\mathbb{R}[S]}_{\leq c}$ with the ℓ^p -norm for some $p < 1$, this problem disappears.

One can salvage the definition for the Smith space W_1 (the direct summand of $\mathcal{M}(\mathbb{N} \cup \{\infty\})$). In this case, one simply directly builds the extension \widetilde{W}_1 , as follows:

$$\widetilde{W}_1 = \bigcup_{c>0} \{(x_0, x_1, \dots, y_0, y_1, \dots) \in \prod_{\mathbb{N}} [-c, c] \times \prod_{\mathbb{N}} [-c, c] \mid \sum_n (|x_n| + |y_n - x_n \log |x_n||) \leq c\}.$$

PROPOSITION 5.6. *The condensed set \widetilde{W}_1 has a natural structure of a condensed \mathbb{R} -vector space, and sits in an exact sequence*

$$0 \rightarrow W_1 \rightarrow \widetilde{W}_1 \rightarrow W_1 \rightarrow 0.$$

PROOF. Surjectivity of $\widetilde{W}_1 \rightarrow W_1$ is clear by taking $y_n = x_n \log |x_n|$. To see that it is a condensed \mathbb{R} -vector space, note that stability under addition follows from Lemma 5.3, and one similarly checks stability under scalar multiplication. \square

EXERCISE 5.7. Show that the extension $0 \rightarrow W_1 \rightarrow \widetilde{W}_1 \rightarrow W_1 \rightarrow 0$ has a Banach analogue: An extension $0 \rightarrow \ell^1 \rightarrow \widetilde{\ell}^1 \rightarrow \ell^1 \rightarrow 0$, sitting inside the previous sequence.¹¹

The proposition shows that the class of \mathcal{M} -complete condensed \mathbb{R} -vector spaces is not stable under extensions. Indeed, if \widetilde{W}_1 were \mathcal{M} -complete, then the map $\mathbb{N} \cup \{\infty\} \rightarrow \widetilde{W}_1$ given by sending n to the element with $x_n = 1$ and all other $x_i, y_i = 0$ would extend to a map $\mathcal{M}(\mathbb{N} \cup \{\infty\}) \rightarrow \widetilde{W}_1$ vanishing on ∞ , thus giving a section $W_1 \rightarrow \widetilde{W}_1$; but the extension is nonsplit.

We can now also show that cokernels are not well-behaved. Indeed, consider also the Smith space version W_∞ of ℓ^∞ , given by $\bigcup_{c>0} \prod_{\mathbb{N}} [-c, c]$. There is a natural inclusion $W_1 \subset W_\infty$.

PROPOSITION 5.8. *The map of condensed sets $f : W_1 \rightarrow W_\infty$ given by*

$$(x_0, x_1, \dots) \mapsto (x_0 \log |x_0|, x_1 \log |x_1|, \dots)$$

induces, via projection $W_\infty \rightarrow W_\infty/W_1$, a nonzero map of condensed \mathbb{R} -vector spaces

$$W_1 \rightarrow W_\infty/W_1.$$

¹⁰This was erroneously stated in the lecture.

¹¹See [KP79].

PROOF. Let us check first that it is a map of condensed abelian groups. For this, we need to see that two maps (of condensed sets) $W_1 \times W_1 \rightarrow W_\infty/W_1$ agree. Their difference is the projection along $W_\infty \rightarrow W_\infty/W_1$ of the map $W_1 \times W_1 \rightarrow W_\infty$ sending pairs of (x_0, x_1, \dots) and (y_0, y_1, \dots) to $(x_0 \log |x_0| + y_0 \log |y_0| - (x_0 + y_0) \log |x_0 + y_0|, x_1 \log |x_1| + y_1 \log |y_1| - (x_1 + y_1) \log |x_1 + y_1|, \dots)$, which lies in W_1 by Lemma 5.3. This gives additivity. One checks \mathbb{R} -linearity similarly.

To see that it is nonzero, it suffices to see that the image of W_1 under f is not contained in the subspace $W_1 \subset W_\infty$. But the sequence $z_n = (\frac{1}{n}, \dots, \frac{1}{n}, 0, \dots)$ for varying n (with n occurrences of $\frac{1}{n}$) defines a map $\mathbb{N} \cup \{\infty\} \rightarrow W_1$, whose image under f is the sequence $(-\frac{\log n}{n}, \dots, -\frac{\log n}{n}, 0, \dots)$ which does not have bounded ℓ^1 -norm. \square

Using this map, one can recover the extension $0 \rightarrow W_1 \rightarrow \widetilde{W}_1 \rightarrow W_1 \rightarrow 0$ as the pullback of the canonical extension $0 \rightarrow W_1 \rightarrow W_\infty \rightarrow W_\infty/W_1 \rightarrow 0$.

Moreover, this finally answers the question at the beginning of this lecture: The map $\mathcal{M}(\mathbb{N} \cup \{\infty\}) \rightarrow W_1 \rightarrow W_\infty/W_1$ is zero when restricted to $\mathbb{N} \cup \{\infty\}$, so the cokernel W_∞/W_1 of a map of Smith spaces does not satisfy the property that any map $S \rightarrow W_\infty/W_1$ extends uniquely to a map $\mathcal{M}(S) \rightarrow W_\infty/W_1$.

As a final topic of this lecture, we construct a kind of “real analogue of B_{dR}^+ ”. We note that the extension \widetilde{W}_1 is in fact a flat condensed $\mathbb{R}[\epsilon]$ -module, where $\mathbb{R}[\epsilon] = \mathbb{R}[t]/t^2$, via letting ϵ map sequences $(x_0, x_1, \dots, y_0, y_1, \dots)$ to $(0, 0, \dots, x_0, x_1, \dots)$.

One may wonder whether one can build an infinite self-extension of W_1 's, leading to a flat condensed $\mathbb{R}[[t]]$ -module $W_{1, \mathbb{R}[[t]]}$ with $W_{1, \mathbb{R}[[t]]} \otimes_{\mathbb{R}[[t]]} \mathbb{R} = W_1$. Ideally, one would want to have a similar self-extension of all $\mathcal{M}(S)$, functorial in the profinite set S ; as before, this does not actually work, but will work in an analogous setting as studied in the next lecture.

For the construction, consider the multiplicative map

$$\mathbb{R} \rightarrow \mathbb{R}[[t]]$$

sending $x \in \mathbb{R}$ to the series

$$[x] := x|x|^t = x + x \log |x|t + \frac{1}{2}x \log^2 |x|t^2 + \dots + \frac{1}{n!}x \log^n |x|t^n + \dots$$

Any element of $\mathbb{R}[t]/t^n$ can be written in a unique way in the form

$$[x_0] + [x_1]t + \dots + [x_{n-1}]t^{n-1}$$

with $x_i \in \mathbb{R}$. For $n = 2$, this writes $a + bt$ in the form $[a] + [b - a \log |a|]t$. For $n = 3$, the formula is already rather complicated:

$$a + bt + ct^2 = [a] + [b - a \log |a|]t + [c - (b - a \log |a|) \log |b - a \log |a|| - \frac{1}{2}a \log^2 |a|]t^2.$$

Thus much complexity is hidden in this isomorphism.

LEMMA 5.9. *For any finite set S and real number $c > 0$, let*

$$(\mathbb{R}[t]/t^n)[S]_{\leq c} = \left\{ \left(\sum_{i=0}^{n-1} [x_{i,s}]t^i \right)_s \in \mathbb{R}[t]/t^n[S] \mid \sum_{i,s} |x_{i,s}| \leq c \right\}.$$

Then $(\mathbb{R}[t]/t^n)[S]_{\leq c}$ is stable under multiplication by $[-\epsilon, \epsilon]$ for some $\epsilon = \epsilon(n)$, and there is a constant $C = C(n)$ such that

$$(\mathbb{R}[t]/t^n)[S]_{\leq c} + (\mathbb{R}[t]/t^n)[S]_{\leq c} \subset (\mathbb{R}[t]/t^n)[S]_{\leq Cc}.$$

PROOF. Let us explain the case of addition; we leave the case of scalar multiplication as an exercise. In the given coordinates, there are functions S_0, \dots, S_{n-1} , where S_i is a continuous function of $x_0, \dots, x_i, y_0, \dots, y_i$, such that

$$[x_0] + [x_1]t + \dots + [x_{n-1}]t^{n-1} + [y_0] + [y_1]t + \dots + [y_{n-1}]t^{n-1} = [S_0] + [S_1]t + \dots + [S_{n-1}]t^{n-1} \in \mathbb{R}[t]/t^n.$$

Using that multiplication by $[\lambda]$ for any $\lambda \in \mathbb{R}$ commutes with all operations, one sees that

$$S_i(\lambda x_0, \dots, \lambda x_i, \lambda y_0, \dots, \lambda y_i) = \lambda S_i(x_0, \dots, x_i, y_0, \dots, y_i).$$

This implies that

$$S_i(x_0, \dots, x_i, y_0, \dots, y_i) \leq C_i(|x_0| + \dots + |x_i| + |y_0| + \dots + |y_i|)$$

by taking $C_i > 0$ to be the maximum absolute value taken by S_i on the compact set where all $x_j, y_j \in [-1, 1]$ and at least one of them is ± 1 . This estimate easily implies the desired stability under addition. \square

Ideally, one would like to use the lemma to see that for any profinite set $S = \varprojlim_i S_i$, one can define a condensed $\mathbb{R}[t]/t^n$ -module

$$\mathcal{M}(S, \mathbb{R}[t]/t^n) := \bigcup_{c>0} \varprojlim_i (\mathbb{R}[t]/t^n)[S_i]_{\leq c}.$$

However, as before this construction does not work because the transition maps do not preserve the desired bounds. However, one can build the corresponding infinite self-extension of W_1 .¹²

¹²We believe this is related to the “complex interpolation of Banach spaces” as studied for example in [CCR⁺82], [CSCK15].

6. Lecture VI: Statement of main result

In the last lecture, we saw that a naive adaptation of solid abelian groups to the case of the real numbers, using the spaces $\mathcal{M}(S)$ of signed Radon measures, does not work; but its failure gives rise to interesting phenomena.

In this lecture, however, we want to find a variant that does work. We want to find a full subcategory of condensed \mathbb{R} -vector spaces that is abelian, stable under kernels, cokernels, and extensions, and contains all \mathcal{M} -complete objects (but is still enforcing a nontrivial condition). This forces us to include also all the extensions constructed in the previous lecture. These are not \mathcal{M} -complete; in the setting of extensions of Banach spaces, these extensions are not locally convex anymore.

The following weakening of convexity has been studied in the Banach space literature.

DEFINITION 6.1. For $0 < p \leq 1$, a p -Banach space is a topological \mathbb{R} -vector space V such that there exists a p -norm, i.e. a continuous map

$$\|\cdot\| : V \rightarrow \mathbb{R}_{\geq 0}$$

with the following properties:

- (1) For any $v \in V$, the norm $\|v\| = 0$ if and only if $v = 0$;
- (2) For all $v \in V$ and $a \in \mathbb{R}$, one has $\|av\| = |a|^p \|v\|$;
- (3) For all $v, w \in V$, one has $\|v + w\| \leq \|v\| + \|w\|$;
- (4) The sets $\{v \in V \mid \|v\| < \epsilon\}$ for varying $\epsilon \in \mathbb{R}_{>0}$ define a basis of open neighborhoods of 0;
- (5) For any sequence $v_0, v_1, \dots \in V$ with $\|v_i - v_j\| \rightarrow 0$ as $i, j \rightarrow \infty$, there exists a (necessarily unique) $v \in V$ with $\|v - v_i\| \rightarrow 0$.

Thus, the only difference is in the scaling behaviour. We note that V is a p -Banach then it is a p' -Banach for all $p' \geq p$, as if $\|\cdot\|$ is a p -norm, then $\|\cdot\|^{p'/p}$ is a p' -norm. A quasi-Banach space is a topological \mathbb{R} -vector space that is a p -Banach for some $p > 0$.

A key result is the following.

THEOREM 6.2 ([Kal81]). *Any extension of p -Banach spaces is a p' -Banach for all $p' < p$.*

This suggests that the extension problems we encountered disappear if instead of convexity, we ask for p -convexity¹³ for all $p < 1$. Here p -convexity is the assertion that for all $a_1, \dots, a_n \in \mathbb{R}$ with $|a_1|^p + \dots + |a_n|^p \leq 1$, the ball $\{v \in V \mid \|v\| < \epsilon\}$ is stable under $(v_1, \dots, v_n) \mapsto a_1 v_1 + \dots + a_n v_n$.

Let us do the obvious translation into the condensed world.

DEFINITION 6.3. Let $0 < p \leq 1$ be a real number. For any finite set S and real number $c > 0$, let

$$\mathbb{R}[S]_{\ell^p \leq c} = \{(a_s)_s \in \mathbb{R}[S] \mid \sum_{s \in S} |a_s|^p \leq c\},$$

and for a profinite set $S = \varprojlim_i S_i$, let

$$\mathcal{M}_p(S) = \bigcup_{c>0} \varprojlim_i \mathbb{R}[S_i]_{\ell^p \leq c}.$$

¹³Note that p -convexity for $p < 1$ is weaker than convexity, and might also be called concavity.

For any fixed value of p , we run into the same problem with non-split self-extensions of $\mathcal{M}_p(S)$ as in the last lecture, except that for $p < 1$ things are better-behaved, in that one can define these self-extensions functorially in the profinite set S : For any finite set S , and $n \geq 1$, one can define

$$(\mathbb{R}[T]/T^n)[S]_{\ell^p \leq c} := \left\{ \left(\sum_{i=0}^{n-1} [x_{i,s}] T^i \right)_s \mid \sum_{i,s} |x_{i,s}|^p \leq c \right\}.$$

Then there is a constant $C = C(n, p) < \infty$ such that for any map of finite sets $S \rightarrow S'$, the image of $(\mathbb{R}[T]/T^n)[S]_{\ell^p \leq c}$ is contained in $(\mathbb{R}[T]/T^n)[S']_{\ell^p \leq Cc}$, by Proposition 6.4 below. In particular, the subset $(\mathbb{R}[T]/T^n)[S]_{\ell^p \leq c, \text{stable}} \subset (\mathbb{R}[T]/T^n)[S]_{\ell^p \leq c}$ of all elements whose images under any map $S \rightarrow S'$ are still in the $\ell^p \leq c$ -subspace, is cofinal with the full space, and now is compatible with transition maps. In other words, for a profinite set $S = \varprojlim_i S_i$, we can now functorially define

$$\mathcal{M}_p(S, \mathbb{R}[T]/T^n) = \bigcup_{c>0} \varprojlim_i (\mathbb{R}[T]/T^n)[S_i]_{\ell^p \leq c, \text{stable}}$$

and thus, in the limit over n , define a B_{dR}^+ -theory for any $p < 1$. As we will not have direct use for it (and, just like \mathcal{M}_p itself, it does not define an abelian category of modules), we will not go into more details about this. We will just note that “the” real numbers are not well-defined anymore: To define a category of modules over \mathbb{R} , one needs to specify in addition $0 < p < 1$; and in some sense an infinitesimal deformation of p gives rise to B_{dR}^+ , so there is in some sense a 1-parameter family of versions of the real numbers. We will make this picture more precise later.

We used the following proposition:

PROPOSITION 6.4. *There is a constant $C = C(n, p) < \infty$ such that for all maps $f : S \rightarrow S'$ of finite sets,*

$$f((\mathbb{R}[T]/T^n)[S]_{\ell^p \leq c}) \subset (\mathbb{R}[T]/T^n)[S']_{\ell^p \leq Cc}.$$

PROOF. Decomposing into fibres over S' , we can assume that S' is a point. Note that multiplication by $[\lambda]$ for $\lambda > 0$ maps the $\ell^p \leq c$ -subspace isomorphically to the $\ell^p \leq \lambda^p c$ -subspace. Thus, by rescaling, we can assume that $c = 1$. It remains to see that there is some bounded subset $B \subset \mathbb{R}[T]/T^n$ such that for all finite sets S and all $(\sum_{i=0}^{n-1} [x_{i,s}] T^i)_s \in (\mathbb{R}[T]/T^n)[S]_{\ell^p \leq 1}$, one has

$$\sum_{s \in S} \sum_{i=0}^{n-1} [x_{i,s}] T^i \in B \subset \mathbb{R}[T]/T^n.$$

By induction, such a statement holds true for the sum over $i = 1, \dots, n-1$. Thus, it suffices to see that for all integers $m = |S|$ and all x_1, \dots, x_m with $\sum_{j=1}^m |x_j|^p \leq 1$, one has

$$\sum_{j=1}^m [x_j] \in B' \subset \mathbb{R}[T]/T^n$$

for some bounded subset B' . In other words, we need to see that for all $i = 0, \dots, n-1$, one has

$$\sum_{j=1}^m x_j \log^i |x_j| \leq C_i$$

for some constant C_i . But for all sufficiently small x , one has $\log^i |x| \leq |x|^{p-1}$, so neglecting large x_j (which can only give a bounded contribution), one has $\sum_{j=1}^m x_j \log^i |x_j| \leq \sum_{j=1}^m |x_j|^p \leq 1$. \square

Without further ado, let us now state the first main theorem of this course. Define

$$\mathcal{M}_{<p}(S) = \varinjlim_{p' < p} \mathcal{M}_{p'}(S).$$

THEOREM 6.5. *Fix any $0 < p \leq 1$. For a condensed \mathbb{R} -vector space V , the following conditions are equivalent.*

- (1) *For all $p' < p$ and all maps $f : S \rightarrow V$ from a profinite set S , there is a unique extension to a map*

$$\tilde{f}_{p'} : \mathcal{M}_{p'}(S) \rightarrow V$$

of condensed \mathbb{R} -vector spaces.

- (2) *For all maps $f : S \rightarrow V$ from a profinite set S , there is a unique extension to a map*

$$\tilde{f}_{<p} : \mathcal{M}_{<p}(S) \rightarrow V$$

of condensed \mathbb{R} -vector spaces.

- (3) *One can write V as the cokernel of a map*

$$\bigoplus_i \mathcal{M}_{<p}(S_i) \rightarrow \bigoplus_j \mathcal{M}_{<p}(S'_j).$$

If V satisfies these conditions, we say that V is p -liquid. The class $\text{Liq}_p(\mathbb{R})$ of p -liquid \mathbb{R} -vector spaces is an abelian subcategory stable under all kernels, cokernels, and extensions. It is generated by the compact projective objects $\mathcal{M}_{<p}(S)$, S extremally disconnected. Moreover:

- (1) *The inclusion $\text{Liq}_p(\mathbb{R}) \rightarrow \text{Cond}(\mathbb{R})$ has a left adjoint $V \mapsto V_p^{\text{liq}}$, and there is a (necessarily unique) symmetric monoidal tensor product \otimes_p^{liq} on $\text{Liq}_p(\mathbb{R})$ making $V \mapsto V_p^{\text{liq}}$ symmetric monoidal.*
- (2) *The functor $D(\text{Liq}_p(\mathbb{R})) \rightarrow D(\text{Cond}(\mathbb{R}))$ is fully faithful, and admits a left adjoint, which is the left derived functor $C \mapsto C_p^{\text{Lliq}}$ of $V \mapsto V_p^{\text{liq}}$. There is a unique symmetric monoidal tensor product \otimes_p^{Lliq} on $D(\text{Liq}_p(\mathbb{R}))$ making $C \mapsto C_p^{\text{Lliq}}$ symmetric monoidal; it is the left derived functor of \otimes_p^{liq} .*
- (3) *An object $C \in D(\text{Cond}(\mathbb{R}))$ lies in $D(\text{Liq}_p(\mathbb{R}))$ if and only if all $H^i(C) \in \text{Liq}_p(\mathbb{R})$; in that case, for all profinite sets S and all $p' < p$, one has*

$$R\text{Hom}_{\mathbb{R}}(\mathcal{M}_{p'}(S), C) = R\text{Hom}_{\mathbb{R}}(\mathbb{R}[S], C).$$

In other words, one gets an extremely well-behaved theory of p -liquid \mathbb{R} -vector spaces. In the language of last semester, the present theory defines an analytic ring [Sch19, Definition 7.4, Proposition 7.5]. In fact, by [Sch19, Lemma 5.9, Remark 5.11], all we need to prove is the following result.

THEOREM 6.6. *Let $0 < p \leq 1$, let*

$$g : \bigoplus_i \mathcal{M}_{<p}(S_i) \rightarrow \bigoplus_j \mathcal{M}_{<p}(S'_j)$$

be a map of condensed \mathbb{R} -vector spaces, where all S_i and S'_j are extremally disconnected, and let V be the kernel of g . Then for any $p' < p$ and any profinite set S , the map

$$R\text{Hom}_{\mathbb{R}}(\mathcal{M}_{p'}(S), V) \rightarrow R\text{Hom}_{\mathbb{R}}(\mathbb{R}[S], V)$$

is a quasi-isomorphism in $D(\text{Cond}(\text{Ab}))$.

Unfortunately, at this point, we have to pay a serious prize for going into the non-locally convex setting. More precisely, in order to compute $R\text{Hom}_{\mathbb{R}}(\mathcal{M}_{p'}(S), V)$, i.e. the complex of \mathbb{R} -linear maps $\mathcal{M}_{p'}(S) \rightarrow V$, we first have to understand maps of condensed sets $\mathcal{M}_{p'}(S) \rightarrow V$, i.e.

$$R\Gamma(\mathcal{M}_{p'}(S), V)$$

for V as above. We may restrict to the compact Hausdorff subspace $\mathcal{M}_{p'}(S)_{\leq 1}$. In order for this to be helpful, we need to be able to compute the cohomology

$$H^i(S, V)$$

for compact Hausdorff S with coefficients in spaces V like above; in particular, for p -Banach spaces.

The proof of [Sch19, Theorem 3.3] gives the following result.

THEOREM 6.7. *Let V be a Banach space and let S be a compact Hausdorff space. Then $H^i(S, \underline{V}) = 0$ for $i > 0$. Similarly, if V is a p -Banach for any $p \leq 1$ and S is a profinite set, then $H^i(S, \underline{V}) = 0$ for $i > 0$.*

However, a key step in the proof is the existence of a partition of unity, the application of which uses that V is locally convex critically. In particular, the first part fails if V is only a p -Banach for $p < 1$.¹⁴

This means that for the key computation, we have to further resolve $\mathcal{M}_{p'}(S)_{\leq 1}$ by profinite sets; and we should do so in a way that keeps the resulting resolution sufficiently explicit. In other words, the condensed formalism now forces us to resolve the real numbers, and vector spaces over it, in terms of profinite sets, explicitly!¹⁵

We are now ready for the key turn. We will generalize the present question about the real numbers to an arithmetic ring that is a countable union of profinite subsets.

Let us define this ring. It depends on a real number $0 < r < 1$. We define the condensed ring $\mathbb{Z}((T))_r$ whose S -valued points are

$$\mathbb{Z}((T))_r(S) = \left\{ \sum_{n \in \mathbb{Z}} a_n T^n \mid a_n \in C(S, \mathbb{Z}), \sum_{n \in \mathbb{Z}} |a_n| r^n < \infty \right\} \subset \mathbb{Z}((T))(S).$$

More precisely, this is the increasing union of the subsets

$$\mathbb{Z}((T))_{r, \leq c}(S) = \left\{ \sum_{n \in \mathbb{Z}} a_n T^n \mid a_n \in C(S, \mathbb{Z}), \sum_{n \in \mathbb{Z}} |a_n| r^n \leq c \right\}$$

for varying reals $c > 0$; and here $\sum_{n \in \mathbb{Z}} |a_n| r^n \leq c$ means that for all $s \in S$, $\sum_{n \in \mathbb{Z}} |a_n(s)| r^n \leq c$.

PROPOSITION 6.8. *The condensed set $\mathbb{Z}((T))_{r, \leq c}$ is a profinite set.*

PROOF. As multiplication by T is an isomorphism between $\mathbb{Z}((T))_{r, \leq c}$ and $\mathbb{Z}((T))_{r, \leq rc}$, we can assume that $c \leq 1$. In that case $\mathbb{Z}((T))_{r, \leq c} \subset \mathbb{Z}[[T]] = \prod_{n \geq 0} \mathbb{Z} \cdot T^n$. Now $\mathbb{Z}((T))_{r, \leq c}$ can be written as the inverse limit of

$$\left\{ \sum_{n=0}^m a_n T^n \mid \sum_{n=0}^m |a_n| r^n \leq c \right\} \subset \prod_{n=0}^m \mathbb{Z} \cdot T^n,$$

¹⁴We believe that if S is finite-dimensional, one can prove the same result; but the relevant compact Hausdorff spaces are highly infinite-dimensional.

¹⁵This is a difficulty that is not seen in the classical quasi-Banach space literature; and so Theorem 6.2 and its techniques of proof cannot easily be extended to the present situation.

each of which is a finite set. □

For any finite set S , we now write the free module $\mathbb{Z}((T))_r[S]$ as the increasing union of

$$\mathbb{Z}((T))_r[S]_{\leq c} := \left\{ \sum_{n \in \mathbb{Z}, s \in S} a_{n,s} T^n [s] \mid a_{n,s} \in C(S, \mathbb{Z}), \sum_{n \in \mathbb{Z}, s \in S} |a_{n,s}| r^n \leq c \right\};$$

each of these is again a profinite set. The addition defines maps

$$\mathbb{Z}((T))_r[S]_{\leq c} \times \mathbb{Z}((T))_r[S]_{\leq c'} \rightarrow \mathbb{Z}((T))_r[S]_{\leq c+c'}$$

and everything is covariantly functorial in S . For any profinite set $S = \varprojlim_i S_i$ we define

$$\mathcal{M}(S, \mathbb{Z}((T))_r)_{\leq c} := \varprojlim_i \mathbb{Z}((T))_r[S_i]_{\leq c}$$

which is a profinite set, functorial in S , equipped with an addition map

$$\mathcal{M}(S, \mathbb{Z}((T))_r)_{\leq c} \times \mathcal{M}(S, \mathbb{Z}((T))_r)_{\leq c'} \rightarrow \mathcal{M}(S, \mathbb{Z}((T))_r)_{\leq c+c'}.$$

In particular, the colimit

$$\mathcal{M}(S, \mathbb{Z}((T))_r) = \bigcup_{c > 0} \mathcal{M}(S, \mathbb{Z}((T))_r)_{\leq c}$$

is a condensed abelian group, and in fact one easily sees that it is a $\mathbb{Z}((T))_r$ -module.

In the next lecture, we will prove the following theorem, relating this theory to the various ℓ^p -theories over \mathbb{R} . Part (1) is due to Harbater, [**Har84**, Lemma 1.5].

THEOREM 6.9. *Let $0 < r' < r$ and consider the map*

$$\theta_{r'} : \mathbb{Z}((T))_r \rightarrow \mathbb{R} : \sum a_n T^n \mapsto \sum a_n (r')^n.$$

This map is surjective. Moreover:

- (1) *The kernel of $\theta_{r'}$ is generated by a nonzerodivisor $f_{r'} \in \mathbb{Z}((T))_r$.*
- (2) *For any profinite set S , there is a canonical isomorphism*

$$\mathcal{M}(S, \mathbb{Z}((T))_r) / (f_{r'}) \cong \mathcal{M}_p(S)$$

of condensed $\mathbb{Z}((T))_r / (f_{r'}) = \mathbb{R}$ -modules, where $0 < p < 1$ is chosen so that $(r')^p = r$.

- (3) *More generally, for any profinite set S and any $n \geq 1$, there is a canonical isomorphism*

$$\mathcal{M}(S, \mathbb{Z}((T))_r) / (f_{r'})^n \cong \mathcal{M}_p(S, \mathbb{R}[X]/X^n)$$

of condensed $\mathbb{Z}((T))_r / (f_{r'})^n \cong \mathbb{R}[X]/X^n$ -modules.

In other words, the theory over $\mathbb{Z}((T))_r$ specializes to all different \mathcal{M}_p -theories for $p < 1$, and using infinitesimal variations one even recovers the B_{dR}^+ -theories for all p . In some sense, one could regard $\mathbb{Z}((T))_r$ as some version of Fontaine's A_{inf} .

On the other hand, this again indicates that for a fixed r , the theory will not work, so we pass to a colimit again. Another twist, necessary for the proof, will be to isolate the desired theory within all condensed $\mathbb{Z}[T^{-1}]$ -modules; this is related to the possibility to isolate p -liquid \mathbb{R} -vector spaces inside all condensed abelian groups (i.e., the forgetful functor is fully faithful).

THEOREM 6.10. *Fix any $0 < r < 1$. For a condensed $\mathbb{Z}[T^{-1}]$ -module M , the following conditions are equivalent.*

- (1) For all $1 > r' > r$ and all maps $f : S \rightarrow M$ from a profinite set S , there is a unique extension to a map

$$\tilde{f}_{r'} : \mathcal{M}(S, \mathbb{Z}((T))_{r'}) \rightarrow M$$

of condensed $\mathbb{Z}[T^{-1}]$ -modules.

- (2) For all maps $f : S \rightarrow M$ from a profinite set S , there is a unique extension to a map

$$\tilde{f}_{>r} : \mathcal{M}(S, \mathbb{Z}((T))_{>r}) \rightarrow M$$

of condensed $\mathbb{Z}[T^{-1}]$ -modules.

- (3) One can write M as the cokernel of a map

$$\bigoplus_i \mathcal{M}(S_i, \mathbb{Z}((T))_{>r}) \rightarrow \bigoplus_j \mathcal{M}(S'_j, \mathbb{Z}((T))_{>r}).$$

The class $\text{Liq}_r(\mathbb{Z}[T^{-1}])$ of such condensed $\mathbb{Z}[T^{-1}]$ -modules is an abelian subcategory stable under all kernels, cokernels, and extensions. It is generated by the compact projective objects $\mathcal{M}(S, \mathbb{Z}((T))_{>r})$, S extremally disconnected. Moreover:

- (1) The inclusion $\text{Liq}_r(\mathbb{Z}[T^{-1}]) \rightarrow \text{Cond}(\mathbb{Z}[T^{-1}])$ has a left adjoint $M \mapsto M_r^{\text{liq}}$, and there is a (necessarily unique) symmetric monoidal tensor product \otimes_r^{liq} on $\text{Liq}_r(\mathbb{Z}[T^{-1}])$ making $M \mapsto M_r^{\text{liq}}$ symmetric monoidal.
- (2) The functor $D(\text{Liq}_r(\mathbb{Z}[T^{-1}])) \rightarrow D(\text{Cond}(\mathbb{Z}[T^{-1}]))$ is fully faithful, and admits a left adjoint, which is the left derived functor $C \mapsto C_r^{\text{Lliq}}$ of $M \mapsto M_r^{\text{liq}}$. There is a unique symmetric monoidal tensor product \otimes_r^{Lliq} on $D(\text{Liq}_r(\mathbb{Z}[T^{-1}]))$ making $C \mapsto C_r^{\text{Lliq}}$ symmetric monoidal; it is the left derived functor of \otimes_r^{liq} .
- (3) An object $C \in D(\text{Cond}(\mathbb{Z}[T^{-1}]))$ lies in $D(\text{Liq}_r(\mathbb{Z}[T^{-1}]))$ if and only if all $H^i(C) \in \text{Liq}_r(\mathbb{Z}[T^{-1}])$; in that case, for all profinite sets S and all $1 > r' > r$, one has

$$R\text{Hom}_{\mathbb{Z}[T^{-1}]}(\mathcal{M}(S, \mathbb{Z}((T))_{r'}), C) = R\text{Hom}_{\mathbb{Z}[T^{-1}]}(\mathbb{Z}[T^{-1}][S], C).$$

It follows formally that $\text{Liq}_r(\mathbb{Z}[T^{-1}])$ also sits fully faithfully in $\text{Cond}(\mathbb{Z}((T))_{>r})$ (note that the tensor unit of $\text{Liq}_r(\mathbb{Z}[T^{-1}])$ is $\mathbb{Z}((T))_{>r}$, and that everything holds true with $\mathbb{Z}[T^{-1}]$ replaced by $\mathbb{Z}((T))_{>r}$; we will then also write $\text{Liq}_r(\mathbb{Z}((T))_{>r})$ or simply $\text{Liq}(\mathbb{Z}((T))_{>r})$ for $\text{Liq}_r(\mathbb{Z}[T^{-1}])$. However, the proof will require us to prove the finer statement with base ring $\mathbb{Z}[T^{-1}]$.

Again, it reduces formally to the following assertion.

THEOREM 6.11. *Let K be a condensed $\mathbb{Z}[T^{-1}]$ -module that is the kernel of some map*

$$f : \bigoplus_{i \in I} \mathcal{M}(S_i, \mathbb{Z}((T))_{>r}) \rightarrow \bigoplus_{j \in J} \mathcal{M}(S'_j, \mathbb{Z}((T))_{>r})$$

where all S_i and S'_j are extremally disconnected. Then for all $1 > r' > r$ and all profinite sets S , the map

$$R\text{Hom}_{\mathbb{Z}[T^{-1}]}(\mathcal{M}(S, \mathbb{Z}((T))_{r'}), K) \rightarrow R\text{Hom}_{\mathbb{Z}[T^{-1}]}(\mathbb{Z}[T^{-1}][S], K)$$

in $D(\text{Cond}(\text{Ab}))$ is an isomorphism.

In the next lecture, we will show that $\mathbb{Z}((T))_{>r}$ is a principal ideal domain, and prove Theorem 6.9. In the appendix to this lecture, we explain how the rest of the assertions of this lecture reduce to Theorem 6.11.

Appendix to Lecture VI: Recollections on analytic rings

In [Sch19, Lecture VII], we defined analytic rings, abstracting the examples we have seen.

DEFINITION 6.12 ([Sch19, Definition 7.1, 7.4]). A pre-analytic ring is a condensed ring $\underline{\mathcal{A}}$ together with a functor

$$\{\text{extremally disconnected sets}\} \rightarrow \text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}} : S \mapsto \mathcal{A}[S]$$

to the category of $\underline{\mathcal{A}}$ -modules in condensed abelian groups, taking finite disjoint unions to finite products, together with a natural transformation $S \rightarrow \mathcal{A}[S]$.

An analytic ring is a pre-analytic ring \mathcal{A} such that for all index sets I, J and extremally disconnected sets S_i, S'_j , and all maps

$$f : \bigoplus_i \mathcal{A}[S_i] \rightarrow \bigoplus_j \mathcal{A}[S'_j]$$

of condensed $\underline{\mathcal{A}}$ -modules with kernel K , the map

$$R\text{Hom}_{\underline{\mathcal{A}}}(\mathcal{A}[S], K) \rightarrow R\text{Hom}_{\underline{\mathcal{A}}}(\mathcal{A}[S], K)$$

is an isomorphism in $D(\text{Cond}(\text{Ab}))$, for all extremally disconnected sets S .

REMARK 6.13. The definition of analytic rings was stated in a slightly different, but equivalent way: the current condition implies the one in [Sch19, Definition 7.4] by writing C as the limit of its Postnikov truncations; conversely, use the argument of [Sch19, Lemma 5.10].

Thus, Theorem 6.11 implies that $\underline{\mathcal{A}} = \mathbb{Z}[T^{-1}]$ with $\mathcal{A}[S] = \mathcal{M}(S, \mathbb{Z}((T))_{>r})$ is an analytic ring. (It says slightly more as we allow S profinite there, and make a claim about all $r' < r$; we will explain what extra information this gives.) This immediately gives a lot of information:

PROPOSITION 6.14 ([Sch19, Proposition 7.5]). *Let \mathcal{A} be an analytic ring.*

(i) *The full subcategory*

$$\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}} \subset \text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}}$$

of all $\underline{\mathcal{A}}$ -modules M in condensed abelian groups such that for all extremally disconnected sets S , the map

$$\text{Hom}_{\underline{\mathcal{A}}}(\mathcal{A}[S], M) \rightarrow M(S)$$

is an isomorphism, is an abelian category stable under all limits, colimits, and extensions. The objects $\mathcal{A}[S]$ for S extremally disconnected form a family of compact projective generators. It admits a left adjoint

$$\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}} \rightarrow \text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}} : M \mapsto M \otimes_{\underline{\mathcal{A}}} \mathcal{A}$$

that is the unique colimit-preserving extension of $\underline{\mathcal{A}}[S] \mapsto \mathcal{A}[S]$. If $\underline{\mathcal{A}}$ is commutative, there is a unique symmetric monoidal tensor product $\otimes_{\underline{\mathcal{A}}}$ on $\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}}$ making the functor

$$\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}} \rightarrow \text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}} : M \mapsto M \otimes_{\underline{\mathcal{A}}} \mathcal{A}$$

symmetric monoidal.

(ii) *The functor*

$$D(\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}}) \rightarrow D(\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}})$$

is fully faithful, and its essential image is stable under all limits and colimits and given by those $C \in D(\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}})$ such that for all extremally disconnected S , the map

$$R\text{Hom}_{\underline{\mathcal{A}}}(\mathcal{A}[S], C) \rightarrow R\text{Hom}_{\underline{\mathcal{A}}}(\underline{\mathcal{A}}[S], C)$$

is an isomorphism; in that case, also the $R\text{Hom}$'s agree. An object $C \in D(\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}})$ lies in $D(\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}})$ if and only if all $H^i(C)$ are in $\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}}$. The inclusion $D(\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}}) \subset D(\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}})$ admits a left adjoint

$$D(\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}}) \rightarrow D(\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}}) : C \mapsto C \otimes_{\underline{\mathcal{A}}}^L \mathcal{A}$$

that is the left derived functor of $M \mapsto M \otimes_{\underline{\mathcal{A}}} \mathcal{A}$. If $\underline{\mathcal{A}}$ is commutative, there is a unique symmetric monoidal tensor product $\otimes_{\underline{\mathcal{A}}}^L$ on $D(\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}})$ making the functor

$$D(\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}}) \rightarrow D(\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}}) : C \mapsto C \otimes_{\underline{\mathcal{A}}}^L \mathcal{A}$$

symmetric monoidal.

This shows that Theorem 6.11 implies most of Theorem 6.10. We note one omission in the statement of the proposition: It is not claimed that $\otimes_{\underline{\mathcal{A}}}^L$ is the left derived functor of $\otimes_{\underline{\mathcal{A}}}$. This is equivalent to the assertion that for all extremally disconnected S, S' , the complex

$$\mathcal{A}[S] \otimes_{\underline{\mathcal{A}}}^L \mathcal{A}[T] = \underline{\mathcal{A}}[S \times T] \otimes_{\underline{\mathcal{A}}}^L \mathcal{A}$$

sits in degree 0. But in the situation of Theorem 6.10, we defined $\mathcal{A}[S]$ for all profinite S , and Theorem 6.11 ensures that

$$\underline{\mathcal{A}}[S] \otimes_{\underline{\mathcal{A}}}^L \mathcal{A} = \mathcal{A}[S]$$

for all profinite sets S . Indeed, the right hand side lies in $\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}}$, and the $R\text{Hom}$ into any complex $C \in D(\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}})$ agrees, by representing C by a complex of direct sums of projectives, taking the limit of Postnikov truncations, and using Theorem 6.11.

In Theorem 6.10, we used the strange base ring $\mathbb{Z}[T^{-1}]$. We note that for analytic rings, one can always change the base ring to its natural choice, and maintain an analytic ring:

PROPOSITION 6.15. *Let \mathcal{A} be an analytic ring. Then $\underline{\mathcal{A}}' = \mathcal{A}[*]$ is naturally a condensed ring, and $\mathcal{A}[S]$ is naturally a condensed $\underline{\mathcal{A}}'$ -module for all extremally disconnected S , defining a pre-analytic ring \mathcal{A}' with a map of pre-analytic rings $\mathcal{A} \rightarrow \mathcal{A}'$. The pre-analytic ring \mathcal{A}' is analytic, and the forgetful functor $D(\text{Mod}_{\underline{\mathcal{A}}'}^{\text{cond}}) \rightarrow D(\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}})$ is an equivalence.*

PROOF. Note that $\mathcal{A}[*]$ is the tensor unit in $\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}}$ (as the image of the tensor unit $\underline{\mathcal{A}}[*]$ under the symmetric monoidal functors $\otimes_{\underline{\mathcal{A}}} \mathcal{A}$). One has $\mathcal{A}[*] = \underline{\text{Hom}}_{\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}}}(\mathcal{A}[*], \mathcal{A}[*])$, which is thus naturally a condensed ring, and every object of $\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}}$, in particular $\mathcal{A}[S]$, is naturally a condensed module over it. Now for any K as in the definition of an analytic ring, we have $K \in \text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}}$. Then formally

$$R\text{Hom}_{\mathcal{A}[*]}(\mathcal{A}[S], K) = R\text{Hom}_{\underline{\mathcal{A}}}(\mathcal{A}[*] \otimes_{\underline{\mathcal{A}}}^L \mathcal{A}[S], K)$$

and as $K \in D(\text{Mod}_{\underline{\mathcal{A}}}^{\text{cond}})$, we can replace $\mathcal{A}[*] \otimes_{\underline{\mathcal{A}}}^L \mathcal{A}[S]$ by

$$(\mathcal{A}[*] \otimes_{\underline{\mathcal{A}}}^L \mathcal{A}[S]) \otimes_{\underline{\mathcal{A}}}^L \mathcal{A}.$$

As $\otimes_{\mathcal{A}}^L \mathcal{A}$ is symmetric monoidal, and both tensor factors on the left are already in $D(\text{Mod}_{\mathcal{A}}^{\text{cond}})$, this agrees with

$$\mathcal{A}[*] \otimes_{\mathcal{A}}^L \mathcal{A}[S] = \mathcal{A}[S]$$

as $\mathcal{A}[*]$ is the tensor unit. But finally

$$R\text{Hom}_{\mathcal{A}}(\mathcal{A}[S], K) = R\text{Hom}_{\mathcal{A}}(\underline{\mathcal{A}}[S], K)$$

as \mathcal{A} is an analytic ring, and the latter is formally the same as

$$R\text{Hom}_{\mathcal{A}'}(\underline{\mathcal{A}}'[S], K).$$

It follows that \mathcal{A}' is an analytic ring. The present computation then also proves that $D(\text{Mod}_{\mathcal{A}'}^{\text{cond}}) \rightarrow D(\text{Mod}_{\mathcal{A}}^{\text{cond}})$ is an equivalence. \square

In particular, $\underline{\mathcal{A}} = \mathbb{Z}((T))_{>r}$ with $\mathcal{A}[S] = \mathcal{M}(S, \mathbb{Z}((T))_{>r})$ defines an analytic ring.

Finally, let us prove Theorem 6.6. Given $0 < p' < p \leq 1$, pick $x = \frac{1}{2}$, $r = x^p$ and $r' = x^{p'}$. Then we can take $f_x = 2 - T^{-1}$, and Theorem 6.9 (2) says that

$$\mathcal{M}(S, \mathbb{Z}((T))_{r'}) = \mathcal{M}_{p'}(S)$$

while (by passage to a colimit over all $r' > r$)

$$\mathcal{M}(S, \mathbb{Z}((T))_{>r}) = \mathcal{M}_{<p}(S).$$

We regard \mathbb{R} everywhere as $\mathbb{Z}[T^{-1}]$ -algebra via $T^{-1} \mapsto 2$. Given V as in Theorem 6.6, it is automatic that $V \in \text{Liq}_r(\mathbb{Z}[T^{-1}])$. Using Theorem 6.11, this implies that

$$R\text{Hom}_{\mathbb{Z}[T^{-1}]}(\mathcal{M}(S, \mathbb{Z}((T))_{r'}), V) = R\text{Hom}_{\mathbb{Z}[T^{-1}]}(\mathbb{Z}[T^{-1}][S], V)$$

for all profinite sets S . The right-hand side agrees with $R\text{Hom}_{\mathbb{Z}}(\mathbb{Z}[S], V)$, while the left-hand side agrees with $R\text{Hom}_{\mathbb{Z}}(\mathcal{M}_{p'}(S), V)$ by taking the quotient by $f_x = 2 - T^{-1}$. Thus, this proves a version of Theorem 6.11 where we use the base ring \mathbb{Z} in place of \mathbb{R} .

From this discussion, by taking the colimit over $p' > p$, one deduces that \mathbb{Z} with the modules $\mathcal{M}_{<p}(S)$ defines an analytic ring structure. Using the proposition above, this then also implies that \mathbb{R} with the same modules $\mathcal{M}_{<p}(S)$ defines an analytic ring, and that the forgetful functor $D(\text{Liq}_r(\mathbb{R})) \rightarrow D(\text{Cond}(\text{Ab}))$ is fully faithful. In particular, we see that a liquid \mathbb{R} -vector space admits a unique \mathbb{R} -linear structure.

Finally, the exact statement of Theorem 6.6 follows by observing that

$$R\text{Hom}_{\mathbb{R}}(\mathcal{M}_{p'}(S), V) = R\text{Hom}_{\mathbb{Z}}(\mathbb{R} \otimes_{\mathbb{Z}}^L \mathcal{M}_{p'}(S), V)$$

and the p -liquidification of $\mathbb{R} \otimes_{\mathbb{Z}}^L \mathcal{M}_{p'}(S)$ is the same as the p -liquidification of $\mathcal{M}_{p'}(S)$ (as p -liquidification is symmetric monoidal and the first tensor factor is the p -liquid tensor unit).

7. Lecture VII: $\mathbb{Z}((T))_{>r}$ is a principal ideal domain

In this lecture, we prove the following theorem.

THEOREM 7.1 ([Har84]). *For any real number r with $0 < r < 1$, the ring*

$$\mathbb{Z}((T))_{>r} = \left\{ \sum_{n \gg -\infty} a_n T^n \mid \exists r' > r, |a_n|(r')^n \rightarrow 0 \right\}$$

is a principal ideal domain. The nonzero prime ideals are the following:

- (1) *For any nonzero complex number $x \in \mathbb{C}$ with $|x| \leq r$, the kernel of the map*

$$\mathbb{Z}((T))_{>r} \rightarrow \mathbb{C} : \sum a_n T^n \mapsto \sum a_n x^n,$$

where this kernel depends only on (and determines) x up to complex conjugation;

- (2) *For any prime number p , the ideal (p) ;*

- (3) *For any prime number p and any topologically nilpotent unit $x \in \overline{\mathbb{Q}}_p$, the kernel of the map*

$$\mathbb{Z}((T))_{>r} \rightarrow \overline{\mathbb{Q}}_p : \sum a_n T^n \mapsto \sum a_n x^n,$$

where this kernel depends only on (and determines) the Galois orbit of x .

PROOF. It is clear that the ideals described are prime ideals. First, we check that all of them are principal, and that the respective quotient rings are already fields (so that these ideals are also maximal).

In case (1), assume first that x is real, so $0 < x \leq r$ or $-r \leq x < 0$. The second case reduces to the first under the involution of $\mathbb{Z}((T))_{>r}$ taking T to $-T$, so we assume for concreteness that $0 < x \leq r$. We start by verifying the surjectivity of

$$\mathbb{Z}((T))_{>r} \rightarrow \mathbb{R} : \sum a_n T^n \mapsto \sum a_n x^n.$$

Fix some integer N such that $x \geq \frac{1}{N}$. Then we claim that any element $y \in \mathbb{R}_{\geq 0}$ is represented by a power series with coefficients in $[0, N-1]$. Note that when $x = \frac{1}{N}$, this is precisely an N -adic expansion of y . For the proof, take the maximal n such that $x^n \leq y$, and the maximal integer a_n such that $a_n x^n \leq y$. Then necessarily $a_n \leq N-1$ (otherwise $x^{n+1} \leq y$). Now pass to $y - a_n x^n$ and repeat, noting that the sequence of n 's is strictly decreasing.

To find a generator, we first find a polynomial $g_n \in 1 + T^n \mathbb{R}[T]$ such that inside the disc $\{0 < |y| \leq r\}$, the only zero is x , with multiplicity 1. We argue by induction on $n = 1$; for $n = 1$, we can take $g_1 = 1 - x^{-1}T$. Now write $g_n = 1 + a_n T^n + \dots$. Then, taking any integer $m > |a_n|$, we can take

$$g_{n+1} = g_n(1 - \frac{a_n}{m} T^n)^m.$$

Now choose n large enough so that $r^n < 2(1-r)$ and let $g = g_n$. We can find some $h = 1 + c_n T^n + c_{n+1} T^{n+1} + \dots \in 1 + T^n \mathbb{R}[[T]]$ such that all $|c_i| \leq \frac{1}{2}$ and $f = gh \in 1 + T^n \mathbb{Z}[[T]]$. Note that h is invertible on $\{y \mid 0 < |y| \leq r\}$, as

$$\left| \sum_{i \geq n} c_i y^i \right| \leq \sum_{i \geq n} \frac{1}{2} r^i = \frac{1}{2} \frac{r^n}{1-r} < 1.$$

We claim that f generates the kernel of evaluation at x . Indeed, if $f' \in \mathbb{Z}((T))_{>r}$ vanishes at x , then $\frac{f'}{f} \in \mathbb{Z}((T))$ (as $f \in 1 + T\mathbb{Z}[[T]] \subset \mathbb{Z}[[T]]^\times$) and it still defines a holomorphic function on $\{0 < |y| < r'\}$ for some $r' > r$; thus, $\frac{f'}{f} \in \mathbb{Z}((T))_{>r}$.

Still in case (1), if now x is complex, we argue similarly: First, surjectivity of

$$\mathbb{Z}((T))_{>r} \rightarrow \mathbb{C} : \sum a_n T^n \mapsto \sum a_n x^n$$

can be proved in a similar way, this time allowing coefficients in $[-N, N]$ such that for any element $y \in \mathbb{C}$ with $|y| \leq 1$ there are integers $a_0, a_1 \in [-N, N]$ with $|y - a_0 - a_1 x| \leq |x|^2$. To find a generator, argue as above by finding first a polynomial $g_n \in 1 + T^n \mathbb{R}[T]$ whose only zeroes in $\{0 < |y| \leq r\}$ are x and its complex conjugate \bar{x} with multiplicity 1, starting with $g_1 = (1 - x^{-1}T)(1 - \bar{x}^{-1}T)$. Multiplying by a power series h as before then produces the desired generator $f = gh \in \mathbb{Z}((T))_{>r}$.

In case (2), it is clear that the ideals are principal. We claim that the map

$$\mathbb{Z}((T))_{>r}/p \rightarrow \mathbb{F}_p((T))$$

is an isomorphism. It is clearly injective. For surjectivity, we use that there is a set-theoretic section $\mathbb{F}_p \rightarrow \mathbb{Z}$ with image $\{0, 1, \dots, p-1\}$, so we can always lift to a power series with bounded coefficients a_n . But any power series with bounded coefficients lies in $\mathbb{Z}((T))_{>r}$.

It remains to handle case (3). Let $K \subset \overline{\mathbb{Q}}_p$ be the field generated by x , which is a finite extension of \mathbb{Q}_p ; let d be its degree. First, we check that the map

$$f_x : \mathbb{Z}((T))_{>r} \rightarrow K : \sum a_n T^n \mapsto \sum a_n x^n$$

is surjective. More precisely, if we denote by $\Lambda \subset \mathcal{O}_K$ the \mathbb{Z}_p -lattice generated by x , we claim that the map

$$\mathbb{Z}[[T]]_{>r} \rightarrow \Lambda : \sum a_n T^n \mapsto \sum a_n x^n$$

is surjective. For this, observe that if $x^d = up^m$ for some unit $u \in \mathcal{O}_K$ and integer $m > 0$, then any element of Λ/x^d can be written uniquely as a sum $\sum_{j=0}^{d-1} a_j x^j$ with all $a_j \in \{0, \dots, p^m - 1\}$. It follows that one can reach any element of Λ uniquely by a power series $\sum a_n T^n$ where all $a_n \in \{0, \dots, p^m - 1\}$. Indeed, the projection to Λ/x^d determines the first d coefficients uniquely, and then we can divide by T^d and induct.

Moreover, representing the element $p^m \in \Lambda$ in this way, we get an equation

$$p^m = f_x \left(\sum_{n>0} a_n T^n \right)$$

where all $a_n \in \{0, \dots, p^m - 1\}$. Consider the element

$$g_x = p^m - \sum a_n T^n$$

in the kernel of f_x . Note that g_x may have additional archimedean zeroes. However, as we already have generators for them, we can write $g_x = g'_x g''_x$ where g'_x has no archimedean zeroes, while $g''_x \in 1 + T\mathbb{Z}[[T]]_{>r}$. We claim that g'_x generates the kernel of f_x . Indeed, it clearly lies in the kernel of f_x ; we have to see that if $h \in \mathbb{Z}((T))_{>r}$ is any element in the kernel of f_x , then $\frac{h}{g'_x} \in \mathbb{Q}((T))$ still lies in $\mathbb{Z}((T))_{>r}$. As $g'_x \in p^m + T\mathbb{Z}[[T]]$, the inverse of g'_x has only powers of p in the denominator, so $\frac{h}{g'_x} \in \mathbb{Z}[\frac{1}{p}]((T))$. But it also lies in $\mathbb{Z}_p[[T]]$ as g'_x is a generator of the kernel of $\mathbb{Z}_p[[T]] \rightarrow \mathcal{O}_K$. It remains to see that the convergence condition is satisfied, but for this we simply note that g'_x has

no archimedean zeroes, so the quotient still defines a holomorphic function on $\{0 < |y| < r'\}$ for some $r' > r$.

Finally, we can finish the proof that $\mathbb{Z}((T))_{>r}$ is a principal ideal domain. Take any nonzero element $f \in \mathbb{Z}((T))_{>r}$. We want to see that, up to a unit, it is a product of the generators of the principal ideals listed in the statement. Up to scaling by a unit, it is of the form

$$f = a_0 + \sum_{n>0} a_n T^n$$

where $a_0 > 0$. We can now see that it lies in only finitely many of the principal ideals listed above: This is a standard result in complex analysis for the first points, and for the second and third type of points, only primes p dividing a_0 are relevant. There are only finitely many such, and for any p these primes are all contracted from $\mathbb{Z}_p((T))$, where the relevant finiteness holds. (We also see that it lies in any of these maximal ideals with finite multiplicity.) Dividing by the corresponding generators, we can assume that f does not lie in any of the principal ideals listed in the statement. This implies that $a_0 = 1$ as otherwise there will be a zero at a p -adic place, by the similar result for $\mathbb{Z}_p((T))$. Then $f^{-1} \in 1 + T\mathbb{Z}[[T]]$, and it still has convergence radius $> r$ as f has no zeroes on the closed disc of radius r . Thus, f is invertible, as desired. \square

Now we can prove Theorem 6.9 from the last lecture.

Given any $0 < r' < r < 1$, we consider the map $\mathbb{Z}((T))_r \rightarrow \mathbb{R} : T \mapsto r'$. Applying the previous lemma to $x = r'$, we get some $f = f_{r'} \in \mathbb{Z}((T))_{>r}$ that vanishes only at r' . We claim that the sequence

$$0 \rightarrow \mathbb{Z}((T))_r \xrightarrow{f_{r'}} \mathbb{Z}((T))_r \rightarrow \mathbb{R} \rightarrow 0$$

is exact. Surjectivity was proved above (even $\mathbb{Z}((T))_{>r}$ surjects). On the other hand, if $g \in \mathbb{Z}((T))_r$ vanishes at x , then $gf_{r'}^{-1} \in \mathbb{Z}((T))$ and it follows from a standard complex analysis consideration that it still lies in $\mathbb{Z}((T))_r$.

We want to see that the sequence is still exact as condensed abelian groups, and we want to identify the spaces of measures. For this, we prove the following quantitative version:

PROPOSITION 7.2. *There are constants C_1, \dots, C_4 with the following properties:*

- (1) *If $g \in \mathbb{Z}((T))_{r, \leq c}$ then $f_{r'}g \in \mathbb{Z}((T))_{r, \leq C_1c}$.*
- (2) *Conversely, if $g \in \mathbb{Z}((T))_r$ such that $f_{r'}g \in \mathbb{Z}((T))_{r, \leq c}$, then $g \in \mathbb{Z}((T))_{r, \leq C_2c}$.*
- (3) *Let $0 < p < 1$ such that $(r')^p = r$. If $g \in \mathbb{Z}((T))_{r, \leq c}$, then $g(x) \in \mathbb{R}_{\ell^p \leq C_3c}$.*
- (4) *Conversely, for any $z \in \mathbb{R}_{\ell^p \leq c}$, there is some $g \in \mathbb{Z}((T))_{r, \leq C_4c}$ such that $g(x) = z$.*

The proposition easily implies exactness as condensed abelian groups (think in terms of ind-(compact Hausdorff) sets, and use that for maps of compact Hausdorff spaces, surjectivity can be checked on points).

PROOF. Take C_1 so that $f_{r'} \in \mathbb{Z}((T))_{r, \leq C_1}$, then the claim about C_1 is clear. For the claim about C_2 , note that the norm can be expressed in terms of the restriction to a function on $\{y \mid |r''| \leq |y| \leq |r|\}$ for $r' < r'' < r$. On that strip, $f_{r'}$ is invertible, so picking a bound of the norm for the inverse gives the claim.

For C_3 , let $g = \sum a_n T^n$ with $\sum |a_n| r^n \leq c$. Then

$$|g(x)|^p \leq \sum |a_n|^p (r')^{np} = \sum |a_n|^p r^n \leq \sum |a_n| r^n \leq c,$$

so we can in fact take $C_3 = 1$. Here, we used critically that all a_n are integers, which implies that $|a_n|^p \leq |a_n|$: This is true whenever $a_n = 0$ or $|a_n| \geq 1$ (when $p < 1$).

For C_4 , take any $z \in \mathbb{R}$ with $|z|^p \leq c$ and assume $z \neq 0$. Take n minimal so that $(r')^n \leq |z|$, and write $z = a_n(r')^n + z'$ where $a_n \in \mathbb{Z}$ and $|z'| < (r')^n$. Note that $|a_n| \leq (r')^{-1}$. Continuing, we can write $z = g(x)$ for some $g = a_n T^n + \dots \in \mathbb{Z}((T))$ with $|a_i| \leq (r')^{-1}$ for all i . Then

$$\sum |a_i| r^i \leq (r')^{-1} \frac{r^n}{1-r} \leq \frac{1}{r'(1-r)} |z|^p,$$

so we can take $C_4 = \frac{1}{r'(1-r)}$. \square

From here, by observing that these bounds immediately generalize to finite free modules, one easily deduces Theorem 6.9 (2). For part (3), we need a corresponding version. Note first that

$$\mathbb{Z}((T))_r / (f_{r'})^m \cong \mathbb{R}[X]/X^m$$

where the isomorphism sends T to $[r'] = (r')^{1+X} = r' + r' \log r' X + \dots$. In other words, we have a short exact sequence

$$0 \rightarrow \mathbb{Z}((T))_r \xrightarrow{f_{r'}^m} \mathbb{Z}((T))_r \xrightarrow{\theta_m} \mathbb{R}[X]/X^m \rightarrow 0.$$

PROPOSITION 7.3. *There are constants C_1, \dots, C_4 (depending on m) with the following properties:*

- (1) *If $g \in \mathbb{Z}((T))_{r, \leq c}$ then $f_{r'}^m g \in \mathbb{Z}((T))_{r, \leq C_1 c}$.*
- (2) *Conversely, if $g \in \mathbb{Z}((T))_r$ such that $f_{r'}^m g \in \mathbb{Z}((T))_{r, \leq c}$, then $g \in \mathbb{Z}((T))_{r, \leq C_2 c}$.*
- (3) *If $g \in \mathbb{Z}((T))_{r, \leq c}$, then $\theta_m(g) \in (\mathbb{R}[X]/X^m)_{\ell^p \leq C_3 c}$.*
- (4) *Conversely, for any $z \in (\mathbb{R}[X]/X^m)_{\ell^p \leq c}$, there is some $g \in \mathbb{Z}((T))_{r, \leq C_4 c}$ such that $\theta_m(g) = z$.*

PROOF. Parts (1) and (2) follows inductively from parts (1) and (2) of the previous proposition. It remains to handle part (3): Part (4) then follows by successive approximation (using parts (3) and (4) of the previous proposition, and the other claims already established).

Thus let $g = \sum a_n T^n \in \mathbb{Z}((T))_r$ with $\sum |a_n| r^n \leq c$. Then

$$\theta_m(g) = \sum a_n [r']^n \in \mathbb{R}[X]/X^m.$$

We need to bound this. Recall from Proposition 6.4 that there is some constant C such that for all maps $f : S \rightarrow S'$ of finite sets, one has

$$f((\mathbb{R}[X]/X^m)[S]_{\ell^p \leq c}) \subset (\mathbb{R}[X]/X^m)[S']_{\ell^p \leq Cc}.$$

We claim that we can take $C_3 = C$; equivalently,

$$\sum a_n [r']^n \in (\mathbb{R}[X]/X^m)_{\ell^p \leq Cc}.$$

To check this, we can assume that the sum is finite (as the target is closed). Let S be a finite set of cardinality $\sum |a_n|$. Then we can define an element $\tilde{z} \in (\mathbb{R}[X]/X^m)[S]$ whose coefficients are of the form $\pm [r']^n = [\pm r']^n$, such that the image of \tilde{z} in $\mathbb{R}[X]/X^m$ (the sum of the coefficients) is $\sum a_n [r']^n$. Thus, it suffices to see that $\tilde{z} \in (\mathbb{R}[X]/X^m)[S]_{\ell^p \leq Cc}$. But this is clear: One has to bound the sum of $(r')^{np} = r^n$, where there are $|a_n|$ occurrences of r^n , giving in total at most $\sum |a_n| r^n \leq c$. \square

The lecturer finds the way in which the varying ℓ^p -norms, as well as the subtle B_{dR}^+ -type deformations, arise from the very simple-minded spaces of measures for $\mathbb{Z}((T))_r$ quite striking. The discreteness of \mathbb{Z} is critical for this behaviour: In some sense, the real B_{dR}^+ arises via some kind of discretization.

8. Lecture VIII: Reduction to “Banach spaces”

Fix some $0 < r < 1$. Recall that we want to prove Theorem 6.11:

THEOREM 8.1. *Let K be a condensed $\mathbb{Z}[T^{-1}]$ -module that is the kernel of some map*

$$f : \bigoplus_{i \in I} \mathcal{M}(S_i, \mathbb{Z}((T))_{>r}) \rightarrow \bigoplus_{j \in J} \mathcal{M}(S'_j, \mathbb{Z}((T))_{>r})$$

where all S_i and S'_j are extremally disconnected. Then for all $1 > r' > r$ and all profinite sets S , the map

$$R\mathbf{Hom}_{\mathbb{Z}[T^{-1}]}(\mathcal{M}(S, \mathbb{Z}((T))_{r'}), K) \rightarrow R\mathbf{Hom}_{\mathbb{Z}[T^{-1}]}(\mathbb{Z}[T^{-1}][S], K)$$

in $D(\mathbf{Cond}(\mathbf{Ab}))$ is an isomorphism.

Today, we make some preliminary reductions, and in particular reduce to the case where K is a suitable analogue of a Banach space. First, we note the following simple proposition.

PROPOSITION 8.2. *Let S and S' be two profinite sets and let $f : S \rightarrow \mathcal{M}(S', \mathbb{Z}((T))_r)$ be a map. Then for any $r' \geq r$ less than 1, there is a unique $\mathbb{Z}[T^{-1}]$ -linear map*

$$\mathcal{M}(S, \mathbb{Z}((T))_{r'}) \rightarrow \mathcal{M}(S', \mathbb{Z}((T))_r)$$

extending the given map f .

PROOF. First, we prove uniqueness. Thus assume that some map $g : \mathcal{M}(S, \mathbb{Z}((T))_{r'}) \rightarrow \mathcal{M}(S', \mathbb{Z}((T))_r)$ has trivial restriction to S . By $\mathbb{Z}[T^{-1}]$ -linearity, it vanishes on $\mathbb{Z}[T^{\pm 1}][S]$, but this is dense in $\mathcal{M}(S, \mathbb{Z}((T))_{r'})$ (and everything is quasiseparated). Thus, the map g vanishes.

For existence, we may assume that $r' = r$, and by rescaling that f maps S into $\mathcal{M}(S', \mathbb{Z}((T))_r)_{\leq 1}$. We want to show that there is a map of profinite sets

$$\mathcal{M}(S, \mathbb{Z}((T))_r)_{\leq 1} \rightarrow \mathcal{M}(S', \mathbb{Z}((T))_r)_{\leq 1}$$

extending the given map on the dense subset $\mathbb{Z}[T^{\pm 1}][S]$. Writing S' as an inverse limit of finite sets, this reduces to the case that S' is finite. Then

$$\mathcal{M}(S', \mathbb{Z}((T))_r)_{\leq 1} \subset \prod_{n \geq 0, s' \in S'} \mathbb{Z} \cdot T^n[s'] = \mathbb{Z}[[T]][S'].$$

Moreover, it is defined as an inverse limit of finite subsets of $(\mathbb{Z}[T]/T^m)[S']$, so we can work modulo T^m for some m . Then the map $S \rightarrow \mathcal{M}(S', \mathbb{Z}((T))_r)_{\leq 1} \rightarrow (\mathbb{Z}[T]/T^m)[S']$ factors through some finite quotient $S \rightarrow S_m$, and we can also reduce to the case that S is finite. In that case, the desired map

$$\mathcal{M}(S, \mathbb{Z}((T))_r) \rightarrow \mathcal{M}(S', \mathbb{Z}((T))_r)$$

clearly exists, and is given by $\sum_{s \in S} a_s[s] \mapsto \sum_{s \in S} a_s f(s)$. One verifies that this indeed maps $\mathcal{M}(S, \mathbb{Z}((T))_r)_{\leq 1}$ into $\mathcal{M}(S', \mathbb{Z}((T))_r)_{\leq 1}$. \square

The following proposition is easy to prove by using that we choose exactly $\mathbb{Z}[T^{-1}]$ as the base ring. (It is also true that $\mathcal{M}_p(S)$ is a pseudocoherent \mathbb{R} -module, but this is somewhat more tricky to prove.)

PROPOSITION 8.3. *For any profinite set S , the condensed $\mathbb{Z}[T^{-1}]$ -module $\mathcal{M}(S, \mathbb{Z}((T))_r)$ is pseudocoherent.*

PROOF. Let $M = \mathcal{M}(S, \mathbb{Z}((T))_r)$. We have the Breen-Deligne resolution

$$\dots \rightarrow \bigoplus_{j=1}^{n_i} \mathbb{Z}[M^{r_{ij}}] \rightarrow \dots \rightarrow \mathbb{Z}[M^2] \rightarrow \mathbb{Z}[M] \rightarrow M \rightarrow 0.$$

Here n_i and all r_{ij} are nonnegative integers. Note that M has an endomorphism (multiplication by T^{-1}), and by functoriality in M , the whole sequence is a sequence of condensed $\mathbb{Z}[T^{-1}]$ -modules. Thus, it is enough to see that each $\mathbb{Z}[M^r]$ is pseudocoherent as $\mathbb{Z}[T^{-1}]$ -module. Replacing S by r copies of itself, we can assume that $r = 1$. Now M has a profinite subset $M_{\leq 1} \subset M$ and $M = \bigcup_{n \geq 0} T^{-n} M_{\leq 1}$. In fact, we claim that the sequence

$$0 \rightarrow \mathbb{Z}[M_{\leq r}][T^{-1}] \xrightarrow{T^{-1}-[T^{-1}]} \mathbb{Z}[M_{\leq 1}][T^{-1}] \rightarrow \mathbb{Z}[M] \rightarrow 0$$

is an exact sequence of $\mathbb{Z}[U]$ -modules. This will give the desired result as the other two terms are pseudocoherent $\mathbb{Z}[T^{-1}]$ -modules. But the sequence is the filtered colimit of the sequences

$$0 \rightarrow \bigoplus_{i=0}^{n-1} \mathbb{Z}[M_{\leq r}] \cdot T^{-i} \xrightarrow{T^{-1}-[T^{-1}]} \bigoplus_{i=0}^n \mathbb{Z}[M_{\leq 1}] \cdot T^{-i} \rightarrow \mathbb{Z}[T^{-n} M_{\leq 1}] \rightarrow 0$$

which are exact by induction on n : For $n = 0$, the first term is zero and the second map is an equality, and for $n > 0$ the quotient by the sequence for $n - 1$ (multiplied by T^{-1}) is an isomorphism $\mathbb{Z}[M_{\leq r}] \cong \mathbb{Z}[M_{\leq 1}]$ between the terms of degree 0 (given by multiplication by T^{-1} : $M_{\leq r} \cong M_{\leq 1}$). \square

It follows that $R\mathbf{Hom}_{\mathbb{Z}[T^{-1}]}(\mathcal{M}(S, \mathbb{Z}((T))_r), K)$ commutes with filtered colimits in K . In particular, we can assume that K is the kernel of some map

$$\mathcal{M}(S_1, \mathbb{Z}((T))_{>r}) \rightarrow \bigoplus_{j \in J} \mathcal{M}(S'_j, \mathbb{Z}((T))_{>r}).$$

Note that by Proposition 8.2 and Proposition 8.3, for any $r' > r$ one has

$$\mathrm{Hom}(\mathcal{M}(S_1, \mathbb{Z}((T))_{r'}), \bigoplus_{j \in J} \mathcal{M}(S'_j, \mathbb{Z}((T))_{>r})) = \bigoplus_{j \in J} \mathcal{M}(S'_j, \mathbb{Z}((T))_{>r})(S_1).$$

Passing to the limit $r' \rightarrow r$, this shows that also

$$\mathrm{Hom}(\mathcal{M}(S_1, \mathbb{Z}((T))_{>r}), \bigoplus_{j \in J} \mathcal{M}(S'_j, \mathbb{Z}((T))_{>r})) = \bigoplus_{j \in J} \mathcal{M}(S'_j, \mathbb{Z}((T))_{>r})(S_1).$$

In particular, the given map factors over a finite direct sum, and we may assume that K is the kernel of some map

$$\mathcal{M}(S_1, \mathbb{Z}((T))_{>r}) \rightarrow \mathcal{M}(S_2, \mathbb{Z}((T))_{>r}),$$

and we recall that any such map is uniquely induced by a map $S_1 \rightarrow \mathcal{M}(S_2, \mathbb{Z}((T))_{>r})$. In fact, by a triangle (and noting that $\mathcal{M}(S_1, \mathbb{Z}((T))_{>r})$ itself is a special case of such an image, when taking the identity map), it is enough to prove the similar assertion for the image of the map

$$\mathcal{M}(S_1, \mathbb{Z}((T))_{>r}) \rightarrow \mathcal{M}(S_2, \mathbb{Z}((T))_{>r}).$$

In other words, we are reduced to the following assertion. (We could assume that S_1 and S_2 are extremally disconnected instead of merely profinite, but this will be of no use.)

THEOREM 8.4. *Let S_1 and S_2 be profinite sets and let $f : \mathcal{M}(S_1, \mathbb{Z}((T))_{>r}) \rightarrow \mathcal{M}(S_2, \mathbb{Z}((T))_{>r})$ be a map of condensed $\mathbb{Z}[T^{-1}]$ -modules; these are in bijection with maps $f|_{S_1} : S_1 \rightarrow \mathcal{M}(S_2, \mathbb{Z}((T))_{>r})$. Let M be the image of f . Then for any $r' > r$, the map*

$$R\text{Hom}_{\mathbb{Z}[T^{-1}]}(\mathcal{M}(S, \mathbb{Z}((T))_{r'}), M) \rightarrow R\text{Hom}_{\mathbb{Z}[T^{-1}]}(\mathbb{Z}[T^{-1}][S], M)$$

in $D(\text{Cond}(\text{Ab}))$ is an isomorphism.

We wish to reduce further. The datum is the map $S_1 \rightarrow \mathcal{M}(S_2, \mathbb{Z}((T))_{>r})$. This factors over $\mathcal{M}(S_2, \mathbb{Z}((T))_{>r_0})$ for some $r_0 > r$, and fixing such a factorization, we get modules $M_{r''}$ for all $r_0 \geq r'' > r$ as the image of $\mathcal{M}(S_1, \mathbb{Z}((T))_{r''}) \rightarrow \mathcal{M}(S_2, \mathbb{Z}((T))_{r''})$, and M is the filtered colimit of the $M_{r''}$. Note that $M_{r''}$ can be endowed with subspaces $M_{r'', \leq c}$ by taking the image of $\mathcal{M}(S_1, \mathbb{Z}((T))_{r''})_{\leq c}$. These are the inverse limits of the images $M_{r'', i, \leq c}$ in $\mathcal{M}(S_{2,i}, \mathbb{Z}((T))_{r''})$, where we write S_2 as an inverse limit of finite sets $S_{2,i}$. We need the following proposition.

One can actually identify the module $M_{r'', i}$ that appears, by applying the following proposition with r'' in place of r .

PROPOSITION 8.5. *Let S_1 be a profinite set and let $S_1 \rightarrow \mathbb{Z}((T))_{>r}^n$ be a map for some integer n . Then the image M of*

$$\mathcal{M}(S_1, \mathbb{Z}((T))_r) \rightarrow \mathbb{Z}((T))_r^n$$

is isomorphic (as condensed $\mathbb{Z}[T^{-1}]$ -module) to $\mathbb{Z}((T))_r^{n'}$ for some $n' \leq n$.

PROOF. By Theorem 7.1, the image of

$$\mathcal{M}(S_1, \mathbb{Z}((T))_{>r}) \rightarrow \mathbb{Z}((T))_{>r}^n$$

on underlying modules is isomorphic to $\mathbb{Z}((T))_{>r}^{n'}$ for some $n' \leq n$. Let $M' = \mathbb{Z}((T))_{>r}^{n'}$ as a condensed $\mathbb{Z}[T^{-1}]$ -module, which comes equipped with a map to $\mathbb{Z}((T))_{>r}^n$. This map is injective as both are quasiseparated as condensed sets and the underlying map is injective. Moreover, the quotient of condensed $\mathbb{Z}[T^{-1}]$ -modules

$$\mathbb{Z}((T))_{>r}^n / M'$$

is still quasiseparated, by the structure of finitely generated $\mathbb{Z}((T))_{>r}$ -modules (using Theorem 7.1). It follows that the map $S_1 \rightarrow \mathbb{Z}((T))_{>r}^n$ factors over M' , and replacing $\mathbb{Z}((T))_{>r}^n$ by M' , we may assume that $\mathcal{M}(S_1, \mathbb{Z}((T))_{>r}) \rightarrow \mathbb{Z}((T))_{>r}^n$ is surjective (on underlying modules). In particular, the image contains a basis, which implies that also

$$\mathcal{M}(S, \mathbb{Z}((T))_r) \rightarrow \mathbb{Z}((T))_r^n$$

is surjective as condensed modules, giving the desired result. \square

The next proposition expresses $M_{r''}$ in terms of the $M_{r'', i}$.

PROPOSITION 8.6. *For any r'' , the complex*

$$0 \rightarrow M_{r''} \rightarrow \bigcup_{c>0} \prod_i M_{r'', i, \leq c} \rightarrow \bigcup_{c>0} \prod_{i_0 \rightarrow i_1} M_{r'', i_1, \leq c} \rightarrow \dots \rightarrow \bigcup_{c>0} \prod_{i_0 \rightarrow \dots \rightarrow i_m} M_{r'', i_m, \leq c} \rightarrow \dots$$

of condensed $\mathbb{Z}[T^{-1}]$ -modules is exact. More precisely, taking E -valued points for some extremally disconnected set E , it is exact, and if

$$g \in \ker\left(\prod_{i_0 \rightarrow \dots \rightarrow i_m} M_{r'', i_m, \leq c}(E) \rightarrow \prod_{i_0 \rightarrow \dots \rightarrow i_{m+1}} M_{r'', i_{m+1}}(E)\right)$$

then there exists some $h \in \prod_{i_0 \rightarrow \dots \rightarrow i_{m-1}} M_{r'', i_{m-1}, \leq c}(E)$ with $d(h) = g$.

We refer to the appendix for the construction of the complex.

PROOF. The claim is equivalent to the surjectivity of some map of profinite sets, more precisely that of

$$\begin{aligned} \prod_{i_0 \rightarrow \dots \rightarrow i_{m-1}} M_{r'', i_{m-1}, \leq c} \times \prod_{i_0 \rightarrow \dots \rightarrow i_m} M_{r'', i_m} & \prod_{i_0 \rightarrow \dots \rightarrow i_m} M_{r'', i_m, \leq c} \\ & \rightarrow \ker \left(\prod_{i_0 \rightarrow \dots \rightarrow i_m} M_{r'', i_m, \leq c} \rightarrow \prod_{i_0 \rightarrow \dots \rightarrow i_{m+1}} M_{r'', i_{m+1}} \right). \end{aligned}$$

We can write the index category \mathcal{I} of i 's as a filtered union of finite categories \mathcal{I}_j that admit an initial object. The corresponding map is a cofiltered limit of similar maps for \mathcal{I}_j in place of \mathcal{I} . Therefore it suffices to prove the similar assertion for each \mathcal{I}_j . In that case, the presence of an initial object gives a contracting homotopy, whose explicit description as given in the appendix to this lecture shows that it preserves the $\leq c$ -subspace. \square

In particular, it is enough to prove the following statement.

THEOREM 8.7. *Fix radii $1 > r' > r_0 > r > 0$. Let S_1 be a profinite set and let $f_i : S_1 \rightarrow \mathbb{Z}((T))_{>r_0}^{n_i}$ be maps for some index set $i \in I$. For $r_0 \geq r'' > r$, let $M_{r'', i} \subset \mathbb{Z}((T))_{r''}^{n_i}$ be the image of the induced maps $\mathcal{M}(S_1, \mathbb{Z}((T))_{r''}) \rightarrow \mathbb{Z}((T))_{r''}^{n_i}$, with the subspace $M_{r'', i, \leq c} \subset M_{r'', i}$ defined as the image of $\mathcal{M}(S_1, \mathbb{Z}((T))_{r''})_{\leq c}$.*

Then, letting

$$M = \varinjlim_{r'' > r} \bigcup_{c > 0} \prod_i M_{r'', i, \leq c},$$

the map

$$R\mathrm{Hom}_{\mathbb{Z}[T^{-1}]}(\mathcal{M}(S, \mathbb{Z}((T))_{r'}), M) \rightarrow R\mathrm{Hom}_{\mathbb{Z}[T^{-1}]}(\mathbb{Z}[T^{-1}][S], M)$$

is an isomorphism in $D(\mathrm{Cond}(\mathrm{Ab}))$ for all profinite sets S .

At this point, we will make a change from Smith to Banach spaces: The $M_{r'', i}$ are some kind of Smith spaces, but their norm allows one to define canonical Banach spaces inside them.

DEFINITION 8.8. Let N be a condensed abelian group written as an increasing union of compact Hausdorff subsets $N_{\leq c}$ for $c \geq 0$, satisfying the following conditions:

- (1) One has $N_{\leq 0} = 0$;
- (2) Each $N_{\leq c}$ is symmetric, i.e. $-N_{\leq c} = N_{\leq c}$;
- (3) It is exhaustive: $N = \bigcup_{c > 0} N_{\leq c}$;
- (4) It satisfies the triangle inequality: $N_{\leq c} + N_{\leq c'} \subset N_{\leq c+c'}$;
- (5) It is continuous: $N_{\leq c} = \bigcap_{c' > c} N_{\leq c'}$.

In that situation, we let N^B be the condensed abelian group which takes any profinite set S to the completion of the normed abelian group of locally constant maps from S to N .

In other words, as discussed in the appendix, N^B is the condensed abelian group associated to the normed abstract abelian group $N(*)$.

PROPOSITION 8.9. *There is a unique map $N^B \rightarrow N$ of condensed sets extending the identity on underlying abelian groups. It is an injective map of condensed abelian groups.*

PROOF. Given a map $f : S \rightarrow N^B$, choose a sequence of locally constant maps $f_0, f_1, \dots : S \rightarrow N$ with limit f (by definition $N^B(S)$ is the space of Cauchy sequences modulo the space of null sequences). This defines a map $S \times \mathbb{N} \rightarrow N$, and it is easy to see that it takes image in $N_{\leq c}$ for some c . Let $\Gamma \subset (S \times \mathbb{N}) \times N_{\leq c}$ be its graph, and let $\bar{\Gamma} \subset (S \times (\mathbb{N} \cup \{\infty\})) \times N_{\leq c}$ be the closure. We claim that $\bar{\Gamma} \rightarrow S \times (\mathbb{N} \cup \{\infty\})$ is an isomorphism. As it is a map of compact Hausdorff spaces, it is enough to check bijectivity, so pick some $s \in S$. We need to see that there is a unique preimage of (s, ∞) . Note that existence is clear, as $\bar{\Gamma}$ has dense image, and thus full image. On the other hand, uniqueness is a simple consequence of the Cauchy property and the separatedness $0 = \bigcup_{c>0} N_{\leq c}$ (which follows from condition (1) and (5)).

In particular, restricting to ∞ , we get a map $S \rightarrow N_{\leq c}$, thus defining a map $S \rightarrow N$. It is clear that this is functorial, additive, and independent of the choice of the f_i (as nullsequences will converge to zero), and defines a map of condensed sets $N^B \rightarrow N$ (and is the only possible choice). The rest of the assertions are also easy to verify. \square

In our situation, the following proposition ensures that replacing $M_{r'',i}$ by $M_{r'',i}^B$ is harmless.

PROPOSITION 8.10. *For any $r_0 \geq r'' > r''' > r$, the map $M_{r'',i} \rightarrow M_{r''',i}$ factors over $M_{r''',i}^B$.*

PROOF. By Proposition 8.5, one has $M_{r'',i} \cong \mathbb{Z}((T))_{r''}^{n'_i}$ for some n'_i , and then

$$M_{r''',i} = M_{r'',i} \otimes_{\mathbb{Z}((T))_{r''}} \mathbb{Z}((T))_{r'''} \cong \mathbb{Z}((T))_{r'''}^{n'_i}.$$

Thus, it suffices to see that $\mathbb{Z}((T))_{r''} \rightarrow \mathbb{Z}((T))_{r'''}$ factors over $\mathbb{Z}((T))_{r'''}^B$. It suffices to check this on $\mathbb{Z}((T))_{r'', \leq 1}$, and then it follows from the observation that $\mathbb{Z}((T))_{r'', \leq 1} \cap T^m \mathbb{Z}[[T]]$ maps into $\mathbb{Z}((T))_{r''', \leq (\frac{r'''}{r''})^m}$, so writing any S -valued section of $\mathbb{Z}((T))_{r'', \leq 1}$ as $\sum_{n \geq 0} a_n T^n$ with $a_n \in C(S, \mathbb{Z})$, this sum is actually convergent in $\mathbb{Z}((T))_{r'''}^B$. \square

This implies that

$$\varinjlim_{r'' > r} \bigcup_{c > 0} \prod_i M_{r'',i, \leq c} = \varinjlim_{r'' > r} \bigcup_{c > 0} \prod_i (M_{r'',i, \leq c}^B).$$

After this change, we can actually prove a statement for an individual r'' (which gives the previous statement by passage to the filtered colimit over all $r'' > r$, using Proposition 8.3 again). We are reduced to the following statement (where the r in the present statement is any of the r'' in the previous statement).

THEOREM 8.11. *Fix radii $1 > r' > r > 0$. Let S_1 be a profinite set and let $f_i : S_1 \rightarrow \mathbb{Z}((T))_{>r}^{n_i}$ be maps for some index set $i \in I$. Let $M_i \subset \mathbb{Z}((T))_r^{n_i}$ be the image of the induced maps $\mathcal{M}(S_1, \mathbb{Z}((T))_r) \rightarrow \mathbb{Z}((T))_r^{n_i}$, with the subspace $M_{i, \leq c} \subset M_i$ defined as the image of $\mathcal{M}(S_1, \mathbb{Z}((T))_{\leq c})$. Then, letting*

$$M = \bigcup_{c > 0} \prod_i (M_{i, \leq c}^B),$$

the map

$$R\mathrm{Hom}_{\mathbb{Z}[T^{-1}]}(\mathcal{M}(S, \mathbb{Z}((T))_{r'}), M) \rightarrow R\mathrm{Hom}_{\mathbb{Z}[T^{-1}]}(\mathbb{Z}[T^{-1}][S], M)$$

is an isomorphism in $D(\mathrm{Cond}(\mathrm{Ab}))$ for all profinite sets S .

Note that we are doing something slightly funny here as we are using the procedure $N \mapsto N^B$ inside the product over i : While $\bigcup_{c>0} \prod_i M_{i,\leq c}$ is of Smith-type (countable union of compact Hausdorff), the present construction is a strange mixture.

We will now get rid of the product over i by formulating a quantitative version of the statement for a single i . To show that the internal Hom agrees, we have to see that for any profinite set S' , one has

$$R\mathrm{Hom}_{\mathbb{Z}[T^{-1}]}(\mathcal{M}(S, \mathbb{Z}((T))_{r'})[S']/\mathbb{Z}[T^{-1}][S \times S'], M) = 0.$$

The source is a pseudocoherent $\mathbb{Z}[T^{-1}]$ -module, so we may pick a projective resolution $P_\bullet = P_\bullet(r', S, S')$ of $\mathcal{M}(S, \mathbb{Z}((T))_{r'})[S']/\mathbb{Z}[T^{-1}][S \times S']$, where each $P_i = \mathbb{Z}[T^{-1}][E_i]$ is the free condensed $\mathbb{Z}[T^{-1}]$ -module on some extremally disconnected set E_i . Then we have to see that the corresponding complex

$$0 \rightarrow M(E_0) \rightarrow M(E_1) \rightarrow \dots$$

is acyclic.

Now we observe that M is naturally the union of the subsets $M_{\leq c} = \prod_i (M_{i,\leq c}^B)$ over all $c > 0$. Using this observation, it is enough to prove the following quantitative result.

THEOREM 8.12. *Fix profinite sets S and S' and radii $1 > r' > r > 0$, as well as a projective resolution P_\bullet of $\mathcal{M}(S, \mathbb{Z}((T))_{r'})[S']/\mathbb{Z}[T^{-1}][S \times S']$, where each $P_i = \mathbb{Z}[T^{-1}][E_i]$ is the free condensed $\mathbb{Z}[T^{-1}]$ -module on some extremally disconnected set E_i .*

Then for any m there is some $C_m > 0$ with the following property. For any profinite set S_1 with a map $f : S_1 \rightarrow \mathbb{Z}((T))_{>r}^n$, letting M be the image of the induced map $\mathcal{M}(S_1, \mathbb{Z}((T))_r) \rightarrow \mathbb{Z}((T))_r^n$ with its subspaces $M_{\leq c}$, the complex

$$0 \rightarrow M^B(E_0) \rightarrow M^B(E_1) \rightarrow \dots$$

computing $R\mathrm{Hom}(\mathcal{M}(S, \mathbb{Z}((T))_{r'})[S']/\mathbb{Z}[T^{-1}][S \times S'], M)$ is exact, and if $g \in \ker(M^B(E_m) \rightarrow M^B(E_{m+1}))$ with $\|g\| \leq c$, then there is some $h \in M^B(E_{m-1})$ with $\|h\| \leq C_m c$ such that $d(h) = g$.

Now M^B with its norm is an example of the following structure:

DEFINITION 8.13. An r -normed $\mathbb{Z}[T^{\pm 1}]$ -module is a normed $\mathbb{Z}[T^{\pm 1}]$ -module V satisfying $\|Tv\| = r\|v\|$ for all $v \in V$.

To an r -normed $\mathbb{Z}[T^{\pm 1}]$ -module V , one can associate a condensed $\mathbb{Z}[T^{\pm 1}]$ -module \widehat{V} , as in the appendix. One main reason that we reduced to this ‘‘Banach space’’ setting is that one can use resolutions by profinite sets, instead of extremally disconnected sets, by Proposition 8.19. We end this lecture by reducing to the following statement.

THEOREM 8.14. *Fix radii $1 > r' > r > 0$. Then for all r -normed $\mathbb{Z}[T^{\pm 1}]$ -modules V and all profinite sets S , the map*

$$R\mathrm{Hom}_{\mathbb{Z}[T^{-1}]}(\mathcal{M}(S, \mathbb{Z}((T))_{r'}), \widehat{V}) \rightarrow \widehat{V}(S)$$

is a quasi-isomorphism.

THEOREM 8.14 IMPLIES THEOREM 8.12. Fix the profinite set S and take a projective resolution P_\bullet of $\mathcal{M}(S, \mathbb{Z}((T))_{r'})/\mathbb{Z}[T^{-1}][S]$, so that each $P_i = \mathbb{Z}[T^{-1}][E_i]$ is the free condensed $\mathbb{Z}[T^{-1}]$ -module on an extremally disconnected set E_i .

As $R\Gamma(S, \widehat{V}) = \widehat{V}(S)$ is concentrated in degree 0, Theorem 8.14 implies that the complex

$$0 \rightarrow \widehat{V}(E_0) \rightarrow \widehat{V}(E_1) \rightarrow \dots$$

computing $R\mathrm{Hom}_{\mathbb{Z}[T^{-1}]}(\mathcal{M}(S, \mathbb{Z}((T))_{r'})/\mathbb{Z}[T^{-1}][S], V)$, is acyclic. We claim that this implies that for all $m \geq 0$, there is some constant C_m (independent of V) such that for all $g \in \ker(\widehat{V}(E_m) \rightarrow \widehat{V}(E_{m+1}))$ with $\|g\| \leq c$, there is some $h \in \widehat{V}(E_{m-1})$ with $\|h\| \leq C_m c$ such that $d(h) = g$.

Indeed, assume the contrary. Then we can find a sequence V_1, V_2, \dots of r -normed $\mathbb{Z}[T^{\pm 1}]$ -modules, equipped with norms $\|\cdot\|_i$ on V_i , together with elements $g_i \in \ker(\widehat{V}_i(E_m) \rightarrow \widehat{V}_i(E_{m+1}))$ with $\|g_i\| \leq c$ for some fixed c , such that for each $i = 1, 2, \dots$ there is no $h \in V_i(E_{m-1})$ with $\|h_i\| \leq r^{-2i}c$ such that $d(h_i) = g_i$.

Let V be the direct sum $V_1 \oplus V_2 \oplus \dots$ equipped with the supremum norm; this is again an r -normed $\mathbb{Z}[T^{\pm 1}]$ -module. Moreover, $g = (Tg_1, T^2g_2, \dots)$ defines an element of $\widehat{V}(E_m)$. Using the exactness of the complex for V , we get some $h \in V$ with $d(h) = g$. Then there is some i such that $\|h\| \leq r^{-i}c$, and then the image $T^{-i}h_i \in V_i$ of h satisfies $\|h_i\| \leq r^{-2i}c$ and $d(h_i) = g_i$, which contradicts our assumption.

Now, we also see that for all profinite sets S' , the complex

$$0 \rightarrow \widehat{V}(E_0 \times S') \rightarrow \widehat{V}(E_1 \times S') \rightarrow \dots$$

is exact, and for all $m \geq 0$, there is some constant C_m (independent of V) such that for all $g \in \ker(\widehat{V}(E_m \times S') \rightarrow \widehat{V}(E_{m+1} \times S'))$ with $\|g\| \leq c$, there is some $h \in \widehat{V}(E_{m-1} \times S')$ with $\|h\| \leq C_m c$ such that $d(h) = g$. Indeed, this complex is obtained by completing the complex of locally constant maps from S' into the previous one, which gives the desired bounds by Proposition 8.17.

But this complex computes $R\mathrm{Hom}_{\mathbb{Z}[T^{-1}]}(\mathcal{M}(S, \mathbb{Z}((T))_{r'})[S']/\mathbb{Z}[T^{-1}][S \times S'], V)$, via the resolution $P_\bullet[S']$. This is not quite a resolution by extremally disconnected sets, but one can refine it by one (resolving each term), and use Proposition 8.19. Applying this to $V = M^B$ with its norm, we get the result. \square

REMARK 8.15. Reductions very similar to the ones in this lecture could be done directly in the setting of liquid \mathbb{R} -vector spaces, reducing Theorem 6.6 to the following statement: For any $0 < p' < p \leq 1$ and any p -Banach space V and profinite set S , the map

$$R\mathrm{Hom}_{\mathbb{R}}(\mathcal{M}_{p'}(S), V) \rightarrow V(S)$$

is a quasi-isomorphism. Equivalently, for all $i > 0$,

$$\mathrm{Ext}_{\mathbb{R}}^i(\mathcal{M}_{p'}(S), V) = 0.$$

It is in the proof of this statement that one runs into the issue pointed out after Theorem 6.7.

Appendix to Lecture VIII: Completions of normed abelian groups

DEFINITION 8.16.

- (1) A normed abelian group is an abelian group M equipped with a map

$$\|\cdot\| : M \rightarrow \mathbb{R}_{\geq 0}$$

satisfying $\|0\| = 0$, $\|-m\| = \|m\|$ and $\|m+n\| \leq \|m\| + \|n\|$ for all $m, n \in M$.

- (2) A normed abelian group is separated if $\|m\| = 0$ implies $m = 0$.
 (3) A normed abelian group is complete if it is separated and for every Cauchy sequence (m_0, m_1, \dots) (i.e., $\|m_i - m_j\| \rightarrow 0$ as $i, j \rightarrow \infty$) there is some $m \in M$ with $\|m - m_i\| \rightarrow 0$ as $i \rightarrow \infty$.

As is well-known, the inclusions of complete into separated into all normed abelian groups admit left adjoints: Separation takes M to the quotient of M by $\{m \in M \mid \|m\| = 0\}$; and completion takes M to the quotient of the space of Cauchy sequences (m_0, m_1, \dots) ($\|m_i - m_j\| \rightarrow 0$ as $i, j \rightarrow \infty$) by the space of nullsequences (m_0, m_1, \dots) ($\|m_i\| \rightarrow 0$ as $i \rightarrow \infty$), with the norm defined by declaring $\|(m_0, m_1, \dots)\|$ to be the limit of $\|m_i\|$ as $i \rightarrow \infty$.

We often use the following exactness property:

PROPOSITION 8.17. *Let $M' \xrightarrow{f} M \xrightarrow{g} M''$ be maps of normed abelian groups with $gf = 0$ and such that for all $m \in \ker(M \rightarrow M'')$ there is some $m' \in M'$ with $f(m') = m$ with $\|m'\| \leq C\|m\|$.*

Then $\widehat{M}' \xrightarrow{\widehat{f}} \widehat{M} \xrightarrow{\widehat{g}} \widehat{M}''$ satisfy $\widehat{g}\widehat{f} = 0$ and for all $m \in \ker(\widehat{M} \rightarrow \widehat{M}'')$ and all $\epsilon > 0$ there is some $m' \in \widehat{M}'$ with $\widehat{f}(\widehat{m}') = \widehat{m}$ and $\|m'\| \leq (C + \epsilon)\|m\|$.

PROOF. One can write any element $m \in \widehat{M}$ as a sum $m_0 + m_1 + \dots$ with $m_i \in M$ and $\|m_i\| \leq \epsilon_i$ for $i > 0$ for any given sequence of positive numbers $\epsilon_1 \geq \epsilon_2 \geq \dots$. Indeed, we can inductively choose the m_i so that $\|m - m_0 - \dots - m_i\| \leq \frac{1}{2}\epsilon_{i+1}$, in which case $\|m_i\| \leq \frac{1}{2}(\epsilon_i + \epsilon_{i+1}) \leq \epsilon_i$. Taking the sequence of ϵ_i 's sufficiently small so that $\sum_{i>0} \epsilon_i \leq \frac{\|m\|}{2C}$, we can lift all m_i to m'_i with $\|m'_i\| \leq C\|m_i\|$, and then $m' = m'_0 + m'_1 + \dots$ satisfies

$$\|m'\| \leq \|m'_0\| + C \sum_{i>0} \epsilon_i \leq C\|m_0\| + C \sum_{i>0} \epsilon_i \leq C\|m\| + 2C \sum_{i>0} \epsilon_i \leq (C + \epsilon)\|m\|.$$

□

DEFINITION 8.18. Let M be a normed abelian group. Let \widehat{M} be the condensed abelian group taking any profinite S to the completion of the normed abelian group of locally constant maps from S to M (equipped with the supremum norm).

We note that implicit here is that this actually is a condensed abelian group.

PROPOSITION 8.19. *The condensed abelian group \widehat{M} is canonically identified with the condensed abelian group associated to the topological abelian group \widehat{M}_{top} given by the completion of M equipped with the topology induced by the norm. The norm defines a natural map of condensed sets*

$$\|\cdot\| : \widehat{M} \rightarrow \mathbb{R}_{\geq 0}.$$

Moreover, for any hypercover $S_\bullet \rightarrow S$ of a profinite set S by profinite sets S_i , the complex

$$0 \rightarrow \widehat{M}(S) \rightarrow \widehat{M}(S_0) \rightarrow \widehat{M}(S_1) \rightarrow \dots$$

is exact, and whenever $f \in \ker(\widehat{M}(S_m) \rightarrow \widehat{M}(S_{m+1}))$ with $\|f\| \leq c$, then for any $\epsilon > 0$ there is some $g \in \widehat{M}(S_{m-1})$ with $\|g\| \leq (m+2+\epsilon)c$ such that $d(g) = f$.

PROOF. For the final assertion, follow the proof of [Sch19, Theorem 3.3]: When S and all S_i are finite, the hypercover splits, so a contracting homotopy gives the result with constant $m+2$. In general, write the hypercover as a cofiltered limit of hypercovers of finite sets by finite sets, pass to the filtered colimit, and complete, using Proposition 8.17.

For the identification with the condensed abelian group associated to the topological abelian group \widehat{M}_{top} , note that in the supremum norm any continuous function from S to \widehat{M}_{top} can be approximated by locally constant functions arbitrarily well, and that the space of continuous functions from S to \widehat{M}_{top} is complete with respect to the supremum norm. That $\|\cdot\|$ defines a map of condensed sets $\widehat{M} \rightarrow \mathbb{R}_{\geq 0}$ follows for example from this identification with \widehat{M}_{top} , as the norm is by definition a continuous map $\widehat{M}_{\text{top}} \rightarrow \mathbb{R}_{\geq 0}$. \square

Appendix to Lecture VIII: Derived inverse limits

We recall the following construction of derived inverse limits. Let \mathcal{I} be some index category, and let $F : \mathcal{I} \rightarrow \mathcal{A}$ be a functor to some abelian category \mathcal{A} that admits all colimits and compact projective generators (in particular, infinite products are exact in \mathcal{A}).

We build a cosimplicial object L_F^\bullet of \mathcal{A} whose n -th term is

$$L_F^n = \prod_{i_0 \rightarrow \dots \rightarrow i_n} F(i_n)$$

where the product is over all chains of n composable morphisms $i_0 \rightarrow i_1 \rightarrow \dots \rightarrow i_n$. We note that the index set of the product is equivalently the set of maps $\Delta_{\text{cat}}^n \rightarrow \mathcal{I}$, where Δ_{cat}^n is the category associated to the ordered set $\{0, \dots, n\}$. Moreover, $F(i_n)$ can be understood as the colimit of F restricted to Δ_{cat}^n . With this description, it is clear that a map of simplices $\Delta^n \rightarrow \Delta^m$ induces a map $L_F^n \rightarrow L_F^m$.

Before going on, let us analyze the case that \mathcal{I} has an initial object $i \in \mathcal{I}$. In that case, L_F^\bullet first of all extends to an augmented cosimplicial object $F(i) \rightarrow L_F^\bullet$, and this augmented cosimplicial object has terms

$$\prod_{i=i_{-1} \rightarrow i_0 \rightarrow \dots \rightarrow i_n} F(i_n)$$

for $n = -1, 0, \dots$, which are even functorial in maps $\Delta^{n+1} \rightarrow \Delta^{m+1}$ preserving the initial vertex. Formally (cf. e.g. (the dual of) [Lur09, Lemma 6.1.3.16]), this implies that the complex

$$0 \rightarrow F(i) \rightarrow L_F^0 \rightarrow L_F^1 \rightarrow \dots$$

associated to the augmented cosimplicial object is acyclic. More concretely, this is a direct verification: the contracting homotopy is given by the map $L_F^0 \rightarrow F(i)$ taking the component at i , and the maps $L_F^n \rightarrow L_F^{n-1}$ given by restricting to components with $i = i_0$.

The following proposition is known as the ‘‘Bousfield–Kan formula’’, cf. [BK72, Chapter XI].

PROPOSITION 8.20. *The complex associated to L_F^\bullet by Dold–Kan computes $R\lim_{\mathcal{I}} F$.*

We will prove this only when \mathcal{I} is cofiltered, which is the argument that is used in the main text.

PROOF. It is clear that $\lim_{\mathcal{I}} F$ is the kernel of $L_F^0 \rightarrow L_F^1$. Picking an injective resolution of F (in the category of functors $\mathcal{I} \rightarrow \mathcal{A}$) and using exactness of infinite products, we get a natural map from $R\lim_{\mathcal{I}} F$ to L_F^\bullet . We claim that this is an equivalence.

If \mathcal{I} is cofiltered, we can write \mathcal{I} as a filtered union of finite subcategories \mathcal{I}_j that contain an initial object. Then both sides are equal to the derived cofiltered limit of their variants for the \mathcal{I}_j . Thus, we can assume that \mathcal{I} is finite and has an initial object i . But in this case, we have proved the exactness above. \square

9. Lecture IX: End of proof

Recall that in the last lecture, we reduced Theorem 6.11 to the following result.

THEOREM 9.1. *Fix radii $1 > r' > r > 0$. Then for all r -normed $\mathbb{Z}[T^{\pm 1}]$ -modules V and all profinite sets S , the map*

$$R\mathrm{Hom}_{\mathbb{Z}[T^{-1}]}(\mathcal{M}(S, \mathbb{Z}((T))_{r'}), \widehat{V}) \rightarrow \widehat{V}(S)$$

is a quasi-isomorphism.

We note that actually the case of $\widehat{V} = \mathbb{Z}((T))_r^B$ itself seems to carry all essential difficulty: We believe that any argument one can give in that case will work in general.

In that sense, we have simplified the target of the $R\mathrm{Hom}$ as far as possible. It is time to understand the source. In the last lecture, we observed that the theorem automatically implies a more precise version bounding the norms of preimages of differentials. Our proof will actually go back to such explicit bounds, but for explicit resolutions of the source, which we will now construct. Write

$$\overline{\mathcal{M}}_{r'}(S) := \mathcal{M}(S, \mathbb{Z}((T))_{r'}) / \mathbb{Z}[T^{-1}][S].$$

This is, as a condensed abelian group (but not as $\mathbb{Z}[T^{-1}]$ -module), a direct summand $\mathcal{M}(S, T\mathbb{Z}[[T]]_{r'})$ of $\mathcal{M}(S, \mathbb{Z}((T))_{r'})$, allowing only positive powers of T .¹⁶ To see this, use Proposition 2.1 to see that the contribution to $\mathcal{M}(S, \mathbb{Z}((T))_{r'})$ from nonpositive powers of T is exactly $\mathbb{Z}[T^{-1}][S]$. In particular, we can write

$$\overline{\mathcal{M}}_{r'}(S) = \bigcup_{c>0} \overline{\mathcal{M}}_{r'}(S)_{\leq c},$$

where

$$\overline{\mathcal{M}}_{r'}(S)_{\leq c} = \varprojlim_i \overline{\mathcal{M}}_{r'}(S_i)_{\leq c}$$

when writing S as an inverse limit of finite sets S_i , and for finite S

$$\overline{\mathcal{M}}_{r'}(S)_{\leq c} = \left\{ \left(\sum_{n \geq 1} a_{n,s} T^n \right)_s \mid \sum_{n \geq 1, s \in S} |a_{n,s}| r^n \leq c \right\}.$$

We need to resolve this explicitly as a condensed $\mathbb{Z}[T^{-1}]$ -module. The Breen-Deligne resolution gives us the resolution

$$\dots \rightarrow \mathbb{Z}[\overline{\mathcal{M}}_{r'}(S)^2] \rightarrow \mathbb{Z}[\overline{\mathcal{M}}_{r'}(S)] \rightarrow \overline{\mathcal{M}}_{r'}(S) \rightarrow 0$$

as condensed $\mathbb{Z}[T^{-1}]$ -modules; as in the appendix, we assume (for notational convenience) that each term is of the form $\mathbb{Z}[\overline{\mathcal{M}}_{r'}(S)^a]$ (instead of some finite direct sum of such). Moreover, each term $\mathbb{Z}[\overline{\mathcal{M}}_{r'}(S)^a]$ admits a two-term resolution

$$0 \rightarrow \mathbb{Z}[T^{-1}][\overline{\mathcal{M}}_{r'}(S)^a] \xrightarrow{T^{-1} - [T^{-1}]} \mathbb{Z}[T^{-1}][\overline{\mathcal{M}}_{r'}(S)^a] \rightarrow \mathbb{Z}[\overline{\mathcal{M}}_{r'}(S)^a] \rightarrow 0$$

¹⁶At this point, we critically use that we chose $\mathbb{Z}[T^{-1}]$ as our base ring; already taking $\mathbb{Z}[T^{\pm 1}]$ would destroy this argument.

that is functorial in the $\mathbb{Z}[T^{-1}]$ -module $\mathbb{Z}[\overline{\mathcal{M}}_{r'}(S)^a]$. This gives a resolution of $\overline{\mathcal{M}}_{r'}(S)$ by the double complex

$$\begin{array}{ccc} \dots & \longrightarrow & \mathbb{Z}[T^{-1}][\overline{\mathcal{M}}_{r'}(S)^2] & \longrightarrow & \mathbb{Z}[T^{-1}][\overline{\mathcal{M}}_{r'}(S)] \\ & & \downarrow T^{-1}-[T^{-1}] & & \downarrow T^{-1}-[T^{-1}] \\ \dots & \longrightarrow & \mathbb{Z}[T^{-1}][\overline{\mathcal{M}}_{r'}(S)^2] & \longrightarrow & \mathbb{Z}[T^{-1}][\overline{\mathcal{M}}_{r'}(S)]. \end{array}$$

Unfortunately, the terms are not of the form $\mathbb{Z}[T^{-1}][E_i]$ with profinite E_i . To remedy this, we write the double complex as the filtered colimit of the double complexes

$$(9.1) \quad \begin{array}{ccc} \dots & \longrightarrow & \mathbb{Z}[T^{-1}][\overline{\mathcal{M}}_{r'}(S)_{\leq r'c_1c}^2] & \longrightarrow & \mathbb{Z}[T^{-1}][\overline{\mathcal{M}}_{r'}(S)_{\leq r'c}] \\ & & \downarrow T^{-1}-[T^{-1}] & & \downarrow T^{-1}-[T^{-1}] \\ \dots & \longrightarrow & \mathbb{Z}[T^{-1}][\overline{\mathcal{M}}_{r'}(S)_{\leq c_1c}^2] & \longrightarrow & \mathbb{Z}[T^{-1}][\overline{\mathcal{M}}_{r'}(S)_{\leq c}] \end{array}$$

for varying $c > 0$, where the constants $c_1, \dots > 0$ are fixed and chosen so that the transition maps are well-defined, cf. Lemma 9.11 in the appendix to this lecture.

It is easy to understand what happens to the vertical part under mapping to V :

LEMMA 9.2. *For any r -normed $\mathbb{Z}[T^{\pm 1}]$ -module V , any $c > 0$ and any a , the map*

$$\widehat{V}(\overline{\mathcal{M}}_{r'}(S)_{\leq c}^a) \xrightarrow{T^{-1}-[T^{-1}]^*} \widehat{V}(\overline{\mathcal{M}}_{r'}(S)_{\leq r'c}^a)$$

is surjective, has norm bounded by $r^{-1} + 1$, and for any $f \in \widehat{V}(\overline{\mathcal{M}}_{r'}(S)_{\leq r'c}^a)$ and $\epsilon > 0$ there is some $g \in \widehat{V}(\overline{\mathcal{M}}_{r'}(S)_{\leq c}^a)$ with $f(x) = T^{-1}g(x) - g(T^{-1}x)$ and $\|g\| \leq \frac{r}{1-r}(1 + \epsilon)\|f\|$.

PROOF. Given $f : \overline{\mathcal{M}}_{r'}(S)_{\leq r'c}^a \rightarrow \widehat{V}$, choose an extension to a map $\tilde{f} : \overline{\mathcal{M}}_{r'}(S)^a \rightarrow \widehat{V}$ with $\|\tilde{f}\| \leq (1 + \epsilon)\|f\|$. Such an extension exists: By induction (and using a sequence of ϵ_n 's with $\prod_n(1 + \epsilon_n) \leq 1 + \epsilon$), it suffices to see that for any closed immersion $A \subset B$ of profinite sets and a map $f_A : A \rightarrow \widehat{V}$, there is an extension $f_B : B \rightarrow \widehat{V}$ of f_A with $\|f_B\| \leq (1 + \epsilon)\|f_A\|$. To see this, write f_A as a (fast) convergent sum of maps that factor over a finite quotient of A ; for maps factoring over a finite quotient of A , the extension is clear (and can be done in a norm-preserving way), as any map from A to a finite set can be extended to a map from B to the same finite set.

Given \tilde{f} , we can now define $g : \overline{\mathcal{M}}_{r'}(S)_{\leq c}^a$ by

$$g(x) = T\tilde{f}(x) + T^2\tilde{f}(T^{-1}x) + \dots + T^{n+1}\tilde{f}(T^{-n}x) + \dots \in \widehat{V};$$

then $\|g\| \leq \frac{r}{1-r}\|\tilde{f}\| \leq \frac{r}{1-r}(1 + \epsilon)\|f\|$. □

Let $\widehat{V}(\overline{\mathcal{M}}(S)_{\leq c})^{T^{-1}} \subset \widehat{V}(\overline{\mathcal{M}}(S)_{\leq c})$ be the kernel of this map, with the subspace norm. For varying c , we now get varying normed complexes. We will need to use the following qualitative notion of exactness.

DEFINITION 9.3. For each sufficiently large c (i.e. all $c \geq c_0$ for some $c_0 > 0$), let

$$C_c^\bullet : C_c^0 \rightarrow C_c^1 \rightarrow \dots$$

be a complex of normed abelian groups, and for $c' > c$, let $\text{res}_{c',c}^i : C_{c'}^\bullet \rightarrow C_c^\bullet$ be a map of complexes, satisfying the obvious associativity condition. This datum is admissible if all differentials and maps $\text{res}_{c',c}^i$ are norm-nonincreasing.

For integers $m \geq 0$ and constants $k > 0$, $c'_0 > 0$, the datum $(C_c^\bullet)_c$ is $\leq k$ -exact in degrees $\leq m$ and for $c \geq c'_0$ if the following condition is satisfied. For all $c \geq c'_0$ and all $x \in C_{kc}^i$ with $i \leq m$ there is some $y \in C_c^{i-1}$ (which is defined to be 0 when $i = 0$) such that

$$\|\text{res}_{kc,c}^i(x) - d_c^{i-1}(y)\|_{C_c^i} \leq k \|d_{kc}^i(x)\|_{C_{kc}^{i+1}}.$$

We apply this to

$$C_c^\bullet : \widehat{V}(\overline{\mathcal{M}}_{r'}(S)_{\leq c})^{T^{-1}} \rightarrow \widehat{V}(\overline{\mathcal{M}}_{r'}(S)_{\leq c_1 c}^2)^{T^{-1}} \rightarrow \dots$$

given by mapping (9.1) into V and using Lemma 9.2. Now we state the following result.

THEOREM 9.4. *Fix radii $1 > r' > r > 0$. For any m there is some k and c_0 such that for all profinite sets S and all r -normed $\mathbb{Z}[T^{\pm 1}]$ -modules V , the system of complexes*

$$C_c^\bullet : \widehat{V}(\overline{\mathcal{M}}_{r'}(S)_{\leq c})^{T^{-1}} \rightarrow \widehat{V}(\overline{\mathcal{M}}_{r'}(S)_{\leq c_1 c}^2)^{T^{-1}} \rightarrow \dots$$

is $\leq k$ -exact in degrees $\leq m$ for $c \geq c_0$.

Let us first check that this implies Theorem 9.1.

THEOREM 9.4 IMPLIES THEOREM 9.1. By the preceding discussion, one can compute

$$R \text{Hom}_{\mathbb{Z}[T^{-1}]}(\overline{\mathcal{M}}_{r'}(S), \widehat{V})$$

as the derived inverse limit of C_c^\bullet over all $c > 0$; equivalently, all $c \geq c_0$. Theorem 9.4 implies that for any $m \geq 0$ the pro-system of cohomology groups $H^m(C_c^\bullet)$ is pro-zero (as $H^m(C_{kc}^\bullet) \rightarrow H^m(C_c^\bullet)$ is zero). Thus, the derived inverse limit vanishes, as desired. \square

We remark that Theorem 9.4 reduces formally to the case that S is finite; we make this reduction. We will prove Theorem 9.4 by induction on m . Unfortunately, the induction requires us to prove a stronger statement. This goes as follows. Consider any polyhedral lattice Λ , by which we mean a finite free abelian group equipped with a norm $\|\cdot\|_\Lambda : \Lambda \otimes \mathbb{R} \rightarrow \mathbb{R}$ (so $\Lambda \otimes \mathbb{R}$ is a Banach space) that is given by the supremum of finitely many linear functions on Λ with rational coefficients; equivalently, the “unit ball” $\{\lambda \in \Lambda \otimes \mathbb{R} \mid \|\lambda\|_\Lambda \leq 1\}$ is a rational polyhedron.

Endow $\text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))$ with the subspaces

$$\text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c} = \{f : \Lambda \rightarrow \overline{\mathcal{M}}_{r'}(S) \mid \forall x \in \Lambda, f(x) \leq \overline{\mathcal{M}}_{r'}(S)_{\leq c\|x\|}\}.$$

As Λ is polyhedral, it is enough to check the given condition for finitely many x .

We can then define double complexes like (9.1). Lemma 9.2 stays true with the same constants. Now we claim the following generalization of Theorem 9.4.

THEOREM 9.5. *Fix radii $1 > r' > r > 0$. For any m there is some k such that for all polyhedral lattices Λ there is a constant $c_0(\Lambda) > 0$ such that for all finite sets S and all r -normed $\mathbb{Z}[T^{\pm 1}]$ -modules V , the system of complexes*

$$C_{\Lambda,c}^\bullet : \widehat{V}(\text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c})^{T^{-1}} \rightarrow \widehat{V}(\text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c_1 c}^2)^{T^{-1}} \rightarrow \dots$$

is $\leq k$ -exact in degrees $\leq m$ for $c \geq c_0(\Lambda)$.

We note that the constants c_1, c_2, \dots implicit in the choice of the complex are chosen once and for all (after fixing r and r'), and it can be ensured that the transition maps in the complex are norm-nonincreasing. Indeed, with the c_i chosen as in Lemma 9.11, the maps

$$\widehat{V}(\mathrm{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c_i c}^{a_i}) \rightarrow \widehat{V}(\mathrm{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c_{i+1} c}^{a_{i+1}})$$

will have bounded norm, independently of V (as they are a certain universal finite sum of maps induced by maps between the profinite sets in paranthesis, each of which induces a map of norm bounded by 1), so on the subspace of T^{-1} -invariants, one can shrink the norm down to 1 by shrinking c_{i+1} . We make and fix this choice of the c_i for the statement of Theorem 9.5, and the rest of the proof.

PROPOSITION 9.6. *Fix an integer $m \geq 0$ and a constant k . Then there exists an $\epsilon > 0$ and a constant k_0 , depending (only) on k and m , with the following property.*

Consider an admissible system of double complexes $M_c^{p,q}$, $p, q \geq 0$, $c \geq c_0$, of normed abelian groups as well as some $k' \geq k_0$ and some $H > 0$, such that

$$\begin{array}{ccccccc} M_c^{0,0} & \xrightarrow{d_c^{0,0}} & M_c^{0,1} & \xrightarrow{d_c^{0,1}} & M_c^{0,2} & \xrightarrow{d_c^{0,2}} & \dots \\ \downarrow d_c^{0,0} & & \downarrow d_c^{0,1} & & \downarrow d_c^{0,2} & & \\ M_c^{1,0} & \xrightarrow{d_c^{1,0}} & M_c^{1,1} & \xrightarrow{d_c^{1,1}} & M_c^{1,2} & \xrightarrow{d_c^{1,2}} & \dots \\ \downarrow d_c^{1,0} & & \downarrow d_c^{1,1} & & \downarrow d_c^{1,2} & & \\ M_c^{2,0} & \xrightarrow{d_c^{2,0}} & M_c^{2,1} & \xrightarrow{d_c^{2,1}} & \dots & & \\ \downarrow d_c^{2,0} & & \downarrow d_c^{2,1} & & & & \\ \vdots & & \vdots & & & & \end{array}$$

- (1) for $j = 0, \dots, m$, the columns $M_c^{p,j}$ are $\leq k$ -exact in degrees $\leq m$ for $c \geq c_0$;
- (2) for $i = 0, \dots, m+1$, the rows $M_c^{i,q}$ are $\leq k$ -exact in degrees $\leq m-1$ for $c \geq c_0$;
- (3) for $q = 0, \dots, m$ and $c \geq c_0$, there is a map $h_{k'_c}^q : M_{k'_c}^{0,q+1} \rightarrow M_c^{1,q}$ with

$$\|h_{k'_c}^q(x)\|_{M_c^{1,q}} \leq H \|x\|_{M_{k'_c}^{0,q+1}}$$

for all $x \in M_c^{0,q+1}$, and such that for all $c \geq c_0$ and $x \in M_{k'^2 c}^{0,q}$, one has

$$(9.2) \quad \|\mathrm{res}_{k'^2 c, k'_c}^{1,q}(d^{0,q}(x)) \pm h_{k'^2 c}^q(d_{k'^2 c}^{0,q}(x)) \pm d_{k'_c}^{1,q-1}(h_{k'^2 c}^{q-1}(x))\|_{M_{k'_c}^{1,q}} \leq \epsilon \|\mathrm{res}_{k'^2 c, c}^{0,q}(x)\|_{M_c^{0,q}}.$$

Then the first row is $\leq \max(k'^2, 2k_0 H)$ -exact in degrees $\leq m$ for $c \geq c_0$.

We note that the bound on the homotopy is of a peculiar nature, in that the bound only depends on a deep restriction of x .

PROOF. First, we treat the case $m = 0$. If $m = 0$, we claim that one can take $\epsilon = \frac{1}{2k}$ and $k_0 = k$. We have to prove exactness at the first step. Let $x_{k'^2 c} \in M_{k'^2 c}^{0,0}$ and denote $x_{k'_c} = \mathrm{res}_{k'^2 c, k'_c}^{0,0}(x)$ and $x_c = \mathrm{res}_{k'^2 c, c}^{0,0}(x)$. Then by assumption (1) (and $k' \geq k$), we have

$$\|x_c\|_{M_c^{0,0}} \leq k \|d_{k'_c}^{0,0}(x_{k'_c})\|_{M_{k'_c}^{1,0}}.$$

On the other hand, by (3),

$$\|\text{res}_{k'/2c, k'c}^{1,0}(d_{k'/2c}^{0,0}(x)) \pm h_{k'/2c}^0(d_{k'/2c}^{\prime 0,0}(x))\|_{M_{k'/c}^{1,0}} \leq \epsilon \|x_c\|_{M_c^{0,0}}.$$

In particular, noting that $\text{res}_{k'/2c, k'c}^{1,0}(d_{k'/2c}^{0,0}(x)) = d_{k'/c}^{0,0}(x_{k'c})$, we get

$$\|x_c\|_{M_c^{0,0}} \leq k \|d_{k'/c}^{0,0}(x_{k'c})\|_{M_{k'/c}^{1,0}} \leq k\epsilon \|x_c\|_{M_c^{0,0}} + kH \|d_{k'/2c}^{\prime 0,0}(x)\|_{M_{k'/2c}^{0,1}}.$$

Thus, taking $\epsilon = \frac{1}{2k}$ as promised, this implies

$$\|x_c\|_{M_c^{0,0}} \leq 2kH \|d_{k'/2c}^{\prime 0,0}(x)\|_{M_{k'/2c}^{0,1}}.$$

This gives the desired $\leq \max(k'^2, 2k_0H)$ -exactness in degrees $\leq m$ for $c \geq c_0$.

Now we argue by induction on m . Consider the complex $N^{p,q}$ given by $M^{p,q+1}$ for $q \geq 1$ and $N^{p,0} = M^{p,1}/M^{p,0}$, equipped with the quotient norm. Using the normed version of the snake lemma, Proposition 9.10 in the appendix to this lecture, one checks that this satisfies the assumptions for $m-1$, with k replaced by $k^3 + k$. \square

Finally, we can prove the key combinatorial lemma, ensuring that any element of $\text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))$ can be decomposed into N elements whose norm is roughly $\frac{1}{N}$ of the original element. As preparation, we have the following simple result.

LEMMA 9.7. *Let Λ be a finite free abelian group, let N be a positive integer, and let $\lambda_1, \dots, \lambda_m \in \Lambda$ be elements. Then there is a finite subset $A \subset \Lambda^\vee$ such that for all $x \in \Lambda^\vee = \text{Hom}(\Lambda, \mathbb{Z})$ there is some $x' \in A$ such that $x - x' \in N\Lambda^\vee$ and for all $i = 1, \dots, m$, the numbers $x'(\lambda_i)$ and $(x - x')(\lambda_i)$ have the same sign, i.e. are both nonnegative or both nonpositive.*

PROOF. It suffices to prove the statement for all x such that $\lambda_i(x) \geq 0$ for all i ; indeed, applying this variant to all $\pm\lambda_i$, one gets the full statement.

Thus, consider the submonoid $\Lambda_+^\vee \subset \Lambda^\vee$ of all x that pair nonnegatively with all λ_i . This is a finitely generated monoid by standard results; let y_1, \dots, y_M be a set of generators. Then we can take for A all sums $n_1y_1 + \dots + n_My_M$ where all $n_j \in \{0, \dots, N-1\}$. \square

Now we have the key lemma:

LEMMA 9.8. *Let Λ be a polyhedral lattice. Then for all positive integers N there is a constant d such that for all $c > 0$ one can write any $x \in \text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c}$ as*

$$x = x_1 + \dots + x_N$$

where all $x_i \in \text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c/N+d}$.

PROOF. Pick $\lambda_1, \dots, \lambda_m \in \Lambda$ generating the norm. We fix a finite subset $A \subset \Lambda^\vee$ satisfying the conclusion of the previous lemma. Write

$$x = \sum_{n \geq 1, s \in S} x_{n,s} T^n[s]$$

with $x_{n,s} \in \Lambda^\vee$. Then we can decompose

$$x_{n,s} = Nx_{n,s}^0 + x_{n,s}^1$$

where $x_{n,s}^1 \in A$ and we have the same-sign property of the last lemma. Letting $x^0 = \sum_{n \geq 1, s \in S} x_{n,s}^0 T^n[s]$, we get a decomposition

$$x = Nx^0 + \sum_{a \in A} ax_a$$

with $x_a \in \overline{\mathcal{M}}_{r'}(S)$ (with the property that in the basis given by the $T^n[s]$, all coefficients are 0 or 1). Crucially, we know that for all $i = 1, \dots, m$, we have

$$\|x(\lambda_i)\| = N\|x^0(\lambda_i)\| + \sum_{a \in A} |a(\lambda_i)| \|x_a(\lambda_i)\|$$

by using the same sign property of the decomposition.

Using this decomposition of x , we decompose each term into N summands. This is trivial for the first term Nx^0 , and each summand of the second term reduces to the similar problem for $\Lambda = \mathbb{Z}$. In that case, one can take $d = 1$, as follows by decomposing any sum with terms of size at most 1 into N such partial sums whose sums differ by at most 1. (It follows that in general one can take for d the supremum over all i of $\sum_{a \in A} |a(\lambda_i)|$.) \square

PROOF OF THEOREM 9.5. We argue by induction on m , so assume the result for $m - 1$ (this is no assumption for $m = 0$, so we do not need an induction start). This gives us some $k > 1$ for which the statement of Theorem 9.5 holds true for $m - 1$; if $m = 0$, simply take any $k > 1$. In the proof below, we will increase k further in a way that depends only on m and r . After this modified choice of k , we fix ϵ and k_0 as provided by Proposition 9.6. Moreover, we let k' be the supremum of k_0 and the c'_i from Lemma 9.12 (and 9.13) for $i = 0, \dots, m$. Finally, choose a positive integer b so that $2k'(\frac{r}{r'})^b \leq \epsilon$, and let N be the minimal power of 2 that satisfies

$$k'/N \leq (r')^b.$$

Then in particular $r^b N \leq \frac{2}{k'} (\frac{r}{r'})^b \leq \epsilon$.

We consider the diagonal embedding

$$\Lambda \hookrightarrow \Lambda' = \Lambda^N,$$

where we endow Λ' with the norm

$$\|(\lambda_1, \dots, \lambda_N)\|_{\Lambda'} = \frac{1}{N} (\|\lambda_1\|_{\Lambda} + \dots + \|\lambda_N\|_{\Lambda}).$$

For any $m \geq 1$, let $\Lambda'^{(m)}$ be given by $\Lambda'^m / \Lambda \otimes (\mathbb{Z}^m)_{\sum=0}$; then $\Lambda'^{(\bullet)}$ is cosimplicial polyhedral lattice, the Čech conerve of $\Lambda \rightarrow \Lambda'$. For $m = 0$, we set $\Lambda'^{(0)} = \Lambda$. It is clear that all of these are polyhedral lattices.

In particular, for any $c > 0$, we have

$$\mathrm{Hom}(\Lambda'^{(m)}, \overline{\mathcal{M}}_{r'}(S))_{\leq c} = \mathrm{Hom}(\Lambda', \overline{\mathcal{M}}_{r'}(S))_{\leq c}^{m / \mathrm{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c}},$$

the m -fold fibre product of $\mathrm{Hom}(\Lambda', \overline{\mathcal{M}}_{r'}(S))_{\leq c}$ over $\mathrm{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c}$; and

$$\mathrm{Hom}(\Lambda', \overline{\mathcal{M}}_{r'}(S))_{\leq c} = \mathrm{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c/N}^N,$$

with the map to $\mathrm{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c}$ given by the sum map.

Consider the collection of double complexes $C_{\Lambda'^{(\bullet)}, c}^{\bullet}$ associated to this cosimplicial polyhedral lattice by Dold-Kan. Up to rescaling the norms in the complex for $\Lambda'^{(m)}$ by a universal constant (something like $(m + 2)!$), the differentials are strictly compatible with norms (as they are an

alternating sum of $m + 1$ face maps, all of which are of norm ≤ 1), so this collection of normed double complexes is admissible. By induction, the first condition of Proposition 9.6 is satisfied for all $c \geq c_0$ with c_0 large enough (depending on Λ but not V or S). By Lemma 9.8, and noting that $\text{Hom}(\Lambda'^{(\bullet)}, \overline{\mathcal{M}}_{r'}(S))_{\leq c}$ is the Čech nerve of

$$\text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c/N}^N \xrightarrow{\Sigma} \text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c},$$

also the second condition is satisfied, with k the maximum of the previous k and some constant depending only on m and r , provided we take c_0 large enough so that $(k - 1)r'c_i c_0/N$ is at least the d of Lemma 9.8 for all $i = 0, \dots, m$ (so this choice of c_0 again depends on Λ). Indeed, then one can splice a surjection of profinite sets between the maps

$$\text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c_i c/N}^{Na} \rightarrow \text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c_i c}^a$$

and

$$\text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq k c_i c/N}^{Na} \rightarrow \text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq k c_i c}^a,$$

and so the transition map between the columns of that double complex factors over a similar complex arising from a simplicial hypercover of profinite sets, so the constants are bounded by Proposition 8.19, Lemma 9.2, and Proposition 9.10. At this point, we have finalized our choice of k (and, as promised, this choice depended only on m and r), and so we also finalized the constants ϵ , k' and N from the first paragraph of the proof.

Finally, to check the third condition, we use Lemma 9.13 to find, in degrees $\leq m$, a homotopy between the two maps from the first row

$$\widehat{V}(\text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c})^{T^{-1}} \rightarrow \widehat{V}(\text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c_1 c}^2)^{T^{-1}} \rightarrow \dots$$

to the second row

$$\widehat{V}(\text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c/N}^N)^{T^{-1}} \rightarrow \widehat{V}(\text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c_1 c/N}^{2N})^{T^{-1}} \rightarrow \dots$$

respectively induced by the addition $\text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c/N}^N \rightarrow \text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c}$ (which is the map that forms part of the double complex), and the map that is the sum of the N maps induced by the N projection maps

$$\text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c/N}^N \rightarrow \text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c/N} \subset \text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c}.$$

By Lemma 9.13, we can find this homotopy between the complex for $k'c$ and the complex for c , by our choice of $k' \geq c'_i$ for $i = 0, \dots, m$. As N is fixed, the homotopy is the universal homotopy from Lemma 9.13, and in particular its norm is bounded by some universal constant H .

Finally, it remains to establish the estimate (9.2) on the homotopic map. We note that this takes $x \in \widehat{V}(\text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq k'^2 c_i c}^{a_i})^{T^{-1}}$ (with $i = q$ in the notation of (9.2)) to the element

$$y \in \widehat{V}(\text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq k' c_i c/N}^{Na_i})^{T^{-1}}$$

that is the sum of the N pullbacks along the N projection maps $\text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq k' c_i c/N}^{Na_i} \rightarrow \text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq k'^2 c_i c}^{a_i}$. We note that these actually take image in $\text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c_i c}^{a_i}$ as $N \geq k'$, so this actually gives a well-defined map

$$\widehat{V}(\text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c_i c}^{a_i})^{T^{-1}} \rightarrow \widehat{V}(\text{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq k' c_i c/N}^{Na_i})^{T^{-1}}.$$

We need to see that this map is of norm $\leq \epsilon$. Now note that by our choice of N , we actually have $k'c_i/N \leq (r')^b c_i c$, so this can be written as the composite of the restriction map

$$\widehat{V}(\mathrm{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq c_i c}^{a_i})^{T^{-1}} \rightarrow \widehat{V}(\mathrm{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq (r')^b c_i c}^{a_i})^{T^{-1}}$$

and

$$\widehat{V}(\mathrm{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq (r')^b c_i c}^{a_i})^{T^{-1}} \rightarrow \widehat{V}(\mathrm{Hom}(\Lambda, \overline{\mathcal{M}}_{r'}(S))_{\leq k'c_i c/N}^{Na_i})^{T^{-1}}.$$

The first map has norm exactly r^b , by T^{-1} -invariance, and as multiplication by T scales the norm with a factor of r on \widehat{V} .¹⁷ The second map has norm at most N (as it is a sum of N maps of norm ≤ 1). Thus, the total map has norm $\leq r^b N$. But by our choice of N , we have $r^b N \leq \epsilon$, giving the result.

Thus, we can apply Proposition 9.6, and get the desired $\leq \max(k'^2, 2k_0 H)$ -exactness in degrees $\leq m$ for $c \geq c_0$, where k' , k_0 and H were defined only in terms of k , m , r' and r , while c_0 depends on Λ (but not on V or S). This proves the inductive step. \square

QUESTION 9.9. Can one make the constants explicit, and how large are they?¹⁸ Modulo the Breen-Deligne resolution, all the arguments give in principle explicit constants; and actually the proof of the existence of the Breen-Deligne resolution should be explicit enough to ensure the existence of bounds on the c_i and c'_i .

This completes the proof of all results announced so far.

¹⁷Here is where we use $r' > r$, ensuring different scaling behaviour of the norm on source and target.

¹⁸A back of the envelope calculation seems to suggest that k is roughly doubly exponential in m , and that N has to be taken of roughly the same magnitude.

Appendix to Lecture IX: Some normed homological algebra

In this appendix, we gather a few results about homological algebra with normed abelian groups, the proofs of which are just obtained by keeping track of constants in the standard proofs.

PROPOSITION 9.10. *Let M_c^\bullet and M'_c^\bullet be two admissible collections of complexes of normed abelian groups, where $c \geq c_0$. Let $f_c^\bullet : M_c^\bullet \rightarrow M'_c^\bullet$ be a collection of maps between these collections of complexes that is strictly compatible with the norm and commutes with restriction maps, and assume that it satisfies*

$$\|\text{res}_{k^3c}^i(x)\|_{M_c^i} \leq k \|f_{k^3c}^i(x)\|_{M_{k^3c}^i}$$

for all $i = 0, \dots, m+1$ and all $x \in M_{k^3c}^i$. Let $N_c^\bullet = M'_c^\bullet / M_c^\bullet$ be the collection of quotient complexes, with the quotient norm; this is again an admissible collection of complexes.

Assume that M_c^\bullet and M'_c^\bullet are $\leq k$ -exact in degrees $\leq m$ for $c \geq c_0$. Then N_c^\bullet is $\leq (k^3 + k)$ -exact in degrees $\leq m - 1$ for $c \geq c_0$.

PROOF. Let $n_{k^3c}^i \in N_{k^3c}^i$ for $i \leq m - 1$, with image $n_{k^3c}^{i+1} \in N_{k^3c}^{i+1}$. We will change the subscript when applying restriction maps. Pick any preimage $m_{k^3c}^i \in M_{k^3c}^i$ of $n_{k^3c}^i$, and let $m_{k^3c}^{i+1} \in M_{k^3c}^{i+1}$ be its image. Then we can write

$$m_{k^3c}^{i+1} = f_{k^3c}^{i+1}(m_{k^3c}^{i+1}) + m_{k^3c}^{i+1}$$

with $\|m_{k^3c}^{i+1}\|_{M_{k^3c}^{i+1}} \leq C := \|n_{k^3c}^{i+1}\|_{N_{k^3c}^{i+1}}$ (as we used quotient norms).

Let $m_{k^3c}^{i+2} \in M_{k^3c}^{i+2}$ be the image of $m_{k^3c}^{i+1}$. Applying the differential to the last displayed equation, and using that this kills $m_{k^3c}^{i+1}$, and that $f_{k^3c}^\bullet$ is a map of complexes, we see that

$$f_{k^3c}^{i+2}(m_{k^3c}^{i+2}) = -m_{k^3c}^{i+2},$$

where similarly $m_{k^3c}^{i+2}$ is the differential of $m_{k^3c}^{i+1}$. We get

$$\begin{aligned} \|m_{k^3c}^{i+2}\|_{M_{k^3c}^{i+2}} &\leq k \|f_{k^3c}^{i+2}(m_{k^3c}^{i+2})\|_{M_{k^3c}^{i+2}} = k \|m_{k^3c}^{i+2}\|_{M_{k^3c}^{i+2}} \\ &\leq k \|m_{k^3c}^{i+1}\|_{M_{k^3c}^{i+1}} \leq kC. \end{aligned}$$

On the other hand, we can find some $m_{kc}^i \in M_{kc}^i$ such that

$$\|m_{kc}^{i+1} - d_{kc}^i(m_{kc}^i)\| \leq k \|m_{k^3c}^{i+2}\|_{M_{k^3c}^{i+2}} \leq k^2C.$$

Now let $m_{kc,\text{new}}^i = m_{kc}^i - f_{kc}^i(m_{kc}^i) \in M_{kc}^i$; this is a lift of n_{kc}^i . Then the image $m_{kc,\text{new}}^{i+1}$ in M_{kc}^{i+1} satisfies

$$m_{kc,\text{new}}^{i+1} = m_{kc}^{i+1} - f_{kc}^{i+1}(m_{kc}^{i+1}) + f_{kc}^{i+1}(m_{kc}^{i+1} - d_{kc}^i(m_{kc}^i)) = m_{kc}^{i+1} + f_{kc}^{i+1}(m_{kc}^{i+1} - d_{kc}^i(m_{kc}^i)).$$

In particular,

$$\|m_{kc,\text{new}}^{i+1}\|_{M_{kc}^{i+1}} \leq C + k^2C.$$

Now we can find $m_c^{i-1} \in M_c^{i-1}$ such that

$$\|m_{c,\text{new}}^i - d_c^{i-1}(m_c^{i-1})\|_{M_c^i} \leq k \|m_{kc,\text{new}}^{i+1}\|_{M_{kc}^{i+1}} \leq (k^3 + k)C.$$

In particular, letting $n_c^{i-1} \in N_c^{i-1}$ be the image of m_c^{i-1} , we get

$$\|n_c^i - d_{N,c}^{i-1}(n_c^{i-1})\|_{N_c^i} \leq (k^3 + k)C = (k^3 + k)\|n_{k^3c}^{i+1}\|.$$

This gives exactly the desired result. \square

We need the following results about the Breen-Deligne resolution for normed abelian groups. Let us consider here abelian groups M (in any topos) equipped with an increasing filtration $M_{\leq c} \subset M$ by subobjects indexed by the positive real numbers, such that $0 \in M_{\leq c}$, $-M_{\leq c} = M_{\leq c}$ and $M_{\leq c} + M_{\leq c'} \subset M_{\leq c+c'}$; we need no further conditions. Let us call these pseudo-normed abelian groups.

Fix a choice of a functorial Breen-Deligne resolution

$$C(M) : \dots \rightarrow \mathbb{Z}[M^{a_i}] \rightarrow \dots \rightarrow \mathbb{Z}[M^{a_1}] \rightarrow \mathbb{Z}[M^{a_0}] \rightarrow M \rightarrow 0$$

of an abelian group M ; purely for notational convenience, we can and do assume that each term is of the form $\mathbb{Z}[M^{a_i}]$ (as opposed to a finite direct sum of such). The possibility of doing this follows from the proof of [Sch19, Theorem 4.10], noting that a functor of the form $A \mapsto \mathbb{Z}[A^n] \oplus \mathbb{Z}[A^m]$ admits a surjection from the functor $A \mapsto \mathbb{Z}[A^{n+m}]$.

LEMMA 9.11. *There are universal constants $c_0 = 1, c_1, c_2, \dots$ so that the Breen-Deligne resolution admits the subcomplex*

$$C(M)_{\leq c} : \dots \rightarrow \mathbb{Z}[M_{\leq c_i c}^{a_i}] \rightarrow \dots \rightarrow \mathbb{Z}[M_{\leq c_1 c}^{a_1}] \rightarrow \mathbb{Z}[M_{\leq c}^{a_0}]$$

for all pseudo-normed abelian group objects in any topos as above, and all $c > 0$.

PROOF. Each differential in the Breen-Deligne resolution is a finite sum of maps induced by maps $M^{a_{i+1}} \rightarrow M^{a_i}$ given by some $a_i \times a_{i+1}$ -matrix of integers. Given c_i , one can thus find some c_{i+1} so that $M_{\leq c_{i+1} c}^{a_{i+1}}$ maps into $M_{\leq c_i c}^{a_i}$ for each of those finitely many maps, which gives the claim. \square

We also need some homotopies. More precisely, we start with the following homotopy.

LEMMA 9.12. *For an abelian group M , the maps σ_1, σ_2 from*

$$C(M^2) : \dots \rightarrow \mathbb{Z}[M^{2a_i}] \rightarrow \dots \rightarrow \mathbb{Z}[M^{2a_1}] \rightarrow \mathbb{Z}[M^{2a_0}]$$

to

$$C(M) : \dots \rightarrow \mathbb{Z}[M^{a_i}] \rightarrow \dots \rightarrow \mathbb{Z}[M^{a_1}] \rightarrow \mathbb{Z}[M^{a_0}],$$

induced by addition $M^2 \rightarrow M$, respectively the sum of the two maps induced by two projections $M^2 \rightarrow M$, are homotopic, via some functorial homotopy

$$h_i : \mathbb{Z}[M^{2a_i}] \rightarrow \mathbb{Z}[M^{a_{i+1}}].$$

If M is a pseudo-normed abelian group object in any topos, then σ_1 and σ_2 are well-defined as maps of complexes from

$$C(M^2)_{\leq c/2} : \dots \rightarrow \mathbb{Z}[M_{\leq c_i c/2}^{2a_i}] \rightarrow \dots \rightarrow \mathbb{Z}[M_{\leq c_1 c/2}^{2a_1}] \rightarrow \mathbb{Z}[M_{\leq c/2}^{2a_0}]$$

to

$$C(M)_{\leq c} : \dots \rightarrow \mathbb{Z}[M_{\leq c_i c}^{a_i}] \rightarrow \dots \rightarrow \mathbb{Z}[M_{\leq c_1 c}^{a_1}] \rightarrow \mathbb{Z}[M_{\leq c}^{a_0}]$$

for all $c > 0$. In that case, for all $i \geq 0$ there are universal constants c'_i such that h_i defines well-defined maps

$$\mathbb{Z}[M_{\leq c_i c/2}^{2a_i}] \rightarrow \mathbb{Z}[M_{\leq c'_i c_{i+1} c}^{a_{i+1}}]$$

for all $c > 0$.

PROOF. This is a consequence of the proof of the existence of the Breen-Deligne resolution, proved in the same way as [Sch19, Proposition 4.17]. The existence of the constants c'_i is again formal, as in the last lemma. \square

Now we need the following generalization to adding N elements.

LEMMA 9.13. *Let N be a power of 2. The maps of complexes σ_1, σ_2 from*

$$C(M^N) : \dots \rightarrow \mathbb{Z}[M^{Na_i}] \rightarrow \dots \rightarrow \mathbb{Z}[M^{Na_1}] \rightarrow \mathbb{Z}[M^{Na_0}]$$

to

$$C(M) : \dots \rightarrow \mathbb{Z}[M^{a_i}] \rightarrow \dots \rightarrow \mathbb{Z}[M^{a_1}] \rightarrow \mathbb{Z}[M^{a_0}],$$

induced by addition $M^N \rightarrow M$, respectively the sum of the N maps induced by the N projections $M^N \rightarrow M$, are homotopic, via some functorial homotopy

$$h_i^N : \mathbb{Z}[M^{Na_i}] \rightarrow \mathbb{Z}[M^{a_{i+1}}]$$

which moreover satisfies the following bound, with the same constants c'_0, c'_1, \dots as in the previous lemma:

If M is a pseudo-normed abelian group object in any topos, then σ_1 and σ_2 are well-defined as maps of complexes from

$$C(M^N)_{\leq c/N} : \dots \rightarrow \mathbb{Z}[M_{\leq c_i c/N}^{Na_i}] \rightarrow \dots \rightarrow \mathbb{Z}[M_{\leq c_1 c/N}^{Na_1}] \rightarrow \mathbb{Z}[M_{\leq c/N}^{Na_0}]$$

to

$$C(M)_{\leq c} : \dots \rightarrow \mathbb{Z}[M_{\leq c_i c}^{a_i}] \rightarrow \dots \rightarrow \mathbb{Z}[M_{\leq c_1 c}^{a_1}] \rightarrow \mathbb{Z}[M_{\leq c}^{a_0}]$$

for all $c > 0$. In that case, h_i^N defines well-defined maps

$$\mathbb{Z}[M_{\leq c_i c/N}^{Na_i}] \rightarrow \mathbb{Z}[M_{\leq c'_i c_{i+1} c}^{a_{i+1}}]$$

for all $c > 0$.

PROOF. Let $N = 2^m$. For each $j = 0, \dots, m-1$, the two maps from $C(M^{2^{j+1}})$ to $C(M^{2^j})$ from the previous lemma are homotopic, and we use the homotopy from that lemma. Composing homotopies (which amounts concretely to a certain sum) we get the desired homotopy from $C(M^{2^m})$ to $C(M)$. It follows directly from this construction that the constants c'_i are unchanged. \square

10. Lecture X: Some computations with liquid modules

Now we have done the hard work to prove that there is a well-behaved category of liquid real vector spaces. In the rest of the course, we want to define analytic spaces, and show that various classical objects, like complex-analytic spaces, or rigid-analytic varieties, give examples.

However, we will leave the discussion of analytic spaces for after Christmas, and in this last lecture before Christmas, we simply want to get a feeling for the category of liquid real vector spaces, or liquid $\mathbb{Z}((T))_{>r}$ -modules; some of these computations will also be necessary for the discussion of complex-analytic spaces.

Let us do some basic computations of liquidifications.

PROPOSITION 10.1. *Fix any radius $0 < r < 1$.*

(1) *For any compact Hausdorff space S , let*

$$\mathcal{M}(S, \mathbb{Z}((T))_r) \subset \prod_{n \in \mathbb{Z}} \mathbb{Z}[S] \cdot T^n$$

be the union over all $c > 0$ of the (compact Hausdorff) subspace $\mathcal{M}(S, \mathbb{Z}((T))_r)_{\leq c}$ whose S' -valued points are all maps $S' \rightarrow \prod_{n \in \mathbb{Z}} \mathbb{Z}[S] \cdot T^n$ such that for all $s' \in S'$, the induced element $(\mu_n \cdot T^n)_n$, $\mu_n \in \mathbb{Z}[S]$ satisfies $\mu_n \in \mathbb{Z}[S]_{\leq a_n}$ for some nonnegative integers a_n such that $\sum a_n r^n \leq c$.

This definition agrees with the previous one for profinite sets S , and in general if $S_\bullet \rightarrow S$ is a hypercover of a compact Hausdorff S by profinite sets S_i , then

$$\mathcal{M}(S_\bullet, \mathbb{Z}((T))_r) \rightarrow \mathcal{M}(S, \mathbb{Z}((T))_r)$$

is a resolution. The derived r -liquidification of $\mathbb{Z}[T^{-1}][S]$ is given by

$$\mathcal{M}(S, \mathbb{Z}((T))_{>r}) = \varinjlim_{r' > r} \mathcal{M}(S, \mathbb{Z}((T))_{r'}).$$

(2) *For any compact abelian group A , the derived r -liquidification of $A[T^{-1}]$ is given by $A((T))$.*

By the first part, we can also define $\mathcal{M}_p(S)$ via base change to \mathbb{R} for general compact Hausdorff S ; equivalently, via resolving S by profinite sets S_i and forming the complex $\mathcal{M}_p(S_\bullet)$, which will be quasi-isomorphic to $\mathcal{M}_p(S)$ in degree 0. Then the derived p -liquidification of $\mathbb{R}[S]$ is equal to $\mathcal{M}_{<p}(S)$, and in particular sits in degree 0. We note that it seems slightly tricky to do these things directly over \mathbb{R} .

PROOF. For the first part, note that for profinite S , this is essentially the definition, with Proposition 2.1. For general compact Hausdorff S , consider a simplicial resolution of S by profinite sets S_i , and the corresponding resolution

$$\dots \rightarrow \mathbb{Z}[S_1] \rightarrow \mathbb{Z}[S_0] \rightarrow \mathbb{Z}[S] \rightarrow 0.$$

We consider this as a complex of pseudonormed abelian groups, with the subgroups $\mathbb{Z}[S]_{\leq n}$. We claim that whenever $x \in \mathbb{Z}[S_i]_{\leq n}$ with $d(x) = 0$ then there is some $y \in \mathbb{Z}[S_{i+1}]_{\leq n}$ with $x = d(y)$. Note that this statement depends only on the underlying sets of S and the S_i , and the hypercover $S_\bullet \rightarrow S$ of sets splits; this produces a contracting homotopy which easily gives the statement (the contracting homotopy is like in the second appendix to Lecture VIII). From here it is easy to see that also

$$\mathcal{M}(S_\bullet, \mathbb{Z}((T))_r) \rightarrow \mathcal{M}(S, \mathbb{Z}((T))_r)$$

is a resolution. Passing to the colimit over $r' > r$ gives the result (as $\mathcal{M}(S_\bullet, \mathbb{Z}((T))_{>r})$ is the derived r -liquidification of $\mathbb{Z}[T^{-1}][S_i]$, and thus the complex $\mathcal{M}(S_\bullet, \mathbb{Z}((T))_{>r})$ computes the derived r -liquidification of $\mathbb{Z}[T^{-1}][S]$).

For the second part, we need to see that $A((T))$ is r -liquid and that derived r -liquidification of $B = A((T))/A[T^{-1}]$ vanishes. It is easy to see that any map $S \rightarrow A((T))$ extends to a map $\tilde{f} : \mathcal{M}(S, \mathbb{Z}((T))_r) \rightarrow A((T))$; as $A((T))$ is quasiseparated, uniqueness is automatic, so $A((T))$ is r -liquid.

Now note that $B = A((T))/A[T^{-1}]$ is a compact abelian group with $\mathbb{Z}[T^{-1}]$ -module structure. It can be resolved via the Breen-Deligne resolution, so it suffices to see that for any compact Hausdorff space S with an endomorphism $T^{-1} : S \rightarrow S$, the derived r -liquidification of $\mathbb{Z}[S]$ (considered as $\mathbb{Z}[T^{-1}]$ -module) vanishes. This can be described as the cone of $T^{-1} - [T^{-1}]$ on the derived r -liquidification of $\mathbb{Z}[T^{-1}][S]$, the latter of which is described by the first part. Now $T^{-1} - [T^{-1}]$ has the inverse $T + [T^{-1}]T^2 + [T^{-1}]^2T^3 + \dots$, giving the result. \square

The following corollary generalizes [Sch19, Theorem 4.3 (ii)].

COROLLARY 10.2. *Let V be a p -liquid \mathbb{R} -vector space and let A be a compact abelian group. Then $R\mathbf{H}\mathbf{om}(A, V) = 0$. Equivalently, the derived p -liquidification of A vanishes.*

PROOF. Write $\mathbb{R} = \mathbb{Z}((T))_{>r}/(2 - T^{-1})$ where $r = 2^{-p}$. Then the derived p -liquidification of A is the cone of $2 - T^{-1}$ on the derived r -liquidification of $A[T^{-1}]$. The latter is $A((T))$ by part (2) of the proposition, and $2 - T^{-1}$ is invertible on this (with inverse $T + 2T^2 + 4T^3 + \dots$). \square

REMARK 10.3. In the proof of [Sch19, Theorem 4.3 (ii)], we also used the action of $2 - [2]$ on a Breen-Deligne resolution; this seems closely related to writing $\mathbb{R} = \mathbb{Z}((T))_{>r}/(2 - T^{-1})$.

The corollary implies formally that if V is a p -liquid \mathbb{R} -vector space considered as a trivial representation of a compact abelian group A , then the group cohomology $H^i(A, V)$ vanishes for $i > 0$. (Indeed, by general nonsense there is a spectral sequence starting with $\mathrm{Ext}^i(\bigwedge^j A, V)$ converging to $H^{i+j}(A, V)$.) The proof uses commutativity critically.

QUESTION 10.4. Let G be a compact nonabelian group, say $G = \mathrm{SU}(2)$, and let V be a p -liquid \mathbb{R} -vector space, e.g. a p -Banach. Does $H^i(G, V)$ vanish for $i > 0$?

Here is a related question.

QUESTION 10.5. Consider \mathbb{C} as a condensed ring, and correspondingly the algebraic K -theory $K(\mathbb{C})$ as a condensed spectrum (sending an extremally disconnected S to $K(C(S, \mathbb{C}))$). What is the p -liquidification of $K(\mathbb{C})$? What is the r -liquidification of $K(\mathbb{C})[T^{-1}]$?¹⁹

We note that the solidification can be computed:²⁰

PROPOSITION 10.6. *The solidification of $K(\mathbb{C})$ is ku , connective topological K -theory.*

PROOF. The essential point is that $\mathbb{S}[\mathrm{GL}_n(\mathbb{C})]^\blacksquare \cong \mathbb{S}[|\mathrm{GL}_n(\mathbb{C})|]$ where $|\mathrm{GL}_n(\mathbb{C})|$ is the corresponding anima (cf. [Sch19, Example 6.5]). \square

¹⁹For the latter, we remark that one can define liquidification also over $\mathbb{S}((T))_{>r}$ where \mathbb{S} is the sphere spectrum.

²⁰Again, solidification can be defined over \mathbb{S} .

Of course, solidification forgets all information about real vector spaces, and we want to recover those via p -liquidification. This has the slightly annoying feature that, at least a priori, it depends on p . The r -liquidification of $K(\mathbb{C})[T^{-1}]$ has the virtue that it specializes to all possible p -liquidifications via specializing T to different values; and it remembers the integral information. We note that for $K_1(\mathbb{C}) = \mathbb{C}^\times$, the r -liquidification of $\mathbb{C}^\times[T^{-1}]$ can be computed: Using the decomposition $\mathbb{C}^\times \cong \mathbb{R} \times \mathbb{R}/\mathbb{Z}$, it is

$$\mathbb{C}^\times((T))_{>r} = \mathbb{R}((T))_{>r} \times \mathbb{R}/\mathbb{Z}((T)).$$

On the other hand, let us try to understand the structure of $\mathcal{M}(S, \mathbb{Z}((T))_r)$ and $\mathcal{M}_p(S)$ better.

PROPOSITION 10.7. *Let S be a compact Hausdorff space and $0 < p < 1$. Then for any $\mu \in \mathcal{M}_p(S)$ (element of the underlying set) there is a sequence s_0, s_1, \dots in S and real numbers x_0, x_1, \dots with $\sum |x_n|^p < \infty$, such that*

$$\mu = \sum_{n=0}^{\infty} x_n [s_n].$$

Similarly, for $0 < r < 1$ and any $\mu \in \mathcal{M}(S, \mathbb{Z}((T))_r)$, there is a sequence $s_0, s_1, \dots \in S$ and elements $x_0, x_1, \dots \in \mathbb{Z}((T))_r$ such that $x_n \in \mathbb{Z}((T))_{r, \leq c_n}$ with $\sum c_n < \infty$ and

$$\mu = \sum_{n=0}^{\infty} x_n [s_n].$$

Thus, all p -measures are just countable sums of Dirac measures. In particular, Haar measures which are “equidistributed” over the whole space, will never lie in $\mathcal{M}_p(S)$ for $p < 1$ (except for finite groups). This phenomenon, of measures being countable sums of Dirac measures, actually happens more generally for measures “of bounded entropy”, and this is the minimal condition that the Ribe extension forces on us.

We note that this is only a description of the underlying set: As a condensed set, $\mathcal{M}_p(S)$ is not a filtered colimit over its subspaces for (closures of) countable subspaces of S . Indeed, the map $S \rightarrow \mathcal{M}_p(S)$ does not factor over such a subspace in general.

PROOF. It suffices to handle the assertion over $\mathbb{Z}((T))_r$; the case of $\mathcal{M}_p(S)$ follows via base change. But then there is some $c < \infty$ such that

$$\mu \in \mathcal{M}(S, \mathbb{Z}((T))_r)_{\leq c} \subset \prod_n \mathbb{Z}[S] \cdot T^n,$$

and each coefficients of T^n involves only finitely many elements of S . It follows that one can write $\mu = \sum_{n=0}^{\infty} x_n [s_n]$ for some countable sequence of elements $s_n \in S$, which we assume has no repetitions. The condition $\mu \in \mathcal{M}(S, \mathbb{Z}((T))_r)$ then amounts to the condition that if $x_n \in \mathbb{Z}((T))_{r, \leq c_n}$ with c_n chosen minimal, then $\sum c_n \leq c$. This gives the result. \square

Finally, we use the formalism in the simplest situation of relevance to complex-analytic geometry.

PROPOSITION 10.8. *Let V be the condensed \mathbb{C} -vector space of overconvergent holomorphic functions on the closed unit disc $\{|z| \leq 1\}$. Equivalently,*

$$V = \varinjlim_{r>1} \left\{ \sum a_n z^n \in \mathbb{C}[[z]] \mid \sum |a_n| r^{-n} < \infty \right\}$$

where each term in the filtered colimit is regarded as a Banach space. Then for any $0 < p < 1$, the tensor product

$$V \otimes_{\mathbb{C}, p}^{\text{Liq}} V$$

is given by

$$\varinjlim_{r>1} \left\{ \sum a_n b_m z^n w^m \in \mathbb{C}[[z, w]] \mid \sum |a_n b_m| r^{-n-m} < \infty \right\}$$

of overconvergent holomorphic functions on the 2-dimensional disc $\{|z|, |w| \leq 1\}$.

PROOF. Decompose $\mathcal{M}_{<p}(\mathbb{N} \cup \{\infty\}) = W \oplus \mathbb{R} \cdot [\infty]$. Then W is the union of the spaces $W_{p'}$ for $p' < p$ whose compact subspaces are given by the space of sequences $(x_0, x_1, \dots) \in \mathbb{R}$ with $\sum |x_i|^{p'} \leq c$. The key observation is now that

$$V = \varinjlim_{r>1} W_{\mathbb{C}}$$

along the maps $f_r : W_{\mathbb{C}} \rightarrow V$ sending (x_0, x_1, \dots) to $\sum x_n r^n z^n$; this is a simple verification, based on the observation that multiplication by the sequence $(1, t, t^2, \dots)$ with $t < 1$ takes bounded sequences to ℓ^p -summable sequences for all $0 < p < 1$.

But now the liquid tensor product of $\mathcal{M}_{<p}(\mathbb{N} \cup \{\infty\})$ with itself is

$$\mathcal{M}_{<p}((\mathbb{N} \cup \{\infty\}) \times (\mathbb{N} \cup \{\infty\}))$$

of which $W \otimes_p^{\text{Liq}} W$ is the direct factor whose compact subspaces are the space of sequences $(x_{nm})_{n,m \geq 0} \in \mathbb{R}$ with $\sum |x_{n,m}|^{p'} \leq c$ for some $p' < p$ and $c > 0$. This gives the result by passing to the colimit over r . \square

COROLLARY 10.9. *Let $D_1, D_2 \subset \mathbb{C}$ be two closed discs with empty intersection. Let $\mathcal{O}(D_1)$ and $\mathcal{O}(D_2)$ be the spaces of overconvergent holomorphic functions, considered as condensed $\mathbb{C}[z]$ -modules. Then*

$$\mathcal{O}(D_1) \otimes_{\mathbb{C}[z], p}^{\text{Liq}} \mathcal{O}(D_2) = 0.$$

PROOF. Forming the tensor product over \mathbb{C} , one gets $\mathcal{O}(D_1 \times D_2)$ by the previous proposition. On this space, $z_1 - z_2$ is invertible, so the tensor product over $\mathbb{C}[z]$ vanishes. \square

We see that for these questions, the choice of p is largely irrelevant, and that one gets the expected tensor products.

11. Lecture XI: Towards localization

In the remaining lectures, we will define the category of analytic spaces, and give various examples. Let us at this point remark that the resulting theory of analytic spaces will, notably, be an absolute theory: Usually, one defines complex-analytic spaces or rigid-analytic varieties over some nonarchimedean field K , so works over a fixed base field (and works with K -algebras satisfying some topological finiteness condition). Here, just like with schemes, it will not be necessary to specify any base in advance.²¹

There are two ways to define schemes: As locally ringed topological spaces, or as certain functors on the category of rings. In the latter approach, the essential non-formal input is just the notion of a localization $A \rightarrow A[f^{-1}]$ of rings; from the category of rings with this class of morphisms, one can recover the category of schemes. Indeed, schemes will be a full subcategory of the category of functors from rings to sets; in particular, we ask that they are Zariski sheaves (where a finite family of localizations $A \rightarrow A[f_i^{-1}]$ is a cover if the base change functor from A -modules to $\prod_i A[f_i^{-1}]$ -modules is faithful). An affine scheme is a functor of the form $F_A : B \mapsto \text{Hom}(A, B)$ for some ring A . A map $F \rightarrow G$ of Zariski sheaves is an open immersion if it is injective and for every ring B with a section $s \in G(B)$, corresponding to a map $F_B \rightarrow G$, the fibre product $F \times_G F_B$ is the colimit of F_A over all localizations $B \rightarrow A$ for which $F_A \rightarrow F_B$ factors over $F \times_G F_B$. Finally, a scheme is a functor F that admits an open cover by F_A 's.

We will follow this route for the definition of analytic spaces. Thus, the next goal is to define the correct notion of localization $\mathcal{A} \rightarrow \mathcal{B}$ of analytic rings. The notion is supposed to have the following properties:

- (1) It is stable under base change.
- (2) It is stable under composition.
- (3) It is stable under filtered colimits.

The first condition actually requires some explanation; we will come to this later. Regarding the third condition, note that this means that in the usual algebraic context we allow general localizations $A \rightarrow A[S^{-1}]$ at multiplicative subsets. It turns out that in the analytic category, there is no reasonable “finiteness” one can a priori impose on the localizations, so we will allow general ones.

EXERCISE 11.1. Verify that the above formal definition of schemes also gives a well-defined notion when one allows general localizations $A \rightarrow A[S^{-1}]$ of rings.

Let us first give several examples.

EXAMPLE 11.2.

- (1) If $A \rightarrow B = A[S^{-1}]$ is a usual localization of (discrete) rings, with trivial analytic ring structure, then $A \rightarrow B$ will be allowed.
- (2) The map $(\mathbb{Z}[T], \mathbb{Z})_{\blacksquare} \rightarrow (\mathbb{Z}[T], \mathbb{Z}[T])_{\blacksquare}$, corresponding to localization to the subset $\{|T| \leq 1\}$ of the adic space $\text{Spa}(\mathbb{Z}[T], \mathbb{Z})$.

²¹Let us mention that Ben-Bassat, Kremnitzer, and coauthors, have also recently undertaken a general study of analytic spaces, including some notion of quasicohherent sheaves, cf. e.g. [BBK17]. Their theory is based on working in the symmetric monoidal category of ind-Banach spaces over a fixed base field, roughly in the manner paraphrased in the first lecture (that is the “relative algebraic geometry” of Toën–Vaquié–Vezzosi). In particular, it is a relative theory; but also in the relative situation, our theory is quite different, making use of varying analytic ring structures.

- (3) More generally, if $X = \mathrm{Spa}(A, A^+)$ is a (reasonable) adic space and $U \subset X$ is a rational subset, the map

$$(A, A^+)_{\blacksquare} \rightarrow (\mathcal{O}_X(U), \mathcal{O}_X^+(U))_{\blacksquare}.$$

- (4) In the context of the last lecture, if $D_1 \subset D_2$ is an inclusion of closed discs in \mathbb{C} , then $\mathcal{O}(D_2) \rightarrow \mathcal{O}(D_1)$ (endowed with the analytic ring structure induced from the p -liquid structure on \mathbb{C} for some chosen $0 < p \leq 1$) is a localization.

The parenthetical word “reasonable” occurs here as general adic spaces are not well-behaved: For example, the structure presheaf may fail to be a sheaf. This failure will actually be corrected by the formalism of analytic spaces, and the correction will be by slightly changing the localizations in general.

For example, consider the case $U = \{|f| \leq 1\}$ for some $f \in A$. Then $\mathcal{O}_X(U)$ is by definition the quotient of the convergent power series algebra $A\langle T \rangle$ by the closure of the ideal generated by $T - f$. On the other hand, in practice this ideal is already closed and moreover $T - f$ is a non-zerodivisor (in fact, by [KL19, Lemma 2.4.10], this is automatic if (A, A^+) is sheafy and A has a topologically nilpotent unit, for example is an algebra over a nonarchimedean field). Thus, in this case

$$\mathcal{O}_X(U) = A\langle T \rangle / (T - f) = A\langle T \rangle /^{\mathbb{L}}(T - f) = [A\langle T \rangle \xrightarrow{T-f} A\langle T \rangle].$$

On the other hand, in general both of these properties can fail. Note that algebraically, the most natural object is the complex on the right, and indeed this will be the localization in the world of analytic rings. Working with topological rings, it is impossible to work with the quotient $A\langle T \rangle / (T - f)$ if the ideal $(T - f)$ is not closed. However, passing to the condensed world, the quotient $A\langle T \rangle / (T - f)$ is perfectly well-behaved. If $T - f$ is a zerodivisor, it however becomes necessary to work even with the derived reduction $A\langle T \rangle /^{\mathbb{L}}(T - f)$: A condensed *animated* ring.

We had seen last semester that localizations will in general not be flat, and turn modules concentrated in degree 0 into complexes (in nonnegative homological degrees). The same will happen in general for localizations of analytic rings, so we need to also derive our rings. This is the subject of derived algebraic geometry, which replaces the usual category of rings by its animation, the ∞ -category of animated rings.²²

11.1. Animation. Let us briefly recall the notion of animation.²³ Let \mathcal{C} be a category that admits all small colimits. Recall that an object $X \in \mathcal{C}$ is compact (also called finitely presented) if $\mathrm{Hom}(X, -)$ commutes with filtered colimits. An object $X \in \mathcal{C}$ is projective if $\mathrm{Hom}(X, -)$ commutes with reflexive coequalizers, i.e. colimits along $\Delta_{\leq 1}^{\mathrm{op}}$ (coequalizers of parallel arrows $Y \rightrightarrows Z$ with a simultaneous section $Z \rightarrow Y$ of both maps). Taken together, an object $X \in \mathcal{C}$ is compact projective if $\mathrm{Hom}(X, -)$ commutes with all filtered colimits and reflexive coequalizers; equivalently, it commutes with all (so-called) 1-sifted colimits.

Let $\mathcal{C}^{\mathrm{cp}} \subset \mathcal{C}$ be the full subcategory of compact projective objects. There is a fully faithful embedding $\mathrm{sInd}(\mathcal{C}^{\mathrm{cp}}) \rightarrow \mathcal{C}$ from the 1-sifted Ind-category of $\mathcal{C}^{\mathrm{cp}}$ (the full subcategory of $\mathrm{Fun}((\mathcal{C}^{\mathrm{cp}})^{\mathrm{op}}, \mathrm{Set})$

²²We refer to the work of Toën–Vezzosi, [TV08], and Lurie, [Lur18], for developments of derived algebraic geometry. Note that these sources say “simplicial ring” when they mean “animated ring”.

²³In classical language, this is a non-abelian derived category in the sense of Quillen. It has also been studied by Rosicky, [Ros07], and we follow Lurie, [Lur09, Section 5.5.8], with language that has been coined by Clausen inspired by Beilinson’s [Bei07], and used first in [ČS19].

generated under small 1-sifted colimits by the Yoneda image; equivalently, the category freely generated under small 1-sifted colimits by \mathcal{C}^{cp} . If \mathcal{C} is generated under small colimits by \mathcal{C}^{cp} , then this functor is an equivalence

$$\text{sInd}(\mathcal{C}^{\text{cp}}) \cong \mathcal{C}.$$

If \mathcal{C}^{cp} is small, then

$$\text{sInd}(\mathcal{C}^{\text{cp}}) \subset \text{Fun}((\mathcal{C}^{\text{cp}})^{\text{op}}, \text{Set})$$

is exactly the full subcategory of functors that take finite coproducts in \mathcal{C}^{cp} to products in Set .

EXAMPLE 11.3.

- (1) If $\mathcal{C} = \text{Set}$ is the category of sets, then \mathcal{C}^{cp} is the category of finite sets, which generates \mathcal{C} under small colimits.
- (2) If $\mathcal{C} = \text{Ab}$ is the category of abelian groups, then \mathcal{C}^{cp} is the category of finite free abelian groups, which generates \mathcal{C} under small colimits.
- (3) If $\mathcal{C} = \text{Ring}$ is the category of commutative rings, then \mathcal{C}^{cp} is the category of retracts of polynomial rings $\mathbb{Z}[X_1, \dots, X_n]$, which generates \mathcal{C} under small colimits.
- (4) If $\mathcal{C} = \text{Cond}(\text{Set})$ is the category of condensed sets, then \mathcal{C}^{cp} is the category of extremally disconnected profinite sets, which generates \mathcal{C} under small colimits.²⁴
- (5) If $\mathcal{C} = \text{Cond}(\text{Ab})$ is the category of condensed abelian groups, then \mathcal{C}^{cp} is the category of direct summands of $\mathbb{Z}[S]$ for extremally disconnected S , which generates \mathcal{C} under small colimits.
- (6) If $\mathcal{C} = \text{Cond}(\text{Ring})$ is the category of condensed rings, then \mathcal{C}^{cp} is the category of retracts of $\mathbb{Z}[\mathbb{N}[S]]$ for extremally disconnected S , where $\mathbb{N}[S]$ is free condensed abelian monoid on S and thus $\mathbb{Z}[\mathbb{N}[S]]$ is the free condensed ring on S . Again, \mathcal{C}^{cp} generates \mathcal{C} under small colimits.

DEFINITION 11.4. Let \mathcal{C} be a category that admits all small colimits and is generated under small colimits by \mathcal{C}^{cp} . The animation of \mathcal{C} is the ∞ -category $\text{Ani}(\mathcal{C})$ freely generated under sifted colimits by \mathcal{C}^{cp} .

EXAMPLE 11.5. If $\mathcal{C} = \text{Set}$, then $\text{Ani}(\mathcal{C}) = \text{Ani}(\text{Set}) =: \text{Ani}$ is the ∞ -category of animated sets, or anima for brevity. In standard language, this is the ∞ -category of “spaces”.

Let us describe the nature of the ∞ -category of anima. Any anima has a set of connected components, giving a functor $\pi_0 : \text{Ani} \rightarrow \text{Set}$ (in fact, π_0 is simply given by the universal property of Ani , as Set is a category with all sifted colimits with a functor from finite sets), which has a fully faithful right adjoint $\text{Set} \hookrightarrow \text{Ani}$. Given an anima A with a point $a \in A$ (meaning a map $a : * \rightarrow A$), one can define groups $\pi_i(A, a)$ for $i \geq 1$, which are abelian for $i \geq 2$. The map $a : * \rightarrow A$ is an equivalence if and only if $\pi_0 A$ is a point and $\pi_i(A, a) = 0$ for all $i \geq 1$. An anima is defined to be i -truncated if $\pi_j(A, a) = 0$ for all $a \in A$ and $j > i$. Then A is 0-truncated if and only if it is in the essential image of $\text{Set} \hookrightarrow \text{Ani}$. The inclusion of i -truncated anima into all anima has a left adjoint $\tau_{\leq i}$. For all anima A , the natural map

$$A \rightarrow \lim_i \tau_{\leq i} A$$

is an equivalence; this is the “convergence of the Postnikov tower”. Picking any $a \in A$ and $i \geq 1$, the fibre of $\tau_{\leq i} A \rightarrow \tau_{\leq i-1} A$ over the image of a is an Eilenberg-MacLane anima $K(\pi_i(A, a), i)$.

²⁴In this example (and the ones to follow), \mathcal{C}^{cp} is itself a large category.

Here, an Eilenberg-MacLane anima $K(\pi, i)$, with $i \geq 1$ an integer and π a group that is abelian if $i > 1$, is a pointed connected anima with $\pi_j = 0$ for $j \neq i$ and $\pi_i = \pi$. It is unique up to unique isomorphism. In fact, the ∞ -category of pointed connected anima (A, a) with $\pi_j(A, a) = 0$ for $j \neq i$ is equivalent to the category of groups when $i = 1$, and to the category of abelian groups when $i \geq 2$.

There are several ways to construct $\text{Ani}(\mathcal{C})$. It can be defined as the full sub- ∞ -category of

$$\text{Fun}((\mathcal{C}^{\text{cp}})^{\text{op}}, \text{Ani})$$

generated under sifted colimits by the Yoneda image. If \mathcal{C}^{cp} is small, this agrees with the full sub- ∞ -category of all contravariant functors $F : \mathcal{C}^{\text{cp}} \rightarrow \text{Ani}$ that take finite coproducts to products. On the other hand, recall that all sifted colimits are generated by filtered colimits and geometric realizations of simplicial objects. This makes simplicial objects central. In fact, Ani can be defined as the ∞ -category obtained from the category of simplicial sets sSet by inverting weak equivalences. Similarly, for any ∞ -category \mathcal{C} generated under small colimits by \mathcal{C}^{cp} , one can describe $\text{Ani}(\mathcal{C})$ as the ∞ -category obtained from simplicial objects in \mathcal{C} by inverting weak equivalences.²⁵

EXAMPLE 11.6.

- (1) As indicated above, for $\mathcal{C} = \text{Set}$, one gets the ∞ -category of anima (a.k.a. spaces).
- (2) For $\mathcal{C} = \text{Ab}$, the ∞ -category $\text{Ani}(\text{Ab})$ of animated abelian groups is, by the Dold-Kan equivalence, equivalent to the ∞ -derived category of abelian groups $\mathcal{D}_{\geq 0}(\text{Ab})$ in nonnegative homological degrees. This motivates the term “nonabelian deried category” for the general construction of animation.
- (3) For $\mathcal{C} = \text{Ring}$, one gets the ∞ -category of animated rings.
- (4) For $\mathcal{C} = \text{Cond}(\text{Set})$, one gets the ∞ -category $\text{Ani}(\text{Cond}(\text{Set}))$ of animated condensed sets. The two operations $\text{Ani}(-)$ and $\text{Cond}(-)$ actually commute by Lemma 11.8 below, so this is equivalent the ∞ -category $\text{Cond}(\text{Ani}(\text{Set})) = \text{Cond}(\text{Ani})$ of condensed anima.
- (5) For $\mathcal{C} = \text{Cond}(\text{Ab})$, one gets, by Dold-Kan again, the ∞ -derived category $\mathcal{D}_{\geq 0}(\text{Cond}(\text{Ab}))$ of condensed abelian groups in nonnegative homological degrees.
- (6) For $\mathcal{C} = \text{Cond}(\text{Ring})$, one gets the ∞ -category of animated condensed rings, or equivalently of condensed animated rings.

The operations of animation and passage to condensed objects commute:

DEFINITION 11.7. Let \mathcal{C} be an ∞ -category that admits all small colimits. For any uncountable strong limit cardinal κ , the ∞ -category $\text{Cond}_{\kappa}(\mathcal{C})$ of κ -condensed objects of \mathcal{C} is the category of contravariant functors from κ -small extremally disconnected profinite sets S to \mathcal{C} that take finite coproducts to products.

Moreover, using the fully faithful left adjoints to the forgetful functors,

$$\text{Cond}(\mathcal{C}) := \varinjlim_{\kappa} \text{Cond}_{\kappa}(\mathcal{C}).$$

LEMMA 11.8. *Let \mathcal{C} be a category that is generated under small colimits by \mathcal{C}^{cp} . Then $\text{Cond}(\mathcal{C})$ is still generated under small colimits by its compact projective objects, and there is a natural equivalence of ∞ -categories*

$$\text{Cond}(\text{Ani}(\mathcal{C})) \cong \text{Ani}(\text{Cond}(\mathcal{C})).$$

²⁵We regard it thus as a misnomer to call objects of $\text{Ani}(\text{Ring})$ simplicial rings: One has changed the morphisms drastically.

PROOF. A family of compact projective generators of $\text{Cond}(\mathcal{C})$ is parametrized by pairs (S, X) of an extremally disconnected S and a compact projective object $X \in \mathcal{C}^{\text{cp}}$, given by the sheafification of sending T to the colimit over $\text{Hom}(T, S)$ many copies of X . These objects still define compact projective objects of the ∞ -category $\text{Cond}(\text{Ani}(\mathcal{C}))$, and generate the latter under small colimits. This implies the desired result. \square

We will be interested in condensed animated rings. However, let us make some remarks about the nature of the ∞ -category of condensed anima, as this combines two different flavours of topology.

The traditional approach to homotopy theory is to define the ∞ -category of anima, or “spaces”, by taking the category of CW complexes, and inverting weak equivalences. This becomes potentially very confusing when one is also considering condensed sets. Namely, CW complexes also embed fully faithfully into condensed sets.

Thus, there are two natural functors from CW complexes to condensed anima: A fully faithful functor, via the full subcategory of condensed sets; and another non-faithful functor, factoring over the full- ∞ -subcategory of anima. Let us (abusively) denote the first functor by $X \mapsto X$, and the second by $X \mapsto |X|$. Then S^1 is a physical circle, while $|S^1|$ is some ghostly appearance of a point with an internal automorphism (the anima $B\mathbb{Z}$).

LEMMA 11.9. *Let X be a CW complex (more generally, any condensed set that is a filtered colimit of condensed sets that are built out of pushouts of $S^{n-1} = \partial D^n \hookrightarrow D^n$). There is a universal anima Y with a map $X \rightarrow Y$ of condensed anima. There is a functorial identification $Y \cong |X|$.*

In particular, there is a natural map $X \rightarrow |X|$ of condensed anima. For S^1 , we get a cartesian square (of condensed anima)

$$\begin{array}{ccc} \mathbb{R} & \longrightarrow & * \\ \downarrow & & \downarrow \\ S^1 & \longrightarrow & |S^1| \end{array}$$

involving the universal cover $\mathbb{R} \rightarrow S^1$. If X is a connected CW complex, then $X \times_{|X|} \tau_{\geq 2}|X| \rightarrow X$ is the universal cover of X . When X has higher homotopy groups, then $X \times_{|X|} *$ is an actual condensed anima in that it is nontrivial both in the condensed and in the animated direction. We invite the reader to get a mental image of $S^2 \times_{|S^2|} *$.

PROOF. As X is an iterated pushout of $S^{n-1} = \partial D^n \hookrightarrow D^n$ (and these pushouts are compatible with the fully faithful functors from compactly generated topological spaces to condensed sets to condensed anima), it suffices to prove the existence of Y in those cases; and this reduces, moreover, inductively to the case $X = D^n$. In that case, the natural map $Y \rightarrow \text{Hom}(X, Y)$ is an equivalence for any anima Y . To check this, it suffices to see that it induces a bijection on π_0 : Indeed, applying the statement on the level of π_0 to $\text{Hom}(Z, Y)$ for any other anima Z will then give $\pi_0 \text{Hom}(Z, Y) \cong \pi_0 \text{Hom}(Z, \text{Hom}(X, Y))$ for all Z (as $\text{Hom}(Z, \text{Hom}(X, Y)) \cong \text{Hom}(X, \text{Hom}(Z, Y))$), which means that $Y \rightarrow \text{Hom}(X, Y)$ is an equivalence.

To check that $\pi_0 Y \rightarrow \pi_0 \text{Hom}(D^n, Y)$ is an isomorphism, we pass up the Postnikov tower. If Y is discrete, this follows from connectedness of X . If $Y = BG$ for some discrete group G , then $\text{Hom}(X, Y)$ classifies G -torsors over X (as a condensed set). It is not hard to see that these are locally on X trivial, as any trivialization at a point spreads uniquely to small closed neighborhoods

(after a resolution by extremally disconnected sets, and thus by descent on X). Thus, they are classified by usual G -torsors on the topological space $X = D^n$, and these are all split. If $Y = K(\pi, n)$ for $n > 1$, then $\text{Hom}(X, Y)$ is given by $H^n(X, \pi)$ (cohomology of X as a condensed set). By last semester, this agrees with singular cohomology, and so it vanishes for $X = D^n$.

We see that for $X = D^n$, the map $X \rightarrow |X| = *$ is the universal map to an anima. As the functor sending X to this initial anima commutes with colimits, as does $X \rightarrow |X|$, this gives the identification in general. \square

12. Lecture XII: Localizations

We need to generalize the theory of analytic rings from condensed rings to condensed animated rings. As this works in the context of (still unital) associative rings, let us temporarily work in that setting. It has the advantage that the ∞ -category of animated condensed associative rings can also be described as the ∞ -category of \mathbb{E}_1 -algebras in the symmetric monoidal ∞ -category $\mathcal{D}_{\geq 0}(\text{Cond}(\text{Ab})) = \text{Ani}(\text{Cond}(\text{Ab}))$ of animated condensed abelian groups. Any animated condensed ring \mathcal{A} thus has an ∞ -category $\mathcal{D}_{\geq 0}(\mathcal{A})$ of modules in animated condensed abelian groups. This is a prestable ∞ -category in the sense of Lurie (cf. [Lur18, Appendix C]); in particular, it embeds fully faithfully into a stable ∞ -category $\mathcal{D}(\mathcal{A})$, as the positive part of a t -structure. The heart is equivalent to the category of condensed modules over the condensed ring $\pi_0\mathcal{A}$. (The preceding discussion applies to sheaves on any site.)

The following is a direct generalization of [Sch19, Definition 7.1, 7.4], but we decided to change notation.

DEFINITION 12.1. An analytic animated associative ring is a pair $(\mathcal{A}, \mathcal{M})$ consisting of a condensed animated associative ring \mathcal{A} together with a covariant functor $\mathcal{M} : S \mapsto \mathcal{M}[S]$ from extremally disconnected profinite sets S to condensed animated \mathcal{A} -modules, taking finite coproducts to finite direct sums, together with a natural transformation $S \rightarrow \mathcal{M}[S]$ of condensed anima, with the following property. For any object $C \in \mathcal{D}_{\geq 0}(\mathcal{A})$ that is a sifted colimit of objects of the form $\mathcal{M}[S]$, the natural map

$$\underline{\text{Hom}}_{\mathcal{D}_{\geq 0}(\mathcal{A})}(\mathcal{M}[S'], C) \rightarrow \underline{\text{Hom}}_{\mathcal{D}_{\geq 0}(\mathcal{A})}(\mathcal{A}[S'], C)$$

of condensed anima is an equivalence for all extremally disconnected profinite sets S' .

REMARK 12.2. The letter \mathcal{M} can be thought of as denoting spaces of *measures*, or as denoting extra data on *modules*, singling out the correct category of modules inside all of $\mathcal{D}_{\geq 0}(\mathcal{A})$.

We note that in the condition, the map is actually a map of condensed animated abelian groups, and by embedding into $\mathcal{D}(\mathcal{A})$, we could also use the $R\text{Hom}$ and ask for an isomorphism in $\mathcal{D}(\text{Cond}(\text{Ab}))$, as in [Sch19, Definition 7.4]. (A priori, the condition on $R\text{Hom}$ is stronger, but it follows from the given condition applied to all shifts $C[i]$, $i \geq 0$.)

Apart from language, the only difference with the situation of [Sch19, Definition 7.1, 7.4] is that we allow both \mathcal{A} and $\mathcal{M}[S]$ to be complexes, living in nonnegative homological degrees.

DEFINITION 12.3. Let $(\mathcal{A}, \mathcal{M})$ be an analytic animated associative ring. The ∞ -category $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$ is defined to be the full ∞ -subcategory

$$\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M}) \subset \mathcal{D}_{\geq 0}(\mathcal{A})$$

spanned by all $C \in \mathcal{D}_{\geq 0}(\mathcal{A})$ such that for all extremally disconnected profinite sets S , the map

$$\underline{\text{Hom}}_{\mathcal{D}_{\geq 0}(\mathcal{A})}(\mathcal{M}[S], C) \rightarrow \underline{\text{Hom}}_{\mathcal{D}_{\geq 0}(\mathcal{A})}(\mathcal{A}[S], C)$$

of condensed anima is an equivalence.

PROPOSITION 12.4. *Let $(\mathcal{A}, \mathcal{M})$ be an analytic animated associative ring. The ∞ -category $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$ is generated under sifted colimits by the objects $\mathcal{M}[S]$ for varying extremally disconnected profinite sets S , which are compact projective objects of $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$. The full ∞ -subcategory*

$$\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M}) \subset \mathcal{D}_{\geq 0}(\mathcal{A})$$

is stable under all limits and colimits and admits a left adjoint

$$- \otimes_{\mathcal{A}} (\mathcal{A}, \mathcal{M}) : \mathcal{D}_{\geq 0}(\mathcal{A}) \rightarrow \mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$$

sending $\mathcal{A}[S]$ to $\mathcal{M}[S]$.

The ∞ -category $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$ is prestable. Its heart is the abelian category $\mathcal{D}^{\heartsuit}(\mathcal{A}, \mathcal{M})$ that is the full subcategory of condensed $\pi_0\mathcal{A}$ -modules generated under colimits by $\pi_0\mathcal{M}[S]$ for varying S . An object $C \in \mathcal{D}_{\geq 0}(\mathcal{A})$ lies in $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$ if and only if all $H_i(C)$ lie in $\mathcal{D}^{\heartsuit}(\mathcal{A}, \mathcal{M})$.

If \mathcal{A} has the structure of a condensed animated commutative ring so that $\mathcal{D}_{\geq 0}(\mathcal{A})$ is naturally a symmetric monoidal ∞ -category, there is a unique symmetric monoidal structure on $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$ making $- \otimes_{\mathcal{A}} (\mathcal{A}, \mathcal{M})$ symmetric monoidal.

REMARK 12.5. Passing to “spectrum objects” (a formal procedure that in the current situation recovers $\mathcal{D}(-)$ from $\mathcal{D}_{\geq 0}(-)$), similar statements hold true on the level of $\mathcal{D}(\mathcal{A}, \mathcal{M}) \subset \mathcal{D}(\mathcal{A})$.

A consequence of the proposition is that $\mathcal{M}[S]$ is determined by the full ∞ -subcategory $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M}) \subset \mathcal{D}_{\geq 0}(\mathcal{A})$ as the image of $\mathcal{A}[S]$ under the left adjoint to the full inclusion. Moreover, this ∞ -subcategory is determined already by the collection $\{\pi_0\mathcal{M}[S]\}_S$ of condensed $\pi_0\mathcal{A}$ -modules. In other words, analytic ring structures on \mathcal{A} are completely determined by data on the level of usual abelian categories. In the first appendix to this lecture, we will characterize analytic ring structures from this perspective.

PROOF. By the definition of analytic animated rings, all sifted colimits of objects $\mathcal{M}[S]$ are in $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$. On the other hand, by the definition of $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$, the objects $\mathcal{M}[S]$ are compact projective generators.

By definition of $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$, it is stable under all limits. For the left adjoint, it is clear that it exists on $\mathcal{A}[S]$ with value $\mathcal{M}[S]$ by definition of $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$, and then it also exists for all of their sifted colimits by the assumption that $(\mathcal{A}, \mathcal{M})$ is an analytic ring; but these sifted colimits exhaust $\mathcal{D}_{\geq 0}(\mathcal{A})$. Now the existence of general colimits follows, by forming them first in $\mathcal{D}_{\geq 0}(\mathcal{A})$, and then applying the left adjoint.

It is now formal that $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$ is prestable, cf. [Lur18, Corollary C.1.2.3]. If $C \in \mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$, then also $\tau_{\geq 1}C \in \mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$ as this is the suspension of the loops of C and $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$ is stable under all limits and colimits. Thus, $H_0(C)[0] \in \mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$. The statement about the heart is then formal.

Regarding the symmetric monoidal structure, one possible reference is [NS18, Theorem I.3.6], noting that the condition of being an analytic ring ensures that the kernel of $- \otimes_{\mathcal{A}} (\mathcal{A}, \mathcal{M})$ is a \otimes -ideal (as the kernel is generated under colimits by the cones of $\mathcal{A}[S] \rightarrow \mathcal{M}[S]$, and tensoring them over \mathbb{Z} with $\mathbb{Z}[T]$ for varying T still lies in the kernel, by evaluating the internal Hom at T). \square

The definition of analytic animated associative rings suggests to define a map $(\mathcal{A}, \mathcal{M}) \rightarrow (\mathcal{B}, \mathcal{N})$ of such to be a map of condensed animated associative rings $\mathcal{A} \rightarrow \mathcal{B}$ together with a natural transformation $\mathcal{M}[S] \rightarrow \mathcal{N}[S]$ linear over this map and commuting with the maps from S .

Following the (updated) discussion in [Sch19, Lecture 7], we instead define a map $(\mathcal{A}, \mathcal{M}) \rightarrow (\mathcal{B}, \mathcal{N})$ to be a map $\mathcal{A} \rightarrow \mathcal{B}$ of condensed animated associative rings such that the forgetful functor $\mathcal{D}_{\geq 0}(\mathcal{B}, \mathcal{N}) \rightarrow \mathcal{D}_{\geq 0}(\mathcal{A})$ takes image in $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$. It is enough to check this condition for $\mathcal{N}[S]$, and in fact for $\pi_0\mathcal{N}[S]$, as these generate the abelian heart, and the categories are determined by the heart.

Given a map $(\mathcal{A}, \mathcal{M}) \rightarrow (\mathcal{B}, \mathcal{N})$ of analytic animated associative rings, the maps $S \rightarrow \mathcal{N}[S]$ extend uniquely to maps $\mathcal{M}[S] \rightarrow \mathcal{N}[S]$ linear over $\mathcal{A} \rightarrow \mathcal{B}$, so one gets a necessarily unique map in the naive sense. The converse is also close to being true: If one has functorial maps $\mathcal{M}[S] \rightarrow \mathcal{N}[S]$ linear over $\mathcal{A} \rightarrow \mathcal{B}$ and commuting with the map from S , then this defines a map $(\mathcal{A}, \mathcal{M}) \rightarrow (\mathcal{B}, \mathcal{N})$ under a mild condition. In fact, it is enough to give functorial maps $\pi_0 \mathcal{M}[S] \rightarrow \pi_0 \mathcal{N}[S]$ linear over $\pi_0 \mathcal{A} \rightarrow \pi_0 \mathcal{B}$ such that for all maps $S \rightarrow \pi_0 \mathcal{A}$ from some extremally disconnected profinite set S , the induced diagram

$$\begin{array}{ccc} \pi_0 \mathcal{M}[S] & \longrightarrow & \pi_0 \mathcal{M}[*] \\ \downarrow & & \downarrow \\ \pi_0 \mathcal{N}[S] & \longrightarrow & \pi_0 \mathcal{N}[*] \end{array}$$

commutes, cf. [Sch19, Proposition 7.14] (whose proof immediately passes to the present situation).

PROPOSITION 12.6. *Let $f : (\mathcal{A}, \mathcal{M}) \rightarrow (\mathcal{B}, \mathcal{N})$ be a map of analytic animated associative rings. The forgetful functor $\mathcal{D}_{\geq 0}(\mathcal{B}, \mathcal{N}) \rightarrow \mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$ admits a left adjoint*

$$- \otimes_{(\mathcal{A}, \mathcal{M})} (\mathcal{B}, \mathcal{N}) : \mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M}) \rightarrow \mathcal{D}_{\geq 0}(\mathcal{B}, \mathcal{N})$$

sending $\mathcal{M}[S]$ to $\mathcal{N}[S]$. This functor is naturally symmetric monoidal when \mathcal{A} and \mathcal{B} and $f : \mathcal{A} \rightarrow \mathcal{B}$ have the structure of condensed animated commutative rings (resp. of a map between such).

REMARK 12.7. We should really encode more functoriality, especially regarding composition. This is best done by defining the relevant (co)Cartesian fibrations and straightforward in the present situation.

PROOF. The existence and description of the left adjoint is clear. The symmetric monoidal structure follows from [NS18, Theorem I.3.6] again. \square

One can base change analytic ring structures along maps of condensed rings, as follows. We emphasize that this proposition works only in the animated setting – the relevant base changes will in general be derived, and we have to keep track of the derived structure.

PROPOSITION 12.8. *Let $(\mathcal{A}, \mathcal{M})$ be an analytic animated associative ring and let $g : \mathcal{A} \rightarrow \mathcal{B}$ be a map of condensed animated associative rings. Then the functor*

$$S \mapsto \mathcal{N}[S] := \mathcal{B}[S] \otimes_{\mathcal{A}} (\mathcal{A}, \mathcal{M})$$

defines an analytic animated associative ring $(\mathcal{B}, \mathcal{N})$.

PROOF. Note that all $\mathcal{N}[S]$ are, as \mathcal{A} -modules, by definition in $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$; thus, for any sifted colimit $N \in \mathcal{D}_{\geq 0}(\mathcal{B})$ of these, forgetting to $\mathcal{D}_{\geq 0}(\mathcal{A})$ defines an object of $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$.

We want to prove

$$\underline{\mathrm{Hom}}_{\mathcal{D}_{\geq 0}(\mathcal{B})}(\mathcal{N}[S], N) = \underline{\mathrm{Hom}}_{\mathcal{D}_{\geq 0}(\mathcal{B})}(\mathcal{B}[S], N).$$

From $N \in \mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$ and the definition of $\mathcal{N}[S]$, we know that

$$\underline{\mathrm{Hom}}_{\mathcal{D}_{\geq 0}(\mathcal{A})}(\mathcal{N}[S], N) = \underline{\mathrm{Hom}}_{\mathcal{D}_{\geq 0}(\mathcal{A})}(\mathcal{B}[S], N).$$

We claim that this implies the desired result formally. Indeed, we first rewrite this as

$$\underline{\mathrm{Hom}}_{\mathcal{D}_{\geq 0}(\mathcal{B})}(\mathcal{N}[S] \otimes_{\mathcal{A}} \mathcal{B}, N) = \underline{\mathrm{Hom}}_{\mathcal{D}_{\geq 0}(\mathcal{B})}(\mathcal{B}[S] \otimes_{\mathcal{A}} \mathcal{B}, N)$$

which moreover implies, by taking $\underline{\mathrm{Hom}}_{\mathcal{D}_{\geq 0}(\mathcal{B})}(M, -)$ for varying $M \in \mathcal{D}_{\geq 0}(\mathcal{B})$, that

$$\underline{\mathrm{Hom}}_{\mathcal{D}_{\geq 0}(\mathcal{B})}(\mathcal{N}[S] \otimes_{\mathcal{A}} M, N) = \underline{\mathrm{Hom}}_{\mathcal{D}_{\geq 0}(\mathcal{B})}(\mathcal{B}[S] \otimes_{\mathcal{A}} M, N)$$

for all such M . This can be rewritten as

$$\underline{\mathrm{Hom}}_{\mathcal{D}_{\geq 0}(\mathcal{B})}(\mathcal{N}[S] \otimes_{\mathcal{B}} (\mathcal{B} \otimes_{\mathcal{A}} M), N) = \underline{\mathrm{Hom}}_{\mathcal{D}_{\geq 0}(\mathcal{B})}(\mathcal{B}[S] \otimes_{\mathcal{B}} (\mathcal{B} \otimes_{\mathcal{A}} M), N).$$

But the objects of the form $\mathcal{B} \otimes_{\mathcal{A}} M \in \mathcal{D}_{\geq 0}(\mathcal{B})$ for $M \in \mathcal{D}_{\geq 0}(\mathcal{B})$ generate all of $\mathcal{D}_{\geq 0}(\mathcal{B})$ under colimits; for example, there is a resolution of \mathcal{B} of the form

$$\dots \rightarrow \mathcal{B} \otimes_{\mathcal{A}} \mathcal{B} \otimes_{\mathcal{A}} \mathcal{B} \rightarrow \mathcal{B} \otimes_{\mathcal{A}} \mathcal{B} \rightarrow \mathcal{B}$$

(as this forms a split simplicial diagram). Thus, also

$$\underline{\mathrm{Hom}}_{\mathcal{D}_{\geq 0}(\mathcal{B})}(\mathcal{N}[S], N) = \underline{\mathrm{Hom}}_{\mathcal{D}_{\geq 0}(\mathcal{B})}(\mathcal{B}[S], N).$$

□

In the third appendix to this lecture, we show that if $(\mathcal{A}, \mathcal{M})$ is an analytic animated associative ring, then the map $\mathcal{A} \rightarrow \mathcal{M}[*]$ is naturally a map of condensed animated associative rings; in particular, $\mathcal{M}[*]$ acquires the structure of a condensed animated associative ring. In fact, $\mathcal{M}[*]$ is initial in the ∞ -category of condensed animated associative rings \mathcal{B} under \mathcal{A} with the property that $\mathcal{B} \in \mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M}) \subset \mathcal{D}_{\geq 0}(\mathcal{A})$.

DEFINITION 12.9. An analytic animated associative ring $(\mathcal{A}, \mathcal{M})$ is normalized if the map $\mathcal{A} \rightarrow \mathcal{M}[*]$ is an isomorphism.

The preceding discussion shows that for any analytic animated associative ring $(\mathcal{A}, \mathcal{M})$, there is an initial analytic animated associative ring under it, given by applying the previous proposition to $(\mathcal{A}, \mathcal{M})$ and the map $\mathcal{A} \rightarrow \mathcal{M}[*]$. Normalization does not affect $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$.

In the commutative case, passing to normalized rings is actually more subtle. The problem is that we have to endow $\mathcal{M}[*]$ with the structure of a condensed animated commutative ring, and, as analyzed in the third appendix, this requires that the Sym^n -operations on $\mathcal{D}_{\geq 0}(\mathcal{A})$ descend to $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$. As proved there, this is implied by insisting on the following property in the definition of condensed animated commutative rings $(\mathcal{A}, \mathcal{M})$:

DEFINITION 12.10. An analytic animated commutative ring is a condensed animated commutative ring \mathcal{A} together with a functor $\mathcal{M} : S \mapsto \mathcal{M}[S]$ as above so that $(\mathcal{A}, \mathcal{M})$ is an analytic animated associative ring, and such that for all primes p , the Frobenius map $\phi_p : \mathcal{A} \rightarrow \mathcal{A}/\mathbb{L}p$ induces a map of analytic animated associative rings $\phi_p : (\mathcal{A}, \mathcal{M}) \rightarrow (\mathcal{A}/\mathbb{L}p, \mathcal{M}/\mathbb{L}p)$. A map of analytic animated commutative rings $(\mathcal{A}, \mathcal{M}_{\mathcal{A}}) \rightarrow (\mathcal{B}, \mathcal{M}_{\mathcal{B}})$ is a map of condensed animated commutative rings $\mathcal{A} \rightarrow \mathcal{B}$ that induces a map of analytic animated associative rings.

This condition on Frobenius is analyzed in the second appendix, and shown there to be automatic in a number of situations; it may well be the case that it is automatic in general. By the results of the third appendix, one can normalize analytic animated commutative rings.²⁶

In the following, we will only work with normalized analytic rings.

²⁶If one works in the context of connective \mathbb{E}_{∞} -rings, this subtlety about Frobenius does not come up, as then it is clear that normalizations exist (as the tensor unit in a symmetric monoidal ∞ -category is an \mathbb{E}_{∞} -ring); also all the rest of the discussion of these lectures immediately adapts to that context.

DEFINITION 12.11. Let AnRing be the ∞ -category of normalized analytic animated commutative rings.

Let us analyze this ∞ -category. For the following proposition, it is again critical to work in the animated context.

PROPOSITION 12.12. *The ∞ -category AnRing admits all small colimits. The initial object is \mathbb{Z} with $S \mapsto \mathbb{Z}[S]$. Sifted colimits commute with the functor $(\mathcal{A}, \mathcal{M}) \mapsto \mathcal{M}[S]$ to $\mathcal{D}_{\geq 0}(\text{Cond}(\text{Ab}))$. Pushouts are computed as follows:*

Let $(\mathcal{B}, \mathcal{M}_{\mathcal{B}}) \leftarrow (\mathcal{A}, \mathcal{M}_{\mathcal{A}}) \rightarrow (\mathcal{C}, \mathcal{M}_{\mathcal{C}})$ be a diagram in AnRing . The pushout $(\mathcal{B}, \mathcal{M}_{\mathcal{B}}) \otimes_{(\mathcal{A}, \mathcal{M}_{\mathcal{A}})} (\mathcal{C}, \mathcal{M}_{\mathcal{C}}) = (\mathcal{E}, \mathcal{M}_{\mathcal{E}})$ of this diagram in AnRing exists. It can be defined as the normalization of an analytic ring structure on

$$\mathcal{B} \otimes_{\mathcal{A}} \mathcal{C}.$$

The corresponding functor

$$\mathcal{D}_{\geq 0}(\mathcal{E}, \mathcal{M}_{\mathcal{E}}) \rightarrow \mathcal{D}_{\geq 0}(\mathcal{B} \otimes_{\mathcal{A}} \mathcal{C})$$

is fully faithful, with essential image given by all $C \in \mathcal{D}_{\geq 0}(\mathcal{B} \otimes_{\mathcal{A}} \mathcal{C})$ whose images in $\mathcal{D}_{\geq 0}(\mathcal{B})$ and $\mathcal{D}_{\geq 0}(\mathcal{C})$ lie in $\mathcal{D}_{\geq 0}(\mathcal{B}, \mathcal{M}_{\mathcal{B}})$ and $\mathcal{D}_{\geq 0}(\mathcal{C}, \mathcal{M}_{\mathcal{C}})$.

The left adjoint

$$- \otimes_{\mathcal{B} \otimes_{\mathcal{A}} \mathcal{C}} (\mathcal{E}, \mathcal{M}_{\mathcal{E}}) : \mathcal{D}_{\geq 0}(\mathcal{B} \otimes_{\mathcal{A}} \mathcal{C}) \rightarrow \mathcal{D}_{\geq 0}(\mathcal{E}, \mathcal{M}_{\mathcal{E}})$$

is given by the sequential colimit

$$- \rightarrow - \otimes_{\mathcal{B}} (\mathcal{B}, \mathcal{M}_{\mathcal{B}}) \rightarrow (- \otimes_{\mathcal{B}} (\mathcal{B}, \mathcal{M}_{\mathcal{B}})) \otimes_{\mathcal{C}} (\mathcal{C}, \mathcal{M}_{\mathcal{C}}) \rightarrow ((- \otimes_{\mathcal{B}} (\mathcal{B}, \mathcal{M}_{\mathcal{B}})) \otimes_{\mathcal{C}} (\mathcal{C}, \mathcal{M}_{\mathcal{C}})) \otimes_{\mathcal{B}} (\mathcal{B}, \mathcal{M}_{\mathcal{B}}) \rightarrow \dots$$

(where each functor is considered as an endofunctor of $\mathcal{D}_{\geq 0}(\mathcal{B} \otimes_{\mathcal{A}} \mathcal{C})$).

In particular, pushouts in AnRing are in general subtle, as the operations of “ $\mathcal{M}_{\mathcal{B}}$ -completion” $- \otimes_{\mathcal{B}} (\mathcal{B}, \mathcal{M}_{\mathcal{B}})$ and “ \mathcal{C} -completion” $- \otimes_{\mathcal{C}} (\mathcal{C}, \mathcal{M}_{\mathcal{C}})$ do not in general commute.

PROOF. The statement about the initial object is clear. For the statement about sifted colimits, let $(\mathcal{A}_{\bullet}, \mathcal{M}_{\bullet})$ be any sifted diagram in AnRing .²⁷ Then for any i and S , the sifted colimit of $\mathcal{M}_{\bullet}[S]$ is an object of $\mathcal{D}_{\geq 0}(\text{colim } \mathcal{A}_{\bullet})$ whose restriction to $\mathcal{D}_{\geq 0}(\mathcal{A}_i)$ lies in $\mathcal{D}_{\geq 0}(\mathcal{A}_i, \mathcal{M}_i)$, and thus for any sifted colimit C of $\mathcal{A}_{\bullet}[S]$'s,

$$\underline{\text{Hom}}_{\mathcal{D}_{\geq 0}(\mathcal{A}_i)}(\mathcal{M}_i[S], C) = \underline{\text{Hom}}_{\mathcal{D}_{\geq 0}(\mathcal{A}_i)}(\mathcal{A}_i[S], C)$$

or equivalently

$$\underline{\text{Hom}}_{\mathcal{D}_{\geq 0}(\text{colim } \mathcal{A}_{\bullet})}(\mathcal{M}_i[S] \otimes_{\mathcal{A}_i} \text{colim } \mathcal{A}_{\bullet}, C) = \underline{\text{Hom}}_{\mathcal{D}_{\geq 0}(\text{colim } \mathcal{A}_{\bullet})}(\text{colim } \mathcal{A}_{\bullet}[S], C).$$

Taking the limit over i of this statement and using that the colimit of $\mathcal{M}_i[S] \otimes_{\mathcal{A}_i} \text{colim } \mathcal{A}_{\bullet}$ is just $\text{colim } \mathcal{M}_{\bullet}[S]$ as the colimit is sifted, we get the desired result.

For pushouts, we note that any object in $(\mathcal{E}', \mathcal{M}_{\mathcal{E}'}) \in \text{AnRing}$ with compatible maps from $(\mathcal{B}, \mathcal{M}_{\mathcal{B}}) \leftarrow (\mathcal{A}, \mathcal{M}_{\mathcal{A}}) \rightarrow (\mathcal{C}, \mathcal{M}_{\mathcal{C}})$ will admit a natural map $\mathcal{D}_{\geq 0}(\mathcal{E}', \mathcal{M}_{\mathcal{E}'}) \rightarrow \mathcal{D}_{\geq 0}(\mathcal{B} \otimes_{\mathcal{A}} \mathcal{C})$ whose essential image lies in the full ∞ -subcategory of all objects whose restrictions to the two factors lie in $\mathcal{D}_{\geq 0}(\mathcal{B}, \mathcal{M}_{\mathcal{B}})$ and $\mathcal{D}_{\geq 0}(\mathcal{C}, \mathcal{M}_{\mathcal{C}})$. Thus, we see that it is enough to prove that the recipee in the statement defines an (a priori unnormalized) analytic ring structure on $\mathcal{B} \otimes_{\mathcal{A}} \mathcal{C}$; but this is easy to see (using, for example, Proposition 12.20). Passing to the normalization gives the desired pushout. \square

²⁷On first reading, assume that it is a filtered colimit.

One instance where the base change in AnRing is simple is for induced analytic ring structures as in Proposition 12.8.

Another such instance is the following class of maps.

DEFINITION 12.13. A map $f : (\mathcal{A}, \mathcal{M}_{\mathcal{A}}) \rightarrow (\mathcal{B}, \mathcal{M}_{\mathcal{B}})$ in AnRing is steady if for all maps $g : (\mathcal{A}, \mathcal{M}_{\mathcal{A}}) \rightarrow (\mathcal{C}, \mathcal{M}_{\mathcal{C}})$ in AnRing, the functor $M \mapsto M \otimes_{\mathcal{B}} (\mathcal{B}, \mathcal{M}_{\mathcal{B}})$ preserves the full ∞ -subcategory of $\mathcal{D}_{\geq 0}(\mathcal{C} \otimes_{\mathcal{A}} \mathcal{B})$ of all objects whose restriction to \mathcal{C} lies in $\mathcal{D}_{\geq 0}(\mathcal{C}, \mathcal{M}_{\mathcal{C}})$.

Equivalently, for all $M \in \mathcal{D}_{\geq 0}(\mathcal{C}, \mathcal{M}_{\mathcal{C}})$, the object

$$M \otimes_{\mathcal{A}} (\mathcal{B}, \mathcal{M}_{\mathcal{B}}) = M \otimes_{(\mathcal{A}, \mathcal{M}_{\mathcal{A}})} (\mathcal{B}, \mathcal{M}_{\mathcal{B}})$$

which is a priori an object of $\mathcal{D}_{\geq 0}(\mathcal{C} \otimes_{\mathcal{A}} \mathcal{B})$, lies in $\mathcal{D}_{\geq 0}(\mathcal{C}, \mathcal{M}_{\mathcal{C}})$ when restricted to \mathcal{C} . As its restriction to \mathcal{B} lies in $\mathcal{D}_{\geq 0}(\mathcal{B}, \mathcal{M}_{\mathcal{B}})$ by construction, this will then actually define an object of

$$\mathcal{D}_{\geq 0}((\mathcal{B}, \mathcal{M}_{\mathcal{B}}) \otimes_{(\mathcal{A}, \mathcal{M}_{\mathcal{A}})} (\mathcal{C}, \mathcal{M}_{\mathcal{C}})).$$

In fact, the condition that f is steady means precisely that for all g the colimit

$$- \rightarrow - \otimes_{\mathcal{C}} (\mathcal{C}, \mathcal{M}_{\mathcal{C}}) \rightarrow (- \otimes_{\mathcal{C}} (\mathcal{C}, \mathcal{M}_{\mathcal{C}})) \otimes_{\mathcal{B}} (\mathcal{B}, \mathcal{M}_{\mathcal{B}}) \rightarrow ((- \otimes_{\mathcal{C}} (\mathcal{C}, \mathcal{M}_{\mathcal{C}})) \otimes_{\mathcal{B}} (\mathcal{B}, \mathcal{M}_{\mathcal{B}})) \otimes_{\mathcal{C}} (\mathcal{C}, \mathcal{M}_{\mathcal{C}}) \rightarrow \dots$$

stabilizes at $(- \otimes_{\mathcal{C}} (\mathcal{C}, \mathcal{M}_{\mathcal{C}})) \otimes_{\mathcal{B}} (\mathcal{B}, \mathcal{M}_{\mathcal{B}})$. The pushout $(\mathcal{E}, \mathcal{M}_{\mathcal{E}})$ of $(\mathcal{C}, \mathcal{M}_{\mathcal{C}}) \leftarrow (\mathcal{A}, \mathcal{M}_{\mathcal{A}}) \rightarrow (\mathcal{B}, \mathcal{M}_{\mathcal{B}})$ is then given by the functor

$$\mathcal{M}_{\mathcal{E}} : S \mapsto \mathcal{M}_{\mathcal{C}}[S] \otimes_{(\mathcal{A}, \mathcal{M}_{\mathcal{A}})} (\mathcal{B}, \mathcal{M}_{\mathcal{B}}).$$

Yet another equivalent characterization is the following.

PROPOSITION 12.14. A map $f : (\mathcal{A}, \mathcal{M}_{\mathcal{A}}) \rightarrow (\mathcal{B}, \mathcal{M}_{\mathcal{B}})$ in AnRing is steady if and only if for all pushout diagrams

$$\begin{array}{ccc} (\mathcal{E}, \mathcal{M}_{\mathcal{E}}) & \xleftarrow{\tilde{f}} & (\mathcal{C}, \mathcal{M}_{\mathcal{C}}) \\ \tilde{g} \uparrow & & \uparrow g \\ (\mathcal{B}, \mathcal{M}_{\mathcal{B}}) & \xleftarrow{f} & (\mathcal{A}, \mathcal{M}_{\mathcal{A}}) \end{array}$$

and all $C \in \mathcal{D}(\mathcal{C}, \mathcal{M}_{\mathcal{C}})$, the base change map

$$(C|_{\mathcal{A}}) \otimes_{(\mathcal{A}, \mathcal{M}_{\mathcal{A}})} (\mathcal{B}, \mathcal{M}_{\mathcal{B}}) \rightarrow (C \otimes_{(\mathcal{C}, \mathcal{M}_{\mathcal{C}})} (\mathcal{E}, \mathcal{M}_{\mathcal{E}}))|_{\mathcal{B}}$$

is an isomorphism.

In the geometric language of the next lecture, this means that in the cartesian diagram

$$\begin{array}{ccc} \text{AnSpec}(\mathcal{E}, \mathcal{M}_{\mathcal{E}}) & \xrightarrow{\tilde{f}} & \text{AnSpec}(\mathcal{C}, \mathcal{M}_{\mathcal{C}}) \\ \downarrow \tilde{g} & & \downarrow g \\ \text{AnSpec}(\mathcal{B}, \mathcal{M}_{\mathcal{B}}) & \xrightarrow{f} & \text{AnSpec}(\mathcal{A}, \mathcal{M}_{\mathcal{A}}), \end{array}$$

the map

$$f^* g_* C \rightarrow \tilde{g}_* \tilde{f}^* C$$

is an isomorphism (using the usual notation for pullbacks and pushforwards of quasicohherent sheaves). Thus, pushforward commutes with base change along steady maps, and this property characterizes steady maps.

PROOF. As both sides commute with filtered colimits in C , it suffices to check on $C = \mathcal{M}_C[S]$, which amounts to the condition above. \square

The class of steady maps has good properties.

PROPOSITION 12.15. *The class of steady maps is stable under base change and composition. If a steady map $f : (\mathcal{A}, \mathcal{M}_{\mathcal{A}}) \rightarrow (\mathcal{B}, \mathcal{M}_{\mathcal{B}})$ in AnRing factors over a map $f' : (\mathcal{A}', \mathcal{M}_{\mathcal{A}'}) \rightarrow (\mathcal{B}, \mathcal{M}_{\mathcal{B}})$, then f' is also steady. Moreover, the class of steady maps is closed under all colimits (in $\text{Fun}(\Delta^1, \text{AnRing})$).*

PROOF. Exercise. \square

DEFINITION 12.16. A map $f : (\mathcal{A}, \mathcal{M}_{\mathcal{A}}) \rightarrow (\mathcal{B}, \mathcal{M}_{\mathcal{B}})$ in AnRing is a localization if the forgetful functor $\mathcal{D}_{\geq 0}(\mathcal{B}, \mathcal{M}_{\mathcal{B}}) \rightarrow \mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M}_{\mathcal{A}})$ is fully faithful. The map f is a steady localization if it is a localization and steady.

EXERCISE 12.17. Show that if a map $f : (\mathcal{A}, \mathcal{M}_{\mathcal{A}}) \rightarrow (\mathcal{B}, \mathcal{M}_{\mathcal{B}})$ in AnRing is a localization, then the induced map $(\mathcal{B}, \mathcal{M}_{\mathcal{B}}) \rightarrow (\mathcal{B}, \mathcal{M}_{\mathcal{B}}) \otimes_{(\mathcal{A}, \mathcal{M}_{\mathcal{A}})} (\mathcal{B}, \mathcal{M}_{\mathcal{B}})$ is an isomorphism. (Hint: First establish that any base change of a localization is again a localization, by noting that $(\mathcal{B}, \mathcal{M}_{\mathcal{B}})$ can be regarded as the normalization of a non-normalized analytic ring structure on \mathcal{A} .) If f is steady, prove the converse. (Hint: The condition of fully faithfulness is equivalent to the statement that for all $M \in \mathcal{D}_{\geq 0}(\mathcal{B}, \mathcal{M}_{\mathcal{B}})$, the map $M \rightarrow M \otimes_{(\mathcal{A}, \mathcal{M}_{\mathcal{A}})} (\mathcal{B}, \mathcal{M}_{\mathcal{B}})$ is an equivalence. Rewrite the latter as $M \otimes_{(\mathcal{B}, \mathcal{M}_{\mathcal{B}})} ((\mathcal{B}, \mathcal{M}_{\mathcal{B}}) \otimes_{(\mathcal{A}, \mathcal{M}_{\mathcal{A}})} (\mathcal{B}, \mathcal{M}_{\mathcal{B}}))$.)

With this definition, we have the following descent of modules.

PROPOSITION 12.18. *Let $f_i : (\mathcal{A}, \mathcal{M}) \rightarrow (\mathcal{A}_i, \mathcal{M}_i)$, $i \in I$, be a finite family of steady localizations in AnRing such that the functor $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M}) \rightarrow \prod_i \mathcal{D}_{\geq 0}(\mathcal{A}_i, \mathcal{M}_i)$ is conservative. Let \mathcal{C}_I be the category of nonempty subsets of I ; we get a functor $\mathcal{C}_I \rightarrow \text{AnRing}$ sending any $J \subset I$ to $(\mathcal{A}_J, \mathcal{M}_J) := \otimes_{i \in J, /(\mathcal{A}, \mathcal{M})} (\mathcal{A}_i, \mathcal{M}_i)$.*

Then for any $M \in \mathcal{D}(\mathcal{A}, \mathcal{M})$ the natural map

$$M \rightarrow \lim_{J \in \mathcal{C}_I} (M \otimes_{(\mathcal{A}, \mathcal{M})} (\mathcal{A}_J, \mathcal{M}_J))$$

is an isomorphism, and

$$\mathcal{D}(\mathcal{A}, \mathcal{M}) \rightarrow \lim_{J \in \mathcal{C}_I} \mathcal{D}(\mathcal{A}_J, \mathcal{M}_J)$$

is an equivalence.

PROOF. This is formal, cf. (proof of) [Sch19, Proposition 10.5]. The key point is that steady localizations commute with any base change. We also use that localizations commute with finite limits, for which we have to pass from the prestable ∞ -category $\mathcal{D}_{\geq 0}$ to the stable ∞ -category \mathcal{D} (where they can be reinterpreted as finite colimits). \square

Next week, we will discuss some examples. In particular, we will see that for the analytic rings corresponding to Huber pairs (A, A^+) , the condition of $(A, A^+)_{\blacksquare} \rightarrow (B, B^+)_{\blacksquare}$ being steady is closely related to the condition that $A \rightarrow B$ is adic in Huber's sense, i.e. given compatible rings of definition $A_0 \subset A$, $B_0 \subset B$, $A_0 \rightarrow B_0$, if $I \subset A_0$ is an ideal of definition, then $IB_0 \subset B_0$ is an ideal of definition. Thus the present discussion mirrors a standard discussion on Huber pairs: In that situation, general pushouts do not exist, but they do exist when one of the maps is adic.

Appendix to Lecture XII: Topological invariance of analytic ring structures

In this appendix, we characterize analytic ring structures in terms of the full sub- ∞ -category $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M}) \subset \mathcal{D}_{\geq 0}(\mathcal{A})$. As an application, we prove that analytic ring structures are invariant under nilpotent thickenings. First, we have the following result.

PROPOSITION 12.19. *Let \mathcal{A} be a condensed animated associative ring. The collection of full sub- ∞ -categories $\mathcal{D} \subset \mathcal{D}_{\geq 0}(\mathcal{A})$ stable under all limits and colimits is in natural bijection with the collection of all full subcategories $\mathcal{C} \subset \text{Mod}_{\pi_0 \mathcal{A}}^{\text{cond}}$ stable under all limits, colimits, and extensions, via sending \mathcal{D} to the intersection with $\text{Mod}_{\pi_0 \mathcal{A}}^{\text{cond}}$, and \mathcal{C} to the full sub- ∞ -category \mathcal{D} of all $C \in \mathcal{D}_{\geq 0}(\mathcal{A})$ such that all $H_i(C) \in \mathcal{C}$ for $i \geq 0$.*

PROOF. If $\mathcal{D} \subset \mathcal{D}_{\geq 0}(\mathcal{A})$ is stable under all limits and colimits, then in particular for any $C \in \mathcal{D}$, also $\tau_{\geq 1} C \in \mathcal{D}$ as the suspension of the loops of C , and thus $H_0(C)[0] \in \mathcal{D}$. It follows that defining \mathcal{C} to be the intersection of \mathcal{D} with $\text{Mod}_{\pi_0 \mathcal{A}}^{\text{cond}}$, one has $C \in \mathcal{D}$ if and only if all $H_i(C) \in \mathcal{C}$. Indeed, the forward direction follows by inductively applying this argument, and the converse follows as \mathcal{D} is stable under extensions (as these can be written as cofibers) and (Postnikov) limits. Moreover, stability of \mathcal{D} under limits and colimits implies stability of \mathcal{C} under limits, colimits, and extensions.

Conversely, if $\mathcal{C} \subset \text{Mod}_{\pi_0 \mathcal{A}}^{\text{cond}}$ is stable under limits, colimits, and extensions, then defining \mathcal{D} as in the statement of the proposition, one sees that \mathcal{D} is stable under fibres and cofibres, as well as infinite direct sums or infinite direct products, and thus under all limits and colimits. The intersection of \mathcal{D} with $\text{Mod}_{\pi_0 \mathcal{A}}^{\text{cond}}$ is by definition \mathcal{C} , giving the result. \square

Now we can characterize analytic ring structures.

PROPOSITION 12.20. *Let \mathcal{A} be a condensed animated associative ring. A full sub- ∞ -category $\mathcal{D} \subset \mathcal{D}_{\geq 0}(\mathcal{A})$ is of the form $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$ for a necessarily unique analytic ring structure $(\mathcal{A}, \mathcal{M})$ on \mathcal{A} if and only if satisfies the following conditions:*

- (1) *The sub- ∞ -category $\mathcal{D} \subset \mathcal{D}_{\geq 0}(\mathcal{A})$ is stable under all limits and colimits.*
- (2) *The sub- ∞ -category $\mathcal{D} \subset \mathcal{D}_{\geq 0}(\mathcal{A})$ is stable under $\underline{\text{Hom}}_{\mathcal{D}_{\geq 0}(\text{Cond}(\text{Ab}))}(\mathbb{Z}[S], -)$ for any extremally disconnected profinite set S .*
- (3) *The inclusion $\mathcal{D} \subset \mathcal{D}_{\geq 0}(\mathcal{A})$ admits a left adjoint.*

We remark that the existence of the left adjoint is automatic modulo set-theoretic issues.

PROOF. This is clear in the forward direction. Conversely, we can define $\mathcal{M}[S]$ as the image of $\mathcal{A}[S]$ under the left adjoint (guaranteed by (3)). Then we formally know that for all $C \in \mathcal{D}$,

$$\text{Hom}_{\mathcal{D}_{\geq 0}(\mathcal{A})}(\mathcal{M}[S], C) \rightarrow \text{Hom}_{\mathcal{D}_{\geq 0}(\mathcal{A})}(\mathcal{A}[S], C)$$

is an isomorphism of anima. By condition (2), we actually see that it is an isomorphism of condensed anima. As by (1), \mathcal{D} contains all sifted colimits of $\mathcal{M}[S]$'s, we see that this defines an analytic ring structure. \square

As an application, one can show that analytic ring structures are independent of the animated structure.

PROPOSITION 12.21. *Let $f : \mathcal{A} \rightarrow \mathcal{B}$ be map of condensed animated associative rings such that $\pi_0 \mathcal{A} \rightarrow \pi_0 \mathcal{B}$ is an isomorphism.*

Then there is a bijective correspondence between analytic ring structures $(\mathcal{A}, \mathcal{M})$ on \mathcal{A} and analytic ring structures $(\mathcal{B}, \mathcal{N})$ on \mathcal{B} . In the forward direction, this takes $(\mathcal{A}, \mathcal{M})$ to an induced analytic ring structure

$$\mathcal{N}[S] := \mathcal{B}[S] \otimes_{\mathcal{A}} (\mathcal{A}, \mathcal{M}).$$

REMARK 12.22. Everything applies, with identical proofs, to general \mathbb{E}_1 -algebras in condensed connective spectra. For example one can define a solid analytic ring structure on \mathbb{S} , and for $0 < r < 1$ and r -liquid analytic ring structure on $S[T^{-1}]$ (whose normalization is some condensed \mathbb{E}_∞ -ring $\mathbb{S}((T))_{>r}$ that seems subtle to define directly).

PROOF. By Proposition 12.8, the functor is well-defined. To prove that it this is a bijection, we can reduce to the case where f is the projection $\mathcal{A} \rightarrow \pi_0\mathcal{A}$.

The hearts of $\mathcal{D}_{\geq 0}(\mathcal{A})$ and $\mathcal{D}_{\geq 0}(\pi_0\mathcal{A})$ agree, and the functor identifies the corresponding hearts of $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$ and $\mathcal{D}_{\geq 0}(\pi_0\mathcal{A} \otimes_{\mathcal{A}} (\mathcal{A}, \mathcal{M}))$. To see that this defines the desired bijection, we need to see that for any analytic ring structure $(\pi_0\mathcal{A}, \mathcal{M}')$ on $\pi_0\mathcal{A}$ with corresponding heart $\mathcal{C} \subset \text{Mod}_{\pi_0\mathcal{A}}^{\text{cond}}$, one can define an analytic ring structure $(\mathcal{A}, \mathcal{M})$ such that $C \in \mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$ if and only if all $H_i(C) \in \mathcal{C}$. By Proposition 12.19, this defines some full sub- ∞ -category $\mathcal{D} \subset \mathcal{D}_{\geq 0}(\mathcal{A})$ stable under all limits and colimits. We need to check conditions (2) and (3) of Proposition 12.20. For condition (2), we can assume (by Postnikov limits and filtrations) that $C = X[j]$ for some $X \in \mathcal{C}$ and $j \geq 0$, in which case C comes from a $\pi_0\mathcal{A}$ -module, and the result follows from that case.

For condition (3), we need to see that the functor $M \mapsto M(S)$ on \mathcal{D} is representable. We prove by induction on i that there is some i -truncated $\mathcal{M}_i[S] \in \mathcal{D}$ with a map $\mathcal{A}[S] \rightarrow \mathcal{M}_i[S]$ such that

$$\text{Hom}_{\mathcal{D}}(\mathcal{M}_i[S], M) \rightarrow M(S)$$

is an isomorphism of anima for all i -truncated $M \in \mathcal{D}$. For $i = 0$, we can take $\mathcal{M}_0[S] = \pi_0\mathcal{M}'[S]$. Given $\mathcal{M}_i[S]$, let N_i be the cofiber of $\mathcal{A}[S] \rightarrow \mathcal{M}_i[S]$ in $\mathcal{D}_{\geq 0}(\mathcal{A})$ and let

$$N'_i = N_i \otimes_{\mathcal{A}} (\pi_0\mathcal{A}, \mathcal{M}').$$

As the forgetful functor $\mathcal{D}_{\geq 0}(\pi_0\mathcal{A}, \mathcal{M}') \rightarrow \mathcal{D}_{\geq 0}(\mathcal{A})$ lands in \mathcal{D} , we can regard N'_i as an object of \mathcal{D} . Then we let $\mathcal{M}_{i+1}[S]$ be the fibre of $\mathcal{M}_i[S] \rightarrow N_i \rightarrow N'_i$, and we note that $\mathcal{A}[S] \rightarrow \mathcal{M}_i[S]$ factors naturally over $\mathcal{M}_{i+1}[S]$.

To see that

$$\text{Hom}_{\mathcal{D}}(\mathcal{M}_{i+1}[S], M) \rightarrow M(S)$$

is an isomorphism of anima for all $i + 1$ -truncated $M \in \mathcal{D}$, it suffices to check for $M = X[i + 1]$ with $X \in \mathcal{C}$. Note that for such M , we have the fibre sequence

$$\text{Hom}_{\mathcal{D}_{\geq 0}(\mathcal{A})}(N_i, M) \rightarrow \text{Hom}_{\mathcal{D}}(\mathcal{M}_i[S], M) \rightarrow M(S)$$

where the right-most term is concentrated in degree $i + 1$, while the middle term is isomorphic to $M(S)$ after applying $\tau_{\geq 1}$, by induction. It follows that $\text{Hom}_{\mathcal{D}_{\geq 0}(\mathcal{A})}(N_i, M)$ sits in degree 0. This can also be written as

$$\text{Hom}_{\mathcal{D}_{\geq 0}(\pi_0\mathcal{A})}(N_i \otimes_{\mathcal{A}} \pi_0\mathcal{A}, M)$$

which again for all $M = X[i + 1]$ with $X \in \mathcal{C}$ sits in degree 0. It follows that $N'_i = N_i \otimes_{\mathcal{A}} (\pi_0\mathcal{A}, \mathcal{M}')$ lies in $\mathcal{D}_{\geq i+1}(\pi_0\mathcal{A}, \mathcal{M}')$, and we still have

$$\text{Hom}_{\mathcal{D}_{\geq 0}(\pi_0\mathcal{A})}(N'_i, M) = \text{Hom}_{\mathcal{D}_{\geq 0}(\pi_0\mathcal{A})}(N_i \otimes_{\mathcal{A}} \pi_0\mathcal{A}, M).$$

On the other hand, the map

$$\mathrm{Hom}_{\mathcal{D}_{\geq 0}(\pi_0\mathcal{A})}(N'_i, M) \rightarrow \mathrm{Hom}_{\mathcal{D}_{\geq 0}(\mathcal{A})}(N'_i, M)$$

is also an isomorphism, as $N'_i \in \mathcal{D}_{\geq i+1}$ and M is $i+1$ -truncated.

In summary, we see that the natural map

$$\mathrm{Hom}_{\mathcal{D}_{\geq 0}(\mathcal{A})}(N_i, M) \leftarrow \mathrm{Hom}_{\mathcal{D}}(N'_i, M)$$

is an isomorphism, where both sides are concentrated in degree 0. We defined $\mathcal{M}_{i+1}[S]$ as the fiber of $\mathcal{M}_i[S] \rightarrow N'_i$. Recall that N'_i was in $\mathcal{D}_{\geq i+1}$, in particular its π_0 vanishes. This implies that we get a cofiber sequence

$$\mathrm{Hom}_{\mathcal{D}}(N'_i, M) \rightarrow \mathrm{Hom}_{\mathcal{D}}(\mathcal{M}_i[S], M) \rightarrow \mathrm{Hom}_{\mathcal{D}}(\mathcal{M}_{i+1}[S], M)$$

and comparing with

$$\mathrm{Hom}_{\mathcal{D}_{\geq 0}(\mathcal{A})}(N_i, M) \rightarrow \mathrm{Hom}_{\mathcal{D}}(\mathcal{M}_i[S], M) \rightarrow M(S)$$

gives the result. □

One can moreover prove invariance under nilpotent thickenings.

PROPOSITION 12.23. *Let $f : \mathcal{A} \rightarrow \mathcal{B}$ be a map of condensed animated associative rings such that $\pi_0\mathcal{A} \rightarrow \pi_0\mathcal{B}$ is surjective with nilpotent kernel $I \subset \mathcal{A}$. Then analytic ring structures $(\mathcal{A}, \mathcal{M})$ on \mathcal{A} are in bijection with analytic ring structures $(\mathcal{B}, \mathcal{N})$ on \mathcal{B} , via taking induced analytic ring structures.*

PROOF. By Proposition 12.21, we can assume that \mathcal{A} and \mathcal{B} are 0-truncated. We may also assume that $I^2 = 0$ by induction. Let us define the inverse functor, so take an analytic ring structure $(\mathcal{B}, \mathcal{N})$ on \mathcal{B} , corresponding to some abelian category $\mathcal{C}_{\mathcal{B}} \subset \mathrm{Mod}_{\mathcal{B}}^{\mathrm{cond}}$. Let $\mathcal{C} \subset \mathrm{Mod}_{\mathcal{A}}^{\mathrm{cond}}$ be the subcategory of all $M \in \mathrm{Mod}_{\mathcal{A}}^{\mathrm{cond}}$ such that M/IM and IM , which lie naturally in $\mathrm{Mod}_{\mathcal{B}}^{\mathrm{cond}}$, are in $\mathcal{C}_{\mathcal{B}}$. This contains $\mathcal{C}_{\mathcal{B}}$ and is stable under all limits, colimits, and extensions; it can also be characterized as the category generated by $\mathcal{C}_{\mathcal{B}}$ under extensions. By Proposition 12.19, it corresponds to some full sub- ∞ -category $\mathcal{D} \subset \mathcal{D}_{\geq 0}(\mathcal{A})$. We need to check the criteria of Proposition 12.20. Condition (2) can again be checked for $C = X[j]$ for some $X \in \mathcal{C}$ and $j \geq 0$, where it follows by filtering in terms of two objects from $\mathcal{C}_{\mathcal{B}}$.

It remains to check condition (3). This follows from the argument in the proof of Proposition 12.21 with a slightly refined induction. Namely, besides the class of i -truncated objects of \mathcal{D} , we also consider the class of i -truncated objects $C \in \mathcal{D}$ such that $H_i(C)$ is killed by I . Then we start the induction with this restricted class of 0-truncated objects, then go to all 0-truncated objects, then to the restricted class of 1-truncated objects, to all 1-truncated objects, etc. . □

Appendix to Lecture XII: Frobenius

Fix a prime p . Consider a condensed animated commutative \mathbb{F}_p -algebra \mathcal{A} , equipped as an associative with an analytic ring structure $(\mathcal{A}, \mathcal{M})$. One may wonder whether the Frobenius $\phi : \mathcal{A} \rightarrow \mathcal{A}$ induces a map of analytic rings $(\mathcal{A}, \mathcal{M}) \rightarrow (\mathcal{A}, \mathcal{M})$. This condition is stable under all colimits. The goal of this appendix is to show that it is true in some important cases. We denote by C_p the cyclic group of order p .

PROPOSITION 12.24. *Extend the functor $S \mapsto \mathcal{M}[S]$ to all profinite sets by using simplicial resolutions by extremally disconnected sets. Assume the following condition:*

ASSUMPTION 12.25. *For all profinite sets S with C_p -action and fixed points $S_0 = S^{C_p}$, the natural map*

$$\mathcal{M}[S_0]^{tC_p} \rightarrow \mathcal{M}[S]^{tC_p}$$

is an isomorphism.

Then ϕ induces a map of analytic rings $(\mathcal{A}, \mathcal{M}) \rightarrow (\mathcal{A}, \mathcal{M})$.

Moreover, the assumption is satisfied in the following two situations:

- (1) *if $(\mathcal{A}, \mathcal{M})$ is an algebra over $\mathbb{F}_{p, \blacksquare}$;*
- (2) *if for all profinite sets S , $\mathcal{M}[S]$ is m -truncated for some m .*

By (the version for animated rings of) [Sch19, Proposition 7.14], it is enough to construct functorial ϕ -linear maps

$$\phi_S : \pi_0 \mathcal{M}[S] \rightarrow \pi_0 \mathcal{M}[S]$$

for all extremally disconnected sets S , subject to the condition that for any map $f : T \rightarrow \mathcal{A}$ from an extremally disconnected profinite set, letting $\tilde{f} : \mathcal{M}[T] \rightarrow \mathcal{M}[*]$ be the induced extension, the diagram

$$\begin{array}{ccc} \pi_0 \mathcal{M}[T] & \xrightarrow{\phi_T} & \pi_0 \mathcal{M}[T] \\ \downarrow \pi_0 \tilde{f} & & \downarrow \pi_0 \tilde{f} \\ \pi_0 \mathcal{M}[*] & \xrightarrow{\phi} & \pi_0 \mathcal{M}[*] \end{array}$$

commutes.

To do this, recall that for any abelian group M , there is a natural linear map

$$M \rightarrow \check{H}^0(C_p, M \otimes \dots \otimes M) = \text{cofib}(\text{Nm} : (M \otimes \dots \otimes M)_{C_p} \rightarrow (M \otimes \dots \otimes M)^{C_p}),$$

where $M \otimes \dots \otimes M$ (with p tensor factors) is equipped with the C_p -action permuting the tensor factors cyclically, and we take the 0-th Tate cohomology of the C_p -action, i.e. the cofiber of the norm map from the coinvariants to the invariants. This map is induced by the (non-additive) map

$$M \rightarrow (M \otimes \dots \otimes M)^{C_p} : m \mapsto m \otimes \dots \otimes m,$$

noting that it becomes additive after passing to the cofiber of the norm map.

Applying this to the condensed abelian group $\pi_0 \mathcal{M}[S]$, we get a natural map

$$\pi_0 \mathcal{M}[S] \rightarrow \check{H}^0(C_p, \pi_0 \mathcal{M}[S] \otimes \dots \otimes \pi_0 \mathcal{M}[S])$$

that we can compose with the C_p -equivariant multiplication map

$$\pi_0 \mathcal{M}[S] \otimes \dots \otimes \pi_0 \mathcal{M}[S] \rightarrow \pi_0 \mathcal{M}[S^p]$$

to get a functorial map

$$\pi_0\mathcal{M}[S] \rightarrow \check{H}^0(C_p, \pi_0\mathcal{M}[S^p]).$$

Now note that there is a map $\pi_0\mathcal{M}[S] \rightarrow \pi_0\mathcal{M}[S^p]$ induced by the diagonal inclusion $S \hookrightarrow S^p$. We would like to use that the map $\check{H}^0(C_p, \pi_0\mathcal{M}[S]) \rightarrow \check{H}^0(C_p, \pi_0\mathcal{M}[S^p])$ is an isomorphism. As C_p acts trivially on the p -torsion module $\pi_0\mathcal{M}[S]$, this would give the desired map

$$\pi_0\mathcal{M}[S] \rightarrow \check{H}^0(C_p, \pi_0\mathcal{M}[S]) = \pi_0\mathcal{M}[S].$$

But we have passed to π_0 too early; we need to stay derived for a little longer to get the passage from $\mathcal{M}[S^p]$ to $\mathcal{M}[S]$. For this, recall that one can define Tate cohomology in general for a spectrum X with C_p -action, as

$$X^{tC_p} = \text{cofib}(\text{Nm} : X_{hC_p} \rightarrow X^{hC_p})$$

the cofibre of the norm map from homotopy orbits to homotopy fixed points, cf. e.g. [NS18, Chapter I]. The map $M \rightarrow \check{H}^0(C_p, M \otimes \dots \otimes M)$ generalizes to the following. Let X be any spectrum in the sense of stable homotopy theory, and consider $X \otimes \dots \otimes X$, the p -fold tensor product, with the cyclic C_p -action. Then there is a unique functorial lax symmetric monoidal map

$$X \rightarrow (X \otimes \dots \otimes X)^{tC_p},$$

the so-called Tate diagonal. (This map exists only for spectra, not in $\mathcal{D}(\text{Ab})$.)

Now following the above, we get a functorial map

$$\mathcal{M}[S] \rightarrow (\mathcal{M}[S] \otimes \dots \otimes \mathcal{M}[S])^{tC_p} \rightarrow \mathcal{M}[S^p]^{tC_p}.$$

Again, we have the map $\mathcal{M}[S] \rightarrow \mathcal{M}[S^p]$ induced by the diagonal $S \rightarrow S^p$, and now the assumption guarantees that this induces an isomorphism after applying $-^{tC_p}$.

In total, we have produced a functorial map $\mathcal{M}[S] \rightarrow \mathcal{M}[S]^{tC_p}$. Now is the time to pass to π_0 : First, on the target, where we get a natural map to $(\pi_0\mathcal{M}[S])^{tC_p}$. In the resulting map

$$\mathcal{M}[S] \rightarrow (\pi_0\mathcal{M}[S])^{tC_p}$$

we take π_0 , which gives the desired functorial map

$$\pi_0\mathcal{M}[S] \rightarrow \pi_0\mathcal{M}[S].$$

We need to see that this is linear over $\phi : \pi_0\mathcal{A} \rightarrow \pi_0\mathcal{A}$. But $\mathcal{M}[S] \rightarrow \mathcal{M}[S^p]^{tC_p} \cong \mathcal{M}[S]^{tC_p}$ is linear over the Tate-valued Frobenius $\mathcal{A} \rightarrow \mathcal{A}^{tC_p}$ by construction, and the Tate-valued Frobenius induces the usual Frobenius on π_0 .

It remains to see that for any map $f : T \rightarrow \mathcal{A}$ from some extremally disconnected T , inducing a map $\tilde{f} : \mathcal{M}[T] \rightarrow \mathcal{M}[*]$, the diagram

$$\begin{array}{ccc} \pi_0\mathcal{M}[T] & \xrightarrow{\phi_T} & \pi_0\mathcal{M}[T] \\ \downarrow \pi_0\tilde{f} & & \downarrow \pi_0\tilde{f} \\ \pi_0\mathcal{M}[*] & \xrightarrow{\phi} & \pi_0\mathcal{M}[*] \end{array}$$

commutes. For this, we note that ϕ_T has the property that the composite

$$\pi_0\mathcal{M}[T] \xrightarrow{\phi_T} \pi_0\mathcal{M}[T] \rightarrow \check{H}^0(C_p, \pi_0\mathcal{M}[T^p])$$

is the naive map ϕ'_T constructed in the beginning. Note that $\pi_0\mathcal{M}[*] \rightarrow \check{H}^0(C_p, \pi_0\mathcal{M}[*])$ is an isomorphism. We see that it suffices to see that the diagram

$$\begin{array}{ccc} \pi_0\mathcal{M}[T] & \xrightarrow{\phi'_T} & \check{H}^0(C_p, \pi_0\mathcal{M}[T^p]) \\ \downarrow \pi_0\tilde{f} & & \downarrow \pi_0\tilde{f}_p \\ \pi_0\mathcal{M}[*] & \xrightarrow{\phi} & \check{H}^0(C_p, \pi_0\mathcal{M}[*]) \end{array}$$

commutes, where $\tilde{f}_p : \mathcal{M}[T^p] \rightarrow \mathcal{M}[*]$ is the p -fold tensor product of \tilde{f} in $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$. This is the composite of the commutative diagram

$$\begin{array}{ccc} \pi_0\mathcal{M}[T] & \xrightarrow{\phi'_T} & \check{H}^0(C_p, \pi_0\mathcal{M}[T] \otimes_{\pi_0\mathcal{A}} \dots \otimes_{\pi_0\mathcal{A}} \pi_0\mathcal{M}[T]) \\ \downarrow \pi_0\tilde{f} & & \downarrow \pi_0\tilde{f}^{\otimes p} \\ \pi_0\mathcal{M}[*] & \xrightarrow{\phi} & \check{H}^0(C_p, \pi_0\mathcal{M}[*] \otimes_{\pi_0\mathcal{A}} \dots \otimes_{\pi_0\mathcal{A}} \mathcal{M}[*]) \end{array}$$

(by functoriality of $M \rightarrow \check{H}^0(C_p, M \otimes_A \dots \otimes_A M)$) and the diagram

$$\begin{array}{ccc} \check{H}^0(C_p, \pi_0\mathcal{M}[T] \otimes_{\pi_0\mathcal{A}} \dots \otimes_{\pi_0\mathcal{A}} \pi_0\mathcal{M}[T]) & \longrightarrow & \check{H}^0(C_p, \pi_0\mathcal{M}[T^p]) \\ \downarrow \pi_0\tilde{f}^{\otimes p} & & \downarrow \pi_0\tilde{f}_p \\ \check{H}^0(C_p, \pi_0\mathcal{M}[*] \otimes_{\pi_0\mathcal{A}} \dots \otimes_{\pi_0\mathcal{A}} \mathcal{M}[*]) & \longrightarrow & \check{H}^0(C_p, \pi_0\mathcal{M}[*]). \end{array}$$

For commutativity of this diagram, it suffices to prove that

$$\begin{array}{ccc} \pi_0\mathcal{M}[T] \otimes_{\pi_0\mathcal{A}} \dots \otimes_{\pi_0\mathcal{A}} \pi_0\mathcal{M}[T] & \longrightarrow & \pi_0\mathcal{M}[T^p] \\ \downarrow \pi_0\tilde{f}^{\otimes p} & & \downarrow \pi_0\tilde{f}_p \\ \pi_0\mathcal{M}[*] \otimes_{\pi_0\mathcal{A}} \dots \otimes_{\pi_0\mathcal{A}} \mathcal{M}[*] & \longrightarrow & \pi_0\mathcal{M}[*]. \end{array}$$

commutes. But this is a diagram in condensed $\pi_0\mathcal{A}$ -modules, and the target $\pi_0\mathcal{M}[*]$ lies in $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$, so the result follows from \tilde{f}_p being, by definition, the image of $\tilde{f}^{\otimes p}$ in $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$.

It remains to show that the assumption is satisfied if \mathcal{A} is an algebra over $\mathbb{F}_{p,\blacksquare}$, or if all $\mathcal{M}[S]$ are truncated. For this, we note that we can write the cofiber of

$$\mathcal{M}[S_0]^{tC_p} \rightarrow \mathcal{M}[S]^{tC_p}$$

as the limit of

$$\dots \rightarrow \overline{\mathcal{M}}[\overline{S}][-2n] \xrightarrow{\beta} \dots \xrightarrow{\beta} \overline{\mathcal{M}}[\overline{S}][-2] \xrightarrow{\beta} \overline{\mathcal{M}}[\overline{S}]$$

along a certain map $\beta : \overline{\mathcal{M}}[\overline{S}][-2] \rightarrow \overline{\mathcal{M}}[\overline{S}]$, where \overline{S} is the quotient of the pointed profinite set S/S_0 by the free (in pointed profinite sets) C_p -action, and $\overline{\mathcal{M}}$ is the extension of \mathcal{M} to pointed profinite sets (taking $(T, *)$ to $\mathcal{M}[T]/\mathcal{M}[*]$). This comes from a general formula for X^{tC_p} as a limit of $X_{hC_p}[-2n]$ when $X \in \mathcal{D}(\text{Ab})$.

Moreover, by functoriality of the whole situation, the map β base changes. In the case of $\mathbb{F}_{p,\blacksquare}$, it is equal to 0 as then $\mathcal{M}[S]$ is projective for all profinite sets S ; thus, it is also 0 for any $(\mathcal{A}, \mathcal{M})$ over $\mathbb{F}_{p,\blacksquare}$, and hence the limit above vanishes.

On the other hand, if $\overline{\mathcal{M}}[\overline{S}]$ is truncated, then the limit vanishes as the terms become more and more coconnective.

Appendix to Lecture XII: Normalizations of analytic animated rings

The goal of this appendix is to indicate the proof of the following result.

PROPOSITION 12.26. *Let $(\mathcal{A}, \mathcal{M})$ be an analytic animated associative (resp. commutative) ring. Consider the ∞ -category of condensed animated associative (resp. commutative) rings \mathcal{B} with a map from \mathcal{A} such that $\mathcal{B} \in \mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M}) \subset \mathcal{D}_{\geq 0}(\mathcal{A})$. This has an initial object, whose underlying object in $\mathcal{D}_{\geq 0}(\mathcal{A})$ is $\mathcal{M}[*]$.*

First, we prove this in the associative case. We use that the ∞ -category of condensed animated associative rings \mathcal{B}/\mathcal{A} is monadic over the $\mathcal{D}_{\geq 0}(\mathcal{A})$, as follows from the ∞ -categorical version of the Barr–Beck theorem [?, Theorem 4.7.0.3]. Here, the relevant left adjoint functor can be defined by animation. More precisely, consider the category of pairs of maps of associative rings $A \rightarrow B$, with its forgetful functor to the category of pairs (A, M) of an associative ring A and an A -module M . This admits a left adjoint, sending M to the free associative algebra

$$T_A(M) = A \oplus M \otimes_{\mathbb{Z}} A \oplus M \otimes_{\mathbb{Z}} M \otimes_{\mathbb{Z}} A \oplus \dots$$

Animating this adjunction defines an adjunction between maps of animated associative rings $A \rightarrow B$ and the ∞ -category of pairs of an animated ring A with an animated A -module M . We can then also pass to condensed objects, and to the fibre over the given condensed animated ring \mathcal{A} . This shows that the relevant monad T is given by a monad structure on the functor $M \mapsto \mathcal{A} \oplus M \otimes_{\mathbb{Z}} \mathcal{A} \oplus M \otimes_{\mathbb{Z}} M \otimes_{\mathbb{Z}} \mathcal{A} \oplus \dots$ on $\mathcal{D}_{\geq 0}(\mathcal{A})$.

Now we observe that if $M \rightarrow N$ is a map in $\mathcal{D}_{\geq 0}(\mathcal{A})$ that is sent to an isomorphism in $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$ under $-\otimes_{\mathcal{A}}(\mathcal{A}, \mathcal{M})$, then also $T(M) \rightarrow T(N)$ is sent to an isomorphism in $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$, by the formula for T , and using that the kernel of the localization is stable under $-\otimes_{\mathbb{Z}} C$ for all $C \in \mathcal{D}_{\geq 0}(\text{Cond}(\text{Ab}))$. This implies formally that T descends to a monad $T_{\mathcal{M}}$ on $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$ such that T -algebras whose underlying object lies in $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$ are equivalent to $T_{\mathcal{M}}$ -algebras. Moreover, $-\otimes_{\mathcal{A}}(\mathcal{A}, \mathcal{M})$ defines a left adjoint to the fully faithful forgetful functor from $T_{\mathcal{M}}$ -algebras to T -algebras. This proves the desired result.

In the commutative case, we argue in the same way, but we observe that the relevant monad structure is now induced from the functor

$$M \mapsto A \oplus M \oplus \text{Sym}^2 M \oplus \dots \oplus \text{Sym}^n M \oplus \dots$$

in the case of usual rings and modules. (See [Lur18, Section 25.2.2, Construction 25.2.2.6] for a more careful discussion.) In particular, we note that if \mathcal{A} is a condensed animated commutative ring, then by animating $\text{Sym}^n : (A, M) \mapsto (A, \text{Sym}^n M)$, passing to condensed objects, and the fibre over the condensed animated ring \mathcal{A} , there is a natural functor Sym^n on $\mathcal{D}_{\geq 0}(\mathcal{A})$. In order to argue as above, we need the following lemma.

LEMMA 12.27. *Let $(\mathcal{A}, \mathcal{M})$ be an analytic animated commutative ring. Let $M \rightarrow N$ be a map in $\mathcal{D}_{\geq 0}(\mathcal{A})$ that becomes an isomorphism after $-\otimes_{\mathcal{A}}(\mathcal{A}, \mathcal{M})$. Then $\text{Sym}^n M \rightarrow \text{Sym}^n N$ becomes an isomorphism after $-\otimes_{\mathcal{A}}(\mathcal{A}, \mathcal{M})$ for all $n \geq 0$.*

PROOF. Let Q be the cofibre of $M \rightarrow N$. By [Lur18, Construction 25.2.5.4], $\text{Sym}^n N$ admits a filtration whose graded pieces are

$$\text{Sym}^n M, \text{Sym}^{n-1} M \otimes Q, \dots, \text{Sym}^{n-i} M \otimes \text{Sym}^i Q, \dots, \text{Sym}^n Q.$$

(This can be proved by animating the corresponding statement for usual (finite projective) modules.) Thus, it suffices to see that for all Q in the kernel K of $- \otimes_{\mathcal{A}} (\mathcal{A}, \mathcal{M})$, also $\mathrm{Sym}^i Q$ lies in K for all $i > 0$.

We argue by induction on i . Then we see by induction, using the above filtration, that the composite functor $F : Q \mapsto \mathrm{Sym}^i Q \otimes_{\mathcal{A}} (\mathcal{A}, \mathcal{M})$ from K to $\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$ commutes with cofibres. In particular, $F(Q[n]) \cong F(Q)[n]$ for all n , or

$$\mathrm{Sym}^i Q \otimes_{\mathcal{A}} (\mathcal{A}, \mathcal{M}) \cong \mathrm{Sym}^i(Q[n])[-n] \otimes_{\mathcal{A}} (\mathcal{A}, \mathcal{M})$$

for all n . Passing to the colimit over n , we note that the functor $Q \mapsto \mathrm{colim}_n \mathrm{Sym}^i(Q[n])[-n]$ is an exact endofunctor of $\mathcal{D}_{\geq 0}(\mathcal{A})$. One can show that this functor vanishes unless i is power p^m of a prime p (the first “Goodwillie derivative of Sym^i ”, in this situation going back to Dold–Puppe, [DP61]), and in that case it is filtered by copies of the functor $Q \mapsto Q \otimes_{\mathcal{A}} \mathcal{A}/p$ where the map $\mathcal{A} \rightarrow \mathcal{A}/p$ is the Frobenius $a \mapsto a^{p^m}$. Now the result follows from the definition of analytic animated commutative rings. \square

13. Lecture XIII: Analytic spaces

Finally, we define analytic spaces. In this lecture, we use the term analytic ring to refer to objects of AnRing , i.e. analytic animated commutative rings $(\mathcal{A}, \mathcal{M})$ with $\mathcal{A} = \mathcal{M}[*]$.²⁸

Consider the ∞ -category of functors from AnRing to anima , i.e. the ∞ -category of presheaves of anima on $\text{AnRing}^{\text{op}}$. By the Yoneda lemma, this admits $\text{AnRing}^{\text{op}}$ as a full subcategory, via the functor

$$(\mathcal{A}, \mathcal{M}) \mapsto \text{AnSpec}(\mathcal{A}, \mathcal{M}) : (\mathcal{B}, \mathcal{N}) \mapsto \text{Hom}_{\text{AnRing}}((\mathcal{A}, \mathcal{M}), (\mathcal{B}, \mathcal{N})).$$

We endow $\text{AnRing}^{\text{op}}$ with the Grothendieck topology generated by the families of maps

$$\{\text{AnSpec}(\mathcal{A}_i, \mathcal{M}_i) \rightarrow \text{AnSpec}(\mathcal{A}, \mathcal{M})\}$$

whenever $(\mathcal{A}, \mathcal{M}) \rightarrow (\mathcal{A}_i, \mathcal{M}_i)$ is a finite family of steady localizations such that

$$\mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M}) \rightarrow \prod_i \mathcal{D}_{\geq 0}(\mathcal{A}_i, \mathcal{M}_i)$$

is conservative. We note that pullbacks of such finite families of maps are of the same form.

PROPOSITION 13.1. *For any analytic ring $(\mathcal{A}, \mathcal{M})$, the functor $\text{AnSpec}(\mathcal{A}, \mathcal{M})$ is a sheaf on $\text{AnRing}^{\text{op}}$.*

PROOF. This follows directly from Proposition 12.18. □

If $f : \mathcal{F} \rightarrow \mathcal{G}$ is a map of functors from AnRing to anima , we write $\mathcal{F} \subset \mathcal{G}$ if for all analytic rings $(\mathcal{A}, \mathcal{M})$, all fibres of

$$\mathcal{F}(\mathcal{A}, \mathcal{M}) \rightarrow \mathcal{G}(\mathcal{A}, \mathcal{M})$$

are either empty or contractible.

DEFINITION 13.2. Let $(\mathcal{A}, \mathcal{M})$ be an analytic ring. A steady subspace of $\text{AnSpec}(\mathcal{A}, \mathcal{M})$ is a subfunctor $U \subset \text{AnSpec}(\mathcal{A}, \mathcal{M})$ such that the natural map

$$\text{colim}_{\text{AnSpec}(\mathcal{A}', \mathcal{M}') \subset U} \text{AnSpec}(\mathcal{A}', \mathcal{M}') \rightarrow U$$

is an isomorphism, where the colimit runs over all steady localizations $(\mathcal{A}, \mathcal{M}) \rightarrow (\mathcal{A}', \mathcal{M}')$ for which $\text{AnSpec}(\mathcal{A}', \mathcal{M}') \subset \text{AnSpec}(\mathcal{A}, \mathcal{M})$ factors over U .

REMARK 13.3. As the colimit is taken over a non-filtered index category, one might worry that it may produce undesired outcomes. This is not so: Compute the colimit first as presheaves. Then for all analytic rings $(\mathcal{B}, \mathcal{N})$ with a fixed map to U , the fibre product computes the colimit of a point over all $\text{AnSpec}(\mathcal{A}', \mathcal{M}')$ contained in U and such that $\text{AnSpec}(\mathcal{B}, \mathcal{N}) \rightarrow U$ factors over $\text{AnSpec}(\mathcal{A}', \mathcal{M}')$. If nonempty, index category is cofiltered (as with any two steady localizations, it contains their product), so the geometric realization of the index category is contractible, and hence this colimit is itself a point or empty. Now note that if \mathcal{G} is a sheaf of anima and $\mathcal{F} \subset \mathcal{G}$ is a presheaf of anima (the inclusion meaning that all fibres are empty or contractible), then the sheafification \mathcal{F}' of \mathcal{F} still has the property that $\mathcal{F}' \rightarrow \mathcal{G}$ has empty or contractible fibres; it is the subsheaf of \mathcal{G} of all sections that locally lift to \mathcal{F} . To see this, note that sheafification commutes with taking fibres, so it is enough to recall that the sheafification of a presheaf whose values are empty or a point has values which still are empty or a point.

²⁸One can consider a variant using analytic connective \mathbb{E}_∞ -rings instead.

This argument shows that $U \subset \text{AnSpec}(\mathcal{A}, \mathcal{M})$ is a steady subspace if and only if there is some collection $(\mathcal{A}, \mathcal{M}) \rightarrow (\mathcal{A}_i, \mathcal{M}_i)_i$ of steady localizations such that U is the image of

$$\bigsqcup_i \text{AnSpec}(\mathcal{A}_i, \mathcal{M}_i) \rightarrow \text{AnSpec}(\mathcal{A}, \mathcal{M})(\mathcal{B}, \mathcal{N}).$$

One verifies immediately that pullbacks of steady subspaces are steady, and conversely $U \subset \text{AnSpec}(\mathcal{A}, \mathcal{M})$ is a steady subspace if and only if the pullbacks $U_i \subset \text{AnSpec}(\mathcal{A}_i, \mathcal{M}_i)$ are steady subspaces for all terms in a finite cover of $(\mathcal{A}, \mathcal{M})$ by steady localizations $(\mathcal{A}_i, \mathcal{M}_i)$. Thus, the following definition has reasonable properties.

DEFINITION 13.4. An inclusion $\mathcal{F} \subset \mathcal{G}$ of sheaves of anima on AnRing is a steady subspace if for all analytic rings $(\mathcal{A}, \mathcal{M})$ the fibre product

$$\mathcal{F} \times_{\mathcal{G}} \text{AnSpec}(\mathcal{A}, \mathcal{M}) \rightarrow \text{AnSpec}(\mathcal{A}, \mathcal{M})$$

is a steady subspace.

Finally, we can define analytic spaces.

DEFINITION 13.5. An analytic space is a sheaf of anima X on AnRing such that the map

$$\text{colim}_{\text{AnSpec}(\mathcal{A}, \mathcal{M}) \subset X} \text{AnSpec}(\mathcal{A}, \mathcal{M}) \rightarrow X$$

is an isomorphism, where the colimit runs over all affine steady subspaces $\text{AnSpec}(\mathcal{A}, \mathcal{M}) \subset X$.

Again, one might worry that the colimit does not behave well as the index category is not well-behaved. However, we have the following result, saying that we could equivalently define analytic spaces X as those sheaves that can be covered by steady subspaces of the form $\text{AnSpec}(\mathcal{A}, \mathcal{M}) \subset X$.

PROPOSITION 13.6. *If X is a sheaf of anima on AnRing such that the map*

$$\text{colim}_{\text{AnSpec}(\mathcal{A}, \mathcal{M}) \subset X} \text{AnSpec}(\mathcal{A}, \mathcal{M}) \rightarrow X$$

is surjective, where as above the colimit runs over all affine steady subspaces $\text{AnSpec}(\mathcal{A}, \mathcal{M}) \subset X$, then X is an analytic space, i.e. the map is an isomorphism.

In particular, any steady subspace of an analytic space is again an analytic space.

PROOF. Assume first that X admits an injection into some affine $\tilde{X} = \text{AnSpec}(\tilde{\mathcal{A}}, \tilde{\mathcal{M}})$; let us call such X quasi-affine for the rest of this proof. In that case, fibres products over X can also be computed as fibre products over \tilde{X} . For any $(\mathcal{B}, \mathcal{N})$ mapping to X , the fibre of the displayed map is computed by the colimit over a point over the index category of all $\text{AnSpec}(\mathcal{A}, \mathcal{M}) \subset X$ over which $\text{AnSpec}(\mathcal{B}, \mathcal{N}) \rightarrow X$, which will be cofiltered in this case. Thus, this colimit is either empty or a point, which gives the result in this case.

At this point, we already see that any steady subspace of some quasi-affine X is again an analytic space (and clearly quasi-affine). Now let X be general and consider the colimit

$$\text{colim}_{Y \subset X} Y \rightarrow X$$

where the colimit runs over all quasi-affine analytic spaces Y that are steady subspaces of X . By the same argument as in the first paragraph, this map is an isomorphism. Finally, the map

$$\text{colim}_{\text{AnSpec}(\mathcal{A}, \mathcal{M}) \subset X} \text{AnSpec}(\mathcal{A}, \mathcal{M}) \rightarrow \text{colim}_{Y \subset X} Y$$

is an isomorphism, by comparing both with

$$\operatorname{colim}_{\operatorname{AnSpec}(\mathcal{A}, \mathcal{M}) \subset Y \subset X} \operatorname{AnSpec}(\mathcal{A}, \mathcal{M}) :$$

This is equal to the first, as the set of possible Y is cofiltered, and it is equal to the second as each Y can be written as $\operatorname{colim}_{\operatorname{AnSpec}(\mathcal{A}, \mathcal{M}) \subset Y} \operatorname{AnSpec}(\mathcal{A}, \mathcal{M})$. \square

By Proposition 12.18, we can define an ∞ -category of quasicoherent sheaves on X .

DEFINITION 13.7. Let X be an analytic space. The ∞ -category of quasicoherent sheaves on X is

$$\mathcal{D}(X) := \lim_{\operatorname{AnSpec}(\mathcal{A}, \mathcal{M}) \subset X} \mathcal{D}(\mathcal{A}, \mathcal{M}).$$

Equivalently, the association $(\mathcal{A}, \mathcal{M}) \mapsto \mathcal{D}(\mathcal{A}, \mathcal{M})$ defines a sheaf of ∞ -categories on $\operatorname{AnRing}^{\text{op}}$, and $\mathcal{D}(X)$ is the value of this sheaf on X . In particular:

PROPOSITION 13.8. *Let X be an analytic space and let $U_i \subset X$, $i \in I$, be steady subspaces that cover X (i.e. $\bigsqcup_i U_i \rightarrow X$ is a surjection). Let C_I be the category of finite nonempty subsets of I , and consider the functor taking $J \in C_I$ to $U_J = \bigcap_{i \in J} U_i$. Then*

$$\mathcal{D}(X) \rightarrow \lim_{J \in C_I} \mathcal{D}(U_J)$$

is an equivalence of ∞ -categories.

If $X = \operatorname{AnSpec}(\mathcal{A}, \mathcal{M})$ is affine, then $\mathcal{D}(X) \cong \mathcal{D}(\mathcal{A}, \mathcal{M})$.

We see that there is a well-defined ∞ -category of analytic spaces, defined very analogously to schemes, and it comes with an ∞ -category of quasicoherent sheaves, satisfying descent.

It is high time for some examples. First, we need some examples of steady localizations. All of them will be produced by the following criterion.

PROPOSITION 13.9. *Let $f : (\mathcal{A}, \mathcal{M}) \rightarrow (\mathcal{B}, \mathcal{N})$ be a map of analytic rings. Assume that for any extremally disconnected profinite set S and any $M \in \mathcal{D}_{\geq 0}(\mathcal{A}, \mathcal{M})$, the natural map*

$$(\underline{\operatorname{Hom}}_{\mathcal{D}(\mathcal{A})}(\mathcal{A}[S], M) \otimes_{(\mathcal{A}, \mathcal{M})} (\mathcal{B}, \mathcal{N}))(*) \rightarrow (M \otimes_{(\mathcal{A}, \mathcal{M})} (\mathcal{B}, \mathcal{N}))(S)$$

in $\mathcal{D}(\operatorname{Ab})$ is an equivalence. Then f is steady.

PROOF. Consider any analytic ring $(\mathcal{C}, \mathcal{M}_{\mathcal{C}})$ over $(\mathcal{A}, \mathcal{M})$ and let $M \in \mathcal{D}_{\geq 0}(\mathcal{C}, \mathcal{M}_{\mathcal{C}})$. Then

$$(M \otimes_{(\mathcal{A}, \mathcal{M})} (\mathcal{B}, \mathcal{N}))(S) = (\underline{\operatorname{Hom}}_{\mathcal{D}(\mathcal{A})}(\mathcal{A}[S], M) \otimes_{(\mathcal{A}, \mathcal{M})} (\mathcal{B}, \mathcal{N}))(*) = (\underline{\operatorname{Hom}}_{\mathcal{D}(\mathcal{C}, \mathcal{M}_{\mathcal{C}})}(\mathcal{M}_{\mathcal{C}}[S], M) \otimes_{(\mathcal{A}, \mathcal{M})} (\mathcal{B}, \mathcal{N}))(*) .$$

Varying S , the right-hand side of this formula defines a functor from the ∞ -category of \mathcal{C} -modules isomorphic to some $\mathcal{M}_{\mathcal{C}}[S]$ towards anima, commuting with finite products, and thus an object of $\mathcal{D}_{\geq 0}(\mathcal{C}, \mathcal{M}_{\mathcal{C}})$, that necessarily agrees with $M \otimes_{(\mathcal{A}, \mathcal{M})} (\mathcal{B}, \mathcal{N})$. \square

An interesting question is whether an induced analytic ring structure is steady. This leads to the following class of modules. Here and in the following, we write

$$P^{\vee} = \underline{\operatorname{Hom}}_{\mathcal{D}(\mathcal{A})}(P, \mathcal{A}) \in \mathcal{D}(\mathcal{A})$$

for $P \in \mathcal{D}(\mathcal{A})$.

DEFINITION 13.10. Let $(\mathcal{A}, \mathcal{M})$ be an analytic ring. An object $C \in \mathcal{D}(\mathcal{A}, \mathcal{M})$ is nuclear if for all extremally disconnected profinite sets S , the natural map

$$(\mathcal{A}[S]^\vee \otimes_{(\mathcal{A}, \mathcal{M})} C)(*) \rightarrow C(S)$$

in $\mathcal{D}(\text{Ab})$ is an isomorphism.

Let us characterize this class of modules. For the characterization, the following class of maps is critical.

DEFINITION 13.11. Let $(\mathcal{A}, \mathcal{M})$ be an analytic ring. A map $f : P \rightarrow Q$ between compact objects of $\mathcal{D}(\mathcal{A}, \mathcal{M})$ is trace-class if there is some map

$$g : \mathcal{A} \rightarrow P^\vee \otimes_{(\mathcal{A}, \mathcal{M})} Q$$

such that f is the composite

$$P \xrightarrow{1 \otimes g} P \otimes_{(\mathcal{A}, \mathcal{M})} P^\vee \otimes_{(\mathcal{A}, \mathcal{M})} Q \rightarrow Q$$

where the second map contracts the first two factors.

DEFINITION 13.12. Let $(\mathcal{A}, \mathcal{M})$ be an analytic ring. An object $C \in \mathcal{D}(\mathcal{A}, \mathcal{M})$ is basic nuclear if it can be written as a sequential colimit

$$C = \text{colim}(P_0 \rightarrow P_1 \rightarrow \dots)$$

of compact $P_i \in \mathcal{D}(\mathcal{A}, \mathcal{M})$ along trace-class maps.

PROPOSITION 13.13. *Let $(\mathcal{A}, \mathcal{M})$ be an analytic ring and let $C \in \mathcal{D}(\mathcal{A}, \mathcal{M})$. Then C is nuclear if and only if it can be written as a filtered colimit of basic nuclear objects. The class of basic nuclear objects is stable under all countable colimits, and the class of nuclear objects is stable under all colimits.*

PROOF. Assume first that C is basic nuclear, so C is a colimit of compact P_i along trace-class maps. Choose maps $\mathcal{A} \rightarrow P_i^\vee \otimes_{\mathcal{A}} P_{i+1}$ giving rise to these trace-class maps $P_i \rightarrow P_{i+1}$. Let $Q = \mathcal{A}[S]^\vee$. Then nuclearity means that $(Q^\vee \otimes_{(\mathcal{A}, \mathcal{M})} C)(*) \rightarrow \text{Hom}_{\mathcal{D}(\mathcal{A})}(Q, C)$ is an isomorphism (in $\mathcal{D}(\text{Ab})$). Both sides commute with colimits in C , and there are natural backwards maps

$$\text{Hom}_{\mathcal{D}(\mathcal{A})}(Q, P_i) \rightarrow (\underline{\text{Hom}}_{\mathcal{D}(\mathcal{A})}(Q, P_i) \otimes_{(\mathcal{A}, \mathcal{M})} P_i^\vee \otimes_{(\mathcal{A}, \mathcal{M})} P_{i+1})(*) \rightarrow (Q^\vee \otimes_{(\mathcal{A}, \mathcal{M})} P_{i+1})(*),$$

showing that C is nuclear.

Next, we show that the class of basic nuclear C is stable under finite colimits; equivalently, under passage to cones. Let $f : C \rightarrow C'$ be a map of basic nuclear objects, and write C resp. C' as a filtered colimit of P_i resp. P'_i as in the definition. As all P_i are compact, the map $C \rightarrow C'$ can be realized via compatible maps $P_i \rightarrow P'_i$, up to reindexing the P'_i . Let Q_i be the cone of $P_i \rightarrow P'_i$, so the cone of f is the sequential colimit of the Q_i . We claim that $Q_i \rightarrow Q_{i+2}$ is trace-class for all i . To see this, we need to find a map $\mathcal{A} \rightarrow Q_i^\vee \otimes_{(\mathcal{A}, \mathcal{M})} Q_{i+2}$ defining this map. Note that $Q_i^\vee \otimes_{(\mathcal{A}, \mathcal{M})} Q_{i+2}$ is the fibre of

$$(P'_i)^\vee \otimes_{(\mathcal{A}, \mathcal{M})} Q_{i+2} \rightarrow P_i^\vee \otimes_{(\mathcal{A}, \mathcal{M})} Q_{i+2}$$

and to define a map there, it suffices to exhibit a section of $(P'_i)^\vee \otimes_{(\mathcal{A}, \mathcal{M})} P'_{i+2}$ and a section of $P_i^\vee \otimes_{(\mathcal{A}, \mathcal{M})} P_{i+2}$ whose images in $P_i^\vee \otimes_{(\mathcal{A}, \mathcal{M})} P'_{i+2}$ agree. To define this, pick sections

$$\alpha : \mathcal{A} \rightarrow P_i^\vee \otimes_{(\mathcal{A}, \mathcal{M})} P_{i+1}, \quad \beta : \mathcal{A} \rightarrow (P'_{i+1})^\vee \otimes_{(\mathcal{A}, \mathcal{M})} P'_{i+2}$$

witnessing that $P_i \rightarrow P_{i+1}$ and $P'_{i+1} \rightarrow P'_{i+2}$ are trace-class, and take the induced sections of $P_i^\vee \otimes_{(\mathcal{A}, \mathcal{M})} P_{i+2}$ and $P_i^\vee \otimes_{(\mathcal{A}, \mathcal{M})} P'_{i+2}$.

To see stability of basic nuclear C under all countable colimits, it remains to observe that it is stable under countable direct sums, which is clear from the definition.

It is clear from the definition that the class of all nuclear objects is stable under all colimits. Now let C be a general nuclear object. We can look at the ω_1 -filtered diagram of all basic nuclear C' mapping to C (here we use that basic nuclear objects admit countable colimits), and replacing C by the cone of the colimit of this diagram mapping to C , we can assume that there are no maps from basic nuclear C'' 's to C . We need to see that then $C = 0$. If not, there is some compact P_0 with a nonzero map $P_0 \rightarrow C$. As by nuclearity,

$$(P_0^\vee \otimes_{\mathcal{A}} C)(*) = \text{Hom}(P_0, C),$$

and both sides commute with filtered colimits as functors of C , we can find some compact $P_1 \rightarrow C$ and a section of $(P_0^\vee \otimes_{\mathcal{A}} P_1)(*)$ so that $P_0 \rightarrow C$ is the composite of $P_1 \rightarrow C$ with the corresponding trace-class map $P_0 \rightarrow P_1$. Continuing, we find a nonzero map from some basic nuclear $\text{colim}(P_0 \rightarrow P_1 \rightarrow \dots)$ to C , contradiction. \square

The proposition implies that nuclear modules have stronger properties than a priori guaranteed by the definition:

PROPOSITION 13.14. *Let $(\mathcal{A}, \mathcal{M})$ be an analytic ring and let $C \in \mathcal{D}(\mathcal{A}, \mathcal{M})$ be nuclear. Then for all $M \in \mathcal{D}(\mathcal{A}, \mathcal{M})$ and extremally disconnected profinite sets S , the map*

$$(\underline{\text{Hom}}_{\mathcal{D}(\mathcal{A})}(\mathcal{A}[S], M) \otimes_{(\mathcal{A}, \mathcal{M})} C)(*) \rightarrow (M \otimes_{(\mathcal{A}, \mathcal{M})} C)(S)$$

in $\mathcal{D}(\text{Ab})$ is an isomorphism.

In particular, if $\mathcal{A} \rightarrow \mathcal{B}$ is a map of condensed animated rings such that $\mathcal{B} \in \mathcal{D}(\mathcal{A}, \mathcal{M})$ is nuclear, and $(\mathcal{B}, \mathcal{N})$ is the induced analytic ring structure on \mathcal{B} , then $(\mathcal{A}, \mathcal{M}) \rightarrow (\mathcal{B}, \mathcal{N})$ satisfies the criterion from Proposition 13.9, and hence is steady.

PROOF. As both sides commute with colimits in C , we can assume that C is basic nuclear, and write it as the sequential colimit of compact P_i along trace-class maps $P_i \rightarrow P_{i+1}$. In that case, there are backwards maps

$$\begin{aligned} (M \otimes_{(\mathcal{A}, \mathcal{M})} P_i)(S) &= \underline{\text{Hom}}_{\mathcal{D}(\mathcal{A})}(\mathcal{A}[S], M \otimes_{(\mathcal{A}, \mathcal{M})} P_i)(*) \\ &\rightarrow (\underline{\text{Hom}}_{\mathcal{D}(\mathcal{A})}(\mathcal{A}[S], M \otimes_{(\mathcal{A}, \mathcal{M})} P_i) \otimes_{(\mathcal{A}, \mathcal{M})} P_i^\vee \otimes_{(\mathcal{A}, \mathcal{M})} P_{i+1})(*) \\ &\rightarrow (\underline{\text{Hom}}_{\mathcal{D}(\mathcal{A})}(\mathcal{A}[S], M \otimes_{(\mathcal{A}, \mathcal{M})} P_i \otimes_{(\mathcal{A}, \mathcal{M})} P_i^\vee) \otimes_{(\mathcal{A}, \mathcal{M})} P_{i+1})(*) \\ &\rightarrow (\underline{\text{Hom}}_{\mathcal{D}(\mathcal{A})}(\mathcal{A}[S], M) \otimes_{(\mathcal{A}, \mathcal{M})} P_{i+1})(*) \end{aligned}$$

and so passing to the filtered colimit over the P_i gives the desired equivalence.

The final sentence follows directly. \square

We will now verify this condition in the examples listed two lectures ago.

EXAMPLE 13.15. The following are steady localizations of analytic rings.

- (1) If $A \rightarrow B = A[S^{-1}]$ is a usual localization of (discrete) rings, with trivial analytic ring structure, then $A \rightarrow B$ is a steady localization. Indeed, B , like any discrete module, is nuclear over A .

- (2) The map $(\mathbb{Z}[T], \mathbb{Z})_{\blacksquare} \rightarrow (\mathbb{Z}[T], \mathbb{Z}[T])_{\blacksquare}$, corresponding to localization to the subset $\{|T| \leq 1\}$ of the adic space $\mathrm{Spa}(\mathbb{Z}[T], \mathbb{Z})$. Recall that in this case (cf. [Sch19, Lecture VII]),

$$M \otimes_{(\mathbb{Z}[T], \mathbb{Z})_{\blacksquare}} (\mathbb{Z}[T], \mathbb{Z}[T])_{\blacksquare} = \underline{\mathrm{Hom}}_{\mathcal{D}((\mathbb{Z}[T], \mathbb{Z})_{\blacksquare})}((\mathbb{Z}((T^{-1}))/\mathbb{Z}[T])[-1], M).$$

Then

$$(\underline{\mathrm{Hom}}_{\mathcal{D}(\mathbb{Z}[T])}(\mathbb{Z}[T][S], M) \otimes_{(\mathbb{Z}[T], \mathbb{Z})_{\blacksquare}} (\mathbb{Z}[T], \mathbb{Z}[T])_{\blacksquare})(*) = \mathrm{Hom}_{\mathcal{D}(\mathbb{Z}[T])}((\mathbb{Z}((T^{-1}))/\mathbb{Z}[T])[-1] \otimes_{\mathbb{Z}} \mathbb{Z}_{\blacksquare}[S], M)$$

while also

$$(M \otimes_{(\mathbb{Z}[T], \mathbb{Z})_{\blacksquare}} (\mathbb{Z}[T], \mathbb{Z}[T])_{\blacksquare})(S) = \mathrm{Hom}_{\mathcal{D}((\mathbb{Z}[T], \mathbb{Z})_{\blacksquare})}((\mathbb{Z}((T^{-1}))/\mathbb{Z}[T])[-1] \otimes_{\mathbb{Z}} \mathbb{Z}_{\blacksquare}[S], M),$$

giving the desired result.

- (3) Any rational subset of an adic space is of the form $U = \{|f_i| \leq |g| \neq 0\}$ for finitely many elements $f_1, \dots, f_n, g \in A$ (satisfying some condition). In the present situation of analytic rings, we can realize this localization by first inverting g , and then asking the condition $|\frac{f_i}{g}| \leq 1$. Inverting g is handled by a base change of (1), while asking $|\frac{f_i}{g}| \leq 1$ is handled by a base change of (2).
- (4) If $D_1 \subset D_2$ is an inclusion of closed discs in \mathbb{C} , then $\mathcal{O}(D_2) \rightarrow \mathcal{O}(D_1)$ (endowed with the analytic ring structure induced from the p -liquid structure on \mathbb{C} for some chosen $0 < p \leq 1$) is a steady localization. This follows from nuclearity of $\mathcal{O}(D_2) \rightarrow \mathcal{O}(D_1)$.

Regarding (3), let us at least recall how general Huber pairs fit into the present framework.

PROPOSITION 13.16. *There is a fully faithful functor $(A, A^+) \mapsto (A, A^+)_{\blacksquare}$ from the category of Huber pairs (A, A^+) to the ∞ -category AnRing , defined as follows. For a finitely generated \mathbb{Z} -algebra R , let R_{\blacksquare} be the analytic ring structure on R given by*

$$R_{\blacksquare}[S] = \varprojlim_i R[S_i]$$

for a profinite set $S = \varprojlim_i S_i$. For a general discrete \mathbb{Z} -algebra R , let $R_{\blacksquare} = \varinjlim_{R' \rightarrow R} R'_{\blacksquare}$, where the colimit is over all finitely generated \mathbb{Z} -algebras R' mapping to R . Finally,

$$(A, A^+)_{\blacksquare} = \underline{A} \otimes_{A_{\mathrm{disc}}^+} (A_{\mathrm{disc}}^+)_{\blacksquare}$$

is an induced analytic ring structure.

REMARK 13.17. A priori, this induced analytic ring structure might be animated. However, Andreychev has proved that $(A, A^+)_{\blacksquare}[S]$ is always 0-truncated.

The analytic ring $(A, A^+)_{\blacksquare}$ is always normalized; this follows from \underline{A} being a module over $(A_{\mathrm{disc}}^+)_{\blacksquare}$, which can be deduced by writing it as the limit of A/I^n where $I \subset A_0 \subset A$ is an ideal of definition that is also an R -module, where $R \subset A^+$ is any chosen finitely generated subalgebra.

REMARK 13.18. The second part of a Huber pair is an open and integrally closed subring $A^+ \subset A$ consisting of powerbounded elements. One may wonder how these precise conditions arise. One answer is the following: Consider all possible normalized analytic ring structures on \underline{A} admitting a map from $\mathbb{Z}_{\blacksquare}$ and that can be obtained via filtered colimits from pushouts of the localization $(\mathbb{Z}[T], \mathbb{Z})_{\blacksquare} \rightarrow (\mathbb{Z}[T], \mathbb{Z}[T])_{\blacksquare}$. Then this class is canonically in bijection with the open and integrally closed subrings $A^+ \subset A$ contained in the powerbounded elements. Given $A^+ \subset A$, one allows the localization via the pushout of $(\mathbb{Z}[T], \mathbb{Z})_{\blacksquare} \rightarrow (\mathbb{Z}[T], \mathbb{Z}[T])_{\blacksquare}$ along any map $\mathbb{Z}[T] \rightarrow A$ sending T to an element of A^+ .

PROOF. That the functor is well-defined follows from the results of [Sch19] and the result on induced analytic ring structures. For fully faithfulness, let (A, A^+) and (B, B^+) be Huber pairs. By definition of maps of analytic rings, maps from $(A, A^+)_{\blacksquare}$ to $(B, B^+)_{\blacksquare}$ embed into maps from \underline{A} to \underline{B} . As A and B are metrizable, these agree with maps from A to B . We need to see that a map $A \rightarrow B$ sends A^+ into B^+ if and only if it induces a map of analytic rings $(A, A^+)_{\blacksquare} \rightarrow (B, B^+)_{\blacksquare}$. This is clear in the forward direction.

Thus, assume that $f : A \rightarrow B$ is a map of Huber rings that induces a map of analytic rings $(A, A^+)_{\blacksquare} \rightarrow (B, B^+)_{\blacksquare}$, but does not send A^+ into B^+ . Take any $g \in A^+$ such that $f(g) \notin B^+$. We get a map $(\mathbb{Z}[T], \mathbb{Z}[T]) \rightarrow (A, A^+)$ sending T to g ; precomposing with this map, we may assume that $(A, A^+) = (\mathbb{Z}[T], \mathbb{Z}[T])$. Note that the localization from $(\mathbb{Z}[T], \mathbb{Z})_{\blacksquare}$ to $(\mathbb{Z}[T], \mathbb{Z}[T])_{\blacksquare}$ is obtained by killing the algebra object $\mathbb{Z}((T))^{-1}$. Thus, our assumption is that

$$\mathbb{Z}((T^{-1})) \otimes_{(\mathbb{Z}[T], \mathbb{Z})_{\blacksquare}} (B, B^+)_{\blacksquare} = 0.$$

As asking whether an algebra is just asking whether $1 = 0$ in this algebra, and if this happens in a filtered colimit, it happens at a finite stage, we can then find some finitely generated subalgebra $R \subset B^+$ such that

$$\mathbb{Z}((T^{-1})) \otimes_{(\mathbb{Z}[T], \mathbb{Z})_{\blacksquare}} (B, R)_{\blacksquare} = 0.$$

Using the resolution

$$0 \rightarrow \prod_{\mathbb{N}} \mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}[T] \xrightarrow{1\text{-shift} \otimes T} \prod_{\mathbb{N}} \mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}[T] \rightarrow \mathbb{Z}((T^{-1})) \rightarrow 0,$$

this means that the map

$$\prod_{\mathbb{N}} R \otimes_{R_{\blacksquare}} \underline{B} \xrightarrow{1\text{-shift} \otimes g} \prod_{\mathbb{N}} R \otimes_{R_{\blacksquare}} \underline{B}$$

is an isomorphism. We need to see that this implies that g is integral over R .

In other words, we now start with a finitely generated \mathbb{Z} -algebra R , some Huber R -algebra B , and an element $g \in B$. Choose a ring of definition $B_0 \subset B$ containing R and some ideal of definition $I \subset B_0$. Then in particular $\underline{B} = \lim_n B/I^n$ as condensed R -modules. We want to show that if

$$\prod_{\mathbb{N}} R \otimes_{R_{\blacksquare}} \underline{B} \xrightarrow{1\text{-shift} \otimes g} \prod_{\mathbb{N}} R \otimes_{R_{\blacksquare}} \underline{B}$$

is an isomorphism, then g is integral over $R + I$ (which is a subring of B and contained in B^+). For any n ,

$$\prod_{\mathbb{N}} R \otimes_R B/I^n = \varinjlim_{M \subset B/I^n} \prod_{\mathbb{N}} R \otimes_R M = \varinjlim_{M \subset B/I^n} \prod_{\mathbb{N}} M \subset \prod_{\mathbb{N}} R/I^n$$

where the colimit runs over finitely generated R -submodules M of B/I^n . We see that the inverse limit over all n maps to $\prod_{\mathbb{N}} B$, and the image is contained in sequences (b_0, b_1, \dots) such that for all n , the submodule of B/I^n generated by all b_i is finitely generated.

If the map is surjective, then in particular 1 is in the image, which means that we can find such a sequence (b_0, b_1, \dots) in B satisfying $b_0 = 1$, $b_1 - gb_0 = 0$, $b_2 - gb_1 = 0$, etc., i.e. $b_i = g^i$. This sequence needs to have the property that the R -submodule of B/I generated by all g^i is finitely generated. But this means that there is some relation

$$g^n + r_{n-1}g^{n-1} + \dots + r_0 = 0 \in B/I,$$

which as desired shows that g is integral over $R + I$. \square

In the context of adic spaces, one can show that nuclearity is closely related to adic maps; for example, if $A \rightarrow B$ is a map of Tate-Huber rings (i.e. Huber rings admitting a topologically nilpotent unit), then B is always a nuclear A -module, for any choice of A^+ . On the other hand, if $A \rightarrow B$ is not adic, then B is never nuclear as A -module, and in fact one can show that for any choices of A^+ mapping to B^+ , the map $(A, A^+)_{\blacksquare} \rightarrow (B, B^+)_{\blacksquare}$ will not be steady.

14. Lecture XIV: Varia

In this final lecture, we briefly outline a few more ideas. We do not give proofs here; however, they are not especially difficult. (The hard work was in obtaining the examples of the solid analytic ring structure on \mathbb{Z} last semester, and the liquid analytic ring structures on \mathbb{R} this semester.)

First, one may wonder whether one can define analytic spaces more in the spirit of locally ringed topological spaces. This is possible. If X is an analytic space, then we can consider the site $|X|$ of steady subspaces $U \subset X$ of X . This forms a locale. If $X = \text{AnSpec}(\mathcal{A}, \mathcal{M})$ is affine, then a basis is given by the affine steady subspaces, which are quasicompact and quasiseparated. Thus, Deligne’s theorem on the existence of points implies that, if there is only a set (instead of a proper class) of affine steady subspaces of X , then $|X|$ can be identified with the site of open subsets of a spectral space. For general (non-affine) X , it would then be the case that $|X|$ is a locally spectral space.

Moreover, $|X|$ carries a sheaf \mathcal{O}_X of condensed animated commutative rings (obtained by sheafifying its value on affine subspaces), as well as a sheaf $\mathcal{M}_X[S]$ of condensed animated \mathcal{O}_X -modules, for every extremally disconnected profinite set S (functorial in S , taking finite disjoint unions to finite direct sums). We want to define what “locally ringed” means, i.e. we want an analogue of the condition that the stalks of \mathcal{O}_X are local.

DEFINITION 14.1. A locally analytically ringed locale is a triple $(X, \mathcal{O}_X, \mathcal{M}_X)$ consisting of a locale X , a sheaf of condensed animated commutative rings \mathcal{O}_X on X , and a functor $\mathcal{M}_X : S \rightarrow \mathcal{M}_X[S]$ from extremally disconnected profinite sets S to condensed animated \mathcal{O}_X -modules, taking finite disjoint unions to finite direct sums, subject to the following conditions:

- (1) (“analytically ringed”) there is a basis of subsets $U \subset X$ for which $(\mathcal{O}_X(U), \mathcal{M}_X(U))$ is an analytic ring; on this basis, the restriction maps are maps of analytic rings;
- (2) (“stalks are local”) for every $U \subset X$ with a map $(\mathcal{A}, \mathcal{M}) \rightarrow (\mathcal{O}_X(U), \mathcal{M}_X(U))$ from some analytic ring $(\mathcal{A}, \mathcal{M})$, and every steady cover

$$\bigsqcup_i \text{AnSpec}(\mathcal{A}_i, \mathcal{M}_i) \rightarrow \text{AnSpec}(\mathcal{A}, \mathcal{M})$$

there is a cover of U by $U_i \subset U$ such that $(\mathcal{A}, \mathcal{M}) \rightarrow (\mathcal{O}_X(U_i), \mathcal{M}_X(U_i))$ factors over $(\mathcal{A}_i, \mathcal{M}_i)$.

A map $f : (Y, \mathcal{O}_Y, \mathcal{M}_Y) \rightarrow (X, \mathcal{O}_X, \mathcal{M}_X)$ of locally analytically ringed locales is a map $f : Y \rightarrow X$ of locales together with a map $f^* \mathcal{O}_X \rightarrow \mathcal{O}_Y$ of sheaves of condensed animated commutative rings, with the following properties.

- (1) (“analytically ringed map”) If $V \subset Y$ maps into $U \subset X$, then $(\mathcal{O}_X(U), \mathcal{M}_X(U)) \rightarrow (\mathcal{O}_Y(V), \mathcal{M}_Y(V))$ is a map of analytic rings whenever both are.
- (2) (“maps on local rings local”) Moreover, in the same situation, if there is some steady localization

$$(\mathcal{A}, \mathcal{M}) = (\mathcal{O}_X(U), \mathcal{M}_X(U)) \rightarrow (\mathcal{A}', \mathcal{M}')$$

over which $(\mathcal{A}, \mathcal{M}) \rightarrow (\mathcal{O}_Y(V), \mathcal{M}_Y(V))$ factors, then one can find $U' \subset U$ containing the image of V such that $(\mathcal{A}, \mathcal{M}) \rightarrow (\mathcal{O}_X(U'), \mathcal{M}_X(U'))$ factors over $(\mathcal{A}', \mathcal{M}')$.

It is now not hard to prove the following proposition.

PROPOSITION 14.2. *The functor $X \mapsto (|X|, \mathcal{O}_X, \mathcal{M}_X)$ from analytic spaces to locally analytically ringed locales is well-defined and fully faithful. An object lies in the essential image if and only*

if it is locally isomorphic to the locally analytically ringed locale associated to some analytic ring $(\mathcal{A}, \mathcal{M})$.

Using this language, we can explain how to embed complex-analytic spaces into analytic spaces over \mathbb{C} equipped with its p -liquid analytic ring structure $(\mathbb{C}, \mathcal{M}_{<p})$ (for any fixed $0 < p \leq 1$).

PROPOSITION 14.3. *Let (X, \mathcal{O}_X) be a complex-analytic space; so X is a topological space and \mathcal{O}_X is a sheaf of condensed \mathbb{C} -algebras, locally Zariski closed in an open n -dimensional complex ball. Consider the functor taking any analytic space Y over $(\mathbb{C}, \mathcal{M}_{<p})$ to maps of locales $f : |Y| \rightarrow X$ together with a map of sheaves of condensed animated \mathbb{C} -algebras $f^* \mathcal{O}_X \rightarrow \mathcal{O}_Y$. This functor is representable by an analytic space X_p^{an} over $\text{AnSpec}(\mathbb{C}, \mathcal{M}_{<p})$.*

Let us give an example: Let $X = \{z \mid |z| < 1\}$ be the open unit disc in \mathbb{C} . Let $D_r = \{z \mid |z| \leq r\} \subset X$ be the closed discs of radius $r < 1$. Let $\mathcal{O}(D_r)$ be the condensed \mathbb{C} -algebra of overconvergent holomorphic functions on D_r . This is a nuclear algebra over $(\mathbb{C}, \mathcal{M}_{<p})$. Endow $\mathcal{O}(D_r)$ with the analytic ring structure induced from $(\mathbb{C}, \mathcal{M}_{<p})$. Then X_p^{an} is the increasing union of $\text{AnSpec } \mathcal{O}(D_r)$ over all $r < 1$.

In fact, similar results hold true for real-analytic, smooth, or topological manifolds: If (X, \mathcal{O}_X) is one such, then the functor taking an analytic space Y over $(\mathbb{R}, \mathcal{M}_{<p})$ to maps of locales $f : |Y| \rightarrow X$ together with a map of sheaves of condensed animated \mathbb{R} -algebras $f^* \mathcal{O}_X \rightarrow \mathcal{O}_Y$ is representable by an analytic space X_p^{an} over $\text{AnSpec}(\mathbb{R}, \mathcal{M}_{<p})$. The explicit description of the functor is similar, using the affine analytic spaces associated with overconvergent real-analytic, smooth, or continuous functions on a closed ball.

WARNING 14.4. In the topological case, the functor does not land in steady analytic spaces over $\text{AnSpec}(\mathbb{R}, \mathcal{M}_{<p})$, and the functor does not commute with products (it does in the other cases). The issue is that for example $C(S^1, \mathbb{R}) \otimes_{(\mathbb{R}, \mathcal{M}_{<p})}^L C(S^1, \mathbb{R})$ is not given by $C((S^1)^2, \mathbb{R})$ (and non-steadyness comes from non-nuclearity of $C(S^1, \mathbb{R})$).

We see that analytic spaces over $\text{AnSpec}(\mathbb{R}, \mathcal{M}_{<p})$ are able to handle all flavours of real or complex manifolds simultaneously. In fact, nothing is holding one back from even mixing them, taking products of real-analytic manifolds with topological manifolds, etc. . Even more, one can also mix with algebraic varieties:

PROPOSITION 14.5. *For any analytic ring $(\mathcal{A}, \mathcal{M})$, there is a fully faithful functor $X \mapsto X^{\text{an}}$ from schemes X over the (animated) ring $\mathcal{A}(*)$ to analytic spaces over $\text{AnSpec}(\mathcal{A}, \mathcal{M})$. In fact, X^{an} represents the functor taking an analytic space Y over $\text{AnSpec}(\mathcal{A}, \mathcal{M})$ to maps $f : |Y| \rightarrow X$ together with a map $f^* \mathcal{O}_X \rightarrow \mathcal{O}_Y$ of sheaves of animated $\mathcal{A}(*)$ -algebras that induces local maps on local rings. Moreover, there is a fully faithful functor $\mathcal{D}_{\text{qc}}(X) \hookrightarrow \mathcal{D}(X^{\text{an}})$.*

REMARK 14.6. The analytification of $\mathbb{A}_{\mathbb{C}}^1$ in this sense is $\text{AnSpec}(\mathbb{C}[T], \mathcal{M}_{<p}[T])$, taking the usual ring of polynomials over \mathbb{C} with the condensed structure and the analytic structure induced from $(\mathbb{C}, \mathcal{M}_{<p})$. This is not what is usually regarded as the analytification of $\mathbb{A}_{\mathbb{C}}^1$, which would be the increasing union of open balls of radius r around the origin, letting $r \rightarrow \infty$. In the present situation, this is still a steady subspace of $\text{AnSpec}(\mathbb{C}[T], \mathcal{M}_{<p}[T])$. The latter is however larger, and contains points “infinitesimally close to infinity”.

We see that the present framework of analytic spaces makes it possible to have analytic spaces that are globally algebraic but locally analytic – in the sense that the global functions

of $\text{AnSpec}(\mathbb{C}[T], \mathcal{M}_{<p}[T])$ are just the polynomials $\mathbb{C}[T]$, but locally one gets certain algebras of convergent functions.

Let us now turn to adic spaces.

PROPOSITION 14.7. *Let (A, A^+) be a Huber pair and $X = \text{AnSpec}((A, A^+)_{\blacksquare})$. The underlying locale $|X|$ maps to the topological space $\text{Spa}(A, A^+)$ defined by Huber. Moreover, for any rational subspace $U \subset \text{Spa}(A, A^+)$ with preimage $U' \subset X$, the map*

$$(A, A^+)_{\blacksquare} \rightarrow (\mathcal{O}_X(U'), \mathcal{M}_X(U'))$$

extends uniquely to an isomorphism

$$(\mathcal{O}_{\text{Spa}(A, A^+)}(U), \mathcal{O}_{\text{Spa}(A, A^+)}^+(U))_{\blacksquare} \cong (\mathcal{O}_X(U'), \mathcal{M}_X(U'))$$

in the following cases:

- (1) *if A is discrete;*
- (2) *if A admits a noetherian ring of definition A_0 over which it is finitely generated;*
- (3) *if A is Tate and sheafy.*

REMARK 14.8. Let us quickly indicate the construction of the map $|X| \rightarrow \text{Spa}(A, A^+)$. An element $a \in A$ is defined to be ≤ 1 on some affine subspace $U \subset X$ if and only if the $(\mathcal{O}_X(U), \mathcal{M}_X(U))$ -module $\mathbb{Z}((T^{-1})) \otimes_{\mathbb{Z}[T]} (\mathcal{O}_X(U), \mathcal{M}_X(U))$ vanishes, where T is mapped to a . Extending this definition to local fractions $\frac{a}{b}$, this defines a valuation on A locally on X . This valuation is however a priori not continuous. Let $\text{Spv}(A, A^+)$ be the spectral space of all (not necessarily continuous) valuations $|\cdot|$ on A with $|A^+| \leq 1$ and $|A^{\circ\circ}| < 1$. Then $\text{Spa}(A, A^+) \subset \text{Spv}(A, A^+)$ and there is a retraction, cf. [**Hub93**, Proposition 2.6, Theorem 3.1]. Now the construction above gives a natural map $|X| \rightarrow \text{Spv}(A, A^+)$, functorial in (A, A^+) . Composing with the retraction $\text{Spv}(A, A^+) \rightarrow \text{Spa}(A, A^+)$ gives the desired map $|X| \rightarrow \text{Spa}(A, A^+)$.

We warn the reader that the retraction $\text{Spv}(A, A^+) \rightarrow \text{Spa}(A, A^+)$ is not natural in (A, A^+) for non-adic maps $(A, A^+) \rightarrow (B, B^+)$ (and hence neither is $|X| \rightarrow \text{Spa}(A, A^+)$). For example, pulling back

$$\text{Spa}(\mathbb{Z}[T^{\pm 1}], \mathbb{Z}) \subset \text{Spa}(\mathbb{Z}[T], \mathbb{Z})$$

to $\text{Spa}(\mathbb{Q}_p\langle T \rangle, \mathbb{Z}_p\langle T \rangle)$, one gets a punctured unit disc, which is not quasicompact anymore. In the context of analytic spaces, the pullback would be given by an analytic ring structure on $\mathbb{Q}_p\langle T \rangle[T^{-1}]$, which is not a Huber ring anymore. Relatedly, there are points of the analytic space $\text{AnSpec}((\mathbb{Q}_p\langle T \rangle, \mathbb{Z}_p\langle T \rangle)_{\blacksquare})$ that are “infinitesimally close to the origin”. These will map in $\text{Spa}(\mathbb{Q}_p\langle T \rangle, \mathbb{Z}_p\langle T \rangle)$ to the origin, while their image in $\text{AnSpec}((\mathbb{Z}[T], \mathbb{Z})_{\blacksquare})$ will not map to the origin of $\text{Spa}(\mathbb{Z}[T], \mathbb{Z})$.

A different way to understand the situation is to regard only $\text{Spv}(A, A^+)$ and the map $|X| \rightarrow \text{Spv}(A, A^+)$ as fundamental, and then observe that Huber was only able to define the structure on those rational subsets of $\text{Spv}(A, A^+)$ that arise via pullback from $\text{Spa}(A, A^+)$, as only on those rational subsets one (usually) gets analytic rings corresponding to Huber pairs again.

The proposition says that in cases (1) – (3), Huber has defined the “correct” structure sheaf. From the proposition, one sees that there is a functor from adic spaces glued out of $\text{Spa}(A, A^+)$ ’s with A satisfying one of the given conditions, to analytic spaces.

One can also use the results of these lectures to recover (and slightly generalize) the results on gluing of vector bundles (or coherent sheaves), cf. e.g. [KL19]. For this, note that from Proposition 12.18 one formally gets descent for dualizable objects. Now we have the following result.

THEOREM 14.9. *Let (A, A^+) be a Huber pair. Then the dualizable objects in $\mathcal{D}((A, A^+)_{\blacksquare})$ are equivalent to the perfect complexes of A -modules. A similar result holds true for the analytic spaces associated to topological, smooth or real-analytic manifolds, or complex-analytic spaces: taking an algebra \mathcal{A} of overconvergent continuous, smooth, real-analytic, or complex-analytic functions on some ball, the dualizable objects of $\mathcal{D}((\mathbb{R}, \mathcal{M}_{<p}) \otimes_{\mathbb{R}} \mathcal{A})$ are equivalent to the perfect complexes of $\mathcal{A}(\ast)$ -modules.*

Generally, for any analytic ring $(\mathcal{A}, \mathcal{M})$, the dualizable objects in $\mathcal{D}(\mathcal{A}, \mathcal{M})$ are generated (under cones, shifts, and retracts) by the cones of $1 - f$ on $\mathcal{M}[S]$, ranging over extremally disconnected profinite sets S and trace-class maps $f : \mathcal{M}[S] \rightarrow \mathcal{M}[S]$.

REMARK 14.10. One can show (for any analytic ring $(\mathcal{A}, \mathcal{M})$) that for perfect complexes the perfect amplitude descends, giving in particular descent for vector bundles.

Let us end these lectures by giving one somewhat funny example of an analytic ring; it is essentially the Novikov ring appearing in symplectic geometry.

For any $0 < r < 1$, consider the ring

$$\mathbb{Z}((T^{\mathbb{R}}))_r = \left\{ \sum_{x \in \mathbb{R}} a_x T^x \mid \sum_x |a_x| r^x < \infty \right\},^{29}$$

equipped with the following condensed ring structure. Write

$$\mathbb{Z}((T^{\mathbb{R}}))_r = \bigcup_{c > 0} \mathbb{Z}((T^{\mathbb{R}}))_{r, \leq c}$$

where

$$\mathbb{Z}((T^{\mathbb{R}}))_{r, \leq c} = \left\{ \sum_{x \in \mathbb{R}} a_x T^x \mid \sum_x |a_x| r^x \leq c \right\},$$

which we give the structure of a compact Hausdorff space by writing it as a quotient of the compact Hausdorff space

$$\{(x_1^+, x_1^-, x_2^+, x_2^-, \dots) \mid x_i^+, x_i^- \in [(\log(c) - \log(i))/\log r, \infty], \sum_i (r^{x_i^+} + r^{x_i^-}) \leq c\}$$

via

$$(x_1^+, x_1^-, x_2^+, x_2^-, \dots) \mapsto \sum_i (T^{x_i^+} - T^{x_i^-}).$$

In particular, the map $\mathbb{R} \cup \{\infty\} \rightarrow \mathbb{Z}((T^{\mathbb{R}}))_r : x \mapsto T^x$ defines a map of condensed sets, and hence $0 = T^\infty$ is connected to $1 = T^0$ in $\mathbb{Z}((T^{\mathbb{R}}))_r$. We note that $\mathbb{Z}((T^{\mathbb{R}}))_r$ has a rather peculiar condensed structure, in that a null-sequence x_0, x_1, \dots need not go to 0 in the $\leq c$ -sense: For example, the sequence

$$1 - T, 1 - T^{1/2}, \dots, 1 - T^{1/n}, \dots$$

converges to $1 - T^0 = 1 - 1 = 0$, so is a null-sequence, but none of the terms lies in $\mathbb{Z}((T^{\mathbb{R}}))_{r, \leq 1}$.

²⁹In particular, at most countably many a_x are nonzero.

Again, we can define similarly for any finite set S a subset

$$\mathbb{Z}((T^{\mathbb{R}}))[S]_{r, \leq c} \subset \mathbb{Z}((T^{\mathbb{R}}))[S]_r,$$

and then define for a profinite set $S = \varprojlim_i S_i$

$$\mathcal{M}(S, \mathbb{Z}((T^{\mathbb{R}}))_r)_{\leq c} = \varprojlim_i \mathbb{Z}((T^{\mathbb{R}}))[S_i]_{r, \leq c},$$

and

$$\mathcal{M}(S, \mathbb{Z}((T^{\mathbb{R}}))_r) = \bigcup_{c>0} \mathcal{M}(S, \mathbb{Z}((T^{\mathbb{R}}))_r)_{\leq c}.$$

Finally, we also define

$$\mathbb{Z}((T^{\mathbb{R}}))_{>r} = \bigcup_{r'>r} \mathbb{Z}((T^{\mathbb{R}}))_{r'}$$

and the same for a profinite set S .

Note that there is an action of the condensed semigroup $\mathbb{R}_{\geq 1}$ on $\mathbb{Z}((T^{\mathbb{R}}))_{>r}$ via $T \mapsto T^t$ for $t \in \mathbb{R}_{\geq 1}$. Restricted to a prime $p \in \mathbb{R}_{\geq 1}$, this action defines a Frobenius lift, so $\mathbb{Z}((T^{\mathbb{R}}))_{>r}$ is a novel kind of λ -ring where the individual Frobenius lifts combine into an action of $\mathbb{R}_{\geq 1}$. Note that the $\mathbb{R}_{\geq 1}$ -action actually also induces canonical isomorphisms between $\mathbb{Z}((T^{\mathbb{R}}))_{>r}$ for different values of r , so this ring is actually independent of the choice of r (up to canonical isomorphism).

THEOREM 14.11. *This construction defines an analytic ring structure on $\mathbb{Z}((T^{\mathbb{R}}))_{>r}$. In fact, it is induced from the analytic ring structure on $\mathbb{Z}((T))_{>r}$ via base change along $\mathbb{Z}[T] \rightarrow \mathbb{Z}[T^{\mathbb{R}}]$.*

Analytic geometry over $\mathbb{Z}((T^{\mathbb{R}}))_{>r}$ is then some funny crossover between arithmetic geometry and complex or real-analytic geometry, allowing actions of connected groups like $\mathbb{R}_{\geq 1}$ in characteristic p ; we believe it is unlike anything that has been studied.


What does the corresponding analytic space look like? It maps at least to the Berkovich spectrum of \mathbb{Z} :

PROPOSITION 14.12. *There is a natural map from $|\mathrm{AnSpec}(\mathbb{Z}((T))_{>r})|$ to the Berkovich spectrum $M(\mathbb{Z})$ of \mathbb{Z} , of all multiplicative norms $|\cdot| : \mathbb{Z} \rightarrow \mathbb{R}_{\geq 0}$. Restricting to $|\mathrm{AnSpec}(\mathbb{Z}((T^{\mathbb{R}}))_{>r})|$, the map*

$$|\mathrm{AnSpec}(\mathbb{Z}((T^{\mathbb{R}}))_{>r})| \rightarrow M(\mathbb{Z})$$

is equivariant for the action of $\mathbb{R}_{\geq 1}$ (acting via rescaling the norm on $M(\mathbb{Z})$).

In this picture, the various theories $(\mathbb{R}, \mathcal{M}_{<p})$ map to the point $p \in (0, 1]$ on the half-line corresponding to real valuations. In particular, taking the limit for $p \rightarrow 0$ makes one enter the arithmetic part of the Berkovich space $M(\mathbb{Z})$, so we see that the analytic ring structures on \mathbb{R} are inextricably linked to arithmetic, which arises in the limit $p \rightarrow 0$.

FIGURE 1. The Berkovich space $M(\mathbb{Z})$

Bibliography

- [ASC⁺16] A. Avilés, F. C. Sánchez, J. M. F. Castillo, M. González, and Y. Moreno, *Separably injective Banach spaces*, Lecture Notes in Mathematics, vol. 2132, Springer, [Cham], 2016.
- [BBK17] O. Ben-Bassat and K. Kremnizer, *Non-Archimedean analytic geometry as relative algebraic geometry*, Ann. Fac. Sci. Toulouse Math. (6) **26** (2017), no. 1, 49–126.
- [Bei07] A. Beilinson, *Topological ϵ -factors*, Pure Appl. Math. Q. **3** (2007), no. 1, Special Issue: In honor of Robert D. MacPherson. Part 3, 357–391.
- [BK72] A. K. Bousfield and D. M. Kan, *Homotopy limits, completions and localizations*, Lecture Notes in Mathematics, Vol. 304, Springer-Verlag, Berlin-New York, 1972.
- [BS15] B. Bhatt and P. Scholze, *The pro-étale topology for schemes*, Astérisque (2015), no. 369, 99–201.
- [CC15] Alain Connes and Caterina Consani, *Universal thickening of the field of real numbers*, Advances in the theory of numbers, Fields Inst. Commun., vol. 77, Fields Inst. Res. Math. Sci., Toronto, ON, 2015, pp. 11–74.
- [CCR⁺82] R. R. Coifman, M. Cwikel, R. Rochberg, Y. Sagher, and G. Weiss, *A theory of complex interpolation for families of Banach spaces*, Adv. in Math. **43** (1982), no. 3, 203–229.
- [Cem84] P. Cembranos, *$C(K, E)$ contains a complemented copy of c_0* , Proc. Amer. Math. Soc. **91** (1984), no. 4, 556–558.
- [Con11] Alain Connes, *The Witt construction in characteristic one and quantization*, Noncommutative geometry and global analysis, Contemp. Math., vol. 546, Amer. Math. Soc., Providence, RI, 2011, pp. 83–113. MR 2815131
- [ČS19] K. Česnavičius and P. Scholze, *Purity for flat cohomology*, arXiv:1912.10932, 2019.
- [CSCK15] F. Cabello Sánchez, J. M. F. Castillo, and N. J. Kalton, *Complex interpolation and twisted twisted Hilbert spaces*, Pacific J. Math. **276** (2015), no. 2, 287–307.
- [DFS08] J. Diestel, J. H. Fourie, and J. Swart, *The metric theory of tensor products*, American Mathematical Society, Providence, RI, 2008, Grothendieck’s résumé revisited.
- [DP61] A. Dold and D. Puppe, *Homologie nicht-additiver Funktoren. Anwendungen.*, Ann. Inst. Fourier **11** (1961), 201–312 (German).
- [Enf73] P. Enflo, *A counterexample to the approximation problem in Banach spaces*, Acta Math. **130** (1973), 309–317.
- [Gro55] A. Grothendieck, *Produits tensoriels topologiques et espaces nucléaires*, Mem. Amer. Math. Soc. **No. 16** (1955), Chapter 1: 196 pp.; Chapter 2: 140.
- [Har84] D. Harbater, *Convergent arithmetic power series*, Amer. J. Math. **106** (1984), no. 4, 801–846.
- [Hub93] R. Huber, *Continuous valuations*, Math. Z. **212** (1993), no. 3, 455–477.
- [Kal81] N. J. Kalton, *Convexity, type and the three space problem*, Studia Math. **69** (1980/81), no. 3, 247–287.
- [KL19] K. S. Kedlaya and R. Liu, *Relative p -adic Hodge theory, II: Imperfect period rings*, arXiv:1602.06899v3, 2019.
- [KP79] N. J. Kalton and N. T. Peck, *Twisted sums of sequence spaces and the three space problem*, Trans. Amer. Math. Soc. **255** (1979), 1–30.
- [Lur09] J. Lurie, *Higher topos theory*, Annals of Mathematics Studies, vol. 170, Princeton University Press, Princeton, NJ, 2009.
- [Lur18] ———, *Spectral algebraic geometry*, 2018.
- [NS18] T. Nikolaus and P. Scholze, *On topological cyclic homology*, Acta Math. **221** (2018), no. 2, 203–409.

- [Rib79] M. Ribe, *Examples for the nonlocally convex three space problem*, Proc. Amer. Math. Soc. **73** (1979), no. 3, 351–355.
- [Ros07] J. Rosický, *On homotopy varieties*, Adv. Math. **214** (2007), no. 2, 525–550.
- [Sch19] P. Scholze, *Lectures on Condensed Mathematics*, 2019.
- [Smi52] M. F. Smith, *The Pontrjagin duality theorem in linear spaces*, Ann. of Math. (2) **56** (1952), 248–253.
- [TV08] B. Toën and G. Vezzosi, *Homotopical algebraic geometry. II. Geometric stacks and applications*, Mem. Amer. Math. Soc. **193** (2008), no. 902, x+224. MR 2394633